

Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

- 1 Doonan, S.: Chromatography on hydroxyapatite. *Methods Mol. Biol. (Totowa)*, 59(Protein Purification Protocols) (1996) 211-215; C.A., 125 (1996) 29337a.
- 2 Doonan, S.: Practical column chromatography. *Methods Mol. Biol. (Totowa)*, 59(Protein Purification Protocols) (1996) 381-396; C.A., 125 (1996) 29338b.
- 3 Eksteen, R.: Column selection in high performance liquid chromatography. In: Katz, E.D. (Editor), *High Perform. Liq. Chromatogr.: Princ. Methods Biotechnol.*, Wiley, Chichester, 1996, pp. 95-162; C.A., 125 (1996) 29340w.
- 4 Hunkeler, D., Janko, M. and Berek, D.: Review of critical conditions of adsorption and limiting conditions of solubility in the liquid chromatography of macromolecules. *ACS Symp. Ser.*, 635 (1996) 250-261; C.A., 125 (1996) 59488t - a review with 32 refs.
- 5 Janis, L.J., Kovach, P.M., Riggin, R.M. and Towns, J.K.: Protein liquid chromatographic analysis in biotechnology. *Methods Enzymol.*, 271(High Resolution Separation and Analysis of Biological Macromolecules) (1996) 86-113; C.A., 125 (1996) 215967f - a review with 43 refs.
- 6 Jerabek, K.: Inverse steric exclusion chromatography as a tool for morphology characterization. *ACS Symp. Ser.*, 635 (1996) 211-224; C.A., 125 (1996) 59698m - a review with 20 refs.
- 7 Kalinichev, A.I.: (Nonlinear theory of multicomponent sorption dynamics and chromatography). *Usp. Khim.*, 65 (1996) 103-124; C.A., 125 (1996) 96686b - a review with 224 refs.
- 8 Kasicka, V. (Editor): *Special Issue of the 13th Symposium on Biomedical Applications of Chromatography and Electrophoresis, held September 4-7, 1995, Prague, Czech Republic*. In: *J. Chromatogr. B*, Vol. 681, No. 1, Elsevier, Amsterdam, 1996, 211 pp.; C.A., 125 (1996) 81226c.
- 9 Laubscher, J.C.: (Chromatography). *Schweiz. Lab.-Z.*, 53 (1996) 119-123; C.A., 125 (1996) 47688p - a review with no refs.
- 10 Neefe, D.A. and Bertini, J.L.: Chromatograph applications and problems from a user's standpoint. *Proc. Int. Sch. Hydrocarbon Meas.*, 70th (1995) 509-513; C.A., 125 (1996) 37580q - a review with no refs.
- 11 Neefe, D.A. and Bertini, J.L.: Chromatograph; applications and problems from a user's standpoint. *Proc. Int. Sch. Hydrocarbon Meas.*, 71st (1996) 525-529; C.A., 125 (1996) 37527c - a review with no refs.

- 12 Novotny, M.V.: Microcolumn liquid chromatography in biochemical analysis. *Methods Enzymol.*, 270(High Resolution Separation and Analysis of Biological Macromolecules, Pt. A) (1996) 101-133; C.A., 125 (1996) 162393g - a review with 25 refs.
- 13 Ouchi, G.I.: Really getting to know noise. *LC-GC*, 14 (1996) 472-476; C.A., 125 (1996) 74746u - a review with 6 refs.
- 14 Sheehan, D. and FitzGerald, R.: Ion-exchange chromatography. *Methods Mol. Biol. (Totowa)*, 59(Protein Purification Protocols) (1996) 145-150; C.A., 125 (1996) 29336z.
- 15 Tsuda, T. (Editor): *Electric Field Applications in Chromatography, Industrial and Chemical Processes*. VCH, Weinheim, 1995, 311 p.; C.A., 125 (1996) 146288w.
- 16 Vailaya, A. and Horvath, C.: Retention thermodynamics in hydrophobic interaction chromatography. *Ind. Eng. Chem. Res.*, 35 (1996) 2964-2981; C.A., 125 (1996) 189633x - a review with 166 refs.
- 17 Wen, J., Arakawa, T. and Philo, J.S.: Size-exclusion chromatography with on-line light-scattering, absorbance, and refractive index detectors for studying proteins and their interactions. *Anal. Biochem.*, 240 (1996) 155-166.

See also 21, 34, 64, 67, 68, 69, 85, 89, 99, 103, 107, 118, 120, 123, 125, 140, 146, 147, 150, 155, 165, 173, 178, 181, 186, 190, 242, 268, 276, 277, 290, 352, 354, 410, 413, 465, 475, 476, 478, 496, 521, 522, 530, 542, 548, 551, 562, 565, 640, 646, 647, 663, 665, 668, 678, 679, 690, 692, 695, 716, 721, 735, 742, 745, 749, 753, 762, 806, 829, 950, 959, 968, 973, 982, 983, 985, 987, 989, 1123, 1153, 1172, 1189, 1192, 1219, 1235.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 18 Alltech Associated: Electrochemically regenerated ion suppression (ERIS) improves and simplifies suppressor-based ion chromatography. *Chem. N.Z.*, 60 (1996) 3-5; C.A., 125 (1996) 131004b.
- 19 Bentley, M., Borlak, J. and Tenant, K.: Post-validation problems with incurred samples. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 327-328; C.A., 125 (1996) 269757f.

- 20 Bjoerkqvist, B.: (High concentrations - a piece of cake?). *Kem.-Kemi*, 23 (1996) 83-85; C.A., 125 (1996) 47714a.
- 21 Bruessau, R.J.: Experiences with interlaboratory GPC experiments. *Macromol. Symp.*, 110 (1996) 15-32; C.A., 125 (1996) 184430u - a review with 7 refs.
- 22 Choudhary, G. and Horvath, C.: Ion-exchange chromatography. *Methods Enzymol.*, 270(High Resolution Separation and Analysis of Biological Macromolecules, Pt. A) (1996) 47-82; C.A., 125 (1996) 162453b.
- 23 Corradini, D.: Effects of electrostatic and hydrophobic interactions on the chromatographic behavior of biopolymers in HPLC columns. *Macromol. Symp.*, 110(10th Bratislava Int. Conference on Macromolecules: Chromatography of Polymers and Related Substances, 1995) (1996) 57-63; C.A., 125 (1996) 189731c.
- 24 Dolan, J.W.: Improving an ion-pairing method. *LC-GC Int.*, 9 (1996) 624-628.
- 25 Fornstedt, T., Zhong, G. and Guiochon, G.: Peak tailing and slow mass transfer kinetics in nonlinear chromatography. *J. Chromatogr. A*, 742 (1996) 55-68.
- 26 Fornstedt, T., Zhong, G. and Guiochon, G.: Peak tailing and mass transfer kinetics in linear chromatography. *J. Chromatogr. A*, 741 (1996) 1-12.
- 27 Gargallo, R., Tauler, R., Cuesta-Sánchez, F. and Massart, D.L.: Validation of alternating least-squares multivariate curve resolution for chromatographic resolution and quantitation. *TrAC*, 15 (1996) 279-286.
- 28 Guo, Y., Xiang, B. and An, D.: (Expert system for chromatographic fault diagnosis). *Zhongguo Yaoake Daxue Xuebao*, 27 (1996) 126-128; C.A., 125 (1996) 242047n.
- 29 Kamiyama, F., Yamazaki, K., Kawamura, K. and Kohara, M.: Temperature dependence of retention in reversed-phase liquid chromatography on a porous acrylic support. *Biomed. Chromatogr.*, 10 (1996) 105-110; C.A., 125 (1996) 80886f.
- 30 McDowell, R.D.: How good is your laboratory? *LC-GC Int.*, 9 (1996) 630-636.
- 31 Miao, H., Yu, M. and Hu, S.: Artificial neural networks aided deconvolving overlapped peaks in chromatograms. *J. Chromatogr. A*, 749 (1996) 5-11.
- 32 Ooms, B.: Temperature control in high performance liquid chromatography. *LC-GC Int.*, 9 (1996) 574-585.
- 33 Outinen, K., Vuorela, H. and Hiltunen, R.: Optimization of HPLC separation by computer-assisted optimization methods. Comparison of three methods. *Eur. J. Pharm. Sci.*, 4 (1996) 199-210; C.A., 125 (1996) 49483y.
- 34 Potschka, M.: Inverse size exclusion chromatography and universal calibration. *Macromol. Symp.*, 110.(1996) 121-154; C.A., 125 (1996) 184431v - a review with 135 refs.
- 35 Southern, D.L. and Dolan, J.W.: Solvent changeover and column equilibration. *LC-GC Int.*, 9 (1996) 706-708.
- 36 Strancar, A., Koselj, P., Schwin, H. and Josic, D.: Application of compact porous disks for fast separations of biopolymers and in-process control in biotechnology. *Anal. Chem.*, 68 (1996) 3483-3488.
- 37 Tamura, A., Khademizadeh, M., Razee, S. and Masujima, T.: Direct observation of solutes trapping on a surface-inert reversed phase column. *Chem. Lett.*, (1996) 475-476; C.A., 125 (1996) 80898m.
- 38 Thordarson, E., Palmarsdottir, S., Mathiasson, L. and Jónsson, J.A.: Sample preparation using a miniaturized supported liquid membrane device connected on-line to packed capillary liquid chromatography. *Anal. Chem.*, 68 (1996) 2559-2563.
- 39 Vissers, J.P.C., de Ru, A.H., Ursem, M. and Chervet, J.-P.: Optimised injection techniques for micro and capillary liquid chromatography. *J. Chromatogr. A*, 746 (1996) 1-7.
- 40 Westerlund, D., Fornstedt, T., Thorsteinsdóttir, M. and Zouxiang Y.: Peak compression and deformation caused by large system peaks in HPLC. Part I: Peak compression. *LC-GC Int.*, 9 (1996) 747-760.
- For additional information see C.A.:
125 (1996) 137211p.
- See also 9, 57, 62, 96, 133, 137, 479, 512, 516.
- 2b. Thermodynamics and theoretical relationships*
- 41 Bian, L. and Geng, X.: (Equation describing the relationship between injected amount of solutes and their retention values in liquid chromatography). *Sepu*, 14 (1996) 179-181; C.A., 125 (1996) 52656n.
- 42 Fóti, G., Révesz, G., Hajós, P., Pellaton, G. and sz. Kováts, E.: Classical retention mechanism in ion exchange chromatography. Theory and experiment. *Anal. Chem.*, 68 (1996) 2580-2589.
- 43 Guillaume, Y.C. and Guinchard, C.: Retention mechanism of weak polar solutes in reversed phase liquid chromatography. *Anal. Chem.*, 68 (1996) 2869-2873.
- 44 Hamaker, K.H. and Ladisch, M.R.: Intraparticle flow and plate height effects in liquid chromatography stationary phases. *Sep. Purif. Methods*, 25 (1996) 47-83; C.A., 125 (1996) 171828e.
- 45 Heeter, G.A. and Liapis, A.I.: Effects of structural and kinetic parameters on the performance of chromatographic columns packed with perfusive and purely diffusive adsorbent particles. *J. Chromatogr. A*, 743 (1996) 3-14.
- 46 Jansen, M.L., Hofland, G.W., Houwers, J., Straathof, A.J.J., van der Wielen, L.A.M., Luyben, K.C.A.M. and van den Tweel, W.J.J.: Effect of pH and concentration on column dynamics of weak electrolyte ion exchange. *AIChE J.*, 42 (1996) 1925-1937; C.A., 125 (1996) 89963m.
- 47 Meyer, V.R. and Welsch, T.: Statistical peak overlap - an under-valuated aspect of chromatography. *LC-GC Int.*, 9 (1996) 670-681.
- 48 Potschka, M.: A soft-body theory of size exclusion chromatography. *ACS Symp. Ser.*, 635(Strategies in Size Exclusion Chromatography) (1996) 37-87; C.A., 125 (1996) 109245j.
- 49 Rathore, A.S. and Horváth, C.: Separation parameters via virtual migration distances in high-performance liquid chromatography, capillary zone electrophoresis and electrokinetic chromatography. *J. Chromatogr. A*, 743 (1996) 231-246.
- 50 Watanabe, H., Yokoyama, Y. and Sato, H.: Retention volume of the second system peaks in ion-exchange chromatography with monobasic acid eluent systems. *Anal. Sci.*, 12 (1996) 381-384; C.A., 125 (1996) 103555p.
- For additional information see C.A.:
125 (1996) 68634q.

See also 7, 10, 25, 26, 31, 104, 107, 112, 124, 128, 168, 182, 196, 959, 1001.

2c. Relationship between structure and chromatographic behaviour

- 51 De Beer, J.O., Vandenbroucke, C.V., Massart, D.L. and de Spiegeler, B.M.: Half-fraction and full factorial designs versus central composite design for retention modelling in reversed-phase ion-pair liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 525-541.
- 52 Forgács, E. and Cserháti, T.: Canonical correlation analysis of the relationship between physicochemical parameters of steroid drugs and their retention characteristics in HPLC. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1849-1858.
- 53 Hong, H., Zhou, D., Han, S., Wang, L., Zhang, Z. and Zou, G.: Retention of aromatic sulfur-containing compounds on RP-HPLC: correlation with partition coefficients and molecular connectivity indices. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1617-1629.

See also 54, 815, 1119.

2d. Measurement of physico-chemical and related values

- 54 Jenke, D.R.: Liquid chromatographic estimation of octanol/water partition coefficients with a high efficiency, nonporous, ultrasmall particle size reverse phase stationary phase. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2227-2245.
- 55 Mlynar, J. and Sarkany, S.: Renaissance in ultracentrifugal sedimentation equilibrium calibrations of size exclusion chromatographic elution profiles. *ACS Symp. Ser.*, 635 (1996) 379-400; C.A., 125 (1996) 61221u.
- 56 Mori, S., Kato, H. and Nishimura, Y.: Problems of baseline construction for calculation of molecular weight averages in size exclusion chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2077-2087.
- 57 Solcova, O. and Schneider, P.: Extra-column effects in determination of rare parameters by the chromatographic method. *Collect. Czech. Chem. Commun.*, 61 (1996) 844-855; C.A., 125 (1996) 118459j.
- 58 Wang, J., Sakoda, A. and Suzuki, M.: (Chromatographic evaluation of microbial activity in a biological activated carbon bed). *Kagaku Kogaku Ronbunshu*, 22 (1996) 948-952; C.A., 125 (1996) 94454a.

For additional information see C.A.:

125 (1996) 131298a.

See also 135, 197, 206, 274, 951, 952, 953, 954, 956, 957, 965, 967, 970, 1074, 1100, 1219.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 59 Dolan, J.W.: Injection loop adsorption. *LC-GC Int.*, 9 (1996) 530-533.

- 60 Dolan, J.W.: Injection loop adsorption in liquid chromatography. *LC-GC*, 14 (1996) 562-566; C.A., 125 (1996) 157300v.
- 61 Schaefer, H., Laebli, M. and Zaehner, P.: Device for ion chromatography and process for cyclic regeneration of several suppressors in such a device. *Eur. Pat. Appl. EP 725,272* (Cl. G01N30/96), 7 Aug. 1996, CH Appl. 95/272, 1 Feb. 1995; 14 p.; C.A., 125 (1996) 184401k.

For additional information see C.A.:

125 (1996) 47852u, 61813p, 96954n, 103744z, 103773h, 103778p, 157397g, 171684e.

See also 38, 74, 190, 191.

3b. Detectors and detection reagents

- 62 Acworth, I.N. and Gamache, P.H.: The coulometric electrode array for use in HPLC analysis. Part 1: Theory. *Am. Lab. (Shelton)*, 28 (1996) 33-38; C.A., 125 (1996) 47921r.
- 63 Eberz, G., Rast, H.-G., Burger, K., Kreiss, W. and Weisemann, C.: Bioactive screening by chromatography-bioluminescence coupling. *Chromatographia*, 43 (1996) 5-9.
- 64 Imasaka, T.: Liquid chromatography and capillary electrophoresis with laser fluorometric detection. *Methods Chromatogr.*, 1 (1996) 123-149; C.A., 125 (1996) 103814x - a review with 31 refs.
- 65 Rezai, M.A., Famiglini, G. and Cappiello, A.: Enhanced detection sensitivity by large volume injection in reversed-phase micro-high-performance liquid chromatography. *J. Chromatogr. A*, 742 (1996) 69-78.
- 66 Richards, D.S., Davidson, S.M. and Holt, R.M.: Detection of non-UV-absorbing chiral compounds by high-performance liquid chromatography-mass spectrometry. *J. Chromatogr. A*, 746 (1996) 9-15.
- 67 Szulc, M., Mhatre, R., Mazzeo, J. and Krull, I.S.: Detection in liquid chromatography. *Methods Enzymol.*, 270(High Resolution Separation and Analysis of Biological Macromolecules, Pt. A) (1996) 175-200; C.A., 125 (1996) 215923p - a review with 66 refs.
- 68 Takata, Y.: Liquid chromatography with coulometric detector. *Methods Chromatogr.*, (1996) 43-74; C.A., 125 (1996) 103552k - a review with 45 refs.
- 69 Takeuchi, T.: (Signal enhancement assisted by stationary phases in liquid chromatography). *Dojin News*, 79 (1996) 3-8; C.A., 125 (1996) 103807x - a review with 8 refs.
- 70 Weber, G.: Selective detection of metal species in HPLC and FIA by means of pulsed amperometric detection (PAD). *Fresenius J. Anal. Chem.*, 356 (1996) 242-246.
- 71 Wilke, S., Wang, H., Muraczewska, M. and Müller, H.: Amperometric detection of heavy metal ions in ion pair chromatography at an array of water/nitrobenzene micro interfaces. *Fresenius J. Anal. Chem.*, 356 (1996) 233-236.

For additional information see C.A.:

125 (1996) 74978w, 99677k, 103588b, 103608h.

See also 17, 217, 310, 384, 388, 419, 436, 443, 447, 454, 466, 564, 701, 784, 793, 855, 866, 929, 1062, 1115, 1181, 1200, 1226.

- 3c. *Sorbents and columns, packing procedures*
- 72 Bereznitski, Y., Jaroniec, M. and Kruk, M.: Surface and structural properties of silica gels used in high performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1523-1537.
- 73 Carr, P.W., McCormick, A.V., Annen, M.J., Sun, L. and Brown, J.R.: Synthesis of porous inorganic particles by polymerization-induced colloid aggregation (PICA). *U.S. US 5,540,834* (Cl. 210-198.2; B01D15/08), 30 Jul. 1996, Appl. 294,374, 23 Aug. 1994; 42 p.; C.A., 125 (1996) 146420h.
- 74 Cheon, W.J. and An, H.J.: Chromatographic application of microcolumns made of glass-linked stainless steel tubing. III. Preparation of a 0.3 mm I.D. column. *Bull. Korean Chem. Soc.*, 17 (1996) 539-541; C.A., 125 (1996) 157308d.
- 75 Chriswanto, H. and Wallace, G.G.: Redox chromatography using polypyrrrole as a stationary phase. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2457-2476.
- 76 Dasarathy, Y., Ramberg, M. and Andersson, M.: A systematic approach to screening ion-exchange chromatography media for process development. *BioPharm (Eugene)*, 9 (1996) 42-45; C.A., 125 (1996) 273707g.
- 77 Dunlap, C.J. and Carr, P.W.: Synthesis and chromatographic characterization of dextran-coated zirconia high-performance liquid chromatographic stationary phases. *J. Chromatogr. A*, 746 (1996) 199-210.
- 78 Enomoto, N., Furukawa, S., Ogasawara, Y., Akano, H., Kawamura, Y., Yashima, E. and Okamoto, Y.: Preparation of silica gel-bonded amylose through enzyme-catalyzed polymerization and chiral recognition ability of its phenylcarbamate derivative in HPLC. *Anal. Chem.*, 68 (1996) 2798-2804.
- 79 Etchegaray, A., Carvalho, R.S.H., Boschcov, P. and Nakae, C.R.: Estimation of the pK_a of the amine groups of benzhydrylamine resin and its lysyl derivative by measurement of swelling. *Chromatographia*, 43 (1996) 82-84.
- 80 Farkas, T., Sepaniak, M.J. and Guiuchon, G.: Column radial homogeneity in high-performance liquid chromatography. *J. Chromatogr. A*, 740 (1996) 169-181.
- 81 Feibusch, B.: A new chemically deactivated silica-based reverse phase/ion exchange support. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2315-2328.
- 82 Fields, S.M.: Silica xerogel as a continuous column support for high-performance liquid chromatography. *Anal. Chem.*, 68 (1996) 2709-2712.
- 83 Guan-Sajonz, H. and Guiuchon, G.: Effect of the packing pressure on the performance of C₁₈ reversed-phase liquid chromatographic columns. *J. Chromatogr. A*, 743 (1996) 247-259.
- 84 Haag, I.: (A stationary phase with high applicability and selectivity). *LaborPraxis*, 20 (1996) 24-28; C.A., 124 (1996) 359509k.
- 85 Hagel, L.: Characteristics of modern media for aqueous size exclusion chromatography. *ACS Symp. Ser.*, 635(Strategies in Size Exclusion Chromatography) (1996) 225-248; C.A., 125 (1996) 109107r - a review with 42 refs.
- 86 Hagel, L., Östberg, M. and Andersson, T.: Apparent pore size distributions of chromatography media. *J. Chromatogr. A*, 743 (1996) 33-42.
- 87 Hanson, M. and Unger, K.K.: Evaluation of advanced silica packings. Part 1: Synthesis and characterization of non-porous silica particles for HPLC. *LC-GC Int.*, 9 (1996) 650-656.
- 88 Hanson, M. and Unger, K.K.: Evaluation of advanced silica packings. Part II: Application of non-porous silica particles in HPLC. *LC-GC Int.*, 9 (1996) 741-746.
- 89 Horak, D., Hradil, J. and Benes, M.J.: Packings for size exclusion chromatography: preparation and some properties. *ACS Symp. Ser.*, 635 (1996) 190-210; C.A., 125 (1996) 103545k - a review with 86 refs.
- 90 Jelinek, L., Ohmacht, R. and Pfenninger, A.: Monosized, silica-based microspheres for chromatographic separations. *Am. Biotechnol. Lab.*, 14 (1996) 18-20; C.A., 125 (1996) 136792s.
- 91 Ji, Z., Pinon, D.I. and Miller, L.J.: Development of magnetic beads for rapid and efficient metal-chelate affinity purification. *Anal. Biochem.*, 240 (1996) 197-201.
- 92 Kaltenbach, P., Swedberg, S.A., Witt, K.E., Bek, F. and Mittelstadt, L.S.: Miniaturized planar columns for use in a liquid phase separation apparatus. *PCT Int. Appl. WO 96 12,546* (Cl. B01D15/08), 2 May 1996, US Appl. 326,111, 19 Oct. 1994; 72 p.; C.A., 125 (1996) 37247m.
- 93 Kawabata, M., Ootsuki, K. and Sato, K.: (Fibrous stationary phase for chromatography of viscous substances). *Jpn. Kokai Tokkyo Koho JP 08,170,958* [96,170,958] (Cl. GO1N30/48), 2 Jul. 1996, Appl. 94/315,211, 19 Dec. 1994; 4 pp.; C.A., 125 (1996) 162747a.
- 94 Kirkland, J.J.: Stability of silica-based, monofunctional C18 bonded-phase column packing for HPLC at high pH. *J. Chromatogr. Sci.*, 34 (1996) 309-313.
- 95 Kubota, N., Kounosu, M., Saito, K., Sugita, K., Watanabe, K. and Sugo, T.: Control of phenyl-group site introduced on the graft chain for hydrophobic interaction chromatography. *React. Funct. Polym.*, 29 (1996) 115-122; C.A., 125 (1996) 80874a.
- 96 Levison, P.R. and Pathirana, N.D.: Rapid ion chromatographic method for the quantitative determination of ligand leakage from ion-exchange media. *J. Chromatogr. A*, 742 (1996) 79-86.
- 97 Li, J. and Carr, P.W.: Retention characteristics of polybutadiene-coated zirconia and comparison to conventional bonded phases. *Anal. Chem.*, 68 (1996) 2857-2868.
- 98 Liquan, C., Wei, Z. and Zhong, Z.: Study of ion chromatography with ion-exchange fibers as the stationary phase. *J. Chromatogr. A*, 740 (1996) 195-199.
- 99 Maruska, A., Liesiene, J. and Serys, A.: Regulation of cellulose-based adsorbent granule morphology. *J. Chromatogr. A*, 746 (1996) 147-160 - a review with 58 refs.
- 100 Minakuchi, H., Nakanishi, K., Soga, N., Ishizuka, N. and Tanaka, N.: Octadecylsilylated porous silica rods as separation media for reversed-phase liquid chromatography. *Anal. Chem.*, 68 (1996) 3498-3501.
- 101 Morino, M., Kaneko, H., Takahashi, T. and Iino, M.: Inverse study exclusion chromatography using extraction residues or extracts of coals as a stationary phase. *Energy Fuels*, 10 (1996) 1012-1016; C.A., 125 (1996) 37766e.
- 102 Myers, T.A., Healy, M.G. and Murphy, W.R.: Chitosan beads as support matrixes in dye-ligand affinity chromatography. In: Karnicki, Z.S. (Editor), *Chitin World, [Proc. Int. Conf. Chitin Chitosan]*, 6th 1994, Wirtschaftsverlag NW, Bremerhaven, 1994, pp. 276-283; C.A., 125 (1996) 52657p.
- 103 Oi, N.: Recent progress of chiral stationary phases for HPLC. *Methods Chromatogr.*, 1 (1996) 213-230; C.A., 125 (1996) 103817a - a review with 59 refs.

- 104 Perek, J.J., Matuska, M.T. and Hemphälä, H.: HPLC evaluation of mono-ol, butylphenol and perfluorinated columns prepared via olefin hydrosilation on a silica hydride intermediate. *Chromatographia*, 43 (1996) 10-16.
- 105 Richard, J., Vaslin, S. and Larpent, C.: Glycosylated latex or inorganic oxide particles and their preparation and use for detection of biological materials or adsorption chromatography. *Eur. Pat. Appl. EP 709,402 (Cl. C08F/30)*, 1 May 1996, FR Appl. 94/12,970, 28 Oct. 1994; 17 p.; C.A., 125 (1996) 59448e.
- 106 Ricker, R.D., Sandoval, L.A., Permar, B.J. and Boyes, B.E.: Improved reversed-phase high performance liquid chromatography columns for biopharmaceutical analysis. *J. Pharm. Biomed. Anal.*, 14 (1995) 93-105.
- 107 Rodrigues, A.E.: Intraparticle convection in chromatographic permeable packings. *ACS Symp. Ser.*, 635 (1996) 157-172; C.A., 125 (1996) 36636a - a review with 49 refs.
- 108 Sarker, M. and Guiochon, G.: Consolidation of the packing material in chromatographic columns under dynamic axial compression. III. Effect of the nature of the packing solvent on the consolidation and performance of axial compression columns. *J. Chromatogr. A*, 741 (1996) 165-173.
- 109 Simova, M. and Dimov, N.: Retention on diol column - a case study. *Chromatographia*, 43 (1996) 436-438.
- 110 Spanos, E. and Buchberger, W.: Preparation of inorganic ion exchangers for ion chromatography. *Am. Lab. (Shelton)*, 28 (1996) 18B-18D; C.A., 125 (1996) 131003a.
- 111 Stanley, B.J., Sarker, M. and Guiochon, G.: Consolidation of the packing material in chromatographic columns under dynamic axial compression. IV. Mechanical properties of some packing materials. *J. Chromatogr. A*, 741 (1996) 175-184.
- 112 Steiner, F., Niederländer, C. and Engelhardt, H.: Optimization of alkali-earth cation separation on weak cation exchanger. *Chromatographia*, 43 (1996) 117-123.
- 113 Svec, F. and Frechet, J.M.J.: Molded separation media. An inexpensive efficient, and versatile alternative to packed columns for the fast HPLC separation of peptides, proteins, and synthetic oligomers and polymers. *Macromol. Symp.*, 110 (1996) 203-216; C.A., 125 (1996) 170072s.
- 114 Sviderskii, V.A., Klimenko, V.S. and Klimenko, S.V.: Studies of the textural features of alkoxide gels of silica. *Colloid J.*, 58 (1996) 377-379; C.A., 125 (1996) 178373m.
- 115 Venema, E., Kraak, J.C., Poppe, H. and Tijssen, R.: Packed-column hydrodynamic chromatography using 1-μm non-porous silica particles. *J. Chromatogr. A*, 740 (1996) 159-167.
- 116 Xiao, J. and Meyerhoff, M.E.: Retention behavior of amino acids and peptides on protoporphyrin - silica stationary phases with varying metal ion centres. *Anal. Chem.*, 68 (1996) 2818-2825.
- For additional information see C.A.:
- 125 (1996) 34931u, 47863y, 47924u, 118851f, 124606w, 124655m, 124669u, 146682v, 157233a, 157304z, 157341j, 184385h, 189971f, 269846j.
- See also 6, 17, 29, 44, 54, 129, 156, 157, 158, 167, 169, 170, 171, 173, 321, 385, 419, 420, 431, 440, 463, 471, 514, 515, 516, 526, 535, 539, 541, 544, 545, 546, 548, 576, 645, 747, 786, 983, 988, 1014, 1205, 1236.
- 3d. Quantitative analysis
- 117 Cao, G.M.: Quantitation of unresolved chromatographic peak using overlapping ratio of profiles of standard chromatographic peaks. *J. Chromatogr. A*, 746 (1996) 161-167.
- 118 Jemke, D.R.: Chromatographic method validation: a review of current practices and procedures. II. Ruggedness, revalidation and system suitability. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1873-1891 - a review with 41 refs.
- See also 27, 215, 405, 436, 889, 1080.
- 3e. Preparative scale chromatography
- 119 Abou-Shoer, M.: Preparative block chromatography (PCB) in the analysis of natural products. *Alexandria J. Pharm. Sci.*, 10 (1996) 117-119; C.A., 125 (1996) 242063q.
- 120 Cox, G.B.: The preparative HPLC of biomolecules. In: Katz, E.D. (Editor), *High Perform. Liq. Chromatogr.: Princ. Methods Biotechnol.*, Wiley, Chichester, 1996, pp. 293-349; C.A., 125 (1996) 29281c - a review with 31 refs.
- 121 Felinger, A. and Guiochon, G.: Optimizing experimental conditions in overloaded gradient elution chromatography. *Biotechnol. Prog.*, 12 (1996) 638-644; C.A., 125 (1996) 216001y.
- 122 Fernandez, E.J., Norton, T.T., Jung, W.C. and Tsavalas, J.G.: A column design for reducing viscous fingering in size exclusion chromatography. *Biotechnol. Prog.*, 12 (1996) 480-487; C.A., 125 (1996) 80889j.
- 123 Gagnon, P. and Grund, E.: Large-scale process development for hydrophobic interaction chromatography, part 4: controlling selectivity. *BioPharm (Eugene)*, 9 (1996) 54-64; C.A., 125 (1996) 52542x - a review with 85 refs.
- 124 Samsonov, G.V. and Pisarev, O.A.: High capacity reversible sorption and effective chromatographic preparative bioseparation at low pressure. *Isol. Purif.*, 2 (1996) 93-102; C.A., 125 (1996) 136787u.
- 125 Schulte, M., Kinkel, J.N., Nicoud, R.-M. and Charton, F.: Simulated moving bed chromatography (SMB). An efficient technique to producing optically active compounds on an industrial scale. *Chem.-Ing.-Tech.*, 68 (1996) 670-683; C.A., 125 (1996) 89907w - a review with 31 refs.
- 126 Szepesy, L. and Bekassy, S.: (New unit operation: high performance liquid chromatography (HPLC). Part IV. Development and realization of continuous chromatography). *Magy Kem. Lapja*, 51 (1996) 239-246; C.A., 125 (1996) 118334q.

For additional information see C.A.:

125 (1996) 36625w, 171836f.

See also 80, 174, 233, 236, 412, 478, 515, 537, 589, 592, 618, 674, 679, 780, 982, 1080.

3f. Programmed temperature, pressure, vapors, gradients

127 Ryan, K., Djordjevic, N.M. and Erni, F.: Temperature programming in open tubular liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2089-2099.

See also 960, 961.

4. SPECIAL TECHNIQUES

4a. Automation

See 499, 564, 914.

4b. Computerization and modelling

- 128 Busnell, J.-P., Degoulet, C. and Tasin, J.-F.: Modelling of size exclusion chromatography by Monte Carlo simulation. *ACS Symp. Ser.*, 635 (1996) 52-66; *C.A.*, 125 (1996) 59544h.
- 129 Chizhkov, V.P. and Varivonshik, V.E.: (Principles and theory of two-cycle circulation liquid chromatography). *Izv. Akad. Nauk, Ser. Khim.*, (1993) 1535-1538; *C.A.*, 125 (1996) 89937f.
- 130 Huber, L.: Validation of computerized analytical systems, Part II: Setting specifications, qualifying a vendor, and validating user-contributed application software. *LC-GC Int.*, 9 (1996) 666-669.
- 131 Huber, L.: Validation of computerized analytical systems. Part I: overview, regulations, and terminology. *LC-GC Int.*, 9 (1996) 564-572.
- 132 Jiménez, O., Benito, I. and Marina, M.L.: Prediction and modelling studies for capacity factors of a group of dihydropyridines in micellar liquid chromatography with hybrid eluents. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2477-2495.
- 133 Lukulay, P.H. and McGuffin, V.L.: Experimental and computer simulation studies of solute-solute interactions in liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2039-2058.
- 134 Qian, M.G., Wu, J.-T., Parus, S. and Lubman, D.M.: The development of a data system for a combination of liquid chromatography or capillary electrophoresis with an ion trap storage/reflection time-of-flight mass detector. *Rapid Commun. Mass Spectrom.*, 10 (1996) 1209-1214; *C.A.*, 125 (1996) 162688g.
- 135 Soria, V., Campos, A., Figueroel, J.E., Gomez, C., Porcar, I. and Garcia, R.: Modelling of stationary phase in size exclusion chromatography with binary eluents. *ACS Symp. Ser.*, 635 (1996) 103-126; *C.A.*, 125 (1996) 59545j.
- For additional information see *C.A.*:
125 (1996) 57387k, 124670n.
- See also 28, 31, 33, 42, 46, 50, 141, 149, 184, 187, 527, 536, 829, 996, 1075, 1199.
- 4c. Combination with other physico-chemical techniques (MS, IR etc.)
- 136 Buttler, T., Jarskog, H., Gorton, L., Marko-Varga, G. and Ramemark, L.: The use of microdialysis for sampling in column liquid chromatography. *Am. Lab. (Shelton)*, 26 (1996) 281-28M; *C.A.*, 125 (1996) 109271q.
- 137 Ioffe, B.V. and Vitenberg, D.A.: Determination of molecular formulas of organic compounds with the use of the data of mass-spectrometric and atomic-emission chromatographic detectors. *J. Anal. Chem.*, 51 (1996) 807-812; *C.A.*, 125 (1996) 221023m.
- 138 Isobe, T., Nakayama, H., Shinkai, F., Yamaguchi, M., Kanai, M., Ikezawa, H., Seta, K. and Okuyama, T.: (Molecular anatomy of brain using multi-dimensional HPLC-ESI/MS or MALDI/TOFMS). *Kuromatogurafi*, 17 (1996) 160-161; *C.A.*, 125 (1996) 109548k.
- 139 Ito, S., Tomioka, M., Mimura, T., Kato, Y., Hirabayashi, Y., Takada, Y., Hirabayashi, A. and Sakair, M.: (Development of liquid chromatography/mass spectrometry using sonic spray ionization). *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 21 (1996) 117-118; *C.A.*, 125 (1996) 269805v.
- 140 Lindon, J.C., Nicholson, J.K. and Wilson, I.D.: Direct coupling of chromatographic separations to NMR spectroscopy. *Prog. Nud. Magn. Reson. Spectrosc.*, 29 (1996) 1-49; *C.A.*, 125 (1996) 109127x - a review with 61 refs.
- 141 Link, A.J., Eng, J. and Yates, J.R., III: Analyzing complex biological systems using micro-LC-ESI-MS-MS. *Am. Lab. (Shelton)*, 28 (1996) 27-30; *C.A.*, 125 (1996) 216241b.
- 142 Mason, C., Moore, J., Churchill, M., Solanki, P. and Galloway, W.A.: A strategy for the development of bioanalytical methods using LC-MS. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 137-140; *C.A.*, 125 (1996) 237407a.
- 143 Nguyen Hong, T.D., Jouan, M., Quy Dao, N., Bouraly, M. and Mantisi, F.: Coupling of high-performance liquid chromatography with Raman spectrometry. *J. Chromatogr. A*, 743 (1996) 323-327.
- 144 Noctor, T.A.G., Hemsley, M. and Robson, J.N.: Rapid sample-preparation approaches, notably disk-based SPE, for HPLC-MS-MS(API). *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 213-214; *C.A.*, 125 (1996) 242131k.
- 145 Owen, L.M.W., Rauscher, A.M., Fairweather-Tait, S.J. and Crews, H.M.: Use of HPLC with inductively coupled plasma mass spectrometry (ICP-MS) for trace element speciation studies in biological materials. *Biochem. Soc. Trans.*, 24 (1996) 947-952; *C.A.*, 125 (1996) 269496v.
- 146 Quevauviller, P.: Improvement of quality control of speciation analysis using hyphenated techniques. A decade of progress within the European Community. *J. Chromatogr. A*, 750 (1996) 25-33 - a review with 31 refs.
- 147 Regnier, F. and Huang, G.: Future potential of targeted component analysis by multidimensional liquid chromatography-mass spectrometry. *J. Chromatogr. A*, 750 (1996) 3-10 - a review with 26 refs.
- 148 Seta, K., Kanai, M., Ikezawa, H., Yamaguchi, M., Nakayama, H., Shinkai, F., Isobe, T. and Okuyama, T.: (Packed capillary HPLC/MS system using a conventional HPLC and splitters). *Kuromatogurafi*, 17 (1996) 94-95; *C.A.*, 125 (1996) 109547j.
- 149 Vanslyke, S.J.: Enhancements to multidimensional methods in analytical chemistry (Kalman filter, digital filter, principal components analysis, chromatography). Avail. *Univ. Microfilms Int.*, Order No. DANN051175, 1994, 285 p.; *C.A.*, 125 (1996) 130925r.
- 150 Wolfender, J.L. and Hostettmann, K.: Importance of LC/MS (liquid chromatography/mass spectrometry) in plant analysis. *Spectrosc. Eur.*, 8 (1996) 7-12; *C.A.*, 125 (1996) 242005x - a review with 17 refs.
- 151 Zhai, C.: (LC-coupled with ^1H -NMR spectroscopy - currents status and future prospects). *Bopuxue Zazhi*, 13 (1996) 403-410; *C.A.*, 125 (1996) 157642h.

For additional information see C.A.:
 125 (1996) 47846v, 47942y.

See also 66, 186, 200, 203, 210, 215, 227, 240, 254, 269, 293, 301, 303, 328, 351, 357, 387, 401, 402, 408, 415, 430, 465, 472, 484, 491, 503, 511, 543, 552, 556, 563, 573, 574, 575, 601, 611, 689, 826, 828, 834, 860, 863, 895, 905, 906, 907, 910, 914, 916, 920, 925, 928, 948, 949, 957, 958, 971, 976, 980, 981, 994, 1005, 1006, 1029, 1034, 1037, 1059, 1067, 1069, 1071, 1082, 1128, 1149, 1156, 1157, 1159, 1169, 1172, 1177, 1187, 1189, 1192, 1201, 1206, 1212, 1224, 1228, 1230.

4d. Affinity chromatography (advances)

- 152 Brekkan, E., Lundquist, A. and Lundahl, P.: Immobilized membrane vesicle or proteoliposome affinity chromatography. Frontal analysis of interactions of cytochalasin B and D-glucose with the human red cell glucose transporter. *Biochemistry*, 35 (1996) 12141-12145.
 153 Burton, S.J.: Affinity chromatography: production and regulatory considerations. *Am. Biotechnol. Lab.*, 14 (1996) 64-66; C.A., 125 (1996) 29339c.
 154 Carlsson, J., Mosbach, K. and Buelow, L.: Affinity precipitation and site-specific immobilization of proteins carrying polyhistidine tails. *Biotechnol. Bioeng.*, 51 (1996) 221-228; C.A., 125 (1996) 52850w.
 155 Olson, W.P.: Affinity chromatography. In: Olson, W.P. (Editor), *Sep. Technol.*, Interpharm Press, Buffalo Grove, 1995, pp. 303-351; C.A., 125 (1996) 177515x - a review with many refs.
 156 Staak, C., Salchow, F., Clausen, P.-H. and Luge, E.: Polystyrene as an affinity chromatography matrix for the purification of antibodies. *J. Immunol. Methods*, 194 (1996) 141-146; C.A., 125 (1996) 165224p.
 157 Zachariou, M., Traberso, I., Spiccia, L. and Hearn, M.T.W.: Potentiometric investigations into acid-base and metal ion binding properties of immobilized metal ion affinity chromatographic (IMAC) adsorbents. *J. Phys. Chem.*, 100 (1996) 12680-12690; C.A., 125 (1996) 42569x.
 158 Zusman, I., Zusman, R., Sandler, B., Korol, D., Gurevich, P., Smirnoff, P., Bass, D., Or, A., Shani, A. and Idelevich, E.: Gel fiberglass as a new matrix of affinity chromatography columns for isolation of colon cancer-associated antigens. *Int. J. Oncol.*, 9 (1996) 153-157; C.A., 125 (1996) 137045n.

For additional information see C.A.:

125 (1996) 80912m, 242071r, 242374s.

See also 102, 222, 286, 346, 381, 423, 471, 508, 532, 534, 562, 564, 576, 579, 605, 632, 643, 647, 663, 665, 669, 674, 678, 679, 680, 716, 735, 745, 747, 750, 780, 785.

4f. Trace analysis and preseparation techniques

- 159 Abushamaa, A., Naim, A. and Giese, R.W.: Satellite high-performance liquid chromatography for trace sample cleanup. *J. Chromatogr. A*, 742 (1996) 295-297.

160 Hajouj, Z., Thomas, J. and Siouffi, A.M.: Mixed mode SPE of pollutants in water on mechanically blended silica-based bonded phases. I. Influence of the preparation method. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1937-1945.

161 Majors, R.E.: The changing role of extraction in preparation of solid samples. *LC-GC Int.*, 9 (1996) 638-648.

For additional information see C.A.:

125 (1996) 131206u.

See also 435, 791, 906, 925, 996, 1205, 1237.

4g. Enantiomers, separation

- 162 Ahmed, A., Ibrahim, H., Pastore, F. and Lloyd, D.K.: Relationship between retention and effective selector concentration in affinity capillary electrophoresis and high-performance liquid chromatography. *Anal. Chem.*, 68 (1996) 3270-3273.
 163 Belokon, Yu.N., Kochetkov, K.A., Churkina, T.D., Ikonnikov, N.S., Orlova, S.A., Kuz'mina, N.A. and Bodrov, D.E.: (Asymmetric addition of methyl nitroacetate to crotonaldehyde catalyzed by chiral amines). *Izv. Akad. Nauk, Ser. Khim.*, (1993) 1591-1595; C.A., 125 (1996) 194975q.
 164 Burmester, A. and Jastorff, B.: Enantioseparation in the synthesis of myo-inositol phosphates by high-performance liquid chromatography using a β -cyclodextrin-bonded column. *J. Chromatogr. A*, 749 (1996) 25-32.
 165 Chandwani, O.D. and Dhaneshwar, S.R.: Chiral liquid chromatography: a review. *Indian Drugs*, 33 (1996) 187-192; C.A., 125 (1996) 241998m - a review with 97 refs.
 166 Ficarra, R., Calabro, M.C., Tommasini, S., Constantino, D., Carulli, M., Melardi, S., di Bella, M.R., Casuscelli, F., Romeo, R. and Ficarra, P.: Influence of organic modifier and temperature on the enantiomeric separation of γ -lactams by HPLC. *Chromatographia*, 43 (1996) 365-368.
 167 Harada, K., Yuan, Q., Nakayama, M. and Sugii, A.: Effects of organic modifiers on the chiral recognition by different types of silica-immobilized bovine serum albumin. *J. Chromatogr. A*, 740 (1996) 207-213.
 168 Kurganov, A.: Effect of solute association on the apparent adsorption isotherm. A model of the separation of non-racemic mixtures of enantiomers in achiral chromatographic systems. *Chromatographia*, 43 (1996) 17-24.
 169 Lämmerhofer, M. and Lindner, W.: Quinine and quinidine derivatives as chiral selectors. I. Brush type chiral stationary phases for high-performance liquid chromatography based on cinchonan carbamates and their application as chiral anion exchangers. *J. Chromatogr. A*, 741 (1996) 33-48.
 170 Lin, J.-Y.: Comparison of the chiral resolution on *cis-trans* isomeric chiral stationary phases derived from (S)-1-(naphthyl)ethylamine. *J. Chromatogr. A*, 740 (1996) 183-193.
 171 Ma, J., Chen, L. and He, B.: Synthesis of crosslinked poly(vinyl alcohol) with L-proline pendant as the chiral stationary phase for resolution of amino acid enantiomers. *J. Appl. Polym. Sci.*, 61 (1996) 2029-2034; C.A., 125 (1996) 248373n.
 172 Ma, Y. and Ito, Y.: Chiral separation of enantiomers by high-speed countercurrent chromatography. *PCT Int. Appl. WO 96 18,895* (Cl. GO1N30/42), 20 Jun. 1996, US Appl. 357,845, 16 Dec. 1994; 25 pp.; C.A., 125 (1996) 143307r.

- 173 Oda, Y. and Miwa, T.: Protein-bonded chiral packings and their application. *Methods Chromatogr.*, 1(Advances in Liquid Chromatography) (1996) 255-288; C.A., 125 (1996) 136685j - a review with 108 refs.
- 174 Ohnishi, A., Ichida, A. and Makino, S.: (Liquid chromatographic resolution of optical isomers. Development and characteristics of chiral stationary phases consist of polysaccharide derivatives and application to large scale separation). *Yuki Gosei Kagaku Kyokaishi*, 54 (1996) 344-353; C.A., 125 (1996) 85806q.
- 175 Pirkle, W.H. and Liu, Y.: On the relevance of face-to-edge $\pi\text{-}\pi$ interactions to chiral recognition. *J. Chromatogr. A*, 749 (1996) 19-24.
- 176 Priegnitz, J.W. and McCulloch, B.: Chiral separations by simulated moving-bed chromatography operating at low capacity factor k' values. *U.S. US 5,518,625* (Cl. 210-659; B01D15/08), 21 May 1996, Appl. 387,984, 13 Feb. 1995; 11 p.; C.A., 125 (1996) 62287g.
- 177 Selness, S.R.: Spectroscopic and chromatographic investigations of chiral recognition processes. Avail. Degree-Granting Institution, 1996, 97 pp.; C.A., 125 (1996) 220921r.
- 178 Takahata, H.: (Resolution of excess enantiomers with achiral phase chromatography). *Yuki Gosei Kagaku Kyokaishi*, 54 (1996) 708-711; C.A., 125 (1996) 166947p - a review with 14 refs.
- 179 Tanaka, N., Kimata, K., Kobayashi, M., Hosoya, K. and Araki, T.: (Chromatographic separation based on isotopic chirality). *Kuromatogurafu*, 17 (1996) 150-151; C.A., 125 (1996) 157336m.
- 180 Williams, K.L., Sander, L.C. and Wise, S.A.: Comparison of liquid and supercritical fluid chromatography using naphthylethyl-carbamoylated- β -cyclodextrin chiral stationary phases. *J. Chromatogr. A*, 746 (1996) 91-101.
- 186 Jinno, K. and Fujimoto, C.: Microcolumn liquid chromatography in hyphenated techniques (microcolumn LC/FTIR). *Methods Chromatogr.*, 1 (1996) 199-211; C.A., 125 (1996) 103816z - a review with 37 refs.
- 187 Lavine, B.K., Hendayana, S., He, Y. and Cooper, W.T.: Solid state NMR studies of ionic surfactants adsorbed on cyanopropyl bonded phases: implications for micellar liquid chromatography. *J. Colloid Interface Sci.*, 179 (1996) 341-349; C.A., 125 (1996) 68616k.
- 188 Mori, K. and Tomigahara, M.: (Chromatographic immunoassay). *Jpn. Kokai Tokkyo Koho JP 08,160,043* [96,160,043] (Cl. G01N33/543), 21 Jun. 1996, Appl. 94/305,040, 8 Dec. 1994; 8 pp.; C.A., 125 (1996) 162770c.
- 189 Reschigian, P., Melucci, D. and Torsi, G.: Experimental study on the retention of silica particles in gravitational field-flow fractionation. Effects of the mobile phase composition. *J. Chromatogr. A*, 740 (1996) 245-252.
- 190 Takeuchi, T. and Ishii, D.: Microcolumn liquid chromatography. *Methods Chromatogr.*, 1 (1996) 169-198; C.A., 125 (1996) 103815y - a review with 63 refs.
- 191 Van der Vlis, E., Mazereeuw, M., Tjaden, U.R., Irth, H. and van der Greef, J.: Development of a needle device for on-line electroextraction-liquid chromatography. *J. Chromatogr. A*, 741 (1996) 13-21.
- 192 West, C.M. and Smith, E.L.: Optimization of a semi-batch chromatographic reactor. In: *IChE Res. Event. Eur. Conf. Young Res. Chem. Eng.*, 2nd 1996, Vol. 1, Institution of Chemical Engineers, Rugby, 1996, pp. 148-150; C.A., 125 (1996) 56276m.
- 193 Zou, H., Zhang, Y., Lu, P. and Krull, I.S.: Perfusion immunoaffinity chromatography and its application in analysis and purification of biomolecules. *Biomed. Chromatogr.*, 10 (1996) 122-126; C.A., 125 (1996) 80887g.

For additional information see C.A.:

125 (1996) 32905q, 276351x.

See also 66, 78, 103, 125, 262, 341, 363, 385, 431, 434, 440, 486, 813, 818, 935, 979, 982, 988, 1001, 1002, 1004, 1005, 1012, 1014, 1015, 1019, 1020, 1030, 1057, 1072, 1076, 1081, 1104, 1126, 1203.

4h. Other special techniques

- 181 Bengio, S., Lettner, H. and Jungbauer, A.: (Aseptic chromatography processing: dream or reality?). *Ann. N.Y. Acad. Sci.*, 782(Recombinant DNA Biotechnology III) (1996) 432-440; C.A., 125 (1996) 109126w - a review with 8 refs.
- 182 Brooks, C.A. and Cramer, S.M.: Solute affinity in ion-exchange displacement chromatography. *Chem. Eng. Sci.*, 51 (1996) 3847-3860; C.A., 125 (1996) 109273s.
- 183 Chizhkov, V.P. and Sterkhov, N.V.: (Concepts and theory of pulsed chromatography with reversed mobile-phase flow). *Zh. Fiz. Khim.*, 70 (1996) 542-544; C.A., 125 (1996) 152045f.
- 184 Hameed, M.S. and Abid, M.F.: Unsteady-state model for continuous moving-bed chromatography. *Sep. Sci. Technol.*, 31 (1996) 2065-2087; C.A., 125 (1996) 171843f.
- 185 Jensen, K.D., Williams, S.K.R. and Giddings, J.C.: High-speed particle separation and steric inversion in thin flow field-flow fractionation channels. *J. Chromatogr. A*, 746 (1996) 137-145.

For additional information see C.A.:

125 (1996) 131001y, 146300w.

See also 36, 95, 101, 349, 399, 505, 526, 536, 543, 622, 838, 1204, 1209.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

- 194 Dai, J.-R., Hallock, Y.F., Cardellina, II, J.H., Gray, G.N. and Boyd, M.R.: Triangulynes A-H and triangulynic acid, new cytotoxic polyacetylenes from the marine sponge *Pellina triangulata*. *J. Natural Prod.*, 59 (1996) 860-865.
- 195 Ghazvini, V.N., Hany, R., Jackle, H.W. and Stenhauer, E.: (Determination of conjugated diolefins in gasolines by HPLC). *Erdgas, Kohle*, 112 (1996) 210-212; C.A., 125 (1996) 62690h.

For additional information see C.A.:

125 (1996) 119234u, 119235v.

5b. Cyclic hydrocarbons, fullerenes

- 196 Foster, M.D. and Synovec, R.E.: Reversed phase liquid chromatography of organic hydrocarbons with water as the mobile phase. *Anal. Chem.*, 68 (1996) 2838-2844.

- 197 Garcia, M.A. and Marina, M.L.: Influence of alcohol organic modifiers upon the association constants and retention mechanism for aromatic compounds in micellar liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1757-1776.
- 198 Manoli, E. and Samara, C.: Polycyclic aromatic hydrocarbons in waste waters and sewage sludge: extraction and clean-up for HPLC analysis with fluorescence detection. *Chromatographia*, 43 (1996) 135-142.
- 199 Mata, N.L., Mata, J.R. and Tsin, A.T.C.: Comparison of retinyl ester hydrolase activities in bovine liver and retinal pigment epithelium. *J. Lipid Res.*, 37 (1996) 1947-1952.
- 200 Moret, S., Grob, K. and Conte, L.S.: On-line high-performance liquid chromatography-solvent evaporation-high-performance liquid chromatography-capillary gas chromatography-flame ionisation detection for the analysis of mineral oil polycyclic hydrocarbons in fatty foods. *J. Chromatogr. A*, 750 (1996) 361-368.
- 201 Pothuluri, J.V., Evans, F.E., Heinze, T.M. and Cerniglia, C.E.: Formation of sulfate and glucoside conjugates of benzo[e]pyrene by *Cunninghamella elegans*. *Appl. Microbiol. Biotechnol.*, 45 (1996) 677-683; C.A., 125 (1996) 163007w.
- 202 Rajagopalan, P., Wudl, F., Schinazi, R.F. and Boudinot, F.D.: Pharmacokinetics of a water-soluble fullerene in rats. *Antimicrob. Agents Chemother.*, 40 (1996) 2262-2265; C.A., 125 (1996) 264906u.
- 203 Van Stijn, F., Kerkhoff, M.A.T. and Vandeginste, B.G.M.: Determination of polycyclic aromatic hydrocarbons in edible oils and fats by on-line donor-acceptor complex chromatography and high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 750 (1996) 263-273.

For additional information see C.A.:
125 (1996) 131000x, 146779g.

See also 75, 143, 160, 1202, 1204.

5c. Halogen derivatives

- 204 Chen, B.H., Wang, C.Y. and Chiu, C.P.: Evaluation of analysis of polycyclic aromatic hydrocarbons in meat products by liquid chromatography. *J. Agric. Food Chem.*, 44 (1996) 2244-2251; C.A., 125 (1996) 84940s.

See also 127.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

- 205 Chaffin, J.M., Lin, M.-S., Liu, M., Davison, R.R., Glover, C.J. and Bullin, J.A.: The use of HPLC to determine the saturate content of heavy petroleum products. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1669-1682.

6. ALCOHOLS

- 206 Hilmi, A., Belgsir, E.M., Leger, J.-M. and Laym, C.: Electrocatalytic oxidation of aliphatic diols on platinum. Part 4. A kinetic study by means of chromatographic analysis during prolonged electrolyses. *J. Electroanal. Chem.*, 410 (1996) 101-107; C.A., 125 (1996) 179744v.
- 207 Malandain, H. and Cano, Y.: (Blood alcohol and isopropanol). *Ann. Biol. Clin.*, 54 (1996) 97; C.A., 125 (1996) 135260e.

See also 194, 241, 330, 358.

7. PHENOLS

- 208 Amarowicz, R. and Shahidi, F.: A rapid chromatographic method for separation of individual catechins from green tea. *Food Res. Int.*, 29 (1996) 71-76; C.A., 125 (1996) 165955c.
- 209 Boyce, M.C. and Bennett, I.J.: Complementary role of micellar electrokinetic capillary chromatography and high performance liquid chromatography in the separation of plant phenolics. *Anal. Lett.*, 29 (1996) 1805-1815.
- 210 Chilla, C., Guillén, D.A., Barroso, C.G. and Pérez-Bustamante, J.A.: Automated on-line solid-phase extraction-high-performance liquid chromatography-diode array detection of phenolic compounds in sherry wine. *J. Chromatogr. A*, 750 (1996) 209-214.
- 211 Keinanen, M. and Julkunen-Tiitto, R.: Effect of sample preparation method on birch (*Betula pendula* Roth) leaf phenolics. *J. Agric. Food Chem.*, 44 (1996) 2231-2234; C.A., 125 (1996) 162434w.
- 212 Maiani, G., Azzini, E., Salucci, M., Ghiselli, A., Serafini, M. and Ferro-Luzzi, A.: A new sensitive and specific HPLC method for measuring polyphenol plasma and urinary levels after green tea ingestion. *Spec. Publ.-R. Soc. Chem.*, 181(Natural Antioxidants and Food Quality in Atherosclerosis and Cancer Prevention) (1996) 264-272; C.A., 125 (1996) 273915y.
- 213 Okamura, N., Asai, M., Hine, N. and Yagi, A.: High-performance liquid chromatographic determination of phenolic compounds in Aloe species. *J. Chromatogr. A*, 746 (1996) 225-231.
- 214 Shahrzad, S. and Bitsch, I.: Determination of some pharmacologically active phenolic acids in juices by high-performance liquid chromatography. *J. Chromatogr. A*, 741 (1996) 223-231.

See also 127, 222, 231.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 215 Gobbato, S., Griffini, A., Lolla, E. and Peterlongo, F.: HPLC quantitative analysis of biflavones in *Ginkgo biloba* leaf extracts and their identification by thermospray liquid chromatography-mass spectrometry. *Fitoterapia*, 67 (1996) 152-158; C.A., 125 (1996) 96290t.

- 216 Herrera, J.C., Romero, A.J.R., Crescente, O.E., Acosta, M. and Pekerar, S.: Analysis of 5-hydroxy-7-methoxyflavones by normal-phase high-performance liquid chromatography. *J. Chromatogr. A.*, 740 (1996) 201-206.
- 217 Hollman, P.C.H., van Trijp, J.M.P. and Buysman, M.N.C.P.: Fluorescence detection of flavonols in HPLC by postcolumn chelation with aluminium. *Anal. Chem.*, 68 (1996) 3511-3515.
- 218 Ishii, K., Furuta, T. and Kasuya, Y.: Determination of naringin and naringenin in human plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 683 (1996) 225-229.
- 219 Nguyenle, T., Wang, E. and Cheung, A.P.: An investigation of the extraction and concentration of isoflavones in soy-based products. *J. Pharm. Biomed. Anal.*, 14 (1995) 221-232.
- 220 Nikitina, V.S., Orazov, O.E. and Shendel, G.V.: (Use of Spheron 40 for separation of flavonoids from willow leaves). *Khim. Prir. Soedin.*, (1995) 755-756; *C.A.*, 125 (1996) 96271n.
- 221 Summer, L.W., Paiva, N.L., Dixon, R.A. and Geno, P.W.: High-performance liquid chromatography/continuous-flow liquid secondary ion mass spectrometry of flavonoid glycosides in leguminous plant extracts. *J. Mass Spectrom.*, 31 (1996) 472-485; *C.A.*, 125 (1996) 52609z.
- 222 Teslov, L.S., Merkulova, I.V. and Kuznetsov, P.V.: (Affinity-type polymeric adsorbents in studies on physiologically active substances. VII. Chromatographic separation of total polyphenolic compounds of *Campanula rapunculoides*). *Khim. Prir. Soedin.*, (1995) 811-813; *C.A.*, 125 (1996) 67928b.

For additional information see *C.A.*:
125 (1996) 96352q.

8b. Aflatoxins and other mycotoxins

- 223 Biancardi, A. and Riberzani, A.: Determination of ochratoxin A in cereals and feed by SAX-SPE clean up and LC fluorimetric detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2395-2407.
- 224 El-Sharkawy, S.H., Badria, F.A., Selim, M.I. and Halim, A.F.: Detection and quantitation of mycotoxins in Egyptian food commodities. *J. Environ. Sci. (Mansoura)*, 11 (1996) 165-178; *C.A.*, 125 (1996) 140826y.
- 225 Holcomb, M., Thompson, H.C., Jr., Cooper, W.M. and Hopper, M.L.: SFE extraction of aflatoxins (B₁, B₂, G₁, and G₂) from corn and analysis by HPLC. *J. Supercrit. Fluids*, 9 (1996) 118-121; *C.A.*, 125 (1996) 140827z.
- 226 Kraska, R., Schuhmacher, R., Grasserbauer, M. and Scott, P.M.: Determination of the *Fusarium* mycotoxin beauvericin at µg/kg levels in corn by high-performance liquid chromatography with diode-array detection. *J. Chromatogr. A*, 746 (1996) 233-238.
- 227 Lucacs, Z., Schaper, S., Herderich, M., Schreier, P. and Humpf, H.-U.: Identification and determination of fumonisin FB₁ and FB₂ in corn and corn products by high-performance liquid chromatography-electrospray-ionization tandem mass spectrometry (HPLC-ESI-MS-MS). *Chromatographia*, 43 (1996) 124-128.
- 228 Trucksess, M.W., Ready, D.W.E., Pender, M.K., Ligmond, C.A., Wood, G.E. and Page, S.W.: Determination and survey of deoxynivalenol in white flour, whole wheat flour, and bran. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 883-888.

- 229 Wilkes, J.G., Churchwell, M.L., Billedeau, S.M., Vollmer, D.L., Volmer, D.A., Thompson, H.C., Jr. and Lay, J.O., Jr.: Determination of underivatized fumonisin B₁ and related compounds by HPLC. *Adv. Exp. Med. Biol.*, 392 (Fumonisins in Food) (1996) 93-103; *C.A.*, 125 (1996) 78699d.

See also 1156, 1182.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 230 Farina, A., Doldo, A., Cotichini, V., Rajevic, M., Quaglia, M.G., Mulinacci, N. and Vincieri, F.F.: HPTLC and reflectance mode densitometry of anthocyanins in *Malva sylvestris* L.: a comparison with gradient-elution reversed-phase HPLC. *J. Pharm. Biomed. Anal.*, 14 (1995) 203-211.
- 231 Goto, T., Yoshida, Y., Kiso, M. and Nagashima, H.: Simultaneous analysis of individual catechins and caffeine in green tea. *J. Chromatogr. A*, 749 (1996) 295-299.
- 232 Killard, A.J., O'Kennedy, R. and Bogan, D.P.: Analysis of the glucuronidation of 7-hydroxycoumarin by HPLC. *J. Pharm. Biomed. Anal.*, 14 (1996) 1585-1590.
- 233 McMurrough, I., Madigan, D. and Smyth, M.R.: Semipreparative chromatographic procedure for the isolation of dimeric and trimeric proanthocyanidins from barley. *J. Agric. Food Chem.*, 44 (1996) 1731-1735; *C.A.*, 125 (1996) 84921m.
- 234 Park, S.W., Seo, B.S., Kim, E.H., Kim, D.H. and Paeng, K.-J.: Purification and determination procedure of coumarin derivatives. *J. Forensic Sci.*, 41 (1996) 685-688; *C.A.*, 125 (1996) 160650q.
- 235 Sperker, B., Schick, M. and Kroemer, H.K.: High-performance liquid chromatographic quantification of 4-methylumbelliferyl-α-D-glucuronide as a probe for human β-glucuronidase activity in tissue homogenates. *J. Chromatogr. B*, 685 (1996) 181-184.
- 236 Unterhalt, B. and Jöstingmeier, R.: Neue substituierte Isochromane. *Pharmazie*, 51 (1996) 641-644.

See also 208, 1170, 1207.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 237 Al-Rawithi, S., El-Yazigi, A. and Nicholls, P.J.: Rapid liquid chromatographic analysis of acrolein in plasma following precolumn derivatization. *Pharm. Sci.*, 1 (1995) 201-203; *C.A.*, 125 (1996) 184741w.
- 238 Borner, K., Borner, E., Hartwig, H. and Lode, H.: Fluorimetric detection of non-fluorescent quinolones after post-column photoreaction. *Methodol. Surv. Bioanal. Drugs*, 24 (Biofluid Assay for Peptide-Related and Other Drugs) (1996) 147-150; *C.A.*, 125 (1996) 264837x.
- 239 Coco, F.L., Valentini, C., Novelli, V. and Ceccan, L.: High-performance liquid chromatographic determination of 2-furaldehyde and 5-hydroxymethyl-2-furaldehyde in honey. *J. Chromatogr. A*, 749 (1996) 95-102.
- 240 Cserháti, T. and Forgács, E.: Simultaneous determination of 3,4-dimethoxybenzaldehyde and 3,4-dimethoxyphenylacetone in industrial waste waters by high-performance liquid chromatography-diode array detection. *J. Chromatogr. A*, 750 (1996) 257-264.

- 241 Fields, S.C., Mireles-Lo, L. and Gerwick, B.C.: Hydroxycornexistin: a new phytotoxin from *Paecilomyces variotti*. *J. Natural Prod.*, 59 (1996) 698-700.
- 242 Goj, T.: (Methods of determination of carbonyl compounds occurring in spirit and alcoholic beverages in view of the latest studies. Part 1. Methods recommended by the Polish norms and changes of these norms with ISO international standards). *Przem. Ferment. Owocowo-Warzywny*, 40 (1996) 14-16; C.A., 125 (1996) 165860t - a review with 14 refs.
- 243 Levin, J.-O., Lindahl, R., Heeremans, C.E.M. and van Oosten, K.: Certification of reference materials related to the monitoring of aldehydes in air by derivatization with 2,4-dinitrophenylhydrazine. *Analyst (Cambridge)*, 121 (1996) 1273-1278.
- 244 Lindahl, R., Levin, J.O. and Märtenson, M.: Validation of diffusive sampler for the determination of acetaldehyde in air. *Analyst (Cambridge)*, 121 (1996) 1177-1181.
- 245 Okada, K., Suzuki, K., Kamiya, Y., Zhu, X.F., Fujisaki, S., Nishimura, Y., Nishino, T., Nakagawa, T., Kawamukai, M. and Matsuda, H.: Polyprenyl diphosphate synthase essentially defines the length of the side chain of ubiquinone. *Biochim. Biophys. Acta*, 1302 (1996) 217-223.
- 246 Possanzini, M. and di Palo, V.: Short-term measurements of acrolein in ambient air. *Chromatographia*, 43 (1996) 433-435.
- 247 Potter, W. and Karst, U.: Identification of chemical interferences in aldehyde and ketone determination using dual-wavelength detection. *Anal. Chem.*, 68 (1996) 3354-3358.
- 248 Starratt, A.N. and Lazarovits, G.: High-performance liquid chromatography with photodiode array detection for analysis of the fungal metabolite sclerin. *J. Chromatogr. A*, 741 (1996) 131-134.
- For additional information see C.A.:
125 (1996) 34543u, 96287x, 123856r.
- See also 316, 330, 1167.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 249 Bai, X. and Esko, J.D.: An animal cell mutant defective in heparan sulfate hexuronic acid 2-O-sulfation. *J. Biol. Chem.*, 271 (1996) 17711-17717.
- 250 Bao, Y., Silva, T.M.J., Guerrant, R.L., Lima, A.A.M. and Fox, J.W.: Direct analysis of mannitol, lactulose and glucose in urine samples by high-performance anion-exchange chromatography with pulse amperometric detection. Clinical evaluation of intestinal permeability in human immunodeficiency virus infection. *J. Chromatogr. B*, 685 (1996) 105-112.
- 251 Caron, I., Salvador, A., Elfakir, C., Herbreteau, B. and Dreux, M.: Analysis of partially methylated cyclodextrins by subcritical fluid and liquid chromatography. *J. Chromatogr. A*, 746 (1996) 103-108.
- 252 Choi, O.-K., Kim, C.-G., Kim, Y.-D., Kim, H. and Jo, J.-S.: Determination of 2-keto-L-gulonic, 2-keto-D-gluconic and 2,5-diketo-D-gluconic acids by capillary zone electrophoresis. *J. Chromatogr. A*, 745 (1996) 249-254.
- 253 Del Carmen Molina, M. and Vicente, C.: High-performance liquid chromatographic characterization of two lichen lectins with arginase activity differing in their glycosyl moiety. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2102-2115.
- 254 Field, M., Papac, D. and Jones, A.: The use of high-performance anion-exchange chromatography and matrix-assisted laser desorption/ionization tissue-of-flight mass spectrometry to monitor and identify oligosaccharide degradation. *Anal. Biochem.*, 239 (1996) 92-98.
- 255 Gopal, P.K. and Richardson, R.K.: A rapid and sensitive method for estimation of galactose in Parmesan cheese. *Int. Dairy J.*, 6 (1996) 399-406; C.A., 125 (1996) 84923p.
- 256 Gough, H., Luke, G.A., Beeley, J.A. and Geddes, D.A.M.: Human salivary glucose analysis by high-performance ion-exchange chromatography and pulsed amperometric detection. *Arch. Oral Biol.*, 41 (1996) 141-145; C.A., 125 (1996) 29353c.
- 257 Greve, K.F., Hughes, D.E. and Karger, B.L.: Capillary electrophoretic examination of underivatized oligosaccharide mixtures released from immunoglobulin G antibodies and CTLA4Ig fusion protein. *J. Chromatogr. A*, 749 (1996) 237-245.
- 258 Heinrich, J. and Mischnick, P.: Rapid method for the determination of the substitution pattern of O-methylated 1,4-glucans by high-pH anion-exchange chromatography with pulsed amperometric detection. *J. Chromatogr. A*, 749 (1996) 41-45.
- 259 Hyoeky, G., Heikkilae, H., Kuusma, J., Monten, K.-E. and Paananen, H.: Method for fractionation of sucrose-containing solutions. *PCT Int. Appl. WO 96 10,650* (Cl. C13D3/14), 11 Apr. 1996, FI Appl. 94/4,577, 30 Sep. 1994; 45 p.; C.A., 125 (1996) 36213k.
- 260 Kim, S.-T., Huh, W.-K., Kim, J.-Y., Hwang, S.-W. and Kang, S.-O.: D-Arabinose dehydrogenase and biosynthesis of erythroascorbic acid in *Candida albicans*. *Biochim. Biophys. Acta*, 1297 (1996) 1-8.
- 261 Kimura, T., Shirota, O., Suzuki, A. and Ohtsu, Y.: (High-sensitivity carbohydrate analysis by semi-microcolumn LC with pulsed amperometric detector). *Kuromatografi*, 17 (1996) 156-157; C.A., 125 (1996) 87033j.
- 262 Koimur, M., Lu, B. and Westerlund, D.: High performance liquid chromatography of disaccharides on a porous graphitic carbon column applying post-column derivatization with benzamidine. *Chromatographia*, 43 (1996) 254-260.
- 263 Kunz, C., Rudloff, S., Hintelmann, A., Pohlentz, G. and Egge, H.: High-pH anion-exchange chromatography with pulsed amperometric detection and molar response factors of human milk oligosaccharides. *J. Chromatogr. B*, 685 (1996) 211-221.
- 264 Lines, A.C.: High-performance liquid chromatographic mapping of the oligosaccharides released from the humanised immunoglobulin, CAMPATH™ 1H. *J. Pharm. Biomed. Anal.*, 14 (1996) 601-608.
- 265 MacCarana, M., Sakura, Y., Tawada, A., Yoshida, K. and Lindahl, U.: Domain structure of heparan sulfates from bovine organs. *J. Biol. Chem.*, 271 (1996) 17804-17810.
- 266 Nestorowicz, J., Pierzynowska-Kornik, G. and Zadernowski, R.: (Application of the HPLC method to the determination of glycopyranosides during ripening of fava bean seeds). *Roczn. Panstw. Zakl. Hig.*, 47 (1996) 231-237; C.A., 125 (1996) 109274t.

- 267 Nguyen, K.P.P., Nguyen, N.S., Pham, D.H., Potier, P. and Qui, K.H.: Microquantitative analysis of neutral sugars in the aqueous extract of *Gossypium herbaceum* seeds by gas-liquid chromatography. *Tap Chi, Hoa Hoc.*, 33 (1995) 80-82; C.A., 125 (1996) 136813z.
- 268 Niessen, W.M.A., Schols, H.A., van der Hoeven, R.A.M., Voragen, A.G.J. and van der Greef, J.: Characterization of oligosaccharides derived from plant cell wall polysaccharides by online high-performance anion-exchange chromatography thermospray mass spectrometry. *Mod. Methods Plant Anal.*, 17(Plant Cell Wall Analysis) (1996) 147-163; C.A., 125 (1996) 52553b - a review with 24 refs.
- 269 Sidelmann, U.G., Hansen, S.H., Gavaghan, C., Carless, H.A.J., Lindon, J.C., Farrant, R.D., Wilson, I.D. and Nicholson, J.K.: Measurement of internal acyl migration reaction kinetics using directly coupled HPLC-NMR: application for the positional isomers of synthetic (2-fluorobenzoyl)-D-glucopyranuronic acid. *Anal. Chem.*, 68 (1996) 2564-2572.
- 270 Townsend, R.R., Lipniunas, P.H., Bigge, C., Ventom, A. and Parekh, R.: Multimode high-performance liquid chromatography of fluorescently labeled oligosaccharides from glycoproteins. *Anal. Biochem.*, 239 (1996) 200-207.
- 271 Uozumi, N., Teshima, T., Yamamoto, T., Nishikawa, A., Gao, Y.-E., Miyoshi, E., Gao, C.-X., Noda, K., Islam, K.N., Ihara, Y., Fujii, S., Shiba, T. and Taniguchi, N.: A fluorescent assay method for GDP-L-Fuc:N-acetyl- β -D-glucosaminide α 1-6fucosyltransferase activity, involving high performance liquid chromatography. *J. Biochem. (Tokyo)*, 120 (1996) 385-392.
- 272 Volpi, N.: Inhibition of human leukocyte elastase activity by heparins: influence of charge density. *Biochim. Biophys. Acta*, 1290 (1996) 299-307.
- For additional information see C.A.:
125 (1996) 103901y, 273930z.
- See also 58, 275, 286, 290, 294, 302, 715, 759, 778, 1174.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 273 Adikwu, M.U.: Gel permeation chromatography of a polysaccharide gum from *Mucuna flagellipes*. *Pharmakeutike*, 8 (1995) 151-153; C.A., 125 (1996) 29979m.
- 274 Bahary, W.S. and Hogan, M.P.: Assessment of branching in polymers by size-exclusion chromatography with light scattering, viscosity, and refractive index detection. *Int. J. Polym. Anal. Charact.*, 2 (1996) 121-133; C.A., 125 (1996) 89519w.
- 275 Guile, G.R., Rudd, P.M., Wing, D.R., Prime, S.B. and Dwek, R.A.: A rapid high-resolution high-performance liquid chromatographic method for separating glycan mixtures and analyzing oligosaccharide profiles. *Anal. Biochem.*, 240 (1996) 210-226.
- 276 Henshall, A.: Analysis of starch and other complex carbohydrates by liquid chromatography. *Cereal Foods World*, 41 (1996) 419-424; C.A., 125 (1996) 140714k - a review with 29 refs.
- 277 Hotchkiss, A.T., Jr., El-Bahtmy, K and Fishman, M.L.: Analysis of pectin structure by HPAEC-PAD. *Mod. Methods Plant Anal.*, 17(Plant Cell Wall Analysis) (1996) 129-146; C.A., 125 (1996) 52552a - a review with 85 refs.
- 278 Huang, Y., Washio, Y., Hara, M., Toyoda, H., Koshiishi, I., Toida, T. and Imanari, T.: Simultaneous determination of dermatan sulfate and oversulfated dermatan sulfate in plasma by high-performance liquid chromatography with postcolumn fluorescence derivatization. *Anal. Biochem.*, 240 (1996) 227-234.
- 279 Huber, A. and Eteshola, E.: Characterization of an aqueous-disolved anionic polysaccharide using rheology and size-exclusion chromatography with a differential refractive-index and low-angle laser-light scattering detectors. *Int. J. Polym. Anal. Charact.*, 2 (1996) 203-212; C.A., 125 (1996) 29344a.
- 280 Izawa, M., Takashio, M. and Tamaki, T.: Determination of high molecular weight β -glucan in wort and beer using a post-column calcofluor flow-injection-analysis (FIA) method. *J. Inst. Brew.*, 102 (1996) 183-189; C.A., 125 (1996) 113020z.
- 281 Kaneo, Y., Uemura, T., Tanaka, T. and Kanoh, S.: (Polysaccharides as drug carriers. High-performance size-exclusion chromatographic determination of FITC-labeled dextran and pullulan). *Drug Delivery Syst.*, 11 (1996) 161-168; C.A., 125 (1996) 123554r.
- 282 Malsch, R., Harenberg, J., Piazzolo, L., Huhle, G. and Heene, D.L.: Chromatographic and electrophoretic applications for the analysis of heparin and dermatan sulfate. *J. Chromatogr. B*, 685 (1996) 223-231.
- 283 Mascellani, G., Liverani, L. and Bianchini, P.: Analysis of heparin origin by HPLC quantitation of disaccharide components. *Farmac.*, 51 (1996) 247-254; C.A., 125 (1996) 25522j.
- 284 McCeely, T.B. and Doyle, P.S.: Isolation of lipophosphoglycans from *Leishmania donovani* amastigotes. *Arch. Biochem. Biophys.*, 334 (1996) 1-8.
- 285 Mehrav, R.: Effects of storage and homogenization methods on the hepatic recovery of dextrans determined by size-exclusion chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 801-805.
- 286 Parthasarathy, N., Goldberg, I.J., Sivaram, P. and Wagner, W.D.: Isolation of heparin-derived oligosaccharides containing 2-O-sulfated hexuronic acids, by lipoprotein lipase affinity chromatography. *J. Biochem. Biophys. Methods*, 32 (1996) 27-32; C.A., 125 (1996) 80875b.
- 287 Rossi, A., Bonaventure, J., Delezoide, A.-L., Cetta, G. and Superti-Furga, A.: Undersulfation of proteoglycans synthesized by chondrocytes from a patient with achondrogenesis type 1B homozygous for an L483P substitution in the diastrophic dysplasia sulfate transporter. *J. Biol. Chem.*, 271 (1996) 18456-18464.
- 288 Striegel, A.M. and Timpa, J.D.: Gel permeation chromatography of polysaccharides using universal calibration. *Int. J. Polym. Anal. Charact.*, 2 (1996) 213-220; C.A., 125 (1996) 36192c.
- 289 Striegel, A.M. and Timpa, J.D.: Size exclusion chromatography of polysaccharides in dimethylacetamide-lithium chloride. *ACS Symp. Ser.*, 635 (1996) 366-378; C.A., 125 (1996) 61381w.
- 290 Takahashi, N.: (Current structural analysis of N-linked oligosaccharides and application to clinical medicine). *Seibutsu Butsuri Kagaku*, 40 (1996) 105-110; C.A., 125 (1996) 109192q - a review with 5 refs.
- 291 Walker, A. and Gallagher, J.T.: Structural domains of heparin sulphate for specific recognition of the C-terminal heparin-binding domain of human plasma fibronectin (HEPII). *Biochem. J.*, 317 (1996) 871-877.

- 292 Wang, R., Klegerman, M.E. and Groves, M.J.: Analysis of anti-neoplastic polysaccharides from *Mycobacterium bovis* BCG vaccine by high performance anion exchange chromatography with pulsed amperometric detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2691-2698.

See also 268, 270, 272, 489, 715, 759, 1155.

10c. Glycoproteins and their constituents

- 293 Apffel, A., Chakel, J.A., Hancock, W.S., Souders, C., M'Timkulu, T. and Pungor, E., Jr.: Application of multidimensional affinity high-performance liquid chromatography and electrospray ionization liquid chromatography-mass spectrometry to the characterization of glycosylation in single-chain plasminogen activator. Initial results. *J. Chromatogr. A*, 750 (1996) 35-42.
- 294 Ashton, D.S., Beddell, C.R., Cooper, D.J. and Lines, A.: The examination of the carbohydrates attached to a monoclonal antibody by the techniques of liquid chromatography and mass spectrometry. In: Geisow, M.J. and Epton, R. (Editors), *Perspect. Protein Eng. Complementary Technol., Collect. Pap., Int. Symp., 3rd 1994*, Mayflower Worldwide, Kingswinford, 1995, pp. 304-305; C.A., 125 (1996) 83920y.
- 295 Castillo, G.M., Cummings, J.A., Ngo, C., Yang, W. and Snow, A.D.: Novel purification and detailed characterization of perlecan isolated from the Engelbreth-Holm-Swarm tumor for use in an animal model of fibrillar Ab amyloid persistence in brain. *J. Biochem. (Tokyo)*, 120 (1996) 433-444.
- 296 Hancock, W.S., Apffel, A., Chakel, J., Souders, C., M'Timkulu, T., Pungor, E., Jr. and Guzzetta, A.W.: Reversed-phase peptide mapping of glycoproteins using liquid chromatography/electrospray ionization-mass spectrometry. *Methods Enzymol.*, 271(High Resolution Separation and Analysis of Biological Macromolecules) (1996) 403-427; C.A., 125 (1996) 242339j.
- 297 Heegaard, N.H.H., Heegaard, P.M.H., Roepstorff, P. and Robey, F.A.: Ligand-binding sites in human serum amyloid P component. *Eur. J. Biochem.*, 239 (1996) 850-856.
- 298 Hering, T.M., Kollar, J., Huynh, T.D. and Varelas, J.B.: Purification and characterization of decorin core protein expressed in *Escherichia coli* as maltose-binding protein fusion. *Anal. Biochem.*, 240 (1996) 98-108.
- 299 Hovnenberg, H.W., Davies, J.R. and Carlstedt, I.: Different mucins are produced by the surface epithelium and the submucosa in human trachea: identification of MUC5AC as a major mucin from the goblet cells. *Biochem. J.*, 318 (1996) 319-324.
- 300 Hvarregaard, J., Andersen, M.A., Berglund, L., Rasmussen, J.T. and Petersen, T.E.: Characterization of glycoprotein PAS-6/7 from membranes of bovine milk fat globules. *Eur. J. Biochem.*, 240 (1996) 628-636.
- 301 Iwase, H., Tanaka, A., Hiki, Y., Kokubo, T., Ishii-Karakasa, I., Kobayashi, Y. and Hotta, K.: Estimation of the number of O-linked oligosaccharides per heavy chain of human serum IgA1 by matrix-assisted laser desorption ionization time-of-flight mass spectrometry (MALDI-TOFMS) analysis of the hinge glycopeptide. *J. Biochem. (Tokyo)*, 120 (1996) 393-397.
- 302 Katsumata, T., Noguchi, S., Yonezawa, N., Tanokura, M. and Nakano, M.: Structural characterization of the N-linked carbohydrate chains of the zone pellucida glycoproteins from bovine ovarian and fertilized eggs. *Eur. J. Biochem.*, 240 (1996) 448-453.
- 303 Liu, C.: Trace determination of glycoproteins by mass spectrometry coupled to high performance liquid chromatography and capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9621902, 1996, 180 pp.; C.A., 125 (1996) 136818e.
- 304 Murata, T., Itoh, T., Hayakawa, Y. and Usui, T.: Convenient synthesis of β -(1 \rightarrow 3)-galactosyl disaccharide α -glycoside and its analogs as mimic units of mucin-type carbohydrate. *J. Biochem. (Tokyo)*, 120 (1996) 851-855.
- 305 Murawaki, Y., Ikuta, Y., Idobe, Y. and Kawasaki, H.: Characterization of tissue inhibitor of metalloproteinases-1 in plasma from patients with chronic liver disease. *Clin. Chim. Acta*, 254 (1996) 77-83.
- 306 Ozaki, Y., Omichi, K. and Hase, S.: Preparation of neoglycolipids with a definite sugar chain moiety from glycoproteins. *Anal. Biochem.*, 241 (1996) 151-155.
- 307 Quill, T.A. and Hedrick, J.L.: The fertilization layer mediated block to polyspermy in *Xenopus laevis*: isolation of the cortical granule lectin ligand. *Arch. Biochem. Biophys.*, 333 (1996) 326-332.
- 308 Streets, A.J., Brooks, S.A., Dwek, M.V. and Leathem, A.J.: Identification, purification and analysis of a 55 kDa lectin binding glycoprotein present in breast cancer tissue. *Clin. Chim. Acta*, 254 (1996) 47-61.
- 309 Townsend, R.R., Basa, L.J. and Spellman, M.W.: Identification and characterization of glycopeptides in tryptic maps by high-pH anion-exchange chromatography. *Methods Enzymol.*, 271(High Resolution Separation and Analysis of Biological Macromolecules) (1996) 135-147; C.A., 125 (1996) 269505x.
- 310 Weitzhandler, M., Pohl, C., Rohrer, J., Narayanan, L., Slingsby, R. and Avdalovic, N.: Eliminating amino acid and peptide interference in high-performance anion-exchange pulsed amperometric detection glycoprotein monosaccharide analysis. *Anal. Biochem.*, 241 (1996) 128-134.
- 311 Yamaguchi, H. and Uchida, M.: A chaperone-like function of intramolecular high-mannose chains in the oxidative refolding of bovine pancreatic RNase B. *J. Biochem. (Tokyo)*, 120 (1996) 474-477.

For additional information see C.A.:
125 (1996) 29586f.

See also 17, 253, 264.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 312 Adlof, R.O.: Analysis of fatty acid mono- and diacylglycerol positional isomers by silver ion high-performance liquid chromatography. *J. Chromatogr. A*, 741 (1996) 135-138.
- 313 Bertone, S., Giacomini, M., Ruggiero, C., Piccarolo, C. and Calegari, L.: Automated systems for identification of heterotrophic marine bacteria in the basis of their fatty acid composition. *Appl. Environ. Microbiol.*, 62 (1996) 2122-2132; C.A., 125 (1996) 52646j.

- 314 Blanco, D., Quintanilla, M.E., Mangas, J.J. and Gutierrez, M.D.: Determination of organic acids in apple juice by capillary liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2615-2621.
- 315 Chalgeri, A. and Tan, H.S.I.: Indirect photometric detection for determination of citrate in pharmaceutical matrices by ion chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 835-844.
- 316 Charriere, R., Leiser, W. and Dousse, R.: HPLC methods for the determination of some quinolones in fish and animal tissues. *Mitt. Geb. Lebensmittelunters. Hyg.*, 87 (1996) 223-229; C.A., 125 (1996) 166053u.
- 317 Duffey, P.S., Guthertz, L.S. and Evans, G.C.: Improved rapid identification of mycobacteria by combining solid-phase extraction with high-performance liquid chromatography analysis of BACTEC cultures. *J. Clin. Microbiol.*, 34 (1996) 1939-1943; C.A., 125 (1996) 189718d.
- 318 Esteve-Juan, E., Puchades, R. and Maqueira, A.: Determination of acetate in foods by ion chromatography-flow injection. *South. Braz. J. Chem.*, 3 (1995) 9-22; C.A., 125 (1996) 193762f.
- 319 Gosalbo, L., Fabrias, G. and Camps, F.: Efficient separation of fatty acyl precursors on *Spodoptera littoralis* sex pheromone by reversed-phase high-performance liquid chromatography. *Arch. Insect Biochem. Physiol.*, 33 (1996) 75-81; C.A., 125 (1996) 216010a.
- 320 Holden, A.J., Littlejohn, D. and Fell, G.S.: Determination of citrate in plasma protein solutions by UV-visible spectrophotometry and ion chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 713-719.
- 321 Inoue, Y., Katsumata, Y., Tani, K. and Suzuki, Y.: Evaluation of the influence of the base material loaded with hydrous zirconium oxide gel on the retention behavior of disubstituted benzoic acids. *Chromatographia*, 43 (1996) 79-81.
- 322 Katagiri, M. and Awatsuura, R.: (Investigation of edible insects in the Ina area and comparison of the contents of fatty acids in the imago stage of locusts by various extraction methods). *Iida Joshi Tanki Daigaku Kiyo*, 13 (1996) 82-89; C.A., 125 (1996) 166212v.
- 323 Koide, M., Takahashi, M., Tamabaki, S. and Tagaki, W.: Catalytic effect of dipotassium glycyrrhizinate on the hydrolysis of nonionic ester surfactants. *J. Am. Oil Chem. Soc.*, 73 (1996) 913.
- 324 Kramell, R., Schneider, G. and Miersch, O.: Resolution of racemic jasmmonic acid and abscisic acid by chiral high performance liquid chromatography. *Phytochem. Anal.*, 7 (1996) 209-212; C.A., 125 (1996) 136794u.
- 325 Krzyszowska, A.J., Blaylock, M.J., Vance, G.F. and David, M.B.: Ion-chromatographic analysis of low molecular weight organic acids in Spodosol forest floor solutions. *Soil Sci. Soc. Am. J.*, 60 (1996) 1565-1571; C.A., 125 (1996) 246483z.
- 326 Lee, C., Martin, K.O. and Javitt, N.B.: Bile acid synthesis: 7 α -hydroxylation of intermediates in the sterol 27-hydroxylase metabolic pathway. *J. Lipid Res.*, 37 (1996) 1356-1362.
- 327 Lin, W.-J., Chen, R.R.-L. and Suen, S.-L.: Determination of benzonate and its metabolite in plasma by high performance liquid chromatography. *Chin. Pharm. J. (Taipei)*, 48 (1996) 53-61; C.A., 125 (1996) 104155v.
- 328 MacPherson, J.C., Pavlovich, J.G. and Jacobs, R.S.: Biosynthesis of arachidonic acid metabolites in *Limulus polyphemus* amebocytes: analysis by liquid chromatography-electrospray ionization mass spectrometry. *Biochim. Biophys. Acta*, 1303 (1996) 127-136.
- 329 Mayer, A.F., Anspach, F.B. and Deckwer, W.-D.: Purification of clavulanic acid by ion-pairing systems. *Bioseparation*, 6 (1996) 25-39; C.A., 125 (1996) 242049q.
- 330 Mefford, I.N., Kincl, L., Dykstra, K.H., Simpson, J.T., Markey, S.P., Dietz, S. and Wightman, R.M.: Facile oxidative decarboxylation of 3,4-dihydroxyphenylacetic acid catalyzed by copper and manganese ions. *Biochim. Biophys. Acta*, 1290 (1996) 224-230.
- 331 Moghaddam, M.F., Motoba, K., Borhan, B., Pinot, F. and Hammock, B.D.: Novel metabolic pathways for linoleic and arachidonic acid metabolism. *Biochim. Biophys. Acta*, 1290 (1996) 327-339.
- 332 Nikolova-Damyanova, B., Christie, W.W. and Herslöf, B.: Mechanistic aspects of fatty acid retention in silver ion chromatography. *J. Chromatogr. A*, 749 (1996) 47-54.
- 333 Pinstone, S.N., Clee, S.M., Gangé, S.E., Miao, L., Zhang, H., Stein, E.A. and Hayden, M.R.: A frequently occurring mutation of the lipoprotein lipase gene (AsN291ser) results in altered post-prandial chylomicron triglyceride and retinyl palmitate response in normolipidemic carriers. *J. Lipid Res.*, 37 (1996) 1675-1684.
- 334 Retterstol, K., Woldseth, B. and Christoffersen, B.O.: The metabolism of 22:5(6-) and of docosahexaenoic acid [22:6(3-)] compared in rat hepatocytes. *Biochim. Biophys. Acta*, 1303 (1996) 180-186.
- 335 Ritter, D., Carlson, Ia Donna C., Logan, B.K., Ramos, L.S., Kilburn, J.O. and Coyle, M.B.: Differentiation of *Mycobacterium genavense* and *Mycobacterium simiae* by automated mycolic acid analysis with high-performance liquid chromatography. *J. Clin. Microbiol.*, 34 (1996) 2004-2006; C.A., 125 (1996) 189719e.
- 336 Sagi, K., Kagechika, H., Fukusawa, H., Hashimoto, Y. and Shudo, K.: Deiodotrifluoromethylation of ethyl 3,3',5'-triodothyroacetate. Divergent derivatization based on the combinatorial concept. *Chem. Pharm. Bull.*, 44 (1996) 1273-1275.
- 337 Sala, A., Bolla, M., Zarini, S., Müller-Peddinghaus, R. and Folco, G.: Release of leukotriene A₄ versus leukotriene B₄ from human polymorphonuclear leukocytes. *J. Biol. Chem.*, 271 (1996) 17944-17948.
- 338 Santa, T., Kimoto, K., Fukushima, T., Homma, H. and Imai K.: 4-(N-Hydrazinoformylmethyl-N-methyl)-amino-7-N,N-dimethylaminosulfonyl-2,1,3-benzoxadiazole (DBD-CO-Hz) as a pre-column fluorescence derivatization reagent for carboxylic acids in high-performance liquid chromatography. *Biomed. Chromatogr.*, 10 (1996) 183-185; C.A., 125 (1996) 136801u.
- 339 Schrader, K., Kiehne, A., Engelhardt, U.H. and Maier, H.G.: Determination of chlorogenic acids with lactones in roasted coffee. *J. Sci. Food Agric.*, 71 (1996) 392-398; C.A., 125 (1996) 113026f.
- 340 Ste-Marie, L., Boismenu, D., Vachon, L. and Montgomery, J.: Evaluation of sodium 4-hydroxybenzoate as an hydroxyl radical trap using gas chromatography - mass spectrometry and high-performance liquid chromatography with electrochemical detection. *Anal. Biochem.*, 241 (1996) 67-74.

- 341 Trute, A. and Nahrstedt, A.: Separation of rosmarinic acid enantiomers by three different chromatographic methods (HPLC, CE, GC) and the determination of rosmarinic acid in *Hedera helix* L. *Phytochem. Anal.*, 7 (1996) 204-208; C.A., 125 (1996) 109268u.
- 342 Veit, M., Reverey, H. and Schmidt, M.F.G.: Cytoplasmic tail length influences fatty acid selection for acylation of viral glycoproteins. *Biochem. J.*, 318 (1996) 163-172.
- 343 Venema, D.P., Hollman, P.C.H., Janssen, K.P.L.T.M. and Katan, M.B.: Determination of acetylsalicylic and salicylic acids in foods, using HPLC with fluorescence detection. *J. Agric. Food Chem.*, 44 (1996) 1762-1767; C.A., 125 (1996) 32164x.
- For additional information see C.A.:
125 (1996) 103849n, 177546h.
- See also 81, 194, 214, 222, 248, 346, 370, 373, 421, 691, 966, 1001, 1150, 1166, 1233.
- 11b. *Prostaglandins*
- 344 Ikuta, N., Sugiyama, S., Takagi, K., Hayakawa, T. and Ozawa, T.: Laser high performance liquid chromatography determination of prostaglandins in pleural effusions. *Biochem. Mol. Biol. Int.*, 36 (1995) 521-527; C.A., 125 (1996) 26433t.
- 11c. *Lipids and their constituents*
- 345 Abidi, S.L. and Mounts, T.L.: Reversed-phase high-performance liquid chromatography of molecular species of phospholipid derivatives. *J. Chromatogr. A*, 741 (1996) 213-222.
- 346 Amoroso, A., Washington, B.A., Gutkind, G. and Doyle, R.J.: An affinity chromatographic method for the preparation of bacterial lipoteichoic acids. *J. Microbiol. Methods*, 25 (1996) 19-22; C.A., 125 (1996) 52640c.
- 347 Andersson, A.-S., Rilfors, L., Bergqvist, M., Persson, S. and Lindblom, G.: New aspects on membrane lipid regulation in *Acholeplasma laidlawii* A phase equilibria of monoacyldiglycosyldiacylglycerol. *Biochemistry*, 35 (1996) 11119-11130.
- 348 Arcos, J.A. and Otero, C.: Enzyme, medium, and reaction engineering to desing a low-cost, selective production method for mono- and dioleoylglycerols. *J. Am. Oil Chem. Soc.*, 73 (1996) 673-682.
- 349 Baudimant, G., Maurice, M., Landrein, A., Durand, G. and Durand, P.: Purification of phosphatidylcholine with high content of DHA from squid *Illex argentinus* by countercurrent chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1793-1804.
- 350 Buchanan, S.K. and Walker, J.E.: Large-scale chromatographic purification of F₁F₀-ATPase and complex I from bovine heart mitochondria. *Biochem. J.*, 318 (1996) 343-349.
- 351 Byrdwell, W.C. and Neff, W.E.: Analysis of genetically modified canola varieties by atmospheric pressure chemical ionization mass spectrometric and flame ionization detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2203-2225.
- 352 Christie, W.W.: Separation of phospholipid classes by high-performance liquid chromatography. *Oily Press Lipid Libr.*, 7(Advances in Lipid Methodology-Three) (1996) 77-107; C.A., 125 (1996) 109178q - a review with 129 refs.
- 353 Clay, M.A. and Barter, P.J.: Formation of new-HDL particles from lipid-free apolipoprotein A-I. *J. Lipid Res.*, 37 (1996) 1722-1732.
- 354 Fischer, W.: Molecular analysis of lipid macroamphiphiles by hydrophobic interaction chromatography. *J. Microbiol. Methods*, 25 (1996) 129-144; C.A., 125 (1996) 109176n - a review with 56 refs.
- 355 Grigg, M.E., Gounaris, K. and Selkirk, M.E.: Characterization of a platelet-activating factor acetylhydrolase secreted by the nematode *Nippostrongylus brasiliensis*. *Biochem. J.*, 317 (1996) 541-547.
- 356 Hirano, T. and Takahashi, K.: Improvement in resolution of low melting point triglyceride molecular species on chilled column isocratic liquid chromatography. *J. Chromatogr. Sci.*, 34 (1996) 341-348.
- 357 Ioneda, T. and Ono, S.S.: Chromatographic and mass spectrometric analyses of 1-monomycoloyl glycerol fraction from *Rhodococcus lentifragimentus* as per-O-benzoyl derivatives. *Chem. Phys. Lipids*, 81 (1996) 11-19; C.A., 125 (1996) 242048p.
- 358 Keller, R.K.: Squalene synthase inhibition alters metabolism of nonsterols in rat liver. *Biochim. Biophys. Acta*, 1303 (1996) 169-179.
- 359 Korytowski, W. and Girotti, A.W.: Lipid hydroperoxide analysis by reverse-phase high-performance liquid chromatography with mercury cathode electrochemical detection. In: Favier, A.E. (Editor), *Anal. Free Radicals Biol. Syst.*, Birkhaeuser, Basel, 1995, pp. 165-183; C.A., 125 (1996) 52638h.
- 360 Lipp, M.: Comparison of PLS, PCR and MLR for the quantitative determination of foreign oils and fats in butter fats of several European countries by their triglyceride composition. *Z. Lebensm.-Unters. Forsch.*, 202 (1996) 193-198; C.A., 125 (1996) 32202h.
- 361 List, G.R., Mounts, T.L., Orthoefer, F. and Neff, W.E.: Potential maragrine oils from genetically modified soybeans. *J. Am. Oil Chem. Soc.*, 73 (1996) 729-732.
- 362 Márquez-Ruiz, G., Jorge, N., Martín-Polvillo, M. and Dobarganes, M.C.: Rapid, quantitative determination of polar compounds in fats and oils by solid-phase extraction and size-exclusion chromatography using monostearin as internal standard. *J. Chromatogr. A*, 749 (1996) 55-60.
- 363 Matsuyama, T. and Nakagawa, Y.: Surface-active exolipids: analysis of absolute chemical structures and biological functions. *J. Microbiol. Methods*, 25 (1996) 165-175; C.A., 125 (1996) 136807a.
- 364 Moreau, R.A., Powell, M.J. and Hicks, K.B.: Extraction and quantitative analysis of oil from commercial corn fiber. *J. Agric. Food Chem.*, 44 (1996) 2149-2154; C.A., 125 (1996) 113034g.
- 365 Muethling, J., Spanbroek, R., Peter-Katalinic, J., Hanisch, F.-G., Hanski, C., Hasegawa, A., Unland, F., Lehmann, J., Tschesche, H. and Egge, H.: Isolation and structural characterization of fucosylated gangliosides with linear poly-N-acetyllactosaminyl chains from human granulocytes. *Glycobiology*, 6 (1996) 147-156; C.A., 125 (1996) 52958n.
- 366 Murphy, E.J., Rosenberger, T.A. and Horrocks, L.A.: Separation of neutral lipids by high-performance liquid chromatography: quantification by ultraviolet, light scattering and fluorescence detection. *J. Chromatogr. B*, 685 (1996) 9-14.

- 367 Negroni, E., Chigorno, V., Tettamanti, G. and Sonnino, S.: Evaluation of the efficiency of an assay procedure for gangliosides in human serum. *Glycoconjugate J.*, 13 (1996) 347-352; *C.A.*, 125 (1996) 80902h.
- 368 Norman, H.A., Mischke, C.F., Allen, B. and Vincent, J.S.: Semi-preparative isolation of plant sulfoquinovosyldiacylglycerols by solid phase extraction and HPLC procedures. *J. Lipid Res.*, 37 (1996) 1372-1376.
- 369 Osman, M., Ishigami, Y., Someya, J. and Jensen, H.B.: The bioconversion of ethanol to biosurfactants and dye by a novel coproduction technique. *J. Am. Oil Chem. Soc.*, 73 (1996) 851-856.
- 370 Ritov, V.B., Banni, S., Yalowich, J.C., Day, B.W., Claycamp, H.G., Corongiu, F.P. and Kagan, V.E.: Non-random peroxidation of different classes of membrane phospholipids in live cells detected by metabolically integrated *cis*-parinaric acid. *Biochim. Biophys. Acta*, 1283 (1996) 127-140.
- 371 Stoll, U.: (Identification of molecular species of phospholipids from pig heart by combining chromatographic techniques). *Fette/Lipid*, 98 (1996) 178-182; *C.A.*, 125 (1996) 216022f.
- 372 Suda, Y., Shiyma, T., Yasukochi, T., Kawano, K., Brade, H., Rietschel, E.T. and Kusumoto, S.: Separation of wild-type lipopolysaccharides by centrifugal partition chromatography: isolation of smooth form species. *J. Endotoxin Res.*, 3 (1996) 1-7; *C.A.*, 125 (1996) 29477w.
- 373 Sugiura, T., Kondo, S., Sukagawa, A., Tonegawa, T., Nakane, S., Yamashita, A., Ishima, Y. and Waku, K.: Transacylase-mediated and phosphodiesterase-mediated synthesis of N-arachidonoylethanolamine, an endogenous cannabinoid-receptor ligand, in rat brain microsomes. Comparison with synthesis from free arachidonic acid and ethanolamine. *Eur. J. Biochem.*, 240 (1996) 53-62.
- 374 Tanaka, R., Miyahara, K. and Noda, N.: Zwitterionic digalactosylceramides from the earthworm, *Pheretima asiatica*. *Chem. Pharm. Bull.*, 44 (1996) 1152-1156.
- 375 Wang, Y.H., Hong, C.Y., Chen, C.F. and Tsai, T.H.: Determination of triacylglycerols in *Panax pseudo-ginseng* by HPLC polymeric column. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2497-2503.
- 376 Wang, Z., Gurel, O., Baatz, J.E. and Notter, R.H.: Differential activity and lack of synergy of lung surfactant protein SP-B and AP-C in interaction with phospholipids. *J. Lipid Res.*, 37 (1996) 1749.
- 377 Whitaker, B.D.: Cerebrosides in mature-green and red-ripe bell pepper and tomato fruits. *Phytochemistry*, 42 (1996) 627-632; *C.A.*, 125 (1996) 56728k.
- 378 Yamamoto, M.: (A new method for analysis of phospholipids by high-performance liquid chromatography and biliary phospholipids in patients with hepatolithiasis). *Wakayama Igaku*, 46 (1995) 483-492; *C.A.*, 125 (1996) 162454c.
- For additional information see *C.A.*:
125 (1996) 52658g.
- See also 164.
- 11d. Lipoproteins and their constituents**
- 379 Atalto-Setala, K., Weinstock, P.H., Bisgaier, C.L., Wu, I., Smith, J.D. and Breslow, J.L.: Further characterization of the metabolic properties of triglyceride-rich lipoproteins from human and mouse apoC-III transgenic mice. *J. Lipid Res.*, 37 (1996) 1802-1811.
- 380 Edelstein, C., Italia, J.A., Klezovitch, O. and Scanu, A.M.: Functional and metabolic differences between elastase-generated fragments of human lipoprotein(A) and apolipoprotein(A). *J. Lipid Res.*, 37 (1996) 1786-1801.
- 381 Kitamura, T., Kato, Y., Okazaki, M. and Sasamoto, K.: (Antibody-containing packing material for separating lipoproteins). *Jpn. Kokai Tokkyo Koho JP 08,105,875 [96,105,875] (Cl. G01N30/48)*, 23 Apr. 1996, JP Appl. 94/190,510, 12 Aug. 1994; 6 pp.; *C.A.*, 125 (1996) 81273r.
- 382 Li, D., Devaraj, S., Fuller, C., Bucala, R. and Jialal, I.: Effect of α -tocopherol on LDL oxidation and glycation: *in vitro* and *in vivo* studies. *J. Lipid Res.*, 37 (1996) 1978-1986.
- 383 Masson, D., Athias, A. and Lagrost, L.: Evidence for electronegativity of plasma high density lipoprotein-3 as one major determinant of human cholesterol ester transfer protein activity. *J. Lipid Res.*, 37 (1996) 1579-1590.
- For additional information see *C.A.*:
125 (1996) 109649u.
- See also 353, 1209.
- 12. ORGANIC PEROXIDES**
- See 359.
- 13. STEROIDS**
- 13a. General techniques**
- 384 Hyttäinen, M., Appelblad, P., Pontén, E., Stigbrand, M., Irgum, K. and Jaegfeldt, H.: Trifluoromethanesulfonic acid as a catalyst for the formation of dansylhydrazone derivatives. *J. Chromatogr. A*, 740 (1996) 279-283.
- 385 Kummer, M., Palme, H.-J. and Werner, G.: Resolution of enantiomeric steroids by high-performance liquid chromatography on chiral stationary phases. *J. Chromatogr. A*, 749 (1996) 61-68.
- See also 52.
- 13b. Pregnane and androstane derivatives**
- 386 Cooper, A.E., Gray, A.J., Collington, J., Seddon, H., Beattie, I. and Logan, C.J.: Excretion and metabolism of tipredane, a novel glucocorticoid, in the rat, mouse, monkey, and human. *Drug Metab. Disp.*, 24 (1996) 1071-1080.

- 387 Nakajima, M., Wakabayashi, H., Yamato, S. and Shimada, K.: (Quantification of serum dehydroepiandrosterone sulfate by liquid chromatography/atmospheric pressure chemical ionization). *Bunseki Kagaku*, 45 (1996) 517-522; C.A., 125 (1996) 76658r.
- 388 Shimada, K. and Nakagi, T.: Studies on neurosteroids. IV. Quantitative determination of pregnenolone in rat brains using high-performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2597-2602.
- 389 Sugano, S., Miura, R. and Morishima, N.: Identification of intermediates in the conversion of cholesterol to pregnenolone with a reconstituted cytochrome P-450_{ccc} system: accumulation of the intermediate modulated by the adrenodoxin level. *J. Biochem. (Tokyo)*, 120 (1996) 780-787.

See also 127, 133, 1006.

13c. Estrogens

- 390 Nascimento, E.S., Salvadori, M.C. and Ribeiro-Neto, L.M.: Determination of synthetic estrogens in illegal veterinary formulations by HPTLC and HPLC. *J. Chromatogr. Sci.*, 34 (1996) 330-333.

13d. Sterols

- 391 Akihisa, T., Kimura, Y., Kokke, W.C.M.C., Itoh, T. and Tamura, T.: Eight novel sterols from the roots of *Bryonia dioica* Jacq. *Chem. Pharm. Bull.*, 44 (1996) 1202-1207.
- 392 Biedermann, M., Grob, K., Mariani, C. and Schmidt, J.P.: Detection of desterolized sunflower oil in olive oil through isomerized Δ7-sterols. *Z. Lebensm.-Unters. Forsch.*, 202 (1996) 199-204; C.A., 125 (1996) 56537x.
- 393 Gallina Toschi, T., Bendini, A. and Lercker, G.: Evaluation of 3,5-stigmastadiene content of edible oils: comparison between the traditional capillary gas chromatographic method and the on-line high performance liquid chromatography-capillary gas chromatographic analysis. *Chromatographia*, 43 (1996) 195-199.
- 394 Roccataglia, A.J., Maier, M.S., Seldes, A.M., Pujol, C.A. and Damonte, E.B.: Antiviral sulfated steroids from the ophiuroid *Ophioplocus januarii*. *J. Natural Prod.*, 59 (1996) 887-889.
- 395 Wilson, W.K., Sumpter, E.M., Warren, J.J., Rogers, P.S., Ruan, B. and Schroepfner, G.J.: Analysis of unsaturated C₂₇ sterols by nuclear magnetic resonance spectroscopy. *J. Lipid Res.*, 37 (1996) 1529-1555.

See also 366, 1179.

13e. Bile acids and alcohols

- 396 Fernandez, J.I., Manas, M., Rueda, A., Martinez-Victoria, E. and Naranjo, J.A.: Taurine- and glycine-conjugated bile acids in the preruminant goat: identification by high performance liquid chromatography. *Br. Vet. J.*, 152 (1996) 433-439; C.A., 125 (1996) 189728g.
- 397 Mikami, T., Ohshima, A., Mosbach, E.H., Cohen, B.I., Ayyard, N., Yoshii, M., Ohtani, K., Kihira, K., Schteingart, C.D. and Hoshita, T.: 15α-Hydroxylation of a bile analogue, sodium 3α,7α-dihydro-25, 26-bishomo-5β-cholane-26-sulfonate in the hamster. *J. Lipid Res.*, 37 (1996) 1189-1197.

- 398 Paauw, J.D., van Wyk, L. and Davis, A.T.: Assay for taurine conjugates of bile acids in serum by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 685 (1996) 171-175.

See also 990.

13f. Ecdysones and other insect steroid hormones

- 399 Bathori, M.I.: Droplet counter-current chromatography to isolate ecdysteroids from the herbs of *Silena tatarica* (L.) Pers. *Acta Pharm. Hung.*, 66 (1996) 125-131; C.A., 125 (1996) 95722y.
- 400 Sarker, S.D., Dinan, L.N., Girault, J.-P., Laafont, R. and Waterman, P.G.: Punisterone [(20R,24S)-25-deoxy-11α,20,24-trihydroxyecdystone]: a new phytoecdysteroid from *Blandfordia punicea*. *J. Natural Prod.*, 59 (1996) 789-793.

13g. Other steroids

- 401 Gamoh, K., Abe, H., Shimada, K. and Takatsuto, S.: Liquid chromatography/mass spectrometry with atmospheric pressure chemical ionization of free brassinosteroids. *Rapid Commun. Mass Spectrom.*, 10 (1996) 903-906; C.A., 125 (1996) 109543e.
- 402 Gamoh, K., Prescott, M.C., Goad, L.J. and Takatsuto, S.: Analysis of brassinosteroids by liquid chromatography/mass spectrometry. *Bunseki Kagaku*, 45 (1996) 523-527; C.A., 125 (1996) 103864p.

14. STEROID GLYCOSIDES AND SAPONINS

- 403 Ruiz, R.G., Price, K.R., Rose, M.E., Rhodes, M.J.C. and Fenwick, G.R.: High-performance liquid chromatography of lupin and lentil saponins. *Spec. Publ.-R. Soc. Chem.*, 179(Agri-Food Quality) (1996) 254-257; C.A., 125 (1996) 242051j.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 404 Chen, K.-S., Chang, F.-R., Jong, T.-T. and Wu, Y.-C.: Two novel sesquiterpenes from *Neolitsea parvigemma*. *J. Natural Prod.*, 59 (1996) 804-806.
- 405 Guenther, B. and Wagner, H.: Quantitative determination of triterpenes in extracts and phytopreparations of *Centella asiatica* (L.) Urban. *Phytomedicine*, 3 (1996) 59-65; C.A., 125 (1996) 151272r.
- 406 Inamdar, P.K., Yeole, R.D., Ghogare, A.B. and de Souza, N.J.: Determination of biologically active constituents in *Centella asiatica*. *J. Chromatogr. A*, 742 (1996) 127-130.
- 407 Kim, K.K., Sawa, Y. and Shibata, H.: Hydroxylation of entkaurenoic acid to steviol in *Stevia rebaudiana* Bertoni - purification and partial characterization of the enzyme. *Arch. Biochem. Biophys.*, 332 (1996) 223-230.

- 408 Németh-Kiss, V., Forgács, E., Cserháti, T. and Schmidt, G.: Determination of taxol in *Taxus* species grown in Hungary by high-performance liquid chromatography-diode array detection. Effect of vegetative period. *J. Chromatogr. A*, 750 (1996) 253-256.
- 409 Ney, I., Jaeger, E., Herderich, M., Schreier, P. and Schwab, W.: Analysis of phosphorylated terpenes by electrospray ionization liquid chromatography tandem mass spectrometry. *Phytochem. Anal.*, 7 (1996) 233-236; C.A., 125 (1996) 269779q.
- 410 Theodoridis, G. and Verpoorte, R.: Taxol analysis by high-performance liquid chromatography. *Phytochem. Anal.*, 7 (1996) 169-184; C.A., 125 (1996) 109169n - a review with many refs.
- 411 Vandana, V., Teja, A.S. and Zalkow, L.H.: Supercritical extraction and HPLC analysis of taxol from *Taxus brevifolia* using nitrous oxide and nitrous oxide+ethanol mixtures. *Fluid Phase Equilib.*, 116 (1996) 162-169; C.A., 125 (1996) 29359j.

See also 358, 412, 894.

15b. Essential oils

- 412 Govindachari, T.R., Gopalakrishnan, G. and Suresh, G.: Isolation of various azadirachtins from neem oil by preparative high performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1729-1733.
- 413 Mondello, L., Dugo, G., Dugo, P. and Bartle, K.D.: Online HPLC-HRGC in the analytical chemistry of citrus essential oils. *Perfum. Flavor*, 21 (1996) 25-49; C.A., 125 (1996) 150713 - a review with 48 refs.

For additional information see C.A.:
125 (1996) 157325g.

16. NITRO AND NITROSO COMPOUNDS

- 414 Bodzek, D., Janoszka, B. and Warzecha, L.: The analysis of PAHs, nitrogen derivatives in the sewage sludges of Upper Silesia, Poland. *Water, Air, Soil Pollut.*, 89 (1996) 417-427; C.A., 125 (1996) 94680w.
- 415 Cappiello, A., Famiglini, G., Lombardozzi, A., Massari, A. and Vadalia, G.G.: Electron capture ionization of explosives with a microflow rate particle beam interface. *J. Am. Soc. Mass Spectrom.*, 7 (1996) 753-758; C.A., 125 (1996) 118928m.
- 416 Meng, H.-h. and Caddy, B.: High-performance liquid chromatographic analysis with fluorescence detection of ethyl centralite and 2,4-dinitrotoluene in gunshot residues after derivatization with 9-fluorenyl-methylchloroformate. *J. Forensic Sci.*, 41 (1996) 213-220; C.A., 125 (1996) 78737q.
- 417 Mora Torres, R., Grosset, C., Steiman, R. and Alary, J.: Liquid chromatography study of degradation and metabolism of pentachloronitrobenzene by four soil micromycetes. *Chemosphere*, 33 (1996) 683-692; C.A., 125 (1996) 160507y.
- 418 Yoshikawa, M., Bessho, J. and Arashidani, K.: Separation of major mononitro-aromatic compounds by reversed-phase high-performance liquid chromatography. *Ind. Health*, 34 (1996) 177-183; C.A., 125 (1996) 149567x.

For additional information see C.A.:
125 (1996) 90817y.

See also 247, 419.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 419 Djozan, Dj. and Faraj-Zadeh, M.A.: Liquid chromatographic determination of nitroaromatics in water following solid phase extraction, pre-column reduction and derivatization. *Chromatographia*, 43 (1996) 25-30.
- 420 Martinsen, A., Hyvonen, S. and Gynther, J.: Evaluation of reversed phase columns for the analysis of diphenylmethane derivatives. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2579-2592.

For additional information see C.A.:
125 (1996) 103904b.

See also 81, 96, 170, 259, 426, 792.

17b. Catecholamines and their metabolites

- 421 Alemany, G., Garmundi, A., Rossello, C. and Rial, R.: Simultaneous determination of vanillylmandelic, homovanillic and 5-hydroxyindoleacetic acids in human urine by thin layer chromatography. *Biomed. Chromatogr.*, 10 (1996) 144-145; C.A., 125 (1996) 52648m.
- 422 Bockhardt, A., Krause, I. and Klostermeyer, H.: Determination of biogenic amines by RP-HPLC of the dansyl derivates. *Z. Lebensm.-Unters. Forsch.*, 203 (1996) 65-70; C.A., 125 (1996) 193788u.
- 423 Brandsteterova, E., Kubalec, P., Krajnak, K. and Skacani, I.: SPE-HPLC determination of catecholamines using an affinity principle. *Neoplasma*, 43 (1996) 107-112; C.A., 125 (1996) 212798r.
- 424 Gumbhir, K. and Mason, W.D.: High-performance liquid chromatographic determination of phenylephrine and its conjugates in human plasma using solid-phase extraction and electrochemical detection. *J. Pharm. Biomed. Anal.*, 14 (1996) 623-630.
- 425 Hernandez-Jover, T., Izquierdo-Pulido, M., Veciana-Nogues, M.T. and Vidal-Carou, M.C.: Ion-pair high-performance liquid chromatographic determination of biogenic amines in meat and meat products. *J. Agric. Food Chem.*, 44 (1996) 2217-2222; C.A., 125 (1996) 165923r.
- 426 Holland, L.A.: Development of a comprehensive two-dimensional microcolumn liquid chromatography system for the separation of nanoliter volumes of biological amines. *Diss. Abstr. Int., B*, 57 (1996) 3175; C.A., 125 (1996) 242066t.
- 427 Mahendradatta, M. and Schwedt, G.: (Determination of histamine in foods by capillary electrophoresis and HPLC in comparison). *Dtsch. Lebensm.-Rundsch.*, 92 (1996) 218-221; C.A., 125 (1996) 193792r.
- 428 Yang, D.-Y., Chia, L.-G., Ho, W.-M., Yang, C.-S., Kuo, J.-S. and Cheng, F.-C.: Determination of biogenic amines and their metabolites in body fluids using ultrafiltration and microbore liquid chromatography with a dual amperometric detector. *Am. Lab. (Shelton)*, 28 (1996) 64-67; C.A., 125 (1996) 52611u.

For additional information see C.A.:
125 (1996) 131274q.

17c. Urea and guanidine derivatives

- 429 Anonymous: Analysis of guanidino compounds by protamine column. *Kitasato Igaku*, 26 (1996) 102-103; C.A., 125 (1996) 136806z.
 430 Tanigawa, T., Mizo-oku, Y., Moriguchi, K., Suzuki, T., Osumi, T. and Odomi, M.: Simple and rapid quantitative assay of ¹³C-labelled urea in human serum using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *J. Chromatogr. B*, 683 (1996) 135-142.

17d. Other amine derivatives and amides (excl. peptides)

- 431 Andersson, S., Allenmark, S., Möller, P., Persson, B. and Sanchez, D.: Chromatographic separation of enantiomers on N,N'-diallyl-L-tartardiamide-based network-polymeric chiral stationary phases. *J. Chromatogr. A*, 741 (1996) 23-31.
 432 Matsuzawa, M. and Kawai, H.: (Determination of choline in spices and beans by HPLC with an electrochemical detector). *Shokuhin Eiseigaku Zasshi*, 37 (1996) 72-76; C.A., 125 (1996) 84954z.
 433 Toyota, E., Itoh, K., Sekizaki, H. and Tanizawa, K.: Chromatographic separation of diastereomeric Schiff base copper(II), nickel(II), and zinc(II) chelates from α -amino racemates. *Bioorg. Chem.*, 24 (1996) 150-158; C.A., 125 (1996) 276448j.
 434 Wood, S.A., Pegg, S.N., Simmonds, R.J. and Stevenson, D.: Development of chiral high performance liquid chromatographic assays for the enantiomers of 8-hydroxy-(di-n-propyl-amino)tetralin and three 8-keto-pyrrole-substituted analogues. *J. Pharm. Biomed. Anal.*, 14 (1996) 1591-1599.

See also 166, 247, 1168, 1177.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 435 Avery, N.C.: The use of solid phase extraction cartridges as a pre-fractionation step in the quantitation of intermolecular collagen crosslinks and advanced glycation end-products. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1831-1848.
 436 Bank, R.A., Jansen, E.J., Beekman, B. and te Koppele, J.M.: Amino acid analysis by reverse-phase high-performance liquid chromatography: improved derivatization/and detection conditions with 9-fluorenylmethyl chloroformate. *Anal. Biochem.*, 240 (1996) 1167-1176.
 437 Deulofeu, R., Giralt, M., Aibar, C., Bauchet, C., Varela-Moreiras, G., Casals, F., Chamorro, A., Vila, N., Diaz Cremades, J. and Balles, A.M.: (HPLC determination of homocysteine in plasma. Application to the study of peripheral and cerebrovascular disease). *Rev. Soc. Esp. Bioquim. Clin. Patol. Mol.*, 15 (1996) 77-84; C.A., 125 (1996) 52653j.
 438 Gopalakrishnan, V., Burton, P.J. and Blaschke, T.F.: High-performance liquid chromatographic assay for the quantitation of L-arginine in human plasma. *Anal. Chem.*, 68 (1996) 3520-3523.

- 439 Graham, L.: A comprehensive survey of the acid-stable fluorescent cross-links formed by ribose with basic amino acids, and partial characterization of a novel Maillard cross-link. *Biochim. Biophys. Acta*, 1297 (1996) 9-16.
 440 Hyun, M.H. and Ryoo, J.-J.: Optical resolution of racemic α -amino acids on a dynamic chiral stationary phase derived from (S)-leucinol by ligand exchange chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2635-2648.
 441 Jacob, J.S., Cistola, D.P., Hsu, F.F., Muzaffar, S., Mueller, D.M., Hazen, S.L. and Heinecke, J.W.: Human phagocytes employ the myeloperoxidase-hydrogen peroxide system to synthesize dityrosine, tyrosine, pulcherosine, and isodityrosine by a tyrosyl radical-dependent pathway. *J. Biol. Chem.*, 271 (1996) 19950-19956.
 442 Khedr, A.: High-performance liquid chromatography of α -amino acids and aztreonam on reversed phase columns with aqueous Cu²⁺ as eluent. *Biomed. Chromatogr.*, 10 (1996) 167-171; C.A., 125 (1996) 136800t.
 443 Kirschbaum, J. and Brückner, H.: Amino acid analysis by derivatization with chromogenic 4-phenylazobenzoyloxycarbonyl chloride (PAZ-Cl): comparison of reversed-phases. *Chromatographia*, 43 (1996) 275-278.
 444 Langford, S.A., Sugden, J.K. and Fitzpatrick, R.W.: Detection and determination of the hydrazo and azo photoproducts of 4-aminobenzoic acid by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 1615-1623.
 445 Lapčík, O., Hampl, R., Hill, M., Starka, L., Kasal, A., Pouzar, V. and Putz, Z.: Radioimmunological and chromatographic properties of tyrosine methyl ester conjugates with stereoisomeric steroid carboxy derivatives. *Collect. Czech. Chem. Commun.*, 61 (1996) 799-807; C.A., 125 (1996) 143118e.
 446 Liu, C.-S., Lin, C.-C., Chen, J.-M., Chang, C.-H. and Lo, T.-B.: A convenient amino acid analysis of proteins electroblotted onto polyvinylidene difluoride [PVDF] membrane from sodium dodecyl sulfate-polyacrylamide gel - a direct *in situ* derivatization of amino acids after gas-phase hydrolysis. *J. Chin. Biochem. Soc.*, 22 (1993) 69-76; C.A., 125 (1996) 29462n.
 447 Marioli, J.M. and Sereno, L.E.: Electrochemical detection of undervatized amino acids with a Ni-Cr alloy electrode. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2505-2515.
 448 Sandee, B., Schipper, C.A. and Eertman, R.H.M.: High-performance liquid chromatographic determination of the imino acids (opines) meso-alanopine and D-strombine in muscle extract of invertebrates. *J. Chromatogr. B*, 685 (1996) 176-180.
 449 Shang, S.F. and Wang, H.: Sensitive determination of amino acids in kelp by reversed phase high performance liquid chromatography with precolumn derivatization using phenylisothiocyanate. *Chromatographia*, 43 (1996) 309-312.
 450 Simat, T., van Wickern, B., Eulitz, K. and Steinhart, H.: Contaminants in biotechnologically manufactured L-tryptophan. *J. Chromatogr. B*, 685 (1996) 41-51.
 451 Strydom, D.J.: Amino acid analysis using various carbamate reagents for precolumn derivatization. In: Marshak, D.R. (Editor), *Tech. Protein Chem. VII, [Symp. Protein Soc.]*, 9th 1995, Academic, San Diego, 1996, pp. 331-339; C.A., 125 (1996) 242264f.
 452 Takahashi, M., Hoshino, H., Kushida, K., Kawana, K. and Inoue, T.: Direct quantification of pentosidine in urine and serum by HPLC with column switching. *Clin. Chem. (Washington)*, 42 (1996) 1439-1444.

- 453 Takahashi, M., Terashima, K., Nishijima, M. and Kamata, K.: Separation of carnitine enantiomers as the 9-anthroylnitrile derivatives and high-performance liquid chromatographic analysis on an ovomucoid-conjugated column. *J. Pharm. Biomed. Anal.*, 14 (1996) 1579-1584.
- 454 Takeuchi, T. and Miwa, T.: Selectivity of dansyl amino acids in micellar liquid chromatography with an ion-exchange-induced stationary phase. *Chromatographia*, 43 (1996) 143-148.
- 455 Thompson, M.G., Palmer, R.M., Thom, A., Garden, K., Lobley, G.E. and Calder, G.: N⁶-Methylhistidine turnover in skeletal muscle cells measured by GC-MS. *Am. J. Physiol.*, 270 (1996) C1875-C1879; C.A., 125 (1996) 137125p.
- 456 Yuan, Y., Shen, M. and Li, J.: (Determination of 3-methylhistidine in urine by high performance liquid chromatography-fluorescence detection). *Sepu*, 14 (1996) 274-276; C.A., 125 (1996) 189722a.
- For additional information see C.A.:
125 (1996) 67948h, 222435j.
- See also 75, 105, 116, 167, 169, 170, 171, 486, 821, 1018, 1212.
- 18b. Peptides, peptidic and proteinous hormones, growth factors
- 457 Bach, L.A. and Rechler, M.M.: Measurement of insulin-like growth factor (IGF)-II binding to purified IGF binding proteins 1-6: comparison of charcoal adsorption and high performance size exclusion chromatography. *Biochim. Biophys. Acta*, 1313 (1996) 79-88.
- 458 Battersby, J.E., Canova-Davis, E., Celniker, A.C., Cunningham, B.C., Fletcher, B.L., Garnick, R.L., Gehant, R.L., Hoff, E.R., Hunt, G., et al.: High performance receptor binding chromatography (HPRBC) for human growth hormone (hGH). *Altern. Methods Toxicol. Life Sci.*, 11(The World Congress on Alternatives and Animal Use in the Life Sciences: Education, Research, Testing, 1993) (1995) 261-267; C.A., 125 (1996) 76657q.
- 459 Blishchenko, E.Y., Mernenko, O.A., Yatskin, O.N., Ziganshin, R.N., Philippova, M.M., Karelina, A.A. and Ivanov, V.T.: Neokytorphin and neokytorphin (1-4): cytolytic activity and comparative levels in rat tissues. *Biochem. Biophys. Res. Commun.*, 224 (1996) 721-727.
- 460 Breslav, M., McKinney, A., Becker, J.M. and Naider, F.: Preparation of radiolabeled peptides via an iodine exchange reaction. *Anal. Biochem.*, 239 (1996) 213-217.
- 461 Brovia, V., Ricciardi, A. and Barbeito, L.: N-Acetylaspartylglutamate (NAAG) in human cerebrospinal fluid: determination by high performance liquid chromatography, and influence of biological variables. *Amino Acids*, 9 (1995) 175-184; C.A., 125 (1996) 269514z.
- 462 Chang, J.: The disulfide folding pathway of tick anticoagulant peptide (TAP), a Kunitz-type inhibitor structurally homologous to BPTI. *Biochemistry*, 35 (1996) 11702-11709.
- 463 Cho, B.-Y., Zou, H., Strong, R., Fisher, D.H., Nappier, J. and Krull, I.S.: Immunochromatographic analysis of bovine growth hormone releasing factor involving reversed-phase high-performance liquid chromatography-immunodetection. *J. Chromatogr. A*, 743 (1996) 181-194.

- 464 Dai, Y. and Tang, J.-G.: Characteristic, activity and conformational studies of [A6-Ser, A11-Ser]-insulin. *Biochim. Biophys. Acta*, 1296 (1996) 63-68.
- 465 Desiderio, D.M.: Mass spectrometry, high performance liquid chromatography, and brain peptides. *Biopolymers*, 40 (1996) 257-264; C.A., 125 (1996) 109183n - a review with 45 refs.
- 466 Fujinari, E.M., Manes, J.D. and Bizek, R.: Peptide content determination of crude synthetic peptides by reversed-phase liquid chromatography and nitrogen-specific detection with a chemiluminescent nitrogen detector. *J. Chromatogr. A*, 743 (1996) 85-89.
- 467 Giraud, A., Whitley, J., Shulkes, A. and Parker, L.: The pregnant ovine endometrium constitutively expresses and secretes a highly stable bombesin-like peptide, which shares C-terminal sequence but differs structurally from gastrin-releasing peptide. *Biochim. Biophys. Acta*, 1296 (1996) 189-197.
- 468 Hill, H.M., Robson, J.N., Noctor, T.A.G., Tennant, K., Thomas, D., Clarke, S.D. and Burnett, J.E.C.: Sample preparation and subsequent steps: strategies for peptides and peptidomimetics. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 48-56; C.A., 125 (1996) 264638h.
- 469 Hubert, F., Noël, T. and Roch, P.: A member of the arthropod defensin family from edible Mediterranean mussels (*Mytilus galloprovincialis*). *Eur. J. Biochem.*, 240 (1996) 302-306.
- 470 Khosravi, M.J., Diamandi, A., Mistry, J. and Lee, P.D.K.: Non-competitive ELISA for human serum insulin-like growth factor-I. *Clin. Chem. (Washington)*, 42 (1996) 1147-1154.
- 471 Lakhiani, H., Jozefowicz, J. and Muller, D.: Separation and purification of insulins on coated silica support functionalized with sialic acid by affinity chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2423-2441.
- 472 Lee, T.D., Davis, M.T., Stahl, D.C., Swiderek, K.M., Esworthy, R.S. and Hefta, S.A.: High sensitivity tandem liquid chromatography-mass spectrometry analysis of peptides and proteins. *Nippon Iyo Masu Supekutoro Gakkai Koenshu*, 21 (1996) 17-32; C.A., 125 (1996) 269799w.
- 473 Lu, L. and Lundahl, P.: Quantitative micellar chromatographic analysis of interaction between peptides and sodium dodecyl sulfate micelles. *J. Chromatogr. A*, 741 (1996) 49-57.
- 474 Marcinkiewicz, C., Rosenthal, L., Mosser, D.M., Kunick, T.J. and Niewarowski, S.: Immunological characterization of eristostatin and echistatin binding sites on $\alpha_{IIb}\beta_3$ and $\alpha\beta_3$ integrins. *Biochem. J.*, 317 (1996) 817-825.
- 475 Miller, C. and Rivier, J.: Peptide chemistry: development of high-performance liquid chromatography and capillary zone electrophoresis. *Biopolymers*, 40 (1996) 265-317; C.A., 125 (1996) 109184p - a review with 124 refs.
- 476 Millington, W.R. and Manning, A.B.: Chromatographic methods for analyzing β -endorphin and related peptides. In: Thomas, J.A. (Editor), *Endocr. Methods*, Academic, San Diego, 1996, pp. 281-297; C.A., 125 (1996) 105235h - a review with 42 refs.
- 477 Müller, S., Dunker, R. and Hochhaus, G.: Interspecies comparison of *in vitro* plasma degradation of dynorphin A1-13. *Pharmazie*, 51 (1996) 581-585.
- 478 Nice, E.C.: Micropreparative high-performance liquid chromatography of proteins and peptides: principles and applications. *Biopolymers*, 40 (1996) 319-341; C.A., 125 (1996) 109185q - a review with 151 refs.

- 479 Pearson, J.D. and McCroskey, M.C.: Perfluorinated acid alternatives to trifluoroacetic acid for reversed-phase high-performance liquid chromatography. *J. Chromatogr. A.*, 746 (1996) 277-281.
- 480 Pisarev, O.A. and Samsonov, G.V.: (Role of mobile and stationary functional domains of net structures of carboxylic polyelectrolytes in insulin sorption and reversible desorption kinetics). *Zh. Prikl. Khim. (S.-Peterburg)*, 69 (1996) 299-301; *C.A.*, 125 (1996) 158781q.
- 481 Sanderson, K., Andrén, P.E., Caprioli, R.M. and Nyberg, F.: *In vitro* metabolism of LVV-hemorphin-7 in human plasma studied by reversed-phase high-performance liquid chromatography and micro-electrospray mass spectrometry. *J. Chromatogr. A*, 743 (1996) 207-212.
- 482 Schnapp, D., Kemp, G.D. and Smith, V.J.: Purification and characterization of a proline-rich antibacterial peptide, with sequence similarity to bactenecin-7, from the haemocytes of the shore crab, *Carcinus maenas*. *Eur. J. Biochem.*, 240 (1996) 532-539.
- 483 Seeger, A. and Rinas, U.: Two-step chromatographic procedure for purification of basic fibroblast growth factor from recombinant *Escherichia coli* and characterization of the equilibrium parameters of adsorption. *J. Chromatogr. A*, 746 (1996) 17-24.
- 484 Stahl, D.C., Swiderek, K.M., Davis, M.T. and Lee, T.D.: Data-controlled automation of liquid chromatography/tandem mass spectrometry analysis of peptide mixtures. *J. Am. Soc. Mass Spectrom.*, 7 (1996) 532-540; *C.A.*, 125 (1996) 52929d.
- 485 Taliani, M., Bianchi, E., Narjes, F., Fossatelli, M., Urbani, A., Steinkübler, C., de Francesco, R. and Pessi, A.: A continuous assay of hepatitis C virus protease based on resonance energy transfer depsipeptide substrates. *Anal. Biochem.*, 240 (1996) 60-67.
- 486 Tang, Y., Zukowski, J. and Armstrong, D.W.: Investigation on enantiomeric separations of fluorenylmethoxycarbonyl amino acids and peptides by high-performance liquid chromatography using native cyclodextrins as chiral stationary phases. *J. Chromatogr. A*, 743 (1996) 261-271.
- 487 Vyas, S.B. and Duffy, L.K.: Equilibration of a synthetic Alzheimer β -protein analog, β 22-35, during RP-HPLC. *Protein Pept. Lett.*, 3 (1996) 89-94; *C.A.*, 125 (1996) 79720r.
- 488 Walker, T.A.: Micellar HPLC: investigation of the retention of positively charged peptides using cationic micellar mobile phases. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1715-1727.
- 489 Wen, J., Hsu, E., Kenney, W.C., Philo, J.S., Morris, C.F. and Arakawa, T.: Characterization of keratinocyte growth factor binding to heparin and dextran sulfate. *Arch. Biochem. Biophys.*, 332 (1996) 41-46.
- 490 Xiao, S., Dorris, M.L., Rawitch, A.B. and Taurog, A.: Selectivity in tyrosyl iodination sites in human thyroglobulin. *Arch. Biochem. Biophys.*, 334 (1996) 284-294.
- For additional information see *C.A.*:
125 (1996) 26442v, 131262j, 131263k.
- See also 79, 88, 91, 106, 113, 116, 297, 538, 549, 563, 578, 588, 704, 730, 1033, 1058, 1127, 1188.
- 18c. *Elucidation of structure of proteins and enzymes*
- 491 Banks, J.F., Jr.: High-sensitivity peptide mapping using packed-capillary liquid chromatography and electrospray ionization mass spectrometry. *J. Chromatogr. A*, 743 (1996) 99-104.
- 492 Corradini, D.: Peptide mapping by reversed phase chromatography. *LC-GC Int.*, 9 (1996) 732-740.
- 493 Frenz, J.: Trace analysis peptide mapping by high-performance displacement chromatography. *Methods Enzymol.*, 271(High Resolution Separation and Analysis of Biological Macromolecules) (1996) 486-504; *C.A.*, 125 (1996) 269506y.
- 494 Gegg, C.V., Bowers, K.E. and Matthews, C.R.: A general approach for the design and isolation of protein fragments: the molecular dissection of dihydrofolate reductase. In: Marshak, D.R. (Editor), *Tech. Protein Chem. VII, [Symp. Protein Soc.]*, 9th 1995, Academic, San Diego, 1996, pp. 439-448; *C.A.*, 125 (1996) 242055p.
- 495 Gordon, N., Nadler, T., Blackburn, C., Wang, H. and Speak, J.: Automated peptide mapping using a multidimensional chromatography workstation. *Am. Biotechnol. Lab.*, 14 (1996) 18-20; *C.A.*, 125 (1996) 29311n.
- 496 Hoff, E.R. and Chloupek, R.C.: Analytical peptide mapping of recombinant DNA-derived proteins by reversed-phase high-performance liquid chromatography. *Methods Enzymol.*, 271(High Resolution Separation and Analysis of Biological Macromolecules) (1996) 51-68; *C.A.*, 125 (1996) 269420r - a review with 32 refs.
- 497 Ishii, S., Mizuguchi, H., Nishino, J., Hayashi, H. and Kagamiyama, H.: Functionally important residues of aromatic L-amino acid decarboxylase probed by sequence alignment and site-directed mutagenesis. *J. Biochem. (Tokyo)*, 120 (1996) 369-376.
- 498 McIntosh, D.B., Parrish, J.C. and Wallace, C.J.A.: Definition of a nucleotide binding site on cytochrome c by photoaffinity labeling. *J. Biol. Chem.*, 271 (1996) 18379-18386.
- 499 Nadler, T., Blackburn, C., Mark, J., Gordon, N., Regnier, F.E. and Vella, G.: Automated proteolytic mapping of proteins. *J. Chromatogr. A*, 743 (1996) 91-98.
- 500 Parente, J.A., Jr., Goldenring, J.R., Petropoulos, A.C., Hellman, U. and Chew, C.S.: Purification, cloning, and expression of a novel, endogenous, calcium-sensitive, 28-kDa phosphoprotein. *J. Biol. Chem.*, 271 (1996) 20096-20101.
- 501 Ramalho-Santos, M., Veríssimo, P., Faro, C. and Pires, E.: Action on bovine α_1 -casein of cardosins A and B, aspartic proteinases from the flowers of the cardoon *Cynara cardunculus* L. *Biochim. Biophys. Acta*, 1297 (1996) 83-89.
- 502 Tanji, M., Yakabe, E., Kageyama, T. and Takahashi, K.: The primary structure of the major pepsinogen from the gastric mucosa of tuna stomach. *J. Biochem. (Tokyo)*, 120 (1996) 647-656.
- 503 Vissers, J.P.C., Chervet, J.-P. and Salzmann, J.-P.: Sodium dodecyl sulfate removal from tryptic digest samples for online capillary liquid chromatography/electrospray mass spectrometry. *J. Mass Spectrom.*, 31 (1996) 1021-1027; *C.A.*, 125 (1996) 269790m.
- 504 Zhao, Q., Lecoeur, C., Sannier, F., Garreau, I. and Piot, J.M.: Quantitative determination of aromatic amino acids at protein surface by size exclusion HPLC coupled with second order derivative spectroscopy. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1551-1566.

See also 556, 590, 620, 621, 634, 707.

19. PROTEINS

19a. General techniques

- 505 Andrews, B.A. and Asenjo, J.A.: Protein partitioning equilibrium between the aqueous poly(ethylene glycol) and salt phases and the solid protein phase in poly(ethylene glycol)-salt two-phase systems. *J. Chromatogr. B*, 685 (1996) 15-20.
- 506 Azarkan, M., Maes, D., Bouckaert, J., Thi, M.-H.D., Wyns, L. and Looze, Y.: Thiol pegylation facilitates purification of chymopain leading to diffraction studies at 1.4 Å resolution. *J. Chromatogr. A*, 749 (1996) 69-72.
- 507 Basak, S.K.: Electrically induced transport properties for protein separation by chromatography. Avail. *Univ. Microfilms Int.*, Order DA9622666, 1995, 180 pp.; C.A., 125 (1996) 136817d.
- 508 Bayer, E.A., Morag, E., Wilchek, M., Lamed, R. and Shoham, Y.: Modified cellulose-binding domain (CBD) proteins and their uses in affinity chromatography, immunoassays, enzyme reactors, and drug delivery. *PCT Int. Appl. WO 96 13,524* (Cl. C07K17/06), 9 May 1996, IL Appl. 111,415, 27 Oct. 1994; 53 pp.; C.A., 125 (1996) 81254k.
- 509 Castellanos-Serra, L.R., Fernandez-Patron, C., Hardy, E. and Huerta, V.: A procedure for protein elution from reverse-stained polyacrylamide gels applicable at the low picomole level: an alternative route to the preparation of low abundance proteins for microanalysis. *Electrophoresis (Weinheim)*, 17 (1996) 1564-1572.
- 510 Chen, G. and Li, H.: (Thermal behavior of proteins in hydrophobic-interaction chromatography). *Sepu*, 14 (1996) 172-175; C.A., 125 (1996) 52655m.
- 511 Clarke, S.D., Hill, H.M., Noctor, T.A.G. and Thomas, D.: Matrix-related modification of ionization in bioanalytical liquid chromatography-atmospheric pressure ionization tandem mass spectrometry. *Pharm. Sci.*, 2 (1996) 203-207; C.A., 125 (1996) 269811u.
- 512 Dale, C.J. and Lyddiatt, A.: High performance size exclusion chromatography media as versatile chromatographic supports for the analysis of process feedstocks. In: *IChemE Res. Event, Eur. Conf. Young Res. Chem. Eng.*, 2nd 1996, Vol. 1, Institution of Chemical Engineers, Rugby, 1996, pp. 280-282; C.A., 125 (1996) 56232b.
- 513 Desnoyers, S., Bourassa, S. and Poirier, G.G.: High-performance electrophoresis chromatography. *Methods Mol. Biol. (Totowa)*, 59(Protein Purification Protocols) (1996) 371-380; C.A., 125 (1996) 29449p.
- 514 Dunlap, C.J. and Carr, P.W.: The effect of mobile phase on protein retention using carboxymethyl dextran-coated zirconia stationary phases. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2059-2076.
- 515 Fernandez, M.A. and Carta, G.: Characterization of protein adsorption by composite silica-polyacrylamide gel anion exchangers. I. Equilibrium and mass transfer in agitated contactors. *J. Chromatogr. A*, 746 (1996) 169-183.
- 516 Fernandez, M.A., Laughinghouse, W.S. and Carta, G.: Characterization of protein adsorption by composite silica-polyacrylamide gel anion exchangers. II. Mass transfer in packed columns and predictability of breakthrough behavior. *J. Chromatogr. A*, 746 (1996) 185-198.
- 517 Finette, G.M.S., Mao, Q.-M. and Hearn, M.T.W.: Studies on the expansion characteristics of fluidised beds with silica-based adsorbents used in protein purification. *J. Chromatogr. A*, 743 (1996) 57-73.
- 518 Gagnon, P., Godfrey, B. and Ladd, D.: Method for obtaining unique selectivities in ion-exchange chromatography by addition of organic polymers to the mobile phase. *J. Chromatogr. A*, 743 (1996) 51-55.
- 519 Gallant, S.R., Sunnum, S. and Cramer, S.M.: Modeling gradient elution of proteins in ion-exchange chromatography. *AIChE J.*, 42 (1996) 2511-2520; C.A., 125 (1996) 189730b.
- 520 Guo, W., Shang, Z., Yu, Y. and Zhou, L.: (Preparation and application of the cartridges of dye membrane affinity chromatography). *Sepu*, 14 (1996) 168-171; C.A., 125 (1996) 52654k.
- 521 Hatano, H.: Advances in chromatography for technological analysis; chemical speciation and chemical characterization. *Methods Chromatogr.*, 1 (1996) 1-10; C.A., 125 (1996) 103811u - a review with 32 refs.
- 522 Holtzhauer, M., et al.: (Chromatography in protein research). In: Holtzhauer, M. (Editor), *Methoden Proteinanal.*, Springer, Berlin, 1996, pp. 7-45; C.A., 125 (1996) 215935u - a review with 15 refs.
- 523 Hortacsu, A. and McCoy, B.J.: Chromatographic separation of dynamically interchanging protein isomers. *Isol. Purif.*, 2 (1996) 133-148; C.A., 125 (1996) 136788v.
- 524 Hossain, M.M. and Do, D.D.: Displacement chromatography of a binary mixture of proteins - effect of protein aggregation. *Isol. Purif.*, 2 (1996) 149-164; C.A., 125 (1996) 136789w.
- 525 Hossain, M.M. and Do, D.D.: Displacement chromatography of mixtures containing a protein and an impurity. Effect of protein denaturation. *Isol. Purif.*, 2 (1996) 201-216; C.A., 125 (1996) 216023g.
- 526 Johns, M.A., Rosengarten, L.K., Jackson, M. and Regnier, F.E.: Enzyme-linked immunosorbent assays in a chromatographic format. *J. Chromatogr. A*, 743 (1996) 195-206.
- 527 Jungbauer, A. and Kaltenbrunner, O.: Fundamental questions in optimizing ion-exchange chromatography of proteins using computer-aided process design. *Biotechnol. Bioeng.*, 52 (1996) 223-236; C.A., 125 (1996) 269512x.
- 528 Kellner, R.: Fragmentation of proteins within a polyacrylamide matrix. *Biochimica*, No. 2 (1995) 19-21; C.A., 125 (1996) 109448c.
- 529 Kubota, N., Miura, S., Saito, K., Sugita, K., Watanabe, K. and Sugo, T.: Comparison of protein adsorption by anion-exchange interaction onto porous hollow-fiber membrane and gel bead-packed bed. *J. Membr. Sci.*, 117 (1996) 135-142; C.A., 125 (1996) 189883d.
- 530 Le Maire, M., Ghazi, A. and Moller, J.V.: Gel chromatography as an analytical tool for characterization of size and molecular mass of proteins. *ACS Symp. Ser.*, 635(Strategies in Size Exclusion Chromatography) (1996) 36-51; C.A., 125 (1996) 109106q - a review with 35 refs.

- 531 Levison, P.R., Mumford, C. and Streater, M.: Studies on the comparative performance of low pressure ion-exchange chromatography media for protein separation. *Spec. Publ.-R. Soc. Chem.*, 182(Ion Exchange Developments and Applications) (1996) 478-485; C.A., 125 (1996) 216003a.
- 532 Ljunglöf, A. and Hjorth, R.: Confocal microscopy as a tool for studying protein adsorption to chromatographic matrices. *J. Chromatogr. A*, 743 (1996) 75-83.
- 533 Loidl-Stahlhofen, A., Kaufmann, S., Braunschweig, T. and Bayeरl, T.M.: The thermodynamic control of protein binding to lipid bilayers for protein chromatography. *Nat. Biotechnol.*, 14 (1996) 999-1002; C.A., 125 (1996) 162440v.
- 534 Mao, Q.M. and Hearn, M.T.W.: Optimization of affinity and ion-exchange chromatographic processes for the purification of proteins. *Biotechnol. Bioeng.*, 52 (1996) 204-222; C.A., 125 (1996) 269511w.
- 535 McCoy, M., Kalghatgi, K., Regnier, F.E. and Afeyan, N.: Perfusion chromatography - characterization of column packings for chromatography of proteins. *J. Chromatogr. A*, 743 (1996) 221-229.
- 536 Mistry, S.L., Kaul, A., Merchuk, J.C. and Asenjo, J.A.: Mathematical modelling and computer simulation of aqueous two-phase continuous protein extraction. *J. Chromatogr. A*, 741 (1996) 151-163.
- 537 Mitra, G., Wong, M., Bettencourt, J., Tsay, G. and Ng, P.: Protein purification in preparative scale of mammalian cell culture-derived products: strategies for extra high purity. *Ann. N.Y. Acad. Sci.*, 782(Recombinant DNA Biotechnology III) (1996) 422-431; C.A., 125 (1996) 109258r.
- 538 Moore, A.W., Jr., Larmann, J.P., Jr., Lemmo, A.V. and Jorgenson, J.W.: Two-dimensional liquid chromatography-capillary electrophoresis techniques for analysis of proteins and peptides. *Methods Enzymol.*, 270(High Resolution Separation and Analysis of Biological Macromolecules, Pt. A) (1996) 401-419; C.A., 125 (1996) 162709q.
- 539 Nash, D.C., McCreath, G.E. and Chase, H.A.: Modification of polystyrenic matrixes for the purification of proteins. In: *IChemE Res. Event. Eur. Conf. Young Res. Chem. Eng.*, 2nd 1996, Vol. 1, Institution of Chemical Engineers, Rugby, 1996, pp. 142-144; C.A., 125 (1996) 52631a.
- 540 Noordeloos, P.: A high-performance chromatography system for developing and optimizing purification methods. *Am. Biotechnol. Lab.*, 14 (1996) 25-26; C.A., 125 (1996) 136739e.
- 541 Ogawa, T. and Hiraide, T.: Effect of pH of gradient elution of different proteins on two types of ceramic hydroxyapatite. *Am. Lab. (Shelton)*, 28 (1996) 17I, 17J, 17L; C.A., 125 (1996) 136791r.
- 542 Paliwal, S.K., de Frutos, M. and Regnier, F.E.: Rapid separations of proteins by liquid chromatography. *Methods Enzymol.*, 270(High Resolution Separation and Analysis of Biological Macromolecules, Pt. A) (1996) 133-151; C.A., 125 (1996) 215922n - a review with 31 refs.
- 543 Qian, M.G., Zhen, K., Chen, Y., Chang, C.L., Hanash, S.M. and Lubman, D.M.: Capillary liquid chromatography coupled with an ion trap storage/reflectron time-of-flight mass spectrometer for structural confirmation of three recombinant protein isoforms. *Rapid Commun. Mass Spectrom.*, 10 (1996) 1079-1087; C.A., 125 (1996) 137143t.
- 544 Ratnayake, C.K. and Regnier, F.E.: Lateral interaction between electrostatically adsorbed and covalently immobilized proteins on the surface of cation-exchange sorbents. *J. Chromatogr. A*, 743 (1996) 25-32.
- 545 Ratnayake, C.K. and Regnier, F.E.: Study of protein binding to a silica support with a polymeric cation-exchange coating. *J. Chromatogr. A*, 743 (1996) 15-23.
- 546 Ricker, R.D. and Sandoval, L.A.: Fast, reproducible size-exclusion chromatography of biological macromolecules. *J. Chromatogr. A*, 743 (1996) 43-50.
- 547 Romano, V.A., Ebeyer, T. and Dubin, P.L.: Influence of net protein charge and stationary phase charge on protein retention in size exclusion chromatography. *ACS Symp. Ser.*, 635(Strategies in Size Exclusion Chromatography) (1996) 88-102; C.A., 125 (1996) 109246k.
- 548 Santambien, P., Bengio, S. and Boschetti, E.: (Validation and characterization of chromatographic supports for protein purification). *Ann. Pharm. Fr.*, 54 (1996) 119-125; C.A., 125 (1996) 41918y - a review with 12 refs.
- 549 Schepky, A.G., Meinhardt, G., Austermann, S., Adermann, K., Schulz-Knappe, P., Forssmann, W.-G. and Hass, R.: Specific determination of tyrosine-phosphorylated proteins and peptides by differential iodination. *J. Chromatogr. A*, 743 (1996) 273-282.
- 550 Schlittler, M.R., Foy, B.A., Triska, J.J., Violand, B.N. and Bachman, G.L.: Development and optimization of SE-HPLC method for proteins using organic mobile phases. In: Marshak, D.R. (Editor), *Tech. Protein Chem. VII, [Symp. Protein Soc.]*, 9th 1995, Academic, San Diego, 1996, pp. 299-304; C.A., 125 (1996) 242054n.
- 551 Swiderek, K.M., Lee, T.D. and Shively, J.E.: Trace structural analysis of proteins. *Methods Enzymol.*, 271(High Resolution Separation and Analysis of Biological Macromolecules) (1996) 68-86; C.A., 125 (1996) 269421s - a review with 9 refs.
- 552 Taniguchi, H.: (Capillary LC/MS analysis of posttranslational and posttranscriptional modifications of proteins and nucleic acids). *J. Mass Spectrom. Soc. Jpn.*, 44 (1996) 443-457; C.A., 125 (1996) 136795v.
- 553 Wen, J., Arakawa, T., Talvenheimo, J., Welcher, A.A., Horan, T., Kita, Y., Tseng, J., Nicolson, M. and Philo, J.S.: A light scattering/size exclusion chromatography method for studying the stoichiometry of a protein-protein complex. In: Marshak, D.R. (Editor), *Tech. Protein Chem. VII, [Symp. Protein Soc.]*, 9th 1995, Academic, San Diego, 1996, pp. 23-31; C.A., 125 (1996) 242053m.
- 554 Wu, S.-L. and Karger, B.L.: Hydrophobic interaction chromatography of proteins. *Methods Enzymol.*, 270(High Resolution Separation and Analysis of Biological Macromolecules, Pt. A) (1996) 27-47; C.A., 125 (1996) 162452a.

For additional information see C.A.:
125 (1996) 109251h.

See also 5, 14, 17, 91, 105, 106, 113, 154, 181, 193, 463, 472, 478.

19b. Proteins of cells, viruses and subcellular particles

- 555 Makifuchi, S., Takeuchi, M., Kawahire, S., Odani, S., Ichimura, T., Omata, S. and Horigome, T.: Characterization of p92, karyopherin β , co-purified with N-acetylglucosamine-bearing nucleoporins from rat liver nuclear envelopes. *J. Biochem. (Tokyo)*, 120 (1996) 716-724.
- 556 McAtee, C.P., Zhang, Y., Yarbough, P.O., Fuerst, T.R., Stone, K.L., Samander, S. and Williams, K.R.: Purification and characterization of a recombinant hepatitis E protein vaccine candidate by liquid chromatography-mass spectrometry. *J. Chromatogr. B*, 685 (1996) 91-104.
- 557 Park, S.-J., Lee, Y.-M., Yoon, K.-H., Lim, K.-J. and Kwon, Y.-S.: Process for purifying hepatitis B viral surface antigen comprising preS2 peptide. *Eur. Pat. Appl. EP 716,094 (Cl. C07K14/02)*, 12 Jun. 1996, KR Appl. 9,433,594, 10 Dec. 1994; 13 pp.; C.A., 125 (1996) 84648c.
- 558 Schlaif, G., Goeddecke, M., Wolff, J.R., Felgenhauer, K. and Maeder, M.: Large-scale purification of synaptophysin and quantification with a newly established enzyme-linked immunosorbent assay. *Biol. Chem.*, 377 (1996) 591-597; C.A., 125 (1996) 269708r.
- 559 Takada, K., Hibi, N., Tsukada, Y., Shibasaki, T. and Ohkawa, K.: Ability of ubiquitin radioimmunoassay to discriminate between monoubiquitin and multi-ubiquitin chains. *Biochim. Biophys. Acta*, 1290 (1996) 282-288.
- 560 Wolff, P., Tanaka, A.M., Chenker, E., Cabrera-Crespo, J., Raw, I. and Ho, P.L.: Purification of fibroblast growth factor-2 from human placenta using tri(n-butyl)phosphate and sodium cholate. *Biochimie*, 78 (1996) 190-194; C.A., 125 (1996) 186018q.

See also 500, 561, 695.

19c. Proteins synthesized by genetic manipulation, monoclonal antibodies

- 561 Flaminio, G., Monti, E., Gargiulo, F., Varinacci, C., Magni, P., Martinelli, F., Caimi, L., Caruso, A., Preti, A. and Turano, A.: Purification of recombinant LamB proteins using continuous elution electrophoresis: a comparison with immunoaffinity chromatography. *Biochem. Mol. Biol. Int.*, 36 (1995) 1255-1261; C.A., 125 (1996) 52804j.
- 562 Hentz, N.G., Vukasinovic, V., Diez, S., Valiente, M. and Daunert, S.: Affinity chromatography for recombinant proteins. In: Butterfield, D.A. (Editor), *Biofunct. Membr. [Proc. Int. Conf.]*, 1995, Plenum, New York, 1996, pp. 73-81; C.A., 125 (1996) 136644v - a review with 57 refs.
- 563 Li, S., Patapoff, T.W., Overcashier, D., Hsu, C., Nguyen, T.H. and Borchardt, R.T.: Effects of reducing sugars on the chemical stability of human relaxin in the lyophilized state. *J. Pharm. Sci.*, 85 (1996) 873-877.
- 564 Miller, K.J. and Herman, A.C.: Affinity chromatography with immunochemical detection applied to the analysis of human methionyl granulocyte colony stimulating factor in serum. *Anal. Chem.*, 68 (1996) 3077-3082.
- 565 Olson, W.P.: Chromatographic purification of recombinant protein products. In: Olson, W.P. (Editor), *Sep. Technol.*, Interpharm Press, Buffalo Grove, 1995, pp. 271-301; C.A., 125 (1996) 177092g - a review with many refs.

- 566 Sabater, M., Heilmann, S., Froekjaer, H., Biedermann, K. and Emborg, C.: The use of antibodies for characterization and quantification of a recombinant protein. *Ann. N.Y. Acad. Sci.*, 782(Recombinant DNA Biotechnology III) (1996) 462-477; C.A., 125 (1996) 109494q.
- 567 Tebbe, H., Buentemeyer, H., Luetkemeyer, D. and Lehmann, J.: Rapid HPLC quantification of recombinant human antithrombin III during production and purification. In: Beauverie, E.C., Griffiths, J.B. and Zeijlemaker, W.P. (Editors), *Anim. Cell Technol.: Dev. 21st Century, [Proc. Meet.] 1994*, Kluwer, Dordrecht, 1995, pp. 455-459; C.A., 125 (1996) 31971w.
- 568 Thompson, S.A., Harris, A., Hoang, D., Ferrer, M. and Johnson, G.R.: COOH-Terminal extended recombinant amphiregulin with bioactivity comparable with naturally derived growth factor. *J. Biol. Chem.*, 271 (1996) 17927-17931.
- 569 Wang, Y., Liu, Q., Hua, B., Zhu, Y., Wang, Y. and Xiu, J.: (Purification of bacterially expressed human anti-HBs Fab using metal chelated chromatography). *Zhonghua Weishengwuxue He Manyixue Zazhi*, 16 (1996) 104-107; C.A., 125 (1996) 165239x.

See also 483, 590, 627, 638.

19d. Microbial and plant proteins

- 570 Enguita, F.J., de la Fuente, J.L., Martin, J.F. and Liras, P.: An inducible expression system of histidine-tagged proteins in *Streptomyces lividans* for one-step purification by Ni²⁺ affinity chromatography. *FEMS Microbiol. Lett.*, 137 (1996) 135-140; C.A., 125 (1996) 77602e.
- 571 Hinode, D., Grenier, D. and Mayrand, D.: A general procedure for the isolation of heat-shock proteins from periodontopathogenic bacteria. *J. Microbiol. Methods*, 25 (1996) 349-355; C.A., 125 (1996) 189903k.
- 572 Krawczyk, B. and Kur, J.: Purification of IHF-like protein from gram-negative bacteria in one chromatographic step. *Acta Biochim. Pol.*, 43 (1996) 379-382; C.A., 125 (1996) 269501t.
- 573 Liang, X., Zheng, K., Qian, M.G. and Lubman, D.M.: Determination of bacterial protein profiles by matrix-assisted laser desorption/ionization mass spectrometry with high-performance liquid chromatography. *Rapid Commun. Mass Spectrom.*, 10 (1996) 1219-1226; C.A., 125 (1996) 162689h.
- 574 Quandroni, M., Staudemann, W., Kertesz, M. and James, P.: Analysis of global responses by protein and peptide fingerprinting of proteins isolated by two-dimensional gel electrophoresis. Application to the sulfate-starving response of *Escherichia coli*. *Eur. J. Biochem.*, 239 (1996) 773-781.

See also 298, 1151, 1152, 1155.

19e. Proteins of blood, serum and blood cells

- 575 Allanson, J.P., Biddlecombe, R.A., Jones, A.E. and Pleasance, S.: The use of automated solid phase extraction in the '96 well' format for high throughput bioanalysis using liquid chromatography coupled to tandem mass spectrometry. *Rapid Commun. Mass Spectrom.*, 10 (1996) 811-816; C.A., 125 (1996) 67958m.

- 576 Björklund, M. and Hearn, T.W.: Characterisation of silica-based heparin-affinity adsorbents through column chromatography of plasma fractions containing thrombin. *J. Chromatogr. A*, 743 (1996) 145-162.
- 577 Blizzard, C.D., Garramone, S., Gordon, S. and Gammell, M.: A method for rapid purification of antibody conjugates. *Am. Biotechnol. Lab.*, 14 (1996) 49-50; *C.A.*, 125 (1996) 83937j.
- 578 Brunati, A.M., Bordin, L., Clari, G. and Moret, V.: The Lyn-catalyzed Tyr phosphorylation of the transmembrane band-3 protein of human erythrocytes. *Eur. J. Biochem.*, 240 (1995) 394-399.
- 579 Chu, C. and Sun, X.: (Purification of human red blood cell group-A protein by affinity chromatography). *Zhongguo Mianyixue Zazhi*, 12 (1996) 31-33; *C.A.*, 125 (1996) 192966p.
- 580 Cuny, C., Pham, L., Kramp, W., Sharp, T. and Soriano, T.F.: Evaluation of a two-site immunoradiometric assay for measuring noncomplexed (free) prostate-specific antigen. *Clin. Chem. (Washington)*, 42 (1996) 1243-1249.
- 581 Dalençon, F., Rosilio, V., Puisieux, F., Baszkin, A. and Wautier, J.-L.: Liposomes bearing platelet proteins: a model for surface functions studies. *Biochim. Biophys. Acta*, 1302 (1996) 241-248.
- 582 Hikawa, A., Nomata, Y., Suzuki, T., Ozasa, H. and Yamada, O.: Soluble transferrin receptor-transferrin complex in serum: measurement by latex agglutination nephelometric immunoassay. *Clin. Chim. Acta*, 254 (1996) 159-172.
- 583 Kunimatsu, M., Sasaki, M., Takagi, O. and Yamamoto, N.: (Purification of IgY antibody). *Jpn. Kokai Tokkyo Koho JP 08,188,599 [96,188,599]* (Cl. C07K16/02), 23 Jul. 1996, Appl. 95/16,388, 9 Jan. 1995; 9 pp.; *C.A.*, 125 (1996) 165704v.
- 584 Mashiko, H., Kato, K., Fujii, K.-i. and Takahashi, H.: Studies on factor XII in porcine plasma: purification and its conversion to activated form by porcine plasma kallikrein. *Biochim. Biophys. Acta*, 1296 (1996) 198-206.
- 585 McCarthy, E., Vella, G., Mhatre, R. and Lim, Y.-P.: Rapid purification and monitoring of immunoglobulin M from ascites by perfusion ion-exchange chromatography. *J. Chromatogr. A*, 743 (1996) 163-170.
- 586 Nishikawa, T., Akiyoshi, K. and Sunamoto, J.: Macromolecular complexation between bovine serum albumin and the self-assembled hydrogel nanoparticle of hydrophobized polysaccharides. *J. Am. Chem. Soc.*, 118 (1996) 6110-6115; *C.A.*, 125 (1996) 80892e.
- 587 Shan, J.: HPLC with online enzymic detection: I. determination of α -fetoprotein in serum: II. separation of aspartate aminotransferase isoenzymes and differentiation of liver and heart cytoplasmic aspartate aminotransferase. *Avail. Univ. Microfilms Int.*, Order No. DA9623958, 1996, 176 pp.; *C.A.*, 125 (1996) 136816c.
- 588 Tang, Z.-H., Luo, L.-H., Yu, G.-H., Shi, F.-X., Ding, S.-B. and Xu, X.-Y.: (Renaturation of recombinant interleukin-2 during purification by ion exchange chromatography). *Shengwu Huaxue Zazhi*, 12 (1996) 341-344; *C.A.*, 125 (1996) 165232q.
- 589 Tharakan, J. and Belizaire, M.: Comparison of axial and radial flow columns for immunoaffinity chromatography of factor IX. *Isol. Purif.*, 2 (1996) 103-112; *C.A.*, 125 (1996) 140579v.
- 590 Tyutyulkova, S., Gao, Q.-S., Thompson, A., Rennard, S. and Paul, S.: Efficient vasoactive intestinal polypeptide hydrolyzing autoantibody light chains selected by phage display. *Biochim. Biophys. Acta*, 1316 (1996) 217-223.
- 591 Uchida, T., Kozuma, T., Yasukawa, K. and Shima, K.: Glycated albumin analyzer. *Methods Chromatogr.*, 1(Advances in Liquid Chromatography) (1996) 33-41; *C.A.*, 125 (1996) 136751c.
- 592 Vazquez-Moreno, L. and Plata, M. del Carmen Candia: Scale up isolation of immunoglobulins from pig serum by immobilized metal affinity chromatography. *J. Food Biochem.*, 19 (1996) 367-380; *C.A.*, 125 (1996) 80909r.
- 593 Yang, Y.-B. and Harrison, K.: Influence of column type and chromatographic conditions on the ion-exchange chromatography of immunoglobulins. *J. Chromatogr. A*, 743 (1996) 171-180.
- For additional information see *C.A.*:
125 (1996) 81289a, 189974j, 193477s, 219620k.
- See also 156, 634.
- 19f. Structural and muscle proteins**
- 594 Salih, E., Zhou, H. and Glimcher, M.J.: Phosphorylation of purified bovine bone sialoprotein and osteopontin by protein kinases. *J. Biol. Chem.*, 271 (1996) 16897-16905.
- 595 Trout, G.R. and Gutzke, D.A.: A simple, rapid preparative method for isolating and purifying oxymyoglobin. *Meat Sci.*, 43 (1996) 1-13; *C.A.*, 125 (1996) 137164a.
- 596 Zhou, H., Huiatt, T.W., Robson, R.M., Sernett, S.W. and Graves, D.J.: Characterization of ADP-ribosylation sites on desmin and restoration of desmin intermediate filament assembly by de-ADP-ribosylation. *Arch. Biochem. Biophys.*, 334 (1996) 214-222.
- For additional information see *C.A.*:
125 (1996) 29350z, 105738z.
- See also 635, 733.
- 19g. Protamines, histones and other nuclear proteins**
- 597 Ballestar, E., Abad, C. and Franco, L.: Core histones are glutaminyl substrates for tissue transglutaminase. *J. Biol. Chem.*, 271 (1996) 18817-18824.
- 598 Diaz-Jullien, C., Pérez-Estévez, A., Covelo, G. and Freire, M.: Prothymosin α binds histones *in vitro* and shows activity in nucleosome assembly assay. *Biochim. Biophys. Acta*, 1296 (1996) 219-227.
- 599 Lindner, H., Sarg, B., Meraner, C. and Helliger, W.: Separation of acetylated core histones by hydrophilic-interaction liquid chromatography. *J. Chromatogr. A*, 743 (1996) 137-144.
- See also 555.
- 19h. Chromoproteins and metalloproteins**
- 600 Ait-Oukhatar, N., Bureau, F., Boudey, M., Place, C., Jauzac, P., Drosdowsky, M.A., Arhan, P. and Bougle, D.: Determination of metallothioneins by HPLC: application to zinc metabolism. *Ann. Biol. Clin.*, 54 (1996) 87-90; *C.A.*, 125 (1996) 109244h.
- 601 Beattie, J.H., Lomax, J.A., Richards, M.P., Self, R., Pesch, R. and Muenster, H.: Metallothionein isoform analysis by liquid chromatography-electrospray ionization mass spectrometry. *Biochim. Soc. Trans.*, 24 (1996) 220S; *C.A.*, 125 (1996) 81160b.

- 602 Bon, C.H., Revenant, M.C., Sotta, C., Mailliavin, A., Bannier, E., Goujon, R., Lestra, C. and Garcia, I.: (Multicenter evaluation of the Abbott IMx glycated hemoglobin assay). *Ann. Biol. Clin.*, 54 (1996) 151-157; *C.A.*, 125 (1996) 136814a.
- 603 Bonifacio, M.D., Steeves, T., Saunders, D.M. and Sinosich, M.J.: Isolation of ER⁷² from guinea pig term placentas using heparin-Sepharose affinity chromatography. *Biochem. Mol. Biol. Int.*, 36 (1995) 1143-1152; *C.A.*, 125 (1996) 80888h.
- 604 De Tata, V., Novelli, M., Bombara, M., Masiello, P., Cappuccio, A., Innocenti, B. and Bergamini, E.: Determination of glycated hemoglobins in the rat: Comparison between two different chromatographic methods and application in experimental diabetes. *Res. Exp. Med.*, 196 (1996) 9-16; *C.A.*, 125 (1996) 216019k.
- 605 Dou, H., Muneo, M. and Takehiko, Y.: (High-performance immobilized metal ion affinity chromatographic separation of four types of molecules of transferrin). *Sepu*, 14 (1996) 176-178; *C.A.*, 125 (1996) 80905m.
- 606 Espinoza, E.O., Kirms, M.A. and Filipek, M.S.: Identification and quantitation of source from hemoglobin of blood and blood mixtures by high performance liquid chromatography. *J. Forensic Sci.*, 41 (1996) 804-811; *C.A.*, 125 (1996) 214384b.
- 607 Kopp, H.P., Festa, A., Hopmeier, P. and Scherthaner, G.: (Evaluation of a new method for measurement of glycated hemoglobin with monoclonal antibodies). *Wien. Klin. Wochenschr.*, 108 (1996) 16-19; *C.A.*, 125 (1996) 81133v.
- 608 Koskinen, L.K.: Specificity of hemoglobin A_{1c} measurement by cation exchange liquid chromatography. Evaluation of a Mono S column method. *Alin. Chim. Acta*, 253 (1996) 159-169.
- 609 Kroes, S.J., Hoitink, C.W.G., Andrew, C.R., Ai, J., Sanders-Loehr, J., Messerschmidt, A., Hagen, W.R. and Canters, G.W.: The mutation Met121→His creates a type-1.5 copper site in *Alcaligenes denitrificans* azurin. *Eur. J. Biochem.*, 240 (1996) 342-351.
- 610 Matsubara, K., Yamaki, M., Nagayama, K., Imai, K., Ishii, H., Gotoh, T. and Ebina, S.: Wheat germ agglutinin-reactive chains of giant hemoglobin from the polychaete *Perinereis aibuhensis*. *Biochim. Biophys. Acta*, 1290 (1996) 215-223.
- 611 Osipov, A.N., Gorbunov, N.V., Day, B.W., Elsayed, N.M. and Kagan, V.E.: Electron spin resonance and mass spectral analysis of interactions of ferrylhemoglobin and ferrylmyoglobin with nitric oxide. *Methods Enzymol.*, 268(Nitric Oxide, Part A) (1996) 193-203; *C.A.*, 125 (1996) 269597d.
- 612 Plura, D.H., Wiffen, D.E., Ashraf, S. and Magnin, A.A.: Purification of hemoglobin by displacement chromatography. *U.S. US 5,545,328* (Cl. 210-635; B01D15/08), 13 Aug. 1996, CA Appl. 2,106,612, 21 Sep. 1993; 13 pp.; *C.A.*, 125 (1996) 162746z.
- 613 Quesada, A.R., Byrnes, R.W., Krezoski, S.O. and Petering, D.H.: Direct reaction of H₂O₂ with sulphydryl groups in HL-60 cells: zinc-metallothionein and other sites. *Arch. Biochem. Biophys.*, 334 (1996) 241-250.
- 614 Shinogi, M., Nishinaga, K. and Yokoyama, I.: Measurement of metallothionein in rat tissues by a chromatographic method using the fluorometry of thiol groups. *Biol. Pharm. Bull.*, 19 (1996) 911-914; *C.A.*, 125 (1996) 136804x.
- 615 Simonsson, P., Lindberg, S. and Alling, C.: Carbohydrate-deficient transferrin measured by high-performance liquid chromatography and CDTeC immunoassay. *Alcohol Alcohol.*, 31 (1996) 397-402; *C.A.*, 125 (1996) 269509b.
- 616 Tomita, S., Okuyama, E., Ohnishi, T. and Ichikawa, Y.: Characteristic properties of a retinoic acid synthetic cytochrome P-450 purified from liver microsomes of 3-methylcholanthrene-induced rats. *Biochim. Biophys. Acta*, 1290 (1996) 273-281.
- 617 Yuasa, H.J., Green, B.N., Takagi, T., Suzuki, N., Vinogradov, S.N. and Suzuki, T.: Electrospray ionization mass spectrometric composition of the 400 kDa hemoglobin from the pogonophoran *Oligobrachia mashikoi* and the primary structures of three major globin chains. *Biochim. Biophys. Acta*, 1296 (1996) 235-244.
- For additional information see *C.A.*:
125 (1996) 137209u.
- See also 498, 682.
- 19i. Proteins of glands, gland products, various zymogens (incl. milk proteins)
- 618 Calvete, J.J., Varela, P.F., Sanz, L., Romero, A., Mann, K. and Toepfer-Petersen, E.: A procedure for the large-scale isolation of major bovine seminal plasma proteins. *Protein Expression Purif.*, 8 (1996) 48-56; *C.A.*, 125 (1996) 136820z.
- 619 Hamako, J., Matsui, T., Suzuki, M., Ito, M., Makita, K., Fujimura, Y., Ozeki, Y. and Titani, K.: Purification and characterization of bitiscetin, a novel von Willebrand factor modulator protein from *Bitis arietans* snake venom. *Biochem. Biophys. Res. Commun.*, 226 (1996) 273-279.
- 620 Hirashima, A., Athauda, S.B.P. and Takahashi, K.: Purification and characterization of turtle pepsinogen and pepsin. *J. Biochem. (Tokyo)*, 120 (1996) 407-414.
- 621 Minkiewicz, P., Slanger, C.J., Lagerwerf, F.M., Haverkamp, J., Rollema, H.S. and Visser, S.: Reversed-phase high-performance liquid chromatographic separation of bovine κ-casein macropeptides and characterization of isolated fractions. *J. Chromatogr. A*, 743 (1996) 123-135.
- 622 Mourad, M., Bascul, A., Delmas, H., Wilhelm, A.M. and Gros, B.: (Recovery of whey proteins in a fluidized bed: effect of solid-phase mixing). *Recents Prog. Genie Procedes*, 9(42, Genie des Procedes Complexes) (1995) 483-488; *C.A.*, 125 (1996) 84961z.
- For additional information see *C.A.*:
125 (1996) 29351a.
- See also 501, 596, 631, 1173.
- 19j. Proteins of brain, cerebrospinal fluid and eye
- 623 Urden, G., Thoernwall, M., Lyrenaes, S., Lindstroem, L. and Nyberg, F.: Classification of CSF samples from normal and post-partum psychotic women using chromatographic profiles with bilinear projections: a multivariate approach. *Biomed. Chromatogr.*, 10 (1996) 149-154; *C.A.*, 125 (1996) 109269v.
- See also 730.

19k. *Proteins of neoplastic tissue and transformed cells*

- 624 Hagiya, Y., Fuako, H., Ueshima, S., Okada, K., Inufusa, H., Kamiishi, H. and Matsuo, O.: Enhanced urokinase-type plasminogen activator activity by extracellular matrix protein obtained from highly metastatic human lung adenocarcinoma cell line. *Clin. Chim. Acta.*, 253 (1996) 37-50.
- 625 Zusman, I., Zusman, R., Sandler, B., Korol, D., Gurevich, P., Smirnoff, P., Bass, D., Or, A., Shani, A., et al.: HPLC determination of serum levels of soluble p53 antigen as a new method for colon cancer detection, and its clinical implication. *Int. J. Oncol.*, 8 (1996) 1249-1256; C.A., 125 (1996) 54997y.

See also 158, 308, 613.

19l. *Specific binding and receptor proteins*

- 626 Bar-Peled, M. and Raikhel, N.V.: A method for isolation and purification of specific antibodies to a protein fused to the GST. *Anal. Biochem.*, 241 (1996) 140-142.
- 627 Ettner, N., Müller, G., Berens, C., Backes, H., Schnappinger, D., Schreppel, T., Pfeiderer, K. and Hillen, W.: Fast large-scale purification of tetracycline repressor variants from overproducing *Escherichia coli* strains. *J. Chromatogr. A*, 742 (1996) 95-105.
- 628 Houen, G., Högdall, E.V., Barkholt, V. and Norskov, L.: Lactoferrin: similarity to diamine oxidase and purification by aminoxyhexyl affinity chromatography. *Eur. J. Biochem.*, 241 (1996) 303-308.
- 629 Ibarrola, I., Andrés, M., Marino, A., Macarulla, J.M. and Trueba, M.: Purification of a cortisol binding protein from hepatic plasma membrane. *Biochim. Biophys. Acta*, 1284 (1996) 41-46.
- 630 Kihlberg, B.-M., Sjöholm, A.G., Björck, L. and Sjöbring, U.: Characterization of the binding properties of protein LG, an immunoglobulin-binding hybrid protein. *Eur. J. Biochem.*, 240 (1996) 556-563.
- 631 Majumdar, K.C., Shetty, S., Wadhwa, R., Bhaskar, S., Ganesam, M. and Singh, L.: Detection and purification of sequence-specific DNA binding protein. *Anal. Biochem.*, 241 (1996) 23-29.
- 632 Mazoy, R., Vazquez, F. and Lemos, M.L.: Isolation of heme-binding proteins from *Vibrio anguillarum* using affinity chromatography. *FEMS Microbiol. Lett.*, 141 (1996) 19-23; C.A., 125 (1996) 108141k.
- 633 Meyer, C., Schmid, R., Scriba, P.C. and Wehling, M.: Purification and partial sequencing of high-affinity progesterone-binding site(s) from porcine liver membranes. *Eur. J. Biochem.*, 239 (1996) 726-731.
- 634 Nakano, Y., Tobe, T., Choi-Miura, N.-H., Mazda, T. and Tomita, M.: Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma. *J. Biochem. (Tokyo)*, 120 (1996) 803-812.
- 635 Remppis, A., Greten, T., Schäfer, B.W., Hunziker, P., Erne, P., Katus, H.A. and Heizmann, C.W.: Altered expression of the Ca^{2+} -binding protein S100A1 in human cardiomyopathy. *Biochim. Biophys. Acta*, 1313 (1996) 253-257.
- 636 Roeth, P.J. and Easterbrook-Smith, S.B.: C1q is a nucleotide binding protein and is responsible for the ability of clusterin preparations to promote immune complex formation. *Biochim. Biophys. Acta*, 1297 (1996) 159-166.

- 637 Tucker, J. and Grisshammer, R.: Purification of a rat neurotsin receptor expressed in *Escherichia coli*. *Biochem. J.*, 317 (1996) 891-899.

See also 733.

19m. *Urinary proteins*

- 638 Okutani, R., Itoh, Y., Yamada, T., Yamaguchi, T., Singh, G., Yagisawa, H. and Kawai, T.: Preparation and characterization of human recombinant protein 1/Clara cell M_r 10000 protein. *Eur. J. Clin. Chem. Clin. Biochem.*, 34 (1996) 691-696.
- 639 Astafieva, I.V., Eberlein, G.A. and Wang, Y.J.: Absolute on-line molecular mass analysis of basic fibroblast growth factor and its multimers by reversed-phase liquid chromatography with multi-angle laser light scattering detection. *J. Chromatogr. A*, 740 (1996) 215-229.
- 640 Awade, A.C.: On hen egg fractionation. Applications of liquid chromatography to the isolation and the purification of hen egg white and egg yolk proteins. *Z. Lebensm.-Unters. Forsch.*, 202 (1996) 1-14; C.A., 125 (1996) 56459y - a review with 108 refs.
- 641 Bayer, E.A., Ehrlich-Rogozinski, S. and Wilchek, M.: Sodium dodecyl sulfate-polyacrylamide gel electrophoretic method for assessing the quaternary state and comparative thermostability of avidin and streptavidin. *Electrophoresis (Weinheim)*, 17 (1996) 1319-1324.
- 642 Hattori, A., Kuroki, Y., Sohma, H., Ogasawara, Y. and Akino, T.: Human surfactant protein A with two distinct oligomeric structures which exhibit different capacities to interact with alveolar type II cells. *Biochem. J.*, 317 (1996) 939-944.
- 643 Katoh, T., Kikuchi, N., Nagata, K. and Yoshida, N.: A mutant trypsin-like enzyme from *Streptomyces fradiae*, created by site-directed mutagenesis, improves affinity chromatography for protein trypsin inhibitors. *Appl. Microbiol. Biotechnol.*, 46 (1996) 15-21; C.A., 125 (1996) 215500y.
- 644 Khan, M.A. and Addo-Yobo, F.: Modeling ion exchange chromatography of egg white proteins for process control purposes. *IChemE Res. Event. Eur. Conf. Young Res. Chem. Eng.*, 2nd 1996, Vol. 1, Institution of Chemical Engineers, Rugby, 1996, pp. 133-135; C.A., 125 (1996) 32168b.

For additional information see C.A.:
125 (1996) 189973h.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

- 645 Bryjak, M. and Trochimczuk, A.: New chromatographic materials for enzyme separation. *Environ. Prot. Eng.*, 20 (1994, Pub. 1995) 43-51; C.A., 125 (1996) 80124f.
- 646 Jansen, E.H.J.M., Laan, C.A. and de Fluiter, P.: Advances in sample preparation, electrophoretic separation and detection methods for rat cytochrome P450 enzymes. *J. Chromatogr. B*, 684 (1996) 133-145 - a review with 50 refs.

- 647 Roos, P.H.: Chromatographic separation and behavior of microsomal cytochrome P450 and cytochrome b₅. *J. Chromatogr.*, 684 (1996) 107-131 - a review with 154 refs.

See also 706.

20a. Oxidoreductases

- 648 Apanovitch, D. and Walz, F.G., Jr.: Adult-male-specific S-warfarin (11S-OH) and progesterone (20 β -OH) keto-reductases in rat hepatic microsomes are not identical. *Biochim. Biophys. Acta*, 1291 (1996) 16-26.
- 649 Bauer, I., Max, N., Fetzner, S. and Lingens, F.: 2,4-Dioxygenases catalyzing N-terocyclic-ring cleavage and formation of carbon monoxide. Purification and some properties of 1H-3-hydroxy-4-oxoquininaline 2,4-dioxygenase from *Arthrobacter* sp. Rü61a and comparison with 1H-3-hydroxy-4-oxoquinoline 2,4-dioxygenase from *Pseudomonas putida* 33/1. *Eur. J. Biochem.*, 240 (1996) 576-583.
- 650 Bayer, M., Walter, K. and Simon, H.: Purification and partial characterization of a reversible artificial mediator accepting NADH oxidoreductase from *Clostridium thermoaceticum*. *Eur. J. Biochem.*, 239 (1996) 686-691.
- 651 Braun, H., Lichter, A. and Häberlein, I.: Kinetic evidence for protein complexes between thioredoxin and NADP-malate dehydrogenase and presence of a thioredoxin binding site at the N-terminus of the enzyme. *Eur. J. Biochem.*, 240 (1996) 781-788.
- 652 Brinckmann, R., Topp, M.S., Zalán, I., Heydeck, D., Ludwig, P., Kühn, H., Berdel, W.E. and Habenicht, A.J.R.: Regulation of 15-lipoxygenase expression in lung epithelial cells by interleukin-4. *Biochem. J.*, 318 (1996) 305-312.
- 653 Bullock, J., Chowdhury, S. and Johnston, D.: Characterization of poly(ethylene glycol)-modified superoxide dismutase: comparison of capillary electrophoresis and matrix-assisted laser desorption/ionization mass spectrometry. *Anal. Chem.*, 68 (1996) 3258-3264.
- 654 Converso, D.A. and Fernández, M.E.: Ca²⁺ Activation of wheat peroxidase: a possible physiological mechanism of control. *Arch. Biochem. Biophys.*, 333 (1996) 59-65.
- 655 During, A., Nagao, A., Hoshino, C. and Terao, J.: Assay of β -carotene 15,15'-dioxygenase activity by reverse-phase high-performance liquid chromatography. *Anal. Biochem.*, 241 (1996) 199-205.
- 656 Gleason, F.K.: Glucose-6-phosphate dehydrogenase from the cyanobacterium, *Anabaena* sp. PCC 7120; purification and kinetics of redox modulation. *Arch. Biochem. Biophys.*, 334 (1996) 277-283.
- 657 Grayson, D.A., Tewari, Y.B., Mayhew, M.P., Vilker, V.L. and Goldberg, R.N.: Tetralin as a substrate for camphor (cytochrome P450) 5-monooxygenase. *Arch. Biochem. Biophys.*, 332 (1996) 239-247.
- 658 Grün, F., Noy, N., Hämmерling, U. and Buck, J.: Purification, cloning, and bacterial expression of retinol dehydratase from *Spodoptera frugiperda*. *J. Biol. Chem.*, 271 (1996) 16135-16138.
- 659 Haining, R.L., Hunter, A.P., Veronese, M.E., Trager, W.F. and Rettie, A.E.: Allelic variants of human cytochrome P450 2C9: baculovirus-mediated expression, purification, structural characterization, substrate stereoselectivity, and prochiral selectivity on the wild type and I359L mutant forms. *Arch. Biochem. Biophys.*, 333 (1996) 447-458.
- 660 Ishida, T., Narita, M., Nozaki, M. and Horiike, K.: Selective cleavage and modification of the intersubunit disulfide bonds of bovine dopamine β -monooxygenase: conversion of tetramer to active dimer. *J. Biochem. (Tokyo)*, 120 (1996) 346-352.
- 661 Jensen, W.A., Armstrong, J.M.D., de Giorgio, J. and Hearn, M.T.W.: Stability studies on pig heart mitochondrial malate dehydrogenase: the effect of salts and amino acids. *Biochim. Biophys. Acta*, 1296 (1996) 23-34.
- 662 Kim, E.-J., Kim, H.-P., Hah, Y.C. and Roe, J.-H.: Differential expression of superoxide dismutases containing Ni and Fe/Zn in *Streptomyces coelicolor*. *Eur. J. Biochem.*, 241 (1996) 178-185.
- 663 Kopperschläger, G. and Kirchberger, J.: Methods for the separation of lactate dehydrogenases and clinical significance of the enzyme. *J. Chromatogr. B*, 684 (1996) 25-49 - a review with 154 refs.
- 664 Krause, R.J., Ripp, S.L., Sausen, P.J., Overby, L.H., Philpot, R.M. and Elfarra, A.A.: Characterization of the methionine S-oxidase activity of rat liver and kidney microsomes: immunochemical and kinetic evidence for FMO3 being the major catalyst. *Arch. Biochem. Biophys.*, 333 (1996) 109-116.
- 665 Lin, S.-X., Xie, B., Li, B. and Zhou, M.: Chromatography of hydroxysteroid dehydrogenases. *J. Chromatogr. B*, 684 (1996) 99-105 - a review with 35 refs.
- 666 McManaman, J.L., Shellman, V., Wright, R.M. and Repine, J.E.: Purification of rat liver xanthine oxidase and xanthine dehydrogenase by affinity chromatography on benzamidine-Sepharose. *Arch. Biochem. Biophys.*, 332 (1996) 135-141.
- 667 Mendoza-Hernández, G. and Rendón, J.L.: Human placental estradiol 17 β -dehydrogenase: structural and catalytic changes during urea denaturation. *Biochim. Biophys. Acta*, 1297 (1996) 219-227.
- 668 Michalski, W.P.: Chromatographic and electrophoretic methods for analysis of superoxide dismutases. *J. Chromatogr. B*, 684 (1996) 59-75 - a review with 159 refs.
- 669 O'Carra, P., Griffin, T., O'Flaherty, M., Kelly, N. and Mulcahy, P.: Further studies on the bioaffinity chromatography of NAD⁺-dependent dehydrogenases using the locking-on effect. *Biochim. Biophys. Acta*, 1297 (1996) 235-243.
- 670 O'Shannessy, K., Scoble, J. and Scopes, R.K.: A simple and economical procedure for purification of muscle lactate dehydrogenase by batch dye-ligand adsorption. *Bioseparation*, 6 (1996) 77-80; *C.A.*, 125 (1996) 189082s.
- 671 Périé, F.H., Sheng, D. and Gold, M.H.: Purification and characterization of two manganese peroxidase isozymes from the white-rot basidiomycete *Dichomitus squalens*. *Biochim. Biophys. Acta*, 1297 (1996) 139-148.
- 672 Pikuleva, I.A., Björkhem, I. and Waterman, M.R.: Studies of distant members of the P450 superfamily (P450scC and P450c27) by random chimeragenesis. *Arch. Biochem. Biophys.*, 334 (1996) 183-192.

- 673 Ren, S., Liu, H., Licad, E. and Correia, M.A.: Expression of rat liver tryptophan 2,3-dioxygenase in *Escherichia coli*: structural and functional characterization of the purified enzyme. *Arch. Biochem. Biophys.*, 333 (1996) 96-102.
- 674 Schaefer, A., Bieg, S., Huwig, A., Kohring, G.-W. and Giffhorn, F.: Purification by immunoaffinity chromatography, characterization, and structural analysis of a thermostable pyranose oxidase from the white rot fungus *Phlebiopsis gigantea*. *Appl. Environ. Microbiol.*, 62 (1996) 2586-2592; C.A., 125 (1996) 52097f.
- 675 Scheibe, R., Baumann, E., Backhausen, J.E., Rak, C. and Vetter, S.: C-Terminal truncation of spinach chloroplast NAD(P)⁺-dependent glyceraldehyde-3-phosphate dehydrogenase prevents inactivation and reaggregation. *Biochim. Biophys. Acta*, 1296 (1996) 228-234.
- 676 Schmitt, U., Jahnke, K., Rosenbaum, K., Cook, P.F. and Schnackerz, K.D.: Purification and characterization of dihydropyrimidine dehydrogenase from *Alcaligenes entrophus*. *Arch. Biochem. Biophys.*, 332 (1996) 175-182.
- 677 Shao, T., Liu, Y. and Shao, C.: (Purification of pig copper,zinc-peroxide dismutase by immobilized metal ion affinity chromatography). *Sepu*, 14 (1996) 218-221; C.A., 125 (1996) 52061q.
- 678 Toribio, F., Alhamá, J. and López-Barea, J.: Methods for chromatographic and electrophoretic separation and assay of NADP⁺ oxidoreductases. *J. Chromatogr. B*, 684 (1996) 1-23 - a review with 214 refs.
- 679 Toribio, F., Martínez-Lara, E., Pascual, P. and López-Barea, J.: Methods for purification of glutathione peroxidase and related enzymes. *J. Chromatogr. B*, 684 (1996) 77-97 - a review with 227 refs.
- 680 Tsui, H.-T., Mock, W.-Y., Lau, K.-K. and Fong, W.-P.: Proteolytic activation of grass carp (*Ctenopharyngodon idellus*) liver alcohol dehydrogenase. *Biochim. Biophys. Acta*, 1296 (1996) 41-46.
- 681 Youn, H.-D., Youn, H., Lee, J.-W., Yim, Y.-I., Lee, J.K., Hah, Y.C. and Kang, S.-O.: Unique isozymes of superoxide dismutase in *Streptomyces griseus*. *Arch. Biochem. Biophys.*, 334 (1996) 341-348.
- 682 Yousafzai, F.K., Buck, M. and Smith, B.E.: Isolation and characterization of nitrogenase MoFe protein from the mutant strain pHK17 of *Klebsiella pneumoniae* in which two bridging cysteine residues of the P-clusters are replaced by the non-coordinating amino acid alanine. *Biochem. J.*, 318 (1996) 111-118.
- 683 Zenno, S., Koike, H., Tanokura, M. and Saigo, K.: Gene cloning, purification, and characterization of NfsB, a minor oxygen-insensitive nitroreductase from *Escherichia coli*, similar in biochemical properties to FRase I, the major flavin reductase in *Vibrio fischeri*. *J. Biochem. (Tokyo)*, 120 (1996) 736-744.
- For additional information see C.A.:
125 (1996) 136093q.
- See also 494, 749, 769, 841.
- 20b. *Transferases (excl. E.C. 2.7.-.)*
- 684 Borgeraas, J., Nislen, K. and Stenersen, J.: Methods for purification of glutathione transferases in the earthworm genus (*Eisenia*) and their characterization. *Comp. Biochem. Physiol. C: Pharmacol., Toxicol. Endocrinol.*, 114C (1996) 129-140; C.A., 125 (1996) 136105v.
- 685 Gandy, J., Primiano, T., Novak, R.F., Kelce, W.R. and York, J.L.: Differential expression of glutathione S-transferase isoforms in compartments of the testis and segments of epididymis of the rat. *Drug Metab. Disp.*, 24 (1996) 725-733.
- 686 Homma, H., Ogawa, K., Hirono, K., Morioka, Y., Hirota, M., Tanahashi, I. and Matsui, M.: Site-directed mutagenesis of rat hepatic hydroxysteroid sulfotransferases. *Biochim. Biophys. Acta*, 1296 (1996) 159-166.
- 687 Jensen, K.F. and Mygind, B.: Different oligomeric states are involved in the allosteric behavior of uracil phosphoribosyltransferase from *Escherichia coli*. *Eur. J. Biochem.*, 240 (1996) 637-645.
- 688 Klebl, B.M. and Pette, D.: A fluorometric assay for measurement of mono-ADP-ribosyltransferase activity. *Anal. Biochem.*, 239 (1996) 145-152.
- 689 Rouimi, P., Amglade, P., Debrauwer, L. and Tulliez, J.: Characterization of pig liver glutathione S-transferases using HPLC-electrospray ionization mass spectrometry. *Biochem. J.*, 317 (1996) 879-884.
- 690 Tilgmann, C. and Ulmanen, I.: Purification methods of mammalian catechol-O-methyltransferases. *J. Chromatogr. B*, 684 (1996) 147-161 - a review with 110 refs.
- 691 Tuomainen, P., Reenilä, I. and Männistö, P.T.: Validation of assay of catechol-O-methyltransferase activity in human erythrocytes. *J. Pharm. Biomed. Anal.*, 14 (1996) 515-523.
- 692 Wilhelm, B., Meinhardt, A. and Seitz, J.: Transglutaminases: purification and activity assays. *J. Chromatogr. B*, 684 (1996) 163-177 - a review with 79 refs.
- For additional information see C.A.:
125 (1996) 108572b.
- See also 271.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.)*
- 693 Adamikova, L., Resnick, R.J. and Tomaska, L.: Enrichment of yeast protein tyrosine kinase activity by substrate affinity chromatography. *Yeast*, 12 (1996) 833-838; C.A., 125 (1996) 108565b.
- 694 Bönisch, H., Backmann, J., Kath, T., Naumann, D. and Schäfer, G.: Adenylate kinase from *Sulfolobus acidocaldarius*: expression in *Escherichia coli* and characterization of Fourier transform infrared spectroscopy. *Arch. Biochem. Biophys.*, 333 (1996) 75-84.
- 695 Boutin, J.A.: Tyrosine protein kinase assays. *J. Chromatogr. B*, 684 (1996) 179-199 - a review with 160 refs.
- 696 Brunati, A.M., James, P., Guerra, B., Ruzzene, M., Donella-Deana, A. and Pinna, L.A.: The spleen protein-tyrosine kinase TPK-IIb is highly similar to the catalytic domain of p72^{syk}. *Eur. J. Biochem.*, 240 (1996) 400-407.

- 697 Cheng, H., Bjorge, J.D., Aebersold, R., Fujita, D.J. and Wang, J.H.: Purification of bovine thymus cytosolic C-terminal Src kinase (CSK) and demonstration of differential efficiencies of phosphorylation and inactivation of p56^{lyn} and pp60^{c-src} by CSK. *Biochemistry*, 35 (1996) 11874-11887.
- 698 Drepper, A. and Pape, H.: Acarbose 7-phosphotransferase from *Actinoplasma* sp.: purification, properties and possible physiological function. *J. Antibiot.*, 49 (1996) 664-668.
- 699 Gatti, A., Wang, X. and Robinson, P.J.: Protein kinase C- α is multiply phosphorylated in response to phorbol ester stimulation of PC12 cells. *Biochim. Biophys. Acta*, 1313 (1996) 111-118.
- 700 Harada, S., Karino, A., Shimoyama, Y., Shamsa, F. and Ohtsuki, K.: Identification of glycorrhizin-binding protein kinase and casein kinase II and characterization of its associated phosphate receptors in mouse liver. *Biochem. Biophys. Res. Commun.*, 227 (1996) 102-109.
- 701 Haystead, T.A.J.: γ -Phosphate-linked adenosine 5'-triphosphate Sepharose for affinity chromatography of protein kinases. *U.S. US 5,536,822* (Cl. 536-26.26; C07H19/20), 16 Jul. 1996, Appl. 207,632, 9 Mar. 1994; 16 pp.; *C.A.*, 125 (1996) 136426a.
- 702 Hu, Z. and Plaxton, W.C.: Purification and characterization of cytosolic pyruvate kinase from leaves of the castor oil plant. *Arch. Biochem. Biophys.*, 333 (1996) 298-307.
- 703 Schachter, J.B., Lester, D.S. and Alkon, D.L.: Synergistic activation of protein kinase C by arachidonic acid and diacylglycerols *in vitro*: generation of a stable membrane-bound, cofactor-independent state of protein kinase C activity. *Biochim. Biophys. Acta*, 1291 (1996) 167-176.
- 704 Schaloske, R., Biethinger, M., Fothergill-Gilmore, L.A. and Hofer, H.W.: Activation by phosphorylation of phosphofructokinase from the annelid *Lumbricus terrestris* and comparison of phosphorylated sites in invertebrate phosphofructokinases. *Biochem. J.*, 317 (1996) 377-383.
- 705 Weijland, A., Neubauer, G., Courtneidge, S.A., Mann, M., Wierenga, R.K. and Superti-Furga, G.: The purification and characterization of the catalytic domain of Src expressed in *Schizosaccharomyces pombe*. Comparison with unphosphorylated and tyrosine phosphorylated species. *Eur. J. Biochem.*, 240 (1996) 756-754.
- See also 311.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.)*
- 706 Antikainen, M., Suurinkeroinen, L., Jauhainen, M., Ehnholm, C. and Taskinen, M.-R.: Development and evaluation of an ELISA method for the determination of lipoprotein lipase mass concentration - comparison with a commercial, one-step enzyme immunoassay. *Eur. J. Clin. Chem. Clin. Biochem.*, 34 (1996) 547-553.
- 707 Caselli, A., Cirri, P., Bonifacio, S., Manao, G., Camici, G., Capugli, G., Moretti, G. and Ramponi, G.: Identity of zinc ion-dependent acid phosphatase from bovine brain and *myo*-inositol 1-phosphatase. *Biochim. Biophys. Acta*, 1290 (1996) 241-249.
- 708 Cashikar, A.G. and Rao, N.M.: Role of the intersubunit disulfide bond in the unfolding pathway of dimeric red kidney bean purple acid phosphatase. *Biochim. Biophys. Acta*, 1296 (1996) 76-84.
- 709 Ghazaleh, F.A., Omburo, G.A. and Colman, R.W.: Evidence for the presence of essential histidine and cysteine residues in platelet cGMP-inhibited phosphodiesterase. *Biochem. J.*, 317 (1996) 495-501.
- 710 Hwang, S.C., Jhon, D., Bae, Y.S., Kim, J.H. and Rhee, S.G.: Activation of phospholipase C- γ by the concerted action of tau proteins and arachidonic acid. *J. Biol. Chem.*, 271 (1996) 18342-18349.
- 711 Lin, S.F., Lee, J.C. and Chiou, C.M.: Purification and characterization of a lipase from *Neurospora* sp. TT-241. *J. Am. Oil Chem. Soc.*, 73 (1996) 739-745.
- 712 Maeda, S. and Kashiwabara, Y.: Purification and properties of a membrane-bound phospholipase B from *Mycobacterium lepraeumum*. *Biochim. Biophys. Acta*, 1303 (1996) 31-38.
- 713 O'Neill, F.H., Christov, L.P., Botes, P.J. and Prior, B.A.: Rapid and simple assay for feruloyl and p-coumaroyl esterases. *World J. Microbiol. Biotechnol.*, 12 (1996) 239-242; *C.A.*, 125 (1996) 10843n.
- 714 Pete, M.J. and Exton, J.H.: Purification of a lysophospholipase from bovine brain that selectively deacylates arachidonoyl-substituted lysophosphatidylcholine. *J. Biol. Chem.*, 271 (1996) 18114-18121.
- 715 Sugahara, K. and Kojima, T.: Specific studies of bacterial sulfatases by means of structurally defined sulfated oligosaccharides isolated from shark cartilage chondroitin sulfate D. *Eur. J. Biochem.*, 239 (1996) 865-870.

See also 705.

- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.)*
- 716 Brena, B.M., Pazos, C., Franco-Fraguas, L. and Batista-Viera, F.: Chromatographic methods for amylases. *J. Chromatogr. B*, 684 (1996) 217-237 - a review with 123 refs.
- 717 Chen, J.-s., Saxton, J., Hemming, F.W. and Peberdy, J.F.: Purification and partial characterization of the high and low molecular weight form (S- and F-form) of invertase secreted by *Aspergillus nidulans*. *Biochim. Biophys. Acta*, 1296 (1996) 207-218.
- 718 Kester, H.C.M., Kusters-van Someren, M.A., Müller, Y. and Visser, J.: Primary structure and characterization of an exopolygalacturonase from *Aspergillus tubingensis*. *Eur. J. Biochem.*, 240 (1996) 738-746.
- 719 Maeda, K.: Effective method for purification of lysozyme from human urine. *J. Chromatogr. B*, 685 (1996) 21-25.
- 720 Mikhaylova, M., Wiederschain, G., Mikhaylov, V. and Aerts, J.M.F.G.: The enzymatic hydrolysis of 6-acylamino-4-methylumbelliferyl- β -D-glucosides: identification of a novel human acid β -glucosidase. *Biochim. Biophys. Acta*, 1317 (1996) 71-79.
- 721 Moriyama, T. and Ikeda, H.: Hydrolases acting on glycosidic bonds: chromatographic and electrophoretic separations. *J. Chromatogr. B*, 684 (1996) 201-216 - a review with 199 refs.
- 722 Rao, M.N., Kembhavi, A. and Pant, A.: Implication of tryptophan and histidine in the active site of endo-polygalacturonase from *Aspergillus ustus*: elucidation of the reaction mechanism. *Biochim. Biophys. Acta*, 1296 (1996) 167-173.
- 723 Takesue, Y. and Takesue, S.: Purification and characterization of α -glucosidase complex from the intestine of the frog, *Rana japonica*. *Biochim. Biophys. Acta*, 1296 (1996) 152-158.

- 724 Terebiznik, M.R., Pilosof, A.M.R. and Moreno, S.: Effective purification procedure of *Aspergillus oryzae* α -amylase from solid state fermentation cultures including concanavalin A-Sepharose. *J. Food Biochem.*, 19 (1996) 341-354; C.A., 125 (1996) 52076y.
- 20f. Other hydrolases
- 725 Beharry, S. and Bragg, P.D.: The bound adenine nucleotides of purified bovine mitochondrial ATP synthase. *Eur. J. Biochem.*, 240 (1996) 165-172.
- 726 Chen, D., Dai, D. and Liu, B.: (RP-HPLC method for determination of $\text{Ca}^{2+}, \text{Mg}^{2+}$ -ATPase activity). *Zhongguo Yaoke Daxue Xuebao*, 27 (1996) 99-101; C.A., 125 (1996) 241451c.
- 727 Cowan, D.A. and Daniel, R.M.: Rapid purification of two thermostable proteinases using dye-ligand chromatography. *J. Biochem. Biophys. Methods*, 32 (1996) 1-6; C.A., 125 (1996) 52018f.
- 728 Daniel, R., Kokel, B., Caminade, E., Martel, A. and le Goffic, F.: Assay of *Escherichia coli* dihydroorotate with enantiomeric substrate: practical preparation of carbamyl L-aspartate and high-performance liquid chromatography analysis of catalysis product. *Anal. Biochem.*, 239 (1996) 130-135.
- 729 Dawson, J.A., Mallonee, D.H., Björkhem, I. and Hylemon, P.B.: Expression and characterization of a C_{24} bile acid 7α -dehydroxidase from *Eubacterium sp.* strain VPI 12708 in *Escherichia coli*. *J. Lipid Res.*, 37 (1996) 1258.
- 730 Eriksson, U., Andrén, P.E., Caprioli, R.M. and Nyberg, F.: Reversed-phase high-performance liquid chromatography combined with tandem mass spectrometry in studies of a substance P-converting enzyme from human cerebrospinal fluid. *J. Chromatogr. A*, 743 (1996) 213-220.
- 731 García-Fuentes, L., Ortiz, E., Jara, V. and Barón, C.: Preparative isolation of a soluble form of bovine lung angiotensin converting enzyme by affinity and size exclusion chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2443-2456.
- 732 Gerentes, L., Kessler, N., Thomas, G. and Aymard, M.: Simultaneous purification of influenza haemagglutinin and neuraminidase proteins by immunochromatography. *J. Virol. Methods*, 58 (1996) 155-165; C.A., 125 (1996) 52860z.
- 733 Hembrough, T.A., Kralovics, K.R. and Gonias, S.L.: Cytokeratin 8 released by breast carcinoma cells *in vitro* binds plasminogen and tissue-type plasminogen activator and promotes plasminogen activation. *Biochem. J.*, 317 (1996) 763-769.
- 734 Ho, H.-C., Chen, L.-Y. and Liao, T.-H.: Identification of a fungal protein of *Syncephalastrum racemosum* as aspartic proteinase. *Arch. Biochem. Biophys.*, 334 (1996) 97-103.
- 735 Ibrahim-Granet, O. and Bertrand, O.: Separation of proteases: old and new approaches. *J. Chromatogr. B*, 684 (1996) 239-263 - a review with 161 refs.
- 736 Jiang, D., Liu, F., Zhao, X., Tian, X., Huang, R., Pan, Y. and Yu, P.: (Application of affinity chromatography for purification of earthworm fibrinolytic enzymes). *Zhongguo Yiiao Gongye Zazhi*, 27 (1996) 147-149; C.A., 125 (1996) 241508b.
- 737 Kobayashi, T., Shinozaki, A., Momoi, T., Arahat, K. and Tsukahara, T.: Identification of an interleukin-1 β converting enzyme-like activity that increases upon treatment of P19 cells with retinoic acid as the proteasome. *J. Biochem. (Tokyo)*, 120 (1996) 699-704.
- 738 Laethem, R.M., Blumenkopf, T.A., Cory, M., Elwell, L., Moxhan, C.P., Ray, P.H., Walton, L.M. and Smith, G.K.: Expression and characterization of human pancreatic preprocarboxypeptidase A1 and preprocarboxypeptidase A2. *Arch. Biochem. Biophys.*, 332 (1996) 8-18.
- 739 Lian, W., Wu, D., Konings, W.N., Mierau, I. and Hersh, L.B.: Heterologous expression and characterization of recombinant *Lactococcus lactis* neutral endopeptidase (neprylisin). *Arch. Biochem. Biophys.*, 333 (1996) 121-126.
- 740 Luderer-Gnach, M., Liebig, H.-D., Sommergruber, W., Voss, T., Fessl, F., Skern, T. and Kuechler, E.: Human rhinovirus 2A proteinase mutant and its second-site revertants. *Biochem. J.*, 318 (1996) 213-218.
- 741 Matveyev, V.V., Usmanova, A.M., Morozova, A.V., Collins, J.H. and Khaitina, S.Y.: Purification and characterization of the proteinase ECP 32 from *Escherichia coli* A2 strain. *Biochim. Biophys. Acta*, 1296 (1996) 55-62.
- 742 Meng, Q.C. and Oparil, S.: Purification and assay methods for angiotensin-converting enzyme. *J. Chromatogr. A*, 743 (1996) 105-122 - a review with 113 refs.
- 743 Misaka, T., Kuroda, M., Iwabuchi, K., Abe, K. and Arai, S.: Soyacystatin, a novel cysteine proteinase inhibitor in soybean, is distinct in protein structure and gene organization from other cystatins of animal and plant origin. *Eur. J. Biochem.*, 240 (1996) 609-614.
- 744 Okumura, Y., Kudoh, A., Takashima, M., Inoue, M., Sakai, K. and Kido, H.: Purification and characterization of a novel isoform of mast cell tryptase from rat tongue. *J. Biochem. (Tokyo)*, 120 (1996) 856-864.
- 745 Raspi, G.: Kallikrein and kallikrein-like proteinases: purification and determination by chromatographic and electrophoretic methods. *J. Chromatogr. B*, 684 (1996) 265-287 - a review with 251 refs.
- 746 Sarkar, D. and DasGupta, C.: Characterization of a molten globule intermediate during GdnHCl-induced unfolding of RTEM β -lactamase from *Escherichia coli*. *Biochim. Biophys. Acta*, 1296 (1996) 85-94.
- 747 Shi, Y.-C., Jiang, Y.-M., Sui, D.-X., Li, Y.-L., Chen, T., Ma, L. and Ding, Z.-T.: Affinity chromatography of trypsin using chitosan as ligand support. *J. Chromatogr. A*, 742 (1996) 107-112.
- 748 Soleihac, J.-M., Cornille, F., Martin, L., Lenoir, C., Fournié-Zaluski, M.-C. and Roques, B.P.: A sensitive and rapid fluorescence-based assay for determination of tetanus toxin peptidase activity. *Anal. Biochem.*, 241 (1996) 120-127.
- 749 Werner, E.R., Werner-Felmayer, G. and Wachter, H.: High-performance liquid chromatographic methods for the quantification of tetrahydrobiopterin biosynthetic enzymes. *J. Chromatogr. B*, 684 (1996) 51-58 - a review with 65 refs.
- 750 Zembower, D.E., Neudauer, C.L., Wick, M.J. and Ames, M.M.: Versatile synthetic ligands for affinity chromatography of serine proteinases. *Int. J. Pept. Protein Res.*, 47 (1996) 405-413; C.A., 125 (1996) 80119h.

For additional information see C.A.:
125 (1996) 108439p.

See also 502, 506, 620.

20g. Lyases

- 751 Adam, K.-P., Crock, J. and Croteau, R.: Partial purification and characterization of a monoterpene cyclase, limonene synthase, from the liverwort *Ricciocarpus natans*. *Arch. Biochem. Biophys.*, 332 (1996) 352-356.
- 752 Anagnostopoulos, C.G. and Kyriakidis, D.A.: Regulation of the *Escherichia coli* biosynthetic ornithine decarboxylase activity by phosphorylation and nucleotides. *Biochim. Biophys. Acta*, 1297 (1996) 228-234.
- 753 Bergenhem, N.: Chromatographic and electrophoretic methods related to the carbonic anhydrase isozymes. *J. Chromatogr. B*, 684 (1996) 289-305 - a review with 121 refs.
- 754 De Montigny, C. and Sygusch, J.: Functional characterization of an extreme thermophilic class I fructose-1,6-bisphosphate aldolase. *Eur. J. Biochem.*, 241 (1996) 243-248.
- 755 Dessauer, C.W. and Gilman, A.G.: Purification and characterization of a soluble form of mammalian adenylyl cyclase. *J. Biol. Chem.*, 271 (1996) 16967-16974.
- 756 Ferreira, R.M.B., Franco, E. and Teixeira, A.R.N.: Covalent dimerization of ribulose bisphosphate carboxylase subunits by UV radiation. *Biochem. J.*, 318 (1996) 227-234.
- 757 Kawabe, J., Toya, Y., Schwencke, C., Oka, N., Ebina, T. and Ishikawa, Y.: Soluble adenylyl cyclase from *Spodoptera frugiperda* (Sf9) cells. Purification and biochemical characterization. *J. Biol. Chem.*, 271 (1996) 20132-20137.
- 758 Lechner, T., Lusser, A., Brosch, G., Eberharter, A., Goralik-Schramel, M. and Loidl, P.: A comparative study of histone deacetylases of plant, fungal and vertebrate cells. *Biochim. Biophys. Acta*, 1296 (1996) 181-188.
- 759 Sugahara, K., Nadanaka, S., Takeda, K. and Kojima, T.: Structural analysis of unsaturated hexasaccharides isolated from shark cartilage chondroitin sulfate D that are substrates for the exolytic action of chondroitin ABC lyase. *Eur. J. Biochem.*, 239 (1996) 871-880.
- 760 Tholl, D., Ober, D., Martin, W., Kellermann, J. and Hartmann, T.: Purification, molecular cloning and expression in *Escherichia coli* of homospermidine synthase from *Rhodopseudomonas viridis*. *Eur. J. Biochem.*, 240 (1996) 373-379.
- 761 Yaguchi, T., Oguni, A., Ouchiyama, N., Igarashi, Y. and Kodama, T.: A non-radioisotope anion-exchange chromatographic method to measure the CO₂/O₂ specificity factor for ribulose bisphosphate carboxylase/oxygenase. *Biosci., Biotechnol., Biochem.*, 60 (1996) 942-944; C.A., 125 (1996) 52004y.

See also 497, 749, 769.

20h. Isomerases

- 762 Knudsen, B.R., Straub, T. and Boege, F.: Separation and functional analysis of eukaryotic DNA topoisomerases by chromatography and electrophoresis. *J. Chromatogr. B*, 684 (1996) 307-321 - a review with 101 refs.

20i. Ligases

- 763 Diczfalusi, M.A. and Alexson, S.E.H.: Isolation and characterization of novel long-chain acyl-CoA thioesterase/carboxylesterase isoenzymes from *Candida rugosa*. *Arch. Biochem. Biophys.*, 334 (1996) 104-112.

- 764 Kounga, K., Song, S., Haslam, G.C. and Himes, R.H.: Site-directed mutagenesis of putative catalytic and nucleotide binding sites in N¹⁰-formyltetrahydrofolate synthetase. *Biochim. Biophys. Acta*, 1296 (1996) 112-120.
- 765 Maeck, G.: (Anion-exchange chromatography of glutamine synthetase isoforms from sugar beet leaves). *BIOspektrum*, 2 (1996) 63-64; C.A., 125 (1996) 136021q.
- 766 Rivoal, J., Dynford, R., Plaxton, W.C. and Turpin, D.H.: Purification and properties of four phosphoenolpyruvate carboxylase isoforms from the green alga *Selenastrum minutum*: evidence that association of the 102-kDa catalytic subunit with unrelated polypeptides may modify the physical and kinetic properties of the enzyme. *Arch. Biochem. Biophys.*, 332 (1996) 47-57.
- 20j. Complex mixtures and incompletely identified enzymes
- 767 Bronnenmeier, K., Adelsberger, H., Lottspeich, F. and Staudenbauer, W.L.: Affinity purification of cellulose-binding enzymes of *Clostridium stercorarium*. *Bioseparation*, 6 (1996) 41-45; C.A., 125 (1996) 189054j.
- 768 Dyer, D. and Stoddard, B.: Purification of a bifunctional tetrahydrofolate-dependent enzyme for x-ray crystallography studies. *Biochemica*, (1996) 22-24; C.A., 125 (1996) 28779j.
- 769 Jiang, L.L., Miyazawa, S. and Hashimoto, T.: Purification and properties of rat D-3-hydroxyacyl-CoA dehydratase/D-3-hydroxyacyl-CoA dehydratase/D-3-hydroxyacyl-CoA dehydrogenase bifunctional protein. *J. Biochem. (Tokyo)*, 120 (1996) 633-641.
- 770 Lingner, J. and Cech, T.R.: Purification of telomerase from *Eu- plotes aediculatus*: requirement of a primer 3' overhang. *Proc. Natl. Acad. Sci. U.S.A.*, 93 (1996) 10712-10717; C.A., 125 (1996) 241521a.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. Purines, pyrimidines, nucleosides, nucleotides
- 771 Chaudhary, A.K., Reddy, G.R., Blair, I.A. and Marnett, L.J.: Characterization of an N⁶-oxopropenyl-2'-deoxyadenosine adduct in malondialdehyde-modified DNA using liquid chromatography/electrospray ionization tandem mass spectrometry. *Carcinogenesis*, 17 (1996) 1167-1170; C.A., 125 (1996) 81159h.
- 772 Henle, E.S., Luo, Y., Gassmann, W. and Linn, S.: Oxidative damage to DNA constituents by iron-mediated Fenton reactions. The deoxyguanosine family. *J. Biol. Chem.*, 271 (1996) 21177-21186.
- 773 Keough, T., Shaffer, J.D., Lacey, M.P., Riley, T.A., Marvin, W.B., Scurria, M.A., Hasselfield, J.A. and Hesselberth, E.P.: Detailed characterization of antisense DNA oligonucleotides. *Anal. Chem.*, 68 (1996) 2305-2312.
- 774 Luo, Y., Henle, E.S. and Linn, S.: Oxidative damage to DNA constituents by iron-mediated Fenton reactions. The deoxycytidine family. *J. Biol. Chem.*, 271 (1996) 21167-21176.
- 775 Pandya, G.A. and Moriya, M.: 1,N⁶-Ethenodeoxyadenosine, a DNA adduct highly mutagenic in mammalian cells. *Biochemistry*, 35 (1996) 11487-11492.

- 776 Takenaka, S., Masuda, W., Tsuyama, S., Tamura, Y., Miyatake, K. and Nakano, Y.: Purification and characterization of arginine:mono-ADP-ribosylhydrolase from *Euglena gracilis* Z. *J. Biochem. (Tokyo)*, 120 (1996) 792-796.
- 777 Tonello, G., Daglio, M., Zaccarelli, N., Sotofattori, E., Mazzei, M. and Balbi, A.: Characterization and quantitation of the active polynucleotide fraction (PDRN) from human placenta, a tissue repair stimulating agent. *J. Pharm. Biomed. Anal.*, 14 (1996) 1555-1560.
- 778 Zeng, Y., Shabalina, Y., Szymilo, T., Pastiszak, I., Drake, R.R. and Elbein, A.D.: Synthesis of aryl azide derivatives of UDP-GlcNAc and UDP-GalNAc and their use for the affinity labeling of glycosyltransferases and the UDP-HexNAc pyrophosphorylase. *Anal. Biochem.*, 239 (1996) 99-106.

For additional information see C.A.:
125 (1996) 29519m.

See also 783, 798, 801, 1092, 1160.

21b. Nucleic acids, RNA

- 779 Chinali, G.: A procedure for isolation of pure complexes of acylated and unacylated tRNA^{Pro} with ribosome from *Escherichia coli*. *Anal. Biochem.*, 241 (1996) 30-34.
- 780 Jarrett, H.W.: Hybrid selection with cDNA-silica. *J. Chromatogr. A*, 742 (1996) 87-94.
- 781 Kanduc, D., Mosavi, A., Bracalello, A. and de Benedetto, C.: Isolation of rat liver tRNA^{Asn}, tRNA^{Leu}, and tRNA^{Ser} isoacceptor species by HPLC. *Biochem. Arch.*, 12 (1996) 27-32; C.A., 125 (1996) 189727f.
- 782 Li, Z., Shalom, A., Huang, Y., Mak, J., Arts, E., Wainberg, M.A. and Kleiman, L.: Multiple forms of tRNA^{Lys3} in HIV-1. *Biochem. Biophys. Res. Commun.*, 227 (1996) 530-540.
- 783 Rogers, M.J., Brown, R.J., Hodkin, V., Blackburn, G.M., Russel, R.G.G. and Watts, D.J.: Bisphosphonates are incorporated into adenine nucleotides by human aminoacyl-tRNA synthetase enzymes. *Biochem. Biophys. Res. Commun.*, 224 (1996) 863-869.

See also 512, 552.

21c. Nucleic acids, DNA

- 784 Kaur, H. and Halliwell, B.: Measurement of oxidized and methylated DNA bases by HPLC with electrochemical detection. *Biochem. J.*, 318 (1996) 21-23.
- 785 Min, C. and Verdine, G.L.: Immobilized metal affinity chromatography of DNA. *Nucleic Acids Res.*, 24 (1996) 3806-3810; C.A., 125 (1996) 266773x.
- 786 Pedersen, N.E.: Spin-column chromatography for DNA purification. *Anal. Biochem.*, 239 (1996) 117-118.
- 787 Yamakawa, H., Higashino, K.-i. and Ohara, O.: Sequence-development DNA separation by anion-exchange high-performance liquid chromatography. *Anal. Biochem.*, 240 (1996) 242-250.

For additional information see C.A.:
125 (1996) 216353q.

See also 512, 552.

21e. Structural studies on DNA and DNA mapping

- 788 Huber, C.G. and Berti, G.N.: Detection of partial denaturation in AT-rich DNA fragments by ion-pair reversed-phase chromatography. *Anal. Chem.*, 68 (1996) 2959-2965.

For additional information see C.A.:
125 (1996) 240235y.

22. ALKALOIDS

- 789 Bertoni, G., di Palo, V., Tappa, R. and Possanzini, M.: Fast determination of nicotine and 3-ethenylpyridine in indoor environments. *Chromatographia*, 43 (1996) 296-300.
- 790 Bickel, U., Schumacher, O.P., Kang, Y.-S. and Voigt, K.: Poor permeability of morphine 3-glucuronide and morphine 6-glucuronide through the blood-brain barrier in the rat. *J. Pharmacol. Exp. Ther.*, 278 (1996) 107-113.
- 791 Bogusz, M.J., Maier, R.-D., Schwy-Bochat, K.-H. and Kohls, U.: Applicability of various brands of mixed-phase extraction columns for opiate extraction from blood and serum. *J. Chromatogr. B*, 683 (1996) 177-188.
- 792 Callaway, J.C., Raymon, L.P., Hearn, W.L., McKenna, D.J., Grob, C.S., Brito, G.S. and Mash, D.C.: Quantitation of N,N-dimethyltryptamine and harmala alkaloids in human plasma after oral dosing with ayahuasca. *J. Anal. Toxicol.*, 20 (1996) 492-497; C.A., 125 (1996) 240391w.
- 793 Cepas, J., Silva, M. and Pérez-Bendito, D.: Integrated derivatization-chemiluminescence detection system for the determination of β-carboline alkaloids by high-performance liquid chromatography. *J. Chromatogr. A*, 749 (1996) 73-80.
- 794 Dash, A.K. and Wong, S.-T.: Liquid chromatographic method for the determination of nicotine in pharmaceutical formulations. *J. Chromatogr. A*, 749 (1996) 81-85.
- 795 Edwards, E.J. and Cobb, A.H.: Improved high-performance liquid chromatographic method for the analysis of potato (*Solanum tuberosum*) glycoalkaloids. *J. Agric. Food Chem.*, 44 (1996) 2212-2216; C.A., 125 (1996) 165925t.
- 796 Glass, R.L. and Johnson, M.B.: Analysis of cuscohygrine in coca leaves by high performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 177-1784.
- 797 Maya, M.T., Pais, J.P., Araujo, H.M. and Morais, J.A.: Determination of apovincaminic acid in human plasma by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 617-622.
- 798 Meyer, A., Ngiruwonsanga, T. and Henze, G.: Determination of adenine, caffeine, theophylline and theobromine by HPLC with amperometric detection. *Fresenius J. Anal. Chem.*, 356 (1996) 284-287.
- 799 Papadoyannis, I.N., Samanidou, V.F. and Georga, K.A.: Solid-phase extraction study and photodiode array RP-HPLC analysis of xanthine derivatives in human biological fluids. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2559-2578.
- 800 Perez-Martinez, I., Sagrado, S. and Medina-Hernandez, M.J.: Determination of theophylline in pharmaceuticals by micellar liquid chromatography and spectrophotometric detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1957-1966.

- 801 Rasmussen, B.B. and Broesen, K.: Determination of urinary metabolites of caffeine for the assessment of cytochrome P4501A2, xanthine oxidase, and N-acetyltransferase activity in humans. *Ther. Drug Monit.*, 18 (1996) 254-262; C.A., 125 (1996) 75211c.
- 802 Tsuchiya, H., Sato, M., Hayashi, H., Kato, H., Kureishi, H. and Hayashi, T.: Simultaneous determination of tetrahydro- β -carbolines and β -carbolines. *Chromatographia*, 43 (1996) 419-425.
- 803 Xu, A., Havel, J., Linderholm, K. and Hulse, J.: Development and validation of an LC/MS/MS method for the determination of L-hyosciamine in human plasma. *J. Pharm. Biomed. Anal.*, 14 (1995) 33-42.

See also 75, 231, 1025, 1059, 1136, 1141, 1157, 1158.

23. OTHER SUBSTANCES CONTAINING HETERO CYCLIC NITROGEN

23a. *Porphyrins and other pyrroles*

See 1052.

23b. *Bile pigments*

- 804 Gordon, E.R., Seligson, D. and Flye, M.W.: Serum bilirubin pigments covalently linked to albumin. *Arch. Pathol. Lab. Med.*, 120 (1996) 648-653; C.A., 125 (1996) 216000x.
- 805 Mizobe, M., Kondo, F., Toyoshima, C., Kumamoto, K., Terada, T. and Nasu, H.: Rapid analysis of four bilirubins in domestic animal sera using high-performance liquid chromatography. *J. Vet. Med. Sci.*, 58 (1996) 495-499; C.A., 125 (1996) 189726e.

23c. *Indole derivatives and plant hormones (gibberellins)*

- 806 Araki, J. and Tabei, H.: (The introduction of highly sensitive analysis of serotonin and dopamine with microbore HPLC using radial flow cell). *Kuromatogurafi*, 17 (1996) 96-97; C.A., 125 (1996) 96243e - a review with no refs.
- 807 Colgan, S.T., Haggan, G.R. and Reed, R.H.: Assay and purity evaluation of 5-chlorooxindole by liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 825-833.
- 808 Hariharan, M., Naickar, J.S. and Halibrand, A.S.: Melatonin determination in human urine by liquid chromatography using fluorometric detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1745-1756.
- 809 Mochida, I., Sakanishi, K., Senzaki, T. and Furumoto, M.: (Method for chromatographic recovery of indole from indole-containing oil by adsorption on ion exchange resin using super critical carbon dioxide). *Jpn. Kokai Tokkyo Koho JP 08,198,848 [96,198,848]* (Cl. C07D209/04), 6 Aug. 1996, Appl. 95/10,090, 25 Jan. 1995; 4 pp.; C.A., 125 (1996) 247612c.
- 810 Navas Diaz, A., Garcia Pareja, A. and Garcia Sanchez, F.: Normal-phase liquid chromatography of plant hormones using reversed cholic acid micelles as the mobile phase. *Anal. Chem.*, 68 (1996) 3029-3032.
- 811 Pezzella, A., Napolitano, A., d'Ischia, M. and Prota, G.: Oxidative polymerization of 5,6-dihydroxyindole-2-carboxylic acid to melanin: a new insight. *Tetrahedron*, 52 (1996) 7913-7920; C.A., 125 (1996) 86385p.

For additional information see C.A.:
125 (1996) 29310m.

See also 1007.

23d. *Pyridine derivatives*

- 812 James, I.T., Walne, A.J. and Perrett, D.: Automated solid-phase extraction with high-performance liquid chromatography analysis of pyridinium crosslinks. *Anal. Biochem.*, 240 (1996) 29-36.

See also 132.

23e. *Other N-heterocyclic compounds*

- 813 Deng, Y., Maruyama, W., Yamamura, H., Kawai, M., Dostert, P. and Naoi, M.: Mechanism of enantioseparation of salsolinols, endogenous neurotoxins in human brain, with ion-pair chromatography using β -cyclodextrin as a mobile phase additive. *Anal. Chem.*, 68 (1996) 2826-2831.
- 814 Karamanos, N.K. and Tsegenidis, T.: Analysis of carcinogenic heterocyclic amines by high performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2247-2256.
- 815 Longhi, M., Linares, M. and de Bertorello, M.M.: High performance liquid chromatography of isoxazolyl-naphthoquinones: a comparison between experimental and theoretical lipophilicity. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1947-1956.
- 816 Magaletta, R.L. and Ho, C.-T.: Effect of roasting time and temperature on the generation of nonvolatile (polyhydroxyalkyl)pyrazine compounds in peanuts, as determined by high-performance liquid chromatography. *J. Agric. Food Chem.*, 44 (1996) 2136-2142; C.A., 125 (1996) 165927v.
- 817 Nakagawa, H., Nihonmatsu, N., Ohta, S. and Hirobe, M.: Effects of new endogenous nonprotein amino acids, 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid derivatives, on behavior of mice. *Biochem. Biophys. Res. Commun.*, 225 (1996) 1027-1034.
- 818 Stelzer, A., Meusel, M., Junghänel, J. and Jira, T.: HPLC-Untersuchungen chiraler Triazolophthalazine. *Pharmazie*, 51 (1996) 647-650.
- 819 Tambe, A.S.: High-performance liquid chromatographic determination of cyclohexylhydantoin and cyclohexylidenehydantoin formed during the synthesis of phenylhydantoin from hydantoin and cyclohexanone. *J. Chromatogr. A*, 740 (1996) 284-288.

See also 169, 414, 939, 1017, 1089.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 820 Jensen, D., Pinhack, H. and Reichert, A.: (Determination of element-specific halides and sulphur contents in organic matrixes). *GfT Fachz. Lab.*, 40 (1996) 404-408; C.A., 125 (1996) 47912p - a review with 16 refs.

- 821 Koyama, H., Sugioka, N., Hirata, I., Ohta, T. and Kishimoto, H.: Determination of L-buthionin (SR)-sulfoximine, gamma-glutamylcysteine synthetase inhibitor in rat plasma with HPLC after prelabeling with dansyl chloride. *J. Chromatogr. Sci.*, 34 (1996) 326-320.
- 822 Lambopoulos, N., Cardwell, T.J., Caridi, D. and Marriott, P.J.: Separation of zinc dialkyldithiophosphates in lubricating oil additives by normal-phase high-performance liquid chromatography. *J. Chromatogr. A*, 749 (1996) 87-94.
- 823 Prestera, T., Fahey, J.W., Holtzclaw, W.D., Abeygunawardana, C., Kachinski, J.L. and Talalay, P.: Comprehensive chromatographic and spectroscopic methods for the separation and identification of intact glucosinolates. *Anal. Biochem.*, 239 (1996) 168-179.
- 824 Shada, D.M., Wong, C.F., Elrod, L., Jr., Morley, J.A. and Gay, C.M.: Determination of 1-benzol[b]thien-2-ylethanone and related impurities by high performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 501-510.

For additional information see C.A.:
125 (1996) 47935y, 146638k.

See also 53, 964, 1187.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 825 Brearley, C.A. and Hanke, D.E.: Inositol phosphates in barley (*Hordeum vulgare* L.) aleurone tissue are stereochemically similar to the products of breakdown of InsP₆ *in vitro* by wheat-bran phytase. *Biochem. J.*, 318 (1996) 279-286.

See also 164, 371, 409, 704, 822.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 826 Cairns, W.R.L., Hill, S.J. and Ebdon, L.: Directly coupled high performance liquid chromatography - inductively coupled plasma-mass spectrometry for the determination of organometallic species in tea. *Microchem. J.*, 54 (1996) 88-110; C.A., 125 (1996) 219894c.
- 827 Liu, Y.-H., Yu, Z.-Y., Huo, S.-Q. and Wu, Y.-J.: (Chromatographic features and IR spectra of ferrocenylimines and their cyclomer-cured derivatives). *Gaoeng Xuexiao Huaxue Xuebao*, 17 (1996) 901-905; C.A., 125 (1996) 248042d.

For additional information see C.A.:
125 (1996) 32156w.

See also 146, 822, 940, 1212.

26b. Boranes, silanes and related non-metallic compounds

- 828 Zhang, X., Cornelis, R., de Kimpe, J., Mees, L., Vanderbiesen, V., de Cubber, A. and Vanholder, R.: Accumulation of arsenic species in serum of patients with chronic renal disease. *Clin. Chem. (Washington)*, 42 (1996) 1231-1237.

26c. Coordination compounds

- 829 Basova, E.M., Ivanov, V.M., Bol'shova, T.A., Shapovalova, E.N. and Sphigun, O.A.: (Ion-pair reversed-phase chromatography of metal chelate. Chromatographic behavior, retentive models, and analytical application). *Vestn. Mosk. Univ., Ser. 2: Khim.*, 37 (1996) 3-14; C.A., 125 (1996) 131233a - a review with 36 refs.
- 830 Li, C.-H., Hor, T.S.A. and Lee, H.K.: High-performance liquid chromatography of the coordination isomers of triphenylphosphine-substituted homo- and hetero-bimetallic carbonyl complexes of manganese and rhenium. *J. Chromatogr. A*, 740 (1996) 231-236.
- 831 Wang, Y.-M. and Cheng, T.-H.: Detection of iron(III) chelates separated by reversed-phase high-performance liquid chromatography. *Gaoxiong Yixue Zazhi*, 12 (1996) 267-273; C.A., 125 (1996) 109247m.
- 832 Yang, W., Li, C., Gao, J., Kang, J. and Wang, J.: Composition determination of rare earth complexes by HPLC. *Anal. Lett.*, 29 (1996) 1637-1645.

For additional information see C.A.:
125 (1996) 184457h.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 833 Acar, J. and Gokmen, V.: A simple HPLC method for the determination of total vitamin C in fruit juices and drinks. *Fruit Process.*, 6 (1996) 198-201; C.A., 125 (1996) 56511j.
- 834 Albert, K., Schlotterbeck, G., Tseng, L.-H. and Braumann, U.: Application of on-line capillary high-performance liquid chromatography-nuclear magnetic resonance spectrometry coupling for the analysis of vitamin A derivatives. *J. Chromatogr. A*, 750 (1996) 303-309.
- 835 Blanco, D., Llaneza, M.B. and Gutierrez, M.D.: A paired ion liquid chromatographic method for thiamine determination in selected foods. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2155-2164.
- 836 Cavazzini, D., Galdieri, M. and Ottanello, S.: Retinoic acid synthesis in the somatic cells of rat seminiferous tubules. *Biochim. Biophys. Acta*, 1313 (1996) 139-145.
- 837 Disdier, B., Bun, H., Catalin, J. and Durand, A.: Simultaneous determination of all-trans-, 13-cis-, 9-cis-retinoic acid and their 4-oxo-metabolites in plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 683 (1996) 143-154.
- 838 Galuba, G.A. and Gogolewski, M.: The comparison of SFC and HPLC techniques utilized for separation of tocopherols in Tenox GT-2. *Chem. Anal. (Warsaw)*, 41 (1996) 737-741; C.A., 125 (1996) 245897a.
- 839 Habif, Turgan, Ozmen and Bayindir: Determination of vitamin E in serum, a comparison of colorimetry and HPLC. *Turk. J. Med. Sci.*, 26 (1996) 277-279; C.A., 125 (1996) 137104f.

- 840 Hewavitharana, A.K., van Brakel, A.S. and Harnett, M.: Simultaneous liquid chromatographic determination of vitamins A, E and β -carotene in common dairy foods. *Int. Dairy J.*, 6 (1996) 613-624; C.A., 125 (1996) 113037k.
- 841 Iino, T., Sawada, H., Tsusué, M. and Takikawa, S.-I.: Discovery of a new tetrahydrobiopterin-synthesizing enzyme in the lemon mutant of the silkworm *Bombyx mori*. *Biochim. Biophys. Acta*, 1297 (1996) 191-199.
- 842 Klyuev, S.A.: (Modified method for the determination of vitamins A and E by reversed-phase HPLC). *Khim.-Farm. Zh.*, 30 (1996) 52-53; C.A., 125 (1996) 67950c.
- 843 Klyuev, S.A.: Determination of vitamins A and E by high-performance liquid chromatography with preliminary equilibration of the distribution between two immiscible liquid phases. *J. Anal. Chem.*, 51 (1996) 884-886; C.A., 125 (1996) 219890y.
- 844 Lanvers, C., Hempel, G., Blaschke, G. and Boos, J.: Simultaneous determination of all-trans-, 13-cis- and 9-cis-retinoic acid, their 4-oxo metabolites and all-trans-retinol in human plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 685 (1996) 233-240.
- 845 Li, S., Barua, A.B. and Huselton, C.A.: Quantification of retinoyl- β -glucuronides in rat urine by reversed-phase high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 683 (1996) 155-162.
- 846 Liu, Q., Scheller, K.K. and Schaefer, D.M.: Technical note: a simplified procedure for vitamin E determination in beef muscle. *J. Anim. Sci.*, 74 (1996) 2406-2410; C.A., 125 (1996) 273922y.
- 847 Luccock, M.D., Daskalakis, I., Schorah, C.J., Levene, M.I. and Hartley, R.: Analysis and biochemistry of blood folate. *Biochem. Mol. Med.*, 58 (1996) 93-112; C.A., 125 (1996) 109252j.
- 848 Ma, Q., Zhou, H., Li, X. and Xu, D.: (Determination of carotenoids in human serum by non aqueous reversed-phase liquid chromatography). *Sepu*, 14 (1996) 277-279; C.A., 125 (1996) 136819f.
- 849 Margolis, S.A. and Duewer, D.L.: Measurement of ascorbic acid in human plasma and serum: stability, intralaboratory repeatability, and interlaboratory reproducibility. *Clin. Chem. (Washington)*, 42 (1996) 1257-1262.
- 850 Nagy, E., Mezey, G. and Kemecsei, E.: Development and validation of a high-performance liquid chromatographic method for the determination of folic acid. *Acta Pharm. Hung.*, 66 (1996) 181-184; C.A., 125 (1996) 242050h.
- 851 Nguyen, T.X.Q., le Bouil, A. and Allain, P.: Simultaneous determination of vitamins A and E in pediatric plasma by HPLC. *J. Pharm. Clin.*, 14 (1995) 269-271; C.A., 125 (1996) 184763e.
- 852 Ohta, K. and Harada, K.: (Studies on the measurement of ascorbic acid and dehydroascorbic acid in tea plants). *Nippon Nogei Kagaku Kaishi*, 70 (1996) 873-882; C.A., 125 (1996) 162710h.
- 853 Sakai, T., Murata, H. and Ito, T.: High-performance liquid chromatographic analysis of ascorbyl-2-phosphate in fish tissues. *J. Chromatogr. B*, 685 (1996) 196-198.
- 854 Ziouzenkova, O., Winkelhofer-Roob, B.M., Puhl, H., Roob, J.M. and Estebauer, H.: Lack of correlations between the α -tocopherol content of plasma and LDL, but high correlation for γ -tocopherol and carotenoids. *J. Lipid Res.*, 37 (1996) 1936-1946.

See also 358, 382, 655, 749, 947, 1176, 1179.

28. ANTIBIOTICS

- 855 Adams, E., Schepers, R., Roets, E. and Hoogmartens, J.: Determination of neomycin sulfate by liquid chromatography with pulsed electrochemical detection. *J. Chromatogr. A*, 741 (1996) 233-240.
- 856 Arcelloni, C., Basile, M., Vaiani, R., Bonini, P. and Paroni, R.: Determination of ceftazidime concentration in Mueller Hinton agar by high-performance liquid chromatography. *J. Chromatogr. A*, 742 (1996) 121-126.
- 857 Barzic, M.-R. and Guittet, E.: Structure and activity of persicomycins, toxins produced by a *Pseudomonas syringae* pv. *persicae/Prunus persica* isolate. *Eur. J. Biochem.*, 239 (1996) 702-709.
- 858 Beconi-Barker, M.G., Roof, R.D., Vidmar, T.J., Hornish, R.E., Smith, E.B., Gatchell, C.L. and Gibertson, T.J.: Ceftiofur sodium: absorption, distribution, metabolism, and excretion in target animals and its determination by high-performance liquid chromatography. *ACS Symp. Ser.*, 636(Veterinary Drug Residues) (1996) 70-84; C.A., 125 (1996) 237465t.
- 859 Benaji, B., Dine, T., Luyckx, M., Gressier, B., Brunet, C., Goudaliez, F., Cazin, M. and Cazin, J.C.: Stability and compatibility studies of zorubicin in intravenous fluids and PVC infusion bags. *J. Pharm. Biomed. Anal.*, 14 (1996) 695-705.
- 860 Carson, M.C.: Confirmation of multiple tetracycline residues in milk by liquid chromatography/particle beam mass spectrometry. In: Enne, G., Kuiper, H.A. and Valentini, A. (Editors), *Residues Vet. Drugs Mycotoxins Animm. Prod.*, Wageningen Pers., Wageningen, 1996, pp. 72-75; C.A., 125 (1996) 193687k.
- 861 Carson, M.C., Righter, H.F. and Wagner, D.D.: Detection of orally administered oxytetracycline and chlortetracycline in bovine milk by a multiresidue screening assay (Charm II) and by LC. In: Enne, G., Kuiper, H.A. and Wagner, D.D. (Editors), *Residues Vet. Drugs Mycotoxins Animm. Prod.*, Wageningen Pers., Wageningen, 1996, pp. 76-80; C.A., 125 (1996) 193688m.
- 862 Chen, B.-M., Xia, L.-W. and Xia, X.-B.: Determination of mitomycin C in rabbit ocular tissue after topical administration by high performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2659-2667.
- 863 Cruz, R., Arias, M.S., Arias, M.E. and Soliveri, J.: APHE-3 and APHE-4, two new pyrazoloisoquinolinone antibiotics produced by *Streptoverticillium friseocarneum* NCIM 40447. *J. Antibiot.*, 49 (1996) 700-702.
- 864 Cui, C.-B., Kakeya, H., Okada, G., Onose, R. and Osada, H.: Novel mammalian cell cycle inhibitors, tryptostatins A, B and other diketopiperazines produced by *Aspergillus fumigatus*. I. Taxonomy, fermentation, isolation and biological properties. *J. Antibiot.*, 49 (1996) 527-533.
- 865 Decolin, D., Gavriloff, C., Archimbaud, P., Leroy, P. and Nicolas, A.: (Analysis of antibiotic residues on edible tissues by HPLC-fluorimetry). *Ann. Falsif. Expert. Chim. Toxicol.*, 89 (1996) 11-23; C.A., 125 (1996) 273923z.

- 866 Fiedler, H.-P., Nega, M., Pfefferle, C., Groth, I., Kempter, C., Stephan, H.O. and Metzger, J.W.: Kanchanamycins, new polyol macrolide antibiotics produced by *Streptomyces olivaceus* tū 4018. I. Taxonomy, fermentation, isolation and biological activities. *J. Antibiot.*, 49 (1996) 758-764.
- 867 Funabashi, Y., Hakoda, S., Watomi, N., Koyama, K., Tanida, S. and Harada, S.: Microbial conversion of EM523, gastrointestinal motor stimulating agents. *J. Antibiot.*, 49 (1996) 802-810.
- 868 Funabashi, Y., Maeshiba, Y., Inatomi, N., Tanayama, S., Harada, S., Itoh, Z. and Omura, S.: Bioactive metabolites of EM574 and EM523, erythromycin derivatives having strong gastrointestinal motor stimulating activity. *J. Antibiot.*, 49 (1996) 794-801.
- 869 Gallo, P., de Grado, G. and Serpe, L.: Chloramphenicol detection in feed by means of chromatographic techniques. *Ital. J. Food Sci.*, 8 (1996) 49-56; *C.A.*, 125 (1996) 113012y.
- 870 Gonschior, A.-K., Christians, U., Winkler, M., Linck, A., Baumann, J. and Sewing, K.-F.: Tacrolimus (FK 506) metabolite patterns in blood from liver and kidney transplant patients. *Clin. Chem. (Washington)*, 42 (1996) 1426-1432.
- 871 Johnson, R.A., Marshall, V.P., Li, G.P., Sih, J.C., Cialdella, J.I., Liggett, W.F. and Nidy, E.G.: Microbiological N-deoxygenation and C-oxygenation of pioglitazone-N-oxide in a single fermentation. *J. Antibiot.*, 49 (1996) 788-793.
- 872 Kimura, K.-i., Takahashi, H., Miyata, N., Yoshihama, M. and Uromoto, M.: New piericidin antibiotics, 7-demethylpiericidin A₁ and 7-demethyl-3-rhamnopiericidin A₁. *J. Antibiot.*, 49 (1996) 697-699.
- 873 Kitahashi, T. and Ohba, Y.: Determination of vancomycin in serum using high-performance liquid chromatography with serum direct injection. *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 25 (1996) 104-108; *C.A.*, 125 (1996) 157653n.
- 874 Kobayashi, H., Namikoshi, M., Yoshimoto, T. and Yokoki, T.: A screening method for antimitotic and antifungal substances using conidia of *Pyricularia oryzae*, modification and application to tropical marine fungi. *J. Antibiot.*, 49 (1996) 873-879.
- 875 Kuno, F., Otaguro, K., Shiomi, K., Iwai, Y. and Omura, S.: Arisugacins A and B, novel and selective acetylcholinesterase inhibitors from *Penicillium* sp. FO-4259. I. Screening, taxonomy, fermentation, isolation and biological activity. *J. Antibiot.*, 49 (1996) 742-747.
- 876 Lacroix, P.M., Curran, N.M. and Sears, R.W.: High-pressure liquid chromatographic methods for ciprofloxacin hydrochloride and related compounds in raw materials. *J. Pharm. Biomed. Anal.*, 14 (1996) 641-654.
- 877 Li, K., Wang, P., Yuan, Y. and Liu, X.: Determination of cyclosporin-A in human whole blood by reversed phase liquid chromatography with single-step extraction. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2179-2188.
- 878 Liu, W.T., Tamolang, M.B., Pang, H., Ren, Y. and Wong, P.Y.: Semipreparative chromatographic separation of cyclosporin G metabolites generated by microsomes from rabbits treated with rifampicin. *J. Pharmacol. Toxicol. Methods*, 35 (1996) 121-129; *C.A.*, 125 (1996) 48318m.
- 879 Lundell, J., Ruohonen, J., Aaltonen, O., Alkio, M., Hase, A. and Suortti, T.: Process for preparation of pure cyclosporin chromatographically using an eluent consisting essentially of high pressure carbon dioxide. *PCT Int. Appl.* WO 96 27,607 (Cl. C07K1/18), 12 Sep. 1996, FI Appl. 95/983, 3 Mar. 1995; 26 pp.; *C.A.*, 125 (1996) 273718m.
- 880 Luo, W., Hansen, E.B., Jr., Ang, C.Y.W. and Thompson, H.C., Jr.: Determination of lincomycin residue in salmon tissues by ion-pair reversed-phase liquid chromatography with electrochemical detection. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 839-843.
- 881 Maertlbauer, E., Dietrich, R. and Usleber, E.: Immunoaffinity chromatography as a tool for the analysis of antibiotics and sulfonamides. *ACS Symp. Ser.*, 636(Veterinary Drug Residues) (1996) 121-131; *C.A.*, 125 (1996) 193766k.
- 882 Matsuoka, M., Maki, T., Banno, K. and Sato, T.: Liquid chromatographic analysis of penem antibiotic FCE22101 in biological fluids with a liquid-liquid extraction procedure. *J. Chromatogr. B*, 685 (1996) 273-279.
- 883 Mayer, A.F. and Deckwer, W.-D.: Ion-pair adsorption chromatography for process purposes. Basic equilibrium studies for the recovery of clavulanic acid by using quaternary ammonium salts. *J. Chromatogr. A*, 741 (1996) 185-203.
- 884 McNeilly, P.J., Reeves, V.B. and Deveau, E.J.I.: Determination of ceftiofur in bovine milk by liquid chromatography. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 844-847.
- 885 Moreno, S., Roca, M. and del Pozo, A.: (Stability of vancomycin eyedrops measured by high-resolution liquid chromatography). *Cienc. Pharm.*, 6 (1996) 77-81; *C.A.*, 125 (1996) 123533h.
- 886 Mosts, W.A. and Harik-Khan, R.: Rapid determination of tetracycline antibiotics in milk and tissues using ion-pairing high-performance liquid chromatography. *ACS Symp. Ser.*, 636(Veterinary Drug Residues) (1996) 85-95; *C.A.*, 125 (1996) 193649z.
- 887 Murakami, Y., Kato, S., Nakajima, M., Matsuoka, M., Kawai, H., Shin-Ya, K. and Seto, H.: Formobactin, a novel free radical scavenging and neuronal cell protecting substance from *Noocardia* sp. *J. Antibiot.*, 49 (1996) 839-845.
- 888 Nasr, M.M. and Tschappeler, T.J.: High performance liquid chromatographic analysis of erythromycin. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2329-2348.
- 889 Okamoto, Y., Itoh, K., Namiki, Y., Matsushita, J., Fujioka, M. and Yasuda, T.: Method development for the determination of cefdinir and its related substances by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 739-748.
- 890 Quarta, C., Borghi, A., Zerilli, L.F., De Pietro, M.T., Ferrari, P., Trani, A. and Lancini, G.C.: Isolation and structure determination of a novel complex of the teicoplanin family. *J. Antibiot.*, 49 (1996) 644-650.
- 891 Schroeder, D.R., Colson, K.L., Klohr, S.E., Lee, M.S., Matson, J.A., Brinen, L.S. and Clardy, J.: Pyrrolosporin A, a new antitumor antibiotic from *Micromonospora* sp. C39217-R109-7 II. Isolation, physico-chemical properties, spectroscopic study and X-ray analysis. *J. Antibiot.*, 49 (1996) 865-872.
- 892 Shirnada, K.-i., Morino, T., Masuda, A., Sato, M., Kitagawa, M. and Saito, S.: Absolute structural determination of stevastelin B. *J. Antibiot.*, 49 (1996) 569-574.
- 893 Shindo, Y., Hayashi, H., Ando, M. and Tatebayashi, T.: Mutagenicity study of the new cephalosporin antibiotic cafiditone pivoxil. *Arzneim.-Forsch.*, 46 (1996) 815-821.
- 894 Stefanelli, S., Sponga, F., Ferrari, P., Sottani, C., Corti, E., Brunati, C. and Islam, K.: Inhibitors of myo-inositol monophosphatase, ATCC 20928 factors A and C. Isolation, physico-chemical characterization and biological properties. *J. Antibiot.*, 49 (1996) 611-616.

- 895 Streit, F., Christians, U., Schiebel, H.-M., Napoli, K.L., Ernst, L., Linck, A., Kahan, B.D. and Sewing, K.-F.: Sensitive and specific quantification of sirolimus (rapamycin) and its metabolites in blood of kidney graft recipients by HPLC/electrospray-mass spectrometry. *Clin. Chem. (Washington)*, 42 (1996) 1417-1425.
- 896 Takahashi, S., Hashimoto, R., Hamano, K., Suzuki, T. and Nakagawa, A.: Melanoxal, new melanin biosynthesis inhibitor discovered by using the larval haemolymph of the silkworm, *Boomyx mori*. Production, isolation, structural elucidation, and biological properties. *J. Antibiot.*, 49 (1996) 513-518.
- 897 Tanimoto, T., Onodera, K., Hosoya, T., Takamatsu, Y., Kinoshita, T., Tago, K., Kogen, H., Fujioka, T., Hamano, K. and Tsujita, Y.: Schizostatin, a novel squalene synthase inhibitor produced by the mushroom, *Schizophyllum commune*. I. Taxonomy, fermentation, isolation, physico-chemical properties and biological activities. *J. Antibiot.*, 49 (1996) 617-623.
- 898 Tsuda, M., Muraoka, Y., Someno, T., Nagai, M., Aoyagi, T. and Takeuchi, T.: Poststatin, a new inhibitor of prolyl endopeptidase VI. Endopeptidase inhibitory activity of poststatin analogues containing pyrrolidine ring. *J. Antibiot.*, 49 (1996) 900-908.
- 899 Ueki, M., Abe, K., Hanafi, M., Shibata, K., Tanaka, T. and Taniguchi, M.: UK-2A, B, C and D, novel antifungal antibiotics from *Streptomyces* sp. 517-02. I. Fermentation, isolation, and biological properties. *J. Antibiot.*, 49 (1996) 639-643.
- 900 Yongxin, Z., Roets, E., Moreno, M.L., Porqueras, E. and Hoogmartens, N.: Evaluation of LC methods for the separation of amoxicillin and its related substances. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1893-1908.
- 901 Zhu, Y., Roets, E., Ni, Z., Moreno, M.L., Porqueras, E. and Hoogmartens, J.: Evaluation of liquid chromatography methods for the separation of ampicillin and its related substances. *J. Pharm. Biomed. Anal.*, 14 (1996) 631-639.
- 902 Zhu, Z., Xuan, D.-W. and Nightingale, C.H.: Modification of an ion-paired high pressure liquid chromatography for ticarcillin in human serum. *Zhongguo Yaoli Xuebao*, 17 (1996) 395-398; C.A., 125 (1996) 211675m.
- 903 Zhuo, H., Cao, W. and Lin, S.: (Determination of cefaclor in plasma by HPLC and its pharmacokinetics). *Zhongguo Yiyuan Yaoxue Zazhi*, 16 (1996) 99-101; C.A., 125 (1996) 184768k.

For additional information see C.A.:
125 (1996) 67969r, 96294x.

See also 698, 977.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 904 Chatani, Y., Ohfuki, M., Chikamoto, T. and Adachi, T.: (Simultaneous determination of fungicides, organophosphorus pesticides, pyrethroid pesticides in citrus fruits. (II). Determination of thiabendazole, benomyl, imazalil and 2,4-D). *Kyoto-fu Hoken Kankyo Kenkyusho Nenpo*, 40 (1995) 12-17; C.A., 125 (1996) 32154u.

- 905 Slobodnik, J., Öztezkizan, Ö., Lingeman, H. and Brinkman, U.A.T.: Solid-phase extraction of polar pesticides from environmental water samples on graphitized carbon and Empore-activated carbon disks and on-line coupling to octadecyl-bonded silica analytical columns. *J. Chromatogr. A*, 750 (1996) 227-238.
- 906 Slobodnik, J., Hogenboom, A.C., Vreuls, J.J., Rontree, J.A., van Baar, B.L.M., Niessen, W.M.A. and Brinkman, U.A.T.: Trace-level determination of pesticide residues using on-line solid-phase extraction-column liquid chromatography with atmospheric pressure ionization mass spectrometric and tandem mass spectrometric detection. *J. Chromatogr. A*, 741 (1996) 59-74.
- 907 Vreuls, J.J., Swen, R.J.J., Goudriaan, V.P., Kerkhoff, M.A.T., Jongenotter, G.A. and Brinkman, U.A.T.: Automated on-line gel permeation chromatography-gas chromatography for the determination of organophosphorus pesticides in olive oil. *J. Chromatogr. A*, 750 (1996) 275-286.
- 908 Wells, M.J.M. and Stearman, G.K.: Coordinating supercritical fluid and solid-phase extraction with chromatographic and immunoassay analysis of herbicides. *ACS Symp. Ser.*, 630(Herbicide Metabolites in Surface Water and Groundwater) (1996) 18-33; C.A., 125 (1996) 79214d.

For additional information see C.A.:
125 (1996) 188308h.

See also 65.

29b. Chlorinated insecticides

- 909 Bahri, U.C. and Jain, C.L.: Simultaneous determination of DDT (dichloro diphenyl trichloro ethane) and monochloro benzene in water and waste water from insecticide manufacturing industry by high performance liquid chromatography. *Pollut. Res.*, 15 (1996) 67-70; C.A., 125 (1996) 95256z.
- 910 Bruner, F., Berloni, A. and Palma, P.: Determination of acidic chlorinated pesticides in water: comparison of official EPA method 515.1 and liquid-solid extraction HPLC/UV and HPLC/PBMS analyses. *Chromatographia*, 43 (1996) 279-284.
- 911 Hormazábal, B. and Yndestad, M.: High-performance liquid chromatographic determination of teflubenzuron and diflubenzuron in fish tissues of farmed fish for residue studies. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2603-2614.
- 912 Parrilla, P. and Martinez Vidal, J.L.: HPLC determination of pesticides in green bean samples after SPE clean-up. *Chromatographia*, 43 (1996) 265-270.

See also 204, 1148, 1161.

29c. Phosphorus insecticides

- 913 Delhom, N., Balant, Y., Ader, J.C. and Lattes, A.: High performance liquid chromatographic determination of diazinon in polymeric matrix. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1735-1743.

- 914 Lacorte, S. and Barceló, D.: Determination of parts per trillion levels of organophosphorus pesticides in groundwater by automated on-line liquid-solid extraction followed by liquid chromatography/atmospheric pressure chemical ionization mass spectrometry using positive and negative ion modes of operation. *Anal. Chem.*, 68 (1996) 2464-2470.

See also 907, 912.

29d. Carbamates

- 915 Conaway, J.E., Jr.: Determination of methomyl in insecticidal formulations by reversed-phase liquid chromatography: collaborative study. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 941-948.

- 916 Di Corcia, A., Crescenzi, C., Lagana, A. and Sebastiani, E.: Evaluation of a method based on liquid chromatography/electrospray/mass spectrometry for analyzing carbamate insecticides in fruits and vegetables. *J. Agric. Food Chem.*, 44 (1996) 1930-1938; C.A., 125 (1996) 32178e.

- 917 Galeano Díaz, T., Guiberteau, A., Salinas, F. and Ortíz, J.M.: Rapid and sensitive determination of carbaryl, carbofuran and fenobucarb by liquid chromatography with electrochemical detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2681-2690.

- 918 Slobodnik, J., Hoekstra-Oussoren, S.J.F., Jager, M.E., Honing, M., van Baar, B.L.M. and Brinkman, U.A.T.: On-line solid-phase extraction-liquid chromatography-particle beam mass spectrometry and gas chromatography-mass spectrometry of carbamate pesticides. *Analyst (Cambridge)*, 121 (1996) 1327-1334.

See also 912.

29e. Herbicides

- 919 Baskaran, S., Lauren, D.R. and Holland, P.T.: High-performance liquid chromatographic determination of flumetsulam, a newly developed sulfonamide herbicide in soil. *J. Chromatogr. A*, 746 (1996) 25-30.

- 920 Cai, Z., Monson, S.J. and Spalding, R.F.: Determination of atrazine and hydroxyatrazine in agricultural runoff waters by liquid chromatography and fast atom bombardment-high-resolution-mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 929-935.

- 921 Field, J.A. and Monohan, K.: Chlorinated acid herbicides in water by strong anion-exchange disk extraction and in-vial elution and derivatization. *J. Chromatogr. A*, 741 (1996) 85-90.

- 922 Gennaro, M.C., Giacosa, D., Abrigo, C. and Simonetti, S.: Separation of a mixture of hydrophilic and hydrophobic herbicides by ion-interaction reversed-phase HPLC. *J. Chromatogr. Sci.*, 34 (1996) 334-340.

- 923 Gökmén, V. and Acar, J.: A study on the possibility of using HPLC for the determination of 2,4-D in tomatoes. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1917-1926.

- 924 Lang, D., Griegee, D., Grothusen, A., Saalfrank, R.W. and Böcker, R.H.: *In vitro* metabolism of atrazine, terbutylazine, ametryne, and terbutryne in rats, pigs, and humans. *Drug Metab. Disp.*, 24 (1996) 859-865.

- 925 Li, L.Y.T., Campbell, D.A., Bennett, P.K. and Henion, J.: Acceptance criteria for ultratrace HPLC - tandem mass spectrometry: qualitative and quantitative determination of sulfonylurea herbicides in soil. *Anal. Chem.*, 68 (1996) 3397-3404.

- 926 Sellero, I.S., Cruz, A., Fernández, P. and López-Rivadulla, M.: Liquid chromatographic method for simultaneous determination of paraquat and diquat in plasma, urine and vitreous humour. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2009-2024.

- 927 Sen, N.P. and Baddoo, P.A.: Determination of glyphosate as N-nitroso derivative by high performance liquid chromatography with chemiluminescence detection. *Int. J. Environ. Anal. Chem.*, 63 (1996) 107-117; C.A., 125 (1996) 84942u.

- 928 Stout, S.J., daCunha, A.R., Picard, G.L. and Safarpour, M.M.: Rapid, direct determination of imidazolinone herbicides in water at the 1 ppb level by liquid chromatography/electrospray ionization mass spectrometry and tandem mass spectrometry. *J. Agric. Food Chem.*, 44 (1996) 2182-2186; C.A., 125 (1996) 79229n.

See also 908, 912.

29f. Fungicides

- 929 Fernandez, C., Reviejo, A.J., Polo, L.M. and Pingarron, J.M.: HPLC-electrochemical detection with graphite-poly(tetrafluoroethylene) electrode determination of the fungicides thiram and disulfiram. *Talanta*, 43 (1996) 1341-1348; C.A., 125 (1996) 140814t.

- 930 Papadopoulou-Mourkidou, E., Patsias, J., Papadopoulos, G. and Galanis, C.: Liquid chromatographic determination of fentin acetate in fentin and Maneb formulations. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 829-832.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 931 Gagliardi, L., De Orsi, D., Cavazzuti, G., Multari, G. and Tonelli, D.: HPLC determination of Rhodamine B (C.I. 45170) in cosmetic products. *Chromatographia*, 43 (1996) 76-78.

- 932 Kai, S., Nikkawa, T., Takahashi, A., Koizumi, A., Hyoudou, Y., Suzuki, S. and Nakazawa, H.: (Preparation of sample solution for determination of acid coal-tar dyes in confectionery by the matrix solid-phase dispersion method). *Shokuhin Eiseigaku Zasshi*, 37 (1996) 146-150; C.A., 125 (1996) 113044k.

- 933 Kimoto, K., Gohda, R., Murayama, K., Santa, T., Fukushima, T., Homma, H. and Imai, K.: Sensitive detection of near-infrared fluorescent dyes using high-performance liquid chromatography with peroxyoxalate chemiluminescence detection system. *Biomed. Chromatogr.*, 10 (1996) 189-190; C.A., 125 (1996) 136802v.

- 934 Okamoto, T., Isozaki, A. and Nagashima, H.: (Determination of iodine in food colors by ion chromatography on ceramic carbon column). *Bunseki Kagaku*, 45 (1996) 717-721; C.A., 125 (1996) 113042h.

- 935 Rang, K., Isaksson, R. and Sandstroem, J.: Chromatographic enantiomer separation and circular dichroism spectra of chiral rhodanines. *J. Chem. Soc. Perkin Trans. 2*, (1996) 1493-1496; C.A., 125 (1996) 157353q.
- 936 Ueno, E., Nakashima, H., Takakura, K., Okazaki, K., Nagai, H., Oka, H., Ikai, Y., Hayakawa, J. and Ishikawa, N.: (Identification of non-permitted dye orange II used in a pickled vegetable). *Shokuhin Eiseigaku Zasshi*, 37 (1996) 226-230; C.A., 125 (1996) 274167t.

For additional information see C.A.:
125 (1996) 57368e, 103695j, 145046d.

See also 832.

30b. Chloroplast and other natural pigments

- 937 Brink, A., Triebkorn, S. and Schmidkunz, H.: (Column chromatography - separation of leaf pigments under elevated pressure with a self-made reversed phase in a school experiment). *Naturwiss. Unterr. Chem.*, 7 (1996) 16-20; C.A., 125 (1996) 246768w.
- 938 Croce, R., Breton, J. and Bassi, R.: Conformational changes induced by phosphorylation in the CP29 subunit of photosystem II. *Biochemistry*, 35 (1996) 11142-11148.
- 939 D'Ischia, M., Costantini, C. and Prota, G.: Lipofuscin-like pigments by autoxidation of polyunsaturated fatty acids in the presence of amine neurotransmitters: the role of malondialdehyde. *Biochim. Biophys. Acta*, 1290 (1996) 319-326.
- 940 Inoue, H., Nonomura, Y., Ashizawa, A., Tajima, I., Nomura, N., Nakamura, N. and Yoshioka, N.: Preparation and separation of metallobacteriochlorophyll a by reversed-phase high-performance liquid chromatography. *Chromatographia*, 43 (1996) 361-364.
- 941 Jagannadham, M.V., Narayanan, K., Rao, C.N. and Shivaji, S.: *In vivo* characteristics and localisation of carotenoid pigments in psychotropic and mesophilic *Micrococcus roseus* using photoacoustic spectroscopy. *Biochem. Biophys. Res. Commun.*, 227 (1996) 221-226.
- 942 Kuntcheva, M.J. and Obretenov, T.D.: Isolation and characterization of melanoidins from beer. *Z. Lebensm.-Unters. Forsch.*, 202 (1996) 238-243; C.A., 125 (1996) 56538y.
- 943 Liu, J. and Chen, L.: (Separation and determination of capsaicin in red pepper pigment by reduced pressure column chromatography and spectrophotometry). *Huaxue Yanjiu Yu Yingyong*, 7 (1995) 318-320; C.A., 125 (1996) 56515p.
- 944 Noell, G.N. and Kalinowski, H.-O.: Identification of retinol isomers by high-performance liquid chromatography not unequivocal up to now. *Vision Res.*, 36 (1996) 1887-1892; C.A., 125 (1996) 162445a.
- 945 Rouseff, R., Raley, L. and Hofsmoer, H.-J.: Application of diode array detection with a C-30 reversed phase column for the separation and identification of saponified orange juice carotenoids. *J. Agric. Food Chem.*, 44 (1996) 2176-2181; C.A., 125 (1996) 84939y.
- 946 Takaichi, S., Sandmann, G., Schnurr, G., Satomi, Y., Suzuki, A. and Misawa, N.: The carotenoid 7,8-dihydro- ψ end group can be cyclized by the lycopene cyclases from the bacterium *Erwinia uredovora* and the higher plant *Capsicum annuum*. *Eur. J. Biochem.*, 241 (1996) 291-296.

- 947 Van Breemen, R.B. and Huang, C.-R.: High-performance liquid chromatography-electrospray mass spectrometry of retinoids. *FASER J.*, 10 (1996) 1098-1101; C.A., 125 (1996) 137172b.
- 948 Van Breemen, R.B., Huang, C.-R., Tan, Y., Sander, L.C. and Schilling, A.B.: Liquid chromatography/mass spectrometry of carotenoids using atmospheric pressure chemical ionization. *J. Mass Spectrom.*, 31 (1996) 975-981; C.A., 125 (1996) 273905v.

See also 844, 1167, 1179.

31. PLASTICS AND THEIR INTERMEDIATES

- 950 Belenkii, B.G., Gankina, E.S., Kasalainen, G.E. and Tennikov, M.B.: Nonstandard methods based on size exclusion chromatography principles. "Critical" chromatography, high-performance membrane chromatography, thin-layer chromatography, and microbore size exclusion chromatography. *ACS Symp. Ser.*, 635 (1996) 274-326; C.A., 125 (1996) 59489n - a review with 55 refs.
- 951 Bruna, J.M.: (New analytical techniques for polymers. Part III. Characterization in size-exclusion chromatography (SEC) with multiple detectors). *Rev. Plast. Mod.*, 71 (1996) 493-497; C.A., 125 (1996) 143869a.
- 952 Chance, R.R., Baniukiewicz, S.P., Mintz, D., Ver Strate, G. and Hadjichristidis, N.: Characterization of low-molecular-weight polymers: failure of universal calibration in size exclusion chromatography. *Int. J. Polym. Anal. Charact.*, 1 (1995) 3-34; C.A., 125 (1996) 115706b.
- 953 Daoust, D., Bebelman, S., Godard, P., Coisme, J.M. and Strazielle, C.: Molecular characterization of poly(*p*-phenylene-sulphide) (PPS) from size exclusion chromatography of a modified PPS and dilute solution properties. *Polymer*, 37 (1996) 3879-3888; C.A., 125 (1996) 169275d.
- 954 Dayal, U., Mathur, A.B., Jain, R.C. and Shashikant, B.R.: High temperature SEC calibration for polypropylene using a broad dispersity standard and evaluation of its Mark-Houwink constants. *Macromol. Rep.*, A33 (1996) 243-250; C.A., 125 (1996) 115712a.
- 955 Dias, M.L., Azuma, C. and Mano, E.B.: Size exclusion behavior of polymers in amide solvents. Part 4. Elution characteristics of acrylonitrile polymers in *N,N*-dimethylacetamide. *Polym. Bull. (Berlin)*, 36 (1996) 729-735; C.A., 125 (1996) 59550g.
- 956 Hernandez-Barajas, J. and Hunkeler, D.: Copolymers of acrylamide and quaternary ammonium cationic monomers: characterization by HPLC and copolymer composition control. *Ber. Bunsen-Ges.*, 100 (1996) 723-729; C.A., 125 (1996) 87674u.
- 957 Just, U. and Krueger, R.-P.: Investigations of chemical heterogeneity in poly(dimethyl)siloxanes) using SFC, HPLC, and MALDI-MS. In: Auner, N. and Weis, J. (Editors), *Organosilicon Chem. II*, 2nd 1994, VCH, Weinheim, 1996, pp. 625-632; C.A., 125 (1996) 144773b.
- 958 Krueger, R.P., Much, H. and Schulz, G.: (Determination of polymer heterogeneity by coupling of HPLC and MALDI-TOF-MS). *GIT Fachz. Lab.*, 40 (1996) 398-402; C.A., 125 (1996) 34520j.

- 959 Lochmuller, C.H., Jaing, C., Liu, Q., Antonucci, V. and Elomaa, M.: High-performance liquid chromatography of polymers: retention mechanisms and recent advances. *Crit. Rev. Anal. Chem.*, 26 (1996) 29-59; *C.A.*, 125 (1996) 115624y - a review with 76 refs.
- 960 Meehan, E., Reid, S., Samois, E. and Dawkins, J.V.: Liquid chromatography techniques for characterizing poly(vinyl alcohol). *Macromol. Symp.*, 110 (1996) 65-80; *C.A.*, 125 (1996) 169137k.
- 961 Meehan, E., Reid, S.P., Warner, F.P., Patterson, M.D. and Dawkins, J.V.: Compositional heterogeneity of poly(vinyl alcohol): characterization by liquid chromatography techniques. *ACS Symp. Ser.*, 635 (1996) 262-273; *C.A.*, 125 (1996) 34544v.
- 962 Mendichi, R., Rizzo, V., Gigli, M. and Giacometti Schieroni, A.: Molecular characterization of polymeric antitumor drug carriers by size exclusion chromatography and universal calibration. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1591-1605.
- 963 Mori, S.: Compositional characterization of styrene copolymers by non-exclusion liquid chromatography. *Macromol. Symp.*, 110 (1996) 87-95; *C.A.*, 125 (1996) 143754j.
- 964 Mori, S. and Oosaki, T.: Comparison of secondary effects in aqueous size exclusion chromatography between sodium poly(styrenesulphonate) compounds of different sulphonations. Preliminary study. *ACS Symp. Ser.*, 635 (1996) 347-350; *C.A.*, 125 (1996) 115703y.
- 965 Netopilik, M., Bohdanecký, M. and Kratochvíl, P.: Influence of chemical heterogeneity of copolymers on the separation process in size-exclusion chromatography. *Macromolecules*, 29 (1996) 6023-6030; *C.A.*, 125 (1996) 143747j.
- 966 Philipsen, H.J.A., Klumperman, B. and German, A.L.: Characterization of low-molar-mass polymers by gradient polymer elution chromatography. I. Practical parameters and applications of the analysis of polyester resins under reversed phase conditions. *J. Chromatogr. A*, 746 (1996) 211-224.
- 967 Radke, W. and Mueller, A.H.E.: Molecular weights of copolymers obtained by gel permeation chromatography-light scattering. *ACS Symp. Ser.*, 635 (1996) 2-6; *C.A.*, 125 (1996) 59599e.
- 968 Rissler, K.: High-performance liquid chromatography and detection of polyethers and their mono(carboxy)alkyl and -arylalkyl substituted derivatives. *J. Chromatogr. A*, 742 (1996) 1-54 - a review with 187 refs.
- 969 Sato, H., Ogino, K., Darwint, T. and Kiyokawa, I.: Separation of copolymers by adsorption liquid chromatography. *Macromol. Symp.*, 110 (1996) 177-189; *C.A.*, 125 (1996) 143758p.
- 970 Trathnigg, B., Thamer, D. and Yan, X.: Analysis of polyethers by isocratic HPLC with universal detectors. IV. Comparison of size exclusion chromatography and liquid adsorption chromatography for the analysis of poly(propylene glycols). *Int. J. Polym. Anal. Charact.*, 1 (1995) 35-47; *C.A.*, 125 (1996) 87683w.
- 971 Trathnigg, B. and Maier, B.: Characterization of polyethers using different methods. SFC, LAC and SEC with different detectors, and MALDI-TOF-MS. *Macromol. Symp.*, 110 (1996) 231-240; *C.A.*, 125 (1996) 143760h.
- 972 Vilenchik, L.Z., Barman, S. and Pathak, C.P.: Method for studying micelle formation using size exclusion chromatography light scattering and UV spectroscopy. *ACS Symp. Ser.*, 635 (1996) 328-346; *C.A.*, 125 (1996) 59547m.
- See also 56, 113, 115, 135.
- ### 32. DRUG ANALYSIS
- #### 32a. Drug analysis, general techniques
- 973 Anderson, D.J.: Direct-injection HPLC: chromatography of biofluids without protein removal. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 183-193; *C.A.*, 125 (1996) 241979f - a review with 33 refs.
- 974 Berridge, J.C.: Impurities in drug substances and drug products: new approaches to quantification and qualification. *J. Pharm. Biomed. Anal.*, 14 (1995) 7-12.
- 975 Braggio, S. and Barnaby, R.J.: An HPLC assay directly on plasma using an ISRP. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 158; *C.A.*, 125 (1996) 211678q.
- 976 Crowe, E.A., Roberts, J.K. and Smith, R.J.: ^1H And ^{19}F LC/NMR: application to the identification of impurities in compounds of pharmaceutical interest. *Pharm. Sci.*, 1 (1995) 103-105; *C.A.*, 125 (1996) 177522x.
- 977 Cserhati, T. and Oros, G.: Determination of hydrophobicity parameters of antibiotics by reversed-phase chromatography. The effect of support. *Biomed. Chromatogr.*, 10 (1996) 117-121; *C.A.*, 125 (1996) 48371y.
- 978 Earley, R.A. and Tini, L.P.: Versatile multidimensional chromatographic system for drug discovery as exemplified by the analysis of a nonpeptidic inhibitor of human leukocyte elastase. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2527-2540.
- 979 Ecker, G., Mohr, E., Geyer, R. and Fleischhacker, W.: Enantioseparation of propafenone-type modulators of multidrug resistance on cyclodextrin based chiral stationary phases. *Sci. Pharm.*, 64 (1996) 1-11; *C.A.*, 125 (1996) 114414t.
- 980 Goldthwaite, C.A., Jr., Hsieh, F.Y., Womble, S.W., Nobos, B.J., Blair, I.A., Klunk, L.J. and Mayol, R.F.: Liquid chromatography/chemical reaction interface mass spectrometry as an alternative to radioisotopes for quantitative drug metabolism studies. *Anal. Chem.*, 68 (1996) 2996-3001.
- 981 Görög, S., Bihari, M., Csizer, E., Dravetz, F., Gazdag, M. and Herenyi, B.: Estimation of impurity profiles of drugs and related materials. Part 14: The role of HPLC/diode array UV spectroscopy in the identification of minor components (impurities, degradation products, metabolites) in various matrices. *J. Pharm. Biomed. Anal.*, 14 (1995) 85-92.
- 982 Grill, C.M., Kern, J.R. and Perrin, S.R.: Preparative separation of enantiomers. In: Olson, W.P. (Editor), *Sep. Technol.*, Interpharm Press, Buffalo Grove, 1995, pp. 231-269; *C.A.*, 125 (1996) 177091f - a review with refs.
- 983 Haginaka, J.: (Development of HPLC packing materials for drug analysis and their applications). *Yakugaku Zasshi*, 116 (1996) 473-490; *C.A.*, 125 (1996) 104048n - a review with 46 refs.
- 984 Heizmann, P., Gora, R., Korner, J. and Maier, S.: Hemolytic plasma: unexpected problems with a validated assay in routine analysis. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 304-311; *C.A.*, 125 (1996) 242060m.

- 985 Kirkland, J.J.: Practical method development strategy for reversed-phase HPLC of ionizable compounds. *LC-GC*, 14 (1996) 486-500; C.A., 125 (1996) 41919z - a review with 27 refs.
- 986 Lough, W.J., Groves, S.-J., Law, B., Maltas, J., Mills, M.J. and Saeed, M.: Drug bioanalysis made easy? *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 142-146; C.A., 125 (1996) 237408b.
- 987 MacDonald, A.: An automated system for tablet dissolution sampling and LC analysis. *Am. Lab. (Shelton)*, 28 (1996) 29-32; C.A., 125 (1996) 41920t - a review with 2 refs.
- 988 Oliveros, L., Franco, P., Senso, A. and Minguillon, C.: Bonded cellulose-derived HPLC chiral stationary phases; examples of resolution of some racemic drugs. *Am. Biotechnol. Lab.*, 14 (1996) 34-35; C.A., 125 (1996) 151298d.
- 989 Qin, Z.-I.: Advances in biopharmaceutical analysis in the People's Republic of China: 1993-1995. *J. Pharm. Biomed. Anal.*, 14 (1996) 1395-1403 - a review with 65 refs.
- 990 Schwarz, M.A., Neubert, R.H.H. and Dongowski, G.: Characterization of interactions between bile salts and drugs by micellar electrokinetic capillary chromatography. *Pharm. Res.*, 13 (1996) 1174-1180; C.A., 125 (1996) 184744z.
- 991 Shea, J.A., Shamrock, W.F., Abboud, C.A., Woodeshick, R.W., Nguyen, L.-Q.S., Rubino, J.T. and Segretario, J.: Validation of cleaning procedures for highly potent drugs. I. Losoxanthrone. *Pharm. Dev. Technol.*, 1 (1996) 69-75; C.A., 125 (1996) 67666q.
- 992 Sidelmann, U., Hansen, S.H., Gavaghan, C., Nicholls, A.W., Carless, H.A.J., Lindon, J.C., Wilson, I.D. and Nicholson, J.K.: Development of a simple liquid chromatographic method for the separation of mixtures of positional isomers and anomers of synthetic 2-, 3- and 4-fluorobenzoic acid glucuronides formed via acyl migration reactions. *J. Chromatogr. B*, 685 (1996) 113-122.
- 993 Thoma, K. and Kübler, N.: Einfluss der Wellenlänge auf die Photozersetzung von Arzneistoffen. *Pharmazie*, 51 (1996) 660-664.
- 994 Van der Hoeven, R.A.M., Reeuwijk, H.J.E.M., Tjaden, U.R. and van der Greef, J.: Analysis of quaternary ammonium drugs by thermospray liquid chromatography-mass spectrometry using a resin-based stationary phase. *J. Chromatogr. A*, 741 (1996) 75-84.
- 995 Volford, O., Vamos, J. and Takacs, M.: The significance of different chromatographic methods in the European pharmacopeia. *Int. J. Pharm.*, 137 (1996) 111-115; C.A., 125 (1996) 96254j.
- 996 Welsh, W.J., Lin, W., Tersigni, S.H., Collantes, E., Duta, R., Carey, M.S., Zielinski, W.L., Brower, J., Spencer, J.A. and Layloff, T.P.: Pharmaceutical fingerprinting: evaluation of neural networks and chemometric techniques for distinguishing among same-product manufacturers. *Anal. Chem.*, 68 (1996) 3473-3482.
- 997 Zhang, Y., Zeng, C.-M., Li, Y.-M., Hjertén, S. and Lundahl, P.: Immobilized liposome chromatography of drugs on capillary continuous beds for model analysis of drug-membrane interactions. *J. Chromatogr. A*, 749 (1996) 13-18.
- 32b. *Antirheumatics and antiinflammatory drugs*
- 998 Calucci, G., Mazzeo, P. and Palumbo, G.: Simultaneous determination of lomafloxacin, fenbufen and felbinac in human plasma using high performance liquid chromatography. *Chromatographia*, 43 (1996) 261-264.
- 999 Carlucci, G. and Mazzeo, P.: High-performance liquid chromatographic determination of 1-benzyl-1H-indazol-3-ol in benzodamine in pharmaceutical formulations. *J. Pharm. Biomed. Anal.*, 14 (1996) 655-657.
- 1000 Geisslinger, G., Bocker, R. and Levy, M.: High-performance liquid-chromatographic analysis of dipyrone metabolites to study their formation in human liver microsomes. *Pharm. Res.*, 13 (1996) 1272-1275; C.A., 125 (1996) 157646n.
- 1001 Gilar, M., Tesarova, E. and Deyl, Z.: (Influence of mobile phase composition on retention and enantioseparation of 2-arylpropanoic acids in HPLC on a β -cyclodextrin stationary phase). *Chem. Listy*, 90 (1996) 461-466; C.A., 125 (1996) 123934q.
- 1002 Herber, B., Bueschges, R. and Spahn-Langguth, H.: Fluorescent carbazoles as chiral derivatizing agents in the analysis of drug enantiomers: isocyanates deriving from carprofen and N-methylcarprofen. In: Daldrup, T. and Musshoff, F. (Editors), *Chem. Spuren Verkehrsunfaellen, Beifr. Symp. Ges. Toxikol. Forensische Chem.*, Verlag Dr. Dieter Helm, Heppenheim, 1995, pp. 136-141; C.A., 125 (1996) 96318h.
- 1003 Klimes, J., Sochor, J. and Sedlacek, J.: Solid-phase extraction and HPLC analysis of kebuzone and its metabolites in blood. *Pharmazie*, 51 (1996) 651-653.
- 1004 Lau, Y.Y.: Determination of ibuprofen enantiomers in human plasma by derivatization and high performance liquid chromatography with fluorescence detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2143-2153.
- 1005 Lenz, E.M., Greatbanks, D., Wilson, I.D., Spraul, M., Hofmann, M., Troke, J., Lindon, J.C. and Nicholson, J.K.: Direct characterization of drug glucuronide isomers in human urine by HPLC-NMR spectroscopy: application to the positional isomers of 6,11-dihydro-11-oxodibenzo[b,e]oxepin-2-acetic acid glucuronide. *Anal. Chem.*, 68 (1996) 2832-2837.
- 1006 Li, Y.N. (Benny), Tattam, B., Brown, K.F. and Seale, J.P.: Determination of epimers 22R and 22S of budesonide in human plasma by high-performance liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *J. Chromatogr. B*, 683 (1996) 259-268.
- 1007 Lin, C., Hsieh, J.Y.-K., Matuszewski, K. and Dobrinska, M.R.: An automated sample preparation and high performance liquid chromatographic method for the determination of MK-591, a novel leukotriene biosynthesis inhibitor, in human plasma. *J. Pharm. Biomed. Anal.*, 14 (1996) 1601-1608.
- 1008 McGurk, K.A., Remmel, R.P., Hosagrahara, V.P., Tosh, D. and Burchell, B.: Reactivity of mefenamic acid 1-O-acyl glucuronide with proteins *in vitro* and *ex vivo*. *Drug Metab. Disp.*, 24 (1996) 842-849.
- 1009 Palumbo, G., Carlucci, G., Mazzeo, P., Frieri, G., Pimpo, M.T. and Fanini, D.: Simultaneous determination of 5-aminosalicylic acid, acetyl-5-aminosalicylic acid and 2,5-dihydroxybenzoic acid in endoscopic intestinal biopsy samples in humans by high-performance liquid chromatography with electrochemical detection. *J. Pharm. Biomed. Anal.*, 14 (1995) 175-180.

For additional information see C.A.:

125 (1996) 96274r, 96317g.

See also 142, 144, 155, 169, 180, 214, 281, 431, 1002, 1058, 1140, 1240.

- 1010 Parimoo, P., Bharathi, A. and Atul, B.V.: Determination of propyphenazone and ketoprofen in drug preparation by high performance liquid chromatography. *East Pharm.*, 39 (1996) 131-132; *C.A.*, 125 (1996) 96340j.
- 1011 Ray, G.F., Lanman, R.C., Fu, C.J., Paranka, N.S., Pamukcu, R. and Wheeler, S.C.: Determination of FGN-1 (an active metabolite of sulindac) in human plasma, urine, and feces by HPLC. *J. Pharm. Biomed. Anal.*, 14 (1995) 213-220.
- 1012 Tan, S.C., Stevens, N., de Biasi, V. and Hutt, A.J.: Chiral-HPLC resolution of carboxyibuprofen, a major urinary metabolite of ibuprofen. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 154-157; *C.A.*, 125 (1996) 211677p.
- 1013 Taylor, M.R., Westwood, S.A., Woodward, K. and Perrett, D.: Analysis of aspirin metabolites in urine and plasma by MECC with direct sample introduction. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 288-298; *C.A.*, 125 (1996) 237414a.
- See also 1020, 1047.
- 32c. *Autonomic and cardiovascular drugs*
- 1014 Booth, T.D. and Wainer, I.W.: Mechanistic investigation into the enantioselective separation of mexiletine and related compounds, chromatographed on an amylose tris(3,5-dimethylphenylcarbamate) chiral stationary phase. *J. Chromatogr. A*, 741 (1996) 205-211.
- 1015 Caccamese, S., Chillemi, R. and Principato, G.: Fluorenone 1,4-dihydropyridine derivatives with cardiac depressant activity: Enantiomeric separation by chiral HPLC and conformational aspects. *Chirality*, 8 (1996) 281-290; *C.A.*, 125 (1996) 123854p.
- 1016 Ceccato, A., Chiap, P., Hubert, P., Toussaint, B. and Crommen, J.: Automated determination of verapamil and norverapamil in human plasma with on-line coupling of dialysis to high-performance liquid chromatography and fluorometric detection. *J. Chromatogr. A*, 750 (1996) 351-360.
- 1017 Chen, L.-S., Fujitaki, J.M. and Dixon, R.: A sensitive assay for the aminoimidazole-containing drug GP531 in plasma using liquid chromatography with amperometric electrochemical detection: a new class of electroactive compounds. *J. Pharm. Biomed. Anal.*, 14 (1996) 1535-1538.
- 1018 Cornelius, K.E., Fadayel, G.M., Baron, B.M., Schmidt, C.J., Haeghe, K.D. and Chen, T.-M.: Analysis of (R)-4-oxo-5-phosphononorvaline (MDL 100,453) in rat plasma and brain dialysate using liquid chromatography after derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate. *J. Pharm. Biomed. Anal.*, 14 (1995) 143-150.
- 1019 Ishii, K., Wakamoto, S., Nakai, H. and Sato, T.: Assay of four stereoisomers of desacetyl diltiazem hydrochloride. Application to *in vitro* chiral inversion studies. *Chromatographia*, 43 (1996) 413-418.
- 1020 Kukushima, T., Santa, T., Homma, H. and Imai, K.: (Enantioselective separation of drugs on chiral stationary phases in HPLC. Effect of mobile phase composition on the enantioselective separation and the detection). *Kuromatogurafu*, 17 (1996) 152-153; *C.A.*, 125 (1996) 123881v.
- 1021 Maguregui, M.I., Alonso, R.M. and Jiménez, R.M.: A rapid quantitative analysis of the β -blocker timolol in human urine by HPLC-electrochemical detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1643-1652.
- 1022 Morris, R.G., Saccoia, N.C. and Jones, T.E.: Modified liquid chromatographic assay for diltiazem and metabolites in human plasma. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2385-2394.
- 1023 Nakakura, M., Kato, Y., Hayakawa, E., Ito, K. and Kuroda, T.: Electrochemical stability of desmopressin acetate during iontophoresis. *Chem. Pharm. Bull.*, 44 (1996) 1238-1241.
- 1024 Nayak, V.G., Belapure, S.G., Gaitonde, C.D. and Sule, A.A.: Determination of salmeterol in metered-dose and dry-powder inhalers by reversed-phase high performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 511-513.
- 1025 Perez-Martinez, I., Sagrado, S. and Medina-Hernández, M.J.: Chromatographic determination of caffeine in pharmaceutical formulations using micellar mobile phases. *Chromatographia*, 43 (1996) 149-152.
- 1026 Shekher, A. and Sankaranarayanan, A.: Reversed-phase high performance liquid chromatography (HPLC) assay of verapamil in plasma. *Indian Drugs*, 33 (1996) 259-260; *C.A.*, 125 (1996) 264867g.
- 1027 Tsai, T.H., Kuo, K.W., Cheng, F.C. and Chen, C.F.: Determination of scopolamine in rat plasma by liquid chromatography and its application to pharmacokinetics. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2649-2657.
- 1028 Tsai, T.H., Tsai, W.J., Chou, C.J. and Chen, C.F.: Inhibition by honokiol of collagen-induced platelet 5-HT release. *Mol. Pharmacol.*, 50 (1996) 326-333; *C.A.*, 125 (1996) 185357n.
- 1029 Van Vyncht, G., Preece, S., Gaspar, P., Maghuijn-Rogister, G. and DePauw, E.: Gas and liquid chromatography coupled to tandem mass spectrometry for the multiresidue analysis of β -agonists in biological matrices. *J. Chromatogr. A*, 750 (1996) 43-49.
- 1030 Verbesselt, R., Zugravu, A., Tjandramaga, T.B. and de Schepper, P.J.: Liquid chromatographic determination of total celiprolol or (S)-celiprolol and (R)-celiprolol simultaneously in human plasma. *J. Chromatogr. B*, 683 (1996) 231-236.
- 1031 Weindlmayr-Goettel, M., Weberhofer, G., Gilly, H. and Kress, H.G.: Determination of mivacurium in plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 685 (1996) 123-127.
- 1032 Williams, R.C., Alasdair, M.S., Fasone, V.L., Boucher, R.J. and Edwards, J.F.: Comparison of liquid chromatography, capillary electrophoresis and super-critical fluid chromatography in the determination of Losartan Potassium drug substance in Cozaar tablets. *J. Pharm. Biomed. Anal.*, 14 (1996) 1539-1546.
- 1033 Wolzman, S.J., Chen, J.-G., Weber, S.G. and Tolley, J.O.: Determination of the pharmaceutical peptide TP9201 by post-column reaction with copper(II) followed by electrochemical detection. *J. Pharm. Biomed. Anal.*, 14 (1995) 155-164.
- 1034 Yamamoto, Y., Sakai, H., Tanaka, H., Hayakawa, H., Mai, M., Segawa, K. and Nakashima, Y.: (Quantitative determination of T-588 and its metabolites in human serum by LC/ESI/MS). *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 21 (1996) 223-226; *C.A.*, 125 (1996) 264873f.

- 1035 Yasuda, T., Tanaka, M. and Iba, K.: Quantitative determination of amlodipine in serum by liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry. *J. Mass Spectrom.*, 31 (1996) 879-884; C.A., 125 (1996) 184735x.

For additional information see C.A.:
125 (1996) 48249q, 67952e, 151290v.

See also 424.

32d. Central nervous system drugs

- 1036 Abdel-Moety, E.M., Al-Rashood, K.A., Rauf, A. and Khattab, N.A.: Photostability-indicating HPLC method for determination of trifluoperazine in bulk form and pharmaceutical formulations. *J. Pharm. Biomed. Anal.*, 14 (1996) 1639-1644.
- 1037 Ackermann, B.L., Gillespie, T.A., Regg, B.T., Austin, K.F. and Coutant, J.E.: Application of packed capillary liquid chromatography/mass spectrometry with electrospray ionization to the study of human biotransformation of the anti-emetic drug dolasetron. *J. Mass Spectrom.*, 31 (1996) 681-689; C.A., 125 (1996) 104163w.
- 1038 Akerman, K.K., Jolkonen, J., Parviainen, M. and Penttilä, I.: Analysis of low-dose benzodiazepines by HPLC with automated solid-phase extraction. *Clin. Chem. (Washington)*, 42 (1996) 1412-1416.
- 1039 Bernal, J.L., Del Nozal, M.J., Velsco, H. and Toribio, L.: HPLC versus SFC for the determination of salbutamol sulphate and its impurities in pharmaceuticals. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1579-1589.
- 1040 Cassel, G.E. and Fosbraey, P.: Measurement of the oxime HI-6 after peripheral administration in tandem with neurotransmitter levels in striatal dialyzates: effects of soman intoxication. *J. Pharmacol. Toxicol. Methods*, 35 (1996) 159-166; C.A., 125 (1996) 48255p.
- 1041 Chiu, J.A. and Franklin, R.B.: Analysis and pharmacokinetics of olanzapine (LY170053) and two metabolites in rat plasma using reversed-phase HPLC with electrochemical detection. *J. Pharm. Biomed. Anal.*, 14 (1996) 609-615.
- 1042 Chollet, D., Castella, E., Combe, P. and Arnera, V.: High-speed liquid chromatographic method for the monitoring of carbamazepine and its active metabolite, carbamazepine-10,11-epoxide, in human plasma. *J. Chromatogr. B*, 683 (1996) 237-243.
- 1043 De Orsi, D., Gagliardi, L. and Tonelli, D.: High performance liquid chromatographic determination of prozapine in pharmaceutical formulations. *J. Pharm. Biomed. Anal.*, 14 (1996) 1635-1638.
- 1044 Dessalles, M.C., Gasdeblay, S. and Mahuzier, G.: (An immunoassay kit for the determination of methadone in serum. Comparison with liquid chromatography). *Ann. Biol. Clin.*, 54 (1996) 203-209; C.A., 125 (1996) 160522z.
- 1045 Doheny, M.H., O'Connell, M.T. and Patsalos, P.N.: A high-performance liquid-chromatographic microanalytical procedure for the rapid estimation of piracetam in plasma or cerebrospinal fluid. *J. Pharm. Pharmacol.*, 48 (1996) 514-516; C.A., 125 (1996) 104121f.
- 1046 Dunne, M. and Andrew, P.: Fully automated assay for the determination of sumatriptan in human serum using solid-phase extraction and high-performance liquid chromatography with electrochemical detection. *J. Pharm. Biomed. Anal.*, 14 (1996) 721-726.
- 1047 Ensslin, H.K., Maurer, H.H., Gouzoulis, E., Hermle, L. and Kovar, K.-A.: Metabolism of racemic 3,4-methylenedioxymethamphetamine in humans. Isolation, identification, quantitation, and synthesis of urinary metabolites. *Drug Metab. Disp.*, 24 (1996) 813-820.
- 1048 Essien, H., Lai, S.J., Binder, S.R. and King, D.L.: Use of direct-probe mass spectrometry as a toxicology confirmation method for demoxepam in urine following high-performance liquid chromatography. *J. Chromatogr. B*, 683 (1996) 199-208.
- 1049 Feierman, D.E. and Lasker, J.M.: Metabolism of fentanyl, a synthetic opioid analgesic, by human liver microsomes. *Drug Metab. Disp.*, 24 (1996) 932-939.
- 1050 Feng, C., Liu, Y.-t. and Luo, Y.: (High-performance liquid-chromatographic analysis of tricyclic antidepressants in blood). *Sepu*, 14 (1996) 282-284; C.A., 125 (1996) 157643j.
- 1051 Forsslund, E., Eriksson, A.-S. and Beck, O.: Liquid chromatographic determination of plasma lamotrigine in pediatric samples. *J. Pharm. Biomed. Anal.*, 14 (1996) 755-758.
- 1052 Gavrilin, M.V., Kompantseva, E.V. and Airapetova, A.Yu.: (HPLC analysis of diazoline). *Khim.-Farm. Zh.*, 30 (1996) 50-51; C.A., 125 (1996) 67979u.
- 1053 Ghahramani, P. and Lennard, M.S.: Quantitative analysis of amitriptyline and nortriptyline in human plasma and liver microsomal preparations by high-performance liquid chromatography. *J. Chromatogr. B*, 685 (1996) 307-313.
- 1054 Haertter, S., Baier, D., Dingemanse, J., Ziegler, G. and Hiemke, C.: Automated determination of dextromethorphan and its main metabolites in human plasma by high-performance liquid chromatography and column switching. *Ther. Drug Monit.*, 18 (1996) 297-303; C.A., 125 (1996) 75214f.
- 1055 Halldin, M.M., Bredberg, E., Angelin, B., Arvidsson, T., Askemark, Y., Elofsson, S. and Widman, M.: Metabolism and excretion of ropivacaine in human. *Drug Metab. Disp.*, 24 (1996) 962-968.
- 1056 Hariharan, U., Hariharan, M., Naickar, J.S. and Tandon, R.: Determination of clozapine and its two major metabolites in human serum by liquid chromatography using ultraviolet detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2409-2417.
- 1057 Haupt, D.: Determination of citalopram enantiomers in human plasma by liquid chromatographic separation on a Chiral-AGP column. *J. Chromatogr. B*, 685 (1996) 299-305.
- 1058 Hinton, J.P., Pablo, J.C., Bjorge, S.M., Hoffman, K., Jennings, K. and Wright, S.S.: Three complementary liquid chromatographic methods for determination of the peptoid cholecystokinin-B antagonist, CI-988, in rat plasma. *J. Pharm. Biomed. Anal.*, 14 (1996) 815-824.
- 1059 Hoja, H., Marquet, P., Verneuil, B., Lofti, H., Dupuy, J.L., Dreyfuss, M.F. and Lachatre, G.: (Buprenorphine and norbuprenorphine determination in urine and serum by liquid chromatography-mass spectrometry with electrospray-type ionization). *Analisis*, 24 (1996) 104-107; C.A., 125 (1996) 123931m.

- 1060 Hui, Y.-h. and Marsh, K.C.: Sensitive detection of selected cholinergic channel activators derivatized with 7-fluoro-4-nitrobenzo-2-oxa-1,3-diazole. *J. Pharm. Biomed. Anal.*, 14 (1995) 131-142.
- 1061 Jiang, Q., Walton, N.Y., Gunawan, S. and Treiman, D.M.: High-performance liquid chromatographic determination of midazolam in rat brain. *J. Chromatogr. B*, 683 (1996) 276-280.
- 1062 Johnson, J.T., Oldham, S.W., Lantz, R.J. and DeLong, A.F.: High performance liquid chromatographic method for the determination of duloxetine and desmethyl duloxetine in human plasma. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1631-1641.
- 1063 King, D.T., Stewart, J.T. and Venkateshwaran, T.G.: HPLC determination of propofol-tiopental sodium and propofol-on-dasetron mixtures. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2285-2294.
- 1064 Kondo, T., Yoshida, K. and Tanayama, S.: Absorption of the anxiolytic pazinaclone in animals as a criterion for species selection for toxicity studies. *Arzneim.-Forsch.*, 46 (1996) 839-843.
- 1065 Kumar, K., Ballantyne, J.A. and Baker, A.B.: A sensitive assay for the simultaneous measurement of alfentanil and fentanyl in plasma. *J. Pharm. Biomed. Anal.*, 14 (1996) 667-673.
- 1066 Kumar, K., Ballantyne, J.A. and Baker, A.B.: Determination of droperidol in plasma by liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 1529-1533.
- 1067 Lau, Y.Y., Anderson, P.H. and Talaat, R.: High sensitivity high performance liquid chromatography electrospray tandem mass spectrometry determination of terfenadine in human plasma. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2669-2679.
- 1068 Lau, Y.Y., Hanson, G.D. and Ichhpuhani, N.: Determination of pramipexole (U-98,528) in human plasma and urine by high-performance liquid chromatography with electrochemical and ultraviolet detection. *J. Chromatogr. B*, 683 (1996) 217-223.
- 1069 Lau, Y.Y., Selenka, J.M., Hanson, G.D., Talaat, R. and Ichhpuhani, N.: Determination of pramipexole (U-98,528) in human plasma by high-performance liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry. *J. Chromatogr. B*, 683 (1996) 209-216.
- 1070 Lin, G., Chu, K.-W., Damani, L.A. and Hawes, E.M.: Identification of lactams as *in vitro* metabolites of piperidine-type phenothiazine antipsychotic drugs. *J. Pharm. Biomed. Anal.*, 14 (1996) 727-738.
- 1071 Lin, G., McKay, G. and Midha, K.K.: Characterization of metabolites of clozapine N-oxide in the rat by micro-column high performance liquid chromatography/mass spectrometry with electrospray interface. *J. Pharm. Biomed. Anal.*, 14 (1996) 1561-1577.
- 1072 Lin, S., Brower, P.L. and Winkle, D.D.: Chiral HPLC separations for process development of S-(+)-isobutyl GABA, a potential anti-epileptic agent. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2699-2708.
- 1073 Lugo, S.I. and Eddington, N.D.: Rapid method for the quantitation of mivacurium isomers in human and dog plasma by using liquid chromatography with fluorescence detection: Application to pharmacokinetic studies. *J. Pharm. Biomed. Anal.*, 14 (1996) 675-683.
- 1074 Moreau, S., Couder, P., Rubat, C., Albuisson, E. and Couquelet, J.: Synthesis of peripherally acting analgesic 3-arylpiperazinyl-5-benzyl-pyridazines. *Arzneim.-Forsch.*, 46 (1996) 800-805.
- 1075 Olsen, B.A. and Argentine, M.D.: HPLC method development for duloxetine hydrochloride using a combination of computer-based solvent strength optimization and solvent selectivity mixture design. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1993-2007.
- 1076 Pichini, S., Pacifici, R., Altieri, I., Pellegrini, M. and Zuccaro, P.: Stereoselective determination of fluoxetine and norfluoxetine enantiomers in plasma samples by high-performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1927-1935.
- 1077 Pleasance, S., Fraser, I.J., Jones, A.E., Allanson, J.A. and Sadra, P.: Determination of the 5-HT receptor agonist 311C90 in human plasma by LC-MS-MS. *Methodol. Surv. Bioanal. Drugs*, 24(Biofluid Assay for Peptide-Related and Other Drugs) (1996) 118-125; C.A., 125 (1996) 264835v.
- 1078 Stanke, F., Jourdin, N., Lauby, V. and Bessard, G.: Zopiclone and zolpidem quantification in human plasma by high performance liquid chromatography with photodiode-array detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2623-2633.
- 1079 Steger, P.J.K., Martinelli, E.F. and Muhlebach, S.F.: Stability of high-dose morphine chloride injection upon heat sterilization: Comparison of UV-spectroscopy and HPLC. *J. Chin. Pharm. Ther.*, 21 (1996) 73-78; C.A., 125 (1996) 123505a.
- 1080 Sultan, S.M. and El-Mubarak, A.H.: High-performance liquid chromatographic method for the separation and quantification of some psychotherapeutic benzodiazepines optimized by the modified simplex procedure. *Talanta*, 43 (1996) 569-576; C.A., 126 (1996) 41949j.
- 1081 Surapaneni, S. and Khalil, S.K.W.: Resolution of terfenadine enantiomers by reversed phase-high performance liquid chromatography using β -cyclodextrin as mobile phase additive. *J. Pharm. Biomed. Anal.*, 14 (1996) 1631-1634.
- 1082 Tatsuno, M., Nishikawa, M., Tsuchihashi, H., Igarashi, K., Kasuya, F. and Fukui, M.: (Thermospray and atmospheric pressure chemical ionization liquid chromatographic-mass spectrometric analysis of benzodiazepines and their metabolites). *Jpn. J. Toxicol. Environ. Health*, 42 (1996) 248-256; C.A., 125 (1996) 240436q.
- 1083 Torchin, C.D., McNeilly, P.J., Kapetanovic, I.M., Strong, J.-M. and Kupferberg, H.J.: Stereoselective metabolism of a new anticonvulsant candidate, losigamone, by human liver microsomes. *Drug Metab. Disp.*, 24 (1996) 1002-1008.
- 1084 Trocewicz, J., Suprynowicz, Z. and Markowicz, J.: Determination of diphrenan in urine by a supported liquid membrane technique and liquid chromatography-electrochemical detection. *J. Chromatogr. B*, 685 (1996) 129-134.
- 1085 Vielhascher, E., Spatzenegger, M., Mayrhofer, A., Klinger, P. and Jäger, W.: Metabolism of dextromethorphan in human liver microsomes: a rapid HPLC assay to monitor cytochrome P450 2D6 activity. *Pharmazie*, 51 (1996) 586-588.
- 1086 Wan, J., Xia, H. and Zhou, H.: (Determination of diazepam and demethyldiazepam in human plasma by high performance liquid chromatography). *Sepu*, 14 (1996) 303-305; C.A., 125 (1996) 184738a.
- 1087 Waters, R.K., Jee, R.D., Watt, R.A. and Moffat, A.C.: Correlation of physicochemical parameters of 28 barbiturates to their retention properties on an ODS reversed-phase HPLC system. *Pharm. Sci.*, 1 (1995) 593-596; C.A., 125 (1996) 151297c.

For additional information see C.A.:
125 (1996) 123886a.

See also 81, 434, 477, 797, 800, 824, 980, 1008, 1010, 1025, 1141, 1142, 1146.

32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 1088 Akhtar, M.J., Khan, S., Roy, I.M. and Jafri, I.A.: High performance liquid chromatographic determination of phenoxetol, methyl paraben, ethyl paraben, *n*-propyl paraben, iso-butyl paraben, *n*-butyl paraben and croconazole.HCl. *J. Pharm. Biomed. Anal.*, 14 (1996) 1609-1613.
- 1089 Alasandro, M.: Separation of diastereoisomers of DuP 105, a novel oxazolidinone antibacterial agent, by supercritical fluid chromatography on a Chiracel OD column. *J. Pharm. Biomed. Anal.*, 14 (1996) 807-814.
- 1090 Argekar, A.P., Kapadia, S.U. and Raj, S.V.: Simultaneous determination of norfloxacin and tinidazole in tablets by reverse phase high performance liquid chromatography (RP-HPLC). *Anal. Lett.*, 29 (1996) 1539-1549.
- 1091 Augustijns, P., Annaert, P., Adriaens, S., de Clercq, E. and Kinget, R.: High speed HPLC determination of bis(pivaloyloxymethyl)-PMEA and its degradation products, mono(POM)-PMEA and PMEA. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2271-2283.
- 1092 Campbell, D.A., Shah, V.R., Srinivas, N.R. and Shyu, W.C.: High-performance liquid chromatographic analysis of 2'-fluoro-2',3'-dideoxyadenosine and 2'-fluoro-2',3'-dideoxyinosine in dog plasma and urine. *J. Pharm. Sci.*, 85 (1996) 890-892.
- 1093 Cundy, K.C., Bidgood, A.M., Lynch, G., Shaw, J.-P., Griffin, L. and Lee, W.A.: Pharmacokinetics, bioavailability, metabolism, and tissue distribution of cidofovir (HPMPC) and cyclic HPMPC in rats. *Drug Metab. Disp.*, 24 (1996) 745-752.
- 1094 Dhaliwal, K. and Theobald, A.E.: The determination of paraben preservatives in syrup formulations. *Pharm. Sci.*, 1 (1995) 99-101; *C.A.*, 125 (1996) 177572p.
- 1095 Dusi, G., Curatolo, M., Fierro, A. and Faggionato, E.: Determination of the antiparasitic drug ivermectin in liver, muscle and fat tissue samples from swine, cattle, horses and sheep using HPLC with fluorescence detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1607-1616.
- 1096 Gehring, T.A., Rushing, L.G., Churchwell, M.I., Doerge, D.R., McErlane, K.M. and Thompson, H.C., Jr.: HPLC determination of sulfadiazine residues in coho salmon (*Oncorhynchus kisutch*) with confirmation by liquid chromatography with atmospheric pressure chemical ionization mass spectrometry. *J. Agric. Food Chem.*, 44 (1996) 3164-3169; *C.A.*, 125 (1996) 219862r.
- 1097 Hormazabal, V. and Yndestad, M.: Rapid assay for the determination of residues of amprolium and ethopabate in chicken meat by HPLC. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2517-2525.
- 1098 Iosifidou, E.G. and Haagsma, N.: High-performance liquid chromatographic determination of fenbendazole and its metabolites, sulphoxide and sulphone, in fish muscle tissue. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1819-1830.
- 1099 Jemal, M. and Hawthorne, D.: Determination of BMS-186318 in dog, rat and monkey plasma by liquid chromatography-ion-spray mass spectrometry. *J. Pharm. Biomed. Anal.*, 14 (1995) 43-48.
- 1100 Knaub, S.R., Chang, M.F., Lunte, C.E., Toppe, E.M. and Riley, C.M.: Automated analytical systems for drug development studies. Part IV. A microdialysis system to study the partitioning of lemofoxacin across an erythrocyte membrane *in vitro*. *J. Pharm. Biomed. Anal.*, 14 (1995) 121-129.
- 1101 Kobylinska, M., Kobylinska, K. and Sobik, B.: High-performance liquid chromatographic analysis for the determination of miconazole in human plasma using solid-phase extraction. *J. Chromatogr. B*, 685 (1996) 191-195.
- 1102 Kolawole, J.A., Mustapha, A., Abdul-Aguye, I. and Ochekpe, N.: Effect of temperature on chlorproguanil and proguanil hydrochloride solutions: a chemical stability study. *J. Pharm. Biomed. Anal.*, 14 (1996) 685-693.
- 1103 Krol, G.J., Beck, G.W. and Benham, T.: HPLC analysis of ciprofloxacin and ciprofloxacin metabolites in body fluids. *J. Pharm. Biomed. Anal.*, 14 (1995) 181-190.
- 1104 Lanchote, V.L., Bonato, P.S., Dreossi, S.A.C., Goncalves, P.V.B., Cesarino, E.J. and Bertucci, C.: High-performance liquid chromatographic determination of mexiletine enantiomers in plasma using direct and indirect enantioselective separations. *J. Chromatogr. B*, 685 (1996) 281-289.
- 1105 Liard, F., Ghiro, E., Paris, W. and Yaokim, C.: Determination of the HIV protease inhibitor BILA 2185 BS in rat plasma by liquid-liquid extraction and high performance liquid chromatography photodiode array detector. *J. Pharm. Biomed. Anal.*, 14 (1995) 151-154.
- 1106 Morris, D.M., Hussey, B.K., Guerin, S.L. and Selinger, K.A.: Determination of the novel sialic acid analog GG167 (GR121167X) in human urine by liquid chromatography: direct injection with column switching. *J. Pharm. Biomed. Anal.*, 14 (1995) 191-201.
- 1107 Polikandritou Lambros, M.P., Abbas, S.A. and Bourne, D.W.A.: New high-performance liquid chromatographic method for amphotericin B analysis using an internal standard. *J. Chromatogr. B*, 685 (1996) 135-140.
- 1108 Ricci, M.C. and Cross, R.F.: High performance liquid chromatographic analysis of sulphonamides and dihydrofolate reductase inhibitors. III. The effect of a competing base, and separations with an ion pairing agent. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2257-2270.
- 1109 Takeba, K., Itoh, T., Matsumoto, M., Nakazawa, H. and Tanabe, S.: Simultaneous determination of five fasciolicides in milk by liquid chromatography with electrochemical detection. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 848-852.
- 1110 Von Heeren, F., Tanner, R., Theurillat, R. and Thormann, W.: Determination of fluconazole in human plasma by micellar electrokinetic capillary chromatography with detection at 190 nm. *J. Chromatogr. A*, 745 (1996) 165-172.

See also 394, 815, 870, 877, 881, 895, 1181.

32f. Cytostatics

- 1111 Cummings, J., Meikle, I., Macpherson, J.S. and Smyth, J.F.: Determination of the novel topoisomerase I inhibitor NU/ICRF 505 and its major metabolite in plasma, tissue and tumour by high-performance liquid chromatography. *J. Chromatogr. B*, 685 (1996) 159-164.

- 1112 Huebert, N.D., Schwartz, J.-J., Zeidler, L., Schwach, V. and Haegle, K.D.: Simultaneous measurement of dolasetron and its major metabolite, MDL 74,156, in human plasma and urine. *J. Chromatogr. B*, 685 (1996) 291-297.
- 1113 Kakchenko, O.I., Golub, V.A. and Zavatskaja, I.V.: HPLC and GLC determination of residual solvents in busulphan. *J. Pharm. Biomed. Anal.*, 14 (1995) 107-111.
- 1114 Koskinen, M., Elo, H., Lukkari, P. and Riekolla, M.-L.: Determination of the antileukemic drug mitoguazone and seven other closely related bis(amidinohydrazones) in human blood serum by high-performance liquid chromatography. *J. Chromatogr. B*, 685 (1996) 141-149.
- 1115 Liang, X.-Z., Palmeier, R.K. and Lunte, C.E.: Dual-electrode amperometric detection for the determination of SR4233 and its metabolites with microbore liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1995) 113-119.
- 1116 Liu, W.-W., Liu, C.-H. and Kuo, S.-C.: HPLC assay of N-ethyl-phenylacetamide in serum and pharmacokinetics in rabbits. *Chin. Pharm. J. (Taipei)*, 48 (1996) 149-155; C.A., 125 (1996) 211666j.
- 1117 Mandl, A. and Lindner, W.: Improved detection of leucovorin in mixed folates and antifolates by reversed-phase liquid chromatography and on-line post-column UV irradiation. *Chromatographia*, 43 (1996) 327-330.
- 1118 Marfil, F., Pineau, V., Sioufi, A. and Godbillon, J.: High-performance liquid chromatography of the aromatase inhibitor, letrozole, and its metabolite in biological fluids with automated liquid-solid extraction and fluorescence detection. *J. Chromatogr. B*, 683 (1996) 251-258.
- 1119 Martinsen, A. and Gynther, J.: Retention behaviour of triphenylethylene derivatives in reverse phase liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2117-2128.
- 1120 Soltis, M.J., Yeh, H.J., Cole, K.A., Whittaker, N., Wersto, R.P. and Kohn, E.C.: Identification and characterization of human metabolites of CAI [5-amino-1-(4'-chlorobenzoyl)-3,5-dichlorobenzyl]-1,2,3-triazole-4-carboxamide]. *Drug Metab. Disp.*, 24 (1996) 799-806.
- 1121 Synold, T.W., Xi, B., Newman, E.M., Muggia, F.M. and Doroshow, J.H.: Simple and sensitive method for the quantitative analysis of lometrexol in plasma using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 683 (1996) 245-249.
- 1122 Van Gijn, R., Havik, E., Boven, E., Vermorken, J.B., tan Bokkel Huinink, W.W., van Tellingen, O. and Beijnen, J.H.: High-performance liquid chromatographic analysis of the new antitumour drug N-benzoylstauropospirone (CGP 41 251) and four potential metabolites in micro-volumes of plasma. *J. Pharm. Biomed. Anal.*, 14 (1995) 165-174.
- 1123 Volkov, S.K.: (Application of high-performance liquid chromatography for the determination of vinblastine and vincristine). *Khim.-Farm.-Zh.*, 30 (1996) 58-62; C.A., 125 (1996) 67908v - a review with 49 refs.
- 1124 Volkov, S.K. and Grodnitskaya, E.I.: (Determination of vincristine in formulations). *Khim.-Pharm. Zh.*, 30 (1996) 58-59; C.A., 125 (1996) 151274t.
- 1125 Yu, Z. and Westerlund, D.: Ion-pair chromatography of methotrexate in a column-switching system using an alkyl-diolsilica precolumn for direct injection of plasma. *J. Chromatogr. A*, 742 (1996) 113-120.
- For additional information see C.A.:
125 (1996) 123939v, 212667x.
- See also 232, 292, 410, 821, 859, 1169.
- 32g. Other drug categories
- 1126 Alfredson, T.V., Towne, R., Elliott, M., Griffin, B., Abubakari, A., Dyson, N. and Kertesz, D.J.: Enantioselectivity of azalanstat and its ketal tosylate intermediate in chiral high performance liquid chromatography separations. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1653-1668.
- 1127 Babizhayev, M.A., Yernakova, V.N., Sakina, N.L., Evstigneeva, R.P., Rozhkova, E.A. and Zheltukhina, G.A.: N⁶-Acetylcarnosine is a prodrug of L-carnosine in ophthalmic application as antioxidant. *Clin. Chim. Acta*, 254 (1996) 1-21.
- 1128 Casals, I., Reixach, M., Amat, J., Fuentes, M. and Serra-Majem, L.: Quantification of cyclamate and cyclohexylamine in urine samples using high-performance liquid chromatography with trinitrobenzenesulfonic acid pre-column derivatization. *J. Chromatogr. A*, 750 (1996) 397-402.
- 1129 Fan, T.Y., Wall, G.M., Sternitzke, K., Bass, L., Morton, A.B. and Muegge, E.: Improved high-performance liquid chromatographic determination of pilocarpine and its degradation products in ophthalmic solutions. Importance of octadecylsilane column choice. *J. Chromatogr. A*, 740 (1996) 289-295.
- 1130 González Gigosos, P., Fente Sampayo, C.A., Franco Abuín, C., Vázquez Belda, B., Quinto Fernández, E. and Cepeda Sáez, A.: Liquid chromatographic analysis of clenbuterol residues in food products of animal origin (cow liver) using diphasic analysis extraction. *Chromatographia*, 43 (1996) 271-274.
- 1131 Hashimoto, Y.: (LC/MS or GC/MS?). *Jasco Rep.*, 38 (1996) 20-26; C.A., 125 (1996) 25435h.
- 1132 Hashizume, T. and Myake, Y.: (Separation of aromatic fluoro derivatives from nitro derivatives by liquid chromatography as intermediates for preparation of diagnostic agents for PET). *Jpn. Kokai Tokkyo JP 08 71,413 [96 71,413]* (Cl. B01J20/10), 19 Mar. 1996, JP Appl. 94/157,097, 8 Jul. 1994; 10 pp.; C.A., 125 (1996) 29267c.
- 1133 Husain, S., Khalid, S., Nagaraju, V. and Rao, R.N.: High-performance liquid chromatographic separation and determination of small amounts of process impurities of famotidine in bulk drugs and formulations. *J. Chromatogr. A*, 743 (1996) 328-334.
- 1134 Lee, C. and Tannock, I.: Pharmacokinetic studies of amiloride and its analogs using reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 685 (1996) 151-157.
- 1135 Ohtani, H., Kotaki, H., Sawada, Y. and Iga, T.: Quantitative determination of epinastine in plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 683 (1996) 281-284.
- 1136 Priston, M.J. and Sewell, G.J.: The analysis of apomorphine formulations of ambulatory infusions. *Pharm. Sci.*, 1 (1995) 91-94; C.A., 125 (1996) 177542d.
- 1137 Tagliaro, F., Moffa, M., de Battisti, Z., Smith, F.P. and Gentile, M.: High-performance liquid chromatographic determination of levodropazine in human plasma with fluorometric detection. *J. Chromatogr. B*, 685 (1996) 165-170.
- 1138 William, M.G., Toth, L.N., Padbury, G.E. and Zhong, W.Z.: Determination of CS-045 (U-87326) in rat serum using LCEC. *Curr. Sep.*, 15 (1996) 13-17; C.A., 125 (1996) 237419f.

For additional information see C.A.:
125 (1996) 67102c.

See also 51, 281, 327, 444, 453, 794, 878, 879, 985, 1006, 1188, 1190, 1191.

32h. Toxicological and forensic applications

- 1139 Akasaka, K., Ohrii, H., Meguro, H. and Yasumoto, T.: Determination of dinophysistoxin-3 by the LC-LC method with fluorometric detection. *Anal. Sci.*, 12 (1996) 557-560; C.A., 125 (1996) 21435m.
- 1140 Balikova, M.: Application of HPLC with photodiode array detection for systematic toxicological analyses of drug groups. In: Jacob, B. and Bonte, W. (Editors), *Adv. Forensic Sci., Proc. Meet. Int. Assoc. Forensic Sci., 13th 1995*, Verlag Dr. Koester, Berlin, 1995, pp.72-76; C.A., 125 (1996) 50947.
- 1141 Barat, S.A., Kardos, S.A. and Abdel-Rahman, M.S.: Development and validation of a high-performance liquid chromatography method for the determination of cocaine, its metabolites and lidocaine. *J. Appl. Toxicol.*, 16 (1996) 215-219; C.A., 125 (1996) 78745r.
- 1142 Bianchi, V., Mazza, L., Macchì, C. and Donzelli, G.: (11-Nor- Δ^9 -tetrahydrocannabinol-9-carboxylic acid: a new HPLC method for the urinary confirmation of cannabis derivative). *G. Ital. Chim. Clin.*, 20 (1995) 227-232; C.A., 125 (1996) 50985v.
- 1143 Clanwaert, K.M., van Bockelaer, J.F., Lambert, W.E. and de Leenheer, A.P.: Analysis of cocaine, benzoylecgonine, and co-acaethylene in urine by HPLC with diode array detection. *Anal. Chem.*, 68 (1996) 3021-3028.
- 1144 Cocker, J., Nutley, B.P. and Wilson, H.K.: Methylene bis(2-chloroaniline) (MbOCA): towards a biological monitoring guidance value. *Biomarkers*, 1 (1996) 185-189; C.A., 125 (1996) 214351p.
- 1145 Dorsey, J.: Thermodynamically correct bioavailability estimations. *Report*, 1995, (AFOSR-TR-96-0133; Order No. AD-A305825), 4 pp.; C.A., 125 (1996) 214501n.
- 1146 Fisher, D.H., Broady, M.I. and Fisher, L.M.: Quantification of 9-carboxy-11-nor- Δ^9 -tetrahydrocannabinol in urine using brominated 9-carboxy-11-nor- Δ^9 -tetrahydrocannabinol as the internal standard and high-performance liquid chromatography with electrochemical detection. *Biomed. Chromatogr.*, 10 (1996) 161-166; C.A., 125 (1996) 160530a.
- 1147 Flynn, K. and Flynn, K.J.: An automated HPLC method for the rapid analysis of paralytic shellfish toxins from dinoflagellates and bacteria using precolumn oxidation at low temperature. *J. Exp. Mar. Biol. Ecol.*, 197 (1996) 145-157; C.A., 125 (1996) 107176p.
- 1148 Haglund, P.: Isolation of PCB atropisomers for toxicological testing using chiral high-performance liquid chromatography. *Organohalogen Compd.*, 23(Dioxin 95, 15th International Symposium on Chlorinated Dioxins and Related Compounds, 1995) (1995) 35-40; C.A., 125 (1996) 107213y.
- 1149 Hua, Y., Lu, W., Henry, M.S., Pierce, R.H. and Cole, R.B.: On-line liquid chromatography-electrospray ionization mass spectrometry for determination of the brevetoxin profile in natural "red tide" algae blooms. *J. Chromatogr. A*, 750 (1996) 115-125.
- 1150 Kelly, S.S., Bishop, A.G., Carmody, E.P. and James, K.J.: Isolation of dinophysistoxin-2 and the high-performance liquid chromatographic analysis of diarrhetic shellfish toxins using derivatisation with 1-bromoacetylpyrene. *J. Chromatogr. A*, 749 (1996) 33-40.
- 1151 List, W., Schueler, E. and Lenz, U.: (A method for the purification of filamentous hemagglutinin and pertussis toxin). *Ger. DE 19,512,346 (Cl. C07K14/235)*, 13 Jun. 1996, Appl. 19,512,346, 1 Apr. 1995; 4 pp.; C.A., 125 (1996) 107645r.
- 1152 Lopes, H.R., Noleto, A.L.S. and Bergdoll, M.S.: Purification of staphylococcal enterotoxins B and C₂ by dye ligand chromatography-production of antisera. *Rev. Microbiol.*, 27 (1996) 52-56; C.A., 125 (1996) 107487r.
- 1153 Luckas, B.: (Determination of algal toxins from seafood). *G/T Fachz. Lab.*, 40 (1996) 355-359; C.A., 125 (1996) 56465x - a review with 15 refs.
- 1154 Mao, I.F., Chen, M.L. and Lo, E.W.: Simultaneous determination of urinary metabolites of toluene, xylene, styrene and ethyl benzene by solid-phase extraction technique and high-performance liquid chromatographic/photo diode array detection. *Int. J. Environ. Anal. Chem.*, 64 (1996) 1-9; C.A., 125 (1996) 240365r.
- 1155 Morita, H., Urashima, Y., Tanaka, S., Aketagawa, J., Tamura, H. and Kumazawa, Y.: (Simultaneous estimation of endotoxin and (1-3)- β -D-glucan with an automated kinetic-chromogenic assay using wellreader). *Bokin Bobai*, 24 (1996) 467-475; C.A., 125 (1996) 107205x.
- 1156 Musser, S.M.: Quantitation and identification of fumonisins by liquid chromatography/mass spectrometry. *Adv. Exp. Med. Biol.*, 392(Fumonisins in Food) (1996) 65-74; C.A., 125 (1996) 107169p.
- 1157 Phillips, D.L., Tebbett, I.R. and Bertholf, R.L.: Comparison of HPLC and GC-MS for measurement of cocaine and metabolites in human urine. *J. Anal. Toxicol.*, 20 (1996) 305-308; C.A., 125 (1996) 187734g.
- 1158 Rashid, B.A., Kwasowski, P. and Stevenson, D.: The development of a single-step solid-phase immuno-extraction for morphine. *Pharm. Sci.*, 2 (1996) 115-116; C.A., 125 (1996) 214366x.
- 1159 Rustichelli, C., Ferioli, V., Vezzalini, F., Rossi, M.C. and Gamberini, G.: Simultaneous separation and identification of hashish constituents by coupled liquid chromatography-mass spectrometry. *Chromatographia*, 43 (1996) 129-134.
- 1160 Sugie, H., Nishikawa, T. and Funao, T.: Identification of bloodstains as antemortem or postmortem: quantitation of nucleotides, nucleosides and bases in bloodstains by high-performance liquid chromatography. In: Jacob, B. and Bonte, W. (Editors), *Adv. Forensic Sci., Proc. Meet. Int. Assoc. Forensic Sci., 13th 1995*, Verlag Dr. Koester, Berlin, 1995, pp. 212-276; C.A., 125 (1996) 50936e.
- 1161 Williams, L.L. and Giesey, J.P.: Characterization studies of a semi-automated separation method for analysis of non-ortho substituted polychlorinated biphenyl (PCB) congeners in environmental samples. *Toxicol. Environ. Chem.*, 51 (1995) 229-241; C.A., 125 (1996) 107197w.
- 1162 Wu, M. and Aderjan, R.: The effect of chromatographic conditions on the retention indices of forensically relevant substances in reversed-phase HPLC. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1967-1991.

- 1163 Zook, C.M., Patel, P.M., LaCourse, W.R. and Ralapati, S.: Characterization of tobacco products by high-performance anion exchange chromatography-pulsed amperometric detection. *J. Agric. Food Chem.*, 44 (1996) 1773-1779; C.A., 125 (1996) 30276m.

See also 390, 606, 792, 828, 926, 1040, 1082.

32i. Plant extracts

- 1164 Bos, R., Woerdenbag, H.J., Hendricks, H., Zwaving, J.H., de Smet, P.A.G.M., Tittel, G., Wikstrom, H.V. and Scheffer, J.J.C.: Analytical aspects of phytotherapeutic valeren preparations. *Phytochem. Anal.*, 7 (1996) 143-151; C.A., 125 (1996) 52637g.

- 1165 Dou, J., Khan, I.A., McChesney, J.D. and Burandt, C.L., Jr.: Qualitative and quantitative high performance liquid chromatographic analysis of quassinooids in Simaroubaceae plants. *Phytochem. Anal.*, 7 (1996) 192-200; C.A., 125 (1996) 96288y.

- 1166 Gobbato, S. and Lolla, E.: A new HPLC method for the analysis of valerenic acids in *Valeriana officinalis* extracts. *Fitoterapia*, 67 (1996) 159-162; C.A., 125 (1996) 96291u.

- 1167 Krizsán, K., Szókán, G., Tóth, Z.A., Hollósy, F., László, M. and Khlafulla, A.: HPLC analysis of anthraquinone derivatives in madder root (*Rubia tinctorum*) and its cell cultures. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2295-2314.

- 1168 Perry, N.B., Burgess, E.J., Lorimer, S.D. and van Klink, J.W.: Fatty acid anilides as internal standards for high performance liquid chromatographic analyses of *Valeriana officinalis* L. and other medicinal plants. *Phytochem. Anal.*, 7 (1996) 263-268; C.A., 125 (1996) 269503v.

- 1169 Poon, G.K., Bloomer, J.W., Clarke, S.E. and Maltais, J.: Rapid screening of taxol metabolites in human microsomes by liquid chromatography/electrospray ionization mass spectrometry. *Rapid Commun. Mass Spectrom.*, 10 (1996) 1165-1168; C.A., 125 (1996) 184734w.

- 1170 Tsai, T.H., Tsai, T.R., Chen, C.C. and Chen, C.F.: Pharmacokinetics of osthol in rat plasma using high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 14 (1996) 749-753.

For additional information see C.A.:

125 (1996) 96350n.

See also 209, 213, 222, 391, 405, 406, 408, 413, 810, 943.

33. CLINICO-CHEMICAL APPLICATIONS

- 33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 1171 Turhan, N., Coker, C., Hamulu, F., Elmaci, S., Yilmaz, C., Ersöz, B. and Islegen, C.: Glucose metabolism and catecholamine responses during physical exercise in non-insulin-dependent diabetes. *Eur. J. Clin. Chem. Clin. Biochem.*, 34 (1996) 683-689.

See also 250, 305, 308, 437, 438, 452, 580, 582, 625, 808, 844, 854, 990, 1221.

34. FOOD ANALYSIS

34a. General papers and reviews

- 1172 Mondello, L., Dufo, G. and Bartle, K.D.: Online microbore high performance liquid chromatography-capillary gas chromatography for food and water analyses. A review. *J. Microcolumn Sep.*, 8 (1996) 275-310; C.A., 125 (1996) 74752t.

See also 640.

34b. Complex mixtures (single compounds by cross-reference only)

- 1173 De Noni, I., Tirelli, A. and Masotti, F.: (Detection of cow milk in non-bovine cheese by HPLC of whey protein: application to goat milk cheese). *Sci. Tex. Latt.-Casearia*, 47 (1996) 7-17; C.A., 125 (1996) 113183e.

- 1174 González-Andrade, J.I., Romero, C., Morales, F.J. and Jiménez-Pérez, S.: An improved method for determination of the activity of β-galactosidase in yoghurt by high-performance liquid chromatography. *Chromatographia*, 43 (1996) 85-88.

- 1175 Imperato, A.: (Isolation of olive oil waxes using column chromatography: Note on Document IV of EEC Regulation 2568/91). *Riv. Ital. Sostanze Grasse*, 73 (1996) 265-266; C.A., 125 (1996) 193953u.

- 1176 Konings, E.J.M., Roomans, H.H.S. and Beljaars, P.R.: Liquid chromatographic determination of tocopherols and tocotrienols in margarine, infant foods, and vegetables. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 902-906.

- 1177 Lewis, D.J., Barnes, K.A., Wilkinson, K., Thorpe, S.A., Reynolds, S.L. and Startin, J.R.: Extraction of maleic hydrazide residues from potato crisps and their determination using high-performance liquid chromatography with UV and atmospheric pressure chemical ionisation mass spectrometric detection. *J. Chromatogr. A*, 750 (1996) 391-396.

- 1178 Low, N.H.: Food authenticity analysis by anion-exchange liquid chromatography. *Am. Lab. (Shelton)*, 28 (1996) 35M-35N, 35P, 35R, 35T, 35V-35W; C.A., 125 (1996) 56476b.

- 1179 Manzi, P., Panfili, G. and Pizzoferrato, L.: Normal and reversed-phase HPLC for more complete evaluation of tocopherols, retinols, carotenes and sterols in dairy products. *Chromatographia*, 43 (1996) 89-91.

- 1180 Monteiro, M., Nerin, C. and Reyes, F.G.R.: Determination of UV stabilizers in PET bottles by high performance-size exclusion chromatography. *Food Addit. Contam.*, 13 (1996) 575-586; C.A., 125 (1996) 56509q.

- 1181 Simeonidou, E.J., Botsoglou, N.A., Psomas, I.E. and Fletouris, D.J.: Liquid chromatographic analysis of multiple sulfonamide residues in chicken muscle using pre-column derivatization and fluorescence detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 2349-2364.

- 1182 Sutikno, Abouzied, M.M., Azcona-Olivera, J.I., Hart, L.P. and Pestka, J.J.: Detection of fumonisins in Fusarium cultures, corn, and corn products by polyclonal antibody-based ELISA: Relation to fumonisin B₁ detection by liquid chromatography. *J. Food Prot.*, 59 (1996) 645-651; C.A., 125 (1996) 140787m.

- 1183 Timmermann, D.G.: (Detection of added protein hydrolyzates in foods). *Ger. DE 19,516,077 (Cl. G01N33/02)*, 18 Jul. 1996, Appl 19,516,077, 5 May 1995; 11 pp.; C.A., 125 (1996) 113355n.
- 1184 Van Boven, M., Daenens, P., Tytgat, J. and Cokelaere, M.: Determination of simmondsins and simmondsin ferulates in jojoba meal and feed by high-performance liquid chromatography. *J. Agric. Food Chem.*, 44 (1996) 2239-2243; C.A., 125 (1996) 140773d.
- 1185 Wilke, J. and Parthey, B.: (Process approximative beer mash analysis). *Monatsschr. Brauwiss.*, 49 (1996) 115-118; C.A., 125 (1996) 56468a.
- 1186 Yan, Z. and Weijun, M.: Simultaneous multicomponent determination of food by column-switching ion chromatography. *Anal. Lett.*, 29 (1996) 1817-1828; C.A., 125 (1996) 140819y.

For additional information see C.A.:
125 (1996) 56544x.

See also 200, 203, 204, 210, 214, 223, 228, 255, 263, 280, 314, 318, 339, 343, 348, 360, 361, 377, 392, 393, 425, 621, 644, 816, 833, 835, 884, 904, 907, 911, 912, 916, 923, 932, 934, 942, 943, 1095, 1097, 1098, 1109, 1153, 1235.

34c. Organoleptically important compounds (flavors, odors, volatiles)

- 1187 Ferary, S. and Auger, J.: What is the true odour of cut Allium? Complementarity of various hyphenated methods: gas chromatography-mass spectrometry and high-performance liquid chromatography-mass spectrometry with particle beam and atmospheric pressure ionization interfaces in sulphenic acids rearrangement components discrimination. *J. Chromatogr. A*, 750 (1996) 63-74.
- 1188 Galletti, G.C., Bocchini, P., Gioacchini, A.M. and Roda, A.: Analysis of the artificial sweetener aspartame by means of liquid chromatography/electrospray mass spectrometry. *Rapid Commun. Mass Spectrom.*, 10 (1996) 1153-1155; C.A., 125 (1996) 140770a.
- 1189 Herderich, M., Roscher, R. and Schreier, P.: Application of atmospheric pressure ionization liquid chromatography-tandem mass spectrometry for the analysis of flavor precursors. *ACS Symp. Ser.*, 637(Biotechnology for Improved Foods and Flavors) (1996) 261-271; C.A., 125 (1996) 273753u - a review with 16 refs.
- 1190 Moriyasu, T., Nakazato, M., Kobayashi, C., Kikuchi, Y., Hayano, K. and Tamura, Y.: (Determination of acesulfame K, saccharin, and aspartame in various foods by HPLC). *Shokuhin Eiseigaku Zasshi*, 37 (1996) 91-96; C.A., 125 (1996) 84955a.
- 1191 Szymczyk, K. and Czerwinski, L.: (HPLC determination of aspartame and acesulfam-K in processed fruit products). *Roczn. Panstw. Zakl. Hig.*, 46 (1995) 373-381; C.A., 125 (1996) 193703n.

For additional information see C.A.:
125 (1996) 216301w.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

- 1192 Barcelo, D. (Editor): *Applications of LC-MS in Environmental Chemistry*. *J. Chromatogr. Libr.*, Vol. 59, Elsevier, Amsterdam, 1996, 543 p.; C.A., 125 (1996) 47957g.

See also 146.

35b. Air pollution (complex mixtures; single compounds by cross-reference only)

- 1193 Tanaka, K. and Haddad, P.: Acid rain monitoring by ion-exclusion-cation-exchange chromatography. *TrAC*, 15 (1996) 266-273.

See also 243, 244, 246, 418, 789.

35c. Water pollution (complex mixtures; single compounds by cross-reference only)

- 1194 Huber, S.A. and Frimmel, F.H.: (Size-exclusion chromatography with organic carbon detection (LC-OCD). A fast and reliable method for the characterization of hydrophilic organic matter in natural waters). *Vom Wasser*, 86 (1996) 277-290; C.A., 125 (1996) 95309u.

1195 Lau, S.L.: Fractionation of extractable organics in urban runoff for toxicity identification. Avail. *Univ. Microfilms Int.*, Order No. DA9616448, 1996, 182 p.; C.A., 125 (1996) 66906f.

- 1196 Van Soest, R.E.J., Chervet, J.P., Ursem, M. and Suijlen, J.M.: Analysis of fluorescent water tracers using on-line preconcentration in micro HPLC. *LC-GC Int.*, 9 (1996) 586-593.

For additional information see C.A.:

125 (1996) 41246c.

See also 160, 198, 240, 414, 905, 906, 909, 910, 914, 917, 920, 1149, 1172, 1211, 1224, 1230.

35d. Soil pollution (complex mixtures; single compounds by cross-reference only)

See 919, 925.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

- 1197 Garcia, M.T., Ribosa, I., Leal, J.S. and Hreczuch, W.: Effect of homolog distribution on the toxicity of alcohol ethoxylates. *J. Am. Oil Chem. Soc.*, 73 (1996) 903-906.

- 1198 Heinig, K., Vogt, C. and Werner, G.: Separation of ionic and neutral surfactants by capillary electrophoresis and high-performance liquid chromatography. *J. Chromatogr. A*, 745 (1996) 281-292.

- 1199 Moskvin, L.N., Simon, J., Loeffler, P., Michailova, N.V. and Nicolaevna, D.N.: Photometric determination of anionic surfactants with a flow-injection analyser that includes a chromatomembrane cell for sample preconcentration by liquid-liquid solvent extraction. *Talanta*, 43 (1996) 819-824; C.A., 125 (1996) 171520y.
- 1200 Truchan, J., Rasmussen, H.T., Omelchenko, N. and McPherson, B.P.: Analysis of cationic surfactants in household products by high performance liquid chromatography with nitrogen chemiluminescence detection. *J. Liq. Chromatogr. Relat. Technol.*, 19 (1996) 1785-1792.

36b. Antioxidants and preservatives

- 1201 Maillard, M.N., Giampaoli, P. and Cuvelier, M.E.: Atmospheric pressure chemical ionization (APCI) liquid chromatography-mass spectrometry: characterization of natural antioxidants. *Talanta*, 43 (1996) 339-347; C.A., 125 (1996) 84910g.

See also 1088, 1094, 1180.

36c. Complex mixtures, technical products and unidentified compounds

- 1202 Barbieri, M. and Macherpa, A.: (HPLC determination of denaturants in automotive petroleum fuels). *Riv. Combust.*, 49 (1995) 427-435; C.A., 125 (1996) 37698j.
- 1203 Farid, N.A., Schreiner, K.M., Coleman, N.L., van Lier, R.B.L. and Wrighton, S.A.: Stereoselective disposition of the enantiomers of the benzoquinolinone LY191704, a human type I 5 α -reductase inhibitor. Differences between rats and dogs. *Drug Metab. Disp.*, 24 (1996) 1162-1165.
- 1204 Kaneko, H., Morino, M., Takanohashi, T. and Iino, M.: Study of the interaction of aromatic compounds with coals by inverse liquid chromatography. *Energy Fuels*, 10 (1996) 1017-1021; C.A., 125 (1996) 37767f.
- 1205 Mohite, B.S.: Recent developments of very simple separation methods for major and trace elements from geological samples using poly(dibenzo-18-crown-6). In: Mehrotra, S.P. and Sjekhar, R. (Editors), *Miner Process: Recent Adv. Future Trends, Proc. Conf.*, Allied Publishers, New Delhi, 1995, pp. 890-895; C.A., 125 (1996) 131005c.
- 1206 Schoenmakers, P., Blomberg, J. and Kerkvliet, S.: Sophisticated separation methods and the oil industry. *LC-GC Int.*, 9 (1996) 718-724.
- 1207 Tang, H.R., Hancock, R.A. and Covington, A.D.: Studies of vegetable tannins. Characterization of commercial chestnut tannin extract. *J. Soc. Leather Technol. Chem.*, 80 (1996) 15-24; C.A., 125 (1996) 29984j.
- 1208 Yuan, J.H., Birkmeier, J., Yang, D.C., Hribar, J.D., Liu, N., Bible, R., Hajdu, E., Rock, M. and Schoenhard, G.: Isolation and identification of metabolites of leukotriene A₄ hydrolase inhibitor SC-57461 in rats. *Drug Metab. Disp.*, 24 (1996) 1124-1133.

For additional information see C.A.:

125 (1996) 103899d, 112195y, 157303y.

See also 822, 1194, 1195.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 1209 Li, P. and Giddings, J.C.: Isolation and measurement of colloids in human plasma by membrane-selective flow field-flow fractionation: lipoproteins and pharmaceutical colloids. *J. Pharm. Sci.*, 85 (1996) 895-898.

See also 729, 972.

38. INORGANIC COMPOUNDS

38a. Cations

- 1210 Bandyopadhyay, A. and Roy, U.S.: Extraction chromatographic studies of zirconium(IV) with *n*-capric acid. *J. Indian Chem. Soc.*, 73 (1996) 177-180; C.A., 125 (1996) 47716c.
- 1211 Duff, M.C. and Amrhein, C.: Method for the separation of uranium(IV) and (VI) oxidation states in natural waters. *J. Chromatogr. A*, 743 (1996) 335-340.
- 1212 Gilon, N., Potin-Gautier, M. and Astruc, M.: Optimization of the determination of inorganic and organic selenium species using high-performance liquid chromatography-electrothermal atomic absorption spectrometry. *J. Chromatogr. A*, 750 (1996) 327-334.
- 1213 Hajos, P., Revesz, G., Horvath, O., Peear, J. and Sarzanini, C.: The simultaneous analysis of metal-EDTA complexes and inorganic anions by suppressed ion chromatography. *J. Chromatogr. Sci.*, 34 (1996) 291-299.
- 1214 Hao, F., Paull, B. and Haddad, P.R.: Determination of trace levels of thorium(IV) and uranyl by reversed-phase chromatography with on-line preconcentration and ligand exchange. *J. Chromatogr. A*, 749 (1996) 103-113.
- 1215 Hawkes, W.C. and Kutnink, M.A.: High-performance liquid chromatographic-fluorescence determination of traces of selenium in biological materials. *Anal. Biochem.*, 241 (1996) 206-211.
- 1216 Ichinoki, S., Yamazaki, M., Omura, M. and Nagai, R.: (Simultaneous determination of heavy metals at ppm level in bovine liver by automated reversed-phase HPLC involving precolumn chelation with hexamethylenedithiocarbamate and online column enrichment). *Bunseki Kagaku*, 45 (1996) 837-844; C.A., 125 (1996) 242069w.
- 1217 Janos, P. and Aczel, P.: Ion chromatographic separation of selenite and selenate using a polyanionic eluent. *J. Chromatogr. A*, 749 (1996) 115-122.
- 1218 Khuhawar, M.Y. and Lanjwani, S.N.: Simultaneous solvent extraction and high-performance liquid chromatographic determination of uranium, iron, nickel and copper in mineral ore samples and phosphate rock residues using N,N'-ethylene-bis(salicylaldimine) as complexing reagent. *J. Chromatogr. A*, 740 (1996) 296-301.
- 1219 Koppen, P.A.M.V., Kemper, P.R. and Bowers, M.T.: Electronic state effects in sigma bond activation by first row transition metal ions: the ion chromatography technique. *Understanding Chem. React.*, 15 (1996) 157-196; C.A., 125 (1996) 68681c - a review with 90 refs.

- 1220 Lajnwani, S.N., Mahar, K.P. and Channer, A.H.: Simultaneous determination of cobalt, copper, iron and vanadium in crude petroleum oils by HPLC. *Chromatographia*, 43 (1996) 431-432.
- 1221 Lee, B.-L., Chua, L.-H., Ong, H.-Y., Yang, H.G., Wu, J. and Ong, C.-N.: Determination of serum and urinary aluminium by HPLC with fluorometric detection of Al-lumogallion complex. *Clin. Chem. (Washington)*, 42 (1996) 1405-1411.
- 1222 Moreira, I.S. and Santiago, M.O.: Identification of [(bpy)₂Ru(Im)(H₂O)]²⁺ complex as a reaction intermediate by reversed-phase high-performance liquid chromatography. *Chromatographia*, 43 (1996) 322-326.
- 1223 Motyleva, S.M. and Sosnina, M.V.: (Determination of heavy metals content of black currant berries and leaves by HPLC). *S-kh. Biol.*, (1996) 67-71; C.A., 125 (1996) 216015f.
- 1224 Pantsar-Kallio, M. and Manninen, P.K.G.: Speciation of chromium in waste waters by coupled column ion chromatography-inductively coupled plasma mass spectrometry. *J. Chromatogr. A*, 750 (1996) 89-95.
- 1225 Pirogov, A.V., Semenova, S.N., Obrezkov, O.N. and Shpigun, O.A.: The use of pentachlorophenol as an eluent preservative in the ion-chromatographic determination of transition metals. *LC-GC Int.*, 9 (1996) 594-596.
- 1226 Prat, M.D., Compañó, R., Granados, M. and Miralles, E.: Liquid chromatographic determination of gallium and indium with fluorimetric detection. *J. Chromatogr. A*, 746 (1996) 239-245.
- 1227 Song, M., Ma, H.M., Huang, Y.X. and Liang, S.Q.: Determination of speciation of magnesium(II) in albumen by gel permeation chromatography. *Chin. Chem. Lett.*, 7 (1996) 823-824; C.A., 125 (1996) 273929f.
- 1228 Teräshde, P., Pantsar-Kallio, M. and Manninen, P.K.G.: Simultaneous determination of arsenic species by ion chromatography-inductively coupled plasma mass spectrometry. *J. Chromatogr. A*, 750 (1996) 83-88.
- For additional information see C.A.:
125 (1996) 74809s, 103804u, 184349z.
- See also 71, 110, 112, 145, 832, 940.
- 38b. Anions**
- 1229 Barman, B.N.: Determination of elemental sulfur by size-exclusion chromatography. Optimization and petrochemical applications. *J. Chromatogr. A*, 740 (1996) 237-244.
- 1230 Charles, L., Pepin, D. and Casetta, B.: Electrospray ion chromatography-tandem mass spectrometry of bromate at sub-ppb levels in water. *Anal. Chem.*, 68 (1996) 2554-2558.
- 1231 Jain, A., Chaurasia, A., Sahasrabuddhey, B. and Verma, K.K.: Determination of bromide in complex matrices by pre-column derivatization linked to solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr. A*, 746 (1996) 31-41.
- 1232 Kobayashi, M.: Study of precise pigment composition of photosystem I-type reaction centers by means of normal-phase HPLC. *J. Planta Res.*, 109 (1996) 223-230; C.A., 125 (1996) 216009g.
- 1233 Magyani, O.F.: Separation of inorganic anions and carboxylic acids using sulphonated acids as eluents. *Philipp. J. Sci.*, 125 (1996) 37-52; C.A., 125 (1996) 130989q.
- 1234 Miura, Y., Kimura, M., Ochi, M. and Koh, T.: Effect of eluents on suppressed ion chromatography using anion-exchange column of dionex HPIC-AS5. *Proc. Sch. Sci. Tokai Univ.*, 31 (1996) 83-90; C.A., 125 (1996) 74803k.
- 1235 Miyazaki, M. and Yamamoto, A.: (Recent development of ion chromatography to the analysis of anions in foodstuffs). *Shokuhin Eiseigaku Zasshi*, 37 (1996) 65-71; C.A., 125 (1996) 56461t - a review with 68 refs.
- 1236 Munae, E., Zein, R., Takeuchi, T. and Miwa, T.: Microcolumn ion chromatography of inorganic anions using bovine serum albumin stationary phase with indirect photometric detection. *Chromatographia*, 43 (1996) 304-308.
- 1237 Roeder, V. and Jardy, A.: (Determination of anionic impurities in acids and concentrated inorganic salts by ion chromatography. II. Case of acetic acid: in line elimination of the matrix and preconcentration of anion traces). *Analisis*, 24 (1996) 43-48; C.A., 125 (1996) 157109q.
- 1238 Warner, C.R., Daniels, D.H., Frank, L., Jr. and Diachenko, G.W.: Measurement of bromate in bottled water by high-performance liquid chromatography with post-column flow reactor detection. *Food Addit. Contam.*, 13 (1996) 633-638; C.A., 125 (1996) 193710n.

For additional information see C.A.:

125 (1996) 32191d, 47820g, 62472p, 74851z, 130993m, 184348y.

See also 98, 110, 820, 934, 1193, 1213.

38d. Volatile inorganic compounds

- 1239 Stratford, M.R.L., Dennis, M.F., Tozer, G.M., Prise, V.E. and Everett, S.A.: Measurement of nitric oxide in biological materials by high performance ion chromatography and EPR spectroscopy: a comparative study. *Portland Press Proc.*, 10(Biology of Nitric Oxide, Part 5) (1996) 175; C.A., 125 (1996) 52936d.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 1240 Abramson, F.P., Teffera, Y., Kusmierz, J., Steenwyk, R.C. and Pearson, P.G.: Replacing ¹⁴C with stable isotopes in drug metabolism studies. *Drug Metab. Disp.*, 24 (1996) 697-701.

For additional information see C.A.:

125 (1996) 98000x.

See also 179, 976.