

COMPOUND INDEX

J. CHROMATOGR. A VOLS. 752–800

For all compound groups: see also sub-groups and individual compounds
For all compound sub-groups: see also main groups and individual compounds
For all individual compounds: see also compound groups and sub-groups

A**A-2545**

Validated liquid chromatographic method for the determination of N-3-(2,2,5,5-tetramethyl-3-pyrrolin-3-carboxamidopropylphthalimide hydrochloride), a novel antiarrhythmic agent in human plasma 797(1998)265

Abietic acid

High-performance liquid chromatographic determination of dehydroabietic and abietic acids in traditional Chinese medications 763(1997)221

Acaricides

Retention profiles of some commercial pesticides, pyrethroid and acaricide residues and their application to tomato and parsley plants 760(1997)179

Influence of solvent and storage conditions on the stability of acaricide standard stock solutions 765(1997)109

Acesulfame-K

Separation and determination of four artificial sweeteners and citric acid by high-performance anion-exchange chromatography 771(1997)135

Acetaldehyde

Study of the acetaldehyde induced polymerisation of flavan-3-ols by liquid chromatography–ion spray mass spectrometry 752(1996)85

Acetaminophen

see also Paracetamol

¹³C Isotopic analysis of an acetaminophen and diacetylmorphine mixture 756(1996)203

Quantitation of acetaminophen and salicylic acid in plasma using capillary electrophoresis without sample pretreatment. Improvement of precision 768(1997)125

Effect of pH and ions in the sample on stacking in capillary electrophoresis 781(1997)35

Acetates

Gas chromatographic determination of vapour pressures of pheromone-like compounds. IV. Acetates, a reinvestigation 759(1997)93

Practical capillary electrophoresis method for the quantitation of the acetate counter-ion in a novel antifungal lipopeptide 763(1997)279

Acetonitrile

Simultaneous use of urea and acetonitrile as organic modifiers for optimization of resolution in micellar electrokinetic chromatography 792(1997)157

Acetophenone

Temperature gradients in preparative high-performance liquid chromatography columns 769(1997)109

Resolution studies on counter-current chromatography using supercritical fluid carbon dioxide 790(1997)31

Acetylalanylalanine

Vacancy affinity capillary electrophoresis, a new method for measuring association constants 775(1997)313

Acetylated cyclodextrins

Effect of β -cyclodextrin acetylation on the resolution of phenethylamines with capillary electrophoresis and nuclear magnetic resonance spectroscopy 758(1997)277

Acetylcysteine

Determination of glutathione, cysteine and N-acetylcysteine in rabbit eye tissues using high-performance liquid chromatography and post-column derivatization with 5,5'-dithiobis(2-nitrobenzoic acid) 778(1997)347

Determination of amino acids by high-performance capillary electrophoresis with on-line mode in-capillary derivatization 787(1997)253

Comparison of the stability and UV and fluorescence characteristics of the *o*-phthaldialdehyde/3-mercaptopropionic acid and *o*-phthaldialdehyde/N-acetyl-L-cysteine reagents and those of their amino acid derivatives 798(1998)37

Acetyldigitoxin

Separation of cardiac glycosides by micellar electrokinetic chromatography and microemulsion electrokinetic chromatography 779(1997)227

Acetyldigoxin

Separation of cardiac glycosides by micellar electrokinetic chromatography and microemulsion electrokinetic chromatography 779(1997)227

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Acetylene

Effect of addition of a graphitized carbon black trap to a glass beads trap on the cryoconcentration of some non-methane hydrocarbons in ambient air 790(1997)117

Acetylprocainamide

Direct determination of procainamide and N-acetylprocainamide by capillary zone electrophoresis in pharmaceutical formulations and urine 772(1997)271

α_1 -Acid glycoprotein

Identification of α_1 -acid glycoprotein (orosomucoid) in human synovial fluid by capillary electrophoresis 772(1997)305

Acids

see also Carboxylic acids, Fatty acids, Organic acids, and Weak acids

Migration behaviour of monovalent weak acids in micellar electrokinetic chromatography. Mobility model versus retention model 765(1997)295

Enantiomer separation of acidic racemates by capillary electrophoresis using cationic and amphoteric β -cyclodextrins as chiral selectors 781(1997)151

Separation of polar solutes by packed column supercritical fluid chromatography (Review) 785(1997)3

Acridines

Clean-up and analysis of carbazole and acridine type polycyclic aromatic nitrogen heterocyclics in complex sample matrices 790(1997)73

Acridinium esters

Optimization of post-column chemiluminescent detection for low-molecular-mass conjugates of acridinium esters 765(1997)145

Acridinylaminopropranolol

Fluorescence detection in packed-column supercritical fluid chromatographic separations 798(1998)203

Acrylamide

Separation of polyelectrolytes of variable compositions by free-zone capillary electrophoresis 766(1997)233

Rigid porous polyacrylamide-based monolithic columns containing butyl methacrylate as a separation medium for the rapid hydrophobic interaction chromatography of proteins 775(1997)65

Acrylamides, N-substituted

Performance of a series of novel N-substituted acrylamides in capillary electrophoresis of DNA fragments 756(1996)255

Acrylamidomethylpropanesulfonate

Separation of polyelectrolytes of variable compositions by free-zone capillary electrophoresis 766(1997)233

Acrylates

Supercritical fluid chromatography-proton nuclear magnetic resonance spectroscopy coupling (Review) 785(1997)65

Acrylic acid

Selection of a precursor monomer for the introduction of affinity ligands onto a porous membrane by radiation-induced graft polymerization 758(1997)209

N-Acryloylaminopropanol

Performance of a series of novel N-substituted acrylamides in capillary electrophoresis of DNA fragments 756(1996)255

Actinides

Use of ion chromatography for the determination of fission products and actinides in nuclear applications (Review) 789(1997)369

Acylcarnitines

Separation of carnitine and acylcarnitines by capillary electrophoresis 781(1997)475

Adenosine triphosphate

On-line electro dialysis–capillary zone electrophoresis of adenosine triphosphate and inositol phosphates 764(1997)135

Development of a capillary electrophoresis method with indirect photometric detection for the determination of anions related to anticaries 781(1997)457

Adipic acid polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. II. Solubility effects in the analysis of polyester resins under reversed-phase conditions 761(1997)147

Agarose

Direct measurements of convective fluid velocities in superporous agarose beads 795(1998)199

Aglycones

High-performance liquid chromatographic determination of flavonoids in *Betula pendula* and *Betula pubescens* leaves 793(1998)370

Alachlor

Comparative degradation kinetics of alachlor in water by photocatalysis with FeCl₃, TiO₂ and photolysis, studied by solid-phase disk extraction followed by gas chromatographic techniques 754(1996)187

Alanine

Computerized optimization of selectivity for direct capillary gas chromatographic multicomponent separations of enantiomers 779(1997)253

Alanine, dansyl-

Indirect time-resolved fluorescence detection of both non-fluorescent and fluorescent compounds separated by high-performance liquid chromatography 787(1997)261

Albumin

Analysis of recombinant human erythropoietin in drug formulations by high-performance capillary electrophoresis 759(1997)177

Chiral resolution of pantoprazole sodium and related sulfoxides by complex formation with bovine serum albumin in capillary electrophoresis 759(1997)185

Preparative desalting of bovine serum albumin by continuous annular chromatography 763(1997)49

Comparative studies on the isothermal characteristics of proteins adsorbed under batch equilibrium conditions to ion-exchange, immobilised metal ion affinity and dye affinity matrices with different ionic strength and temperature conditions 763(1997)71

Identification of proteins by capillary electrophoresis–tandem mass spectrometry. Evaluation of an on-line solid-phase extraction device 763(1997)295

Effect of mobile phase composition on the binding kinetics of chiral solutes on a protein-based high-performance liquid chromatography column. Interactions of D- and L-tryptophan with immobilized human serum albumin 766(1997)15

Effect of organic modifiers on retention and enantiomeric separations by capillary electrophoresis with human serum albumin as a chiral selector in solution 766(1997)237

Site accessibility and the pH dependence of the saturation capacity of a highly cross-linked matrix. Immobilized metal affinity chromatography of bovine serum albumin on Chelating Superose 767(1997)11

Enantioselectivity of bovine serum albumin-bonded columns produced with isolated protein fragments. II. Characterization of protein fragments and chiral binding sites 769(1997)215

Characterisation of human serum albumin heterogeneity by capillary zone electrophoresis and electrospray ionization mass spectrometry 772(1997)235

Model discrimination and estimation of the intraparticle mass transfer parameters for the adsorption of bovine serum albumin onto porous adsorbent particles by the use of experimental frontal analysis data 776(1997)3

Study of the binding of antibiotics to human serum albumin by charge-transfer chromatography 776(1997)31

Fluidization characteristics of and protein adsorption on fluoride-modified porous zirconium oxide particles 776(1997)179

Comparison of five methods for the study of drug–protein binding in affinity capillary electrophoresis 777(1997)311

Recycling electrophoretic separations: modeling of isotachopheresis and isoelectric focusing 779(1997)165

Characterization of antigen–antibody complexes by size-exclusion chromatography coupled with low-angle light-scattering photometry and viscometry 787(1997)101

Modeling column regeneration effects on ion-exchange chromatography 791(1997)99

Importance of heat of adsorption in modeling protein equilibria for overloaded chromatography 796(1998)141

Electrostatic effects on protein partitioning in size-exclusion chromatography and membrane ultrafiltration 796(1998)229

Alcohol ethoxylates

Separation and determination of oligomers and homologues of aliphatic alcohol ethoxylates in textile lubricants and lubricant emulsion by high-performance liquid chromatography 771(1997)145

Alcohol polyethoxylates

High-performance liquid chromatographic–fluorescence determination of aliphatic alcohol polyethoxylates and poly(ethylene glycol)s in aqueous samples 756(1996)159

Alcohols

Isolation of three triterpenes and several aliphatic alcohols from *Crataegus monogyna* Jacq 767(1997)340

Silicalite as a sorbent for solid-phase extraction 771(1997)45

Gas–liquid chromatography–mass spectrometry of primary and secondary fatty alcohols and diols as their *tert.*-butyldimethylsilyl derivatives 771(1997)191

Non-aqueous capillary electrophoretic separation of Brønsted acids as heteroconjugated anions 771(1997)275

Capillary gas chromatography–mass spectrometry of lower oxyethylenated aliphatic alcohols 773(1997)219

New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259

Retention behavior of nitrite on a reversed-phase column with an acidic aqueous eluent containing alcohol as a modifier 789(1997)207

Aldehydes

see also Carbonyl compounds

High-resolution gas chromatography–nitrogen–phosphorous detection of saturated volatile aldehydes derivatized with 2-hydrizinobenzothiazole 752(1996)209

Separation of phenolic aldehydes, ketones and acids from lignin degradation by capillary zone electrophoresis 755(1996)89

Impinger sampling coupled to high-performance liquid chromatography by a modified autoinjector interface 762(1997)187

Silicalite as a sorbent for solid-phase extraction 771(1997)45

Characterisation of aldehydic acids in used and unused frying oils 776(1997)245

Qualitative and quantitative high-performance liquid chromatographic analysis of aldehydes in Brazilian sugar cane spirits and other distilled alcoholic beverages 782(1997)13

Non-porous silica for ultrafast reversed-phase high-performance liquid chromatographic separation of aldehyde and ketone 2,4-dinitrophenylhydrazones 786(1997)47

High-performance liquid chromatographic peak identification of 2,4-dinitrophenylhydrazine derivatives of lipid peroxidation aldehydes by photodiode array detection 798(1998)117

Alditols

Anion-exchange chromatography with electrochemical detection of alditols and sugars at a Cu₂O–carbon composite electrode 773(1997)115

Separation of alditols of interest in food products by high-performance anion-exchange chromatography with pulsed amperometric detection 791(1997)343

Aldoses

Unusual separation of 1-phenyl-3-methyl-5-pyrazolone derivatives of aldoses by capillary zone electrophoresis 791(1997)307

Separation of aldose enantiomers by capillary electrophoresis in the presence of optically active N-dodecoxycarbonylvalines 792(1997)385

Aldrin

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

Aliphatic solutes

Study of retention in reversed-phase liquid chromatography using linear solvation energy relationships. I. The stationary phase 752(1996)1

Alkali metals

Histidine as a dipolar eluent component in cation chromatography 789(1997)141

Cationization in electrospray microcolumn liquid chromatography–mass spectrometry 794(1998)359

Alkaline earth metals

Histidine as a dipolar eluent component in cation chromatography 789(1997)141

- Determination of calcium and magnesium in water samples by high-performance liquid chromatography on a graphitic stationary phase with a mobile phase containing *o*-cresolphthalein complexone 789(1997)329
- Role of ligand purity in separations of alkaline earth metals as arsenazo I complexes by capillary zone electrophoresis 793(1998)177
- Alkaloids**
pH-Zone-refining countercurrent chromatography 753(1996)1
- Liquid chromatographic–electrospray mass spectrometric analysis of *Coptidis Rhizoma* 755(1996)19
- Determination of the alkaloids in *Evodiae fructus* by capillary electrophoresis 755(1996)113
- Ergot alkaloids as chiral selectors in capillary electrophoresis. Determination of the separation mechanism 755(1996)251
- Quantification of vincristine and vinblastine in *Catharanthus roseus* plants by capillary zone electrophoresis 755(1996)281
- Liquid chromatographic separation of the alkaloids in *coptis*–*evodia* herb couple 756(1996)137
- Lesser alkaloids of cocaine-bearing plants. III. 2-Carbomethoxy-3-oxo substituted tropane esters: detection and gas chromatographic–mass spectrometric characterization of new minor alkaloids found in South American *Erythroxylum coca* var. *coca* 756(1996)185
- Improved separation of six harmine alkaloids by high-performance capillary electrophoresis 761(1997)297
- High-performance liquid chromatography–electrospray ionisation–tandem mass spectrometry for the analysis of 1,2,3,4-tetrahydro- β -carboline derivatives 767(1997)101
- General approach for the analysis of various alkaloid classes using capillary electrophoresis and capillary electrophoresis–mass spectrometry 767(1997)263
- Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277
- Determination of atropine in pharmaceutical preparations by liquid chromatography with fluorescence detection 775(1997)137
- Application of micellar electrokinetic capillary chromatography to the analysis of illicit drug seizures 780(1997)265
- Determination of berberine and palmatine in *Phellodendri Cortex* using ion-pair supercritical fluid chromatography on-line coupled with ion-pair supercritical fluid extraction by on-column trapping 786(1997)371
- Erratum to “General approach for the analysis of various alkaloid classes using capillary electrophoresis and capillary electrophoresis–mass–spectrometry”. [J. Chromatogr. A, 767 (1997) 263–276] 786(1997)384
- Gas chromatographic analysis of indole alkaloids from *Tabernaemontana hilariana* 788(1997)204
- Improved detection of alkaloids in crude extracts applying capillary electrophoresis with field amplified sample injection 791(1997)323
- Capillary electrophoretic separation of herbicidal enantiomers applying ergot alkaloids 791(1997)339
- Analysis of traditional Chinese anticancer drugs by capillary electrophoresis 795(1998)388
- Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337
- Alkanes**
Clean-up procedures for the unbiased estimation of C_{37} alkenone sea surface temperatures and terrigenous *n*-alkane inputs in paleoceanography 757(1997)145
- Pressure drop effects in packed capillary column supercritical fluid chromatography 758(1997)117
- Hold-up time in gas chromatography. I. New approach to its estimation 760(1997)219
- Unified retention concept – statistical treatment of Kováts retention index 764(1997)257
- Optimization of the dwell-time for the quantitative analysis by gas chromatography–mass spectrometry 765(1997)241
- Integration of the equation of peak motion in programmed–pressure and –temperature gas chromatography 766(1997)147
- The hold-up time in gas chromatography. II. Validation of the estimation based on the concept of a zero carbon atoms alkane 767(1997)127
- Occurrence and determination of organic pollutants in aerosol, precipitation, and sediment samples collected at Lake Balaton 774(1997)349
- Evaluation of a micro gas chromatographic technique for environmental analyses of CO_2 and C_1 – C_6 alkanes 775(1997)243
- High speed solvating gas chromatography using packed capillaries containing sub-5 μm particles 778(1997)31
- Use of a thermally insulated column for improved speed, efficiency and resolution in packed-column supercritical fluid chromatography 785(1997)129

Supercritical fluid chromatograph for studies of retention and efficiency with a brief study of *n*-alkanes showing evidence of significant temperature drop with pressure drop 785(1997)135

Comparison of supercritical fluid extraction and Soxhlet extraction of organic compounds from carpet samples 787(1997)171

Fast monitoring of C₂₀–C₁₆₀ crude oil alkanes by size-exclusion chromatography–evaporative light scattering detection performed with silica columns 788(1997)63

Novel preconcentration technique for on-line coupling to high-speed narrow-bore capillary gas chromatography: sample enrichment by equilibrium (ab)sorption. II. Coupling to a portable micro gas chromatograph 791(1997)187

Determination of organic molecular markers in marine aerosols and sediments: one-step flash chromatography compound class fractionation and capillary gas chromatographic analysis 799(1998)215

Alkanes, branched

Evaluation of theoretical models of non electrolyte solutions in the prediction of Kováts retention indices of branched alkanes in alkane stationary phases 755(1996)49

Alkanesulfonates

Alkanesulfonate homologous series as eluent components in anion chromatography. Novel eluent selection system for monovalent analytes 771(1997)23

Alkannin derivatives

Quantitative determination of naphthaquinones of *Arnebia densiflora* (Nordm.) Ledeb. by an improved high-performance liquid chromatographic method 782(1997)133

Alkanols

Determination of organic molecular markers in marine aerosols and sediments: one-step flash chromatography compound class fractionation and capillary gas chromatographic analysis 799(1998)215

Alkenes

Influence of conformation on the accuracy of quantitative–structure retention relationship calculations in gas chromatography 773(1997)368

Alkenones

Clean-up procedures for the unbiased estimation of C₃₇ alkenone sea surface temperatures and terrigenous *n*-alkane inputs in paleoceanography 757(1997)145

Alkyl aryl ketones

Determination of diffusion coefficients and separation numbers in micellar electrokinetic chromatography 766(1997)187

Alkylbenzenes

Effect of skeleton size on the performance of octadecylsilylated continuous porous silica columns in reversed-phase liquid chromatography 762(1997)135

Application of 0.5- μ m porous silanized silica beads in electrochromatography 786(1997)229

Temperature programming and gradient elution in reversed-phase chromatography with packed capillary columns 788(1997)51

Novel polymeric resins for anion-exchange chromatography 793(1998)231

Comparative studies of chromatographic properties of silica-based amide-bonded phases under hydro–organic conditions 797(1998)103

Alkylbenzenesulfonates

High-performance liquid chromatographic analysis of a linear alkylbenzenesulfonate and its environmental biodegradation metabolites 760(1997)285

Determination of alkylphenols, alkylphenolethoxylates and linear alkylbenzenesulfonates in sediments by accelerated solvent extraction and supercritical fluid extraction 775(1997)187

Alkyl bromides

Use of fluorine-containing stationary phases for the separation of an alkyl bromide from its hydrocarbon analog by high-performance liquid chromatography 762(1997)159

Alkylglucosides

Capillary electrophoresis of herbicides. II. Evaluation of alkylglucoside chiral surfactants in the enantiomeric separation of phenoxy acid herbicides 757(1997)263

Alkylglycosides

Porous graphitized carbon and octadecyl-silica columns in the separation of some alkylglycoside detergents 782(1997)191

Alkyllead compounds

Speciation of trimethyllead in rainwater by gas chromatography–mass spectrometry after ethylation with sodium tetraethylborate. Comparison with other alkylation methods 779(1997)299

Alkylphenolethoxylates

Determination of alkylphenols, alkylphenolethoxylates and linear alkylbenzenesulfonates in sediments by accelerated solvent extraction and supercritical fluid extraction 775(1997)187

Alkylphenols

Determination of alkylphenols, alkylphenolethoxylates and linear alkylbenzenesulfonates in sediments by accelerated solvent extraction and supercritical fluid extraction 775(1997)187

Novel polymeric resins for anion-exchange chromatography 793(1998)231

Alkyl phenyl ketones

Selectivity of polymeric and polymer-supported pseudo-stationary phases in micellar electrokinetic chromatography 792(1997)105

Alkylphosphonic acids

Application of liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry, and tandem mass spectrometry, to the analysis and identification of degradation products of chemical warfare agents 759(1997)79

Capillary electrophoresis separation of alkylphosphonic acid monoesters with indirect ultraviolet detection 779(1997)245

Alkylpoly(ethylene glycol) ethers

Capillary gas chromatography–mass spectrometry of lower oxyethylenated aliphatic alcohols 773(1997)219

Capillary gas chromatography of *n*-butyl and isobutyl-, *n*-amyl and isoamyl polyethylene glycol ethers and their derivatives 800(1998)305

Alkylsulfonates

Capillary electrophoresis of cationic and anionic surfactants with indirect conductivity detection 781(1997)533

Alkyltrimethylammonium bromide

Quantitation of alkyltrimethylammonium bromides in Bayer process liquors by gas chromatography and gas chromatography–mass spectrometry 765(1997)287

Alkynes

Normal-phase and reversed-phase high-performance liquid chromatographic behaviour and particle beam mass spectrometric characterization of alkyne-substituted iron–carbonyl clusters 789(1997)461

Allenic acid

Predictable chromatographic separations of enantiomers: aryl allenic acids and their derivatives 761(1997)65

Allyl bromide

Bifunctional etherification of a bead cellulose for ligand attachment with allyl bromide and allyl glycidyl ether 775(1997)29

Allyl glycidyl ether

Bifunctional etherification of a bead cellulose for ligand attachment with allyl bromide and allyl glycidyl ether 775(1997)29

Allylic alcohols

Predictive strategies for determining retention indices of some allylic alcohols and their esters by gas chromatography 766(1997)141

Allyl terguride

Reversed determination of the formation constants of 1-allyl terguride with mandelic acid optical isomers using capillary electrophoresis 772(1997)179

Alprazolam

Simulations of liquid chromatography–diode array detector data including instrumental artefacts for the evaluation of mixture analysis techniques 758(1997)1

Use of simulated liquid chromatography–diode array detection data for the definition of a guide curve in peak purity assessment by spectral comparison 786(1997)1

Alprenolol

Robustness of the chromatographic separation of alprenolol and related substances using a silica-based stationary phase and selective retention of metoprolol and related substances on a porous graphitic carbon stationary phase 797(1998)75

Alternariol

Gas chromatography–mass spectrometry of *Alternaria* mycotoxins 765(1997)255

Aluminium

Fluorometric determination of serum and urinary aluminium with 8-quinolinol by kinetic-differentiation-mode micellar chromatography 789(1997)361

Amaranth

Identification and determination of red dyes in confectionery by ion-interaction high-performance liquid chromatography 767(1997)87

Amides

Reaction of fatty acid amides and ethanolamides with trimethylsulfonium hydroxide 773(1997)239

Amidines

Identification of hexahydroimidazo[1,2-*a*]pyrazine-3,6-diones and hexahydroimidazo[1,2-*a*]imidazo[1,2-*d*]pyrazine-3,8-diones, unusual products of silica-catalyzed amino acid thermal condensation and products of their thermal decomposition using coupled high-performance liquid chromatography–particle beam mass spectrometry and gas chromatography–Fourier transform infrared spectroscopy–mass spectrometry 776(1997)255

Amines

see also Biogenic amines and Catecholamines

pH-Dependent elution behaviour of meleme in high-performance cation-exchange chromatography 752(1996)287

Determination of aromatic amines at trace levels by derivatization with heptafluorobutyric anhydride and gas chromatography–electron-capture negative-ion chemical ionization mass spectrometry 753(1996)91

Separation of enantiomers and isomers of amino compounds by capillary electrophoresis and high-performance liquid chromatography utilizing crown ethers 757(1997)225

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

- Potentiometric detection of aliphatic amines by flow injection analysis and ion-interaction chromatography with a metallic copper electrode 758(1997)227
- Rapid determination of amphetamines by high-performance liquid chromatography with UV detection 761(1997)71
- Separation of primary and secondary amines as their sulfonamide derivatives by reversed-phase high-performance liquid chromatography 763(1997)173
- Analysis of foods for heterocyclic aromatic amine carcinogens by solid-phase extraction and high-performance liquid chromatography 763(1997)179
- Analysis of primary and secondary aliphatic amines in waste water and surface water by gas chromatography–mass spectrometry after derivatization with 2,4-dinitrofluorobenzene or benzenesulfonyl chloride 764(1997)85
- Determination of heterocyclic aromatic amines by micellar electrokinetic chromatography with amperometric detection 765(1997)329
- Determination of twenty eight biogenic amines and amino acids during wine aging by micellar electrokinetic chromatography and laser-induced fluorescence detection 765(1997)337
- Determination of residual amines used in bulk drug synthesis by pre-column derivatization with 3,5-dinitrobenzoyl chloride and high-performance liquid chromatography 766(1997)77
- Adsorption liquid chromatography on silica for the chiral separation of amino acids and asymmetric amines derivatized with optically active N- α -9-Fmoc-amino acid-N-carboxyanhydrides 767(1997)69
- Analysis of heterocyclic amines as their N-dimethylaminomethylene derivatives by gas chromatography with nitrogen–phosphorus selective detection 767(1997)187
- Electroosmotic flow mobilities in open and packed capillaries with spermine and other amine flow modifiers 768(1997)3
- Determination of heterocyclic aromatic amines by capillary zone electrophoresis in a meat extract 769(1997)293
- Optimization of resolution in micellar electrokinetic chromatography via computer-aided variation of concentrations of sodium dodecyl sulfate and acetonitrile as modifier 772(1997)27
- Determination of amines in air and water using derivatization combined with solid-phase microextraction 773(1997)249
- Liquid chromatography–electrospray mass spectrometry with in-source fragmentation for the identification and quantification of fourteen mutagenic amines in beef extracts 775(1997)125
- High-performance liquid chromatography–atmospheric pressure negative chemical ionisation mass spectrometry of 2,4-dinitrophenyl derivatives of amines. Applications in epoxy chemistry 777(1997)193
- Liquid chromatography–atmospheric-pressure chemical ionization mass spectrometry as a routine method for the analysis of mutagenic amines in beef extracts 778(1997)207
- Synthesis and characterization of a strong cation exchanger based on polymer-coated silica for high-performance liquid chromatography 779(1997)113
- Achievement of enantioselectivity of non-polar primary amines by a non-chiral crown ether 781(1997)129
- New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259
- Determination of histamine and some other amines by high-performance capillary electrophoresis with on-line mode in-capillary derivatization 790(1997)177
- Highly efficient separation of amines by electrokinetic chromatography using resorcarene-octacarboxylic acids as pseudostationary phases 792(1997)143
- Simultaneous use of urea and acetonitrile as organic modifiers for optimization of resolution in micellar electrokinetic chromatography 792(1997)157
- Mechanism of capillary electrophoresis enantioseparations using a combination of an achiral crown ether plus cyclodextrins 793(1998)115
- Improved procedure for *n*-hexyl chloroformate-mediated derivatization of highly hydrophilic substances directly in water: hydroxyaminic compounds 793(1998)307
- Liquid chromatography with luminol-based electrochemiluminescence detection. Determination of histamine 799(1998)57
- Amines, aromatic**
Determination of aromatic amines by solid-phase microextraction and gas chromatography–mass spectrometry in water samples 791(1997)221
- Amines, heterocyclic**
Analysis of heterocyclic amines as their N-dimethylaminomethylene derivatives by gas chromatography with nitrogen–phosphorus selective detection 767(1997)187
- Methods for the determination of mutagenic heterocyclic amines and their applications in environmental analysis 774(1997)121
- Liquid chromatography–electrospray mass spectrometry with in-source fragmentation for the identification and quantification of fourteen mutagenic amines in beef extracts 775(1997)125

Liquid chromatography–atmospheric-pressure chemical ionization mass spectrometry as a routine method for the analysis of mutagenic amines in beef extracts 778(1997)207

Amines, heterocyclic aromatic

Analysis of foods for heterocyclic aromatic amine carcinogens by solid-phase extraction and high-performance liquid chromatography 763(1997)179

Determination of heterocyclic aromatic amines by micellar electrokinetic chromatography with amperometric detection 765(1997)329

Analysis of heterocyclic aromatic amines in wine by high-performance liquid chromatography–electrospray tandem mass spectrometry 791(1997)71

Amino acids

pH-Zone-refining countercurrent chromatography 753(1996)1

Chiral crown ether-anchored polysiloxanes as capillary gas chromatography stationary phases 753(1996)269

Enhanced temporal resolution for the microdialysis monitoring of catecholamines and excitatory amino acids using capillary electrophoresis with laser-induced fluorescence detection. Analytical developments and in vitro validations 755(1996)99

Ergot alkaloids as chiral selectors in capillary electrophoresis. Determination of the separation mechanism 755(1996)251

Rapid and simultaneous analysis of protein and non-protein amino acids as N(O,S)-isobutoxycarbonyl methyl ester derivatives by capillary gas chromatography 758(1997)167

Quantitative analysis of fluorinated ethylchloroformate derivatives of protein amino acids and hydrolysis products of small peptides using chemical ionization gas chromatography–mass spectrometry 759(1997)111

Enantioseparation of 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate-derivatized-amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives. I. Discussion of optimum separation conditions 759(1997)157

Enantioseparation of 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate-derivatized amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives. II. Evaluation of complexation constants 759(1997)167

Study of retention, efficiency and selectivity in chiral ligand-exchange chromatography with a dynamically coated stationary phase 761(1997)79

Partitioning of hydrophobic amino acids and oligopeptides in aqueous two-phase system containing self-aggregating block copolymer. Effects of temperature, salts and surfactants 761(1997)91

High-performance liquid chromatographic methods for separation of enantiomers of alicyclic β -amino acids 761(1997)103

Tryptophan analysis in peptides and proteins, mainly by liquid chromatography 763(1997)1

Using quaternary high-performance liquid chromatography eluent systems for separating 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate-derivatized amino acid mixtures 763(1997)11

Determination of molecular-mass distribution of food-grade protein hydrolyzates by size-exclusion chromatography and chemiluminescent nitrogen detection 763(1997)323

High sample throughput phosphoamino acid analysis of proteins separated by one- and two-dimensional gel electrophoresis 764(1997)201

Chiral chromatography of amino acids on porous graphitic carbon coated with a series of *N*-substituted *L*-phenylalanine selectors. Effect of the anchor molecule on enantioselectivity 765(1997)187

Determination of twenty eight biogenic amines and amino acids during wine aging by micellar electrokinetic chromatography and laser-induced fluorescence detection 765(1997)337

Nuclear magnetic resonance monitoring of centrifugal partition chromatography in pH-zone-refining mode 766(1997)255

Adsorption liquid chromatography on silica for the chiral separation of amino acids and asymmetric amines derivatized with optically active *N*- α -9-Fmoc-amino acid-*N*-carboxyanhydrides 767(1997)69

Mobility measurements on dansylated amino acids 768(1997)153

Determination of cysteine by capillary zone electrophoresis with end-column amperometric detection at a gold/mercury amalgam microelectrode without deoxygenation 769(1997)307

Analysis of aliphatic carboxylic acids and amino acids in effluents of landfills, composting plants and fermentation plants by ion-exclusion and ion-exchange chromatography 770(1997)229

Application of a colour analyzer in quantitative thin-layer chromatography 771(1997)251

Optical resolution of phenylthiohydantoin-amino acids by capillary electrophoresis for protein sequencing 771(1997)311

Molecularly imprinted ligand-exchange adsorbents for the chiral separation of underivatized amino acids 775(1997)51

- Enhanced capillary zone electrophoretic separation of dinitrophenyl-amino acid derivatives through control of electroosmotic flow by the buffer cation 775(1997)327
- Determination of available lysine in infant milk formulae by high-performance liquid chromatography 778(1997)235
- Study of binding media in works of art by gas chromatographic analysis of amino acids and fatty acids derivatized with ethyl chloroformate 778(1997)373
- Preparative isolation of nine indispensable amino acids from a single sample 778(1997)383
- Microcolumn liquid chromatography: instrumentation, detection and applications 779(1997)1
- Ultrasensitive near-infrared laser-induced fluorescence detection in capillary electrophoresis using a diode laser and avalanche photodiode 779(1997)185
- Computerized optimization of selectivity for direct capillary gas chromatographic multicomponent separations of enantiomers 779(1997)253
- Molecular recognition with micellar and micelle-like aggregates in aqueous media 780(1997)179
- Micelles for signal enhancement and novel selectivity of dansyl amino acids 780(1997)219
- Role of alkyl and aryl substituents in chiral ligand exchange chromatography of amino acids. Study using porous graphitic carbon coated with N-substituted-L-proline selectors 786(1997)249
- Enantiomeric determination of amino compounds with high sensitivity using the chiral reagents (+)- and (-)-1-(9-anthryl)-2-propyl chloroformate 786(1997)347
- New 1,3,5-triazine based chiral stationary phase for the high-performance liquid chromatographic separation of enantiomers 786(1997)355
- Gas chromatography-chemical ionization mass spectrometry in amino acid analysis of pyoverdins 787(1997)195
- Identification of nonprotein amino acids from cycad seeds as N-ethoxycarbonyl ethyl ester derivatives by positive chemical-ionization gas chromatography-mass spectrometry 787(1997)288
- Determination of amino acids by high-performance capillary electrophoresis with on-line mode in-capillary derivatization 787(1997)253
- Analysis of organic and inorganic selenium anions by ion chromatography-inductively coupled plasma atomic emission spectroscopy 788(1997)105
- Empirical equation for the accurate prediction of retention in planar chromatography 788(1997)207
- Selectivity control in the non-aqueous capillary electrophoretic separation of amino acids 790(1997)185
- Chiral separation principles in capillary electrophoresis 792(1997)179
- Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309
- Influence of the degree of substitution of cyclodextrin sulfobutyl ether derivatives on enantioselective separations by electrokinetic chromatography 792(1997)379
- High-performance liquid chromatographic separation of enantiomers of unusual amino acids on a teicoplanin chiral stationary phase 793(1998)283
- Enantiomeric separation of amino acids by capillary electrophoresis with α -cyclodextrin 793(1998)414
- Indirect separation of chiral proteinogenic α -amino acids using the fluorescence active (1R,2R)-N-[(2-isothiocyanato)cyclohexyl]-6-methoxy-4-quinolinylamide) as chiral derivatizing agent. A comparison 795(1998)251
- Effect of the hydrolysis method on the determination of the amino acid composition of proteins 795(1998)263
- Chromatographic resolution of synthetically useful chiral glycine derivatives by high-performance liquid chromatography 796(1998)299
- Chromatographic methods for monitoring the optical isomers of unusual aromatic amino acids 797(1998)165
- High-performance liquid chromatographic methods for separation of the isomers of β -amino acids possessing bicyclo[2.2.1]heptane or heptene skeletons 797(1998)177
- Automated amino acid determination by high-performance liquid chromatography with 2-(9-anthryl)ethyl chloroformate as precolumn reagent 798(1998)1
- Fully automated precolumn derivatization, on-line dialysis and high-performance liquid chromatographic analysis of amino acids in food, beverages and feedstuff 798(1998)9
- Simultaneous determination of total and extracellular concentrations of the amino acid neurotransmitters in cat visual cortex by microbore liquid chromatography and electrochemical detection 798(1998)19
- Urinary excretion measurement of cysteine and homocysteine in the form of their S-pyridinium derivatives by high-performance liquid chromatography with ultraviolet detection 798(1998)27

Comparison of the stability and UV and fluorescence characteristics of the *o*-phthalaldehyde/3-mercaptopropionic acid and *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents and those of their amino acid derivatives 798(1998)37

Improved ion chromatography–integrated pulsed amperometric detection method for the evaluation of biogenic amines in food of vegetable or animal origin and in fermented foods 798(1998)109

Capillary electrophoresis and electrochemical detection with a conventional detector cell 798(1998)289

Separation and direct UV detection of sugars by capillary electrophoresis using chelation of copper(II) 799(1998)283

Separation of amino acid and peptide stereoisomers by nonionic micelle-mediated capillary electrophoresis after chiral derivatization 800(1998)345

Amino acids, dansyl derivatives

Sensitive absorbance detection method for capillary electrophoresis based on laser wave-mixing 773(1997)291

Amino acids, PTH derivatives

Optical resolution of phenylthiohydantoin-amino acids by capillary electrophoresis and identification of the phenylthiohydantoin-D-amino acid residue of [D-Ala²]-methionine enkephalin 752(1996)279

Amino alcohols

Enantiomeric separation of diols and β -amino alcohols by chiral stationary phase derived from (*R,R*)-tartaramide 757(1997)73

Aminoantipyrine

Simultaneous determination of chlorine dioxide and hypochlorite in water by high-performance liquid chromatography 796(1998)397

Amino-N-(4-(1,2-benzisothiazol-3-yl)-1-piperazinyl)butyl benzamide

Use of liquid chromatography–tandem mass spectrometry for the quantitative and qualitative analysis of an antipsychotic agent and its metabolites in human plasma and urine 794(1998)27

N-(4-Aminobenzoyl)-L-glutamic acid

Capillary zone electrophoresis of oligosaccharides derivatized with N-(4-aminobenzoyl)-L-glutamic acid for ultraviolet absorbance detection 757(1997)215

Amino(bis)phosphonates

Derivatization of pamidronate and other amino(bis)phosphonates with different isothiocyanates prior to ion-pair liquid chromatography 782(1997)211

2-Aminobutanoic acid

Computerized optimization of selectivity for direct capillary gas chromatographic multicomponent separations of enantiomers 779(1997)253

Aminobutylethylisoluminol isothiocyanate

Liquid chromatography with luminol-based electrochemiluminescence detection. Determination of histamine 799(1998)57

Aminocresols

Separation of enantiomers and isomers of amino compounds by capillary electrophoresis and high-performance liquid chromatography utilizing crown ethers 757(1997)225

Aminocyclopropanecarboxylic acid

Quantitative analysis of 1-aminocyclopropane-1-carboxylic acid by liquid chromatography coupled to electrospray tandem mass spectrometry 775(1997)143

2-Amino-1,2-diphenylethanol

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Aminoglutethimide

N-(Polyethyleneglycol monomethyl ether)-*N*-methylmorpholinium-based background electrolytes in capillary electrophoresis 763(1997)253

Applications of simulated moving-bed chromatography to the separation of the enantiomers of chiral drugs 769(1997)101

Aminoisobutyric acid, binaphthyl substituted

High-performance liquid chromatographic separation of enantiomers of 1,1'-binaphthyl-substituted α -aminoisobutyric acid 790(1997)41

7-Amino-4-methylcoumarine (Coumarine 440)

Highly sensitive detection using laser two-photon excited fluorescence in capillary electrophoresis 765(1997)315

Aminonaphthalenedisulfonic acid

Capillary electrophoresis of carboxylated carbohydrates. IV. Adjusting the separation selectivity of derivatized carboxylated carbohydrates by controlling the electrolyte ionic strength at subambient temperature and in the absence of electroosmotic flow 792(1997)75

Aminonaphthalenesulfonic acid

Capillary electrophoretic analysis of cyclodextrins with dynamic fluorescence labeling and detection 800(1998)333

Aminonaphthalene trisulfonate

Characterization of dextrans by size-exclusion chromatography on unmodified silica gel columns, with light-scattering detection, and capillary electrophoresis with laser-induced fluorescence detection 790(1997)93

Aminophenols

Separation of enantiomers and isomers of amino compounds by capillary electrophoresis and high-performance liquid chromatography utilizing crown ethers 757(1997)225

2-Amino-4-picoline

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Aminopolycarboxylic acids

Studies on the combination of ion chromatography–particle-beam mass spectrometry with capillary columns 770(1997)59

Capillary electrophoresis in metal analysis. Investigations of multi-elemental separation of metal chelates with aminopolycarboxylic acids 773(1997)351

Aminopyridine

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

5-Aminosalicylic acid

Simple liquid chromatographic method for the analysis of 5-aminosalicylic acid and its degradation product 765(1997)233

4-(Aminosulfonyl)-7-fluoro-2,1,3-benzoxadiazole

Removal of the fluorescent 4-(aminosulfonyl)-2,1,3-benzoxadiazole label from cysteine-containing peptides 798(1998)47

Amlodipine

Effect of charged and uncharged chiral additives on the resolution of amlodipine enantiomers in liquid chromatography and capillary electrophoresis 797(1998)187

Ammonium

Nonaqueous anion-exchange chromatography. II. Changing anion-exchange selectivity by resin surface complex formation of crown ethers 758(1997)29

Water clusters studied by electrospray mass spectrometry 777(1997)155

Column switching for difficult cation separations 789(1997)149

Amoxicillin

Development of a simple liquid chromatographic method with UV and mass spectrometric detection for the separation of substances related to amoxicillin sodium 797(1998)311

Amphetamines

Rapid determination of amphetamines by high-performance liquid chromatography with UV detection 761(1997)71

Urinary excretion of deprenyl metabolites 762(1997)321

Application of micellar electrokinetic capillary chromatography to the analysis of illicit drug seizures 780(1997)265

Sensitive and selective determination of methylenedioxyated amphetamines by high-performance liquid chromatography with fluorimetric detection 787(1997)137

Determination of methamphetamine and related compounds by capillary electrophoresis with UV and laser-induced fluorescence detection 798(1998)325

Amphiphiles

Theoretical aspects on the regulation of electroosmotic flow in capillary electrophoresis by adding charged amphiphiles 776(1997)311

Amylase inhibitors

Detection of electrophoretically separated amylase inhibitors in starch–polyacrylamide gels 752(1996)261

Characterization of protein glycoforms by capillary-zone electrophoresis–nanoelectrospray mass spectrometry 794(1998)327

Amylopectin

Calibration of a size-exclusion chromatography system using fractions with defined amylopectin unit chains 768(1997)325

Quantitative analysis of amylopectin unit chains by means of high-performance anion-exchange chromatography with pulsed amperometric detection 800(1998)199

Amylose

3,5-Dimethylphenylcarbamates of amylose, chitosan and cellulose bonded on silica gel. Comparison of their chiral recognition abilities as high-performance liquid chromatography chiral stationary phases 796(1998)265

Approach to the thermodynamics of enantiomer separation by gas chromatography. Enantioselectivity between the chiral inhalation anesthetics enflurane, isoflurane and desflurane and a diluted γ -cyclodextrin derivative 757(1997)119

Anaesthetics

Preparative enantiomer separation of the inhalation anesthetics enflurane, isoflurane and desflurane by gas chromatography on a derivatized γ -cyclodextrin stationary phase 769(1997)119

Determination of sedatives and anesthetics in plasma by liquid chromatography–mass spectrometry with a desalting system 797(1998)227

Anatoxins

Sensitive determination of anatoxin-a, homoanatoxin-a and their degradation products by liquid chromatography with fluorimetric detection 798(1998)147

Angiotensin I

Cyclodextrin aided separation of peptides and proteins by capillary zone electrophoresis 796(1998)367

Anilines

Chiral crown ether-anchored polysiloxanes as capillary gas chromatography stationary phases 753(1996)269

Combined use of temperature and solvent strength in reversed-phase gradient elution. III. Selectivity for ionizable samples as a function of sample type and pH 756(1996)51

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Simultaneous separation of metal ions and neutral organics by reversed-phase ion-pair liquid chromatography 776(1997)221

Determination of aromatic amines by solid-phase microextraction and gas chromatography–mass spectrometry in water samples 791(1997)221

Effects of tetraalkylammonium salts on the micellar electrokinetic chromatography of aniline and substituted anilines 799(1998)289

Anions

see also Inorganic anions and Organic anions

Ion chromatography methods for the simultaneous determination of mineral anions in plant sap 752(1996)298

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. I. Large-volume stacking with polarity switching 757(1997)281

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. II. Large-volume stacking without polarity switching 757(1997)291

Capillary electrophoretic determination of inorganic and organic anions using 2,6-pyridinedicarboxylic acid: effect of electrolyte's complexing ability 767(1997)223

Determination of small anions by capillary electrophoresis using indirect UV detection with sulphonated nitrosonaphthol dyes 767(1997)293

Calculation of detection limits for a single-laboratory ion-chromatographic method to determine parts-per-trillion ions in ultrapure water 770(1997)105

High-performance ion chromatography applied to free-radical mechanisms in drug design. The problem of ion analysis at high ionic strengths 770(1997)273

Determination and prediction of transfer ratios for anions in capillary zone electrophoresis with indirect UV detection 770(1997)291

Alkanesulfonate homologous series as eluent components in anion chromatography. Novel eluent selection system for monovalent analytes 771(1997)23

Non-aqueous capillary electrophoretic separation of Brønsted acids as heteroconjugated anions 771(1997)275

Comparisons of the mobilities of salt ions obtained by the moving boundary method and two empirical equations in capillary electrophoresis 771(1997)374

Modified electroosmotic flow by cationic surfactant additives in capillary electrophoresis. Evaluation of electrolyte systems for anion analysis 772(1997)171

Development of new methods for the analysis of single cloud-, fog- and raindrops by capillary electrophoresis 775(1997)372

Capillary electrophoresis as a routine industrial tool for quantitative analytical testing. Determination of sodium dimethyldithiocarbamate in effluents 781(1997)541

Flow injection–capillary electrophoresis coupling to automate on-line sample treatment for the determination of inorganic ions in waters 791(1997)279

Ultrasonic effects on ion-exchange chromatographic retention 793(1998)365

Studies on elution conditions for the determination of anions by suppressed ion-interaction chromatography using a graphitized carbon column 800(1998)239

Anisole

Influence of the concentration of adsorbate and modifier in the mobile phase on retention in high-performance liquid chromatography 797(1998)3

Anthocyanins

Preparative separation of anthocyanins by gradient elution centrifugal partition chromatography 763(1997)345

Separation of blackcurrant anthocyanins by capillary zone electrophoresis 799(1998)321

1-Anthrolylnitrile

Determination of atropine in pharmaceutical preparations by liquid chromatography with fluorescence detection 775(1997)137

1-(9-Anthryl)-2-propyl chloroformate

Enantiomeric determination of amino compounds with high sensitivity using the chiral reagents (+)- and (–)-1-(9-anthryl)-2-propyl chloroformate 786(1997)347

Anthryldiazomethane

Optimization of sample cleanup procedure for determination of diarrhetic shellfish poisoning toxins by use of experimental design 764(1997)223

Anthrylethyl chloroformate

Automated amino acid determination by high-performance liquid chromatography with 2-(9-anthryl)ethyl chloroformate as precolumn reagent 798(1998)1

Anthryltrifluoroethanol

Influence of the degree of substitution of cyclodextrin sulfobutyl ether derivatives on enantioselective separations by electrokinetic chromatography 792(1997)379

Anti-AIDS drugs

Application of spherical and other polymers in capillary zone electrophoresis: separation of antiviral drugs and deoxyribonucleoside phosphates by different principles 756(1996)263

New procedure for the use of high-performance liquid chromatography–electrospray ionization mass spectrometry for the analysis of nucleotides and oligonucleotides 777(1997)3

Antibacterial agents

Identification of an anti-bacterial agent in toothpaste via liquid chromatography–Fourier transform infrared spectrometry mobile phase elimination 755(1996)211

Optimization of capillary electrophoretic separation of quinolone antibacterials using the overlapping resolution mapping scheme 766(1997)215

Determination of antibacterial quaternary ammonium compounds in lozenges by capillary electrophoresis 798(1998)335

Antibiotics

Liquid chromatography of spiramycin on poly(styrene–divinylbenzene) 764(1997)43

Analysis of kanamycin sulfate by liquid chromatography with pulsed electrochemical detection 766(1997)133

Optimisation of a simultaneous separation of sulphonamides, dihydrofolate reductase inhibitors and β -lactam antibiotics by capillary electrophoresis 768(1997)97

Evaluation of liquid chromatography methods for the analysis of benzylpenicillin and its related substances 773(1997)147

Study of the binding of antibiotics to human serum albumin by charge-transfer chromatography 776(1997)31

Use of metal complexation in non-aqueous capillary electrophoresis systems for the separation and improved detection of tetracyclines 779(1997)235

Micellar electrokinetic capillary chromatography for the separation of phenoxymethylpenicillin and related substances 781(1997)417

Determination of the chemotherapeutic quinolonic and cinolonic derivatives in urine by high-performance liquid chromatography with ultraviolet and fluorescence detection in series 787(1997)119

Dual-mode high-speed counter-current chromatography: retention, resolution and examples 790(1997)17

Analysis of benzylpenicillin by capillary electrophoresis 792(1997)83

Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309

Development of a simple liquid chromatographic method with UV and mass spectrometric detection for the separation of substances related to amoxicillin sodium 797(1998)311

Determination of flumequine and oxolinic acid in fortified chicken tissue using on-line dialysis and high-performance liquid chromatography with fluorescence detection 799(1998)349

Residual solvents determination in the antibiotic L-749,345 by static headspace gas chromatography 800(1998)275

Antibodies

see also Monoclonal antibodies

Role of high-performance liquid chromatographic protein analysis in developing fermentation processes for recombinant human growth hormone, relaxin, antibody fragments and lymphotoxin 763(1997)31

Characterization of antigen–antibody complexes by size-exclusion chromatography coupled with low-angle light-scattering photometry and viscometry 787(1997)101

Antidepressants, tricyclic

Stability of silica-based, endcapped columns with pH 7 and 11 mobile phases for reversed-phase high-performance liquid chromatography 762(1997)97

Retention characteristics and selected applications of cyclic siloxane-based octadecylsilyl bonded phases in reversed-phase high-performance liquid chromatography 773(1997)53

Antigens

Use of monoclonal antibodies for weak affinity chromatography 758(1997)199

Kinetic analysis of the interaction between the monoclonal antibody A33 and its colonic epithelial antigen by the use of an optical biosensor. A comparison of immobilisation strategies 776(1997)15

Automated microanalysis using magnetic beads with commercial capillary electrophoretic instrumentation 781(1997)197

Characterization of antigen–antibody complexes by size-exclusion chromatography coupled with low-angle light-scattering photometry and viscometry 787(1997)101

Micro-sequencing strategies for the human A33 antigen, a novel surface glycoprotein of human gastrointestinal epithelium 798(1998)91

Antihistamines

Comparative capillary electrophoresis and NMR studies of enantioseparation of dimethindene with cyclodextrins 798(1998)315

Anti-inflammatory drugs, non-steroidal

Separation of ibuprofen, codeine phosphate, their degradation products and impurities by capillary electrophoresis. I. Method development and optimization with fractional factorial design 798(1998)307

Antimony

Selective determination of Sb(III) by gas chromatography–quartz furnace atomic absorption spectrometry after derivatization with triphenylmagnesium bromide 764(1997)169

Antioxidants

Identification of antioxidative compounds in plant beverages by capillary electrophoresis with the marker index technique 781(1997)523

Reversed-phase high-performance liquid chromatographic separations of tocopherols 782(1997)25

Antipsychotic agent

Use of liquid chromatography–tandem mass spectrometry for the quantitative and qualitative analysis of an antipsychotic agent and its metabolites in human plasma and urine 794(1998)27

Antipyrine

Characterization of ⁶⁰Co γ -radiation induced radical products of antipyrine by means of high-performance liquid chromatography, mass spectrometry, capillary zone electrophoresis, micellar electrokinetic capillary chromatography and nuclear magnetic resonance spectrometry 788(1997)95

Antiviral drugs

Application of spherical and other polymers in capillary zone electrophoresis: separation of antiviral drugs and deoxyribonucleoside phosphates by different principles 756(1996)263

Apolipoproteins

Rapid reversed-phase high-performance liquid chromatography method for quantitation, at high pH, of the recombinant apolipo-protein A-IMilano in *Escherichia coli* fermentation broth 798(1998)83

Aporphine alkaloids

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Arachidonic acid

Liquid chromatography–mass spectrometry with on-line solid-phase extraction by a restricted-access C₁₈ precolumn for direct plasma and urine injection 762(1997)193

Arbutin

Influence of differences between sample and mobile phase viscosities on the shape of chromatographic elution profiles 775(1997)73

Arenes

Unified retention concept – statistical treatment of Kováts retention index 764(1997)257

Aroclors

Chromatography pattern recognition of Aroclors using iterative probabilistic neural networks 775(1997)231

Aromatic compounds

see also Hydrocarbons, aromatic

Study of retention in reversed-phase liquid chromatography using linear solvation energy relationships. I. The stationary phase 752(1996)1

High-temperature open tubular liquid chromatography coupled to atmospheric pressure chemical ionisation mass spectrometry 755(1996)153

Refractive-index detection by interferometric backscatter in packed-capillary high-performance liquid chromatography 762(1997)219

New selectivity in electrokinetic chromatography using a polymeric dye as novel separation carrier 792(1997)151

Optimization of a supercritical fluid chromatograph–atmospheric pressure chemical ionization mass spectrometer interface using an ion trap and two quadrupole mass spectrometers 800(1998)39

Aromoline

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Arpromidine

Chiral separation of 3-phenyl-3-(2-pyridyl)propylamines, and analogous guanidines and guanidine-N-carboxylic acid esters with high-performance liquid chromatography and capillary zone electrophoresis 793(1998)77

Arsenazo I

Role of ligand purity in separations of alkaline earth metals as arsenazo I complexes by capillary zone electrophoresis 793(1998)177

Arsenic

Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217

Simultaneous determination of toxic arsenic and chromium species in water samples by ion chromatography–inductively coupled plasma mass spectrometry 779(1997)139

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

Arsenic acid

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. I. Large-volume stacking with polarity switching 757(1997)281

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. II. Large-volume stacking without polarity switching 757(1997)291

Arsenic compounds

High-performance liquid chromatography–hydride generation–atomic fluorescence spectroscopic determination of arsenic species in water 756(1996)292

Speciation of arsenic compounds by using ion-pair chromatography with atomic spectrometry and mass spectrometry detection 764(1997)55

Liquid chromatography and mass spectrometry for the speciation of arsenic animal feed additives 764(1997)211

Ion-exchange separation of eight arsenic compounds by high-performance liquid chromatography–UV decomposition–hydride generation–atomic fluorescence spectrometry and stability tests for food treatment procedures 789(1997)339

Arsenobetaine

Ion-exchange separation of eight arsenic compounds by high-performance liquid chromatography–UV decomposition–hydride generation–atomic fluorescence spectrometry and stability tests for food treatment procedures 789(1997)339

Artemisinic acid

Extraction of artemisinin and artemisinic acid from *Artemisia annua* L. using supercritical carbon dioxide 785(1997)353

Artemisinin

Extraction of artemisinin and artemisinic acid from *Artemisia annua* L. using supercritical carbon dioxide 785(1997)353

Artificial sweeteners

Separation and determination of four artificial sweeteners and citric acid by high-performance anion-exchange chromatography 771(1997)135

Determination of aspartame and phenylalanine in diet soft drinks by high-performance liquid chromatography with direct spectrofluorimetric detection 773(1997)163

Determination of aspartame by high-performance capillary electrophoresis 781(1997)423

Separation of alditols of interest in food products by high-performance anion-exchange chromatography with pulsed amperometric detection 791(1997)343

Arylhdyantoins

Facile and predictable means of separating the enantiomers of 5-arylhdyantoins 790(1997)65

Aryloxyphenoxypropionic acids

Determination of aryloxyphenoxypropionic acid herbicides in water using different solid-phase extraction procedures and liquid chromatography–diode array detection 796(1998)309

Aryloxypropionic acids

Enantiomeric separations of halogen-substituted 2-aryloxypropionic acids by high-performance liquid chromatography on a terguride-based chiral stationary phase 756(1996)119

Arylpropionic acids

Enantiomeric resolution study by capillary electrophoresis. Selection of the appropriate chiral selector 772(1997)185

Separation of profen enantiomers by capillary electrophoresis using cyclodextrins as chiral selectors 793(1998)165

Chiral capillary electrophoresis–electrospray mass spectrometry coupling using vancomycin as chiral selector 800(1998)69

N-Arylthiazolin-2(thi)ones

Investigation into the chiral recognition mechanism of N-arylthiazolin-2(thi)one atropisomers on Chiralcel OJ by factorial design and lipophilicity approaches 761(1997)129

Aryltrialkylammonium chlorides

N-(Polyethyleneglycol monomethyl ether)-N-methylmorpholinium-based background electrolytes in capillary electrophoresis 763(1997)253

Ascorbate

Analysis of ascorbate and dehydroascorbate in plant extracts by high-resolution selected ion monitoring gas chromatography–mass spectrometry 782(1997)95

Ascorbic acid

see also Vitamins

Simultaneous determination of qualitatively important components in green tea infusions using capillary electrophoresis 758(1997)332

Determination of ascorbic acid in vegetables by capillary zone electrophoresis 772(1997)313

Determination of L-ascorbic acid in foods by capillary zone electrophoresis 781(1997)435

Ascorbigen

Comparison of thin-layer chromatography and overpressured layer chromatographic techniques for the separation of ascorbigen and 1'-methylascorbigen 764(1997)103

Aspartame

Separation and determination of four artificial sweeteners and citric acid by high-performance anion-exchange chromatography 771(1997)135

Determination of aspartame and phenylalanine in diet soft drinks by high-performance liquid chromatography with direct spectrofluorimetric detection 773(1997)163

Determination of aspartame by high-performance capillary electrophoresis 781(1997)423

Analysis of beverages by capillary electrophoresis 781(1997)481

Aspartate

Enhanced temporal resolution for the microdialysis monitoring of catecholamines and excitatory amino acids using capillary electrophoresis with laser-induced fluorescence detection. Analytical developments and in vitro validations 755(1996)99

Atenolol

Direct injection of large volumes of plasma/serum on a new biocompatible extraction column for the determination of atenolol, propranolol and ibuprofen. Mechanisms for the improvement of chromatographic performance 797(1998)251

Atrazine

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Microbore liquid chromatographic-mass spectrometric determination of atrazine and its major hydroxylated degradate in water at parts-per-trillion levels using electrospray 753(1996)243

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography-Fourier-transform infrared spectrometry 756(1996)145

New and simple method for the analysis of the glutathione adduct of atrazine 795(1998)53

High sample throughput flow immunoassay utilising restricted access columns for the separation of bound and free label 800(1998)219

Atropine

High-performance liquid chromatographic determination of atropine and atropine-like alkaloids in pharmaceutical preparations with indirect conductometric detection 766(1997)270

Determination of atropine in pharmaceutical preparations by liquid chromatography with fluorescence detection 775(1997)137

Auxins

Determination of indole-3-acetic acid in the Gulf of Gdańsk by high-performance liquid chromatography of its 4-methyl-7-methoxycoumarin derivative 766(1997)261

Liquid chromatographic determination of β -naphthoxyacetic acid in tomatoes 798(1998)167

Azadirachtin

Comparison of chromatographic systems for triterpenoids from Neem (*Azadirachta indica*) seeds 761(1997)53

Azinphosmethyl

Gas chromatographic determination of airborne residues of azinphosmethyl and azinphosmethyl-oxon by cool on-column injection 775(1997)307

Azo dyes

Speciation of chromium dyes by high-performance liquid chromatography with inductively coupled plasma mass spectrometric detection 773(1997)157

Azo dyes, sulphonated

Determination of mono- and disulphonated azo dyes by liquid chromatography-atmospheric pressure ionization mass spectrometry 777(1997)177

B

Bacitracin

Effect of foaming power and foam stability on continuous concentration with foam counter-current chromatography 791(1997)53

Bafilomycins

Cationization in electrospray microcolumn liquid chromatography-mass spectrometry 794(1998)359

Barbiturates

Retention characteristics and selected applications of cyclic siloxane-based octadecylsilyl bonded phases in reversed-phase high-performance liquid chromatography 773(1997)53

Determination of barbiturates by solid-phase microextraction (SPME) and ion trap gas chromatography-mass spectrometry 777(1997)275

Superheated water as an eluent for reversed-phase high-performance liquid chromatography 785(1997)49

Barium

Determination of barium and strontium by capillary zone electrophoresis using an electrolyte containing sulfonazo III 767(1997)303

Basic compounds

Comparative evaluation of bonded-silica reversed-phase columns for high-performance liquid chromatography using strongly basic compounds and alternative organic modifiers buffered at acid pH 769(1997)169

Separation of polar solutes by packed column supercritical fluid chromatography (Review) 785(1997)3

Influence of sample mass on the performance of reversed-phase columns in the analysis of strongly basic compounds by high-performance liquid chromatography 793(1998)31

Reversed-phase high-performance liquid chromatography of basic compounds at pH 11 with silica-based column packings 797(1998)111

Basic drugs

Combined use of temperature and solvent strength in reversed-phase gradient elution. III. Selectivity for ionizable samples as a function of sample type and pH 756(1996)51

Monitoring of new silica-based reversed-phase stationary phases for the liquid chromatographic analysis of basic pharmaceuticals using principal components analysis 765(1997)157

Beclinazone

Liquid chromatographic separation of the enantiomers of beclinazone and its potential impurities 769(1997)231

Benomyl

High-performance liquid chromatographic determination of benomyl and carbendazim residues in apiarian samples 787(1997)129

Benzalkonium chloride

Analysis of benzalkonium chloride in the effluent from European hospitals by solid-phase extraction and high-performance liquid chromatography with post-column ion-pairing and fluorescence detection 774(1997)281

Synthesis and characterization of a strong cation exchanger based on polymer-coated silica for high-performance liquid chromatography 779(1997)113

Determination of antibacterial quaternary ammonium compounds in lozenges by capillary electrophoresis 798(1998)335

Benzenecarboxylic acids

Ion-exclusion chromatography of benzenecarboxylic acids on an unmodified silica-gel column 782(1997)33

Benzenes

Headspace gas chromatographic determination of the plant cuticle-air partition coefficients for monocyclic aromatic hydrocarbons as environmental compartment 774(1997)213

Analytical methods for the determination of organochlorine compounds. Application to environmental samples in the Slovak Republic 774(1997)333

Capacity ratios in supercritical fluid chromatography. Effect of mobile and stationary phases on hexasubstituted benzenes 776(1997)305

Evaluation of environmental levels of aromatic hydrocarbons in gasoline service stations by gas chromatography 778(1997)263

Retention modeling in micellar liquid chromatography 780(1997)149

Resorcinol derivative used as a new stationary phase for capillary gas chromatography 787(1997)161

Selectivity of polymeric and polymer-supported pseudo-stationary phases in micellar electrokinetic chromatography 792(1997)105

Effect of polystyrene-divinylbenzene resin sulfonation on solute retention in high-performance liquid chromatography 797(1998)139

Determination of volatile organic compounds in marine biota 799(1998)233

Benzenesulfonyl chloride

Analysis of primary and secondary aliphatic amines in waste water and surface water by gas chromatography-mass spectrometry after derivatization with 2,4-dinitrofluorobenzene or benzenesulfonyl chloride 764(1997)85

Benzenetetracarboxylic acid

Dynamic control and indirect absorption detection for high-speed capillary electrophoretic separation of organic acids 800(1998)339

Benzilic acid

Application of liquid chromatography-atmospheric pressure chemical ionisation mass spectrometry, and tandem mass spectrometry, to the analysis and identification of degradation products of chemical warfare agents 759(1997)79

Benzimidazoles

Simplified reversed-phase conditions for the determination of benzimidazole fungicides in fruits by high-performance liquid chromatography with UV detection 778(1997)95

Benzoates

Influence of sample plug width in capillary electrochromatography 779(1997)155

Analysis of beverages by capillary electrophoresis
781(1997)481

Benzodiazepine receptors

Size-exclusion chromatographic reconstitution of the bovine brain benzodiazepine receptor. Effects of lipid environment on the binding characteristics 776(1997)101

Benzodiazepines

Migration behaviour of benzodiazepines in micellar electrokinetic chromatography 756(1996)279

On-column enantiomerization of 3-hydroxybenzodiazepines during chiral liquid chromatography with optical rotation detection 767(1997)93

Gradient capillary electrochromatography of drug mixtures with UV and electrospray ionisation mass spectrometric detection 768(1997)89

Erratum to "On-column enantiomerization of 3-hydroxybenzodiazepines during chiral liquid chromatography with optical rotation detection". [J. Chromatogr. A, 767 (1997) 93–100] 782(1997)296

Regioselectively carbamoylated polysaccharides for the separation of enantiomers in high-performance liquid chromatography 786(1997)13

Separations of benzodiazepines using electrochemically modulated liquid chromatography. Efficient separations from changes in the voltage applied to a porous graphitic carbon stationary phase 793(1998)204

Solvent versatility of bonded cellulose-derived chiral stationary phases for high-performance liquid chromatography and its consequences in column loadability 793(1998)239

Benzoic acids

Comparative investigation of UV, electrochemical and particle beam mass spectrometric detection for the high-performance liquid chromatographic determination of benzoic and cinnamic acids and of their corresponding phenolic acids 753(1996)157

Combined use of temperature and solvent strength in reversed-phase gradient elution. III. Selectivity for ionizable samples as a function of sample type and pH 756(1996)51

Determination of preservatives in food products by cyclodextrin-modified capillary electrophoresis with multiwavelength detection 768(1997)334

Ultra-trace determination of fluorinated aromatic carboxylic acids in aqueous reservoir fluids using solid-phase extraction in combination with gas chromatography–mass spectrometry 793(1998)297

Chromatographic peak profile of ionogenic analytes upon elution with unbuffered eluents 797(1998)57

Benzoin

Effect of organic modifiers on retention and enantiomeric separations by capillary electrophoresis with human serum albumin as a chiral selector in solution 766(1997)237

Benzophenone

Resolution studies on counter-current chromatography using supercritical fluid carbon dioxide 790(1997)31

Benzosuberan derivatives

Chiral separations of indan, tetralin and benzosuberan derivatives by capillary electrophoresis 793(1998)135

Benzo[a]pyrene

Enrichment of benzo[a]pyrene in smoked food products and determination by high-performance liquid chromatography–fluorescence detection 753(1996)207

On-line preconcentration and high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbon–DNA adducts using copper phthalocyanine trisulfonic acid as a group-specific adsorbent 755(1996)205

Benzo[a]pyrene sulfates

Separation of benzo[a]pyrene sulfate isomers by reversed-phase liquid chromatography 753(1996)201

Benzoylphenylalanine

Enantiomeric properties of human albumin immobilized on porous silica supports coated with polymethacryloyl chloride 776(1997)37

N-Benzyl-N-alkyl-N,N-dimethylammonium chloride

Analysis of cationic surfactants by capillary electrophoresis 757(1997)275

Benzylamine

Capillary zone electrophoretic separation of biogenic amines: influence of organic modifier 755(1996)142

N-Benzylmethylamine

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Benzylpenicillin

Evaluation of liquid chromatography methods for the analysis of benzylpenicillin and its related substances 773(1997)147

Analysis of benzylpenicillin by capillary electrophoresis 792(1997)83

Berberamine

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Berberine

Determination of berberine and palmatine in *Phellodendri Cortex* using ion-pair supercritical fluid chromatography on-line coupled with ion-pair supercritical fluid extraction by on-column trapping 786(1997)371

Analysis of traditional Chinese anticancer drugs by capillary electrophoresis 795(1998)388

Bergapten

Retention of some components in supercritical fluid chromatography and application to bergamot peel oil fractionation 771(1997)241

 β -Agonists

Detection of clenbuterol in bovine retinal tissue by high-performance liquid chromatography with electrochemical detection 762(1997)275

Rapid pharmacokinetic screening of salbutamol in plasma samples by column-switching high-performance liquid chromatography–electrospray mass spectrometry 777(1997)67

 β -Blockers

Capillary zone electrophoretic separation of β -blockers using citrate buffer at low pH 753(1996)133

Chiral separations of β -blocking drug substances using the Pirkle-type α -Burke I chiral stationary phase 757(1997)65

Direct liquid chromatographic enantioseparation of sotalol and other β -blockers using an α_1 -acid glycoprotein-based chiral stationary phase 760(1997)193

On-line column-switching high-performance liquid chromatography analysis of cardiovascular drugs in serum with automated sample clean-up and zone-cutting technique to perform chiral separation 762(1997)235

Liquid chromatographic procedure for the evaluation of β -blockers in pharmaceuticals using hybrid micellar mobile phases 765(1997)221

Migration behavior and selectivity of β -blockers in micellar electrokinetic chromatography. Influence of micelle concentration of cationic surfactants 775(1997)349

Capillary zone electrophoretic separation of β -blockers using citrate buffer at low pH 776(1997)361

Reply to “Capillary zone electrophoretic separation of β -blockers using citrate buffer at low pH” 776(1997)362

Bonded cellulose-derived high-performance liquid chromatography chiral stationary phases. III. Effect of the reticulation of the cellulose derivative on performance 791(1997)37

Solvent versatility of bonded cellulose-derived chiral stationary phases for high-performance liquid chromatography and its consequences in column loadability 793(1998)239

Enantiomeric separation of propranolol and selected metabolites by using capillary electrophoresis with hydroxypropyl- β -cyclodextrin as chiral selector 793(1998)357

Robustness of the chromatographic separation of alprenolol and related substances using a silica-based stationary phase and selective retention of metoprolol and related substances on a porous graphitic carbon stationary phase 797(1998)75

Direct injection of large volumes of plasma/serum on a new biocompatible extraction column for the determination of atenolol, propranolol and ibuprofen. Mechanisms for the improvement of chromatographic performance 797(1998)251

Fluorescence detection in packed-column supercritical fluid chromatographic separations 798(1998)203

Evaluation of the immobilized artificial membrane phosphatidylcholine. Drug discovery column for high-performance liquid chromatographic screening of drug–membrane interactions 800(1998)161

Bicycloheptane amino acids

High-performance liquid chromatographic methods for separation of the isomers of β -amino acids possessing bicyclo[2.2.1]heptane or heptene skeletons 797(1998)177

Bifenthrin

Enantiomeric resolution of pyrethroids by high-performance liquid chromatography with diode-laser polarimetric detection 754(1996)97

Bile acids

Capillary gas–liquid chromatography of acetate–methyl esters of bile acids 766(1997)286

Microcolumn liquid chromatography: instrumentation, detection and applications 779(1997)1

Bile salts

Bile salt micellar electrokinetic chromatography of bilirubin and related compounds 779(1997)205

Investigation of the interactions between drugs and mixed bile salt/lecithin micelles. A characterization by micellar affinity capillary electrophoresis (MACE). Part III 781(1997)377

Biliproteins

Chromatographic purification of biliproteins from *Spirulina platensis*. High-performance liquid chromatographic separation of their α and β subunits 778(1997)441

Bilirubins

Bile salt micellar electrokinetic chromatography of bilirubin and related compounds 779(1997)205

High resolution multichannel fluorescence detection for capillary electrophoresis. Application to multicomponent analysis 788(1997)173

Biliverdin

Bile salt micellar electrokinetic chromatography of bilirubin and related compounds 779(1997)205

Binaphthol

Modeling, simulation and operation of a simulated moving bed for continuous chromatographic separation of 1,1'-bi-2-naphthol enantiomers 769(1997)25

Influence of the degree of substitution of cyclodextrin sulfobutyl ether derivatives on enantioselective separations by electrokinetic chromatography 792(1997)379

Binaphthyl diyl hydrogenphosphate

Enantioseparation using selected polysaccharides as chiral buffer additives in capillary electrophoresis 773(1997)331

Biochanin A

Development and validation of an isocratic high-performance liquid chromatographic method for quantitative determination of phytoestrogens in soya bean 795(1998)377

Biochenin

Linearity and the limitations of least squares calibration 762(1997)73

Biogenic amines

see also Amines and Catecholamines

Capillary zone electrophoretic separation of biogenic amines: influence of organic modifier 755(1996)142

Determination of twenty eight biogenic amines and amino acids during wine aging by micellar electrokinetic chromatography and laser-induced fluorescence detection 765(1997)337

Improved ion chromatography-integrated pulsed amperometric detection method for the evaluation of biogenic amines in food of vegetable or animal origin and in fermented foods 798(1998)109

Biotin

Applying hollow fibres for separating free and bound label in continuous-flow immunochemical detection 755(1996)179

Implementation of affinity solid-phases in continuous-flow biochemical detection 776(1997)169

Theoretical concepts of on-line liquid chromatographic-biochemical detection systems. II. Detection systems based on labelled affinity proteins 787(1997)37

Bisbenzylisoquinolines

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Bismuth

High-performance chelation ion chromatography for the determination of traces of bismuth in lead by means of a novel hypercrosslinked polystyrene resin 789(1997)389

Bisphenol-A, dipropoxylated, polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. II. Solubility effects in the analysis of polyester resins under reversed-phase conditions 761(1997)147

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions 775(1997)157

Bisphenol A diglycidyl ether

Identification of potential migrants from a vinylic organosol varnish by gas chromatography-mass spectrometry and liquid chromatography-mass spectrometry 771(1997)366

Bis(1-phenylethyl) dithioacetals

Determination of the absolute configuration of sugar residues using gas chromatography. Method with potential for elimination of references 769(1997)253

Bisphosphonates

Determination of impurities in clodronic acid by anion-exchange chromatography 770(1997)261

Bitter acids

Preparative separation of bitter acids from hop extracts by centrifugal partition chromatography 771(1997)71

Block copolymers

Theory of liquid adsorption chromatography of macromolecules applied to polyethylene glycol and fatty alcohol ethoxylates 791(1997)21

BODIPY-hydrazide

Hydrazide as a ligand moiety in immobilized metal ion affinity chromatography. Separation of BO-IMI and BODIPY-hydrazide 777(1997)261

Boiling phase

Letter to the editor 771(1997)385

Boldine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Boranes

High-performance liquid chromatographic separations of boron-cluster compounds (Review) 789(1997)497

Borate

Chiral resolution of diols by capillary electrophoresis using borate-cyclodextrin complexation 761(1997)269

BO-IMI

Hydrazide as a ligand moiety in immobilized metal ion affinity chromatography. Separation of BO-IMI and BODIPY-hydrazide 777(1997)261

Bradykinins

Determination of peptides by high-performance liquid chromatography with laser-induced fluorescence detection 763(1997)23

Analysis of bioactive peptides by liquid chromatography-high-resolution electrospray mass spectrometry 767(1997)77

Brij 35

Separation of unsaturated fatty acids and related isomeric hydroperoxides by micellar electrokinetic chromatography 781(1997)215

Composition analysis of detergents of the polyoxyethylene type: comparison of thin-layer chromatography, reversed-phase chromatography and matrix-assisted laser desorption/ionization mass spectrometry 791(1997)245

Bromacil

Direct determination of bromacil and diuron residues in environmental water samples by coupled-column liquid chromatography and large-volume injection 761(1997)322

Bromate

Determination of bromate in drinking waters by ion chromatography with inductively coupled plasma mass spectrometric detection 753(1996)261

Correlation and digital signal processing techniques for the reduction of detection limits of bromate and bromide in model water samples by ion chromatography with direct ultraviolet detection 789(1997)247

Specific determination of bromate and iodate in ozonized water by ion chromatography with postcolumn derivatization and inductively-coupled plasma mass spectrometry 789(1997)259

Bromide

Simultaneous determination of bromide and iodide ions by capillary isotachopheresis using quaternary ammonium salts 760(1997)253

Correlation and digital signal processing techniques for the reduction of detection limits of bromate and bromide in model water samples by ion chromatography with direct ultraviolet detection 789(1997)247

Bromine

Determination of chlorine and bromine in rocks by alkaline fusion with ion chromatography detection 770(1997)93

Bromoacetylpyrene

Improvement on sample clean-up for high-performance liquid chromatographic-fluorimetric determination of diarrhetic shellfish toxins using 1-bromoacetylpyrene 793(1998)63

Bromocriptine

High-temperature open tubular liquid chromatography coupled to atmospheric pressure chemical ionisation mass spectrometry 755(1996)153

Bromodeoxyuridine

New hydrolytic products detected in aqueous solutions of bromodeoxyuridine 797(1998)283

2-[(5-Bromo-2-pyridyl)azo]-5-diethylaminophenol

On-line pretreatment and determination of Pb, Cu and Cd at the $\mu\text{g l}^{-1}$ level in drinking water by chelation ion chromatography 800(1998)247

N-Bromosuccinimide

High-density ligand attachment to brominated allyl matrices and application to mixed mode chromatography of chymosin 775(1997)39

Bropirimine

Assay of bropirimine in rat plasma by means of robotic solid-phase extraction and high-performance liquid chromatography 762(1997)243

Bupivacaine

Capillary electrochromatography with molecular imprint-based selectivity for enantiomer separation of local anaesthetics 792(1997)401

Buspirone

Measurement and pharmacokinetic analysis of buspirone by means of brain microdialysis coupled to high-performance liquid chromatography with electrochemical detection 762(1997)269

High-performance liquid chromatographic method with coulometric detection for the determination of buspirone in human plasma by means of a column-switching technique 797(1998)221

Validated liquid chromatographic method for the determination of N-3-(2,2,5,5-tetramethyl-3-pyrrolin-3-carboxamidopropylphthalimide hydrochloride), a novel antiarrhythmic agent in human plasma 797(1998)265

Butanediol polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions 775(1997)157

Butanone

Glucose metabolism and bioreduction of 2-butanone by *Candida utilis* studied by means of ion-exchange chromatography 763(1997)205

Butyldimethylsilyl ethers

Gas-liquid chromatography-mass spectrometry of hydroxy fatty acids as their methyl esters *tert.*-butyldimethylsilyl ethers 793(1998)91

Butylmercaptan

Gas chromatographic–mass spectrometric analysis of mercaptan odorants in liquefied petroleum gas and liquefied natural gas 791(1997)367

Butyl methacrylate

Rigid porous polyacrylamide-based monolithic columns containing butyl methacrylate as a separation medium for the rapid hydrophobic interaction chromatography of proteins 775(1997)65

Butyrospermol

Isolation of three triterpenes and several aliphatic alcohols from *Crataegus monogyna* Jacq 767(1997)340

C**C2B8**

Validation of a capillary isoelectric focusing method for the recombinant monoclonal antibody C2B8 800(1998)355

Cadaverine

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

Determination of histamine and some other amines by high-performance capillary electrophoresis with on-line mode in-capillary derivatization 790(1997)177

Use of 2-chloroethylnitrosourea, a new type of pre-column derivatizing agent for the measurement of biogenic amines, by high-performance liquid chromatography with ultraviolet detection 795(1998)239

Cadmium

Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217

Rapid assay for the analysis of class III metallothioneins by capillary zone electrophoresis in crude extracts of microalgal cells exposed to cadmium 775(1997)339

Cibacron Blue F3GA-attached polyvinylbutyral microbeads as novel magnetic sorbents for removal of Cu(II), Cd(II) and Pb(II) ions 793(1998)47

On-line pretreatment and determination of Pb, Cu and Cd at the $\mu\text{g l}^{-1}$ level in drinking water by chelation ion chromatography 800(1998)247

Cadmium sulfide

Analysis of colloids. IX. Investigation of the electrical double layer of colloidal inorganic nanometer-particles by size-exclusion chromatography 773(1997)179

Caffeine

Fiber optic-based interface for on-line selective photometric determinations in solid samples by supercritical fluid extraction 753(1996)299

Characterization and impact of the temporal behavior of the electroosmotic flow in capillary isoelectric focusing with electroosmotic zone displacement 756(1996)219

Simultaneous determination of qualitatively important components in green tea infusions using capillary electrophoresis 758(1997)332

Analysis of beverages by capillary electrophoresis 781(1997)481

On-line micellar electrokinetic chromatography–mass spectrometry with a high-molecular-mass surfactant 794(1998)317

Calcium

Determination of free calcium in vegetables by capillary zone electrophoresis 759(1997)211

Determination of Na^+ , K^+ , Mg^{2+} and Ca^{2+} in mixtures of seawater and formation water by capillary electrophoresis 767(1997)311

Validation of a capillary electrophoresis procedure for the determination of calcium in calcium acamprostate 772(1997)265

Simple recognition of similar samples for the ion-chromatographic determination of the main cations 789(1997)323

Determination of calcium and magnesium in water samples by high-performance liquid chromatography on a graphitic stationary phase with a mobile phase containing *o*-cresolphthalein complexone 789(1997)329

Calcium acamprostate

Validation of a capillary electrophoresis procedure for the determination of calcium in calcium acamprostate 772(1997)265

Canrenone

High-performance liquid chromatography methods for the separation and quantitation of spironolactone and its degradation products in aqueous formulations and of its metabolites in rat serum 797(1998)271

Caproic acid

Chemically removable derivatization reagent for liquid chromatography: 2-(2-naphthoxy)ethyl 2-[1-(4-benzyl)piperazyl]ethanesulfonate 769(1997)201

Capsanthin

Isocratic non-aqueous reversed-phase high-performance liquid chromatographic separation of capsanthin and capsorubin in red peppers (*Capsicum annuum* L.), paprika and oleoresin 757(1997)89

Capsorubin

Isocratic non-aqueous reversed-phase high-performance liquid chromatographic separation of capsanthin and capsorubin in red peppers (*Capsicum annuum* L.), paprika and oleoresin 757(1997)89

Captafol

Determination of the fungicides folpet, captan and captafol by cloud-point preconcentration and high-performance liquid chromatography with electrochemical detection 754(1996)85

Captan

Determination of the fungicides folpet, captan and captafol by cloud-point preconcentration and high-performance liquid chromatography with electrochemical detection 754(1996)85

Caraboranes

High-performance liquid chromatographic enantiomeric resolution in the ten-vertex carborane series. Comparison of acetyl- and native β -cyclodextrin bonded chiral stationary phases 793(1998)249

Carbamates

Erratum to "Gas chromatographic-mass spectrometric study of photodegradation of carbamate pesticides". [J. Chromatogr. A, 738 (1996) 225–231] 761(1997)341

Development of a new high-performance liquid chromatography method to analyse N-methylcarbamate insecticides by a simple post-column derivatization system and fluorescence detection 778(1997)103

Capillary electrophoresis as a routine industrial tool for quantitative analytical testing. Determination of sodium dimethyldithiocarbamate in effluents 781(1997)541

Study of the feasibility of the use of activated carbon membranes for the on-line cleanup of vegetable extracts in the determination of N-methylcarbamate pesticides by liquid chromatography 788(1997)87

Resolution and quantitative determination of coeluted pesticide mixtures in liquid chromatography–thermospray mass spectrometry by multivariate curve resolution 795(1998)3

Comparison of different clean-up procedures for the determination of N-methylcarbamate insecticides in vegetable matrices by high-performance liquid chromatography with UV detection 795(1998)43

Carbamazepine

Purity control of carbamazepine by micellar electrokinetic chromatography 768(1997)113

Carbaryl

Simultaneous quantification of insecticides including carbaryl in drinking water by gas chromatography using dual electron-capture and nitrogen–phosphorus detection 769(1997)285

Carbazoles

Clean-up and analysis of carbazole and acridine type polycyclic aromatic nitrogen heterocyclics in complex sample matrices 790(1997)73

Carbendazim

Chromatographic analysis of imazalil and carbendazim in fruits. Method validation and residue monitoring program 1995 765(1997)91

Simplified reversed-phase conditions for the determination of benzimidazole fungicides in fruits by high-performance liquid chromatography with UV detection 778(1997)95

High-performance liquid chromatographic determination of benomyl and carbendazim residues in apiarian samples 787(1997)129

Determination of carbendazim in soil and lake water by immunoaffinity extraction and coupled-column liquid chromatography–tandem mass spectrometry 791(1997)119

Carbofuran

Resolution and quantitative determination of coeluted pesticide mixtures in liquid chromatography–thermospray mass spectrometry by multivariate curve resolution 795(1998)3

Carbohydrates

see also Disaccharides, Monosaccharides, Oligosaccharides, Polysaccharides, Saccharides, Sugars and Special Issue Index

Simultaneous determination of soluble sugars and organic acids as their trimethylsilyl derivatives in apricot fruits by gas–liquid chromatography 758(1997)99

Use of monoclonal antibodies for weak affinity chromatography 758(1997)199

Analysis of monosaccharide composition by capillary electrophoresis 763(1997)271

Simultaneous amperometric and potentiometric detection of sugars, polyols and carboxylic acids in flow systems using copper wire electrodes 766(1997)27

Continuous chromatographic separation process: simulated moving bed allowing simultaneous withdrawal of three fractions 770(1997)39

On-line electro dialysis–capillary zone electrophoresis–mass spectrometry of inositol phosphates in complex matrices 777(1997)51

Subcritical fluid chromatography of monosaccharides and polyols using silica and trimethylsilyl columns 785(1997)195

Sub-minute separations of organic and inorganic anions with co-electroosmotic capillary electrophoresis 787(1997)243

On-line electrochemical detection of carbohydrates coupled with the periodate oxidation 790(1997)1

Capillary electrophoresis of carboxylated carbohydrates. IV. Adjusting the separation selectivity of derivatized carboxylated carbohydrates by controlling the electrolyte ionic strength at subambient temperature and in the absence of electroosmotic flow 792(1997)75

Separation of aldose enantiomers by capillary electrophoresis in the presence of optically active N-dodecoxycarbonylvalines 792(1997)385

Carbolines

High-performance liquid chromatography–electrospray ionisation–tandem mass spectrometry for the analysis of 1,2,3,4-tetrahydro- β -carboline derivatives 767(1997)101

Carbon 13

Influence of gas chromatographic parameters on measurement of $^{13}\text{C}/^{12}\text{C}$ isotope ratios by gas–liquid chromatography–combustion isotope ratio mass spectrometry. I 752(1996)233

Carbonate

Carbonate interferences by ion chromatographic determination of anions in mineral waters 764(1997)249

Carbon dioxide

Multi-channel on-line chromatographic analysis system for quick air–fuel ratio measurement 752(1996)183

Liquid and subcritical CO_2 separations of enantiomers on a broadly applicable polysiloxane chiral stationary phase 753(1996)109

Chemiluminescence nitrogen detection for packed-column supercritical fluid chromatography with methanol modified carbon dioxide 757(1997)183

Evaluation of a micro gas chromatographic technique for environmental analyses of CO_2 and C_1 – C_6 alkanes 775(1997)243

Supercritical CO_2 assisted liquid extraction of nitroaromatic and polycyclic aromatic compounds in soil 785(1997)227

Semi-continuous extraction/purification of lipids by means of supercritical fluids 785(1997)337

Interfering contaminants in carbon dioxide solvent used in the supercritical fluid extraction of polychlorinated biphenyls 799(1998)259

Carbon monoxide

Gas chromatography of Titan's atmosphere. VIII. Analysis of permanent gases with carbon molecular sieve packed capillary columns 761(1997)195

Carbonyl compounds

see also Aldehydes and Ketones

UV detection of derivatized carbonyl compounds in rain samples in capillary electrophoresis using sample stacking and a Z-shaped flow cell 767(1997)241

Microemulsion electrokinetic chromatography of diphenylhydrazones of dicarbonyl sugars 772(1997)297

Comparison of extraction methods and detection systems in the gas chromatographic analysis of volatile carbonyl compounds 779(1997)360

New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259

Normal-phase and reversed-phase high-performance liquid chromatographic behaviour and particle beam mass spectrometric characterization of alkyne-substituted iron–carbonyl clusters 789(1997)461

Determination of organic molecular markers in marine aerosols and sediments: one-step flash chromatography compound class fractionation and capillary gas chromatographic analysis 799(1998)215

Carboranes

High-performance liquid chromatographic separations of boron-cluster compounds (Review) 789(1997)497

High-performance liquid chromatographic enantiomeric resolution in the ten-vertex carborane series. Comparison of acetyl- and native β -cyclodextrin bonded chiral stationary phases 793(1998)249

Carboxyethylgermanium sesquioxide

Separation and determination of inorganic germanium and β -carboxyethylgermanium sesquioxide by high-performance ion-exclusion chromatography 789(1997)403

Carboxylates

Divalent pairing ion for ion-interaction chromatography of sulphonates and carboxylates 770(1997)51

Carboxylic acids

see also Acids, Fatty acids and Organic acids

Mixed-mode liquid chromatography of carboxylic acids and inorganic anions on a latex-based pellicular stationary phase 753(1996)253

Use of modifier as trapping fluid in preparative supercritical fluid chromatography 753(1996)306

High-speed separation of carboxylic acids by co-electroosmotic capillary electrophoresis with direct and indirect UV detection 758(1997)263

Direct liquid chromatographic enantioseparation of sotalol and other β -blockers using an α_1 -acid glycoprotein-based chiral stationary phase 760(1997)193

Presence of tetrahydro- β -carboline-3-carboxylic acids in foods by gas chromatography–mass spectrometry as their *N*-methoxycarbonyl methyl ester derivatives 765(1997)265

Simultaneous amperometric and potentiometric detection of sugars, polyols and carboxylic acids in flow systems using copper wire electrodes 766(1997)27

Determination of fermenting acids in silage by capillary electrophoresis 766(1997)197

Ion chromatographic separation of carboxylic acids. Prediction of retention data 770(1997)13

Retention behavior of carboxylic acids on highly cross-linked poly(styrene–divinylbenzene)-based and silica-based cation exchangers 770(1997)23

Ion-exclusion chromatography of aliphatic carboxylic acids on a cation-exchange resin by elution with polyvinyl alcohol 770(1997)211

Analysis of aliphatic carboxylic acids and amino acids in effluents of landfills, composting plants and fermentation plants by ion-exclusion and ion-exchange chromatography 770(1997)229

Non-aqueous capillary electrophoretic separation of Brønsted acids as heteroconjugated anions 771(1997)275

Capillary zone electrophoresis of hydroxynaphthalenecarboxylic acids. Purity monitoring of β -hydroxynaphthoic acid in industry 772(1997)397

Quantitative analysis of 1-aminocyclopropane-1-carboxylic acid by liquid chromatography coupled to electrospray tandem mass spectrometry 775(1997)143

Comparison of capillary electrophoresis systems for the reliable identification of some carboxylic acids using the marker technique 781(1997)239

Ion-exclusion chromatography of benzenecarboxylic acids on an unmodified silica-gel column 782(1997)33

New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259

Sub-minute separations of organic and inorganic anions with co-electroosmotic capillary electrophoresis 787(1997)243

Determination of organic and inorganic anions in insect haemolymph and midgut contents by ion chromatography 789(1997)279

Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309

Ultra-trace determination of fluorinated aromatic carboxylic acids in aqueous reservoir fluids using solid-phase extraction in combination with gas chromatography–mass spectrometry 793(1998)297

Systematic approach to the separation of mono- and hydroxycarboxylic acids in environmental samples by ion chromatography and capillary electrophoresis 796(1998)335

Carboxylic acids, perfluorinated

Reversed-phase liquid chromatographic separation of platinum(II) complexes of methionine- and histidine-containing peptides using perfluorinated carboxylic acids as ion-pairing reagents 761(1997)115

Carboxymethylated- β -cyclodextrin

Cyclodextrin aided separation of peptides and proteins by capillary zone electrophoresis 796(1998)367

Cardenolides

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Cardiac glycosides

Separation of cardiac glycosides by micellar electrokinetic chromatography and microemulsion electrokinetic chromatography 779(1997)227

Carnitine

Separation of carnitine and acylcarnitines by capillary electrophoresis 781(1997)475

Carotenes

Synthesis and characterization of extended length alkyl stationary phases for liquid chromatography with application to the separation of carotenoid isomers 753(1996)37

Supercritical fluid extraction and reversed-phase liquid chromatography methods for vitamin A and β -carotene. Heterogeneous distribution of vitamin A in the liver 762(1997)201

Concurrent liquid chromatographic separation and photodiode array detection of retinol, tocopherols, all-*trans*- α -carotene, all-*trans*- β -carotene and the mono-*cis* isomers of β -carotene in extracts of human plasma 787(1997)111

Carotenoids

Synthesis and characterization of extended length alkyl stationary phases for liquid chromatography with application to the separation of carotenoid isomers 753(1996)37

Combined use of temperature and solvent strength in reversed-phase gradient elution. IV. Selectivity for neutral (non-ionized) samples as a function of sample type and other separation conditions 756(1996)63

Temperature dependence of carotenoids on C₁₈, C₃₀ and C₃₄ bonded stationary phases 757(1997)29

Isocratic non-aqueous reversed-phase high-performance liquid chromatographic separation of capsanthin and capsorubin in red peppers (*Capsicum annuum* L.), paprika and oleoresin 757(1997)89

Use of high-precision gas isotope ratio mass spectrometry to determine natural abundance ¹³C in lutein isolated from C₃ and C₄ plant sources 800(1998)51

Carvone

Quantification of carvone, cineole, perillaldehyde, perillyl alcohol and sobrerol by isocratic high-performance liquid chromatography 793(1998)71

Casein hydrolyzates

Determination of molecular-mass distribution of food-grade protein hydrolyzates by size-exclusion chromatography and chemiluminescent nitrogen detection 763(1997)323

Caseinoglycomacropeptide

Development of a capillary zone electrophoresis method for caseinoglycomacropeptide determination 790(1997)195

Caseins

Capillary electrophoretic analysis of genetic variants of milk proteins from different species 768(1997)47

Capillary electrophoresis of peptides in isoelectric buffers 772(1997)203

Characterization of protein glycoforms by capillary-zone electrophoresis–nanoelectrospray mass spectrometry 794(1998)327

Catechins

Study of the acetaldehyde induced polymerisation of flavan-3-ols by liquid chromatography–ion spray mass spectrometry 752(1996)85

Simultaneous determination of qualitatively important components in green tea infusions using capillary electrophoresis 758(1997)332

Selection of column and gradient elution system for the separation of catechins in green tea using high-performance liquid chromatography 793(1998)265

Analysis of catechins in tea extracts by liquid chromatography–electrospray ionization mass spectrometry 794(1998)63

Separation of tea polyphenols using micellar electrokinetic chromatography with diode array detection 799(1998)309

Catechol

Off-column chemiluminescence detection in capillary electrophoresis 771(1997)379

Catecholamines

see also Amines and Biogenic amines

pH-Zone-refining countercurrent chromatography 753(1996)1

Enhanced temporal resolution for the microdialysis monitoring of catecholamines and excitatory amino acids using capillary electrophoresis with laser-induced fluorescence detection. Analytical developments and in vitro validations 755(1996)99

Off-column chemiluminescence detection in capillary electrophoresis 771(1997)379

Capillary electrophoresis and electrochemical detection with a conventional detector cell 798(1998)289

Cations

see also Inorganic cations and Metal cations

Simultaneous determination of common mono- and divalent cations by ion chromatography with an unmodified silica gel column 752(1996)167

Determination of Na⁺, K⁺, Mg²⁺ and Ca²⁺ in mixtures of seawater and formation water by capillary electrophoresis 767(1997)311

Practical aspects of suppressed ion chromatography for cations in human serum 770(1997)137

Comparisons of the mobilities of salt ions obtained by the moving boundary method and two empirical equations in capillary electrophoresis 771(1997)374

Water clusters studied by electrospray mass spectrometry 777(1997)155

Simple recognition of similar samples for the ion-chromatographic determination of the main cations 789(1997)323

Flow injection–capillary electrophoresis coupling to automate on-line sample treatment for the determination of inorganic ions in waters 791(1997)279

Ceftiofur

Quantitative determination of ceftiofur in milk by liquid chromatography–electrospray mass spectrometry 794(1998)57

Cellulose

Bifunctional etherification of a bead cellulose for ligand attachment with allyl bromide and allyl glycidyl ether 775(1997)29

Capillary electrochromatography: normal-phase mode using silica gel and cellulose-based packing materials 782(1997)167

3,5-Dimethylphenylcarbamates of amylose, chitosan and cellulose bonded on silica gel. Comparison of their chiral recognition abilities as high-performance liquid chromatography chiral stationary phases 796(1998)265

Cellulose, fibrous

Evaluation of a fibrous cellulose drying agent in supercritical fluid extraction and pressurized liquid extraction of diverse pesticides 785(1997)313

Cellulose, modified

Effects of flow-rates and sample concentration on the molar mass characterisation of modified celluloses using asymmetrical flow field-flow fractionation–multi-angle light scattering 791(1997)135

Cellulose tris(4-methylbenzoate)

Non-linear structure–enantioselective retention relationships in a homologous series of 1,4-disubstituted piperazine derivatives 788(1997)81

Cetylpyridinium chloride

Determination of antibacterial quaternary ammonium compounds in lozenges by capillary electrophoresis 798(1998)335

Chaperonins

Purification of chaperonins from thermophilic bacteria and archaea 773(1997)131

Charged solutes

Ideal model of chromatography applied to charged solutes in reversed-phase liquid chromatography 761(1997)3

Chitosan

3,5-Dimethylphenylcarbamates of amylose, chitosan and cellulose bonded on silica gel. Comparison of their chiral recognition abilities as high-performance liquid chromatography chiral stationary phases 796(1998)265

Chloralose

Identification of pesticide poisoning in wildlife 754(1996)463

Chlorbromuron

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Chlordane

High-resolution gas chromatographic test for the characterisation of enantioselective separation of organochlorine compounds. Application to *tert*-butyldimethylsilyl β -cyclodextrin 775(1997)275

Chloride

Ion chromatography methods for the simultaneous determination of mineral anions in plant sap 752(1996)298

Indirect photometric detection of inorganic anions in microcolumn ion chromatography using octadecylsilica immobilized with bovine serum albumin as stationary phase 755(1996)37

Theoretical study of the elution profile of a high-concentration sample anion in non-suppressed ion chromatography 789(1997)43

Ion chromatographic separation of metallic ions on a bonded poly- β -diketone-type non-cyclic crown ether stationary phase 789(1997)135

Analysis of difficult samples by capillary zone electrophoresis. I. Simultaneous separation of detrimental anions in concrete digested with concentrated nitric acid 793(1998)187

Chlorine

Determination of chlorine and bromine in rocks by alkaline fusion with ion chromatography detection 770(1997)93

Chlorine-containing anions

Separation of chlorine-containing anions by ion chromatography and capillary electrophoresis 777(1997)375

Chlorine dioxide

Simultaneous determination of chlorine dioxide and hypochlorite in water by high-performance liquid chromatography 796(1998)397

Chloro(dichloromethyl)hydroxyfuranone

New derivatization method for determination of 3-chloro-4(dichloromethyl)-5-hydroxy-2(5H)-furanone in water 790(1997)242

Chloroalkanes

New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259

Chlorobenzenes

Analysis of chlorobenzenes in soils by headspace solid-phase microextraction and gas chromatography–ion trap mass spectrometry 771(1997)181

Chlorobiphenyls

Development of a thermal desorption modulator for gas chromatography 767(1997)137

Chloro complexes

Determination of platinum, palladium, osmium, iridium, rhodium and gold as chloro complexes by capillary zone electrophoresis 772(1997)347

Speciation of platinum–chloro complexes and their hydrolysis products by ion chromatography. Determination of platinum oxidation states 775(1997)197

2-Chloroethylnitrosourea

Use of 2-chloroethylnitrosourea, a new type of pre-column derivatizing agent for the measurement of biogenic amines, by high-performance liquid chromatography with ultraviolet detection 795(1998)239

Chlorogenic acid

New methods for spectrometric peak purity analysis in chromatography 800(1998)121

6-Chloronicotinic acid

Resolution of imidacloprid pesticide and its metabolite 6-chloronicotinic acid using cross-sections of spectrochromatograms obtained by high-performance liquid chromatography with diode-array detection 799(1998)149

Chlorophenols

Determination of chlorophenols by micellar electrokinetic chromatography with electrochemical detection 757(1997)203

Capillary electrophoretic separation of chlorophenols using amperometric detection 761(1997)259

Supercritical fluid and enhanced-fluidity liquid extraction of phenolics from river sediment 764(1997)265

Determination of phenols by solid-phase microextraction 767(1997)171

Effect of cationic polymer on the separation of phenols by capillary electrophoresis 768(1997)161

Analytical methods for the determination of organochlorine compounds. Application to environmental samples in the Slovak Republic 774(1997)333

Determination of phenolic pollutants in drinking water by capillary electrophoresis in the sample stacking mode 778(1997)279

Determination of priority phenolic compounds in water and industrial effluents by polymeric liquid–solid extraction cartridges using automated sample preparation with extraction columns and liquid chromatography. Use of liquid–solid extraction cartridges for stabilization of phenols 778(1997)301

Determination of polar priority phenols at parts per trillion levels in water using on-line liquid–solid extraction followed by liquid chromatography with coulometric detection 778(1997)313

A simple means of generating pH gradients in capillary zone electrophoresis 781(1997)3

Evaluation of two solid-phase extraction procedures for the preconcentration of chlorophenols in drinking water 786(1997)285

Liquid chromatography–atmospheric pressure ionization mass spectrometry for the determination of chloro- and nitrophenolic compounds in tap water and sea water 787(1997)79

Capillary electrophoresis and electrochemical detection with a conventional detector cell 798(1998)289

Chlorophylls

Combined use of temperature and solvent strength in reversed-phase gradient elution. III. Selectivity for ionizable samples as a function of sample type and pH 756(1996)51

Chloropropanediol

Determination of 1,3-dichloro-2-propanol and 3-chloro-1,2-propanediol in papers treated with polyamidoamine–epichlorohydrin wet-strength resins by gas chromatography–mass spectrometry using selective ion monitoring 788(1997)195

Chlorotrifluoroethylene

Interfering contaminants in carbon dioxide solvent used in the supercritical fluid extraction of polychlorinated biphenyls 799(1998)259

Chlorpheniramine

Development and validation of a high-performance liquid chromatography method for the determination of cold relief ingredients in chewing gum 775(1997)179

Separation of basic drug enantiomers by capillary zone electrophoresis using ovoglycoprotein as a chiral selector 782(1997)281

Chlorpyrifos

Automated sample clean-up and fractionation of chlorpyrifos, chlorpyrifos-methyl and metabolites in mussels using normal-phase liquid chromatography 778(1997)151

Chlortoluron

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Cholesterol oxides

Procedure for the determination of eight cholesterol oxides in poultry meat using on-column and solvent venting capillary gas chromatography 764(1997)279

Evaluation of particle beam high-performance liquid chromatography–mass spectrometry for analysis of cholesterol oxides 794(1998)253

Cholic acid

Bile salt micellar electrokinetic chromatography of bilirubin and related compounds 779(1997)205

Choline

Analysis of tetramine in sea snails by capillary electrophoresis–tandem mass spectrometry 781(1997)555

Choline glycolate

Analysis of tetramine in sea snails by capillary electrophoresis–tandem mass spectrometry 781(1997)555

Chorsulfuron

Determination of chorsulfuron and tribenuron-methyl residues in agricultural soils 778(1997)119

Chromium

Speciation of chromium dyes by high-performance liquid chromatography with inductively coupled plasma mass spectrometric detection 773(1997)157

Speciation of chromium from industrial wastes and incinerated sludges 774(1997)363

Simultaneous determination of toxic arsenic and chromium species in water samples by ion chromatography–inductively coupled plasma mass spectrometry 779(1997)139

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

Liquid chromatographic separation of aqueous species of Cr(VI) and Cr(III) 789(1997)469

Electrophoretic studies on mixed complexes. Metal–methionine–penicillamine system 793(1998)220

Chromium complexes

Chromatographic behaviour of the chromium(III) complex of 1-phenyl-3-methyl-4-benzoyl-5-pyrazolone 789(1997)479

Chymosin

High-density ligand attachment to brominated allyl matrices and application to mixed mode chromatography of chymosin 775(1997)39

Cibacron Blue

Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217

Dye–ligand and metal chelate poly(2-hydroxyethylmethacrylate) membranes for affinity separation of proteins 799(1998)83

Cineole

Quantification of carvone, cineole, perillaldehyde, perillyl alcohol and sabinol by isocratic high-performance liquid chromatography 793(1998)71

Cinnamaldehyde

Determination of cinnamaldehyde, cinnamic acid, paeoniflorin, glycyrrhizin and [6]-gingerol in the traditional Chinese medicinal preparation Kuei-chih-tang by cyclodextrin-modified micellar electrokinetic chromatography 771(1997)267

Cinnamic acid

Comparative investigation of UV, electrochemical and particle beam mass spectrometric detection for the high-performance liquid chromatographic determination of benzoic and cinnamic acids and of their corresponding phenolic acids 753(1996)157

Determination of cinnamaldehyde, cinnamic acid, paeoniflorin, glycyrrhizin and [6]-gingerol in the traditional Chinese medicinal preparation Kuei-chih-tang by cyclodextrin-modified micellar electrokinetic chromatography 771(1997)267

Cinoxacin

Determination of the chemotherapeutic quinolonic and cinolonic derivatives in urine by high-performance liquid chromatography with ultraviolet and fluorescence detection in series 787(1997)119

Citrate

Capillary zone electrophoretic separation of β -blockers using citrate buffer at low pH 753(1996)133

Migration behavior and separation of sulfonamides in capillary zone electrophoresis. III. Citrate buffer as a background electrolyte 768(1997)105

Citric acid

High-speed separation of carboxylic acids by co-electroosmotic capillary electrophoresis with direct and indirect UV detection 758(1997)263

New approach to determining each component of a two-component overlapping peak by single-column anion ion chromatography 769(1997)333

Separation and determination of four artificial sweeteners and citric acid by high-performance anion-exchange chromatography 771(1997)135

Gas chromatographic determination of organic acids from fruit juices by combined resin mediated methylation and extraction in supercritical carbon dioxide 785(1997)251

Clenbuterol

Detection of clenbuterol in bovine retinal tissue by high-performance liquid chromatography with electrochemical detection 762(1997)275

Effect of on-capillary large volume sample stacking on limits of detection in the capillary zone electrophoretic determination of selected drugs, dyes and metal chelates 772(1997)161

Clodronic acid

Determination of impurities in clodronic acid by anion-exchange chromatography 770(1997)261

Clopyralid

Analysis of the herbicide clopyralid in cultivated soils
754(1996)265

Clotting factor VIII

Application of compact porous tubes for preparative isolation of clotting factor VIII from human plasma 760(1997)117

Clotting factor IX

Size-exclusion chromatography of plasma proteins with high molecular masses 796(1998)289

Clotting factor XIII

Size-exclusion chromatography of plasma proteins with high molecular masses 796(1998)289

Cobalt complexes

Chromatographic separation and characterization of the photoproduct of tris(L-cysteinesulfinate-N,S)cobaltate(III)
789(1997)273

Cocaine

Lesser alkaloids of cocaine-bearing plants. III. 2-Carbomethoxy-3-oxo substituted tropane esters: detection and gas chromatographic–mass spectrometric characterization of new minor alkaloids found in South American *Erythroxylum coca* var. *coca* 756(1996)185

Comparative determination of total isomeric truxillines in illicit, refined, South American cocaine hydrochloride using capillary gas chromatography–electron capture detection 756(1996)193

Application of micellar electrokinetic capillary chromatography to the analysis of illicit drug seizures 780(1997)265

Codeine

Determination of morphine and its analogues in urine by on-line coupled reversed-phase liquid chromatography–gas chromatography with on-line derivatization 771(1997)360

Effect of on-capillary large volume sample stacking on limits of detection in the capillary zone electrophoretic determination of selected drugs, dyes and metal chelates 772(1997)161

Codeine phosphate

Separation of ibuprofen, codeine phosphate, their degradation products and impurities by capillary electrophoresis. I. Method development and optimization with fractional factorial design 798(1998)307

Coenzyme A esters

Isocratic high-performance liquid chromatography of coenzyme A esters involved in the metabolism of 3S-hydroxy-3-methylglutaryl-coenzyme A. Detection of related enzyme activities in *Catharanthus roseus* plant cell cultures 752(1996)123

Collagenase

Micropreparation of tissue collagenase fragments of type I collagen in the form of surfactant–peptide complexes and their identification by capillary electrophoresis and partial sequencing 796(1998)181

Collagens

Semipreparative isolation of collagen types I, II, III and V by sodium dodecyl sulfate–polyacrylamide gel electrophoresis and electroelution 758(1997)313

Using quaternary high-performance liquid chromatography eluent systems for separating 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate-derivatized amino acid mixtures 763(1997)11

Capillary electrophoresis of large cyanogen bromide peptides of fibre-forming collagens with special reference to cross-linking 772(1997)213

Micropreparation of tissue collagenase fragments of type I collagen in the form of surfactant–peptide complexes and their identification by capillary electrophoresis and partial sequencing 796(1998)181

2,4,6-Collidine

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Colorflamine

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Conalbumin

Development of a high-performance capillary isoelectric focusing technique with application to studies of microheterogeneity in chicken conalbumin 757(1997)247

Conjugated linoleic acid

Fractionation of *cis*- and *trans*-oleic, linoleic, and conjugated linoleic fatty acid methyl esters by silver ion high-performance liquid chromatography 799(1998)329

Copper

Potentiometric detection of aliphatic amines by flow injection analysis and ion-interaction chromatography with a metallic copper electrode 758(1997)227

Cibacron Blue F3GA-attached polyvinylbutyral microbeads as novel magnetic sorbents for removal of Cu(II), Cd(II) and Pb(II) ions 793(1998)47

Reversed-phase high-performance liquid chromatography of iron(II) and copper(II) chelates with 4,7-diphenyl-1,10-phenanthroline disulfonate 799(1998)171

Separation and direct UV detection of sugars by capillary electrophoresis using chelation of copper(II) 799(1998)283

On-line pretreatment and determination of Pb, Cu and Cd at the $\mu\text{g l}^{-1}$ level in drinking water by chelation ion chromatography 800(1998)247

Copper complexes

Reversed-phase ion-interaction chromatography of Cu(I)-cyanide complexes 770(1997)165

Detection of glycosaminoglycans as a copper(II) complex in high-performance liquid chromatography 787(1997)266

Copper(I)-cyanide

Ion chromatographic analysis of cyanate in gold processing samples containing large concentrations of copper(I) and other metallo-cyanide complexes 770(1997)175

Copper phthalocyanine trisulfonate

On-line preconcentration and high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbon-DNA adducts using copper phthalocyanine trisulfonic acid as a group-specific adsorbent 755(1996)205

5-Corononylpentylsilyl silica

Selectivity of electron-donor- and electron-acceptor-bonded silica packing materials for hydrophobic environmental contaminants in polar and non-polar eluents 786(1997)237

Corticosteroids

Gradient capillary electrochromatography of drug mixtures with UV and electrospray ionisation mass spectrometric detection 768(1997)89

Retention behavior of synthetic corticosteroids in packed-column supercritical fluid chromatography 773(1997)277

Immunoaffinity chromatography combined on-line with high-performance liquid chromatography-mass spectrometry for the determination of corticosteroids 794(1998)37

Cortisol

Liquid chromatography-mass spectrometry with on-line solid-phase extraction by a restricted-access C_{18} precolumn for direct plasma and urine injection 762(1997)193

Coumarin

Application of high-speed counter-current chromatography to the separation of coumarin and related compounds 759(1997)47

CP-93,393

Assay and purity evaluation of CP-93,393-1 by reversed-phase liquid chromatography. A snapshot of current practices for liquid chromatography methods development and validation 764(1997)233

Creatinine

Quantification of urinary uric acid in the presence of thymol and thimerosal by high-performance liquid chromatography 763(1997)187

Cresolphthalein complexone

Determination of calcium and magnesium in water samples by high-performance liquid chromatography on a graphitic stationary phase with a mobile phase containing *o*-cresolphthalein complexone 789(1997)329

Cresols

Effects of the gradient profile on the production rate in reversed-phase gradient elution overloaded chromatography 760(1997)25

Fitting competitive adsorption isotherms to the experimental distribution data in reversed-phase systems 762(1997)3

Evaluation of supercritical fluid extraction, microwave-assisted extraction and sonication in the determination of some phenolic compounds from various soil matrices 774(1997)243

Optimization of the recovery yield and of the production rate in overloaded gradient-elution reversed-phase chromatography 796(1998)115

Crown ethers

Chiral crown ether-anchored polysiloxanes as capillary gas chromatography stationary phases 753(1996)269

Elution delay of fluoride by chromatography of anions on silica gels coated with crown ether dyes 755(1996)43

Separation of enantiomers and isomers of amino compounds by capillary electrophoresis and high-performance liquid chromatography utilizing crown ethers 757(1997)225

Nonaqueous anion-exchange chromatography. II. Changing anion-exchange selectivity by resin surface complex formation of crown ethers 758(1997)29

Achievement of enantioselectivity of non-polar primary amines by a non-chiral crown ether 781(1997)129

Mechanism of capillary electrophoresis enantioseparations using a combination of an achiral crown ether plus cyclodextrins 793(1998)115

Cyanate

Ion chromatographic analysis of cyanate in gold processing samples containing large concentrations of copper(I) and other metallo-cyanide complexes 770(1997)175

Cyanazine

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Cyanide complexes

Reversed-phase ion-interaction chromatography of Cu(I)-cyanide complexes 770(1997)165

Capillary electrophoretic determination of cyanide leaching solutions from automobile catalytic converters 778(1997)397

Cyanobacterial toxins

Determination of some physicochemical parameters of microcystins (cyanobacterial toxins) and trace level analysis in environmental samples using liquid chromatography 799(1998)155

Cyanocarbons

Cyanocarbons as ligands for electron donor acceptor chromatography of human serum proteins 753(1996)57

Cyanocobalamin

Determination of cyanocobalamin in foods by high-performance liquid chromatography with visible detection after solid-phase extraction and membrane filtration for the precolumn separation of lipophilic species 771(1997)127

Cyanolipids

Analysis of cyanolipids and triacylglycerols from sapindaceae seed oils with high-temperature gas chromatography and high-temperature gas chromatography-chemical ionization mass spectrometry 787(1997)181

Cyclamate

Separation and determination of four artificial sweeteners and citric acid by high-performance anion-exchange chromatography 771(1997)135

Cycloalkanes

Unified retention concept – statistical treatment of Kováts retention index 764(1997)257

Cycloartenol

Isolation of three triterpenes and several aliphatic alcohols from *Crataegus monogyna* Jacq 767(1997)340

Cyclodextrins

Separation of chiral polychlorinated biphenyls by micellar electrokinetic chromatography using β - and γ -cyclodextrin mixtures in the separation buffer 752(1996)265

Comparison of γ -cyclodextrin sulfobutyl ether and unmodified γ -cyclodextrin as chiral selectors in capillary electrophoresis 755(1996)81

Effect of β -cyclodextrin acetylation on the resolution of phenethylamines with capillary electrophoresis and nuclear magnetic resonance spectroscopy 758(1997)277

Chiral separation of monoterpenes using mixtures of sulfated β -cyclodextrins and α -cyclodextrin as chiral additives in the reversed-polarity capillary electrophoresis mode 759(1997)149

Enantioseparation of 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate-derivatized-amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives. I. Discussion of optimum separation conditions 759(1997)157

Enantioseparation of 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate-derivatized amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives. II. Evaluation of complexation constants 759(1997)167

Chiral separation of neutral species by capillary electrophoresis. Evaluation of a theoretical model 761(1997)249

Chiral resolution of diols by capillary electrophoresis using borate-cyclodextrin complexation 761(1997)269

Ruggedness of enantiomeric separation by capillary electrophoresis and high-performance liquid chromatography with methylated cyclodextrins as chiral selectors 763(1997)139

Chiral recognition and racemic resolution of drug enantiomers by electrophoresis based on host-guest complexation 763(1997)261

Migration order reversal of enantiomers in capillary electrophoretic separation 764(1997)151

Systematic approach to links between separations in capillary electrophoresis and liquid chromatography. IV. Application of binding constant-retention factor relationship to the separation of 2-, 3- and 4-methylbenzoate anions using β -cyclodextrin as selector 768(1997)29

Enantioseparation in capillary electrophoresis using 2-hydroxypropyltrimethylammonium salt of β -cyclodextrin as a chiral selector 771(1997)259

Micellar electrokinetic capillary chromatographic separation of polychlorinated biphenyl congeners 778(1997)77

Achievement of enantioselectivity of non-polar primary amines by a non-chiral crown ether 781(1997)129

Enantiomer separation of acidic racemates by capillary electrophoresis using cationic and amphoteric β -cyclodextrins as chiral selectors 781(1997)151

Chiral discrimination by high-performance liquid chromatography with joint use of two cyclodextrin additives 782(1997)1

Systematic approach to cost- and time-effective method development with a starter kit for chiral separations by capillary electrophoresis 782(1997)257

Thermodynamic study of the retention behaviour of selected macrocycles using reversed-phase high-performance thin-layer chromatography plates and methanol–water mobile phases 787(1997)227

Characterization of modified cyclodextrins applied in capillary electrophoresis and high-performance liquid chromatography as chiral selectors by matrix-assisted laser desorption ionization curved field reflectron mass spectrometry 791(1997)299

Electrokinetic chromatography without electroosmotic flow 792(1997)125

Controlling enantioselectivity in chiral capillary electrophoresis with inclusion–complexation 792(1997)227

Separation selectivity in chiral capillary electrophoresis with charged selectors 792(1997)269

Separation selectivity in chiral and achiral capillary electrophoresis with mixed cyclodextrins 792(1997)297

Thermodynamics of chiral selectivity in capillary electrophoresis: separation of ibuprofen enantiomers with β -cyclodextrin 792(1997)371

Influence of the degree of substitution of cyclodextrin sulfobutyl ether derivatives on enantioselective separations by electrokinetic chromatography 792(1997)379

Enhancement of the separation selectivity of a group of polycyclic aromatic hydrocarbons using mixed cyclodextrin-modified micellar electrokinetic chromatography 792(1997)411

Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers 792(1997)419

Achiral selectivity in cyclodextrin-modified capillary electrophoresis 792(1997)431

Determination of impurities in α -cyclodextrin by gradient high-performance liquid chromatography with pulsed amperometric detection 793(1998)57

Separation of enantiomers of drugs by capillary electrophoresis. V. Hydroxypropyl- α -cyclodextrin as chiral solvating agent 793(1998)153

Cationization in electrospray microcolumn liquid chromatography–mass spectrometry 794(1998)359

Chiral resolution of functionalized piperidine enantiomers by capillary electrophoresis with native, alkylated and anionic β -cyclodextrins 796(1998)375

Screening of cyclodextrins by nuclear magnetic resonance for the design of chiral capillary electrophoresis separations 797(1998)149

Effect of charged and uncharged chiral additives on the resolution of amlodipine enantiomers in liquid chromatography and capillary electrophoresis 797(1998)187

Use of cationic cyclodextrin for enantioseparation by capillary electrophoresis 798(1998)275

Use of chiral and achiral ion-pairing reagents in combination with cyclodextrins in capillary electrophoresis 798(1998)281

Comparative capillary electrophoresis and NMR studies of enantioseparation of dimethindene with cyclodextrins 798(1998)315

Chiral capillary electrophoresis–electrospray mass spectrometry coupling with charged cyclodextrin derivatives as chiral selectors 800(1998)77

$^2\text{H}_2\text{O}$ effects on the inclusional complexation of β -cyclodextrin with sodium 2-naphthalenesulfonate in capillary electrophoresis and UV spectrophotometry 800(1998)327

Capillary electrophoretic analysis of cyclodextrins with dynamic fluorescence labeling and detection 800(1998)333

Cyclodrine

Use of chiral and achiral ion-pairing reagents in combination with cyclodextrins in capillary electrophoresis 798(1998)281

Cyclohexane

Quantitative injections or calibrations in gas chromatography. Anomaly associated with syringe type 773(1997)372

Cyclohexane-1,2-diaminetetraacetic acid

Advanced possibilities on multi-element separation and detection of metal ions by capillary zone electrophoresis using precapillary complexation. I. Separation aspects 756(1996)300

Cyclohexanol

Quantitative injections or calibrations in gas chromatography. Anomaly associated with syringe type 773(1997)372

Cyclosporins

High-performance liquid chromatographic analysis of cyclosporin A and its oral solutions 759(1997)217

Cymene

Supercritical fluid chromatography–mass spectrometry of thyme extracts (*Thymus vulgaris* L.) 773(1997)377

Cypermethrin

Separation and analysis of the diastereomers and enantiomers of cypermethrin and related compounds 777(1997)363

Cysteine

Theory concerning the current for an end-column amperometric detector with a disk working electrode in capillary zone electrophoresis 765(1997)307

Determination of cysteine by capillary zone electrophoresis with end-column amperometric detection at a gold/mercury amalgam microelectrode without deoxygenation 769(1997)307

Determination of glutathione, cysteine and N-acetylcysteine in rabbit eye tissues using high-performance liquid chromatography and post-column derivatization with 5,5'-dithiobis(2-nitrobenzoic acid) 778(1997)347

Urinary excretion measurement of cysteine and homocysteine in the form of their S-pyridinium derivatives by high-performance liquid chromatography with ultraviolet detection 798(1998)27

Removal of the fluorescent 4-(aminosulfonyl)-2,1,3-benzoxadiazole label from cysteine-containing peptides 798(1998)47

Cytochalasin B

D-Glucose, forskolin and cytochalasin B affinities for the glucose transporter Glut1. Study of pH and reconstitution effects by biomembrane affinity chromatography 776(1997)81

Glucose affinity for the glucose transporter Glut1 in native or reconstituted lipid bilayers. Temperature-dependence study by biomembrane affinity chromatography 776(1997)87

Cytochromes

Productivity and operating regimes in protein chromatography using low-molecular-mass displacers 771(1997)9

Protein analysis by packed capillary liquid chromatography with electrospray ionization and time-of-flight mass spectrometry detection 786(1997)67

Cytokinin ribosides

Capillary electrophoresis of cytokinins and cytokinin ribosides 764(1997)331

Cytokinins

Capillary electrophoresis of cytokinins and cytokinin ribosides 764(1997)331

D

2,4-D

Determination of chlorophenoxy herbicides in waters by capillary gas chromatography with ion trap detection 758(1997)324

Daidzein

Development and validation of an isocratic high-performance liquid chromatographic method for quantitative determination of phytoestrogens in soya bean 795(1998)377

Dalargin

Use of capillary electrophoresis and high-performance liquid chromatography for monitoring of glycosylation of the peptides dalargin and desmopressin 761(1997)285

DDD

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

DDE

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

DDT

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

DDT congeners

Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers 792(1997)419

Debrisoquine

Separation and quantitation of debrisoquine and 4-hydroxydebrisoquine in human urine by capillary electrophoresis and high-performance liquid chromatography 778(1997)389

Decanediol polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions 775(1997)157

N⁺-*n*-Decyl-L-histidine

Study of retention, efficiency and selectivity in chiral ligand-exchange chromatography with a dynamically coated stationary phase 761(1997)79

Dehydroabietic acid

High-performance liquid chromatographic determination of dehydroabietic and abietic acids in traditional Chinese medications 763(1997)221

Dehydroascorbate

Analysis of ascorbate and dehydroascorbate in plant extracts by high-resolution selected ion monitoring gas chromatography-mass spectrometry 782(1997)95

Dendrimers

Selectivity of polymeric and polymer-supported pseudo-stationary phases in micellar electrokinetic chromatography 792(1997)105

Deoxynivalenol

Determination of trichothecene mycotoxins in wheat by use of supercritical fluid extraction and high-performance liquid chromatography with diode array detection or gas chromatography with electron capture detection 795(1998)297

Deoxy pyridinoline

Automated on-line solid-phase extraction and high-performance liquid chromatographic analysis of total and free pyridinium crosslinks in serum 798(1998)159

Deoxyribonuclease

Purification of a Streptococcal deoxyribonuclease by affinity chromatography based on a DNA–cellulose matrix 788(1997)75

Deoxyribonucleosides

Application of mixed mobile phases and a step gradient method in capillary electrochromatography for the separation of isomeric polycyclic aromatic hydrocarbon–deoxyribonucleoside adduct mixtures prepared in vitro 781(1997)327

Comparison of retention models for the dependence of retention factors on mobile phase composition in reversed-phase high-performance liquid chromatography 797(1998)23

Deoxyribonucleotides

Prediction of the migration behavior of analytes in capillary electrophoresis based on three fundamental parameters 781(1997)23

Deprenyl metabolites

Urinary excretion of deprenyl metabolites 762(1997)321

Dequalinium chloride

Determination of antibacterial quaternary ammonium compounds in lozenges by capillary electrophoresis 798(1998)335

Dequest

Analysis of amine-containing phosphonates in detergent powders by anion-exchange chromatography with pulsed amperometric detection 771(1997)155

Desethyloxybutynin

Chiral bioanalysis by normal phase high-performance liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 759(1997)65

Desflurane

Approach to the thermodynamics of enantiomer separation by gas chromatography. Enantioselectivity between the chiral inhalation anesthetics enflurane, isoflurane and desflurane and a diluted γ -cyclodextrin derivative 757(1997)119

Preparative enantiomer separation of the inhalation anesthetics enflurane, isoflurane and desflurane by gas chromatography on a derivatized γ -cyclodextrin stationary phase 769(1997)119

Deslanoside

Separation of cardiac glycosides by micellar electrokinetic chromatography and microemulsion electrokinetic chromatography 779(1997)227

Desmopressin

Use of capillary electrophoresis and high-performance liquid chromatography for monitoring of glycosylation of the peptides dalargin and desmopressin 761(1997)285

Detergents

see also Surfactants

Analysis of amine-containing phosphonates in detergent powders by anion-exchange chromatography with pulsed amperometric detection 771(1997)155

Porous graphitized carbon and octadecyl-silica columns in the separation of some alkylglycoside detergents 782(1997)191

Composition analysis of detergents of the polyoxyethylene type: comparison of thin-layer chromatography, reversed-phase chromatography and matrix-assisted laser desorption/ionization mass spectrometry 791(1997)245

Dexamethasone

Immunoaffinity chromatography combined on-line with high-performance liquid chromatography–mass spectrometry for the determination of corticosteroids 794(1998)37

Dexmedetomidine

Analytical method development for the simultaneous quantitation of dexmedetomidine and three potential metabolites in plasma 762(1997)281

Dextrans

Fast molecular mass and size characterization of polysaccharides using asymmetrical flow field-flow fractionation–multiangle light scattering 760(1997)205

Interaction between tryptophan residues and hydrophobically modified dextran. Effect on partitioning of peptides and proteins in aqueous two-phase systems 766(1997)99

Characterization of dextrans by size-exclusion chromatography on unmodified silica gel columns, with light-scattering detection, and capillary electrophoresis with laser-induced fluorescence detection 790(1997)93

Diacetylmorphine

¹³C Isotopic analysis of an acetaminophen and diacetylmorphine mixture 756(1996)203

Diacylglycerols

Non-aqueous reversed-phase high-performance liquid chromatography of synthetic triacylglycerols and diacylglycerols 782(1997)41

Diadzen

Linearity and the limitations of least squares calibration
762(1997)73

Diamines

New sorbents for removal of heavy metal ions: diamine-glow-discharge treated polyhydroxyethylmethacrylate microspheres
773(1997)169

Diaminobenzenesulfonic acid

Monitoring the products of condensation of 4-nitrobenzoyl chloride and 2,4-diaminobenzenesulphonic acid by ion-pair high-performance liquid chromatography 777(1997)370

Diaminodibenzanilide-4-sulfonic acid

Monitoring the products of condensation of 4-nitrobenzoyl chloride and 2,4-diaminobenzenesulphonic acid by ion-pair high-performance liquid chromatography 777(1997)370

1,2-Diaminopropane-N,N,N',N'-tetraacetic acid

Laser induced resonance energy transfer – a novel approach towards achieving high sensitivity in capillary electrophoresis. Part B: Applications for environmental testing 767(1997)217

Dibutyl butyl phosphonate

Novel calibration technique for headspace analysis of semivolatiles compounds 793(1998)397

Dibutyltin

Speciation of organo-tin compounds using liquid chromatography–atmospheric pressure ionisation mass spectrometry and liquid chromatography–inductively coupled plasma mass spectrometry as complementary techniques
794(1998)211

Dicarboxyimides

Application of porous carbon for solid-phase extraction of dicarboxyimide fungicide residues from wines in combination with high-resolution capillary gas chromatography and gas chromatography–mass spectrometry 754(1996)445

Dichlorobenzenes

Gas-phase electron diffraction as a detection method in capillary gas chromatography 793(1998)209

Dichloropropanol

Determination of 1,3-dichloro-2-propanol and 3-chloro-1,2-propanediol in papers treated with polyamidoamine–epichlorohydrin wet-strength resins by gas chromatography–mass spectrometry using selective ion monitoring
788(1997)195

Dichlorosilane

Measurement of trace phosphorus in dichlorosilane by high-temperature hydrogen reduction gas chromatography
757(1997)319

Dichlorprop

Simultaneous determination and enantiomeric resolution of mecoprop and dichlorprop in soil samples by high-performance liquid chromatography and gas chromatography–mass spectrometry 799(1998)355

Diclofop

Enantiomeric and isomeric separation of herbicides using cyclodextrin-modified capillary zone electrophoresis
790(1997)225

Dieldrin

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants
769(1997)275

Diesel fuel

Recovery of diesel fuel from soil by supercritical fluid extraction–gas chromatography 790(1997)125

7-Diethylaminocoumarine-3-carboxylic acid succinimidyl ester

Highly sensitive detection using laser two-photon excited fluorescence in capillary electrophoresis 765(1997)315

Diethylenetriamine pentaacetic acids

Purification of *p*-nitrobenzyl C-functionalized diethylenetriamine pentaacetic acids for clinical applications using anion-exchange chromatography 771(1997)63

Di(2-ethylhexyl)phosphate

Separation of samarium, gadolinium, terbium, dysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

Diethyl phthalate

Temperature gradients in preparative high-performance liquid chromatography columns 769(1997)109

Diflubenzuron

Determination of diflubenzuron in apples by gas chromatography 765(1997)85

Diflufenican

Comparison between accelerated solvent extraction and traditional extraction methods for the analysis of the herbicide diflufenican in soil 765(1997)121

2-Difluoromethyl-DL-ornithine

Analysis of 2-difluoromethyl-DL-ornithine in human plasma, cerebrospinal fluid and urine by cation-exchange high-performance liquid chromatography 762(1997)293

Digalactosyldiacylglycerols

Semi-continuous extraction/purification of lipids by means of supercritical fluids 785(1997)337

Digoxin

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Separation of cardiac glycosides by micellar electrokinetic chromatography and microemulsion electrokinetic chromatography 779(1997)227

Dihydrocarbamazepine

Purity control of carbamazepine by micellar electrokinetic chromatography 768(1997)113

Dihydrocodeine

Determination of morphine and its analogues in urine by on-line coupled reversed-phase liquid chromatography–gas chromatography with on-line derivatization 771(1997)360

Dihydroimidazoles

Chromatographic resolution of synthetically useful chiral glycine derivatives by high-performance liquid chromatography 796(1998)299

Dihydropyranoimidazopyridines

Enantiomeric resolution on Chiral-AGP with the aid of experimental design. Unusual effects of mobile phase pH and column temperature 763(1997)105

Dihydropyridines

Influence of chromatographic descriptors on enantioresolution of a dihydropyridine and structurally related compounds 763(1997)115

Retention modeling in micellar liquid chromatography 780(1997)149

Dihydroxydithiane

Process clearance of dithiothreitol monitored by reversed-phase high-performance liquid chromatography with fluorescence detection. Assay development and validation 775(1997)151

Dihydroxynaphthalene

On-line concentration of neutral analytes for micellar electrokinetic chromatography. IV. Field-enhanced sample injection 798(1998)251

Dihydroxythiobenzanilides

Influence of modifier and molecular structure of some dihydroxythiobenzanilides on retention in reversed-phase high-performance thin-layer chromatography 791(1997)237

Diisopropylaminoethanol

Application of liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry, and tandem mass spectrometry, to the analysis and identification of degradation products of chemical warfare agents 759(1997)79

 β -Diketonates

Investigation of adducts of lanthanide and uranium β -diketonates with organophosphorus Lewis bases by supercritical fluid chromatography 793(1998)107

 β -Diketones

Supercritical fluid extraction: metals as complexes (Review) 785(1997)369

Diltiazem

On-line column-switching high-performance liquid chromatography analysis of cardiovascular drugs in serum with automated sample clean-up and zone-cutting technique to perform chiral separation 762(1997)235

Enantioseparation using selected polysaccharides as chiral buffer additives in capillary electrophoresis 773(1997)331

Diltiazem hydrochloride

Chiral resolution of four optical isomers of diltiazem hydrochloride on Chiralcel columns by packed-column supercritical fluid chromatography 785(1997)185

Dimethindene

Comparative capillary electrophoresis and NMR studies of enantioseparation of dimethindene with cyclodextrins 798(1998)315

Dimethoxyanthracene-2-sulfonate

Analysis of benzalkonium chloride in the effluent from European hospitals by solid-phase extraction and high-performance liquid chromatography with post-column ion-pairing and fluorescence detection 774(1997)281

N-Dimethylaminomethylene

Analysis of heterocyclic amines as their N-dimethylaminomethylene derivatives by gas chromatography with nitrogen–phosphorus selective detection 767(1997)187

N,N-Dimethylaniline

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Dimethylarsinic acid

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. I. Large-volume stacking with polarity switching 757(1997)281

Dimethylbenzanthracene

Isolation and recovery of 9,10-dimethyl-1,2-benzanthracene from animal bedding by reversed-phase solid-phase extraction combined with high-performance liquid chromatography 774(1997)223

3,5-Dimethylphenylcarbamate

3,5-Dimethylphenylcarbamates of amylose, chitosan and cellulose bonded on silica gel. Comparison of their chiral recognition abilities as high-performance liquid chromatography chiral stationary phases 796(1998)265

Dimethylquinoline

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Dimethyl sulfide

Deposition parameters of air pollutants on solid surfaces, measured in the presence of surface and gaseous reactions, with a simultaneous determination of the experimental isotherms 775(1997)211

Gas chromatographic–mass spectrometric analysis of mercaptan odorants in liquefied petroleum gas and liquefied natural gas 791(1997)367

2,4-Dinitroaniline

Influence of the capillary dimensions on the performance of the preconcentration technique based on parallel current chromatography 791(1997)163

Dinitrofluorobenzene

Analysis of primary and secondary aliphatic amines in waste water and surface water by gas chromatography–mass spectrometry after derivatization with 2,4-dinitrofluorobenzene or benzenesulfonyl chloride 764(1997)85

Dinitrophenylhydrazine

Impinger sampling coupled to high-performance liquid chromatography by a modified autoinjector interface 762(1997)187

Dinitrophenylhydrazines

High-performance liquid chromatographic peak identification of 2,4-dinitrophenylhydrazine derivatives of lipid peroxidation aldehydes by photodiode array detection 798(1998)117

2,4-Dinitrophenylhydrazone

Qualitative and quantitative high-performance liquid chromatographic analysis of aldehydes in Brazilian sugar cane spirits and other distilled alcoholic beverages 782(1997)13

Dinophysistoxin-2

Liquid chromatography with fluorimetric, mass spectrometric and tandem mass spectrometric detection for the investigation of the seafood-toxin-producing phytoplankton, *Dinophysis acuta* 777(1997)213

Dinophysistoxins

Optimization of sample cleanup procedure for determination of diarrhetic shellfish poisoning toxins by use of experimental design 764(1997)223

Improvement on sample clean-up for high-performance liquid chromatographic–fluorimetric determination of diarrhetic shellfish toxins using 1-bromoacetylpyrene 793(1998)63

Diocetyl sulfosuccinate

Determination of berberine and palmatine in *Phellodendri Cortex* using ion-pair supercritical fluid chromatography on-line coupled with ion-pair supercritical fluid extraction by on-column trapping 786(1997)371

Diols

Enantiomeric separation of diols and β -amino alcohols by chiral stationary phase derived from (*R,R*)-tartramide 757(1997)73

Determination of the diol content of chromatographic supports by capillary electrophoresis 758(1997)255

Chiral resolution of diols by capillary electrophoresis using borate–cyclodextrin complexation 761(1997)269

Gas–liquid chromatography–mass spectrometry of primary and secondary fatty alcohols and diols as their *tert.*-butyldimethylsilyl derivatives 771(1997)191

Dioxane

Rapid determination of 1,4-dioxane in water by solid-phase extraction and gas chromatography–mass spectrometry 787(1997)283

4,7-Diphenyl-1,10-phenanthroline disulfonate

Reversed-phase high-performance liquid chromatography of iron(II) and copper(II) chelates with 4,7-diphenyl-1,10-phenanthroline disulfonate 799(1998)171

Diphenylethanediamine

Diphenylethanediamine derivatives as chiral selectors. VIII. Influence of the second amido function on the high-performance liquid chromatographic enantioseparation characteristics of (*N*-3,5-dinitrobenzoyl)-diphenylethanediamine based chiral stationary phases 799(1998)67

Diphenylpentimamine

Diphenylethanediamine derivatives as chiral selectors. VIII. Influence of the second amido function on the high-performance liquid chromatographic enantioseparation characteristics of (*N*-3,5-dinitrobenzoyl)-diphenylethanediamine based chiral stationary phases 799(1998)67

Diphenylpropenoic acid esters

Chromatographic behaviour of diastereoisomers. XIII. Adsorptivity of esters of *Z*- and *E*-2,3-diphenylpropenoic acids and similar compounds on silica in terms of the Snyder theory 758(1997)135

Diphenyltin

Speciation of organo-tin compounds using liquid chromatography–atmospheric pressure ionisation mass spectrometry and liquid chromatography–inductively coupled plasma mass spectrometry as complementary techniques 794(1998)211

Diphosphine

Direct enantiomeric resolution of diphosphine and diphosphine oxide ligands by high-performance liquid chromatography 795(1998)289

Diphosphine oxide

Direct enantiomeric resolution of diphosphine and diphosphine oxide ligands by high-performance liquid chromatography 795(1998)289

Disaccharides

see also Saccharides

Analysis of a monophosphoryl lipid A immunostimulant preparation from *Salmonella minnesota* R595 by high-performance liquid chromatography 767(1997)53

Disinfectants

Analysis of benzalkonium chloride in the effluent from European hospitals by solid-phase extraction and high-performance liquid chromatography with post-column ion-pairing and fluorescence detection 774(1997)281

Diterpenoids

Analysis of diterpenoids from *Sideritis* species by reversed-phase high-performance liquid chromatography 778(1997)421

Dithiocarbamates

Supercritical fluid extraction: metals as complexes (Review) 785(1997)369

Dithiothreitol

Process clearance of dithiothreitol monitored by reversed-phase high-performance liquid chromatography with fluorescence detection. Assay development and validation 775(1997)151

Dithizone sulfonate

Capillary electrophoresis determination of methylmercury in fish and crab meat after extraction as the dithizone sulphonate complex 791(1997)333

Dithizone sulfonate complexes

Development of a capillary electrophoretic method for the separation and determination of trace inorganic and organomercury species utilizing the formation of highly absorbing water soluble dithizone sulphonate complexes 765(1997)345

Diuron

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Direct determination of bromacil and diuron residues in environmental water samples by coupled-column liquid chromatography and large-volume injection 761(1997)322

Divinylsulfone

Assay for determining the number of reactive groups on gels used in affinity chromatography and its application to the optimisation of the epichlorohydrin and divinylsulfone activation reactions 752(1996)67

DNA

On-line preconcentration and high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbon–DNA adducts using copper phthalocyanine trisulfonic acid as a group-specific adsorbent 755(1996)205

Performance of a series of novel N-substituted acrylamides in capillary electrophoresis of DNA fragments 756(1996)255

Simultaneous separation and genetic typing of four short tandem repeat loci by capillary electrophoresis 768(1997)135

Analysis of DNA fragmentation using a dynamic size-sieving polymer solution in capillary electrophoresis 771(1997)319

Electromigration behaviour of DNA molecules at the free electrolyte–polymer solution interface 772(1997)243

Comparison of resolution of double-stranded and single-stranded DNA in capillary electrophoresis 772(1997)255

Application of capillary electrophoresis–fluorescence line-narrowing spectroscopy for on-line spectral characterization of closely related analytes 781(1997)73

Rapid sizing of polymorphic microsatellite markers by capillary array electrophoresis 781(1997)295

Method for detection of epsilon-secondary structure in the precore region of human hepatitis B virus DNA using a fluorescence-based polymerase chain reaction–single-strand-conformation polymorphism technique with capillary electrophoresis 781(1997)307

Separation of DNA sequencing fragments up to 1000 bases by using poly(ethylene oxide)-filled capillary electrophoresis 781(1997)315

Application of mixed mobile phases and a step gradient method in capillary electrochromatography for the separation of isomeric polycyclic aromatic hydrocarbon–deoxyribonucleoside adduct mixtures prepared in vitro 781(1997)327

DNA fragment analysis by capillary polymer sieving electrophoresis using poly(acryloylaminoethoxyethanol)-coated capillaries 781(1997)335

Separations of DNA fragments by capillary electrophoresis in N-substituted polyacrylamides 781(1997)347

Characterization of TO-PRO-3 as an intercalator for double-stranded DNA analysis with red diode laser-induced fluorescence detection 781(1997)357

Capillary electrophoresis for the detection of known point mutations by single-nucleotide primer extension and laser-induced fluorescence detection 781(1997)367

DNA adducts

Analysis of DNA adducts by capillary methods coupled to mass spectrometry: a perspective 794(1998)97

DNA oligomer

Ketone electrophores and an olefin-release group electrophore-labeled DNA oligomer. Detection via electron capture 764(1997)95

Dodecamolybdophosphoric acid

Aminosilica chemically modified with dodecamolybdophosphoric acid as stationary phase in high-performance liquid chromatography 796(1998)259

Dodecanedioic acid polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions 775(1997)157

Dodecoxycarbonylvaline

Separation of aldose enantiomers by capillary electrophoresis in the presence of optically active N-dodecoxycarbonylvalines 792(1997)385

Dodecyl sulfate

Migration behaviour of micelle counterions in micellar electrokinetic chromatography: influence on micelle mobility, efficiency and selectivity 764(1997)127

DOPA

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Dopamine

Enhanced temporal resolution for the microdialysis monitoring of catecholamines and excitatory amino acids using capillary electrophoresis with laser-induced fluorescence detection. Analytical developments and in vitro validations 755(1996)99

Doxazosin

Chiral bioanalysis by normal phase high-performance liquid chromatography-atmospheric pressure ionization tandem mass spectrometry 759(1997)65

Drugs

see also Basic drugs and Special Issue Index

Enantiomer separation on a Chirasil-Dex-polymer-coated stationary phase by conventional and micro-packed high-performance liquid chromatography 755(1996)299

Screening and identification of drugs in human hair by high-performance liquid chromatography-photodiode-array UV detection and gas chromatography-mass spectrometry after solid-phase extraction. A powerful tool in forensic medicine 762(1997)251

Use of high-performance liquid chromatography with photodiode-array UV detection for the creation of a 600-compound library. Application to forensic toxicology 763(1997)149

Chiral recognition and racemic resolution of drug enantiomers by electrophoresis based on host-guest complexation 763(1997)261

High-performance ion chromatography applied to free-radical mechanisms in drug design. The problem of ion analysis at high ionic strengths 770(1997)273

Separation and quantitation of debrisoquine and 4-hydroxydebrisoquine in human urine by capillary electrophoresis and high-performance liquid chromatography 778(1997)389

Investigation of the interactions between drugs and mixed bile salt/lecithin micelles. A characterization by micellar affinity capillary electrophoresis (MACE). Part III 781(1997)377

Chiral separation principles in capillary electrophoresis 792(1997)179

Determination of drugs in biological fluids by high-performance liquid chromatography with on-line sample processing 797(1998)203

Drugs, chiral

Separation of enantiomers of drugs by capillary electrophoresis. V. Hydroxypropyl- α -cyclodextrin as chiral solvating agent 793(1998)153

Drugs, non-basic

Combined use of temperature and solvent strength in reversed-phase gradient elution. IV. Selectivity for neutral (non-ionized) samples as a function of sample type and other separation conditions 756(1996)63

DU-6859

Novel quartz flow-cell as a post-column photochemical reactor for high-performance liquid chromatography 800(1998)377

DUP105

Application of a modified central composite design to optimize the capillary electrochromatographic separation of related S-oxidation compounds 769(1997)145

Dyes

see also Azo dyes

Determination of single-stranded oligodeoxynucleotides by capillary gel electrophoresis with laser induced fluorescence and on column derivatization 755(1996)271

Screening of a large number of dyes for the separation of human immunoglobulin G2 from the other immunoglobulin G subclasses. Immunoglobulin G2 enrichment on immobilized Procion Yellow HE-4R 766(1997)49

Identification and determination of red dyes in confectionery by ion-interaction high-performance liquid chromatography 767(1997)87

Analysis of sulphonated phthalocyanine dyes by capillary electrophoresis 770(1997)345

Application of a colour analyzer in quantitative thin-layer chromatography 771(1997)251

Effect of on-capillary large volume sample stacking on limits of detection in the capillary zone electrophoretic determination of selected drugs, dyes and metal chelates 772(1997)161

Speciation of chromium dyes by high-performance liquid chromatography with inductively coupled plasma mass spectrometric detection 773(1997)157

Determination of mono- and disulphonated azo dyes by liquid chromatography–atmospheric pressure ionization mass spectrometry 777(1997)177

Monitoring the products of condensation of 4-nitrobenzoyl chloride and 2,4-diaminobenzenesulphonic acid by ion-pair high-performance liquid chromatography 777(1997)370

Separation, comparison and identification of fountain pen inks by capillary electrophoresis with UV–visible and fluorescence detection and by proton-induced X-ray emission 781(1997)391

Effect of foaming power and foam stability on continuous concentration with foam counter-current chromatography 791(1997)53

High-performance liquid chromatography determination of direct and temporary dyes in natural hair colourings 796(1998)319

Development of system suitability tests for ion-interaction chromatography of colorants on reversed-phase packing materials 797(1998)65

Dynorphins

Electrospray ionization mass spectrometric and liquid chromatographic–mass spectrometric studies on the metabolism of synthetic dynorphin A peptides in brain tissue in vitro and in vivo 800(1998)59

E

Ecdysteroids

High-performance liquid chromatography coupled to nuclear magnetic resonance spectroscopy. Application to the ecdysteroids of *Silene otites* 799(1998)333

Econazole

Liquid chromatographic separation of the enantiomers of beclonazole and its potential impurities 769(1997)231

Ecstasy

Sensitive and selective determination of methylenedioxyated amphetamines by high-performance liquid chromatography with fluorimetric detection 787(1997)137

EDTA–iron(III) complex

Trace determination of iron in water at the $\mu\text{g/l}$ level by on-line coupling of capillary isotachopheresis and capillary zone electrophoresis with UV detection of the EDTA–Fe(III) complex 757(1997)297

Eicosanoids

Identification of eicosanoids in the red algae, *Gracilaria asiatica*, using high-performance liquid chromatography and electropray ionization mass spectrometry 795(1998)227

Endorhamnosidase, P22 tailspike

Monitoring the refolding pathway for a large multimeric protein using capillary zone electrophoresis 769(1997)315

β -Endorphin

Quantitative analysis of methionine enkephalin and β -endorphin in the pituitary by liquid secondary ion mass spectrometry and tandem mass spectrometry 794(1998)85

Endosulfan

Analysis of lindane, α - and β -endosulfan and endosulfan sulfate in greenhouse air by gas chromatography 765(1997)99

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

Application of C_{18} disks followed by gas chromatography techniques to degradation kinetics, stability and monitoring of endosulfan in water 795(1998)93

Endrin

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

Enkephalins

Optical resolution of phenylthiohydantoin-amino acids by capillary electrophoresis and identification of the phenylthiohydantoin-D-amino acid residue of [D-Ala²]-methionine enkephalin 752(1996)279

Determination of peptides by high-performance liquid chromatography with laser-induced fluorescence detection 763(1997)23

Gradient high-performance liquid chromatographic analysis of enkephalin peptides, their metabolites and enzyme inhibitors using combined ultraviolet and electrochemical detection. I. Optimization of separation and detection 766(1997)85

Quantitative analysis of methionine enkephalin and β -endorphin in the pituitary by liquid secondary ion mass spectrometry and tandem mass spectrometry 794(1998)85

Enzyme inhibitors

Determination of an in vivo metabolite of a human immunodeficiency virus protease inhibitor in human plasma by high-performance liquid chromatography with tandem mass spectrometry 762(1997)311

Urinary excretion of deprenyl metabolites 762(1997)321

Migration order reversal of enantiomers in capillary electrophoretic separation 764(1997)151

Gradient high-performance liquid chromatographic analysis of enkephalin peptides, their metabolites and enzyme inhibitors using combined ultraviolet and electrochemical detection. I. Optimization of separation and detection 766(1997)85

Optimisation of a simultaneous separation of sulphonamides, dihydrofolate reductase inhibitors and β -lactam antibiotics by capillary electrophoresis 768(1997)97

Pharmacokinetic screening for the selection of new drug discovery candidates is greatly enhanced through the use of liquid chromatography-atmospheric pressure ionization tandem mass spectrometry 777(1997)61

Affinity chromatography of thiol ester-containing proteins 799(1998)139

Chromatographic purification of human α_1 proteinase inhibitor from dissolved Cohn fraction IV-1 paste 800(1998)207

Enzymes

see also Proteins

Isocratic high-performance liquid chromatography of coenzyme A esters involved in the metabolism of 3*S*-hydroxy-3-methylglutaryl-coenzyme A. Detection of related enzyme activities in *Catharanthus roseus* plant cell cultures 752(1996)123

Detection of electrophoretically separated amylase inhibitors in starch-polyacrylamide gels 752(1996)261

New class of poly(vinyl alcohol) polymers as column-chromatography stationary phases for *Candida rugosa* lipase isoforms separation 753(1996)47

Large-scale high-performance liquid chromatography of enzymes for food applications 760(1997)81

High-performance liquid chromatography quantitation of limonin D-ring lactone hydrolase and limonoate dehydrogenase activities 760(1997)173

Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67

Monitoring the refolding pathway for a large multimeric protein using capillary zone electrophoresis 769(1997)315

In vitro glucuronidation of 7-hydroxycoumarin and determination of 7-hydroxycoumarin and 7-hydroxycoumarin glucuronide by capillary electrophoresis 772(1997)321

Automated microanalysis using magnetic beads with commercial capillary electrophoretic instrumentation 781(1997)197

Quaternary structural analysis of nucleoside diphosphate kinases using capillary electrophoresis 781(1997)251

Application of capillary electrophoresis to determination of enzyme activity and other properties of phosphatidylinositol-specific phospholipase C 781(1997)263

Study of the enzymatic transformation of fluorescently labeled oligosaccharides in human epidermoid cells using capillary electrophoresis with laser-induced fluorescence detection 781(1997)515

Purification of a Streptococcal deoxyribonuclease by affinity chromatography based on a DNA-cellulose matrix 788(1997)75

Preliminary study on hydrophobic interaction chromatography of *Chromobacterium viscosum* lipase on polypropylene glycol immobilized on Sepharose 796(1998)177

Micropreparation of tissue collagenase fragments of type I collagen in the form of surfactant-peptide complexes and their identification by capillary electrophoresis and partial sequencing 796(1998)181

High-performance liquid chromatography study of stereospecific microsomal enzymes catalysing the reduction of a potential cytostatic drug, oracin. Interspecies comparison 797(1998)197

Dye-ligand and metal chelate poly(2-hydroxyethylmethacrylate) membranes for affinity separation of proteins 799(1998)83

Isolation of starch branching enzyme I from potato using γ -cyclodextrin affinity chromatography 800(1998)382

Eperisone

Separation of basic drug enantiomers by capillary zone electrophoresis using ovoglycoprotein as a chiral selector 782(1997)281

Ephedrine

Analysis of basic pharmaceuticals by capillary electrophoresis in coated capillaries and on-line mass spectrometric detection 752(1996)271

Epicatechin

Study of the acetaldehyde induced polymerisation of flavan-3-ols by liquid chromatography-ion spray mass spectrometry 752(1996)85

New methods for spectrometric peak purity analysis in chromatography 800(1998)121

Epichlorohydrin

Assay for determining the number of reactive groups on gels used in affinity chromatography and its application to the optimisation of the epichlorohydrin and divinylsulfone activation reactions 752(1996)67

Epoxy hardeners

High-performance liquid chromatography-atmospheric pressure negative chemical ionisation mass spectrometry of 2,4-dinitrophenyl derivatives of amines. Applications in epoxy chemistry 777(1997)193

Epoxy resins

Identification of potential migrants from a vinyllic organosol varnish by gas chromatography-mass spectrometry and liquid chromatography-mass spectrometry 771(1997)366

High-performance liquid chromatography-atmospheric pressure negative chemical ionisation mass spectrometry of 2,4-dinitrophenyl derivatives of amines. Applications in epoxy chemistry 777(1997)193

Erbium

Separation of samarium, gadolinium, terbium, dysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

Ergot alkaloids

Capillary electrophoretic separation of herbicidal enantiomers applying ergot alkaloids 791(1997)339

Erythropoietin

Analysis of recombinant human erythropoietin in drug formulations by high-performance capillary electrophoresis 759(1997)177

Chromatographic separation of recombinant human erythropoietin isoforms 791(1997)109

Esculin

Application of high-speed counter-current chromatography to the separation of coumarin and related compounds 759(1997)47

Esters

Silicalite as a sorbent for solid-phase extraction 771(1997)45

New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259

Estradiol

Theoretical concepts of on-line liquid chromatographic-biochemical detection systems. I. Detection systems based on labelled ligands 787(1997)27

Estrogens

Theoretical concepts of on-line liquid chromatographic-biochemical detection systems. I. Detection systems based on labelled ligands 787(1997)27

Development and validation of an isocratic high-performance liquid chromatographic method for quantitative determination of phytoestrogens in soya bean 795(1998)377

On-line concentration of neutral analytes for micellar electrokinetic chromatography. IV. Field-enhanced sample injection 798(1998)251

Ethane

Effect of addition of a graphitized carbon black trap to a glass beads trap on the cryoconcentration of some non-methane hydrocarbons in ambient air 790(1997)117

Ethanol

Ethanol-sodium chloride-phosphate mobile phase for size-exclusion chromatography of poly(ethylene glycol) modified proteins 763(1997)337

Ethanolamines

Application of liquid chromatography-atmospheric pressure chemical ionisation mass spectrometry, and tandem mass spectrometry, to the analysis and identification of degradation products of chemical warfare agents 759(1997)79

Ethers

Chemically bonded chelates as selective complexing sorbents for gas chromatography v. Silica chemically modified by Cu(II) complexes via amino groups 757(1997)137

Ethoxylated dodecyl alcohols

High-performance liquid chromatographic analysis of polydisperse ethoxylated non-ionic surfactants in aqueous samples 752(1996)155

Ethoxylates

Determination of alkylphenols, alkylphenoethoxylates and linear alkylbenzenesulfonates in sediments by accelerated solvent extraction and supercritical fluid extraction 775(1997)187

Theory of liquid adsorption chromatography of macromolecules applied to polyethylene glycol and fatty alcohol ethoxylates 791(1997)21

Analysis of ethoxylated polymers by capillary electrophoresis in UV-transparent polymer networks and by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry 800(1998)13

Ethylamine

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

N-Ethylaniline

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Ethylbenzene

Headspace gas chromatographic determination of the plant cuticle-air partition coefficients for monocyclic aromatic hydrocarbons as environmental compartment 774(1997)213

Ethyl chloroformate

Quantitative analysis of fluorinated ethylchloroformate derivatives of protein amino acids and hydrolysis products of small peptides using chemical ionization gas chromatography-mass spectrometry 759(1997)111

Study of binding media in works of art by gas chromatographic analysis of amino acids and fatty acids derivatized with ethyl chloroformate 778(1997)373

Ethylene

Effect of addition of a graphitized carbon black trap to a glass beads trap on the cryoconcentration of some non-methane hydrocarbons in ambient air 790(1997)117

Ethylenediaminedi(hydroxyphenylacetic) acids

Chromatographic determination of commercial Fe(III) chelates of ethylenediaminetetraacetic acid, ethylenediaminedi(*o*-hydroxyphenylacetic) acid and ethylenediaminedi(*o*-hydroxy-*p*-methylphenylacetic) acid 789(1997)453

Ethylenediaminetetraacetic acid

Laser induced resonance energy transfer – a novel approach towards achieving high sensitivity in capillary electrophoresis. Part B: Applications for environmental testing 767(1997)217

Chromatographic determination of commercial Fe(III) chelates of ethylenediaminetetraacetic acid, ethylenediaminedi(*o*-hydroxyphenylacetic) acid and ethylenediaminedi(*o*-hydroxy-*p*-methylphenylacetic) acid 789(1997)453

Ethylene glycol

Separation of free and glucuronidated opioids by capillary electrophoresis in aqueous, binary and micellar media 764(1997)157

Ethylenethiourea

Determination of ethylenethiourea in food commodities by a two-step derivatization method and gas chromatography with electron-capture and nitrogen-phosphorus detection 765(1997)31

Ethylmorphine

Determination of morphine and its analogues in urine by on-line coupled reversed-phase liquid chromatography-gas chromatography with on-line derivatization 771(1997)360

Ethylphosphonic acid monoesters

Capillary electrophoresis separation of alkylphosphonic acid monoesters with indirect ultraviolet detection 779(1997)245

Etodolac

Chiral capillary electrophoresis-electrospray mass spectrometry coupling using vancomycin as chiral selector 800(1998)69

Eugenol

Analysis of volatile components of oak wood by solvent extraction and direct thermal desorption-gas chromatography-mass spectrometry 778(1997)427

Europium chelates

Indirect time-resolved fluorescence detection of both non-fluorescent and fluorescent compounds separated by high-performance liquid chromatography 787(1997)261

Explosives

Rapid on-line chromatographic determination of trace-level munitions in aqueous samples 753(1996)81

Trace analysis of military high explosives (2,4,6-trinitrotoluene and hexahydro-1,3,5-trinitro-1,3,5-triazine) in agricultural crops 775(1997)117

F**Fat**

Development of a supercritical fluid extraction method for determination of lipid classes and total fat in meats and its comparison with conventional methods 785(1997)345

Fatty acid amides

Reaction of fatty acid amides and ethanolamides with trimethylsulfonium hydroxide 773(1997)239

Fatty acid esters

see also Propanediol fatty acid esters

Gas chromatography-mass spectrometry coupled with pseudo-Sadler retention indices, for the identification of components in the essential oil of *Curcuma longa* L 760(1997)303

Fatty acid ethanolamides

Reaction of fatty acid amides and ethanolamides with trimethylsulfonium hydroxide 773(1997)239

Fatty acid ethyl esters

Determination of free fatty acids and their ethyl esters in musts and wines 776(1997)283

Supercritical fluid simulated moving bed chromatography 786(1997)309

Fatty acid methyl esters

Influence of gas chromatographic parameters on measurement of $^{13}\text{C}/^{12}\text{C}$ isotope ratios by gas-liquid chromatography-combustion isotope ratio mass spectrometry. I 752(1996)233

Combined use of temperature and solvent strength in reversed-phase gradient elution. IV. Selectivity for neutral (non-ionized) samples as a function of sample type and other separation conditions 756(1996)63

Quantitative evaluation of carbon isotopic fractionation during reversed-phase high-performance liquid chromatography 757(1997)307

Rapid procedure for the isolation and analysis of fatty acid and fatty alcohol fractions from wax esters of marine zooplankton 760(1997)309

Normal-phase separation effects with lipids on a silver ion high-performance liquid chromatography column 764(1997)337

Determination of free fatty acids and their ethyl esters in musts and wines 776(1997)283

Enantiomeric separation of various lipoxygenase derived monohydroxy polyunsaturated fatty acid methyl esters by high-performance liquid chromatography 790(1997)235

Fractionation of *cis*- and *trans*-oleic, linoleic, and conjugated linoleic fatty acid methyl esters by silver ion high-performance liquid chromatography 799(1998)329

Fatty acids

see also Acids, Carboxylic acids and Organic acids

Cell lipids of the *Candida lipolytica* yeast grown on methanol 755(1996)75

Stereospecific analysis of the triacylglycerol fraction and linear discriminant analysis in a climatic differentiation of Umbrian extra-virgin olive oils 758(1997)109

High-performance liquid chromatographic analysis of free long chain fatty acids produced during lipolysis by anaerobic digester sludge 760(1997)271

Gas chromatography-mass spectrometry coupled with pseudo-Sadtler retention indices, for the identification of components in the essential oil of *Curcuma longa* L 760(1997)303

Molar enthalpy and molar volume of methylene and benzene homologues in reversed-phase liquid chromatography 762(1997)35

Determination of higher fatty acids in oils by high-performance liquid chromatography with electrochemical detection 764(1997)177

Separation of isomeric hydroperoxides of unsaturated fatty acids by capillary electrophoresis 767(1997)249

Use of capillary gas chromatography for determining the hydrogenation level of edible oils 767(1997)335

Determination of anions in landfill leachates by ion chromatography 770(1997)203

Determination of long-chained fatty acids using non-aqueous capillary electrophoresis and indirect UV detection 771(1997)301

Characterisation of aldehydic acids in used and unused frying oils 776(1997)245

Determination of free fatty acids and their ethyl esters in musts and wines 776(1997)283

Study of binding media in works of art by gas chromatographic analysis of amino acids and fatty acids derivatized with ethyl chloroformate 778(1997)373

Comparison of two methods for the determination of fatty acid profiles in plasma and erythrocytes 778(1997)435

Preparation and liquid chromatographic analysis of propanediol fatty acid esters 779(1997)337

Separation of unsaturated fatty acids and related isomeric hydroperoxides by micellar electrokinetic chromatography 781(1997)215

Short chain fatty acids analysis by capillary electrophoresis and indirect UV detection or laser-induced fluorescence 781(1997)233

Capillary electrophoresis and indirect UV detection as a fast and simple analytical tool for bacterial taxonomy 781(1997)497

Coupling chemical derivatization reactions with supercritical fluid extraction (Review) 785(1997)239

Dual-mode high-speed counter-current chromatography: retention, resolution and examples 790(1997)17

Micellar electrokinetic chromatography of long chain saturated and unsaturated free fatty acids with neutral micelles. Considerations regarding selectivity and resolution optimization 792(1997)165

Gas-liquid chromatography-mass spectrometry of hydroxy fatty acids as their methyl esters *tert.*-butyldimethylsilyl ethers 793(1998)91

Silver ion high-performance liquid chromatography of isomeric *cis*- and *trans*-octadecenoic acids. Effect of the ester moiety and mobile phase composition 793(1998)275

Analysis of plant oils by subcritical fluid chromatography using pattern fitting 800(1998)317

Fatty acids, hydroxy

Determination of β -hydroxy fatty acids in sewage sludge by using selected ion monitoring 775(1997)287

Gas-liquid chromatography-mass spectrometry of hydroxy fatty acids as their methyl esters *tert.*-butyldimethylsilyl ethers 793(1998)91

Fatty alcohol ethoxylates

Theory of liquid adsorption chromatography of macromolecules applied to polyethylene glycol and fatty alcohol ethoxylates 791(1997)21

Fatty alcohols

Rapid procedure for the isolation and analysis of fatty acid and fatty alcohol fractions from wax esters of marine zooplankton 760(1997)309

Gas-liquid chromatography-mass spectrometry of primary and secondary fatty alcohols and diols as their *tert.*-butyldimethylsilyl derivatives 771(1997)191

Fenamiphos

Synthesis and use of deuterated fenamiphos and its metabolites as internal standards for mass spectrometric analysis in water 754(1996)207

Fenbendazole

Determination of fenbendazole, praziquantel and pyrantel pamoate in dog plasma by high-performance liquid chromatography 797(1998)237

Fenfluramine

Training software for chiral separations in capillary electrophoresis 772(1997)195

Fenoprofen

Training software for chiral separations in capillary electrophoresis 772(1997)195

Separation of profen enantiomers by capillary electrophoresis using cyclodextrins as chiral selectors 793(1998)165

Fenoxaprop

Capillary electrophoretic separation of herbicidal enantiomers applying ergot alkaloids 791(1997)339

Fenpropathrin

Enantiomeric resolution of pyrethroids by high-performance liquid chromatography with diode-laser polarimetric detection 754(1996)97

Fenthion

Identification of fenthion and temephos and their transformation products in water by high-performance liquid chromatography with diode array detection and atmospheric pressure chemical ionization mass spectrometric detection 777(1997)99

Ferulic acid

Micellar electrokinetic chromatography as an alternative to high-performance liquid chromatography for separation and determination of phenolic compounds in Japanese spirituous liquor 793(1998)409

Fibrinogen

Multiple peaks induced by domain-specific binding of fibrinogen in anion-exchange high-performance liquid chromatography 753(1996)63

Gradient chromatofocusing high-performance liquid chromatography. II. Theoretical aspects 762(1997)47

Fibrinogen degradation products

Gradient chromatofocusing high-performance liquid chromatography. I. Practical aspects 762(1997)207

Ficoll

Observation of Ficoll charge using size-exclusion chromatography 800(1998)181

Flamprop

Capillary electrophoretic separation of herbicidal enantiomers applying ergot alkaloids 791(1997)339

Flavan-3-ols

Study of the acetaldehyde induced polymerisation of flavan-3-ols by liquid chromatography-ion spray mass spectrometry 752(1996)85

Flavanone glycosides

Simultaneous separation of flavanone glycosides and polymethoxylated flavones in citrus juices using liquid chromatography 800(1998)171

Flavanones

Quantitative analysis of flavonols, flavones, and flavanones in fruits, vegetables and beverages by high-performance liquid chromatography with photo-diode array and mass spectrometric detection 799(1998)101

Simultaneous separation of flavanone glycosides and polymethoxylated flavones in citrus juices using liquid chromatography 800(1998)171

Flavans

High-performance liquid chromatographic determination of flavan fingerprints in plant extracts 775(1997)359

Flavones

Separation of naphthalene and flavone derivatives by micellar electrokinetic chromatography with double- and triple-chain surfactants 755(1996)147

Development and validation of an isocratic high-performance liquid chromatographic method for quantitative determination of phytoestrogens in soya bean 795(1998)377

Quantitative analysis of flavonols, flavones, and flavanones in fruits, vegetables and beverages by high-performance liquid chromatography with photo-diode array and mass spectrometric detection 799(1998)101

Flavonoids

Analysis of flavonoids from red clover by liquid chromatography–electrospray mass spectrometry 755(1996)127

Quantitative analysis of flavonoids by reversed-phase high-performance liquid chromatography 761(1997)315

Preparative separation of anthocyanins by gradient elution centrifugal partition chromatography 763(1997)345

Prediction of the gas chromatographic relative retention times of flavonoids from molecular structure 771(1997)213

High-performance liquid chromatographic determination of flavan fingerprints in plant extracts 775(1997)359

Mass spectrometric characterization of flavonoids in extracts from *Passiflora incarnata* 777(1997)223

Identification of antioxidative compounds in plant beverages by capillary electrophoresis with the marker index technique 781(1997)523

High-performance liquid chromatography–electrospray mass spectrometry in phytochemical analysis of sour orange (*Citrus aurantium* L.) 791(1997)127

Separation of hydroxylated and methoxylated flavonoids by micellar electrokinetic capillary chromatography. Determination of analyte partition coefficients between aqueous and sodium dodecyl sulfate micellar phases 791(1997)289

High-performance liquid chromatographic determination of flavonoids in *Betula pendula* and *Betula pubescens* leaves 793(1998)370

Overview of the applications of liquid chromatography–mass spectrometry interfacing systems in food analysis: naturally occurring substances in food 794(1998)263

Prediction of migration behaviour of flavonoids in capillary zone electrophoresis by means of topological indices 798(1998)233

Analysis of secreted flavonoids of *Cistus ladanifer* L. by high-performance liquid chromatography–particle beam mass spectrometry 799(1998)111

Separation of tea polyphenols using micellar electrokinetic chromatography with diode array detection 799(1998)309

Simultaneous separation of flavanone glycosides and polymethoxylated flavones in citrus juices using liquid chromatography 800(1998)171

Flavonols

Quantitative analysis of flavonols, flavones, and flavanones in fruits, vegetables and beverages by high-performance liquid chromatography with photo-diode array and mass spectrometric detection 799(1998)101

Fluazifop

Capillary electrophoretic separation of herbicidal enantiomers applying ergot alkaloids 791(1997)339

Flumequine

Determination of flumequine and oxolinic acid in fortified chicken tissue using on-line dialysis and high-performance liquid chromatography with fluorescence detection 799(1998)349

Flumetasone

Immunoaffinity chromatography combined on-line with high-performance liquid chromatography–mass spectrometry for the determination of corticosteroids 794(1998)37

Flunixin glucuronide

Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67

Fluoranthene

Influence of the capillary dimensions on the performance of the preconcentration technique based on parallel current chromatography 791(1997)163

N- α -9-Fluorenylmethoxycarbonyl-amino acid-N-carboxy anhydrides

Adsorption liquid chromatography on silica for the chiral separation of amino acids and asymmetric amines derivatized with optically active N- α -9-Fmoc-amino acid-N-carboxyanhydrides 767(1997)69

Fluorescein

Implementation of affinity solid-phases in continuous-flow biochemical detection 776(1997)169

High resolution multichannel fluorescence detection for capillary electrophoresis. Application to multicomponent analysis 788(1997)173

Fluoride

Elution delay of fluoride by chromatography of anions on silica gels coated with crown ether dyes 755(1996)43

Simultaneous determination of monofluorophosphate and fluoride in toothpaste by capillary electrophoresis 765(1997)353

Ion chromatographic procedures for analysis of total fluoride content in dentifrices 771(1997)355

Development of a capillary electrophoresis method with indirect photometric detection for the determination of anions related to anticaries 781(1997)457

Fluorinated compounds

Sol-gel-derived fluorinated stationary phase for open tubular electrochromatography 773(1997)65

Fluorobenzoic acids

Ultra-trace determination of fluorinated aromatic carboxylic acids in aqueous reservoir fluids using solid-phase extraction in combination with gas chromatography-mass spectrometry 793(1998)297

Fluoronaphthalene

Determination of 1,3-dichloro-2-propanol and 3-chloro-1,2-propanediol in papers treated with polyamidoamine-epichlorohydrin wet-strength resins by gas chromatography-mass spectrometry using selective ion monitoring 788(1997)195

Fluorophosphates

Ion chromatographic procedures for analysis of total fluoride content in dentifrices 771(1997)355

Fluoroquinolones

Determination of flumequine and oxolinic acid in fortified chicken tissue using on-line dialysis and high-performance liquid chromatography with fluorescence detection 799(1998)349

Fluorosurfactants

Capillary electrophoretic separation of acidic and basic proteins in the presence of cationic and anionic fluorosurfactants 757(1997)255

Flurazepam

Effect of on-capillary large volume sample stacking on limits of detection in the capillary zone electrophoretic determination of selected drugs, dyes and metal chelates 772(1997)161

Folpet

Determination of the fungicides folpet, captan and captafol by cloud-point preconcentration and high-performance liquid chromatography with electrochemical detection 754(1996)85

Resolution of folpet, procymidone and triazophos in high-performance liquid chromatography-diode array detection by using partial least squares calibration to cross-sections of spectrochromatograms 778(1997)183

Formoterol

Applications of simulated moving-bed chromatography to the separation of the enantiomers of chiral drugs 769(1997)101

Forskolin

D-Glucose, forskolin and cytochalasin B affinities for the glucose transporter Glut1. Study of pH and reconstitution effects by biomembrane affinity chromatography 776(1997)81

Separation of forskolin derivatives by automated multiple development high-performance thin-layer chromatography 791(1997)231

Fructose

Parameter estimation for the simulation of liquid chromatography 769(1997)59

Analysis of selected alkaloids and sugars in tobacco extract 775(1997)101

Fullerenes

Optimization of the high-performance liquid chromatographic separation of fullerenes using 1-methylnaphthalene as the mobile phase on a tetraphenylporphyrin-silica stationary phase 786(1997)31

Enantiomer separation of [60]fullerene derivatives by micro-column high-performance liquid chromatography using (*R*)-(-)-2-(2,4,5,7-tetranitro-9-fluorenylideneaminoxy) propionic acid as chiral stationary phase 791(1997)65

Fulvic acids

Comparative study for the separation of aquatic humic substances by electrophoresis 777(1997)355

Fumagillin

Monitoring of selected metabolites and biotransformation products from fermentation broths by high-performance liquid chromatography 763(1997)165

Fumaric acid

Gas chromatographic determination of organic acids from fruit juices by combined resin mediated methylation and extraction in supercritical carbon dioxide 785(1997)251

Fumarprotocetraric acid

Quantitative determination of protolicheterinic- and fumarprotocetraric acids in *Cetraria islandica* by high-performance liquid chromatography 757(1997)303

Fumonisin

Determination of mycotoxins produced by *Fusarium* isolates from banana fruits by capillary gas chromatography and high-performance liquid chromatography 778(1997)363

Furaltadone

Determination of nitrofurans residues in avian eggs by liquid chromatography-UV photodiode array detection and confirmation by liquid chromatography-ionspray mass spectrometry 777(1997)201

Furanochromones

Determination of furanochromones and pyranocoumarins in drugs and *Ammi visnaga* fruits by combined solid-phase extraction–high-performance liquid chromatography and thin-layer chromatography–high-performance liquid chromatography 797(1998)305

Furanone

New derivatization method for determination of 3-chloro-4(dichloromethyl)-5-hydroxy-2(5H)-furanone in water 790(1997)242

Furans

Gas chromatographic separation of volatile furans from birchwood smoke 753(1996)151

Furazolidone

Determination of furazolidone in animal feeds using liquid chromatography with UV and thermospray mass spectrometric detection 771(1997)349

Determination of nitrofurans residues in avian eggs by liquid chromatography–UV photodiode array detection and confirmation by liquid chromatography–ionspray mass spectrometry 777(1997)201

Fusion proteins

Conformational transformations of the fusion protein BR96 sFv-PE40 as monitored by micellar electrokinetic capillary chromatography and circular dichroism 766(1997)205

G**Gadolinium**

Separation of samarium, gadolinium, terbium, dysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

Gangliosides

Analysis of gangliosides by capillary zone electrophoresis and capillary zone electrophoresis–electrospray mass spectrometry 779(1997)195

Capillary electrophoresis of carboxylated carbohydrates. IV. Adjusting the separation selectivity of derivatized carboxylated carbohydrates by controlling the electrolyte ionic strength at subambient temperature and in the absence of electroosmotic flow 792(1997)75

Gases

see also Permanent gases

Helium direct current discharge ionization detector for gas chromatography 761(1997)169

Genistein

Development and validation of an isocratic high-performance liquid chromatographic method for quantitative determination of phytoestrogens in soya bean 795(1998)377

Genistin

Linearity and the limitations of least squares calibration 762(1997)73

Germanium

Direct determination of trimethylgermanium in water by on-column capillary gas chromatography with flame photometric detection using quartz surface-induced germanium emission 759(1997)119

Separation and determination of inorganic germanium and β -carboxyethylgermanium sesquioxide by high-performance ion-exclusion chromatography 789(1997)403

Gibberellins

Reversed-phase C_{18} high-performance liquid chromatography of gibberellins GA_3 and GA_1 782(1997)137

Erratum to “Reversed-phase C_{18} high-performance liquid chromatography of gibberellins GA_3 and GA_1 ”. [J. Chromatogr. A, 782 (1997) 137–139] 795(1998)399

Gingerol

Determination of cinnamaldehyde, cinnamic acid, paeoniflorin, glycyrrhizin and [6]-gingerol in the traditional Chinese medicinal preparation Kuei-chih-tang by cyclodextrin-modified micellar electrokinetic chromatography 771(1997)267

High-performance liquid chromatography–electrospray mass spectrometric analysis of pungent constituents of ginger 796(1998)327

Ginsenosides

Reversed-phase high-performance liquid chromatographic determination of ginsenosides of *Panax quinquefolium* 755(1996)11

Globin peptides

Generation of tryptic maps of α - and β -globin chains by capillary electrophoresis in isoelectric buffers 791(1997)313

Globulins

Rapid separation of soybean globulins by reversed-phase high-performance liquid chromatography 758(1997)75

Glucodigifucoside

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Glucosylated cardenolides

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Glucogitoroside

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Glucose

Glucose metabolism and bioreduction of 2-butanone by *Candida utilis* studied by means of ion-exchange chromatography 763(1997)205

Parameter estimation for the simulation of liquid chromatography 769(1997)59

Analysis of selected alkaloids and sugars in tobacco extract 775(1997)101

D-Glucose, forskolin and cytochalasin B affinities for the glucose transporter Glut1. Study of pH and reconstitution effects by biomembrane affinity chromatography 776(1997)81

Glucose affinity for the glucose transporter Glut1 in native or reconstituted lipid bilayers. Temperature-dependence study by biomembrane affinity chromatography 776(1997)87

Glucosinolates

Determination of glucosinolates in domestic and wild mustard by high-performance liquid chromatography with confirmation by electrospray mass spectrometry and photodiode-array detection 767(1997)43

Overview of the applications of liquid chromatography–mass spectrometry interfacing systems in food analysis: naturally occurring substances in food 794(1998)263

Glucuronidase

Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67

In vitro glucuronidation of 7-hydroxycoumarin and determination of 7-hydroxycoumarin and 7-hydroxycoumarin glucuronide by capillary electrophoresis 772(1997)321

Glucuronides

Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67

In vitro glucuronidation of 7-hydroxycoumarin and determination of 7-hydroxycoumarin and 7-hydroxycoumarin glucuronide by capillary electrophoresis 772(1997)321

Analysis of catechol-type glucuronides in urine samples by liquid chromatography–electrospray ionization–tandem mass spectrometry 794(1998)75

Glufosinate

Selective analysis of the herbicides glyphosate and aminomethylphosphonic acid in water by on-line solid-phase extraction–high-performance liquid chromatography–electrospray ionization mass spectrometry 794(1998)187

Glut1

D-Glucose, forskolin and cytochalasin B affinities for the glucose transporter Glut1. Study of pH and reconstitution effects by biomembrane affinity chromatography 776(1997)81

Glucose affinity for the glucose transporter Glut1 in native or reconstituted lipid bilayers. Temperature-dependence study by biomembrane affinity chromatography 776(1997)87

Glutamate

Enhanced temporal resolution for the microdialysis monitoring of catecholamines and excitatory amino acids using capillary electrophoresis with laser-induced fluorescence detection. Analytical developments and in vitro validations 755(1996)99

Glutamic acids

Enantioseparation of 3-phenylacetyl-amino-2,6-piperidinedione and related chiral compounds 752(1996)93

Glutamines

Enantioseparation of 3-phenylacetyl-amino-2,6-piperidinedione and related chiral compounds 752(1996)93

Glutathione

Determination of glutathione, cysteine and N-acetylcysteine in rabbit eye tissues using high-performance liquid chromatography and post-column derivatization with 5,5'-dithiobis(2-nitrobenzoic acid) 778(1997)347

Glutathione derivatives

New and simple method for the analysis of the glutathione adduct of atrazine 795(1998)53

Glyburide

Application of liquid chromatography–mass spectrometryⁿ analyses to the characterization of novel glyburide metabolites formed in vitro 794(1998)15

Glycans

Exoglycosidase matrix-mediated sequencing of a complex glycan pool by capillary electrophoresis 781(1997)547

Detection of glycosaminoglycans as a copper(II) complex in high-performance liquid chromatography 787(1997)266

Glycerides

Quantitative aspects of triglyceride analysis by gas–liquid chromatography using a short metal capillary column 773(1997)233

Glycidyl methacrylate

Selection of a precursor monomer for the introduction of affinity ligands onto a porous membrane by radiation-induced graft polymerization 758(1997)209

Glycines

High-performance liquid chromatographic separation of enantiomers of 1,1'-binaphthyl-substituted α -aminoisobutyric acid 790(1997)41

Chromatographic resolution of synthetically useful chiral glycine derivatives by high-performance liquid chromatography 796(1998)299

Glycolic acid

Determination of glycolic acid in cosmetic products by solid-phase extraction and reversed-phase ion-pair high-performance liquid chromatography 795(1998)219

Glycolipids

Separation and quantitation of glycolipids as penetration modifiers in human skin using high-performance liquid chromatography-mass spectrometry with electrospray ionization 766(1997)71

Glycols

Effect of polystyrene-divinylbenzene resin sulfonation on solute retention in high-performance liquid chromatography 797(1998)139

Glycoproteins

Analysis of recombinant human erythropoietin in drug formulations by high-performance capillary electrophoresis 759(1997)177

Analysis of monosaccharide composition by capillary electrophoresis 763(1997)271

Study of haptoglobin-hemoglobin complexes by titration curves, capillary electrophoresis and capillary isoelectric focusing 767(1997)255

Critical conditions for separating the microheterogeneous components of glycoproteins by capillary electrophoresis 768(1997)39

Identification of α_1 -acid glycoprotein (orosomucoid) in human synovial fluid by capillary electrophoresis 772(1997)305

Interactions of the human class II major histocompatibility complex protein HLA-DR4 with a peptide ligand demonstrated by affinity capillary electrophoresis 781(1997)91

Characterization of protein glycoforms by capillary-zone electrophoresis-nanoelectrospray mass spectrometry 794(1998)327

Micro-sequencing strategies for the human A33 antigen, a novel surface glycoprotein of human gastrointestinal epithelium 798(1998)91

Glycosaminoglycans

Ion-pair high-performance liquid chromatography for determining disaccharide composition in heparin and heparan sulphate 765(1997)169

Coupling high-performance size exclusion and ion chromatography for the analysis of low-molecular-mass heparin 775(1997)91

Detection of glycosaminoglycans as a copper(II) complex in high-performance liquid chromatography 787(1997)266

Glycosides

Preparative separation of anthocyanins by gradient elution centrifugal partition chromatography 763(1997)345

Fractionation of glycoside precursors of aroma in grapes and wine 778(1997)219

Capillary electrophoretic analysis of salicin in *Salix* spp 781(1997)487

Porous graphitized carbon and octadecyl-silica columns in the separation of some alkylglycoside detergents 782(1997)191

High-performance liquid chromatographic determination of flavonoids in *Betula pendula* and *Betula pubescens* leaves 793(1998)370

Simultaneous separation of flavanone glycosides and polymethoxylated flavones in citrus juices using liquid chromatography 800(1998)171

Glycosphingolipids

Analysis of gangliosides by capillary zone electrophoresis and capillary zone electrophoresis-electrospray mass spectrometry 779(1997)195

Glycyrrhizin

Determination of cinnamaldehyde, cinnamic acid, paeoniflorin, glycyrrhizin and [6]-gingerol in the traditional Chinese medicinal preparation Kuei-chih-tang by cyclodextrin-modified micellar electrokinetic chromatography 771(1997)267

Glyphosate

Selective analysis of the herbicides glyphosate and aminomethylphosphonic acid in water by on-line solid-phase extraction-high-performance liquid chromatography-electrospray ionization mass spectrometry 794(1998)187

Gold cyanide complexes

Determination of metal cyanide complexes in gold processing solutions by capillary electrophoresis 767(1997)319

Goserelin

Capillary zone electrophoresis-mass spectrometry as a tool in the stability research of the luteinising hormone-releasing hormone analogue goserelin 776(1997)319

Growth factor inhibitors

Use of fluorine-containing stationary phases for the separation of an alkyl bromide from its hydrocarbon analog by high-performance liquid chromatography 762(1997)159

Growth hormones

see also Hormones

Role of high-performance liquid chromatographic protein analysis in developing fermentation processes for recombinant human growth hormone, relaxin, antibody fragments and lymphotoxin 763(1997)31

Immobilized metal affinity chromatography of human growth hormone. Effect of ligand density 764(1997)27

Analysis of recombinant human growth hormone directly in osmotic shock fluids 782(1997)199

Guaifenesin

Applications of simulated moving-bed chromatography to the separation of the enantiomers of chiral drugs 769(1997)101

Simulated moving bed chromatographic resolution of a chiral antitussive 796(1998)239

Guanidines

Chiral separation of 3-phenyl-3-(2-pyridyl)propylamines, and analogous guanidines and guanidine-N-carboxylic acid esters with high-performance liquid chromatography and capillary zone electrophoresis 793(1998)77

Guanylin

Matrix-assisted laser desorption/ionisation mass spectrometry guided purification of human guanylin from blood ultrafiltrate 776(1997)139

Gysprosium

Separation of samarium, gadolinium, terbium, gysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

H

Haemoglobin

Study of haptoglobin-hemoglobin complexes by titration curves, capillary electrophoresis and capillary isoelectric focusing 767(1997)255

Strategies to improve performance of capillary isoelectric focusing 772(1997)145

Recycling electrophoretic separations: modeling of isotachopheresis and isoelectric focusing 779(1997)165

Haemorphins

Rapid analysis of endogenous LVV-hemorphin-7 in cerebrospinal fluid by size-exclusion chromatography and electrospray ionization mass spectrometry 777(1997)41

Halocarbons

see Organochlorine compounds

Haloethers

Determination of haloethers in water by solid-phase microextraction 769(1997)239

Halogen anions

Specific determination of bromate and iodate in ozonized water by ion chromatography with postcolumn derivatization and inductively-coupled plasma mass spectrometry 789(1997)259

Halogenated compounds

Enantiomeric separations of halogen-substituted 2-aryloxypropionic acids by high-performance liquid chromatography on a terguride-based chiral stationary phase 756(1996)119

Determination of volatile organic compounds in water by purge-and-trap gas chromatography coupled to atomic emission detection 768(1997)259

Halossifop

Capillary electrophoretic separation of herbicidal enantiomers applying ergot alkaloids 791(1997)339

Haptoglobin

Study of haptoglobin-hemoglobin complexes by titration curves, capillary electrophoresis and capillary isoelectric focusing 767(1997)255

Harmane alkaloids

Improved separation of six harmane alkaloids by high-performance capillary electrophoresis 761(1997)297

Heavy metals

see also Metals

New sorbents for removal of heavy metal ions: diamine-glow-discharge treated polyhydroxyethylmethacrylate microspheres 773(1997)169

Determination of polycyclic aromatic compounds and heavy metals in sludges from biological sewage treatment plants 774(1997)177

Applications of column liquid chromatography to inorganic analysis in agricultural research (Review) 789(1997)519

Cibacron Blue F3GA-attached polyvinylbutyral microbeads as novel magnetic sorbents for removal of Cu(II), Cd(II) and Pb(II) ions 793(1998)47

α -Helical coiled-coil peptides

Effect of mobile phase on the oligomerization state of α -helical coiled-coil peptides during high-performance size-exclusion chromatography 791(1997)85

Heparan sulphate

Ion-pair high-performance liquid chromatography for determining disaccharide composition in heparin and heparan sulphate 765(1997)169

Heparin

Ion-pair high-performance liquid chromatography for determining disaccharide composition in heparin and heparan sulphate 765(1997)169

Application of a β -cyclodextrin sulfate-immobilized precolumn to selective on-line enrichment and separation of heparin-binding proteins by column-switching high-performance liquid chromatography 769(1997)209

Coupling high-performance size exclusion and ion chromatography for the analysis of low-molecular-mass heparin 775(1997)91

Heparin lyases

Ion-pair high-performance liquid chromatography for determining disaccharide composition in heparin and heparan sulphate 765(1997)169

Heptachlor

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

Heroin

^{13}C Isotopic analysis of an acetaminophen and diacetylmorphine mixture 756(1996)203

Micellar electrokinetic chromatography of charged and neutral drugs in acidic running buffers containing a zwitterionic surfactant, sulfonic acids or sodium dodecyl sulphate. Separation of heroin, basic by-products and adulterants 760(1997)245

Determination of morphine and its analogues in urine by on-line coupled reversed-phase liquid chromatography–gas chromatography with on-line derivatization 771(1997)360

Application of micellar electrokinetic capillary chromatography to the analysis of illicit drug seizures 780(1997)265

Heteroboranes

High-performance liquid chromatographic separations of boron-cluster compounds (Review) 789(1997)497

Hexachlorobenzene

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

Hexachlorocyclohexanes

Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275

Retention time repeatability as a function of the injection automatism in the analysis of trace organochlorinated compounds with high-resolution gas chromatography 778(1997)23

Hexachlorocyclohexanes in human blood serum 778(1997)87

Hexachloronaphthalenes

Selectivity of electron-donor- and electron-acceptor-bonded silica packing materials for hydrophobic environmental contaminants in polar and non-polar eluents 786(1997)237

Hexahydro-1,3,5-trinitro-1,3,5-triazine

Trace analysis of military high explosives (2,4,6-trinitrotoluene and hexahydro-1,3,5-trinitro-1,3,5-triazine) in agricultural crops 775(1997)117

Hexahydroimidazopyrazinediones

Identification of hexahydroimidazo[1,2-*a*]pyrazine-3,6-diones and hexahydroimidazo[1,2-*a*]imidazo[1,2-*d*]pyrazine-3,8-diones, unusual products of silica-catalyzed amino acid thermal condensation and products of their thermal decomposition using coupled high-performance liquid chromatography–particle beam mass spectrometry and gas chromatography–Fourier transform infrared spectroscopy–mass spectrometry 776(1997)255

Hexahydrotrinitrotriazine

Rapid on-line chromatographic determination of trace-level munitions in aqueous samples 753(1996)81

Hexazinone

Determination of hexazinone and its metabolites in groundwater by capillary electrophoresis 793(1998)349

Hexobarbital

Ruggedness of enantiomeric separation by capillary electrophoresis and high-performance liquid chromatography with methylated cyclodextrins as chiral selectors 763(1997)139

Hexyl chloroformate

Improved procedure for *n*-hexyl chloroformate-mediated derivatization of highly hydrophilic substances directly in water: hydroxyaminic compounds 793(1998)307

Hippuric acid

On-line isotachopheresis–capillary zone electrophoresis versus sample self stacking capillary zone electrophoresis. Analysis of hippurate in serum 772(1997)283

Histamine

Capillary zone electrophoretic separation of biogenic amines: influence of organic modifier 755(1996)142

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

Determination of histamine and some other amines by high-performance capillary electrophoresis with on-line mode in-capillary derivatization 790(1997)177

Use of 2-chloroethylnitrosourea, a new type of pre-column derivatizing agent for the measurement of biogenic amines, by high-performance liquid chromatography with ultraviolet detection 795(1998)239

Liquid chromatography with luminol-based electrochemiluminescence detection. Determination of histamine 799(1998)57

Histamine agonists

Chiral separation of 3-phenyl-3-(2-pyridyl)propylamines, and analogous guanidines and guanidine-N-carboxylic acid esters with high-performance liquid chromatography and capillary zone electrophoresis 793(1998)77

Histidine

Gas chromatography–chemical ionization mass spectrometry in amino acid analysis of pyoverdins 787(1997)195

Histidine as a dipolar eluent component in cation chromatography 789(1997)141

Histones

Use of reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene columns for the rapid separation and purification of acid-soluble nuclear proteins 763(1997)65

Application of hydrophilic-interaction liquid chromatography to the separation of phosphorylated H1 histones 782(1997)55

HLA-DR4

Interactions of the human class II major histocompatibility complex protein HLA-DR4 with a peptide ligand demonstrated by affinity capillary electrophoresis 781(1997)91

HMG14

Use of reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene columns for the rapid separation and purification of acid-soluble nuclear proteins 763(1997)65

HMG17

Use of reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene columns for the rapid separation and purification of acid-soluble nuclear proteins 763(1997)65

Homatropine

N-(Polyethyleneglycol monomethyl ether)-N-methylmorpholinium-based background electrolytes in capillary electrophoresis 763(1997)253

High-performance liquid chromatographic determination of atropine and atropine-like alkaloids in pharmaceutical preparations with indirect conductometric detection 766(1997)270

Training software for chiral separations in capillary electrophoresis 772(1997)195

Homoanatoxins

Sensitive determination of anatoxin-a, homoanatoxin-a and their degradation products by liquid chromatography with fluorimetric detection 798(1998)147

Homoaromoline

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Homocysteine

Quantitation of total homocysteine in human plasma by derivatization to its N(O,S)-propoxycarbonyl propyl ester and gas chromatography–mass spectrometry analysis 776(1997)342

Urinary excretion measurement of cysteine and homocysteine in the form of their S-pyridinium derivatives by high-performance liquid chromatography with ultraviolet detection 798(1998)27

Honokiol

Determination of magnolol and honokiol in Magnoliae Cortex using supercritical fluid chromatography on-line coupled with supercritical fluid extraction by on-column trapping 786(1997)366

Hormones

see also Growth hormones and Plant hormones

Use of capillary electrophoresis and high-performance liquid chromatography for monitoring of glycosylation of the peptides dalargin and desmopressin 761(1997)285

Matrix-assisted laser desorption/ionisation mass spectrometry guided purification of human guanylin from blood ultrafiltrate 776(1997)139

Quantitation of insulin by capillary electrophoresis and high-performance liquid chromatography. Method comparison and validation 781(1997)445

Anionic–zwitterionic mixed micelles in the micellar electrokinetic separation of clinically relevant steroids on a fused-silica capillary 781(1997)467

Analysis of recombinant human growth hormone directly in osmotic shock fluids 782(1997)199

Humic acids

Determination of terrestrial markers in marine environments by gas chromatography–mass-selective detection compared to high-performance liquid chromatography–fluorescence detection 758(1997)243

Solid-phase microextraction of volatile organic compounds. Estimation of the sorption equilibrium from the Kováts index, effect of salinity and humic acids and the study of the kinetics by the development of an "agitated/static layer" model 761(1997)205

Polyacrylamide gel electrophoresis of soil humic acid fractionated by size-exclusion chromatography and ultrafiltration 767(1997)285

Comparative study for the separation of aquatic humic substances by electrophoresis 777(1997)355

Humulones

Preparative separation of bitter acids from hop extracts by centrifugal partition chromatography 771(1997)71

Hyaluronic acid

High-performance capillary electrophoresis of hyaluronic acid: determination of its amount and molecular mass 768(1997)295

Hydantoins

Facile and predictable means of separating the enantiomers of 5-arylhydantoins 790(1997)65

Hydrazides

Hydrazide as a ligand moiety in immobilized metal ion affinity chromatography. Separation of BO-IMI and BODIPY-hydrazide 777(1997)261

Hydrazinobenzothiazole

High-resolution gas chromatography-nitrogen-phosphorous detection of saturated volatile aldehydes derivatized with 2-hydrazinobenzothiazole 752(1996)209

Hydrocarbons

see also Volatile organic compounds

Chemically bonded chelates as selective complexing sorbents for gas chromatography v. Silica chemically modified by Cu(II) complexes via amino groups 757(1997)137

Solid-phase microextraction of volatile organic compounds. Estimation of the sorption equilibrium from the Kováts index, effect of salinity and humic acids and the study of the kinetics by the development of an "agitated/static layer" model 761(1997)205

Relationship between stationary and mobile phase composition and its influence on retention factors of aromatic hydrocarbons in reversed-phase high-performance liquid chromatography 762(1997)27

Integration of the equation of peak motion in programmed-pressure and -temperature gas chromatography 766(1997)147

Chemically bonded chelates as selective complexing sorbents for gas chromatography. VI. Modification of silica with NiCl₂ and CoCl₂ via β -diketonate groups 773(1997)209

Chromatographic study of the influence of nitrogen dioxide on the reactions between volatile hydrocarbons and inorganic pigments 776(1997)275

Refrigerated multibed adsorption in sampling and analysis of atmospheric light hydrocarbons at ppb (v/v) and sub-ppb (v/v) concentrations 778(1997)269

Application of Kováts retention indices for investigation of adsorption properties of activated carbons 782(1997)87

Determination of total petroleum hydrocarbons in soil by on-line supercritical fluid extraction-infrared spectroscopy using a fiber-optic transmission cell and a simple filter spectrometer 785(1997)269

Intermolecular interactions in adsorbate-carbon systems of various nano- and adsorption textures 786(1997)117

Hydrocarbons, aromatic

see also Monocyclic aromatic hydrocarbons

Comparative studies of chromatographic properties of silica-based amide-bonded phases under hydro-organic conditions 797(1998)103

Hydrocarbons, gaseous

Deposition parameters of air pollutants on solid surfaces, measured in the presence of surface and gaseous reactions, with a simultaneous determination of the experimental isotherms 775(1997)211

Hydrochlorothiazide

Use of simulated liquid chromatography-diode array detection data for the definition of a guide curve in peak purity assessment by spectral comparison 786(1997)1

Hydrochlorthiazide

Simulations of liquid chromatography-diode array detector data including instrumental artefacts for the evaluation of mixture analysis techniques 758(1997)1

Hydronium ion

Water clusters studied by electrospray mass spectrometry 777(1997)155

Hydroperoxides

Separation of isomeric hydroperoxides of unsaturated fatty acids by capillary electrophoresis 767(1997)249

Separation of unsaturated fatty acids and related isomeric hydroperoxides by micellar electrokinetic chromatography 781(1997)215

Hydroxamic acids

New fluorinated hydroxamic acid reagents for the extraction of metal ions with supercritical CO₂ 770(1997)85

Hydroxanthommatin

Detection of the oxidative products of 3-hydroxykynurenine using high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–electron spin resonance spectrometry and high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–mass spectrometry 773(1997)23

Hydroxyaminic compounds

Improved procedure for *n*-hexyl chloroformate-mediated derivatization of highly hydrophilic substances directly in water: hydroxyaminic compounds 793(1998)307

Hydroxyarginine

High-performance ion chromatography applied to free-radical mechanisms in drug design. The problem of ion analysis at high ionic strengths 770(1997)273

Hydroxybenzaldehydes

Determination of hydroxycinnamic acids, hydroxybenzoic acids, hydroxybenzaldehydes, hydroxybenzyl alcohols and their glucosides by high-performance liquid chromatography 756(1996)129

Hydroxybenzoates

Determination of preservatives in food products by cyclodextrin-modified capillary electrophoresis with multiwavelength detection 768(1997)334

Separation of neutral substances by non-aqueous capillary electrophoresis through interactions with cationic additives 792(1997)475

Hydroxybenzoic acid

Determination of hydroxycinnamic acids, hydroxybenzoic acids, hydroxybenzaldehydes, hydroxybenzyl alcohols and their glucosides by high-performance liquid chromatography 756(1996)129

Hydroxybenzyl alcohols

Determination of hydroxycinnamic acids, hydroxybenzoic acids, hydroxybenzaldehydes, hydroxybenzyl alcohols and their glucosides by high-performance liquid chromatography 756(1996)129

Hydroxycinnamic acids

Determination of hydroxycinnamic acids, hydroxybenzoic acids, hydroxybenzaldehydes, hydroxybenzyl alcohols and their glucosides by high-performance liquid chromatography 756(1996)129

Application of high-speed counter-current chromatography to the separation of coumarin and related compounds 759(1997)47

Hydroxycoumarin

In vitro glucuronidation of 7-hydroxycoumarin and determination of 7-hydroxycoumarin and 7-hydroxycoumarin glucuronide by capillary electrophoresis 772(1997)321

Hydroxycoumarin glucuronide

In vitro glucuronidation of 7-hydroxycoumarin and determination of 7-hydroxycoumarin and 7-hydroxycoumarin glucuronide by capillary electrophoresis 772(1997)321

Hydroxydebrisoquine

Separation and quantitation of debrisoquine and 4-hydroxydebrisoquine in human urine by capillary electrophoresis and high-performance liquid chromatography 778(1997)389

Hydroxydeoxyuridine

New hydrolytic products detected in aqueous solutions of bromodeoxyuridine 797(1998)283

3-(2-Hydroxyethyl)indole

Analysis of ethereal extracts of wines and other alcoholic beverages by high-performance liquid chromatography with microbore columns 782(1997)219

Hydroxyethylidenediphosphonic acid

Characterization of metal complexes of pharmaceutical interest by capillary electrophoresis with element sensitive detection 781(1997)407

Hydroxyguanidine

High-performance ion chromatography applied to free-radical mechanisms in drug design. The problem of ion analysis at high ionic strengths 770(1997)273

Hydroxyhistidine

Gas chromatography–chemical ionization mass spectrometry in amino acid analysis of pyoverdins 787(1997)195

3-Hydroxykynurenine

Detection of the oxidative products of 3-hydroxykynurenine using high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–electron spin resonance spectrometry and high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–mass spectrometry 773(1997)23

Hydroxyl radicals

Determination of hydroxyl radicals in an advanced oxidation process with salicylic acid trapping and liquid chromatography 796(1998)283

5-Hydroxymethyl-2-furaldehyde

Analysis of ethereal extracts of wines and other alcoholic beverages by high-performance liquid chromatography with microbore columns 782(1997)219

Hydroxymethylglutaryl-coenzyme A

Isocratic high-performance liquid chromatography of coenzyme A esters involved in the metabolism of 3S-hydroxy-3-methylglutaryl-coenzyme A. Detection of related enzyme activities in *Catharanthus roseus* plant cell cultures 752(1996)123

Hydroxynaphthalenecarboxylic acids

Capillary zone electrophoresis of hydroxynaphthalenecarboxylic acids. Purity monitoring of β -hydroxynaphthoic acid in industry 772(1997)397

2-(4-Hydroxyphenyl)ethanol

Analysis of ethereal extracts of wines and other alcoholic beverages by high-performance liquid chromatography with microbore columns 782(1997)219

Hydroxypipelic acid

Pipelic acid methyl esters as artefacts from the ion-exchange chromatography of *Inga punctata* foliar extracts 766(1997)267

2-Hydroxypropyltrimethylammonium- β -cyclodextrin

Enantioseparation in capillary electrophoresis using 2-hydroxypropyltrimethylammonium salt of β -cyclodextrin as a chiral selector 771(1997)259

Hydroxyquinoline sulfate

High-performance liquid chromatographic assay of 8-hydroxyquinoline sulfate and its stability in immunobiological preparations 798(1998)173

Hydroxyquinolinesulfonic acid

Dynamic control and indirect absorption detection for high-speed capillary electrophoretic separation of organic acids 800(1998)339

Hydroxytetraenoic acids

Identification of eicosanoids in the red algae, *Gracilaria asiatica*, using high-performance liquid chromatography and electrospray ionization mass spectrometry 795(1998)227

Hydroxytryptophan

Determination of γ -radiation induced products in aqueous solutions of tryptophan and synthesis of 4-, 6- and 7-hydroxytryptophan 786(1997)57

Hyoscine

High-performance liquid chromatographic determination of atropine and atropine-like alkaloids in pharmaceutical preparations with indirect conductometric detection 766(1997)270

Hypochlorite

Simultaneous determination of chlorine dioxide and hypochlorite in water by high-performance liquid chromatography 796(1998)397

I**Ibuprofen**

Training software for chiral separations in capillary electrophoresis 772(1997)195

Thermodynamics of chiral selectivity in capillary electrophoresis: separation of ibuprofen enantiomers with β -cyclodextrin 792(1997)371

Separation of profen enantiomers by capillary electrophoresis using cyclodextrins as chiral selectors 793(1998)165

Direct injection of large volumes of plasma/serum on a new biocompatible extraction column for the determination of atenolol, propranolol and ibuprofen. Mechanisms for the improvement of chromatographic performance 797(1998)251

Separation of ibuprofen, codeine phosphate, their degradation products and impurities by capillary electrophoresis. I. Method development and optimization with fractional factorial design 798(1998)307

Chiral capillary electrophoresis–electrospray mass spectrometry coupling using vancomycin as chiral selector 800(1998)69

Imazalil

Chromatographic analysis of imazalil and carbendazim in fruits. Method validation and residue monitoring program 1995 765(1997)91

Imazamethabenz

Enantiomeric and isomeric separation of herbicides using cyclodextrin-modified capillary zone electrophoresis 790(1997)225

Imazaquin

Enantiomeric and isomeric separation of herbicides using cyclodextrin-modified capillary zone electrophoresis 790(1997)225

Imidacloprid

Determination of the insecticide imidacloprid in water and soil using high-performance liquid chromatography 787(1997)271

Resolution of imidacloprid pesticide and its metabolite 6-chloronicotinic acid using cross-sections of spectrochromatograms obtained by high-performance liquid chromatography with diode-array detection 799(1998)149

Imidazolinone herbicides

Development of a method based on liquid chromatography–electrospray mass spectrometry for analyzing imidazolinone herbicides in environmental water at part-per-trillion levels 800(1998)109

Imidodipeptides

Complete resolution of imidodipeptide mixtures in urine of prolidase-deficient patients using micellar electrokinetic chromatography 768(1997)57

A-IMilano

Rapid reversed-phase high-performance liquid chromatography method for quantitation, at high pH, of the recombinant apolipo-protein A-IMilano in *Escherichia coli* fermentation broth 798(1998)83

Immunoglobulins

Post-capillary affinity detection of protein microheterogeneity in capillary zone electrophoresis 759(1997)139

Screening of a large number of dyes for the separation of human immunoglobulin G2 from the other immunoglobulin G subclasses. Immunoglobulin G2 enrichment on immobilized Procion Yellow HE-4R 766(1997)49

Kinetic analysis of the interaction between the monoclonal antibody A33 and its colonic epithelial antigen by the use of an optical biosensor. A comparison of immobilisation strategies 776(1997)15

Conventional capillary electrophoresis in comparison with short-capillary capillary electrophoresis and microfabricated glass chip capillary electrophoresis for the analysis of fluorescein isothiocyanate anti-human immunoglobulin G 781(1997)287

Ligand structure of the divinylsulfone-based T-gel 787(1997)47

On-line post-capillary affinity detection of immunoglobulin G subclasses and monoclonal antibody variants for capillary electrophoresis 790(1997)207

Bovine whey fractionation based on cation-exchange chromatography 795(1998)277

Development of a downstream process for the isolation and separation of monoclonal immunoglobulin A monomers, dimers and polymers from cell culture supernatant 796(1998)165

Fast isolation of protein receptors from streptococci G by means of macroporous affinity discs 798(1998)65

Comparison of the protein adsorption selectivity of salt-promoted agarose-based adsorbents. Hydrophobic, thiophilic and electron donor-acceptor adsorbents 800(1998)151

Indan derivatives

Chiral separations of indan, tetralin and benzosuberane derivatives by capillary electrophoresis 793(1998)135

Indenestrols

Separation of indenestrol A and B isomers and enantiomers by high-performance liquid chromatography 799(1998)117

Indinavir

Enantioseparation of a protease inhibitor, indinavir, by subcritical fluid chromatography 752(1996)101

Determination of an in vivo metabolite of a human immunodeficiency virus protease inhibitor in human plasma by high-performance liquid chromatography with tandem mass spectrometry 762(1997)311

Indium

Selective fluorometric determination of indium(III) by high-performance liquid chromatography with 2-methyl-8-quinolinol based on a ligand-exchange reaction of silanol groups 789(1997)395

Indoleacetic acid

Determination of plant indoles by capillary electrophoresis with amperometric detection 755(1996)289

Determination of indole-3-acetic acid in the Gulf of Gdańsk by high-performance liquid chromatography of its 4-methyl-7-methoxycoumarin derivative 766(1997)261

Determination of indole-3-acetic acid in plant tissues by capillary electrophoresis 779(1997)342

Indoleacetylaspartic acid

Determination of plant indoles by capillary electrophoresis with amperometric detection 755(1996)289

Indoleamines

Microcolumn liquid chromatography: instrumentation, detection and applications 779(1997)1

Indolepyruvic acid

Determination of indole-3-pyruvic acid levels in *Arabidopsis thaliana* by gas chromatography-selected ion monitoring-mass spectrometry 800(1998)101

Indolequinazolines

Determination of the alkaloids in *Evodiae fructus* by capillary electrophoresis 755(1996)113

Indomethacin

Assessment of injection volume limits when using on-column focusing with microbore liquid chromatography 759(1997)1

Inorganic anions

see also Anions, Inorganic ions and Special Issue Index

Mixed-mode liquid chromatography of carboxylic acids and inorganic anions on a latex-based pellicular stationary phase 753(1996)253

Indirect photometric detection of inorganic anions in microcolumn ion chromatography using octadecylsilica immobilized with bovine serum albumin as stationary phase 755(1996)37

Elution delay of fluoride by chromatography of anions on silica gels coated with crown ether dyes 755(1996)43

Nonaqueous anion-exchange chromatography. I. Role of solvation in anion-exchange resin 758(1997)19

Nonaqueous anion-exchange chromatography. II. Changing anion-exchange selectivity by resin surface complex formation of crown ethers 758(1997)29

- Detection of traces of oxidized and reduced sulfur compounds in small samples by combination of different high-performance liquid chromatography methods 760(1997)295
- Gas chromatographic determination of nitrite in water by pre-column formation of 2-phenylphenol with flame ionization detection 760(1997)319
- Determination of anions in rainwater by capillary electrophoresis with conductivity detection 760(1997)326
- Preconcentration determination of inorganic anions and organic acids in power plant waters. Separation optimization through control of column capacity and selectivity 761(1997)163
- Simultaneous determination of organic acids and inorganic anions in tea by ion chromatography 764(1997)341
- Separation behavior of inorganic anions on an octadecyl-bonded silica column coated with cetyltrimethylammonium by phthalate and salicylate eluents 764(1997)346
- Electroosmotic flow mobilities in open and packed capillaries with spermine and other amine flow modifiers 768(1997)3
- Method development and validation for the simultaneous determination of organic and inorganic acids by capillary zone electrophoresis 765(1997)321
- Ion chromatographic separation of carboxylic acids. Prediction of retention data 770(1997)13
- Use of factorial experimental design for the rapid evaluation of main and interactive factors affecting linearity in calibration curves for sulfate analysis by ion chromatography 770(1997)29
- Studies on the combination of ion chromatography-particle-beam mass spectrometry with capillary columns 770(1997)59
- Ultra-pure water for ion chromatography 770(1997)115
- Performance of ion chromatography in the determination of anions and cations in various natural waters with elevated mineralization 770(1997)119
- Analysis of ions in polar ice core samples by use of large injection volumes in ion chromatography 770(1997)125
- Determination of nitrite and nitrate in human serum 770(1997)143
- The role of nitric oxide in cancer. Improved methods for measurement of nitrite and nitrate by high-performance ion chromatography 770(1997)151
- Determination of sulfur oxyanions by ion chromatography on a silica ODS column with tetrapropylammonium salt as an ion-pairing reagent 770(1997)157
- Monitoring the removal of soluble salts from ancient tiles by ion chromatography 770(1997)195
- Determination of anions in landfill leachates by ion chromatography 770(1997)203
- Determination of impurities in clodronic acid by anion-exchange chromatography 770(1997)261
- Comparison of ion chromatography and capillary electrophoresis for the determination of inorganic ions 770(1997)281
- Separation of inorganic anions in acidic solution by capillary electrophoresis 770(1997)301
- Determination of phosphate in natural waters by capillary electrophoresis. Influence of solution composition on migration time and response 770(1997)321
- Capillary zone electrophoretic determination of ionic impurities in silicone products used for electronic applications 770(1997)337
- Routine determination of major anions in atmospheric aerosols by capillary electrophoresis 770(1997)349
- Optimising the determination of nitrate and phosphate in sea water with ion chromatography using experimental design 771(1997)163
- Ion chromatographic procedures for analysis of total fluoride content in dentifrices 771(1997)355
- Determination of low-molecular-mass organic acids in fogwater by ion-exchange chromatography with automatic column switching 773(1997)123
- Ion chromatographic characterization of toxic solutions: analysis and ion chemistry of biological liquids (Review) 774(1997)21
- Thin-layer chromatographic separation, colorimetric determination and recovery of thiocyanate from photogenic waste, river and sea waters 774(1997)373
- Use of cationic polymers for the simultaneous determination of inorganic anions and metal-4-(2-pyridylazo)resorcinolato chelates in kinetic differentiation-mode capillary electrophoresis 776(1997)329
- Separation of chlorine-containing anions by ion chromatography and capillary electrophoresis 777(1997)375
- Micellar chromatography of inorganic compounds 780(1997)343
- Investigation of inorganic ions and organic acids in atmospheric aerosol by capillary electrophoresis 781(1997)223

- Development of a capillary electrophoresis method with indirect photometric detection for the determination of anions related to anticaries 781(1997)457
- Electro-elution ion chromatography: controlling sample retention by electrochemically generating or modifying the mobile phase 782(1997)69
- Electrochemically regenerated ion suppressors, new suppressors for ion chromatography 782(1997)75
- Sub-minute separations of organic and inorganic anions with co-electroosmotic capillary electrophoresis 787(1997)243
- Analysis of organic and inorganic selenium anions by ion chromatography–inductively coupled plasma atomic emission spectroscopy 788(1997)105
- Retention models in ion chromatography: the role of side equilibria in ion-exchange chromatography of inorganic cations and anions (Review) 789(1997)3
- Factors controlling ion-exchange selectivity in suppressed ion chromatography (Review) 789(1997)29
- Theoretical study of the elution profile of a high-concentration sample anion in non-suppressed ion chromatography 789(1997)43
- Applications of a new methacrylate-based anion stationary phase for the separation of inorganic anions 789(1997)127
- Capillary electrophoresis of inorganic anions and its comparison with ion chromatography (Review) 789(1997)169
- Separation and determination of inorganic anions by reversed-phase high-performance liquid chromatography (Review) 789(1997)181
- Anion separation by high-performance ion chromatography coupled with indirect photometric detection and 1,2-dihydroxybenzene-3,5-disulfonate or sodium sulfosalicylate as the eluent 789(1997)195
- Ion chromatography of anions on stationary phases modified with chondroitin sulfate 789(1997)201
- Methods for the determination of nitrite by high-performance liquid chromatography with electrochemical detection (Review) 789(1997)213
- Quantitation of sulfate and thiosulfate in clinical samples by ion chromatography (Review) 789(1997)221
- Specific determination of bromate and iodate in ozonized water by ion chromatography with postcolumn derivatization and inductively-coupled plasma mass spectrometry 789(1997)259
- Determination of organic and inorganic anions in insect haemolymph and midgut contents by ion chromatography 789(1997)279
- Liquid chromatographic separation of aqueous species of Cr(VI) and Cr(III) 789(1997)469
- Applications of column liquid chromatography to inorganic analysis in agricultural research (Review) 789(1997)519
- State-of-the-art ion chromatographic determination of inorganic ions in food 789(1997)529
- Determination of some inorganic species in edible vegetable oils and fats by ion chromatography 789(1997)549
- Analysis of difficult samples by capillary zone electrophoresis. I. Simultaneous separation of detrimental anions in concrete digested with concentrated nitric acid 793(1998)187
- Preparation of monodisperse agglomerated pellicular anion-exchange resins compatible with high-performance liquid chromatography solvents for ion chromatography 793(1998)193
- Novel polymeric resins for anion-exchange chromatography 793(1998)231
- Nitrite/nitrate balance during photoinduced cerebral ischemia in the rat determined by high-performance liquid chromatography with UV and electrochemical detection 798(1998)103
- Simultaneous separation of inorganic anions and cations by capillary zone electrophoresis 800(1998)369
- Inorganic cations**
see also Cations, Inorganic ions, Metal cations and Special Issue Index
- Analysis of individual raindrops by capillary zone electrophoresis 755(1996)121
- Use of mordenite columns in ion-exclusion chromatography 760(1997)292
- Ultra-pure water for ion chromatography 770(1997)115
- Performance of ion chromatography in the determination of anions and cations in various natural waters with elevated mineralization 770(1997)119
- Analysis of ions in polar ice core samples by use of large injection volumes in ion chromatography 770(1997)125
- Monitoring the removal of soluble salts from ancient tiles by ion chromatography 770(1997)195
- Comparison of ion chromatography and capillary electrophoresis for the determination of inorganic ions 770(1997)281

- Capillary zone electrophoretic determination of ionic impurities in silicone products used for electronic applications 770(1997)337
- Ion chromatographic characterization of toxic solutions: analysis and ion chemistry of biological liquids (Review) 774(1997)21
- Thin-layer chromatographic separation, colorimetric determination and recovery of thiocyanate from photogenic waste, river and sea waters 774(1997)373
- Prevention of peak overlap in the capillary electrophoretic separation of inorganic cations by use of a positively charged coated capillary 779(1997)347
- Micellar chromatography of inorganic compounds 780(1997)343
- Investigation of inorganic ions and organic acids in atmospheric aerosol by capillary electrophoresis 781(1997)223
- Electro-elution ion chromatography: controlling sample retention by electrochemically generating or modifying the mobile phase 782(1997)69
- Electrochemically regenerated ion suppressors, new suppressors for ion chromatography 782(1997)75
- Simultaneous determination of six cations in mineral water by single-column ion chromatography 782(1997)140
- Retention models in ion chromatography: the role of side equilibria in ion-exchange chromatography of inorganic cations and anions (Review) 789(1997)3
- Factors controlling ion-exchange selectivity in suppressed ion chromatography (Review) 789(1997)29
- Some examples of the use of amphoteric ion-exchange resins for inorganic separations 789(1997)157
- Separation and detection of vanadate, tungstate, molybdate and chromate ions 789(1997)267
- Separation and determination of inorganic germanium and β -carboxyethylgermanium sesquioxide by high-performance ion-exclusion chromatography 789(1997)403
- Liquid chromatographic separation of aqueous species of Cr(VI) and Cr(III) 789(1997)469
- Applications of column liquid chromatography to inorganic analysis in agricultural research (Review) 789(1997)519
- State-of-the-art ion chromatographic determination of inorganic ions in food 789(1997)529
- Separation and direct UV detection of sugars by capillary electrophoresis using chelation of copper(II) 799(1998)283
- Simultaneous separation of inorganic anions and cations by capillary zone electrophoresis 800(1998)369
- Inorganic ions**
see also Inorganic anions and Inorganic cations
- Isotachophoretic crystallization of ionogenic substances in electrolyte solutions 752(1996)251
- Validation of ion chromatographic methods for the trace analysis of ions in pharmacopoeial grades of water 770(1997)99
- Determination of inorganic ions in oil field waters by single-column ion chromatography 789(1997)569
- Inositol phosphates**
On-line electro dialysis–capillary zone electrophoresis of adenosine triphosphate and inositol phosphates 764(1997)135
- On-line electro dialysis–capillary zone electrophoresis–mass spectrometry of inositol phosphates in complex matrices 777(1997)51
- Three-compartment electro dialysis device for on-line sample clean-up and enrichment prior to capillary electrophoresis 788(1997)165
- Inositols**
Identification of phenolic acids and inositols in balms and tissues from an Egyptian mummy 767(1997)177
- Insulin**
Effect of skeleton size on the performance of octadecylsilylated continuous porous silica columns in reversed-phase liquid chromatography 762(1997)135
- Quantitation of insulin by capillary electrophoresis and high-performance liquid chromatography. Method comparison and validation 781(1997)445
- Theory of the correlation between capillary and free-flow zone electrophoresis and its use for the conversion of analytical capillary separations to continuous free-flow preparative processes. Application to analysis and preparation of fragments of insulin 796(1998)211
- Interleukin**
Selection of optimum affinity tags from a phage-displayed peptide library. Application to immobilized copper(II) affinity chromatography 787(1997)91
- Iodate**
Specific determination of bromate and iodate in ozonized water by ion chromatography with postcolumn derivatization and inductively-coupled plasma mass spectrometry 789(1997)259

Iodide

Simultaneous determination of bromide and iodide ions by capillary isotachopheresis using quaternary ammonium salts 760(1997)253

Iodine-azide

Iodine-azide reagent in detection of thiophosphoryl nucleotides in thin-layer chromatography systems 773(1997)285

Iopamidol

Effect of pH and ions in the sample on stacking in capillary electrophoresis 781(1997)35

Iothalamic acid

Effect of pH and ions in the sample on stacking in capillary electrophoresis 781(1997)35

Irgarol 1051

Ultra-trace-level determination of the antifouling agent Irgarol 1051 by gas chromatography with tandem mass spectrometric detection 766(1997)153

Iron

Trace determination of iron in water at the $\mu\text{g/l}$ level by on-line coupling of capillary isotachopheresis and capillary zone electrophoresis with UV detection of the EDTA-Fe(III) complex 757(1997)297

Chromatographic determination of commercial Fe(III) chelates of ethylenediaminetetraacetic acid, ethylenediaminedi(*o*-hydroxyphenylacetic) acid and ethylenediaminedi(*o*-hydroxy-*p*-methylphenylacetic) acid 789(1997)453

Normal-phase and reversed-phase high-performance liquid chromatographic behaviour and particle beam mass spectrometric characterization of alkyne-substituted iron-carbonyl clusters 789(1997)461

Electrophoretic studies on mixed complexes. Metal-methionine-penicillamine system 793(1998)220

Dye-ligand and metal chelate poly(2-hydroxyethylmethacrylate) membranes for affinity separation of proteins 799(1998)83

Reversed-phase high-performance liquid chromatography of iron(II) and copper(II) chelates with 4,7-diphenyl-1,10-phenanthroline disulfonate 799(1998)171

Isoboldine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Isobutoxycarbonyl methyl esters

Rapid and simultaneous analysis of protein and non-protein amino acids as N(O,S)-isobutoxycarbonyl methyl ester derivatives by capillary gas chromatography 758(1997)167

Isobutylchloroformate

Gas chromatographic-mass spectrometric determination of 4-(5) methylimidazole in ammonia caramel colour using ion-pair extraction and derivatization with isobutylchloroformate 786(1997)299

Isocitric acid

Gas chromatographic determination of organic acids from fruit juices by combined resin mediated methylation and extraction in supercritical carbon dioxide 785(1997)251

Isocorydine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Isoflurane

Approach to the thermodynamics of enantiomer separation by gas chromatography. Enantioselectivity between the chiral inhalation anesthetics enflurane, isoflurane and desflurane and a diluted γ -cyclodextrin derivative 757(1997)119

Preparative enantiomer separation of the inhalation anesthetics enflurane, isoflurane and desflurane by gas chromatography on a derivatized γ -cyclodextrin stationary phase 769(1997)119

Isoinosine

Analysis of traditional Chinese anticancer drugs by capillary electrophoresis 795(1998)388

Isoindoles

Comparison of the stability and UV and fluorescence characteristics of the *o*-phthalaldehyde/3-mercaptopropionic acid and *o*-phthalaldehyde/N-acetyl-L-cysteine reagents and those of their amino acid derivatives 798(1998)37

Isophthalic acid polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. II. Solubility effects in the analysis of polyester resins under reversed-phase conditions 761(1997)147

Isopropyl myristate

Analysis of impurities of isopropyl myristate by gas-liquid chromatography 767(1997)330

Isoquinolines

Inverse gas chromatography in the examination of organic compounds. Polarity and solubility parameters of isoquinoline derivatives 795(1998)349

Isetrandrine

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Isothiocyanates

Derivatization of pamidronate and other amino(bis)phosphonates with different isothiocyanates prior to ion-pair liquid chromatography 782(1997)211

Itaconic acid

High-speed separation of carboxylic acids by co-electroosmotic capillary electrophoresis with direct and indirect UV detection 758(1997)263

K**Kanamycin sulfate**

Analysis of kanamycin sulfate by liquid chromatography with pulsed electrochemical detection 766(1997)133

Kerogens

Programmed-temperature vaporiser injector as a new analytical tool for combined thermal desorption–pyrolysis of solid samples. Application to geochemical analysis 764(1997)73

Ketoamines

Capillary gas chromatographic determination of vanadium in crude petroleum oil using fluorinated ketoamines as derivatizing reagents 766(1997)159

Ketones

see also Carbonyl compounds

Separation of phenolic aldehydes, ketones and acids from lignin degradation by capillary zone electrophoresis 755(1996)89

Impinger sampling coupled to high-performance liquid chromatography by a modified autoinjector interface 762(1997)187

Glucose metabolism and bioreduction of 2-butanone by *Candida utilis* studied by means of ion-exchange chromatography 763(1997)205

Determination of diffusion coefficients and separation numbers in micellar electrokinetic chromatography 766(1997)187

Silicalite as a sorbent for solid-phase extraction 771(1997)45

Polyallylamine-supported pseudo-stationary phases for electrokinetic chromatography. Effect of alkyl chain length of the pseudo-stationary phase and methanol content of aqueous buffer on the separation of hydrophobic compounds 781(1997)139

Non-porous silica for ultrafast reversed-phase high-performance liquid chromatographic separation of aldehyde and ketone 2,4-dinitrophenylhydrazones 786(1997)47

Ketoprofen

Separation of profen enantiomers by capillary electrophoresis using cyclodextrins as chiral selectors 793(1998)165

Chiral and nonchiral determination of ketoprofen in pharmaceuticals by capillary zone electrophoresis 799(1998)301

Khellin

Determination of furanochromones and pyranocoumarins in drugs and *Ammi visnaga* fruits by combined solid-phase extraction–high-performance liquid chromatography and thin-layer chromatography–high-performance liquid chromatography 797(1998)305

Kynurenic acid

Determination of kynurenic acid by capillary electrophoresis with laser-induced fluorescence detection 781(1997)81

L**L-749,345**

Residual solvents determination in the antibiotic L-749,345 by static headspace gas chromatography 800(1998)275

Labdanes

Separation of forskolin derivatives by automated multiple development high-performance thin-layer chromatography 791(1997)231

Lactalbumins

Comparison of anion-exchange and hydroxyapatite displacement chromatography for the isolation of whey proteins 760(1997)125

Assessment of heat-induced denaturation of whey proteins 772(1997)231

Lactams

Optimisation of a simultaneous separation of sulphonamides, dihydrofolate reductase inhibitors and β -lactam antibiotics by capillary electrophoresis 768(1997)97

Lactoferrin

Bovine whey fractionation based on cation-exchange chromatography 795(1998)277

Lactoglobulins

Comparison of anion-exchange and hydroxyapatite displacement chromatography for the isolation of whey proteins 760(1997)125

Complete structure elucidation of a globular protein by particle beam liquid chromatography–Fourier transform infrared spectrometry and electrospray liquid chromatography–mass spectrometry. Sequence and conformation of β -lactoglobulin 763(1997)91

Assessment of heat-induced denaturation of whey proteins 772(1997)231

Multiple peaks in high-performance liquid chromatography of proteins. β -Lactoglobulins eluted in a hydrophobic interaction chromatography system 778(1997)43

Temperature programming and gradient elution in reversed-phase chromatography with packed capillary columns 788(1997)51

Lactones

Enantiodifferentiation of aryl-substituted heterocycles: a mechanistic study using γ -lactones 775(1997)81

Analysis of volatile components of oak wood by solvent extraction and direct thermal desorption–gas chromatography–mass spectrometry 778(1997)427

Packed column supercritical fluid chromatography of sesquiterpene lactones with different carbon skeletons 779(1997)315

Lanatosides

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Lansoprazole

Chiral resolution of pantoprazole sodium and related sulfoxides by complex formation with bovine serum albumin in capillary electrophoresis 759(1997)185

Lantadenes

Reversed-phase high-performance liquid chromatographic separation and quantification of lantadenes using isocratic systems 786(1997)181

Lanthanide complexes

New method to determine stability constants of lanthanide–EDTA complexes by isotachopheresis 786(1997)377

Lanthanides

Separation of samarium, gadolinium, terbium, dysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

Surfactant-sensitized post-column reaction with xylenol orange for the determination of lanthanides by ion chromatography 770(1997)75

Simultaneous separation of yttrium and lanthanide ions by isotachopheresis 772(1997)357

Investigation of adducts of lanthanide and uranium β -diketonates with organophosphorus Lewis bases by supercritical fluid chromatography 793(1998)107

Lauroilsine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Laurotetanine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Lead

Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217

Speciation of trimethyllead in rainwater by gas chromatography–mass spectrometry after ethylation with sodium tetraethylborate. Comparison with other alkylation methods 779(1997)299

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

High-performance chelation ion chromatography for the determination of traces of bismuth in lead by means of a novel hypercrosslinked polystyrene resin 789(1997)389

Cibacron Blue F3GA-attached polyvinylbutyral microbeads as novel magnetic sorbents for removal of Cu(II), Cd(II) and Pb(II) ions 793(1998)47

Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385

On-line pretreatment and determination of Pb, Cu and Cd at the $\mu\text{g l}^{-1}$ level in drinking water by chelation ion chromatography 800(1998)247

Lecithin

Investigation of the interactions between drugs and mixed bile salt/lecithin micelles. A characterization by micellar affinity capillary electrophoresis (MACE). Part III 781(1997)377

Leucine enkephalin

Gradient high-performance liquid chromatographic analysis of enkephalin peptides, their metabolites and enzyme inhibitors using combined ultraviolet and electrochemical detection. I. Optimization of separation and detection 766(1997)85

Leucine enkephalinamide

Gradient high-performance liquid chromatographic analysis of enkephalin peptides, their metabolites and enzyme inhibitors using combined ultraviolet and electrochemical detection. I. Optimization of separation and detection 766(1997)85

Leukotriene antagonists

Separation and quantification of geometric isomers in a leukotriene antagonist using cyclodextrins in capillary electrophoresis. Comparison to high-performance liquid chromatographic methods 768(1997)81

Leukotrienes

General equation for calculating the dissociation constants of polyprotic acids and bases from measured retention factors in high-performance liquid chromatography 762(1997)63

Use of secondary equilibria in reversed-phase high-performance liquid chromatography for the determination of dissociation constants of polyprotic leukotrienes 763(1997)199

Identification of eicosanoids in the red algae, *Gracilaria asiatica*, using high-performance liquid chromatography and electrospray ionization mass spectrometry 795(1998)227

Lidocaine

On-line column-switching high-performance liquid chromatography analysis of cardiovascular drugs in serum with automated sample clean-up and zone-cutting technique to perform chiral separation 762(1997)235

Lignans

Analysis of lignan constituents from *Schisandra chinensis* by liquid chromatography–electrospray mass spectrometry 757(1997)81

neo-Lignans

Determination of magnolol and honokiol in Magnoliae Cortex using supercritical fluid chromatography on-line coupled with supercritical fluid extraction by on-column trapping 786(1997)366

Lignin

Separation of phenolic aldehydes, ketones and acids from lignin degradation by capillary zone electrophoresis 755(1996)89

Size-exclusion chromatography of lignin as ion-pair complex 764(1997)183

Absolute quantitation of lignin pyrolysis products using an internal standard 773(1997)227

Lignosulfonic acids

Size-exclusion chromatography of lignin as ion-pair complex 764(1997)183

Limonin D-ring lactone hydrolase

High-performance liquid chromatography quantitation of limonin D-ring lactone hydrolase and limonoate dehydrogenase activities 760(1997)173

Limonoate dehydrogenase

High-performance liquid chromatography quantitation of limonin D-ring lactone hydrolase and limonoate dehydrogenase activities 760(1997)173

Limonoids

High-performance liquid chromatography quantitation of limonin D-ring lactone hydrolase and limonoate dehydrogenase activities 760(1997)173

Lindane

Analysis of lindane, α - and β -endosulfan and endosulfan sulfate in greenhouse air by gas chromatography 765(1997)99

Lindcarpine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Linoleic acid

Determination of primary oxidation products of linoleic acid and triacylglycerols 767(1997)63

Fractionation of *cis*- and *trans*-oleic, linoleic, and conjugated linoleic fatty acid methyl esters by silver ion high-performance liquid chromatography 799(1998)329

Linuron

Determination of linuron in aqueous soil extracts by high-performance liquid chromatography 754(1996)295

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Lipases

New class of poly(vinyl alcohol) polymers as column-chromatography stationary phases for *Candida rugosa* lipase isoforms separation 753(1996)47

Preliminary study on hydrophobic interaction chromatography of *Chromobacterium viscosum* lipase on polypropylene glycol immobilized on Sepharose 796(1998)177

Lipids

Cell lipids of the *Candida lipolytica* yeast grown on methanol 755(1996)75

Triacylglycerol mixture for testing capillary columns for high-temperature gas chromatography 758(1997)235

Rapid procedure for the isolation and analysis of fatty acid and fatty alcohol fractions from wax esters of marine zooplankton 760(1997)309

Normal-phase separation effects with lipids on a silver ion high-performance liquid chromatography column 764(1997)337

Separation and quantitation of glycolipids as penetration modifiers in human skin using high-performance liquid chromatography–mass spectrometry with electrospray ionization 766(1997)71

Analysis of a monophosphoryl lipid A immunostimulant preparation from *Salmonella minnesota* R595 by high-performance liquid chromatography 767(1997)53

Gas-liquid chromatography-mass spectrometry of primary and secondary fatty alcohols and diols as their *tert.*-butyldimethylsilyl derivatives 771(1997)191

Reversed-phase separations of nitrogenous phospholipids on an octadecanoyl poly(vinyl alcohol) phase 773(1997)93

Quantitative aspects of triglyceride analysis by gas-liquid chromatography using a short metal capillary column 773(1997)233

Characterisation of aldehydic acids in used and unused frying oils 776(1997)245

Distribution of the total unsaturation in lipid components of plasma as a new differential diagnostic method in clinical analysis 777(1997)47

Analysis of gangliosides by capillary zone electrophoresis and capillary zone electrophoresis-electrospray mass spectrometry 779(1997)195

Semi-continuous extraction/purification of lipids by means of supercritical fluids 785(1997)337

Development of a supercritical fluid extraction method for determination of lipid classes and total fat in meats and its comparison with conventional methods 785(1997)345

Analysis of cyanolipids and triacylglycerols from sapindaceae seed oils with high-temperature gas chromatography and high-temperature gas chromatography-chemical ionization mass spectrometry 787(1997)181

Gas-liquid chromatography-mass spectrometry of hydroxy fatty acids as their methyl esters *tert.*-butyldimethylsilyl ethers 793(1998)91

Overview of the applications of liquid chromatography-mass spectrometry interfacing systems in food analysis: naturally occurring substances in food 794(1998)263

Lipopeptides

Practical capillary electrophoresis method for the quantitation of the acetate counter-ion in a novel antifungal lipopeptide 763(1997)279

Lipopolysaccharide

High-performance liquid chromatographic peak identification of 2,4-dinitrophenylhydrazine derivatives of lipid peroxidation aldehydes by photodiode array detection 798(1998)117

Lipoproteins

Rapid reversed-phase high-performance liquid chromatography method for quantitation, at high pH, of the recombinant apolipo-protein A-IMilano in *Escherichia coli* fermentation broth 798(1998)83

Lipoxygenases

Enantiomeric separation of various lipoxygenase derived monohydroxy polyunsaturated fatty acid methyl esters by high-performance liquid chromatography 790(1997)235

Liquid crystals

Letter to the editor 771(1997)385

Lolitrein B

Rapid determination of the mycotoxin lolitrein B in endophyte-infected perennial ryegrass by high-performance thin-layer chromatography. A validated assay 769(1997)343

Lorazepam

On-column enantiomerization of 3-hydroxybenzodiazepines during chiral liquid chromatography with optical rotation detection 767(1997)93

Erratum to "On-column enantiomerization of 3-hydroxybenzodiazepines during chiral liquid chromatography with optical rotation detection". [J. Chromatogr. A, 767 (1997) 93-100] 782(1997)296

Bonded cellulose-derived high-performance liquid chromatography chiral stationary phases. III. Effect of the reticulation of the cellulose derivative on performance 791(1997)37

Lupulones

Preparative separation of bitter acids from hop extracts by centrifugal partition chromatography 771(1997)71

Lutein

Use of high-precision gas isotope ratio mass spectrometry to determine natural abundance ^{13}C in lutein isolated from C_3 and C_4 plant sources 800(1998)51

2,6-Lutidine

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

LY213829

Chiral separation of neutral species by capillary electrophoresis. Evaluation of a theoretical model 761(1997)249

LY297802

Investigation of response factor ruggedness for the determination of drug impurities using high-performance liquid chromatography with ultraviolet detection 762(1997)227

Lymphotoxin

Role of high-performance liquid chromatographic protein analysis in developing fermentation processes for recombinant human growth hormone, relaxin, antibody fragments and lymphotoxin 763(1997)31

Lysergic acid diethylamide

Application of micellar electrokinetic capillary chromatography to the analysis of illicit drug seizures 780(1997)265

Lysine

Determination of available lysine in infant milk formulae by high-performance liquid chromatography 778(1997)235

Lysozyme

Direct purification of lysozyme using continuous counter-current expanded bed adsorption 757(1997)41

Synergistic effects in the processes of protein multicomponent sorption 760(1997)105

Comparative studies on the isothermal characteristics of proteins adsorbed under batch equilibrium conditions to ion-exchange, immobilised metal ion affinity and dye affinity matrices with different ionic strength and temperature conditions 763(1997)71

Studies of the hydrodynamic volume changes that occur during refolding of lysozyme using size-exclusion chromatography 766(1997)109

Productivity and operating regimes in protein chromatography using low-molecular-mass displacers 771(1997)9

M**Macrocyclic compounds**

Supercritical fluid extraction: metals as complexes (Review) 785(1997)369

Magnesium

Determination of Na⁺, K⁺, Mg²⁺ and Ca²⁺ in mixtures of seawater and formation water by capillary electrophoresis 767(1997)311

Simple recognition of similar samples for the ion-chromatographic determination of the main cations 789(1997)323

Determination of calcium and magnesium in water samples by high-performance liquid chromatography on a graphitic stationary phase with a mobile phase containing *o*-cresolphthalein complexone 789(1997)329

Magnolol

Determination of magnolol and honokiol in Magnoliae Cortex using supercritical fluid chromatography on-line coupled with supercritical fluid extraction by on-column trapping 786(1997)366

Malathion

Indirect liquid chromatographic determination of malathion in formulations, based on the formation of palladium(II)-dimethyldithiosphate complex 758(1997)159

Solid-phase microextraction of organophosphorus pesticides from water 759(1997)225

Maleic acid polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions 775(1997)157

Maleic anhydride

Composition and structure analysis of styrene-maleic anhydride copolymer by pyrolysis-gas chromatography 765(1997)279

Malic acid

Gas chromatographic determination of organic acids from fruit juices by combined resin mediated methylation and extraction in supercritical carbon dioxide 785(1997)251

Malonaldehyde

High-performance liquid chromatographic peak identification of 2,4-dinitrophenylhydrazine derivatives of lipid peroxidation aldehydes by photodiode array detection 798(1998)117

Maltose

Analysis of selected alkaloids and sugars in tobacco extract 775(1997)101

Mandelic acid

Reversed determination of the formation constants of 1-allyl terguride with mandelic acid optical isomers using capillary electrophoresis 772(1997)179

Manganese dioxide

Development of a manganese dioxide solid-phase reactor for oxidation of toxins associated with paralytic shellfish poisoning 755(1996)227

Maytansinoids

Normal- and bonded-phase liquid chromatography with photodiode array detection of maytansinoids 763(1997)213

MCPA

Determination of chlorophenoxy herbicides in waters by capillary gas chromatography with ion trap detection 758(1997)324

Mecoprop

Simultaneous determination and enantiomeric resolution of mecoprop and dichlorprop in soil samples by high-performance liquid chromatography and gas chromatography–mass spectrometry 799(1998)355

Medetomidine

Analytical method development for the simultaneous quantitation of dexmedetomidine and three potential metabolites in plasma 762(1997)281

Melamine

pH-Dependent elution behaviour of meleme in high-performance cation-exchange chromatography 752(1996)287

Meleme

pH-Dependent elution behaviour of meleme in high-performance cation-exchange chromatography 752(1996)287

Membrane proteins

see also Proteins

Rapid resolution by reversed-phase high-performance liquid chromatography of the thylakoid membrane proteins of the photosystem II light-harvesting complex 779(1997)131

Mepivacaine

Capillary electrochromatography with molecular imprint-based selectivity for enantiomer separation of local anaesthetics 792(1997)401

Mercaptan

Gas chromatographic–mass spectrometric analysis of mercaptan odorants in liquefied petroleum gas and liquefied natural gas 791(1997)367

Mercaptoalkyl acids

High-density ligand attachment to brominated allyl matrices and application to mixed mode chromatography of chymosin 775(1997)39

Mercaptopropionic acid

Comparison of the stability and UV and fluorescence characteristics of the *o*-phthaldialdehyde/3-mercaptopropionic acid and *o*-phthaldialdehyde/*N*-acetyl-L-cysteine reagents and those of their amino acid derivatives 798(1998)37

Mercury

Reversed-phase high-performance liquid chromatography of pyrrolidinedithiocarbamate complexes of mercuric species using amperometric and coulometric detection 761(1997)139

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385

Mercury compounds

Development of a capillary electrophoretic method for the separation and determination of trace inorganic and organomercury species utilizing the formation of highly absorbing water soluble dithizone sulphonate complexes 765(1997)345

Metal cations

see also Cations and Inorganic cations

Ion-exchange properties of glutamic acid-bonded silica 769(1997)179

Modification of 4-(2-pyridylazo)-resorcinol postcolumn reagent selectivity through competitive equilibria with chelating ligands 770(1997)69

New fluorinated hydroxamic acid reagents for the extraction of metal ions with supercritical CO₂ 770(1997)85

Use of ion chromatography for the determination of transition metals in the control of sewage-treatment-plant and related waters 770(1997)185

Retention behavior of alkali, alkaline earth and transition metal cations in ion chromatography with an unmodified silica gel column 770(1997)219

Electroinjection analysis. Concept, mathematical model and applications 772(1997)115

Separation of metal ions and metal-containing species by micellar electrokinetic capillary chromatography, including utilisation of metal ions in separations of other species 780(1997)329

Sulfoacylated macroporous polystyrene–divinylbenzene: a new type of cation exchanger for the analysis of multivalent metal cations 782(1997)149

Multi-element trace analysis by reversed-phase high-performance liquid chromatography followed by on-line column enrichment as 2-(5-nitro-2-pyridylazo)-5-[*N*-propyl-*N*-(3-sulphopropyl)amino] phenol chelates 788(1997)113

Advances in detection techniques for ion chromatography (Review) 789(1997)67

Histidine as a dipolar eluent component in cation chromatography 789(1997)141

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

Determination of calcium and magnesium in water samples by high-performance liquid chromatography on a graphitic stationary phase with a mobile phase containing *o*-cresolphthalein complexone 789(1997)329

- High-performance chelation ion chromatography. A new dimension in the separation and determination of trace metals (Review) 789(1997)413
- Determination of some inorganic species in edible vegetable oils and fats by ion chromatography 789(1997)549
- Ion chromatography as reference method for serum cations (Review) 789(1997)557
- Simultaneous speciation of aqueous selenium(IV) and selenium(VI) by high-performance liquid chromatography with ultraviolet detection 791(1997)357
- Strategies for selectivity control in capillary electrophoresis of metal species 792(1997)495
- Cibacron Blue F3GA-attached polyvinylbutyral microbeads as novel magnetic sorbents for removal of Cu(II), Cd(II) and Pb(II) ions 793(1998)47
- Role of ligand purity in separations of alkaline earth metals as arsenazo I complexes by capillary zone electrophoresis 793(1998)177
- Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385
- Metal chelates**
- Effect of on-capillary large volume sample stacking on limits of detection in the capillary zone electrophoretic determination of selected drugs, dyes and metal chelates 772(1997)161
- Influence of cationic polymers on separation selectivity in kinetic differentiation mode capillary electrophoresis of metal-4-(2-pyridylazo)resorcinolato chelates 772(1997)339
- Supercritical fluid extraction of metal ions and metal chelates from different environments (Review) 774(1997)37
- Molecularly imprinted ligand-exchange adsorbents for the chiral separation of underivatized amino acids 775(1997)51
- Use of cationic polymers for the simultaneous determination of inorganic anions and metal-4-(2-pyridylazo)resorcinolato chelates in kinetic differentiation-mode capillary electrophoresis 776(1997)329
- Micelle-mediated extraction 780(1997)229
- Multi-element trace analysis by reversed-phase high-performance liquid chromatography followed by on-line column enrichment as 2-(5-nitro-2-pyridylazo)-5-[N-propyl-N-(3-sulphopropyl)amino] phenol chelates 788(1997)113
- Determination and speciation of metals by liquid chromatography (Review) 789(1997)301
- Fluorometric determination of serum and urinary aluminium with 8-quinolinol by kinetic-differentiation-mode micellar chromatography 789(1997)361
- Selective fluorometric determination of indium(III) by high-performance liquid chromatography with 2-methyl-8-quinolinol based on a ligand-exchange reaction of silanol groups 789(1997)395
- Chromatographic determination of commercial Fe(III) chelates of ethylenediaminetetraacetic acid, ethylenediaminedi(*o*-hydroxyphenylacetic) acid and ethylenediaminedi(*o*-hydroxy-*p*-methylphenylacetic) acid 789(1997)453
- Ion-pair reversed-phase high-performance liquid chromatographic separation and determination of ruthenium, rhodium, cobalt and copper as chelates with 1-(2-pyridylazo)-2-naphthol-6-sulfonic acid 789(1997)485
- Dye-ligand and metal chelate poly(2-hydroxyethylmethacrylate) membranes for affinity separation of proteins 799(1998)83
- Reversed-phase high-performance liquid chromatography of iron(II) and copper(II) chelates with 4,7-diphenyl-1,10-phenanthroline disulfonate 799(1998)171
- Molecular characterization of metal-binding polypeptide domains by electrospray ionization mass spectrometry and metal chelate affinity chromatography 800(1998)29
- Metal complexes**
- Advanced possibilities on multi-element separation and detection of metal ions by capillary zone electrophoresis using precapillary complexation. I. Separation aspects 756(1996)300
- Optimisation of selectivity in the separation of metallo-cyanide complexes by ion-interaction liquid chromatography 770(1997)3
- Studies on the combination of ion chromatography-particle-beam mass spectrometry with capillary columns 770(1997)59
- Reversed-phase ion-interaction chromatography of Cu(I)-cyanide complexes 770(1997)165
- Determination of platinum, palladium, osmium, iridium, rhodium and gold as chloro complexes by capillary zone electrophoresis 772(1997)347
- Chemically bonded chelates as selective complexing sorbents for gas chromatography. VI. Modification of silica with NiCl₂ and CoCl₂ via β-diketonate groups 773(1997)209
- Capillary electrophoresis in metal analysis. Investigations of multi-elemental separation of metal chelates with aminopolycarboxylic acids 773(1997)351
- Chromatographic method involving inductively coupled plasma atomic emission spectrometric detection for the study of metal-protein complexes 776(1997)93

- Capillary electrophoretic determination of cyanide leaching solutions from automobile catalytic converters 778(1997)397
- Use of metal complexation in non-aqueous capillary electrophoresis systems for the separation and improved detection of tetracyclines 779(1997)235
- Separation of metal ions and metal-containing species by micellar electrokinetic capillary chromatography, including utilisation of metal ions in separations of other species 780(1997)329
- Determination of kynurenic acid by capillary electrophoresis with laser-induced fluorescence detection 781(1997)81
- Characterization of metal complexes of pharmaceutical interest by capillary electrophoresis with element sensitive detection 781(1997)407
- Liquid chromatographic separation of metal–tetraphenylporphine complexes based on coordination interaction with immobilized neutral ligand in stationary phase 786(1997)269
- Evaluation of capillary zone electrophoresis equilibrium data using the CELET program 786(1997)321
- New method to determine stability constants of lanthanide–EDTA complexes by isotachopheresis 786(1997)377
- Detection of glycosaminoglycans as a copper(II) complex in high-performance liquid chromatography 787(1997)266
- Retention models in ion chromatography: the role of side equilibria in ion-exchange chromatography of inorganic cations and anions (Review) 789(1997)3
- Chromatographic separation and characterization of the photoproduct of tris(L-cysteinesulfinate-N,S)cobaltate(III) 789(1997)273
- Recent applications of high-performance liquid chromatography to the analysis of metal complexes (Review) 789(1997)437
- Chromatographic behaviour of the chromium(III) complex of 1-phenyl-3-methyl-4-benzoyl-5-pyrazolone 789(1997)479
- High-performance liquid chromatographic separations of boron-cluster compounds (Review) 789(1997)497
- Strategies for selectivity control in capillary electrophoresis of metal species 792(1997)495
- Role of ligand purity in separations of alkaline earth metals as arsenazo I complexes by capillary zone electrophoresis 793(1998)177
- Electrophoretic studies on mixed complexes. Metal–methionine–penicillamine system 793(1998)220
- Comparison of prediction power between theoretical and neural-network models in ion–interaction chromatography 799(1998)35
- Metal cyanide complexes**
Determination of metal cyanide complexes in gold processing solutions by capillary electrophoresis 767(1997)319
- Optimisation of selectivity in the separation of metallo–cyanide complexes by ion–interaction liquid chromatography 770(1997)3
- Ion chromatographic analysis of cyanate in gold processing samples containing large concentrations of copper(I) and other metallo–cyanide complexes 770(1997)175
- Determination of metallo–cyanides by capillary electrophoresis after concentration on supported liquid membranes 770(1997)329
- Metaldehyde**
Identification of pesticide poisoning in wildlife 754(1996)463
- Metal ions**
Supercritical fluid extraction of metal ions and metal chelates from different environments (Review) 774(1997)37
- Simultaneous separation of metal ions and neutral organics by reversed-phase ion-pair liquid chromatography 776(1997)221
- Use of 2-quinolyazo compounds for the reversed-phase high-performance liquid chromatographic separation of metal ions 789(1997)491
- Metalloproteins**
Development of a high-performance capillary isoelectric focusing technique with application to studies of microheterogeneity in chicken conalbumin 757(1997)247
- Metallothioneins**
Application of solid-phase extraction to the preconcentration of metallothionein and metals from physiological fluids 766(1997)121
- Chromatographic and hyphenated methods for elemental speciation analysis in environmental media (Review) 774(1997)3
- Rapid assay for the analysis of class III metallothioneins by capillary zone electrophoresis in crude extracts of microalgal cells exposed to cadmium 775(1997)339
- Characterization of metal complexes of pharmaceutical interest by capillary electrophoresis with element sensitive detection 781(1997)407
- On-line solid-phase extraction–capillary electrophoresis for enhanced detection sensitivity and selectivity: application to the analysis of metallothionein isoforms in sheep fetal liver 792(1997)463

Metal oxides

Deposition parameters of air pollutants on solid surfaces, measured in the presence of surface and gaseous reactions, with a simultaneous determination of the experimental isotherms 775(1997)211

Metal oxoanions

Separation and detection of vanadate, tungstate, molybdate and chromate ions 789(1997)267

Metal phthalocyanines

Analysis of sulphonated phthalocyanine dyes by capillary electrophoresis 770(1997)345

Metals

see also Heavy metals and Transition metals

High-performance liquid chromatography–hydride generation–atomic fluorescence spectroscopic determination of arsenic species in water 756(1996)292

Advanced possibilities on multi-element separation and detection of metal ions by capillary zone electrophoresis using precapillary complexation. I. Separation aspects 756(1996)300

Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217

Selective determination of Sb(III) by gas chromatography–quartz furnace atomic absorption spectrometry after derivatization with triphenylmagnesium bromide 764(1997)169

Application of solid-phase extraction to the preconcentration of metallothionein and metals from physiological fluids 766(1997)121

Single-column method of chelation ion chromatography for the analysis of trace metals in complex samples 770(1997)129

Capillary electrophoresis in metal analysis. Investigations of multi-elemental separation of metal chelates with aminopolycarboxylic acids 773(1997)351

Chromatographic and hyphenated methods for elemental speciation analysis in environmental media (Review) 774(1997)3

Speciation of chromium from industrial wastes and incinerated sludges 774(1997)363

Speciation of platinum–chloro complexes and their hydrolysis products by ion chromatography. Determination of platinum oxidation states 775(1997)197

Capillary electrophoretic determination of cyanide leaching solutions from automobile catalytic converters 778(1997)397

Simultaneous determination of toxic arsenic and chromium species in water samples by ion chromatography–inductively coupled plasma mass spectrometry 779(1997)139

Supercritical fluid extraction: metals as complexes (Review) 785(1997)369

Ion chromatographic separation of metallic ions on a bonded poly- β -diketone-type non-cyclic crown ether stationary phase 789(1997)135

High-performance chelation ion chromatography for the determination of traces of bismuth in lead by means of a novel hypercrosslinked polystyrene resin 789(1997)389

Normal-phase and reversed-phase high-performance liquid chromatographic behaviour and particle beam mass spectrometric characterization of alkyne-substituted iron–carbonyl clusters 789(1997)461

Methamphetamines

Determination of methamphetamine and related compounds by capillary electrophoresis with UV and laser-induced fluorescence detection 798(1998)325

Methanol

Cell lipids of the *Candida lipolytica* yeast grown on methanol 755(1996)75

Chemiluminescence nitrogen detection for packed-column supercritical fluid chromatography with methanol modified carbon dioxide 757(1997)183

Trace determination of methanol in water–ethanol solution by derivatization and high-performance liquid chromatography 799(1998)93

Methionine

Electrophoretic studies on mixed complexes. Metal–methionine–penicillamine system 793(1998)220

Methionine enkephalin

Optical resolution of phenylthiohydantoin-amino acids by capillary electrophoresis and identification of the phenylthiohydantoin-D-amino acid residue of [D-Ala²]-methionine enkephalin 752(1996)279

Gradient high-performance liquid chromatographic analysis of enkephalin peptides, their metabolites and enzyme inhibitors using combined ultraviolet and electrochemical detection. I. Optimization of separation and detection 766(1997)85

Methionine enkephalinamide

Gradient high-performance liquid chromatographic analysis of enkephalin peptides, their metabolites and enzyme inhibitors using combined ultraviolet and electrochemical detection. I. Optimization of separation and detection 766(1997)85

N-Methyl-1-(1,3-benzodioxol-5-yl)-2-butanamine

Sensitive and selective determination of methylenedioxyethylated amphetamines by high-performance liquid chromatography with fluorimetric detection 787(1997)137

Methyl ADC-13 enolphosphate diphenyl ester

High-performance liquid chromatographic separation of β -methyl ADC-13 enolphosphate diphenyl ester and its α -methyl diastereomer 762(1997)327

Methylamine

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

Methylascorbigen

Comparison of thin-layer chromatography and overpressured layer chromatographic techniques for the separation of ascorbigen and 1'-methylascorbigen 764(1997)103

Methylaspartate receptor blockers

Determination of several N-methyl-D-aspartate receptor blockers in plasma and brain by a selective high-performance liquid chromatographic method with column switching 797(1998)211

Methylbenzoate

Systematic approach to links between separations in capillary electrophoresis and liquid chromatography. IV. Application of binding constant-retention factor relationship to the separation of 2-, 3- and 4-methylbenzoate anions using β -cyclodextrin as selector 768(1997)29

3-Methylbutamine

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

Methylcarbamates

Recent advances in the residue analysis of N-methylcarbamate pesticides (Review) 754(1996)3

Application of liquid chromatography-atmospheric-pressure chemical-ionization mass spectrometry to pesticide analysis 754(1996)61

Development of a new high-performance liquid chromatography method to analyse N-methylcarbamate insecticides by a simple post-column derivatization system and fluorescence detection 778(1997)103

Methyl cyclohexane

Methyl cyclohexane as a new eluting solvent for the size-exclusion chromatography of polyethylene and polypropylene at 90°C 755(1996)27

24-Methylen-24-dihydrolanosterol

Isolation of three triterpenes and several aliphatic alcohols from *Crataegus monogyna* Jacq 767(1997)340

Methylenebisacrylamide

Rigid porous polyacrylamide-based monolithic columns containing butyl methacrylate as a separation medium for the rapid hydrophobic interaction chromatography of proteins 775(1997)65

Methylenedioxyamphetamine

Sensitive and selective determination of methylenedioxyethylated amphetamines by high-performance liquid chromatography with fluorimetric detection 787(1997)137

Methylenedioxyethamphetamine

Sensitive and selective determination of methylenedioxyethylated amphetamines by high-performance liquid chromatography with fluorimetric detection 787(1997)137

Methylenedioxymethamphetamine

Sensitive and selective determination of methylenedioxyethylated amphetamines by high-performance liquid chromatography with fluorimetric detection 787(1997)137

Methylephedrine

Ruggedness of enantiomeric separation by capillary electrophoresis and high-performance liquid chromatography with methylated cyclodextrins as chiral selectors 763(1997)139

4-(5-Methylimidazole

Gas chromatographic-mass spectrometric determination of 4-(5-methylimidazole in ammonia caramel colour using ion-pair extraction and derivatization with isobutylchloroformate 786(1997)299

N-Methylauroretanine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

N-Methylindocarpine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Methylmercury

Capillary electrophoresis determination of methylmercury in fish and crab meat after extraction as the dithione sulphonate complex 791(1997)333

Methylnaphthalene

Optimization of the high-performance liquid chromatographic separation of fullerenes using 1-methylnaphthalene as the mobile phase on a tetraphenylporphyrin-silica stationary phase 786(1997)31

Methyl palmitate

Quantitative evaluation of carbon isotopic fractionation during reversed-phase high-performance liquid chromatography 757(1997)307

Methylphenols

Phenol and methylphenol isomers determination in soils by in-situ microwave-assisted extraction and derivatisation 757(1997)153

Methylphosphonic acid monoesters

Capillary electrophoresis separation of alkylphosphonic acid monoesters with indirect ultraviolet detection 779(1997)245

Methylphosphonic acids

Analysis of methylphosphonic acid, ethyl methylphosphonic acid and isopropyl methylphosphonic acid at low microgram per liter levels in groundwater 790(1997)143

Methylpyrrolidinone

N-Methyl-2-pyrrolidinone as a mobile phase in the size-exclusion chromatography of coal derivatives 758(1997)65

2-Methylquinoline

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Methylquinolinol

Selective fluorometric determination of indium(III) by high-performance liquid chromatography with 2-methyl-8-quinolinol based on a ligand-exchange reaction of silanol groups 789(1997)395

Metobromuron

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Metolachlor

Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers 792(1997)419

Metoprolol

Robustness of the chromatographic separation of alprenolol and related substances using a silica-based stationary phase and selective retention of metoprolol and related substances on a porous graphitic carbon stationary phase 797(1998)75

Mevinolin

Monitoring of selected metabolites and biotransformation products from fermentation broths by high-performance liquid chromatography 763(1997)165

Mexiletine hydrochloride

Chiral separation of racemic mexiletine hydrochloride using cyclodextrins as chiral additive by capillary electrophoresis 795(1998)394

Miconazole

Liquid chromatographic separation of the enantiomers of beclonazole and its potential impurities 769(1997)231

Microcystins

Effect of foaming power and foam stability on continuous concentration with foam counter-current chromatography 791(1997)53

Determination of some physicochemical parameters of microcystins (cyanobacterial toxins) and trace level analysis in environmental samples using liquid chromatography 799(1998)155

Mirtazapin

Development and validation of a capillary electrophoresis method within a pharmaceutical quality control environment and comparison with high-performance liquid chromatography 773(1997)339

MK-0677

Enantiomeric separation of the novel growth hormone secretagogue MK-0677 by capillary zone electrophoresis 773(1997)311

Molybdate

New approach to determining each component of a two-component overlapping peak by single-column anion ion chromatography 769(1997)333

Monobromobimane

Process clearance of dithiothreitol monitored by reversed-phase high-performance liquid chromatography with fluorescence detection. Assay development and validation 775(1997)151

Monoclonal antibodies

see also Antibodies

Isolation of monoclonal antibodies from cell containing hybridoma broth using a protein A coated adsorbent in expanded beds 752(1996)111

Use of monoclonal antibodies for weak affinity chromatography 758(1997)199

Purification of monoclonal antibodies by simulated moving-bed chromatography 765(1997)201

Purification of *p*-nitrobenzyl C-functionalized diethylenetriamine pentaacetic acids for clinical applications using anion-exchange chromatography 771(1997)63

Kinetic analysis of the interaction between the monoclonal antibody A33 and its colonic epithelial antigen by the use of an optical biosensor. A comparison of immobilisation strategies 776(1997)15

Purification of monoclonal antibodies by epitope and mimotope affinity chromatography 782(1997)49

On-line post-capillary affinity detection of immunoglobulin G subclasses and monoclonal antibody variants for capillary electrophoresis 790(1997)207

Rapid high-performance isoelectric focusing of monoclonal antibodies in uncoated fused-silica capillaries 790(1997)215

Development of a downstream process for the isolation and separation of monoclonal immunoglobulin A monomers, dimers and polymers from cell culture supernatant 796(1998)165

Validation of a capillary isoelectric focusing method for the recombinant monoclonal antibody C2B8 800(1998)355

Monocyclic aromatic hydrocarbons

see also Hydrocarbons, aromatic

Determination of partition coefficients of monocyclic aromatic hydrocarbons between leaf essential oil and air by headspace gas chromatography 765(1997)247

Headspace gas chromatographic determination of the plant cuticle-air partition coefficients for monocyclic aromatic hydrocarbons as environmental compartment 774(1997)213

Monofluorophosphate

Simultaneous determination of monofluorophosphate and fluoride in toothpaste by capillary electrophoresis 765(1997)353

Development of a capillary electrophoresis method with indirect photometric detection for the determination of anions related to anticaries 781(1997)457

Monohydroperoxides

Determination of primary oxidation products of linoleic acid and triacylglycerols 767(1997)63

Monohydroxides

Determination of primary oxidation products of linoleic acid and triacylglycerols 767(1997)63

Monomethylarsonic acid

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. I. Large-volume stacking with polarity switching 757(1997)281

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. II. Large-volume stacking without polarity switching 757(1997)291

Monophenylarsonic acid

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. II. Large-volume stacking without polarity switching 757(1997)291

Monophosphoryl lipid A

Analysis of a monophosphoryl lipid A immunostimulant preparation from *Salmonella minnesota* R595 by high-performance liquid chromatography 767(1997)53

Monosaccharides

see also Saccharides

Analysis of monosaccharide composition by capillary electrophoresis 763(1997)271

Separation of acyclic derivatives of sugar enantiomers using gas chromatography on chiral stationary phases of permethylated α -, β - and γ -cyclodextrin 767(1997)325

Determination of the absolute configuration of sugar residues using gas chromatography. Method with potential for elimination of references 769(1997)253

Subcritical fluid chromatography of monosaccharides and polyols using silica and trimethylsilyl columns 785(1997)195

Monoterpenes

Chiral separation of monoterpenes using mixtures of sulfated β -cyclodextrins and α -cyclodextrin as chiral additives in the reversed-polarity capillary electrophoresis mode 759(1997)149

Quantification of carvone, cineole, perillaldehyde, perillyl alcohol and sobrerol by isocratic high-performance liquid chromatography 793(1998)71

Monuron

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography-Fourier-transform infrared spectrometry 756(1996)145

Morphine

Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67

Determination of morphine and its analogues in urine by on-line coupled reversed-phase liquid chromatography-gas chromatography with on-line derivatization 771(1997)360

Determination of morphine in urine by solid-phase immunoextraction and high-performance liquid chromatography with electrochemical detection 797(1998)245

Muricholic acids

Capillary gas-liquid chromatography of acetate-methyl esters of bile acids 766(1997)286

MX

New derivatization method for determination of 3-chloro-4(dichloromethyl)-5-hydroxy-2(5H)-furanone in water 790(1997)242

Mycotoxins

see also Toxins

Gas chromatography-mass spectrometry of *Alternaria* mycotoxins 765(1997)255

Rapid determination of the mycotoxin lolitrem B in endophyte-infected perennial ryegrass by high-performance thin-layer chromatography. A validated assay 769(1997)343

Determination of mycotoxins produced by *Fusarium* isolates from banana fruits by capillary gas chromatography and high-performance liquid chromatography 778(1997)363

Determination of trichothecene mycotoxins in wheat by use of supercritical fluid extraction and high-performance liquid chromatography with diode array detection or gas chromatography with electron capture detection 795(1998)297

Myotoxins

see also Toxins

Detection of isoforms and isomers of rattlesnake myotoxins by capillary electrophoresis and matrix-assisted laser desorption time-of-flight mass spectrometry 781(1997)429

N

Naphthalene-2,3-dicarboxaldehyde

Enhanced temporal resolution for the microdialysis monitoring of catecholamines and excitatory amino acids using capillary electrophoresis with laser-induced fluorescence detection. Analytical developments and in vitro validations 755(1996)99

Naphthalenes

Separation of naphthalene and flavone derivatives by micellar electrokinetic chromatography with double- and triple-chain surfactants 755(1996)147

Application of particles as pseudo-stationary phases in electrokinetic chromatography 780(1997)63

Comparison of 1,1,1,2-tetrafluoroethane and carbon dioxide-based mobile phases for packed column supercritical fluid chromatography 782(1997)237

Resolution studies on counter-current chromatography using supercritical fluid carbon dioxide 790(1997)31

On-line micellar electrokinetic chromatography–mass spectrometry with a high-molecular-mass surfactant 794(1998)317

Naphthalenesulfonates

Wavelength-resolved laser-induced fluorescence detection in capillary electrophoresis: naphthalenesulphonates in river water 771(1997)331

Naphthalenesulfonic acids

Effects of the working electrolyte (cyclodextrin type and pH) on the separation of aromatic sulphonic acids by capillary zone electrophoresis 772(1997)385

Naphthalenetrisulfonate

Sulfur speciation by capillary electrophoresis with indirect spectrophotometric detection: In search of a suitable carrier electrolyte to maximize sensitivity 770(1997)311

Naphthaquinones

Quantitative determination of naphthaquinones of *Arnebia densiflora* (Nordm.) Ledeb. by an improved high-performance liquid chromatographic method 782(1997)133

Naphthols

On-line concentration of neutral analytes for micellar electrokinetic chromatography. IV. Field-enhanced sample injection 798(1998)251

Naphthoxyacetic acid

Liquid chromatographic determination of β -naphthoxyacetic acid in tomatoes 798(1998)167

Naphthoxyethyl 2-[1-(4-benzyl)piperazyl]ethanesulfonate

Chemically removable derivatization reagent for liquid chromatography: 2-(2-naphthoxy)ethyl 2-[1-(4-benzyl)piperazyl]ethanesulfonate 769(1997)201

1-Naphthylethylamine

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Naphthyl sulfate

Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67

Naproxen

Bonded cellulose-derived high-performance liquid chromatography chiral stationary phases. III. Effect of the reticulation of the cellulose derivative on performance 791(1997)37

Solvent versatility of bonded cellulose-derived chiral stationary phases for high-performance liquid chromatography and its consequences in column loadability 793(1998)239

Narcotine

Effect of on-capillary large volume sample stacking on limits of detection in the capillary zone electrophoretic determination of selected drugs, dyes and metal chelates 772(1997)161

Neohesperidine

Validated high-performance liquid chromatographic method for quantitation of neohesperidine dihydro-chalcone in foodstuffs 758(1997)163

Neomycin sulfate

Productivity and operating regimes in protein chromatography using low-molecular-mass displacers 771(1997)9

Neuropeptides

Quantitative analysis of methionine enkephalin and β -endorphin in the pituitary by liquid secondary ion mass spectrometry and tandem mass spectrometry 794(1998)85

Electrospray ionization mass spectrometric and liquid chromatographic-mass spectrometric studies on the metabolism of synthetic dynorphin A peptides in brain tissue in vitro and in vivo 800(1998)59

Neurotoxins

see also Toxins

Rapid determination of the mycotoxin lolitrem B in endophyte-infected perennial ryegrass by high-performance thin-layer chromatography. A validated assay 769(1997)343

Sensitive determination of anatoxin-a, homoanatoxin-a and their degradation products by liquid chromatography with fluorimetric detection 798(1998)147

Neurotransmitters

Selectivity enhancement for free zone capillary electrophoresis using conventional ion-pairing agents as complexing additives 758(1997)293

Simultaneous determination of total and extracellular concentrations of the amino acid neurotransmitters in cat visual cortex by microbore liquid chromatography and electrochemical detection 798(1998)19

Nitrite/nitrate balance during photoinduced cerebral ischemia in the rat determined by high-performance liquid chromatography with UV and electrochemical detection 798(1998)103

Nicotine

Analysis of selected alkaloids and sugars in tobacco extract 775(1997)101

Nifedipine

Economic approach to robotic sample pretreatment in high-performance liquid chromatography 771(1997)35

Nitrate

Ion chromatography methods for the simultaneous determination of mineral anions in plant sap 752(1996)298

Indirect photometric detection of inorganic anions in microcolumn ion chromatography using octadecylsilica immobilized with bovine serum albumin as stationary phase 755(1996)37

Use of mordenite columns in ion-exclusion chromatography 760(1997)292

Determination of nitrite and nitrate in human serum 770(1997)143

The role of nitric oxide in cancer. Improved methods for measurement of nitrite and nitrate by high-performance ion chromatography 770(1997)151

Optimising the determination of nitrate and phosphate in sea water with ion chromatography using experimental design 771(1997)163

Analysis of nitrate in biological fluids by capillary electrophoresis 781(1997)491

Ion chromatographic separation of metallic ions on a bonded poly- β -diketone-type non-cyclic crown ether stationary phase 789(1997)135

Methods for the determination of nitrite by high-performance liquid chromatography with electrochemical detection (Review) 789(1997)213

Nitrite/nitrate balance during photoinduced cerebral ischemia in the rat determined by high-performance liquid chromatography with UV and electrochemical detection 798(1998)103

Nitrated polynuclear aromatic hydrocarbons

Evidence for the adsorption of nitrated polycyclic aromatic hydrocarbons by tree bark 786(1997)275

Nitric oxide

Determination of nitrite and nitrate in human serum 770(1997)143

The role of nitric oxide in cancer. Improved methods for measurement of nitrite and nitrate by high-performance ion chromatography 770(1997)151

High-performance ion chromatography applied to free-radical mechanisms in drug design. The problem of ion analysis at high ionic strengths 770(1997)273

Nitrite/nitrate balance during photoinduced cerebral ischemia in the rat determined by high-performance liquid chromatography with UV and electrochemical detection 798(1998)103

Nitrilotriacetic acid

Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385

Nitrite

Use of mordenite columns in ion-exclusion chromatography 760(1997)292

Gas chromatographic determination of nitrite in water by pre-column formation of 2-phenylphenol with flame ionization detection 760(1997)319

Determination of nitrite and nitrate in human serum
770(1997)143

The role of nitric oxide in cancer. Improved methods for measurement of nitrite and nitrate by high-performance ion chromatography 770(1997)151

Retention behavior of nitrite on a reversed-phase column with an acidic aqueous eluent containing alcohol as a modifier
789(1997)207

Methods for the determination of nitrite by high-performance liquid chromatography with electrochemical detection (Review)
789(1997)213

Nitrite/nitrate balance during photoinduced cerebral ischemia in the rat determined by high-performance liquid chromatography with UV and electrochemical detection 798(1998)103

Nitroalkanes

Chemically bonded chelates as selective complexing sorbents for gas chromatography v. Silica chemically modified by Cu(II) complexes via amino groups 757(1997)137

Nitroaromatic compounds

Combined use of temperature and solvent strength in reversed-phase gradient elution. IV. Selectivity for neutral (non-ionized) samples as a function of sample type and other separation conditions 756(1996)63

Oxidative and reductive amperometric detection of phenolic and nitroaromatic compounds in packed capillary column supercritical fluid chromatography 785(1997)121

Supercritical CO₂ assisted liquid extraction of nitroaromatic and polycyclic aromatic compounds in soil 785(1997)227

Determination of 4-nitrocatechol in biodegradation samples by gas chromatography–mass spectrometry 793(1998)214

Nitrobenzoic acids

Separation of positional isomers of nitrobenzoic acid by reversed-phase liquid chromatography with 2-propanol–water–acetic acid as eluent 773(1997)365

Nitrobenzoyl chloride

Monitoring the products of condensation of 4-nitrobenzoyl chloride and 2,4-diaminobenzenesulphonic acid by ion-pair high-performance liquid chromatography 777(1997)370

Nitrocatechol

Determination of 4-nitrocatechol in biodegradation samples by gas chromatography–mass spectrometry 793(1998)214

Nitro compounds

Comparison of retention models for the dependence of retention factors on mobile phase composition in reversed-phase high-performance liquid chromatography 797(1998)23

Nitro explosives

Stationary phases with chemically bonded fluorene ligands: A new approach for environmental analysis of π -electron containing solutes 757(1997)3

Nitrofurans

Determination of nitrofurantoin, furazolidone and furaltadone in milk by high-performance liquid chromatography with electrochemical detection 764(1997)243

Determination of nitrofurans residues in avian eggs by liquid chromatography–UV photodiode array detection and confirmation by liquid chromatography–ionspray mass spectrometry 777(1997)201

Nitrofurazone

Determination of nitrofurans residues in avian eggs by liquid chromatography–UV photodiode array detection and confirmation by liquid chromatography–ionspray mass spectrometry 777(1997)201

Nitrogen

Multi-channel on-line chromatographic analysis system for quick air–fuel ratio measurement 752(1996)183

Gas chromatography of Titan's atmosphere. VIII. Analysis of permanent gases with carbon molecular sieve packed capillary columns 761(1997)195

Flow-rate dependence of the height equivalent to a theoretical plate in nitrogen isotope separation by displacement chromatography 793(1998)223

Nitrogen-containing compounds

Prediction of pK_a values of phenolic and nitrogen-containing compounds by computational chemical analysis compared to those measured by liquid chromatography 762(1997)55

Nitrogen dioxide

Chromatographic study of the influence of nitrogen dioxide on the reactions between volatile hydrocarbons and inorganic pigments 776(1997)275

Nitroimidazoles

Management of nonmatrix interfering peaks in a chiral high-performance liquid chromatographic assay produced by solid-phase extraction of rat plasma 763(1997)129

Nitrophenols

On-line solid-phase extraction coupled to supercritical fluid chromatography to determine phenol and nitrophenols in water 755(1996)67

Supercritical fluid and enhanced-fluidity liquid extraction of phenolics from river sediment 764(1997)265

Determination of phenols by solid-phase microextraction 767(1997)171

Effect of cationic polymer on the separation of phenols by capillary electrophoresis 768(1997)161

Displacement electrophoresis of ampholytes in a continuous pH gradient moving in a capillary with a non-constant cross-section 768(1997)283

Determination of nitrophenols by capillary zone electrophoresis in a hydrodynamically closed separation compartment 772(1997)327

Determination of phenolic pollutants in drinking water by capillary electrophoresis in the sample stacking mode 778(1997)279

Determination of priority phenolic compounds in water and industrial effluents by polymeric liquid–solid extraction cartridges using automated sample preparation with extraction columns and liquid chromatography. Use of liquid–solid extraction cartridges for stabilization of phenols 778(1997)301

Determination of polar priority phenols at parts per trillion levels in water using on-line liquid–solid extraction followed by liquid chromatography with coulometric detection 778(1997)313

Liquid chromatography–atmospheric pressure ionization mass spectrometry for the determination of chloro- and nitrophenolic compounds in tap water and sea water 787(1997)79

2-(5-Nitro-2-pyridylazo)-5-[N-propyl-N-(3-sulfopropyl)-amino]phenol

Multi-element trace analysis by reversed-phase high-performance liquid chromatography followed by on-line column enrichment as 2-(5-nitro-2-pyridylazo)-5-[N-propyl-N-(3-sulphopropyl)amino] phenol chelates 788(1997)113

Nitrosamines

Rapid semi-quantitative estimation of N-nitrosodibutylamine and N-nitrosodibenzylamine in smoked hams by solid-phase microextraction followed by gas chromatography–thermal energy analysis 788(1997)131

Nitrosonaphthols

Determination of small anions by capillary electrophoresis using indirect UV detection with sulphonated nitrosonaphthol dyes 767(1997)293

Nitroxide free radicals

Separation of cyclic nitroxide free radicals and their redox forms with dual microelectrochemical detection 771(1997)89

Nonsteroidal anti-inflammatory drugs

Direct injection of large volumes of plasma/serum on a new biocompatible extraction column for the determination of atenolol, propranolol and ibuprofen. Mechanisms for the improvement of chromatographic performance 797(1998)251

Nonylphenol polyethoxylates

Determination of nonylphenol polyethoxylates and their carboxylic acid metabolites in sewage treatment plant sludge by supercritical carbon dioxide extraction 785(1997)385

Noradrenaline

Enhanced temporal resolution for the microdialysis monitoring of catecholamines and excitatory amino acids using capillary electrophoresis with laser-induced fluorescence detection. Analytical developments and in vitro validations 755(1996)99

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Norephedrine

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Nornicotine

Analysis of selected alkaloids and sugars in tobacco extract 775(1997)101

Norobaberine

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Norpredicentrine

Separation of aporphine alkaloids by micellar electrokinetic chromatography 767(1997)277

Norverapamil

Chiral bioanalysis by normal phase high-performance liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 759(1997)65

Nucleic acid bases

Synthesis and characterization of a strong cation exchanger based on polymer-coated silica for high-performance liquid chromatography 779(1997)113

Nucleoside diphosphate kinases

Quaternary structural analysis of nucleoside diphosphate kinases using capillary electrophoresis 781(1997)251

Nucleosides

Quantitation of urinary nucleosides by high-performance liquid chromatography 763(1997)193

Variance contributed by pressure induced injection in capillary electrophoresis 767(1997)205

Application of mixed mobile phases and a step gradient method in capillary electrochromatography for the separation of isomeric polycyclic aromatic hydrocarbon–deoxyribonucleoside adduct mixtures prepared in vitro 781(1997)327

Capillary electrophoresis of urinary normal and modified nucleosides of cancer patients 793(1998)341

Liquid chromatography–mass spectrometry in nucleoside, nucleotide and modified nucleotide characterization 794(1998)109

Analysis of traditional Chinese anticancer drugs by capillary electrophoresis 795(1998)388

New hydrolytic products detected in aqueous solutions of bromodeoxyuridine 797(1998)283

Nucleotides

see also Oligonucleotides

Application of spherical and other polymers in capillary zone electrophoresis: separation of antiviral drugs and deoxyribonucleoside phosphates by different principles 756(1996)263

Variance contributed by pressure induced injection in capillary electrophoresis 767(1997)205

Simple and rapid chromatographic method for the separation of major classes of ribosomal ribomononucleotides 769(1997)225

Iodine–azide reagent in detection of thiophosphoryl nucleotides in thin-layer chromatography systems 773(1997)285

New procedure for the use of high-performance liquid chromatography–electrospray ionization mass spectrometry for the analysis of nucleotides and oligonucleotides 777(1997)3

Separation of ribonucleotides by capillary electrophoresis with multifunctional electrophoretic media of phosphate ammonium salts 779(1997)215

Optimization for the separation of ribonucleotides by capillary electrophoresis at high pH 792(1997)67

Liquid chromatography–mass spectrometry in nucleoside, nucleotide and modified nucleotide characterization 794(1998)109

Erratum to “Separation of ribonucleotides by capillary electrophoresis with multifunctional electrophoretic media of phosphate ammonium salts”. [J. Chromatogr. A, 779 (1997) 215–225] 795(1998)400

O

Obamegine

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Octadecenoic acids

Silver ion high-performance liquid chromatography of isomeric *cis*- and *trans*-octadecenoic acids. Effect of the ester moiety and mobile phase composition 793(1998)275

Octakis(3-O-butanoyl-2,6-di-O-*n*-pentyl)- γ -cyclodextrin

Preparative enantiomer separation of the inhalation anesthetics enflurane, isoflurane and desflurane by gas chromatography on a derivatized γ -cyclodextrin stationary phase 769(1997)119

Octanesulfonic acid

Direct injection of large volumes of plasma/serum on a new biocompatible extraction column for the determination of atenolol, propranolol and ibuprofen. Mechanisms for the improvement of chromatographic performance 797(1998)251

Odorants

Gas chromatographic–mass spectrometric analysis of mercaptan odorants in liquefied petroleum gas and liquefied natural gas 791(1997)367

Odorobioside G

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Ofloxacin

Optical resolution of new quinolone drugs by capillary electrophoresis with ligand-exchange and host–guest interactions 760(1997)235

Okadaic acid

Optimization of sample cleanup procedure for determination of diarrhetic shellfish poisoning toxins by use of experimental design 764(1997)223

Liquid chromatography with fluorimetric, mass spectrometric and tandem mass spectrometric detection for the investigation of the seafood-toxin-producing phytoplankton, *Dinophysis acuta* 777(1997)213

Improvement on sample clean-up for high-performance liquid chromatographic–fluorimetric determination of diarrhetic shellfish toxins using 1-bromoacetylpyrene 793(1998)63

Isolation of a new okadaic acid analogue from phytoplankton implicated in diarrhetic shellfish poisoning 798(1998)137

Olefins

Chemically bonded chelates as selective complexing sorbents for gas chromatography. VI. Modification of silica with NiCl₂ and CoCl₂ via β -diketonate groups 773(1997)209

Oleic acid

Fractionation of *cis*- and *trans*-oleic, linoleic, and conjugated linoleic fatty acid methyl esters by silver ion high-performance liquid chromatography 799(1998)329

Oligogalacturonic acids

Determination of saturated and unsaturated oligogalacturonic acids by means of thin-layer chromatography 756(1996)211

Oligomaltoses

Isotachophoretic focusing of strong electrolytes on the background of carrier ampholytes 798(1998)223

Oligomers

Qualitative and quantitative analysis of styrene and its various oligomers by liquid chromatography 771(1997)111

Oligomers, nitrogen containing

Chemiluminescence nitrogen detection for packed-column supercritical fluid chromatography with methanol modified carbon dioxide 757(1997)183

Oligonucleotides

see also Nucleotides

Determination of single-stranded oligodeoxynucleotides by capillary gel electrophoresis with laser induced fluorescence and on column derivatization 755(1996)271

New procedure for the use of high-performance liquid chromatography–electrospray ionization mass spectrometry for the analysis of nucleotides and oligonucleotides 777(1997)3

Oligopeptides

see also Peptides

Capillary zone electrophoresis with external radial electric field control of electroosmotic flow and its application to the separation of synthetic oligopeptides 772(1997)221

Study of correlations between fine structure of bonded layer and affinity properties of silicas with attached cyclic oligopeptides 777(1997)249

Oligosaccharides

see also Saccharides

Capillary zone electrophoresis of oligosaccharides derivatized with N-(4-aminobenzoyl)-L-glutamic acid for ultraviolet absorbance detection 757(1997)215

Analysis of monosaccharide composition by capillary electrophoresis 763(1997)271

Coupled size-exclusion chromatography–anion-exchange chromatography in the analysis of poly- and oligosaccharides 763(1997)331

Polyacrylamide gradient gel-filled capillaries with low detection background 772(1997)129

Comparison of dynamic fast atom bombardment/liquid secondary ion mass spectrometry and electrospray mass spectrometry coupled to reversed-phase liquid chromatography for the determination of oligosaccharides in seawater 777(1997)141

Study of the enzymatic transformation of fluorescently labeled oligosaccharides in human epidermoid cells using capillary electrophoresis with laser-induced fluorescence detection 781(1997)515

Exoglycosidase matrix-mediated sequencing of a complex glycan pool by capillary electrophoresis 781(1997)547

Determination of impurities in α -cyclodextrin by gradient high-performance liquid chromatography with pulsed amperometric detection 793(1998)57

Overview of the applications of liquid chromatography–mass spectrometry interfacing systems in food analysis: naturally occurring substances in food 794(1998)263

Cationization in electrospray microcolumn liquid chromatography–mass spectrometry 794(1998)359

OliGreen reagent

Determination of single-stranded oligodeoxynucleotides by capillary gel electrophoresis with laser induced fluorescence and on column derivatization 755(1996)271

Omeprazole

Chiral resolution of pantoprazole sodium and related sulfoxides by complex formation with bovine serum albumin in capillary electrophoresis 759(1997)185

Opiates

Micellar electrokinetic chromatography of charged and neutral drugs in acidic running buffers containing a zwitterionic surfactant, sulfonic acids or sodium dodecyl sulphate. Separation of heroin, basic by-products and adulterants 760(1997)245

Separation of free and glucuronidated opioids by capillary electrophoresis in aqueous, binary and micellar media 764(1997)157

Opioids

Separation of free and glucuronidated opioids by capillary electrophoresis in aqueous, binary and micellar media 764(1997)157

Oracin

High-performance liquid chromatography study of stereospecific microsomal enzymes catalysing the reduction of a potential cytostatic drug, oracin. Interspecies comparison 797(1998)197

Orellanine

Novel methods for identification and quantification of the mushroom nephrotoxin orellanine. Thin-layer chromatography and electrophoresis screening of mushrooms with electron spin resonance determination of the toxin 758(1997)145

Organic acids

see also Acids, Carboxylic acids, Fatty acids and Weak acids

Design of double cell quartz crystal detector for ion chromatography and its applications to determination of organic acids in traditional Chinese herb medicine 753(1996)171

Ergot alkaloids as chiral selectors in capillary electrophoresis. Determination of the separation mechanism 755(1996)251

Simultaneous determination of soluble sugars and organic acids as their trimethylsilyl derivatives in apricot fruits by gas-liquid chromatography 758(1997)99

Chromatographic behaviour of diastereoisomers. XIII. Adsorptivity of esters of *Z*- and *E*-2,3-diphenylpropenoic acids and similar compounds on silica in terms of the Snyder theory 758(1997)135

Determination of terrestrial markers in marine environments by gas chromatography-mass-selective detection compared to high-performance liquid chromatography-fluorescence detection 758(1997)243

Selectivity enhancement for free zone capillary electrophoresis using conventional ion-pairing agents as complexing additives 758(1997)293

Application of high-speed counter-current chromatography to the separation of coumarin and related compounds 759(1997)47

Preconcentration determination of inorganic anions and organic acids in power plant waters. Separation optimization through control of column capacity and selectivity 761(1997)163

General equation for calculating the dissociation constants of polyprotic acids and bases from measured retention factors in high-performance liquid chromatography 762(1997)63

Selectivity of alkylamide bonded-phases with respect to organic acids under reversed-phase conditions 762(1997)147

Use of secondary equilibria in reversed-phase high-performance liquid chromatography for the determination of dissociation constants of polyprotic leukotrienes 763(1997)199

High-performance liquid chromatographic determination of dehydroabietic and abietic acids in traditional Chinese medications 763(1997)221

Simultaneous determination of organic acids and inorganic anions in tea by ion chromatography 764(1997)341

Method development and validation for the simultaneous determination of organic and inorganic acids by capillary zone electrophoresis 765(1997)321

Determination of small anions by capillary electrophoresis using indirect UV detection with sulphonated nitrosonaphthol dyes 767(1997)293

Analysis of old brandy and oak extracts by capillary electrophoresis 768(1997)143

Divalent pairing ion for ion-interaction chromatography of sulphonates and carboxylates 770(1997)51

Use of ion chromatography for monitoring microbial spoilage in the fruit juice industry 770(1997)243

Determination of oxalate in parenteral nutrition solutions by capillary electrophoresis 771(1997)285

Fingerprint of anions involving pK_a , mobility and concentration established by using capillary isotachopheresis 772(1997)19

Prediction of zone patterns in capillary zone electrophoresis with conductivity detection. Concept of the zone conductivity diagram 772(1997)63

System peaks in capillary zone electrophoresis. I. Simple model of vacancy electrophoresis 772(1997)73

System peaks in capillary zone electrophoresis. II. Experimental study of vacancy peaks 772(1997)81

Determination of the thermodynamic equilibrium constants of water-insoluble (sparingly soluble) compounds by capillary electrophoresis 772(1997)91

Reversed determination of the formation constants of 1-allyl terguride with mandelic acid optical isomers using capillary electrophoresis 772(1997)179

Determination of low-molecular-mass organic acids in fogwater by ion-exchange chromatography with automatic column switching 773(1997)123

Ion chromatographic characterization of toxic solutions: analysis and ion chemistry of biological liquids (Review) 774(1997)21

Occurrence and determination of organic pollutants in aerosol, precipitation, and sediment samples collected at Lake Balaton 774(1997)349

Analysis of organic acids in industrial samples. Comparison of capillary electrophoresis and ion chromatography 781(1997)205

Investigation of inorganic ions and organic acids in atmospheric aerosol by capillary electrophoresis 781(1997)223

Anion separation by high-performance ion chromatography coupled with indirect photometric detection and 1,2-dihydroxybenzene-3,5-disulfonate or sodium sulfosalicylate as the eluent 789(1997)195

Electroosmotic flow suppressing additives for capillary zone electrophoresis in a hydrodynamically closed separation system 792(1997)483

Dynamic control for the separation of organic acids in capillary electrophoresis 793(1998)145

Quantitative determination of oxalate and other organic acids in drinking water at low $\mu\text{g}/\text{l}$ concentrations 793(1998)198

Determination of glycolic acid in cosmetic products by solid-phase extraction and reversed-phase ion-pair high-performance liquid chromatography 795(1998)219

Use of chiral and achiral ion-pairing reagents in combination with cyclodextrins in capillary electrophoresis 798(1998)281

Determination of organic molecular markers in marine aerosols and sediments: one-step flash chromatography compound class fractionation and capillary gas chromatographic analysis 799(1998)215

Dynamic control and indirect absorption detection for high-speed capillary electrophoretic separation of organic acids 800(1998)339

Organic anions

see also Anions

Determination of anions in landfill leachates by ion chromatography 770(1997)203

Fingerprint of anions involving pK_a , mobility and concentration established by using capillary isotachopheresis 772(1997)19

Organic bases

General equation for calculating the dissociation constants of polyprotic acids and bases from measured retention factors in high-performance liquid chromatography 762(1997)63

Organoantimony compounds

Selective determination of Sb(III) by gas chromatography–quartz furnace atomic absorption spectrometry after derivatization with triphenylmagnesium bromide 764(1997)169

Organoarsenic compounds

High-performance liquid chromatography–hydride generation–atomic fluorescence spectroscopic determination of arsenic species in water 756(1996)292

Organochlorine compounds

Chromatographic techniques in the analysis of organochlorine pesticide residues (Review) 754(1996)33

Matrix effect on supercritical fluid extraction of organochlorine pesticides from sulfur-containing soils 754(1996)285

Determination of polychlorinated biphenyls and chlorinated pesticides in human body fluids and tissues 754(1996)479

Determination of organochlorine compounds in fatty matrices. Application of rapid off-line normal-phase liquid chromatographic clean-up 754(1996)487

Enantiomer separation on a Chirasil-Dex-polymer-coated stationary phase by conventional and micro-packed high-performance liquid chromatography 755(1996)299

Application of solid-phase microextraction to the headspace gas chromatographic analysis of semi-volatile organochlorine contaminants in aqueous matrices 757(1997)173

Elution pattern of planar chlorinated biphenyls and 2,3,7,8-polychlorinated dibenzo-*p*-dioxins/dibenzofurans on chromatographic adsorbents and factors affecting the mechanism of retention. Possibilities of selective separation of both families 759(1997)127

Single-step separation of organochlorine pesticide residues from fatty materials by combined use of solid-matrix partition and C_{18} cartridges 765(1997)61

Analysis of lindane, α - and β -endosulfan and endosulfan sulfate in greenhouse air by gas chromatography 765(1997)99

Determination of pesticides in environmental waters by solid-phase extraction and gas chromatography with electron-capture and mass spectrometry detection 771(1997)221

Simultaneous quantitation of sixteen organochlorine pesticides in drinking waters using automated solid-phase extraction, high-volume injection, high-resolution gas chromatography 773(1997)271

Enantioselective determination of chiral organochlorine compounds in biota by gas chromatography on modified cyclodextrins 774(1997)143

Application of accelerated solvent extraction followed by gas chromatography, high-performance liquid chromatography and gas chromatography–mass spectrometry for the determination of polycyclic aromatic hydrocarbons, chlorinated pesticides and polychlorinated dibenzo-*p*-dioxins and dibenzofurans in solid wastes 774(1997)203

Analysis of toxic wastes in tissues from aquatic species. Applications of matrix solid-phase dispersion 774(1997)287

Analytical methods for the determination of organochlorine compounds. Application to environmental samples in the Slovak Republic 774(1997)333

Purge-and-trap gas chromatography–mass spectrometry in the analysis of volatile organochlorine compounds in water 775(1997)267

High-resolution gas chromatographic test for the characterisation of enantioselective separation of organochlorine compounds. Application to *tert*.-butyldimethylsilyl β -cyclodextrin 775(1997)275

Retention time repeatability as a function of the injection automatism in the analysis of trace organochlorinated compounds with high-resolution gas chromatography 778(1997)23

Hexachlorocyclohexanes in human blood serum 778(1997)87

Supercritical fluid extraction and clean-up of organochlorine pesticides and polychlorinated biphenyls in mussels 790(1997)153

Application of C_{18} disks followed by gas chromatography techniques to degradation kinetics, stability and monitoring of endosulfan in water 795(1998)93

Determination of volatile organic compounds in marine biota 799(1998)233

Organogermanium compounds

Direct determination of trimethylgermanium in water by on-column capillary gas chromatography with flame photometric detection using quartz surface-induced germanium emission 759(1997)119

Separation and determination of inorganic germanium and β -carboxyethylgermanium sesquioxide by high-performance ion-exclusion chromatography 789(1997)403

Organolead compounds

Speciation of trimethyllead in rainwater by gas chromatography–mass spectrometry after ethylation with sodium tetraethylborate. Comparison with other alkylation methods 779(1997)299

Sodium tetra(*n*-propyl)borate: a novel aqueous in situ derivatization reagent for the simultaneous determination of organomercury, -lead and -tin compounds with capillary gas chromatography–inductively coupled plasma mass spectrometry 793(1998)99

Characterization of multicapillary gas chromatography–microwave-induced plasma atomic emission spectrometry for the expeditious analysis for organometallic compounds 795(1998)359

Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385

Organomercury compounds

Reversed-phase high-performance liquid chromatography of pyrrolidinedithiocarbamate complexes of mercuric species using amperometric and coulometric detection 761(1997)139

Development of a capillary electrophoretic method for the separation and determination of trace inorganic and organomercury species utilizing the formation of highly absorbing water soluble dithizone sulphonate complexes 765(1997)345

Interfacing supercritical fluid chromatography with atomic fluorescence spectrometry for the determination of organomercury compounds 786(1997)188

Capillary electrophoresis determination of methylmercury in fish and crab meat after extraction as the dithizone sulphonate complex 791(1997)333

Sodium tetra(*n*-propyl)borate: a novel aqueous in situ derivatization reagent for the simultaneous determination of organomercury, -lead and -tin compounds with capillary gas chromatography–inductively coupled plasma mass spectrometry 793(1998)99

Characterization of multicapillary gas chromatography–microwave-induced plasma atomic emission spectrometry for the expeditious analysis for organometallic compounds 795(1998)359

Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385

Organometallic compounds

Separation of metal ions and metal-containing species by micellar electrokinetic capillary chromatography, including utilisation of metal ions in separations of other species 780(1997)329

Coupling chemical derivatization reactions with supercritical fluid extraction (Review) 785(1997)239

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

Recent applications of high-performance liquid chromatography to the analysis of metal complexes (Review) 789(1997)437

Organophosphorus compounds

Application of liquid chromatography–atmospheric-pressure chemical-ionization mass spectrometry to pesticide analysis 754(1996)61

Synthesis and use of deuterated fenamiphos and its metabolites as internal standards for mass spectrometric analysis in water 754(1996)207

Selective, solid-matrix dispersion extraction of organophosphate pesticide residues from milk 754(1996)497

Solid-phase microextraction of organophosphorus pesticides from water 759(1997)225

Simplified multiresidue method for the determination of organophosphorus insecticides in olive oil 761(1997)327

Enantioseparation of fourteen O-ethyl O-phenyl N-isopropyl phosphoramidothioates by high-performance liquid chromatography on a chiral stationary phase 763(1997)125

Gas chromatographic determination of airborne residues of azinphosmethyl and azinphosmethyl-oxon by cool on-column injection 775(1997)307

Automated sample clean-up and fractionation of chlorpyrifos, chlorpyrifos-methyl and metabolites in mussels using normal-phase liquid chromatography 778(1997)151

Stability of pesticides stored on polymeric solid-phase extraction cartridges 778(1997)161

Use of different surfactants (sodium dodecyl sulfate, bile salts and ionic polymers) in micellar electrokinetic capillary chromatography. Application to the separation of organophosphorus pesticides 778(1997)201

Supercritical fluid extraction: metals as complexes (Review) 785(1997)369

Optimized method for the determination of organophosphorus pesticides in meat and fatty matrices 786(1997)145

Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers 792(1997)419

Investigation of adducts of lanthanide and uranium β -diketonates with organophosphorus Lewis bases by supercritical fluid chromatography 793(1998)107

Analysis of degradation products of organophosphorus chemical warfare agents and related compounds by liquid chromatography–mass spectrometry using electrospray and atmospheric pressure chemical ionisation 794(1998)233

Temperature and extraction voltage effect on fragmentation of organophosphorus pesticides in liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 795(1998)13

Organoselenium compounds

High-performance liquid chromatography of selenoamino acids and organo selenium compounds. Speciation by inductively coupled plasma mass spectrometry 789(1997)349

Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385

Organotin compounds

Reversed-phase ion-pair chromatography with inductively coupled plasma–mass spectrometry detection for the determination of organo-tin compounds in waters 758(1997)85

Chromatographic and hyphenated methods for elemental speciation analysis in environmental media (Review) 774(1997)3

Speciation of organotin compounds in marine biomaterials after basic leaching in a non-focused microwave extractor equipped with pressurized vessels 774(1997)379

Speciation of organotin compounds released from poly(vinyl chloride) at increased temperature by gas chromatography with atomic emission detection 775(1997)295

Interlaboratory evaluation of an off-line supercritical fluid extraction and gas chromatography with atomic emission detection method for the determination of organotin compounds in soil and sediments 785(1997)279

Analytical procedures for the determination of organotin compounds in sediment and biota: a critical review 788(1997)1

Sodium tetra(*n*-propyl)borate: a novel aqueous in situ derivatization reagent for the simultaneous determination of organomercury, -lead and -tin compounds with capillary gas chromatography–inductively coupled plasma mass spectrometry 793(1998)99

Speciation of organo-tin compounds using liquid chromatography–atmospheric pressure ionisation mass spectrometry and liquid chromatography–inductively coupled plasma mass spectrometry as complementary techniques 794(1998)211

Characterization of multicapillary gas chromatography–microwave-induced plasma atomic emission spectrometry for the expeditious analysis for organometallic compounds 795(1998)359

Trace analysis of tributyltin and triphenyltin compounds in sea water by gas chromatography–negative ion chemical ionization mass spectrometry 800(1998)267

Ovalbumin

Critical conditions for separating the microheterogeneous components of glycoproteins by capillary electrophoresis 768(1997)39

Importance of heat of adsorption in modeling protein equilibria for overloaded chromatography 796(1998)141

Ovoglycoprotein

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. II. Comparison of chiral recognition properties with crude ovomucoid 777(1997)241

Separation of basic drug enantiomers by capillary zone electrophoresis using ovoglycoprotein as a chiral selector 782(1997)281

Ovomucoid

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. II. Comparison of chiral recognition properties with crude ovomucoid 777(1997)241

Separation of basic drug enantiomers by capillary zone electrophoresis using ovoglycoprotein as a chiral selector 782(1997)281

Oxalate

Routine determination of major anions in atmospheric aerosols by capillary electrophoresis 770(1997)349

Quantitative determination of oxalate and other organic acids in drinking water at low $\mu\text{g}/\text{l}$ concentrations 793(1998)198

Oxalic acid

Determination of oxalate in parenteral nutrition solutions by capillary electrophoresis 771(1997)285

Oxamniquine

Screening of cyclodextrins by nuclear magnetic resonance for the design of chiral capillary electrophoresis separations 797(1998)149

Oxazepam

On-column enantiomerization of 3-hydroxybenzodiazepines during chiral liquid chromatography with optical rotation detection 767(1997)93

Enantiomeric properties of human albumin immobilized on porous silica supports coated with polymethacryloyl chloride 776(1997)37

Erratum to "On-column enantiomerization of 3-hydroxybenzodiazepines during chiral liquid chromatography with optical rotation detection". [J. Chromatogr. A, 767 (1997) 93–100] 782(1997)296

Oxolinic acid

Determination of flumequine and oxolinic acid in fortified chicken tissue using on-line dialysis and high-performance liquid chromatography with fluorescence detection 799(1998)349

Oxo-oxazolidines

Chromatographic resolution of synthetically useful chiral glycine derivatives by high-performance liquid chromatography 796(1998)299

Oxoprolintane

Direct determination of prolintane and its metabolite oxoprolintane in human urine by capillary zone electrophoresis and β -cyclodextrin-modified micellar electrokinetic chromatography 778(1997)355

Oxybutynin

Chiral bioanalysis by normal phase high-performance liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 759(1997)65

P

Paclitaxel

Determination of a new polymer-bound paclitaxel derivative (PNU 166945), free paclitaxel and 7-epipaclitaxel in dog plasma and urine by reversed-phase high-performance liquid chromatography with UV detection 797(1998)295

Paeoniflorin

Determination of cinnamaldehyde, cinnamic acid, paeoniflorin, glycyrrhizin and [6]-gingerol in the traditional Chinese medicinal preparation Kuei-chih-tang by cyclodextrin-modified micellar electrokinetic chromatography 771(1997)267

Palmitine

Determination of berberine and palmitine in Phellodendri Cortex using ion-pair supercritical fluid chromatography on-line coupled with ion-pair supercritical fluid extraction by on-column trapping 786(1997)371

Pamidronate

Derivatization of pamidronate and other amino(bis)phosphonates with different isothiocyanates prior to ion-pair liquid chromatography 782(1997)211

Pantoprazole

Chiral resolution of pantoprazole sodium and related sulfoxides by complex formation with bovine serum albumin in capillary electrophoresis 759(1997)185

Parabens

Superheated water as an eluent for reversed-phase high-performance liquid chromatography 785(1997)49

Separation of neutral substances by non-aqueous capillary electrophoresis through interactions with cationic additives 792(1997)475

Paracetamol

see also Acetaminophen

^{13}C Isotopic analysis of an acetaminophen and diacetylmorphine mixture 756(1996)203

Development and validation of a high-performance liquid chromatography method for the determination of cold relief ingredients in chewing gum 775(1997)179

Paraquat

Identification of pesticide poisoning in wildlife 754(1996)463

Parathion

Solid-phase microextraction of organophosphorus pesticides from water 759(1997)225

Penicillamine

Electrophoretic studies on mixed complexes. Metal–methionine–penicillamine system 793(1998)220

Penicillins

Micellar electrokinetic capillary chromatography for the separation of phenoxymethylpenicillin and related substances 781(1997)417

Pentaerythritols

Quantitative determination of mono- and dipentaerythritol in synthesis solutions by high-performance liquid chromatography 773(1997)361

Pentafluorobenzylhydroxylamine

Determination of indole-3-pyruvic acid levels in *Arabidopsis thaliana* by gas chromatography–selected ion monitoring–mass spectrometry 800(1998)101

4'-Pentafluorobenzoyloxyacetophenone

Ketone electrophores and an olefin-release group electrophore-labeled DNA oligomer. Detection via electron capture 764(1997)95

Pentafluorophenylhydrazine

Comparison of extraction methods and detection systems in the gas chromatographic analysis of volatile carbonyl compounds 779(1997)360

Determination of indole-3-pyruvic acid levels in *Arabidopsis thaliana* by gas chromatography–selected ion monitoring–mass spectrometry 800(1998)101

Pentamidine

Liquid chromatography–tandem mass spectrometry applied to a study of the metabolism of pentamidine. Discussion of possibilities and problems 777(1997)73

Peptide hormones

Mapping of peptides and protein fragments in human urine using liquid chromatography–mass spectrometry 776(1997)117

Peptides

see also Special Issue Index

pH-Zone-refining countercurrent chromatography 753(1996)1

Combined use of temperature and solvent strength in reversed-phase gradient elution. III. Selectivity for ionizable samples as a function of sample type and pH 756(1996)51

Selectivity enhancement for free zone capillary electrophoresis using conventional ion-pairing agents as complexing additives 758(1997)293

Enantiomeric and diastereomeric separation of di- and tripeptides by capillary electrophoresis 758(1997)303

Quantitative analysis of fluorinated ethylchloroformate derivatives of protein amino acids and hydrolysis products of small peptides using chemical ionization gas chromatography–mass spectrometry 759(1997)111

Partitioning of hydrophobic amino acids and oligopeptides in aqueous two-phase system containing self-aggregating block copolymer. Effects of temperature, salts and surfactants 761(1997)91

Reversed-phase liquid chromatographic separation of platinum(II) complexes of methionine- and histidine-containing peptides using perfluorinated carboxylic acids as ion-pairing reagents 761(1997)115

Use of capillary electrophoresis and high-performance liquid chromatography for monitoring of glycosylation of the peptides dalargin and desmopressin 761(1997)285

Tryptophan analysis in peptides and proteins, mainly by liquid chromatography 763(1997)1

Determination of peptides by high-performance liquid chromatography with laser-induced fluorescence detection 763(1997)23

Practical capillary electrophoresis method for the quantitation of the acetate counter-ion in a novel antifungal lipopeptide 763(1997)279

Analysis of phosphopeptides by capillary electrophoresis and matrix-assisted laser-desorption ionization–time-of-flight mass spectrometry 763(1997)285

Separation of proteins and peptides by capillary electrochromatography in diol- and octadecyl-modified etched capillaries 763(1997)307

Determination of molecular-mass distribution of food-grade protein hydrolyzates by size-exclusion chromatography and chemiluminescent nitrogen detection 763(1997)323

Gradient high-performance liquid chromatographic analysis of enkephalin peptides, their metabolites and enzyme inhibitors using combined ultraviolet and electrochemical detection. I. Optimization of separation and detection 766(1997)85

- Interaction between tryptophan residues and hydrophobically modified dextran. Effect on partitioning of peptides and proteins in aqueous two-phase systems 766(1997)99
- Analysis of bioactive peptides by liquid chromatography–high-resolution electrospray mass spectrometry 767(1997)77
- Complete resolution of imidodipeptide mixtures in urine of prolidase-deficient patients using micellar electrokinetic chromatography 768(1997)57
- Separation of a bioactive cyclic peptide and its oligomeric forms by micellar electrokinetic chromatography 769(1997)301
- Peptide separation by pH-zone-refining countercurrent chromatography 771(1997)81
- Evaluation of pH gradient formation of carrier ampholytes with synthesized isoelectric point markers in capillary isoelectric focusing 772(1997)137
- Capillary electrophoresis of peptides in isoelectric buffers 772(1997)203
- Capillary electrophoresis of large cyanogen bromide peptides of fibre-forming collagens with special reference to cross-linking 772(1997)213
- Capillary zone electrophoresis with external radial electric field control of electroosmotic flow and its application to the separation of synthetic oligopeptides 772(1997)221
- Liquid chromatography on soft packing material, under axial compression. Size-exclusion chromatography of polypeptides 773(1997)103
- Vacancy affinity capillary electrophoresis, a new method for measuring association constants 775(1997)313
- Peptide bank generated by large-scale preparation of circulating human peptides 776(1997)125
- Capillary electrophoresis in the assay of the hydrolysis of glycine-containing peptides by a protease from *Pseudomonas aeruginosa* 776(1997)133
- Matrix-assisted laser desorption/ionisation mass spectrometry guided purification of human guanylin from blood ultrafiltrate 776(1997)139
- Anomalous retention behaviour of peptides on porous graphitized carbon column 776(1997)147
- Use of sodium perchlorate at low pH for peptide separations by reversed-phase liquid chromatography. Influence of perchlorate ion on apparent hydrophilicity of positively charged amino acid side-chains 776(1997)153
- Capillary zone electrophoresis–mass spectrometry as a tool in the stability research of the luteinising hormone-releasing hormone analogue goserelin 776(1997)319
- Rapid analysis of endogenous LVV-hemorphin-7 in cerebrospinal fluid by size-exclusion chromatography and electrospray ionization mass spectrometry 777(1997)41
- Study of correlations between fine structure of bonded layer and affinity properties of silicas with attached cyclic oligopeptides 777(1997)249
- Interactions of the human class II major histocompatibility complex protein HLA-DR4 with a peptide ligand demonstrated by affinity capillary electrophoresis 781(1997)91
- Capillary electrophoresis with electrospray mass spectrometry detection for low-molecular-mass compounds 781(1997)99
- Quantitation of insulin by capillary electrophoresis and high-performance liquid chromatography. Method comparison and validation 781(1997)445
- Enantiomeric separation of a tetrapeptide with cyclodextrin. Extension of the model for chiral capillary electrophoresis by complex formation of one enantiomer molecule with more than one chiral selector molecules 782(1997)271
- Salt effects in capillary zone electrophoresis. I. Dependence of electrophoretic mobilities upon the hydrodynamic radius 786(1997)171
- Enantiomeric determination of amino compounds with high sensitivity using the chiral reagents (+)- and (-)-1-(9-anthryl)-2-propyl chloroformate 786(1997)347
- Displacement chromatography with on-column isomerization 787(1997)1
- Selection of optimum affinity tags from a phage-displayed peptide library. Application to immobilized copper(II) affinity chromatography 787(1997)91
- Development of a capillary zone electrophoresis method for caseinoglycomacropeptide determination 790(1997)195
- Effect of foaming power and foam stability on continuous concentration with foam counter-current chromatography 791(1997)53
- Effect of mobile phase on the oligomerization state of α -helical coiled-coil peptides during high-performance size-exclusion chromatography 791(1997)85
- Generation of tryptic maps of α - and β -globin chains by capillary electrophoresis in isoelectric buffers 791(1997)313

Modification of capillary electrophoresis selectivity in hydro-organic solutions. Dissociation constants and Stokes radius measurements of peptides in water–2,2,2-trifluoroethanol mixtures 792(1997)57

Enantiomeric separation of small chiral peptides by capillary electrophoresis 792(1997)393

Development of liquid chromatography–mass spectrometry using continuous-flow matrix-assisted laser desorption ionization time-of-flight mass spectrometry 794(1998)367

On-line analysis by capillary separations interfaced to an ion trap storage/reflectron time-of-flight mass spectrometer 794(1998)377

Micropreparation of tissue collagenase fragments of type I collagen in the form of surfactant–peptide complexes and their identification by capillary electrophoresis and partial sequencing 796(1998)181

Removal of the fluorescent 4-(aminosulfonyl)-2,1,3-benzoxadiazole label from cysteine-containing peptides 798(1998)47

Sensitive high-resolution analysis of biological molecules by capillary zone electrophoresis coupled with reflecting time-of-flight mass spectrometry 800(1998)1

Substance P and related peptides in porcine cortex: whole tissue and nuclear localization 800(1998)21

Molecular characterization of metal-binding polypeptide domains by electrospray ionization mass spectrometry and metal chelate affinity chromatography 800(1998)29

Electrospray ionization mass spectrometric and liquid chromatographic–mass spectrometric studies on the metabolism of synthetic dynorphin A peptides in brain tissue in vitro and in vivo 800(1998)59

Liquid chromatographic–high-resolution mass spectrometric and tandem mass spectrometric identification of synthetic peptides using electrospray ionization 800(1998)89

Separation of amino acid and peptide stereoisomers by nonionic micelle-mediated capillary electrophoresis after chiral derivatization 800(1998)345

Perfluoropolymers

Preparation and use of ion-exchange chromatographic supports based on perfluoropolymers 773(1997)73

Perfumery compounds

Temperature dependence of the retention index for perfumery compounds on a SE-30 glass capillary column. I. Linear equations 779(1997)287

Perillaldehyde

Quantification of carvone, cineole, perillaldehyde, perillyl alcohol and sobrerol by isocratic high-performance liquid chromatography 793(1998)71

Perillyl alcohol

Quantification of carvone, cineole, perillaldehyde, perillyl alcohol and sobrerol by isocratic high-performance liquid chromatography 793(1998)71

Perloza

High-density ligand attachment to brominated allyl matrices and application to mixed mode chromatography of chymosin 775(1997)39

Permanent gases

see also Gases

Hold-up time in gas chromatography. I. New approach to its estimation 760(1997)219

Gas chromatography of Titan's atmosphere. VIII. Analysis of permanent gases with carbon molecular sieve packed capillary columns 761(1997)195

The hold-up time in gas chromatography. II. Validation of the estimation based on the concept of a zero carbon atoms alkane 767(1997)127

Permethrin

Separation and analysis of the diastereomers and enantiomers of cypermethrin and related compounds 777(1997)363

Peroxyacetyl nitrate

Improvement in the use of capillary columns for ambient air peroxyacetyl nitrate monitoring 786(1997)361

Perthiol drugs

High-performance ion chromatography applied to free-radical mechanisms in drug design. The problem of ion analysis at high ionic strengths 770(1997)273

Pesticides

see also Special Issue Index

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

New design for the on-line solid-phase extraction of pesticides using membrane extraction disk material and liquid chromatography in environmental waters 752(1996)291

Influence of template basicity and hydrophobicity on the molecular recognition properties of molecularly imprinted polymers 753(1996)191

- Microbore liquid chromatographic–mass spectrometric determination of atrazine and its major hydroxylated degradate in water at parts-per-trillion levels using electrospray 753(1996)243
- Recent advances in the residue analysis of N-methylcarbamate pesticides (Review) 754(1996)3
- High-performance separations in the determination of triazine herbicides and their residues (Review) 754(1996)17
- Chromatographic techniques in the analysis of organochlorine pesticide residues (Review) 754(1996)33
- Minimization of solvent consumption in pesticide residue analysis (Review) 754(1996)43
- Screening and analysis of polar pesticides in environmental monitoring programmes by coupled-column liquid chromatography and gas chromatography–mass spectrometry 754(1996)49
- Application of liquid chromatography–atmospheric-pressure chemical-ionization mass spectrometry to pesticide analysis 754(1996)61
- On-line and off-line solid-phase extraction with styrene–divinylbenzene-membrane extraction disks for determining pesticides in water by reversed-phase liquid chromatography–diode-array detection 754(1996)77
- Determination of the fungicides folpet, captan and captafol by cloud-point preconcentration and high-performance liquid chromatography with electrochemical detection 754(1996)85
- Enantiomeric resolution of pyrethroids by high-performance liquid chromatography with diode-laser polarimetric detection 754(1996)97
- Occurrence and determination of pesticides in natural and treated waters (Review) 754(1996)103
- Strategies for chromatographic analysis of pesticide residues in water (Review) 754(1996)125
- Analysis of pesticides in environmental water samples by solid-phase micro-extraction–high-performance liquid chromatography 754(1996)137
- On-line solid-phase extraction coupled to supercritical fluid chromatography with diode array detection for the determination of pesticides in water 754(1996)145
- Monitoring of herbicides in river water by gas chromatography–mass spectrometry and solid-phase extraction 754(1996)159
- Spatial and temporal trends of pesticide residues in water and particulates in the Mississippi River plume and the northwestern Gulf of Mexico 754(1996)169
- Comparative degradation kinetics of alachlor in water by photocatalysis with FeCl₃, TiO₂ and photolysis, studied by solid-phase disk extraction followed by gas chromatographic techniques 754(1996)187
- Supported liquid membranes for sampling and sample preparation of pesticides in water 754(1996)197
- Synthesis and use of deuterated fenamiphos and its metabolites as internal standards for mass spectrometric analysis in water 754(1996)207
- Effect of soil–pesticide interactions on the efficiency of supercritical fluid extraction 754(1996)221
- Analytical method development for the determination of synthetic pyrethroid insecticides in soil by gas chromatography–mass spectrometry operated in negative-ion chemical-ionization mode 754(1996)235
- Possibilities of gas chromatography–atomic emission detection in pesticide multiresidue analysis. Application to herbicide analysis in soils 754(1996)245
- Extraction of triallate from soil with supercritical carbon dioxide and determination by gas chromatography–atomic emission detection. Comparison with a solvent extraction procedure 754(1996)257
- Analysis of the herbicide clopyralid in cultivated soils 754(1996)265
- Application of instrumental thin-layer chromatography and solid-phase extraction to the analyses of pesticide residues in grossly contaminated samples of soil 754(1996)273
- Study of the effect of exogenous organic matter on the mobility of pesticides in soils using soil thin-layer chromatography 754(1996)279
- Matrix effect on supercritical fluid extraction of organochlorine pesticides from sulfur-containing soils 754(1996)285
- Determination of linuron in aqueous soil extracts by high-performance liquid chromatography 754(1996)295
- Determination of pesticide residues in fruit and vegetables (Review) 754(1996)301
- Official multiresidue methods of pesticide analysis in vegetables, fruits and soil (Review) 754(1996)333
- Determination of cereal herbicide residues in environmental samples by gas chromatography (Review) 754(1996)347
- Pesticide residue analyses in plant material by chromatographic methods: clean-up procedures and selective detectors (Review) 754(1996)397

- Pesticide residue analysis in fresh produce by gas chromatography–tandem mass spectrometry 754(1996)411
- Analysis of pesticide residues in drinking water by planar chromatography 754(1996)423
- Complementation of direct-injection high-performance liquid chromatography and enzyme-linked immunosorbent assay for the analysis of thiabendazole in fruit juices and concentrates 754(1996)431
- Optimization of a matrix solid-phase dispersion method for the analysis of pesticide residues in vegetables 754(1996)437
- Application of porous carbon for solid-phase extraction of dicarboximide fungicide residues from wines in combination with high-resolution capillary gas chromatography and gas chromatography–mass spectrometry 754(1996)445
- Off-line high-performance liquid chromatography and solid-phase extraction clean-up for confirmation of pesticide residues in fresh produce by gas chromatography–mass spectrometry 754(1996)455
- Identification of pesticide poisoning in wildlife 754(1996)463
- Determination of polychlorinated biphenyls and chlorinated pesticides in human body fluids and tissues 754(1996)479
- Determination of organochlorine compounds in fatty matrices. Application of rapid off-line normal-phase liquid chromatographic clean-up 754(1996)487
- Selective, solid-matrix dispersion extraction of organophosphate pesticide residues from milk 754(1996)497
- Determination of the fungicide vinclozolin in honey and bee larvae by solid-phase and solvent extraction with gas chromatography and electron-capture and mass spectrometric detection 754(1996)507
- Determination of phenoxy acid herbicides in vegetation, utilising high-resolution gel permeation chromatographic clean-up and methylation with trimethylsilyldiazomethane prior to gas chromatographic analysis with mass-selective detection 755(1996)245
- Enantiomer separation on a Chirasil-Dex-polymer-coated stationary phase by conventional and micro-packed high-performance liquid chromatography 755(1996)299
- Combined use of temperature and solvent strength in reversed-phase gradient elution. III. Selectivity for ionizable samples as a function of sample type and pH 756(1996)51
- Enantiomeric separations of halogen-substituted 2-aryloxypropionic acids by high-performance liquid chromatography on a terguride-based chiral stationary phase 756(1996)119
- Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145
- Application of solid-phase microextraction to the headspace gas chromatographic analysis of semi-volatile organochlorine contaminants in aqueous matrices 757(1997)173
- Capillary electrophoresis of herbicides. II. Evaluation of alkylglucoside chiral surfactants in the enantiomeric separation of phenoxy acid herbicides 757(1997)263
- Indirect liquid chromatographic determination of malathion in formulations, based on the formation of palladium(II)–dimethyldithiosphate complex 758(1997)159
- Determination of chlorophenoxy herbicides in waters by capillary gas chromatography with ion trap detection 758(1997)324
- Prediction of retention in gradient-elution normal-phase high-performance liquid chromatography with binary solvent gradients 759(1997)13
- Rapid analysis of organic microcontaminants in environmental water samples by trace enrichment and liquid chromatography on a single short column 759(1997)55
- Solid-phase microextraction of organophosphorus pesticides from water 759(1997)225
- Retention profiles of some commercial pesticides, pyrethroid and acaricide residues and their application to tomato and parsley plants 760(1997)179
- Preconcentration of triazine herbicides from water by an ion chromatography column and determination by gas chromatography–mass spectrometry 760(1997)314
- Direct determination of bromacil and diuron residues in environmental water samples by coupled-column liquid chromatography and large-volume injection 761(1997)322
- Simplified multiresidue method for the determination of organophosphorus insecticides in olive oil 761(1997)327
- Erratum to ‘‘Gas chromatographic-mass spectrometric study of photodegradation of carbamate pesticides’’. [J. Chromatogr. A, 738 (1996) 225–231] 761(1997)341
- Description and prediction of retention in normal-phase high-performance liquid chromatography with binary and ternary mobile phases 762(1997)15
- Determination of herbicides by reductive amperometric detection in liquid chromatography 763(1997)227

- Adaptation and application of a multi-residue method for the determination of a range of pesticides, including phenoxy acid herbicides in vegetation, used on high-resolution gel permeation chromatographic clean-up and gas chromatographic analysis with mass-selective detection 765(1997)3
- Use of gas-liquid chromatography with electron-capture and thermionic-sensitive detection for the quantitation and identification of pesticide residues 765(1997)13
- Diatomaceous earth-assisted extraction for the multiresidue determination of pesticides 765(1997)23
- Determination of ethylenethiourea in food commodities by a two-step derivatization method and gas chromatography with electron-capture and nitrogen-phosphorus detection 765(1997)31
- Clean-up of aqueous acetone vegetable extracts by solid-matrix partition for pyrethroid residue determination by gas chromatography-electron-capture detection 765(1997)39
- Selective extraction of pyrethroid pesticide residues from milk by solid-matrix dispersion 765(1997)51
- Single-step separation of organochlorine pesticide residues from fatty materials by combined use of solid-matrix partition and C₁₈ cartridges 765(1997)61
- Evaluation of different solid-phase traps for automated collection and clean-up in the analysis of multiple pesticides in fruits and vegetables after supercritical fluid extraction 765(1997)69
- Determination of diflubenzuron in apples by gas chromatography 765(1997)85
- Chromatographic analysis of imazalil and carbendazim in fruits. Method validation and residue monitoring program 1995 765(1997)91
- Analysis of lindane, α - and β -endosulfan and endosulfan sulfate in greenhouse air by gas chromatography 765(1997)99
- Influence of solvent and storage conditions on the stability of acaricide standard stock solutions 765(1997)109
- Adaptation of the internal standard method to a pesticide residues study in/on plants 765(1997)115
- Comparison between accelerated solvent extraction and traditional extraction methods for the analysis of the herbicide diflufenican in soil 765(1997)121
- Simplified determination of combined residues of prochloraz and its metabolites in vegetable, fruit and wheat samples by gas chromatography 765(1997)127
- Ultra-trace-level determination of the antifouling agent Irgarol 1051 by gas chromatography with tandem mass spectrometric detection 766(1997)153
- Analysis of primisulfuron and triasulfuron in water and soil samples by micellar electrokinetic capillary chromatography 766(1997)225
- Direct determination of pyrethrins in pyrethrum extracts by reversed-phase high-performance liquid chromatography with diode-array detection 766(1997)277
- Matrix effects on solid-phase microextraction of organophosphorus pesticides from water 767(1997)195
- Large volume injection of acidic pesticides by reversed-phase micro high-performance liquid chromatography 768(1997)215
- Monitoring of organic micropollutants in surface water by automated on-line trace-enrichment liquid and gas chromatographic systems with ultraviolet diode-array and mass spectrometric detection 768(1997)239
- Extraction and clean-up methods for the determination of organochlorine pesticide residues in medicinal plants 769(1997)275
- Simultaneous quantification of insecticides including carbaryl in drinking water by gas chromatography using dual electron-capture and nitrogen-phosphorus detection 769(1997)285
- Improved chromatographic method for the simultaneous determination of ten phenylurea herbicides and some of their degradation products in soil 769(1997)338
- Relationship between the hydrophobicity and specific hydrophobic surface area of pesticides determined by high-performance liquid chromatography compared with reversed-phase thin-layer chromatography 771(1997)105
- Determination of pesticides in environmental waters by solid-phase extraction and gas chromatography with electron-capture and mass spectrometry detection 771(1997)221
- Simultaneous quantitation of sixteen organochlorine pesticides in drinking waters using automated solid-phase extraction, high-volume injection, high-resolution gas chromatography 773(1997)271
- Enantioselective determination of chiral organochlorine compounds in biota by gas chromatography on modified cyclodextrins 774(1997)143
- Application of accelerated solvent extraction followed by gas chromatography, high-performance liquid chromatography and gas chromatography-mass spectrometry for the determination of polycyclic aromatic hydrocarbons, chlorinated pesticides and polychlorinated dibenzo-*p*-dioxins and dibenzofurans in solid wastes 774(1997)203

- Analysis of toxic wastes in tissues from aquatic species. Applications of matrix solid-phase dispersion 774(1997)287
- Analytical methods for the determination of organochlorine compounds. Application to environmental samples in the Slovak Republic 774(1997)333
- High-resolution gas chromatographic test for the characterisation of enantioselective separation of organochlorine compounds. Application to *tert*.-butyldimethylsilyl β -cyclodextrin 775(1997)275
- Gas chromatographic determination of airborne residues of azinphosmethyl and azinphosmethyl-oxon by cool on-column injection 775(1997)307
- Design of automated solid-phase microextraction for trace analysis of organic compounds in aqueous samples 776(1997)293
- Solid-phase extraction disks for determining pesticides from soil leachates 776(1997)348
- Rapid target analysis of microcontaminants in water by on-line single-short-column liquid chromatography combined with atmospheric pressure chemical ionization tandem mass spectrometry 777(1997)81
- Automated sample preparation with extraction columns by means of anti-isoproturon immunosorbents for the determination of phenylurea herbicides in water followed by liquid chromatography–diode array detection and liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 777(1997)91
- Identification of fenthion and temephos and their transformation products in water by high-performance liquid chromatography with diode array detection and atmospheric pressure chemical ionization mass spectrometric detection 777(1997)99
- Determination of traces of pesticides in water by solid-phase extraction and liquid chromatography–ionspray mass spectrometry 777(1997)115
- Separation and analysis of the diastereomers and enantiomers of cypermethrin and related compounds 777(1997)363
- Retention time repeatability as a function of the injection automatism in the analysis of trace organochlorinated compounds with high-resolution gas chromatography 778(1997)23
- Simplified reversed-phase conditions for the determination of benzimidazole fungicides in fruits by high-performance liquid chromatography with UV detection 778(1997)35
- Development of a new high-performance liquid chromatography method to analyse N-methylcarbamate insecticides by a simple post-column derivatization system and fluorescence detection 778(1997)103
- Matrix effects in the determination of acaricides and fungicides in must by gas chromatography with electron-capture and nitrogen–phosphorus detection 778(1997)111
- Determination of chlorsulfuron and tribenuron-methyl residues in agricultural soils 778(1997)119
- Comparison of octadecylsilica and graphitized carbon black as materials for solid-phase extraction of fungicide and insecticide residues from fruit and vegetables 778(1997)127
- Evaluation of multiwavelength chromatograms for the quantification of mixtures of pesticides by high-performance liquid chromatography–diode array detection with multivariate calibration 778(1997)139
- Automated sample clean-up and fractionation of chlorpyrifos, chlorpyrifos-methyl and metabolites in mussels using normal-phase liquid chromatography 778(1997)151
- Stability of pesticides stored on polymeric solid-phase extraction cartridges 778(1997)161
- New method for the rapid determination of triazine herbicides and some of their main metabolites in water by using coupled-column liquid chromatography and large volume injection 778(1997)171
- Resolution of folpet, procymidone and triazophos in high-performance liquid chromatography–diode array detection by using partial least squares calibration to cross-sections of spectrochromatograms 778(1997)183
- Determination of thiazopyr residues in soil and plants by gas chromatography with nitrogen–phosphorus detection and confirmation by gas chromatography–mass spectrometry 778(1997)193
- Use of different surfactants (sodium dodecyl sulfate, bile salts and ionic polymers) in micellar electrokinetic capillary chromatography. Application to the separation of organophosphorus pesticides 778(1997)201
- Determination of pesticide residues in waters from small loughs by solid-phase extraction and combined use of gas chromatography with electron-capture and nitrogen–phosphorus detection and high-performance liquid chromatography with diode array detection 778(1997)289
- Sulfur-selective chemiluminescence detection with packed column supercritical fluid chromatography 779(1997)307
- Enantiomeric separation of acidic herbicides by capillary electrophoresis using vancomycin as chiral selector 781(1997)503
- Multiresidue analysis of pesticides in fruits and vegetables by gel permeation chromatography followed by gas chromatography with electron-capture and mass spectrometric detection 782(1997)105

- Supercritical fluid extraction of pesticide residues in fortified apple matrices 782(1997)123
- Separation of quaternary ammonium herbicides by capillary electrophoresis with indirect UV detection 782(1997)289
- Coupling chemical derivatization reactions with supercritical fluid extraction (Review) 785(1997)239
- Supercritical fluid extraction of pesticides in foods (Review) 785(1997)289
- Evaluation of a fibrous cellulose drying agent in supercritical fluid extraction and pressurized liquid extraction of diverse pesticides 785(1997)313
- Optimized method for the determination of organophosphorus pesticides in meat and fatty matrices 786(1997)145
- Effects of modifiers, adsorbents and eluents in supercritical fluid extraction of selected pesticides in soil 786(1997)155
- High-performance liquid chromatographic determination of benomyl and carbendazim residues in apian samples 787(1997)129
- Determination of the insecticide imidacloprid in water and soil using high-performance liquid chromatography 787(1997)271
- Study of the feasibility of the use of activated carbon membranes for the on-line cleanup of vegetable extracts in the determination of N-methylcarbamate pesticides by liquid chromatography 788(1997)87
- Application of activated carbon membranes for on-line cleanup of vegetable and fruit extracts in the determination of pesticide multiresidues by gas chromatography with mass selective detection 788(1997)141
- Supercritical fluid extraction and clean-up of organochlorine pesticides and polychlorinated biphenyls in mussels 790(1997)153
- Optimizing recoveries of two chlorotriazine herbicide metabolites and 11 pesticides from aqueous samples using solid-phase extraction and gas chromatography–mass spectrometry 790(1997)161
- Enantiomeric and isomeric separation of herbicides using cyclodextrin-modified capillary zone electrophoresis 790(1997)225
- Characterisation and prediction of retention in isocratic and gradient-elution normal-phase high-performance liquid chromatography on polar bonded stationary phases with binary and ternary solvent systems 791(1997)1
- Chiral separation principles in capillary electrophoresis 792(1997)179
- Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers 792(1997)419
- Comparison of different sorbents for on-line solid-phase extraction of pesticides and phenolic compounds from natural water followed by liquid chromatography 793(1998)257
- Orthogonal array design experiments for optimizing the separation of various pesticides by cyclodextrin-modified micellar electrokinetic chromatography 793(1998)331
- Determination of hexazinone and its metabolites in groundwater by capillary electrophoresis 793(1998)349
- Investigation of photochemical behavior of pesticides in a photolysis reactor coupled on-line with a liquid chromatography–electrospray ionization tandem mass spectrometry system. Application to trace and confirmatory analyses in food samples 794(1998)129
- Comparison of automated on-line solid-phase extraction followed by liquid chromatography–mass spectrometry with atmospheric pressure chemical ionization and particle beam mass spectrometry for the determination of a priority group of pesticides in environmental waters 794(1998)147
- Selective analysis of the herbicides glyphosate and aminomethylphosphonic acid in water by on-line solid-phase extraction–high-performance liquid chromatography–electrospray ionization mass spectrometry 794(1998)187
- Rapid target analysis of microcontaminants in water by on-line single-short-column liquid chromatography combined with atmospheric pressure chemical ionization ion-trap mass spectrometry 794(1998)201
- Resolution and quantitative determination of coeluted pesticide mixtures in liquid chromatography–thermospray mass spectrometry by multivariate curve resolution 795(1998)3
- Temperature and extraction voltage effect on fragmentation of organophosphorus pesticides in liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 795(1998)13
- Effect of the various parameters governing solid-phase microextraction for the trace-determination of pesticides in water 795(1998)27
- Comparison of different clean-up procedures for the determination of N-methylcarbamate insecticides in vegetable matrices by high-performance liquid chromatography with UV detection 795(1998)43
- New and simple method for the analysis of the glutathione adduct of atrazine 795(1998)53
- Stability of polar pesticides on disposable solid-phase extraction precolumns 795(1998)61

Monitoring of pesticides in river water using fully automated on-line solid-phase extraction and liquid chromatography with diode array detection with a novel filtration device 795(1998)71

Multiresidue analysis of pesticides using new laminar extraction disks and liquid chromatography and application to the French priority list 795(1998)83

Application of C₁₈ disks followed by gas chromatography techniques to degradation kinetics, stability and monitoring of endosulfan in water 795(1998)93

Solid-phase microextraction and gas chromatography with mass spectrometric detection for the determination of pesticides in aqueous samples 795(1998)105

Determination of pesticides in high-water-content samples by off-line supercritical fluid extraction–gas chromatography–electron-capture detection 795(1998)117

Dissipation of some citrus selective residual herbicides in an irrigation well 795(1998)125

Inter-laboratory validation of solid-phase microextraction for the determination of triazine herbicides and their degradation products at ng/l level in water samples 795(1998)371

Determination of aryloxyphenoxypropionic acid herbicides in water using different solid-phase extraction procedures and liquid chromatography–diode array detection 796(1998)309

Optimisation of gradient elution in normal-phase high-performance liquid chromatography 797(1998)11

Use of principal component analysis for studying the separation of pesticides on polyethylene-coated silica columns 797(1998)33

Microwave-assisted solvent extraction and reversed-phase liquid chromatography–UV detection for screening soils for sulfonylurea herbicides 798(1998)179

Neural network and experimental design to investigate the effect of five factors in ion-interaction high-performance liquid chromatography 799(1998)47

Resolution of imidacloprid pesticide and its metabolite 6-chloronicotinic acid using cross-sections of spectrochromatograms obtained by high-performance liquid chromatography with diode-array detection 799(1998)149

Capillary supercritical fluid chromatography of pyrethrins and pyrethroids with positive pressure and negative temperature gradients 799(1998)265

Simultaneous determination and enantiomeric resolution of mecoprop and dichlorprop in soil samples by high-performance liquid chromatography and gas chromatography–mass spectrometry 799(1998)355

High sample throughput flow immunoassay utilising restricted access columns for the separation of bound and free label 800(1998)219

Matrix-induced effects: a critical point in the gas chromatographic analysis of pesticide residues 800(1998)283

Stability of pesticides in plant extracts used as calibrants in the gas chromatographic analysis of residues 800(1998)297

Pethidine

Effect of on-capillary large volume sample stacking on limits of detection in the capillary zone electrophoretic determination of selected drugs, dyes and metal chelates 772(1997)161

Phenanthrene

Influence of the capillary dimensions on the performance of the preconcentration technique based on parallel current chromatography 791(1997)163

Phenazines

High-performance liquid chromatographic analysis of *Pseudomonas aeruginosa* phenazines 771(1997)99

Phenethylamines

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

Effect of β -cyclodextrin acetylation on the resolution of phenethylamines with capillary electrophoresis and nuclear magnetic resonance spectroscopy 758(1997)277

Phenolic acids

Comparative investigation of UV, electrochemical and particle beam mass spectrometric detection for the high-performance liquid chromatographic determination of benzoic and cinnamic acids and of their corresponding phenolic acids 753(1996)157

Separation of phenolic aldehydes, ketones and acids from lignin degradation by capillary zone electrophoresis 755(1996)89

Identification of phenolic acids and inositols in balms and tissues from an Egyptian mummy 767(1997)177

Phenolic compounds

Chiral crown ether-anchored polysiloxanes as capillary gas chromatography stationary phases 753(1996)269

Separation of phenolic aldehydes, ketones and acids from lignin degradation by capillary zone electrophoresis 755(1996)89

Procedures for analysing phenolic metabolites of polychlorinated dibenzofurans, -dibenzo-*p*-dioxins and -biphenyls extracted from a microsomal assay: optimising solid-phase adsorption clean-up and derivatisation methods 761(1997)219

Prediction of pK_a values of phenolic and nitrogen-containing compounds by computational chemical analysis compared to those measured by liquid chromatography 762(1997)55

- Supercritical fluid and enhanced-fluidity liquid extraction of phenolics from river sediment 764(1997)265
- Analysis of phenolic compounds in industrial wastewater with high-performance liquid chromatography and post-column reaction detection 769(1997)189
- Chemically modified polymeric resin used as sorbent in a solid-phase extraction process to determine phenolic compounds in water 771(1997)55
- Identification of antioxidative compounds in plant beverages by capillary electrophoresis with the marker index technique 781(1997)523
- Oxidative and reductive amperometric detection of phenolic and nitroaromatic compounds in packed capillary column supercritical fluid chromatography 785(1997)121
- Sub-minute separations of organic and inorganic anions with co-electroosmotic capillary electrophoresis 787(1997)243
- Comparison of different sorbents for on-line solid-phase extraction of pesticides and phenolic compounds from natural water followed by liquid chromatography 793(1998)257
- High-performance liquid chromatographic determination of flavonoids in *Betula pendula* and *Betula pubescens* leaves 793(1998)370
- Micellar electrokinetic chromatography as an alternative to high-performance liquid chromatography for separation and determination of phenolic compounds in Japanese spirituous liquor 793(1998)409
- Overview of the applications of liquid chromatography–mass spectrometry interfacing systems in food analysis: naturally occurring substances in food 794(1998)263
- Separation of blackcurrant anthocyanins by capillary zone electrophoresis 799(1998)321
- Phenols**
- On-line solid-phase extraction coupled to supercritical fluid chromatography to determine phenol and nitrophenols in water 755(1996)67
- Experiences with packed capillary electrochromatography at ambient pressure 755(1996)165
- Examination of a new chromatographic function, based on an exponential resolution term, for use in optimization strategies: application to capillary gas chromatography separation of phenols 755(1996)235
- Phenol and methylphenol isomers determination in soils by in-situ microwave-assisted extraction and derivatisation 757(1997)153
- Trace enrichment of phenols by on-line solid-phase extraction and gas chromatographic determination 757(1997)165
- Effects of the gradient profile on the production rate in reversed-phase gradient elution overloaded chromatography 760(1997)25
- Fitting competitive adsorption isotherms to the experimental distribution data in reversed-phase systems 762(1997)3
- Description and prediction of retention in normal-phase high-performance liquid chromatography with binary and ternary mobile phases 762(1997)15
- Continuous membrane extraction of phenols from crude oils followed by high-performance liquid chromatographic determination with electrochemical detection 766(1997)61
- Determination of phenols by solid-phase microextraction 767(1997)171
- Effect of cationic polymer on the separation of phenols by capillary electrophoresis 768(1997)161
- Phenolic compounds in virgin olive oils. I. Low-wavelength quantitative determination of complex phenols by high-performance liquid chromatography under isocratic elution 768(1997)207
- Analysis of phenolic compounds in effluent by solid-phase extraction and gas chromatography–mass spectrometry with direct on-column benzylation: sensitive negative ion chemical ionisation gas chromatography–mass spectrometry detection of phenyl benzyl ethers 771(1997)203
- Non-aqueous capillary electrophoretic separation of Brønsted acids as heteroconjugated anions 771(1997)275
- Determination of nitrophenols by capillary zone electrophoresis in a hydrodynamically closed separation compartment 772(1997)327
- Modelling gradient elution in centrifugal partition chromatography 773(1997)13
- Evaluation of supercritical fluid extraction, microwave-assisted extraction and sonication in the determination of some phenolic compounds from various soil matrices 774(1997)243
- Analytical methods for the determination of organochlorine compounds. Application to environmental samples in the Slovak Republic 774(1997)333
- Simultaneous separation of metal ions and neutral organics by reversed-phase ion-pair liquid chromatography 776(1997)221
- Determination of phenolic pollutants in drinking water by capillary electrophoresis in the sample stacking mode 778(1997)279

Determination of priority phenolic compounds in water and industrial effluents by polymeric liquid–solid extraction cartridges using automated sample preparation with extraction columns and liquid chromatography. Use of liquid–solid extraction cartridges for stabilization of phenols 778(1997)301

Determination of polar priority phenols at parts per trillion levels in water using on-line liquid–solid extraction followed by liquid chromatography with coulometric detection 778(1997)313

Application of particles as pseudo-stationary phases in electrokinetic chromatography 780(1997)63

A simple means of generating pH gradients in capillary zone electrophoresis 781(1997)3

Superheated water as an eluent for reversed-phase high-performance liquid chromatography 785(1997)49

Coupling chemical derivatization reactions with supercritical fluid extraction (Review) 785(1997)239

New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259

Evaluation of two solid-phase extraction procedures for the preconcentration of chlorophenols in drinking water 786(1997)285

Liquid chromatography–atmospheric pressure ionization mass spectrometry for the determination of chloro- and nitrophenolic compounds in tap water and sea water 787(1997)79

Sample stacking using field-amplified sample injection in capillary zone electrophoresis in the analysis of phenolic compounds 788(1997)185

Influence of the capillary dimensions on the performance of the preconcentration technique based on parallel current chromatography 791(1997)163

Optimization of the recovery yield and of the production rate in overloaded gradient-elution reversed-phase chromatography 796(1998)115

Separation of tea polyphenols using micellar electrokinetic chromatography with diode array detection 799(1998)309

Phenoxy acids

Determination of phenoxy acid herbicides in vegetation, utilising high-resolution gel permeation chromatographic clean-up and methylation with trimethylsilyldiazomethane prior to gas chromatographic analysis with mass-selective detection 755(1996)245

Capillary electrophoresis of herbicides. II. Evaluation of alkylglucoside chiral surfactants in the enantiomeric separation of phenoxy acid herbicides 757(1997)263

Adaptation and application of a multi-residue method for the determination of a range of pesticides, including phenoxy acid herbicides in vegetation, used on high-resolution gel permeation chromatographic clean-up and gas chromatographic analysis with mass-selective detection 765(1997)3

Large volume injection of acidic pesticides by reversed-phase micro high-performance liquid chromatography 768(1997)215

Phenoxy acids, methyl esters

Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers 792(1997)419

Phenoxymethylpenicillin

Micellar electrokinetic capillary chromatography for the separation of phenoxymethylpenicillin and related substances 781(1997)417

Phenylacetylaminopiperidinedione

Enantioseparation of 3-phenylacetyl-amino-2,6-piperidinedione and related chiral compounds 752(1996)93

Phenylalanine

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Partitioning of hydrophobic amino acids and oligopeptides in aqueous two-phase system containing self-aggregating block copolymer. Effects of temperature, salts and surfactants 761(1997)91

Determination of aspartame and phenylalanine in diet soft drinks by high-performance liquid chromatography with direct spectrofluorimetric detection 773(1997)163

Phenylcyclohexenes

Comparison of supercritical fluid extraction and Soxhlet extraction of organic compounds from carpet samples 787(1997)171

Phenylethanethiol

Determination of the absolute configuration of sugar residues using gas chromatography. Method with potential for elimination of references 769(1997)253

Phenylethanol

Extension of a Jovanovic–Freundlich isotherm model to multicomponent adsorption on heterogeneous surfaces 796(1998)15

Phenylethylamines

Capillary zone electrophoretic separation of biogenic amines: influence of organic modifier 755(1996)142

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Determination of methamphetamine and related compounds by capillary electrophoresis with UV and laser-induced fluorescence detection 798(1998)325

1-Phenyl-3-methyl-4-benzoyl-5-pyrazolone

Chromatographic behaviour of the chromium(III) complex of 1-phenyl-3-methyl-4-benzoyl-5-pyrazolone 789(1997)479

Phenylmethylpyrazolone

Separation of aldose enantiomers by capillary electrophoresis in the presence of optically active N-dodecoxycarbonylvalines 792(1997)385

1-Phenyl-3-methyl-5-pyrazolone derivatives

Unusual separation of 1-phenyl-3-methyl-5-pyrazolone derivatives of aldoses by capillary zone electrophoresis 791(1997)307

Phenylphenol

Gas chromatographic determination of nitrite in water by pre-column formation of 2-phenylphenol with flame ionization detection 760(1997)319

Phenylpropanols

Chiral supercritical fluid chromatography of phenylpropanols and related compounds 785(1997)179

Extension of a Jovanovic–Freundlich isotherm model to multicomponent adsorption on heterogeneous surfaces 796(1998)15

Phenylpropionic acid

Comparison of polymer coatings of capillaries for capillary electrophoresis with respect to their applicability to molecular imprinting and electrochromatography 781(1997)43

Phenylpyridylpropylamines

Chiral separation of 3-phenyl-3-(2-pyridyl)propylamines, and analogous guanidines and guanidine-N-carboxylic acid esters with high-performance liquid chromatography and capillary zone electrophoresis 793(1998)77

Phenylthiohydantoin-alanine

Indirect time-resolved fluorescence detection of both non-fluorescent and fluorescent compounds separated by high-performance liquid chromatography 787(1997)261

Phenylureas

Determination of linuron in aqueous soil extracts by high-performance liquid chromatography 754(1996)295

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Prediction of retention in gradient-elution normal-phase high-performance liquid chromatography with binary solvent gradients 759(1997)13

Improved chromatographic method for the simultaneous determination of ten phenylurea herbicides and some of their degradation products in soil 769(1997)338

Rapid target analysis of microcontaminants in water by on-line single-short-column liquid chromatography combined with atmospheric pressure chemical ionization tandem mass spectrometry 777(1997)81

Automated sample preparation with extraction columns by means of anti-isoproturon immunosorbents for the determination of phenylurea herbicides in water followed by liquid chromatography–diode array detection and liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 777(1997)91

Optimisation of gradient elution in normal-phase high-performance liquid chromatography 797(1998)11

Pheromones

Gas chromatographic determination of vapour pressures of pheromone-like compounds. IV. Acetates, a reinvestigation 759(1997)93

Phosphate

Ion chromatography methods for the simultaneous determination of mineral anions in plant sap 752(1996)298

Determination of phosphate in natural waters by capillary electrophoresis. Influence of solution composition on migration time and response 770(1997)321

Optimising the determination of nitrate and phosphate in sea water with ion chromatography using experimental design 771(1997)163

Development of a capillary electrophoresis method with indirect photometric detection for the determination of anions related to anticaries 781(1997)457

Phosphate salts

Chromatographic ion-exchange and change-partner reactions. New mechanisms concerning ion–ion interactions of inorganic compounds in Sephadex G-15 columns (Review) 789(1997)21

Phosphatidylcholine

Reversed-phase separations of nitrogenous phospholipids on an octadecanoyl poly(vinyl alcohol) phase 773(1997)93

Phosphatidylethanolamine

Reversed-phase separations of nitrogenous phospholipids on an octadecanoyl poly(vinyl alcohol) phase 773(1997)93

Phosphatidylinositol

Application of capillary electrophoresis to determination of enzyme activity and other properties of phosphatidylinositol-specific phospholipase C 781(1997)263

Phosphoamino acid

High sample throughput phosphoamino acid analysis of proteins separated by one- and two-dimensional gel electrophoresis 764(1997)201

Phospholipases

Application of capillary electrophoresis to determination of enzyme activity and other properties of phosphatidylinositol-specific phospholipase C 781(1997)263

Phospholipids

Reversed-phase separations of nitrogenous phospholipids on an octadecanoyl poly(vinyl alcohol) phase 773(1997)93

Application of capillary electrophoresis to determination of enzyme activity and other properties of phosphatidylinositol-specific phospholipase C 781(1997)263

Phosphonates

Determination of small anions by capillary electrophoresis using indirect UV detection with sulphonated nitrosonaphthol dyes 767(1997)293

Analysis of amine-containing phosphonates in detergent powders by anion-exchange chromatography with pulsed amperometric detection 771(1997)155

Determination of phosphonates in natural waters by ion-pair high-performance liquid chromatography 773(1997)139

Phosphonic acids

In-line respeciation: an ion-exchange ion chromatographic method applied to the separation of degradation products of chemical warfare nerve agents in soil 770(1997)253

Analysis of degradation products of organophosphorus chemical warfare agents and related compounds by liquid chromatography–mass spectrometry using electrospray and atmospheric pressure chemical ionisation 794(1998)233

Phosphopeptides

Analysis of phosphopeptides by capillary electrophoresis and matrix-assisted laser-desorption ionization–time-of-flight mass spectrometry 763(1997)285

Phosphoramidothioates

Enantioseparation of fourteen O-ethyl O-phenyl N-isopropyl phosphoramidothioates by high-performance liquid chromatography on a chiral stationary phase 763(1997)125

Phosphorus

Measurement of trace phosphorus in dichlorosilane by high-temperature hydrogen reduction gas chromatography 757(1997)319

Phthalaldehyde

Determination of amino acids by high-performance capillary electrophoresis with on-line mode in-capillary derivatization 787(1997)253

Phthalates

Separation behavior of inorganic anions on an octadecyl-bonded silica column coated with cetyltrimethylammonium by phthalate and salicylate eluents 764(1997)346

Supercritical fluid chromatography–proton nuclear magnetic resonance spectroscopy coupling (Review) 785(1997)65

Phthaldialdehyde

Comparison of the stability and UV and fluorescence characteristics of the *o*-phthaldialdehyde/3-mercaptopropionic acid and *o*-phthaldialdehyde/N-acetyl-L-cysteine reagents and those of their amino acid derivatives 798(1998)37

Phthalocyanines

Analysis of sulphonated phthalocyanine dyes by capillary electrophoresis 770(1997)345

Phycobilisomes

Chromatographic purification of biliproteins from *Spirulina platensis*. High-performance liquid chromatographic separation of their α and β subunits 778(1997)441

Phytic acid

Evaluation of phytic acid as a buffer additive for the separation of proteins in capillary electrophoresis 768(1997)307

Phytoestrogens

Development and validation of an isocratic high-performance liquid chromatographic method for quantitative determination of phytoestrogens in soya bean 795(1998)377

Phytosphingosines

Cell lipids of the *Candida lipolytica* yeast grown on methanol 755(1996)75

Picoline

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Picrate

Nonaqueous anion-exchange chromatography. II. Changing anion-exchange selectivity by resin surface complex formation of crown ethers 758(1997)29

Pigments

Chromatographic study of the influence of nitrogen dioxide on the reactions between volatile hydrocarbons and inorganic pigments 776(1997)275

Study of binding media in works of art by gas chromatographic analysis of amino acids and fatty acids derivatized with ethyl chloroformate 778(1997)373

Pinenes

Chiral separation of monoterpenes using mixtures of sulfated β -cyclodextrins and α -cyclodextrin as chiral additives in the reversed-polarity capillary electrophoresis mode 759(1997)149

Decomposition of α -pinene and sabinene on solid sorbents, Tenax TA and Carboxen 786(1997)185

Pipecolic acid methyl esters

Pipecolic acid methyl esters as artefacts from the ion-exchange chromatography of *Inga punctata* foliar extracts 766(1997)267

Piperazineethanols

Displacement electrophoresis of ampholytes in a continuous pH gradient moving in a capillary with a non-constant cross-section 768(1997)283

Piperazines

Relationship between structure and reversed-phase thin-layer chromatographic lipophilicity parameters in a group of piperazine derivatives 766(1997)165

Piperazines, 1,4-disubstituted

Non-linear structure–enantioselective retention relationships in a homologous series of 1,4-disubstituted piperazine derivatives 788(1997)81

Piperidines

Pharmacokinetic screening for the selection of new drug discovery candidates is greatly enhanced through the use of liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 777(1997)61

Chiral resolution of functionalized piperidine enantiomers by capillary electrophoresis with native, alkylated and anionic β -cyclodextrins 796(1998)375

Pirimicarb

Resolution and quantitative determination of coeluted pesticide mixtures in liquid chromatography–thermospray mass spectrometry by multivariate curve resolution 795(1998)3

Plant hormones

Determination of plant indoles by capillary electrophoresis with amperometric detection 755(1996)289

Quantitative analysis of 1-aminocyclopropane-1-carboxylic acid by liquid chromatography coupled to electrospray tandem mass spectrometry 775(1997)143

Reversed-phase C_{18} high-performance liquid chromatography of gibberellins GA_3 and GA_1 782(1997)137

Erratum to “Reversed-phase C_{18} high-performance liquid chromatography of gibberellins GA_3 and GA_1 ”. [J. Chromatogr. A, 782 (1997) 137–139] 795(1998)399

Platinum

Reversed-phase liquid chromatographic separation of platinum(II) complexes of methionine- and histidine-containing peptides using perfluorinated carboxylic acids as ion-pairing reagents 761(1997)115

Speciation of platinum–chloro complexes and their hydrolysis products by ion chromatography. Determination of platinum oxidation states 775(1997)197

Platinum-group metals

Determination of platinum, palladium, osmium, iridium, rhodium and gold as chloro complexes by capillary zone electrophoresis 772(1997)347

Pluronic P105

Partitioning of hydrophobic amino acids and oligopeptides in aqueous two-phase system containing self-aggregating block copolymer. Effects of temperature, salts and surfactants 761(1997)91

Pneumocandins

Practical capillary electrophoresis method for the quantitation of the acetate counter-ion in a novel antifungal lipopeptide 763(1997)279

PNU 166945

Determination of a new polymer-bound paclitaxel derivative (PNU 166945), free paclitaxel and 7-epipaclitaxel in dog plasma and urine by reversed-phase high-performance liquid chromatography with UV detection 797(1998)295

Polar compounds

Separation of polar solutes by packed column supercritical fluid chromatography (Review) 785(1997)3

Polar organic compounds

Analysis of polar organic pollutants in the Elbe river by flow injection analysis and high-performance liquid chromatography with tandem mass spectrometry 777(1997)127

Polyacrylamide

Determination of polyacrylamide in polyvinyl alcohol by pyrolysis–gas chromatography with atomic emission detection 753(1996)101

Polyamines

High-performance liquid chromatographic determination of polyamines in milk as their 9-fluorenylmethoxycarbonyl derivatives using a column-switching technique 791(1997)79

Polyamino acids

Polyacrylamide gradient gel-filled capillaries with low detection background 772(1997)129

Polyaminopolyacetic acids

Characterization of metal complexes of pharmaceutical interest by capillary electrophoresis with element sensitive detection 781(1997)407

Polyanions

Macromolecules in ordered media VIII. High-performance size-exclusion chromatography as a technique for characterizing the interaction between polyanions and cationic liposomes 778(1997)53

Polybrominated biphenyls

Supercritical fluid extraction with solid-phase trapping of chlorinated and brominated pollutants from sediment samples 774(1997)229

Polybrominated diphenyl ethers

Supercritical fluid extraction with solid-phase trapping of chlorinated and brominated pollutants from sediment samples 774(1997)229

Polybutadiene

Factors influencing polybutadiene deposition within porous chromatographic zirconia 791(1997)45

Polychlorinated benzenes

Supercritical fluid extraction with solid-phase trapping of chlorinated and brominated pollutants from sediment samples 774(1997)229

Polychlorinated biphenyls

Application of correlation analysis for identification of polychlorinated biphenyls 752(1996)197

Separation of chiral polychlorinated biphenyls by micellar electrokinetic chromatography using β - and γ -cyclodextrin mixtures in the separation buffer 752(1996)265

Occurrence and determination of pesticides in natural and treated waters (Review) 754(1996)103

Spatial and temporal trends of pesticide residues in water and particulates in the Mississippi River plume and the northwestern Gulf of Mexico 754(1996)169

Determination of polychlorinated biphenyls and chlorinated pesticides in human body fluids and tissues 754(1996)479

Determination of organochlorine compounds in fatty matrices. Application of rapid off-line normal-phase liquid chromatographic clean-up 754(1996)487

Integrating gel permeation chromatography clean-up in the analysis of metabolites of polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans extracted from a microsomal assay. A comparison of different mobile phases 755(1996)57

Elution pattern of planar chlorinated biphenyls and 2,3,7,8-polychlorinated dibenzo-*p*-dioxins/dibenzofurans on chromatographic adsorbents and factors affecting the mechanism of retention. Possibilities of selective separation of both families 759(1997)127

Procedures for analysing phenolic metabolites of polychlorinated dibenzofurans, -dibenzo-*p*-dioxins and -biphenyls extracted from a microsomal assay: optimising solid-phase adsorption clean-up and derivatisation methods 761(1997)219

Development of a thermal desorption modulator for gas chromatography 767(1997)137

Enantiomer separation of selected atropisomeric polychlorinated biphenyls including PCB 144 on *tert*-butyldimethylsilylated β -cyclodextrin 769(1997)247

Separation of extracts from biological tissues into polycyclic aromatic hydrocarbon, polychlorinated biphenyl and polychlorinated dibenzo-*p*-dioxin/polychlorinated dibenzofuran fractions prior to analysis 771(1997)169

Enantioselective determination of chiral organochlorine compounds in biota by gas chromatography on modified cyclodextrins 774(1997)143

Analysis of toxic wastes in tissues from aquatic species. Applications of matrix solid-phase dispersion 774(1997)287

Chromatography pattern recognition of Aroclors using iterative probabilistic neural networks 775(1997)231

Negative chromatographic peaks with oxygen doped electron capture detection of polychlorinated biphenyls 775(1997)368

Micellar electrokinetic capillary chromatographic separation of polychlorinated biphenyl congeners 778(1997)77

Field-portable determination of polychlorinated biphenyls and polynuclear aromatic hydrocarbons in soil using supercritical fluid extraction 785(1997)205

Extraction of polychlorinated biphenyls with water under subcritical conditions 785(1997)219

Selectivity of electron-donor- and electron-acceptor-bonded silica packing materials for hydrophobic environmental contaminants in polar and non-polar eluents 786(1997)237

Gas chromatographic retention index system for polychlorinated biphenyls: possibilities and limitations 787(1997)215

Supercritical fluid extraction and clean-up of organochlorine pesticides and polychlorinated biphenyls in mussels 790(1997)153

Elution order of the 209 polychlorinated biphenyls on a high-temperature capillary column 799(1998)249

Interfering contaminants in carbon dioxide solvent used in the supercritical fluid extraction of polychlorinated biphenyls 799(1998)259

Solid-phase microextraction of polychlorinated biphenyls 800(1998)257

Polychlorinated biphenyls, hydroxylated

Determination of hydroxylated polychlorinated biphenyls by ion trap gas chromatography–tandem mass spectrometry 790(1997)131

Polychlorinated dibenzo-*p*-dioxins

Integrating gel permeation chromatography clean-up in the analysis of metabolites of polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans extracted from a microsomal assay. A comparison of different mobile phases 755(1996)57

Elution pattern of planar chlorinated biphenyls and 2,3,7,8-polychlorinated dibenzo-*p*-dioxins/dibenzofurans on chromatographic adsorbents and factors affecting the mechanism of retention. Possibilities of selective separation of both families 759(1997)127

Procedures for analysing phenolic metabolites of polychlorinated dibenzofurans, -dibenzo-*p*-dioxins and -biphenyls extracted from a microsomal assay: optimising solid-phase adsorption clean-up and derivatisation methods 761(1997)219

Separation of extracts from biological tissues into polycyclic aromatic hydrocarbon, polychlorinated biphenyl and polychlorinated dibenzo-*p*-dioxin/polychlorinated dibenzofuran fractions prior to analysis 771(1997)169

Gas chromatographic analysis of polychlorinated dibenzo-*p*-dioxins and dibenzofurans 774(1997)97

Application of accelerated solvent extraction followed by gas chromatography, high-performance liquid chromatography and gas chromatography–mass spectrometry for the determination of polycyclic aromatic hydrocarbons, chlorinated pesticides and polychlorinated dibenzo-*p*-dioxins and dibenzofurans in solid wastes 774(1997)203

Supercritical CO₂-assisted liquid extraction of polycyclic aromatic hydrocarbons and polychlorinated dibenzo-*p*-dioxins and -furans from solid matrices 777(1997)289

Application of DB-5ms gas chromatography column for the complete assignment of 2,3,7,8-substituted polychlorodibenzo-*p*-dioxins and polychlorodibenzofurans in samples from municipal waste incinerator emissions 786(1997)125

Selectivity of electron-donor- and electron-acceptor-bonded silica packing materials for hydrophobic environmental contaminants in polar and non-polar eluents 786(1997)237

Polychlorinated dibenzofurans

Integrating gel permeation chromatography clean-up in the analysis of metabolites of polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans extracted from a microsomal assay. A comparison of different mobile phases 755(1996)57

Elution pattern of planar chlorinated biphenyls and 2,3,7,8-polychlorinated dibenzo-*p*-dioxins/dibenzofurans on chromatographic adsorbents and factors affecting the mechanism of retention. Possibilities of selective separation of both families 759(1997)127

Procedures for analysing phenolic metabolites of polychlorinated dibenzofurans, -dibenzo-*p*-dioxins and -biphenyls extracted from a microsomal assay: optimising solid-phase adsorption clean-up and derivatisation methods 761(1997)219

Separation of extracts from biological tissues into polycyclic aromatic hydrocarbon, polychlorinated biphenyl and polychlorinated dibenzo-*p*-dioxin/polychlorinated dibenzofuran fractions prior to analysis 771(1997)169

Gas chromatographic analysis of polychlorinated dibenzo-*p*-dioxins and dibenzofurans 774(1997)97

Application of accelerated solvent extraction followed by gas chromatography, high-performance liquid chromatography and gas chromatography–mass spectrometry for the determination of polycyclic aromatic hydrocarbons, chlorinated pesticides and polychlorinated dibenzo-*p*-dioxins and dibenzofurans in solid wastes 774(1997)203

Supercritical CO₂-assisted liquid extraction of polycyclic aromatic hydrocarbons and polychlorinated dibenzo-*p*-dioxins and -furans from solid matrices 777(1997)289

Application of DB-5ms gas chromatography column for the complete assignment of 2,3,7,8-substituted polychlorodibenzo-*p*-dioxins and polychlorodibenzofurans in samples from municipal waste incinerator emissions 786(1997)125

Selectivity of electron-donor- and electron-acceptor-bonded silica packing materials for hydrophobic environmental contaminants in polar and non-polar eluents 786(1997)237

Polychlorinated naphthalenes

Polychlorinated naphthalenes in groundwater samples from the Llobregat aquifer (Spain) 786(1997)135

Polycyclic aromatic hydrocarbons

see Polynuclear aromatic hydrocarbons

Polydimethylsiloxane

Fractionation of polymers using supercritical fluid extraction 785(1997)395

Polyelectrolytes

Separation of polyelectrolytes of variable compositions by free-zone capillary electrophoresis 766(1997)233

Electrophoretic behaviour of fully sulfonated polystyrenes in capillaries filled with entangled polymer solutions 772(1997)369

Polyester oligomers

Separation of polyester oligomers by gradient high-performance liquid chromatography 786(1997)85

Polyesters

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. II. Solubility effects in the analysis of polyester resins under reversed-phase conditions 761(1997)147

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions 775(1997)157

Study on the retention behaviour of low-molar-mass polystyrenes and polyesters in reversed-phase liquid chromatography by evaluation of thermodynamic parameters 790(1997)101

Polyethene

Analysis of odour and taste problems in high-density polyethene 791(1997)213

Polyethers

Liquid adsorption chromatography of polyethers. Theory and experiment 761(1997)21

Liquid chromatography of polyethers using universal detectors. V. Quantitative aspects in the analysis of low-molecular-mass poly(ethylene glycol)s and their derivatives by reversed-phase high-performance liquid chromatography with an evaporative light scattering detector 768(1997)223

Theory of liquid adsorption chromatography of macromolecules applied to polyethylene glycol and fatty alcohol ethoxylates 791(1997)21

Reversed-phase high-performance liquid chromatography of polyethers. Comparison with a theory for flexible-chain macromolecules 798(1998)187

Polyethoxylates

High-performance liquid chromatographic–fluorescence determination of aliphatic alcohol polyethoxylates and poly(ethylene glycol)s in aqueous samples 756(1996)159

Comparison of the retention behavior of polyethoxylated alcohols on porous graphitic carbon and polar as well as apolar bonded-silica phases 797(1998)83

Polyethylene

Methyl cyclohexane as a new eluting solvent for the size-exclusion chromatography of polyethylene and polypropylene at 90°C 755(1996)27

Selection of a precursor monomer for the introduction of affinity ligands onto a porous membrane by radiation-induced graft polymerization 758(1997)209

Poly(ethylene glycol)s

Analysis of basic pharmaceuticals by capillary electrophoresis in coated capillaries and on-line mass spectrometric detection 752(1996)271

High-performance liquid chromatographic–fluorescence determination of aliphatic alcohol polyethoxylates and poly(ethylene glycol)s in aqueous samples 756(1996)159

Liquid adsorption chromatography of polyethers. Theory and experiment 761(1997)21

Liquid chromatography of polyethers using universal detectors. V. Quantitative aspects in the analysis of low-molecular-mass poly(ethylene glycol)s and their derivatives by reversed-phase high-performance liquid chromatography with an evaporative light scattering detector 768(1997)223

Theory of liquid adsorption chromatography of macromolecules applied to polyethylene glycol and fatty alcohol ethoxylates 791(1997)21

Reversed-phase high-performance liquid chromatography of polyethers. Comparison with a theory for flexible-chain macromolecules 798(1998)187

Isotachophoretic focusing of strong electrolytes on the background of carrier ampholytes 798(1998)223

Analysis of ethoxylated polymers by capillary electrophoresis in UV-transparent polymer networks and by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry 800(1998)13

Determination of poly(ethylene glycol)s by both normal-phase and reversed-phase modes of high-performance liquid chromatography 800(1998)231

(Polyethylene glycol monomethyl ether)-N-methylmorpholinium hydroxides

N-(Polyethyleneglycol monomethyl ether)-N-methylmorpholinium-based background electrolytes in capillary electrophoresis 763(1997)253

Polyethyleneimine

Role of polyethyleneimine in the purification of recombinant human tumour necrosis factor beta 760(1997)165

Effect of cationic polymer on the separation of phenols by capillary electrophoresis 768(1997)161

Polyhydroxyethylmethacrylate

Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217

New sorbents for removal of heavy metal ions: diamine-glow-discharge treated polyhydroxyethylmethacrylate microspheres 773(1997)169

Polyimide

Polyimide for deactivating gas chromatography vaporising chambers and fixing packing materials 764(1997)65

Polyisobutene

Fractionation of polymers using supercritical fluid extraction 785(1997)395

Polyisoprenes

Liquid chromatography of polyisoprenes: determination of critical composition using non protic binary and acidic ternary solvent mixtures 791(1997)350

Polymers

Molded continuous poly(styrene-*co*-divinylbenzene) rod as a separation medium for the very fast separation of polymers. Comparison of the chromatographic properties of the monolithic rod with columns packed with porous and non-porous beads in high-performance liquid chromatography of polystyrenes 752(1996)59

Packings of an unidimensional regular pore structure as model packings in size-exclusion and inverse size-exclusion chromatography 753(1996)177

Degradative polymer analysis by chromatography (Review) 756(1996)1

Ruggedness testing of a size-exclusion chromatographic assay for low-molecular-mass polymers 756(1996)89

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. II. Solubility effects in the analysis of polyester resins under reversed-phase conditions 761(1997)147

Polymer characterization using on-line coupling of thermal field flow fractionation and hydrodynamic chromatography 765(1997)135

Separation of polyelectrolytes of variable compositions by free-zone capillary electrophoresis 766(1997)233

Identification of potential migrants from a vinylic organosol varnish by gas chromatography-mass spectrometry and liquid chromatography-mass spectrometry 771(1997)366

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions 775(1997)157

Fractionation of polymers using supercritical fluid extraction 785(1997)395

Quantitation in liquid chromatography of polymers: size-exclusion chromatography with dual detection 786(1997)75

Separation of polyester oligomers by gradient high-performance liquid chromatography 786(1997)85

Interface between pyrolyzer and gas chromatograph. A different configuration of pyrolysis-gas chromatography 786(1997)107

Gel permeation chromatography of polymers degrading randomly in the column. Theoretical treatment and practical aspects 786(1997)209

Interpretation of thermal field-flow fractionation experiments in a tilted channel 788(1997)121

Characterization of dextrans by size-exclusion chromatography on unmodified silica gel columns, with light-scattering detection, and capillary electrophoresis with laser-induced fluorescence detection 790(1997)93

Study on the retention behaviour of low-molar-mass polystyrenes and polyesters in reversed-phase liquid chromatography by evaluation of thermodynamic parameters 790(1997)101

Theory of liquid adsorption chromatography of macromolecules applied to polyethylene glycol and fatty alcohol ethoxylates 791(1997)21

Modification of a commercial pyrolysis probe to improve its temporal and spatial temperature profiles. Performance evaluation using model polymers 791(1997)151

Reversed-phase high-performance liquid chromatography of polyethers. Comparison with a theory for flexible-chain macromolecules 798(1998)187

Analysis of ethoxylated polymers by capillary electrophoresis in UV-transparent polymer networks and by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry 800(1998)13

Determination of poly(ethylene glycol)s by both normal-phase and reversed-phase modes of high-performance liquid chromatography 800(1998)231

Polynuclear aromatic compounds

Investigation of the separation efficiency of hydrophobic compounds in suspension electrokinetic chromatography 768(1997)320

Supercritical CO₂ assisted liquid extraction of nitroaromatic and polycyclic aromatic compounds in soil 785(1997)227

Polynuclear aromatic hydrocarbons

Separation of benzo[*a*]pyrene sulfate isomers by reversed-phase liquid chromatography 753(1996)201

Enrichment of benzo[*a*]pyrene in smoked food products and determination by high-performance liquid chromatography-fluorescence detection 753(1996)207

- Evaluation of selected filters for collection and subsequent supercritical fluid extraction of suspended solids for trace organic analysis 753(1996)279
- Experiences with packed capillary electrochromatography at ambient pressure 755(1996)165
- On-line preconcentration and high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbon-DNA adducts using copper phthalocyanine trisulfonic acid as a group-specific adsorbent 755(1996)205
- Combined use of temperature and solvent strength in reversed-phase gradient elution. IV. Selectivity for neutral (non-ionized) samples as a function of sample type and other separation conditions 756(1996)63
- Stationary phases with chemically bonded fluorene ligands: A new approach for environmental analysis of π -electron containing solutes 757(1997)3
- Temperature dependence of carotenoids on C₁₈, C₃₀ and C₃₄ bonded stationary phases 757(1997)29
- Rapid analysis of organic microcontaminants in environmental water samples by trace enrichment and liquid chromatography on a single short column 759(1997)55
- Relationship between stationary and mobile phase composition and its influence on retention factors of aromatic hydrocarbons in reversed-phase high-performance liquid chromatography 762(1997)27
- Molar enthalpy and molar volume of methylene and benzene homologues in reversed-phase liquid chromatography 762(1997)35
- Separation of hydrophobic compounds by electrokinetic chromatography with tetraalkylammonium ions 764(1997)323
- Preparation of continuous beds for electrochromatography and reversed-phase liquid chromatography of low-molecular-mass compounds 767(1997)33
- The determination of seventeen polycyclic aromatic hydrocarbons in tobacco smoke condensate 767(1997)163
- Separation of extracts from biological tissues into polycyclic aromatic hydrocarbon, polychlorinated biphenyl and polychlorinated dibenzo-*p*-dioxin/polychlorinated dibenzofuran fractions prior to analysis 771(1997)169
- Retention characteristics and selected applications of cyclic siloxane-based octadecylsilyl bonded phases in reversed-phase high-performance liquid chromatography 773(1997)53
- Synthesis and characterization of a novel wall coated capillary column for the separation of polycyclic aromatic hydrocarbons 773(1997)199
- Developments in the analysis of petroleum hydrocarbons in oils, petroleum products and oil-spill-related environmental samples by gas chromatography (Review) 774(1997)51
- Determination of coal tar and creosote constituents in the aquatic environment (Review) 774(1997)79
- Application of planar chromatography to the analysis of polycyclic aromatic hydrocarbons and their derivatives in environmental samples 774(1997)111
- Determination of polycyclic aromatic compounds and heavy metals in sludges from biological sewage treatment plants 774(1997)177
- Separation of hydroxylated polycyclic aromatic hydrocarbons by micellar electrokinetic capillary chromatography 774(1997)193
- Application of accelerated solvent extraction followed by gas chromatography, high-performance liquid chromatography and gas chromatography-mass spectrometry for the determination of polycyclic aromatic hydrocarbons, chlorinated pesticides and polychlorinated dibenzo-*p*-dioxins and dibenzofurans in solid wastes 774(1997)203
- Isolation and recovery of 9,10-dimethyl-1,2-benzanthracene from animal bedding by reversed-phase solid-phase extraction combined with high-performance liquid chromatography 774(1997)223
- Analysis of toxic wastes in tissues from aquatic species. Applications of matrix solid-phase dispersion 774(1997)287
- Occurrence and determination of organic pollutants in aerosol, precipitation, and sediment samples collected at Lake Balaton 774(1997)349
- Using systematic and comparative analytical data to identify the source of an unknown oil on contaminated birds 775(1997)251
- Simultaneous separation of metal ions and neutral organics by reversed-phase ion-pair liquid chromatography 776(1997)221
- Extraction of polycyclic aromatic hydrocarbons from polluted soils with binary and ternary supercritical phases 776(1997)233
- Supercritical fluid chromatography-atmospheric pressure chemical ionisation mass spectrometry for the analysis of hydroxy polycyclic aromatic hydrocarbons 777(1997)167
- Description of the solubilities of solids in supercritical fluids 777(1997)283
- Supercritical CO₂-assisted liquid extraction of polycyclic aromatic hydrocarbons and polychlorinated dibenzo-*p*-dioxins and -furans from solid matrices 777(1997)289

- Determination of polycyclic aromatic hydrocarbons in waters by use of supercritical fluid chromatography coupled on-line to solid-phase extraction with disks 778(1997)321
- Application of solid-phase extraction discs with a glass fiber matrix to fast determination of polycyclic aromatic hydrocarbons in water 778(1997)337
- Characterization of coal-derived pitches as precursors for advanced carbon materials by chromatographic and related techniques 778(1997)403
- Development of fast-scanning fluorescence spectra as a detection system for high-performance liquid chromatography. Determination of polycyclic aromatic hydrocarbons in water samples 779(1997)123
- Influence of sample plug width in capillary electrochromatography 779(1997)155
- Optimization of separation of some polycyclic aromatic compounds by thin-layer chromatography 779(1997)321
- Application of particles as pseudo-stationary phases in electrokinetic chromatography 780(1997)63
- Retention modeling in micellar liquid chromatography 780(1997)149
- Application of capillary electrophoresis–fluorescence line-narrowing spectroscopy for on-line spectral characterization of closely related analytes 781(1997)73
- Polyallylamine-supported pseudo-stationary phases for electrokinetic chromatography. Effect of alkyl chain length of the pseudo-stationary phase and methanol content of aqueous buffer on the separation of hydrophobic compounds 781(1997)139
- Application of mixed mobile phases and a step gradient method in capillary electrochromatography for the separation of isomeric polycyclic aromatic hydrocarbon–deoxyribonucleoside adduct mixtures prepared in vitro 781(1997)327
- Validation studies for a new supercritical fluid extraction method for the isolation of hydrophobic organic compounds from filtered suspended solids 782(1997)247
- Temperature and pressure effects on solubility in supercritical carbon dioxide and retention in supercritical fluid chromatography 785(1997)57
- Field-portable determination of polychlorinated biphenyls and polynuclear aromatic hydrocarbons in soil using supercritical fluid extraction 785(1997)205
- New, high-capacity carboxylic acid functionalized resins for solid-phase extraction of a broad range of organic compounds 786(1997)259
- Evidence for the adsorption of nitrated polycyclic aromatic hydrocarbons by tree bark 786(1997)275
- Clean-up and analysis of carbazole and acridine type polycyclic aromatic nitrogen heterocyclics in complex sample matrices 790(1997)73
- Improved detection of polycyclic aromatic compounds in complex mixtures by liquid chromatographic fractionation on poly(divinylbenzene) prior to gas chromatography–mass spectrometry. Application to the analysis of diesel particulates 791(1997)203
- Extraction of polycyclic aromatic hydrocarbons from contaminated soil using Soxhlet extraction, pressurised and atmospheric microwave-assisted extraction, supercritical fluid extraction and accelerated solvent extraction 791(1997)361
- Selectivity of polymeric and polymer-supported pseudo-stationary phases in micellar electrokinetic chromatography 792(1997)105
- Enhancement of the separation selectivity of a group of polycyclic aromatic hydrocarbons using mixed cyclodextrin-modified micellar electrokinetic chromatography 792(1997)411
- Thin-layer chromatographic separation of selected polyaromatic hydrocarbons on copper phthalocyanine dye impregnated plates 793(1998)403
- Determination of organic molecular markers in marine aerosols and sediments: one-step flash chromatography compound class fractionation and capillary gas chromatographic analysis 799(1998)215
- Quantitative structure–retention relationship studies for predicting the gas chromatography retention indices of polycyclic aromatic hydrocarbons. Quasi-length of carbon chain and pseudo-conjugated system surface 799(1998)361
- Polynuclear aromatic nitrogen heterocyclics**
Clean-up and analysis of carbazole and acridine type polycyclic aromatic nitrogen heterocyclics in complex sample matrices 790(1997)73
- Polynucleotides**
see Nucleotides and Special Issue Index
- Polyols**
Polyol-promoted adsorption of serum proteins to amphiphilic agarose-based adsorbents 764(1997)193
- Simultaneous amperometric and potentiometric detection of sugars, polyols and carboxylic acids in flow systems using copper wire electrodes 766(1997)27
- Separation and determination of pharmaceutically important polyols in dosage forms by capillary isotachopheresis 772(1997)277

Subcritical fluid chromatography of monosaccharides and polyols using silica and trimethylsilyl columns 785(1997)195

Polyoxyethylene

Composition analysis of detergents of the polyoxyethylene type: comparison of thin-layer chromatography, reversed-phase chromatography and matrix-assisted laser desorption/ionization mass spectrometry 791(1997)245

Polyoxypropylene glycol

Dual-mode high-speed counter-current chromatography: retention, resolution and examples 790(1997)17

Polypeptides

Liquid chromatography on soft packing material, under axial compression. Size-exclusion chromatography of polypeptides 773(1997)103

Polyphenolic compounds

Analysis of catechins in tea extracts by liquid chromatography–electrospray ionization mass spectrometry 794(1998)63

Polyphenols

Analysis of old brandy and oak extracts by capillary electrophoresis 768(1997)143

Selection of column and gradient elution system for the separation of catechins in green tea using high-performance liquid chromatography 793(1998)265

Liquid chromatography coupled to mass spectrometry and nuclear magnetic resonance spectroscopy for the screening of plant constituents 794(1998)299

Separation of tea polyphenols using micellar electrokinetic chromatography with diode array detection 799(1998)309

Simultaneous separation of flavanone glycosides and polymethoxylated flavones in citrus juices using liquid chromatography 800(1998)171

Polyphosphates

Single-column ion chromatography of linear polyphosphates 767(1997)107

Capillary electrophoresis of longer-chain polyphosphates 769(1997)349

Polypropylene

Methyl cyclohexane as a new eluting solvent for the size-exclusion chromatography of polyethylene and polypropylene at 90°C 755(1996)27

Poly(propylene glycol)

Reversed-phase high-performance liquid chromatography of polyethers. Comparison with a theory for flexible-chain macromolecules 798(1998)187

Polysaccharides

see also Saccharides

Fast molecular mass and size characterization of polysaccharides using asymmetrical flow field-flow fractionation–multiangle light scattering 760(1997)205

Coupled size-exclusion chromatography–anion-exchange chromatography in the analysis of poly- and oligosaccharides 763(1997)331

Calibration of a size-exclusion chromatography system using fractions with defined amylopectin unit chains 768(1997)325

Enantioseparation using selected polysaccharides as chiral buffer additives in capillary electrophoresis 773(1997)331

Regioselectively carbamoylated polysaccharides for the separation of enantiomers in high-performance liquid chromatography 786(1997)13

Characterization of dextrans by size-exclusion chromatography on unmodified silica gel columns, with light-scattering detection, and capillary electrophoresis with laser-induced fluorescence detection 790(1997)93

Enantioselectivity in chiral capillary electrophoresis with polysaccharides 792(1997)327

Observation of Ficoll charge using size-exclusion chromatography 800(1998)181

Polysorbate 80

Single step method for the accurate concentration determination of polysorbate 80 786(1997)99

Polystyrenes

Molded continuous poly(styrene-*co*-divinylbenzene) rod as a separation medium for the very fast separation of polymers. Comparison of the chromatographic properties of the monolithic rod with columns packed with porous and non-porous beads in high-performance liquid chromatography of polystyrenes 752(1996)59

N-Methyl-2-pyrrolidinone as a mobile phase in the size-exclusion chromatography of coal derivatives 758(1997)65

Study on the retention behaviour of low-molar-mass polystyrenes and polyesters in reversed-phase liquid chromatography by evaluation of thermodynamic parameters 790(1997)101

Polystyrenesulfonates

Electrophoretic behaviour of fully sulfonated polystyrenes in capillaries filled with entangled polymer solutions 772(1997)369

Polystyrene–divinylbenzene polymers

Molded continuous poly(styrene–*co*-divinylbenzene) rod as a separation medium for the very fast separation of polymers. Comparison of the chromatographic properties of the monolithic rod with columns packed with porous and non-porous beads in high-performance liquid chromatography of polystyrenes 752(1996)59

Determination of solute lipophilicity, as log *P*(octanol) and log *P*(alkane) using poly(styrene–divinylbenzene) and immobilised artificial membrane stationary phases in reversed-phase high-performance liquid chromatography 766(1997)35

Analysis of phenolic compounds in effluent by solid-phase extraction and gas chromatography–mass spectrometry with direct on-column benzylation: sensitive negative ion chemical ionisation gas chromatography–mass spectrometry detection of phenyl benzyl ethers 771(1997)203

Sulfoacylated macroporous polystyrene–divinylbenzene: a new type of cation exchanger for the analysis of multivalent metal cations 782(1997)149

Polythionates

Determination of sulfur oxyanions by ion chromatography on a silica ODS column with tetrapropylammonium salt as an ion-pairing reagent 770(1997)157

Poly(vinyl alcohol)

Analysis of basic pharmaceuticals by capillary electrophoresis in coated capillaries and on-line mass spectrometric detection 752(1996)271

Determination of polyacrylamide in polyvinyl alcohol by pyrolysis–gas chromatography with atomic emission detection 753(1996)101

Modification of polystyrenic matrices for the purification of proteins. Effect of the adsorption of poly(vinyl alcohol) on the characteristics of poly(styrene–divinylbenzene) beads for use in affinity chromatography 758(1997)53

Ion-exclusion chromatography of aliphatic carboxylic acids on a cation-exchange resin by elution with polyvinyl alcohol 770(1997)211

Poly(vinyl chloride)

Speciation of organotin compounds released from poly(vinyl chloride) at increased temperature by gas chromatography with atomic emission detection 775(1997)295

Porous graphite carbon

Anomalous retention behaviour of peptides on porous graphitized carbon column 776(1997)147

Potassium

Determination of Na⁺, K⁺, Mg²⁺ and Ca²⁺ in mixtures of seawater and formation water by capillary electrophoresis 767(1997)311

Potassium salts

Chromatographic ion-exchange and charge-partner reactions. New mechanisms concerning ion–ion interactions of inorganic compounds in Sephadex G-15 columns (Review) 789(1997)21

Praziquantel

Determination of fenbendazole, praziquantel and pyrantel pamoate in dog plasma by high-performance liquid chromatography 797(1998)237

Prazosin

Chiral bioanalysis by normal phase high-performance liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 759(1997)65

Enantioselective determination of terazosin in human plasma by normal phase high-performance liquid chromatography–electrospray mass spectrometry 794(1998)45

High-performance liquid chromatographic method with coulometric detection for the determination of buspirone in human plasma by means of a column-switching technique 797(1998)221

Prednisolone

Liquid chromatography–mass spectrometry with on-line solid-phase extraction by a restricted-access C₁₈ precolumn for direct plasma and urine injection 762(1997)193

Primisulfuron

Analysis of primisulfuron and triasulfuron in water and soil samples by micellar electrokinetic capillary chromatography 766(1997)225

Procainamide

Direct determination of procainamide and N-acetylprocainamide by capillary zone electrophoresis in pharmaceutical formulations and urine 772(1997)271

Prochloraz

Simplified determination of combined residues of prochloraz and its metabolites in vegetable, fruit and wheat samples by gas chromatography 765(1997)127

Procymidone

Resolution of folpet, procymidone and triazophos in high-performance liquid chromatography–diode array detection by using partial least squares calibration to cross-sections of spectrochromatograms 778(1997)183

Profens

Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309

Separation of profen enantiomers by capillary electrophoresis using cyclodextrins as chiral selectors 793(1998)165

Chiral and nonchiral determination of ketoprofen in pharmaceuticals by capillary zone electrophoresis 799(1998)301

Progesterone

Capillary electrochromatography of steroids. Increased sensitivity by on-line concentration and comparison with high-performance liquid chromatography 798(1998)259

Proline

Computerized optimization of selectivity for direct capillary gas chromatographic multicomponent separations of enantiomers 779(1997)253

Prolintane

Direct determination of prolintane and its metabolite oxoprolintane in human urine by capillary zone electrophoresis and β -cyclodextrin-modified micellar electrokinetic chromatography 778(1997)355

Prometone

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Propanediol fatty acid esters

Preparation and liquid chromatographic analysis of propanediol fatty acid esters 779(1997)337

Propanol

Description of the retention behaviour in micellar liquid chromatography as a function of pH, surfactant and modifier concentration 769(1997)155

Propranolol

Direct injection of large volumes of plasma/serum on a new biocompatible extraction column for the determination of atenolol, propranolol and ibuprofen. Mechanisms for the improvement of chromatographic performance 797(1998)251

Fluorescence detection in packed-column supercritical fluid chromatographic separations 798(1998)203

Propazine

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Propiomazine

Effect of organic modifiers on retention and enantiomeric separations by capillary electrophoresis with human serum albumin as a chiral selector in solution 766(1997)237

Propoxur

Resolution and quantitative determination of coeluted pesticide mixtures in liquid chromatography–thermospray mass spectrometry by multivariate curve resolution 795(1998)3

N(O,S)-Propoxycarbonyl propyl ester

Quantitation of total homocysteine in human plasma by derivatization to its N(O,S)-propoxycarbonyl propyl ester and gas chromatography–mass spectrometry analysis 776(1997)342

Propranolol

Improved chiral separation using achiral modifiers in cyclodextrin modified capillary zone electrophoresis 773(1997)321

Enantiomeric separation of propranolol and selected metabolites by using capillary electrophoresis with hydroxypropyl- β -cyclodextrin as chiral selector 793(1998)357

Propranolol hydrochloride

Influence of the sample volume and the position of the electrode and the capillary-end in the sample vial on the electrokinetic injection in capillary electrophoresis 788(1997)212

Prostaglandins

Analysis of prostaglandins by micellar electrokinetic capillary chromatography 761(1997)277

Identification of eicosanoids in the red algae, *Gracilaria asiatica*, using high-performance liquid chromatography and electrospray ionization mass spectrometry 795(1998)227

Proteases

Capillary electrophoresis in the assay of the hydrolysis of glycine-containing peptides by a protease from *Pseudomonas aeruginosa* 776(1997)133

Protein A

Isolation of monoclonal antibodies from cell containing hybridoma broth using a protein A coated adsorbent in expanded beds 752(1996)111

Effects of fused tryptophan rich peptides to a recombinant protein A domain on the partitioning in polyethylene glycol–dextran and Ucon–dextran aqueous two-phase systems 756(1996)107

Accelerated recombinant protein purification process development. Automated, robotics-based integration of chromatographic purification and analysis 798(1998)73

Proteinase inhibitor

Chromatographic purification of human α_1 proteinase inhibitor from dissolved Cohn fraction IV-1 paste 800(1998)207

Protein G

Fast isolation of protein receptors from streptococci G by means of macroporous affinity discs 798(1998)65

Proteins

see also Antibodies, Enzymes, Monoclonal antibodies, Immunoglobulins and Special Issue Index

pH-Zone-refining countercurrent chromatography 753(1996)1

- Cyanocarbons as ligands for electron donor acceptor chromatography of human serum proteins 753(1996)57
- Multiple peaks induced by domain-specific binding of fibrinogen in anion-exchange high-performance liquid chromatography 753(1996)63
- Purification of insulin-like growth factor-I and related proteins using underivatized silica 753(1996)73
- Peculiarities of gradient ion-exchange high-performance liquid chromatography of proteins 753(1996)217
- Hydrophobic-interaction chromatography of proteins on continuous beds derivatized with isopropyl groups 753(1996)227
- Dextran-coated silica and its behavior in high-performance size-exclusion chromatography 755(1996)1
- Combined use of temperature and solvent strength in reversed-phase gradient elution. III. Selectivity for ionizable samples as a function of sample type and pH 756(1996)51
- Characterization and impact of the temporal behavior of the electroosmotic flow in capillary isoelectric focusing with electroosmotic zone displacement 756(1996)219
- Adsorption of proteins to fused-silica capillaries as probed by atomic force microscopy 756(1996)233
- Preparation of linear polyacrylamide gel step gradients for capillary electrophoresis 756(1996)245
- Capillary isoelectric focusing: the problem of protein solubility 757(1997)237
- Capillary electrophoretic separation of acidic and basic proteins in the presence of cationic and anionic fluorosurfactants 757(1997)255
- Modification of polystyrenic matrices for the purification of proteins. Effect of the adsorption of poly(vinyl alcohol) on the characteristics of poly(styrene-divinylbenzene) beads for use in affinity chromatography 758(1997)53
- Rapid separation of soybean globulins by reversed-phase high-performance liquid chromatography 758(1997)75
- Selection of a precursor monomer for the introduction of affinity ligands onto a porous membrane by radiation-induced graft polymerization 758(1997)209
- Selectivity enhancement for free zone capillary electrophoresis using conventional ion-pairing agents as complexing additives 758(1997)293
- Semipreparative isolation of collagen types I, II, III and V by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and electroelution 758(1997)313
- Post-capillary affinity detection of protein microheterogeneity in capillary zone electrophoresis 759(1997)139
- Analysis of recombinant human erythropoietin in drug formulations by high-performance capillary electrophoresis 759(1997)177
- Combination of the steric mass action and non-ideal surface solution models for overload protein ion-exchange chromatography 760(1997)89
- Synergistic effects in the processes of protein multicomponent sorption 760(1997)105
- Comparison of anion-exchange and hydroxyapatite displacement chromatography for the isolation of whey proteins 760(1997)125
- Refining the description of protein chromatography 760(1997)139
- Performance comparison of low-pressure ion-exchange chromatography media for protein separation 760(1997)151
- Role of polyethyleneimine in the purification of recombinant human tumour necrosis factor beta 760(1997)165
- Estimation of pore diameter for intraparticle fluid flow in bidisperse porous chromatographic particles 761(1997)35
- Characterisation of silica-based heparin affinity sorbents from equilibrium binding studies on plasma fractions containing thrombin 762(1997)113
- Tryptophan analysis in peptides and proteins, mainly by liquid chromatography 763(1997)1
- Role of high-performance liquid chromatographic protein analysis in developing fermentation processes for recombinant human growth hormone, relaxin, antibody fragments and lymphotoxin 763(1997)31
- Preparative desalting of bovine serum albumin by continuous annular chromatography 763(1997)49
- Ion-exchange high-performance liquid chromatographic separation of protein variants and isoforms on MCI GEL ProtEx stationary phases 763(1997)57
- Use of reversed-phase high-performance liquid chromatography on polystyrene-divinylbenzene columns for the rapid separation and purification of acid-soluble nuclear proteins 763(1997)65
- Comparative studies on the isothermal characteristics of proteins adsorbed under batch equilibrium conditions to ion-exchange, immobilised metal ion affinity and dye affinity matrices with different ionic strength and temperature conditions 763(1997)71

- Complete structure elucidation of a globular protein by particle beam liquid chromatography–Fourier transform infrared spectrometry and electrospray liquid chromatography–mass spectrometry. Sequence and conformation of β -lactoglobulin 763(1997)91
- Identification of proteins by capillary electrophoresis–tandem mass spectrometry. Evaluation of an on-line solid-phase extraction device 763(1997)295
- Separation of proteins and peptides by capillary electrochromatography in diol- and octadecyl-modified etched capillaries 763(1997)307
- Determination of molecular-mass distribution of food-grade protein hydrolyzates by size-exclusion chromatography and chemiluminescent nitrogen detection 763(1997)323
- Ethanol–sodium chloride–phosphate mobile phase for size-exclusion chromatography of poly(ethylene glycol) modified proteins 763(1997)337
- Mass transfer limitations in protein separations using ion-exchange membranes 764(1997)3
- Comment on the theory of protein adsorption on a biospecific rigid matrix 764(1997)21
- Immobilized metal affinity chromatography of human growth hormone. Effect of ligand density 764(1997)27
- Concentrating protein samples for sodium dodecyl sulphate–polyacrylamide gel electrophoresis and isoelectric focusing using protein-blotting membranes 764(1997)143
- Polyol-promoted adsorption of serum proteins to amphiphilic agarose-based adsorbents 764(1997)193
- High sample throughput phosphoamino acid analysis of proteins separated by one- and two-dimensional gel electrophoresis 764(1997)201
- Protein retention in hydrophobic interaction chromatography: modeling variation with buffer ionic strength and column hydrophobicity 766(1997)1
- Screening of a large number of dyes for the separation of human immunoglobulin G2 from the other immunoglobulin G subclasses. Immunoglobulin G2 enrichment on immobilized Procion Yellow HE-4R 766(1997)49
- Interaction between tryptophan residues and hydrophobically modified dextran. Effect on partitioning of peptides and proteins in aqueous two-phase systems 766(1997)99
- Studies of the hydrodynamic volume changes that occur during refolding of lysozyme using size-exclusion chromatography 766(1997)109
- Application of solid-phase extraction to the preconcentration of metallothionein and metals from physiological fluids 766(1997)121
- Conformational transformations of the fusion protein BR96 sFv-PE40 as monitored by micellar electrokinetic capillary chromatography and circular dichroism 766(1997)205
- Site accessibility and the pH dependence of the saturation capacity of a highly cross-linked matrix. Immobilized metal affinity chromatography of bovine serum albumin on Chelating Superose 767(1997)11
- Capillary isoelectric focusing with anionic coated capillaries 767(1997)231
- Study of haptoglobin–hemoglobin complexes by titration curves, capillary electrophoresis and capillary isoelectric focusing 767(1997)255
- Critical conditions for separating the microheterogeneous components of glycoproteins by capillary electrophoresis 768(1997)39
- Capillary electrophoretic analysis of genetic variants of milk proteins from different species 768(1997)47
- Evaluation of phytic acid as a buffer additive for the separation of proteins in capillary electrophoresis 768(1997)307
- Experimental and numerical studies of the chromatofocusing of dilute proteins using retained pH gradients formed on a strong-base anion-exchange column 769(1997)129
- Application of a β -cyclodextrin sulfate-immobilized precolumn to selective on-line enrichment and separation of heparin-binding proteins by column-switching high-performance liquid chromatography 769(1997)209
- Enantioselectivity of bovine serum albumin-bonded columns produced with isolated protein fragments. II. Characterization of protein fragments and chiral binding sites 769(1997)215
- Monitoring the refolding pathway for a large multimeric protein using capillary zone electrophoresis 769(1997)315
- Productivity and operating regimes in protein chromatography using low-molecular-mass displacers 771(1997)9
- Evaluation of pH gradient formation of carrier ampholytes with synthesized isoelectric point markers in capillary isoelectric focusing 772(1997)137
- Strategies to improve performance of capillary isoelectric focusing 772(1997)145
- Assessment of heat-induced denaturation of whey proteins 772(1997)231

- Characterisation of human serum albumin heterogeneity by capillary zone electrophoresis and electrospray ionization mass spectrometry 772(1997)235
- Identification of α_1 -acid glycoprotein (orosomucoid) in human synovial fluid by capillary electrophoresis 772(1997)305
- Preparation and use of ion-exchange chromatographic supports based on perfluoropolymers 773(1997)73
- Purification of chaperonins from thermophilic bacteria and archaea 773(1997)131
- Purification by reflux electrophoresis of whey proteins and of a recombinant protein expressed in *Dictyostelium discoideum* 773(1997)299
- Rigid porous polyacrylamide-based monolithic columns containing butyl methacrylate as a separation medium for the rapid hydrophobic interaction chromatography of proteins 775(1997)65
- Rapid assay for the analysis of class III metallothioneins by capillary zone electrophoresis in crude extracts of microalgal cells exposed to cadmium 775(1997)339
- Model discrimination and estimation of the intraparticle mass transfer parameters for the adsorption of bovine serum albumin onto porous adsorbent particles by the use of experimental frontal analysis data 776(1997)3
- Kinetic analysis of the interaction between the monoclonal antibody A33 and its colonic epithelial antigen by the use of an optical biosensor. A comparison of immobilisation strategies 776(1997)15
- Modification of polystyrenic matrices for the purification of proteins. II. Effect of the degree of glutaraldehyde-poly(vinyl alcohol) crosslinking on various dye ligand chromatography systems 776(1997)55
- Modification of polystyrenic matrices for the purification of proteins. III. Effects of poly(vinyl alcohol) modification on the characteristics of protein adsorption on conventional and perfusion polystyrenic matrices 776(1997)65
- Effect of temperature upon the chromatography of proteins on porous glass, chemically coated with N-isopropylacrylamide copolymer 776(1997)75
- D-Glucose, forskolin and cytochalasin B affinities for the glucose transporter Glut1. Study of pH and reconstitution effects by biomembrane affinity chromatography 776(1997)81
- Chromatographic method involving inductively coupled plasma atomic emission spectrometric detection for the study of metal-protein complexes 776(1997)93
- Size-exclusion chromatographic reconstitution of the bovine brain benzodiazepine receptor. Effects of lipid environment on the binding characteristics 776(1997)101
- Two-dimensional electrophoretic analysis of vesicular and micellar proteins of gallbladder bile 776(1997)109
- Matrix-assisted laser desorption/ionisation mass spectrometry guided purification of human guanylin from blood ultrafiltrate 776(1997)139
- Fluidization characteristics of and protein adsorption on fluoride-modified porous zirconium oxide particles 776(1997)179
- Superporous agarose as an affinity chromatography support 776(1997)197
- Probing protein structure using biochemical and biophysical methods. Proteolysis, matrix-assisted laser desorption/ionization mass spectrometry, high-performance liquid chromatography and size-exclusion chromatography of p21^{Waf1/Cip1/Sdi1} 777(1997)23
- Use of microdialysis for the on-line coupling of capillary isoelectric focusing with electrospray mass spectrometry 777(1997)31
- Multiple peaks in high-performance liquid chromatography of proteins. β -Lactoglobulins eluted in a hydrophobic interaction chromatography system 778(1997)43
- Chromatographic purification of biliproteins from *Spirulina platensis*. High-performance liquid chromatographic separation of their α and β subunits 778(1997)441
- Microcolumn liquid chromatography: instrumentation, detection and applications 779(1997)1
- In situ analysis of protein chromatography and column efficiency using magnetic resonance imaging 779(1997)73
- Rapid resolution by reversed-phase high-performance liquid chromatography of the thylakoid membrane proteins of the photosystem II light-harvesting complex 779(1997)131
- Recycling electrophoretic separations: modeling of isotachopheresis and isoelectric focusing 779(1997)165
- Micelle-mediated extraction 780(1997)229
- Micellar electrokinetic chromatography of proteins 780(1997)285
- Effects of electroosmotic flow on zone mobilization in capillary isoelectric focusing 781(1997)113
- Quaternary structural analysis of nucleoside diphosphate kinases using capillary electrophoresis 781(1997)251

- Microchip-based capillary electrophoresis of human serum proteins 781(1997)271
- Protein band spreading in capillary zone electrophoresis. Effects of sample zone length and presence of polymer 781(1997)277
- Conventional capillary electrophoresis in comparison with short-capillary capillary electrophoresis and microfabricated glass chip capillary electrophoresis for the analysis of fluorescein isothiocyanate anti-human immunoglobulin G 781(1997)287
- Characterization of metal complexes of pharmaceutical interest by capillary electrophoresis with element sensitive detection 781(1997)407
- Detection of isoforms and isomers of rattlesnake myotoxins by capillary electrophoresis and matrix-assisted laser desorption time-of-flight mass spectrometry 781(1997)429
- Application of hydrophilic-interaction liquid chromatography to the separation of phosphorylated H1 histones 782(1997)55
- Analysis of recombinant human growth hormone directly in osmotic shock fluids 782(1997)199
- Protein analysis by packed capillary liquid chromatography with electrospray ionization and time-of-flight mass spectrometry detection 786(1997)67
- Single step method for the accurate concentration determination of polysorbate 80 786(1997)99
- Selection of optimum affinity tags from a phage-displayed peptide library. Application to immobilized copper(II) affinity chromatography 787(1997)91
- Characterization of antigen-antibody complexes by size-exclusion chromatography coupled with low-angle light-scattering photometry and viscometry 787(1997)101
- Temperature programming and gradient elution in reversed-phase chromatography with packed capillary columns 788(1997)51
- Capillary zone electrophoresis with fluid-impervious polymer tubing inside a fused-silica capillary 788(1997)155
- Evaluation of quail egg white riboflavin binding protein as a chiral selector in high-performance liquid chromatography and capillary electrophoresis 790(1997)47
- On-line post-capillary affinity detection of immunoglobulin G subclasses and monoclonal antibody variants for capillary electrophoresis 790(1997)207
- Modeling column regeneration effects on ion-exchange chromatography 791(1997)99
- Selectivity in capillary electrophoresis: the use of proteins 792(1997)349
- Development of liquid chromatography-mass spectrometry using continuous-flow matrix-assisted laser desorption ionization time-of-flight mass spectrometry 794(1998)367
- Non-linear elution effects in split-peak chromatography. II. Role of ligand heterogeneity in solute binding to columns with adsorption-limited kinetics 795(1998)185
- Preparation of magnetic immobilized metal affinity separation media and its use in the isolation of proteins 795(1998)211
- Effect of the hydrolysis method on the determination of the amino acid composition of proteins 795(1998)263
- Bovine whey fractionation based on cation-exchange chromatography 795(1998)277
- Characterisation of ANX Sepharose 4 Fast Flow media 796(1998)129
- Importance of heat of adsorption in modeling protein equilibria for overloaded chromatography 796(1998)141
- Frontal chromatography of proteins. Effect of axial dispersion on column performance 796(1998)157
- Micropreparation of tissue collagenase fragments of type I collagen in the form of surfactant-peptide complexes and their identification by capillary electrophoresis and partial sequencing 796(1998)181
- Electrostatic effects on protein partitioning in size-exclusion chromatography and membrane ultrafiltration 796(1998)229
- Preparation of chromatographic matrices by free radical addition ligand attachment to allyl groups 796(1998)273
- Size-exclusion chromatography of plasma proteins with high molecular masses 796(1998)289
- Effect of porous structure of macroporous polymer supports on resolution in high-performance membrane chromatography of proteins 798(1998)55
- Fast isolation of protein receptors from streptococci G by means of macroporous affinity discs 798(1998)65
- Accelerated recombinant protein purification process development. Automated, robotics-based integration of chromatographic purification and analysis 798(1998)73
- Rapid reversed-phase high-performance liquid chromatography method for quantitation, at high pH, of the recombinant apolipo-protein A-IMilano in *Escherichia coli* fermentation broth 798(1998)83
- Micro-sequencing strategies for the human A33 antigen, a novel surface glycoprotein of human gastrointestinal epithelium 798(1998)91

Dye–ligand and metal chelate poly(2-hydroxyethylmethacrylate) membranes for affinity separation of proteins 799(1998)83

Zn²⁺-selective purification of recombinant proteins from the milk of transgenic animals 799(1998)125

Affinity chromatography of thiol ester-containing proteins 799(1998)139

Sensitive high-resolution analysis of biological molecules by capillary zone electrophoresis coupled with reflecting time-of-flight mass spectrometry 800(1998)1

Molecular characterization of metal-binding polypeptide domains by electrospray ionization mass spectrometry and metal chelate affinity chromatography 800(1998)29

Nanogram scale separations of proteins using capillary high-performance liquid chromatography with fully-automated on-line microfraction collection followed by matrix-assisted laser desorption ionisation time-of-flight mass spectrometry, protein sequencing and Western blot analysis 800(1998)83

Comparison of the protein adsorption selectivity of salt-promoted agarose-based adsorbents. Hydrophobic, thiophilic and electron donor–acceptor adsorbents 800(1998)151

Chromatographic purification of human α_1 proteinase inhibitor from dissolved Cohn fraction IV-1 paste 800(1998)207

Validation of a capillary isoelectric focusing method for the recombinant monoclonal antibody C2B8 800(1998)355

Proteoliposomes

Size-exclusion chromatographic reconstitution of the bovine brain benzodiazepine receptor. Effects of lipid environment on the binding characteristics 776(1997)101

Protoberberine alkaloids

Liquid chromatographic–electrospray mass spectrometric analysis of *Coptidis Rhizoma* 755(1996)19

Protolichesterinic acid

Quantitative determination of protolichesterinic- and fumarprotocetraric acids in *Cetraria islandica* by high-performance liquid chromatography 757(1997)303

Pseudoephedrine

Development and validation of a high-performance liquid chromatography method for the determination of cold relief ingredients in chewing gum 775(1997)179

Psoralens

Retention of some components in supercritical fluid chromatography and application to bergamot peel oil fractionation 771(1997)241

Pullulans

Fast molecular mass and size characterization of polysaccharides using asymmetrical flow field-flow fractionation–multiangle light scattering 760(1997)205

High-performance capillary electrophoresis of hyaluronic acid: determination of its amount and molecular mass 768(1997)295

Observation of Ficoll charge using size-exclusion chromatography 800(1998)181

Purines

Simultaneous determination of purine bases, ribonucleosides and ribonucleotides by capillary electrophoresis–electrochemistry with a copper electrode 760(1997)227

Purpureaglycoside A

Isolation of cardenolides from a Brazilian cultivar of *Digitalis lanata* by rotation locular counter-current chromatography 756(1996)287

Putrescine

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

High-performance liquid chromatographic determination of polyamines in milk as their 9-fluorenylmethoxycarbonyl derivatives using a column-switching technique 791(1997)79

Use of 2-chloroethylnitrosourea, a new type of pre-column derivatizing agent for the measurement of biogenic amines, by high-performance liquid chromatography with ultraviolet detection 795(1998)239

Pyoverdins

Gas chromatography–chemical ionization mass spectrometry in amino acid analysis of pyoverdins 787(1997)195

Pyranocoumarins

Determination of furanochromones and pyranocoumarins in drugs and *Ammi visnaga* fruits by combined solid-phase extraction–high-performance liquid chromatography and thin-layer chromatography–high-performance liquid chromatography 797(1998)305

Pyrantel pamoate

Determination of fenbendazole, praziquantel and pyrantel pamoate in dog plasma by high-performance liquid chromatography 797(1998)237

Pyrazines

Supercritical fluid extraction of pyrazines in roasted cocoa beans. Effect of pod storage period 785(1997)361

Pyrazoles

Retention modeling in micellar liquid chromatography 780(1997)149

2-(1-Pyrenyl)ethylsilyl silica

Selectivity of electron-donor- and electron-acceptor-bonded silica packing materials for hydrophobic environmental contaminants in polar and non-polar eluents 786(1997)237

Pyrethrins

Direct determination of pyrethrins in pyrethrum extracts by reversed-phase high-performance liquid chromatography with diode-array detection 766(1997)277

In situ transesterification of the natural pyrethrins to methyl esters by heterogeneous catalysis using a supercritical fluid extraction system and detection by gas chromatography–mass spectrometry 785(1997)263

Capillary supercritical fluid chromatography of pyrethrins and pyrethroids with positive pressure and negative temperature gradients 799(1998)265

Pyrethroids

Enantiomeric resolution of pyrethroids by high-performance liquid chromatography with diode-laser polarimetric detection 754(1996)97

Analytical method development for the determination of synthetic pyrethroid insecticides in soil by gas chromatography–mass spectrometry operated in negative-ion chemical-ionization mode 754(1996)235

Retention profiles of some commercial pesticides, pyrethroid and acaricide residues and their application to tomato and parsley plants 760(1997)179

Clean-up of aqueous acetone vegetable extracts by solid-matrix partition for pyrethroid residue determination by gas chromatography–electron-capture detection 765(1997)39

Selective extraction of pyrethroid pesticide residues from milk by solid-matrix dispersion 765(1997)51

Separation and analysis of the diastereomers and enantiomers of cypermethrin and related compounds 777(1997)363

Capillary supercritical fluid chromatography of pyrethrins and pyrethroids with positive pressure and negative temperature gradients 799(1998)265

Pyridine

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Synthesis of rhodium(III)–pyridine complexes: an electrophoretic and chromatographic study 769(1997)325

Pyridinecarboxylic acids

Selectivity of alkylamide bonded-phases with respect to organic acids under reversed-phase conditions 762(1997)147

Capillary electrophoretic determination of inorganic and organic anions using 2,6-pyridinedicarboxylic acid: effect of electrolyte's complexing ability 767(1997)223

Pyridinoline

Automated on-line solid-phase extraction and high-performance liquid chromatographic analysis of total and free pyridinium crosslinks in serum 798(1998)159

Pyridoxal

Determination of vitamin B₆ (pyridoxamine, pyridoxal and pyridoxine) in pork meat and pork meat products by liquid chromatography 795(1998)383

Pyridoxamine

Determination of vitamin B₆ (pyridoxamine, pyridoxal and pyridoxine) in pork meat and pork meat products by liquid chromatography 795(1998)383

Pyridoxine

Determination of vitamin B₆ (pyridoxamine, pyridoxal and pyridoxine) in pork meat and pork meat products by liquid chromatography 795(1998)383

Pyridylaminosugar chains

Introduction of a new scale into reversed-phase high-performance liquid chromatography of pyridylamino sugar chains for structural assignment 800(1998)187

Pyridylazonaphtholsulfonic acid

Ion-pair reversed-phase high-performance liquid chromatographic separation and determination of ruthenium, rhodium, cobalt and copper as chelates with 1-(2-pyridylazo)-2-naphthol-6-sulfonic acid 789(1997)485

Pyridylazoresorcinol

Modification of 4-(2-pyridylazo)-resorcinol postcolumn reagent selectivity through competitive equilibria with chelating ligands 770(1997)69

Influence of cationic polymers on separation selectivity in kinetic differentiation mode capillary electrophoresis of metal–4-(2-pyridylazo)resorcinolato chelates 772(1997)339

Use of cationic polymers for the simultaneous determination of inorganic anions and metal–4-(2-pyridylazo)resorcinolato chelates in kinetic differentiation-mode capillary electrophoresis 776(1997)329

 α -(4-Pyridyl-1-oxide)-*N*-*tert*.-butylnitron

Quantitative detection of reduced, radical and oxidized forms of α -(4-pyridyl-1-oxide)-*N*-*tert*.-butylnitron radical adduct using high-performance liquid chromatography with electrochemical detection 753(1996)235

Pyromellitate

Sulfur speciation by capillary electrophoresis with indirect spectrophotometric detection: In search of a suitable carrier electrolyte to maximize sensitivity 770(1997)311

Pyromellitic acid

Single-column ion chromatography of linear polyphosphates 767(1997)107

Pyrrolidinedithiocarbamates

Reversed-phase high-performance liquid chromatography of pyrrolidinedithiocarbamate complexes of mercuric species using amperometric and coulometric detection 761(1997)139

Pyrrolidinones

Chiral resolution of 4-substituted pyrrolidin-2-ones using high-performance liquid chromatography on cellulose and amylose chiral stationary phases 771(1997)343

Q**Quaternary ammonium compounds**

Analysis of cationic surfactants by capillary electrophoresis 757(1997)275

Simultaneous determination of bromide and iodide ions by capillary isotachopheresis using quaternary ammonium salts 760(1997)253

Analysis of benzalkonium chloride in the effluent from European hospitals by solid-phase extraction and high-performance liquid chromatography with post-column ion-pairing and fluorescence detection 774(1997)281

Peak shape distortions during capillary electrophoretic separations of multicomponent samples in two co-ion buffers 781(1997)107

Capillary electrophoresis of cationic and anionic surfactants with indirect conductivity detection 781(1997)533

Analysis of tetramine in sea snails by capillary electrophoresis–tandem mass spectrometry 781(1997)555

Separation of quaternary ammonium herbicides by capillary electrophoresis with indirect UV detection 782(1997)289

Separation of neutral substances by non-aqueous capillary electrophoresis through interactions with cationic additives 792(1997)475

Determination of antibacterial quaternary ammonium compounds in lozenges by capillary electrophoresis 798(1998)335

Quinazoline-(3H,1H)-4-on-2-thiones

Chiral resolution of diols by capillary electrophoresis using borate–cyclodextrin complexation 761(1997)269

Quinoline

Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37

Quinolinesulfonamides

Separation of primary and secondary amines as their sulfonamide derivatives by reversed-phase high-performance liquid chromatography 763(1997)173

Quinolinol

Fluorometric determination of serum and urinary aluminium with 8-quinolinol by kinetic-differentiation-mode micellar chromatography 789(1997)361

Quinolones

Determination of the alkaloids in *Evodiae fructus* by capillary electrophoresis 755(1996)113

Optical resolution of new quinolone drugs by capillary electrophoresis with ligand-exchange and host–guest interactions 760(1997)235

Optimization of capillary electrophoretic separation of quinolone antibacterials using the overlapping resolution mapping scheme 766(1997)215

Quinolonic acids

Determination of the chemotherapeutic quinolonic and cinolonic derivatives in urine by high-performance liquid chromatography with ultraviolet and fluorescence detection in series 787(1997)119

2-Quinolylazo compounds

Use of 2-quinolylazo compounds for the reversed-phase high-performance liquid chromatographic separation of metal ions 789(1997)491

Quinuclidinol

Application of liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry, and tandem mass spectrometry, to the analysis and identification of degradation products of chemical warfare agents 759(1997)79

R

Ranitidine

Optimization of the capillary electrophoresis separation of ranitidine and related compounds 766(1997)245

Ranitidine hydrochloride

Optimisation, validation and application of a capillary electrophoresis method for the determination of ranitidine hydrochloride and related substances 798(1998)297

Rare earth elements

Enrichment separation of rare earth elements by high-speed countercurrent chromatography in a multilayer coiled column 789(1997)381

Rare earth ions

Simultaneous separation of yttrium and lanthanide ions by isotachopheresis 772(1997)357

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

Recombinant proteins

see also Proteins

Analysis of recombinant human erythropoietin in drug formulations by high-performance capillary electrophoresis 759(1997)177

Purification by reflux electrophoresis of whey proteins and of a recombinant protein expressed in *Dictyostelium discoideum* 773(1997)299

Accelerated recombinant protein purification process development. Automated, robotics-based integration of chromatographic purification and analysis 798(1998)73

Relaxin

Role of high-performance liquid chromatographic protein analysis in developing fermentation processes for recombinant human growth hormone, relaxin, antibody fragments and lymphotoxin 763(1997)31

Resorcarene-octacarboxylic acids

Highly efficient separation of amines by electrokinetic chromatography using resorcarene-octacarboxylic acids as pseudostationary phases 792(1997)143

Resorcinol

Fitting competitive adsorption isotherms to the experimental distribution data in reversed-phase systems 762(1997)3

On-line concentration of neutral analytes for micellar electrokinetic chromatography. IV. Field-enhanced sample injection 798(1998)251

Retinoic acid

Comparison of the separation of *cis/trans* isomers of tretinoin with different stationary phases by liquid chromatography–nuclear magnetic resonance coupling 765(1997)207

Retinoids

Separation of retinoids by micellar electrokinetic capillary chromatography 761(1997)307

Retinol

see also Vitamins

Concurrent liquid chromatographic separation and photodiode array detection of retinol, tocopherols, all-*trans*- α -carotene, all-*trans*- β -carotene and the mono-*cis* isomers of β -carotene in extracts of human plasma 787(1997)111

Development of a method for quantitation of retinol and retinyl palmitate in human serum using high-performance liquid chromatography–atmospheric pressure chemical ionization–mass spectrometry 794(1998)245

Retinyl palmitate

Development of a method for quantitation of retinol and retinyl palmitate in human serum using high-performance liquid chromatography–atmospheric pressure chemical ionization–mass spectrometry 794(1998)245

Rhodium(III)

Synthesis of rhodium(III)–pyridine complexes: an electrophoretic and chromatographic study 769(1997)325

Ribonuclease

Synergistic effects in the processes of protein multicomponent sorption 760(1997)105

Ribonucleosides

Simultaneous determination of purine bases, ribonucleosides and ribonucleotides by capillary electrophoresis–electrochemistry with a copper electrode 760(1997)227

Ribonucleotides

Simultaneous determination of purine bases, ribonucleosides and ribonucleotides by capillary electrophoresis–electrochemistry with a copper electrode 760(1997)227

Simple and rapid chromatographic method for the separation of major classes of ribosomal ribomononucleotides 769(1997)225

Separation of ribonucleotides by capillary electrophoresis with multifunctional electrophoretic media of phosphate ammonium salts 779(1997)215

Optimization for the separation of ribonucleotides by capillary electrophoresis at high pH 792(1997)67

Erratum to "Separation of ribonucleotides by capillary electrophoresis with multifunctional electrophoretic media of phosphate ammonium salts". [J. Chromatogr. A, 779 (1997) 215–225] 795(1998)400

Rifampicin

Thermodynamic study of the retention behaviour of selected macrocycles using reversed-phase high-performance thin-layer chromatography plates and methanol–water mobile phases 787(1997)227

Rifamycins

Thermodynamic study of the retention behaviour of selected macrocycles using reversed-phase high-performance thin-layer chromatography plates and methanol–water mobile phases 787(1997)227

Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309

Ristocetin A

Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309

RNA

Simple and rapid chromatographic method for the separation of major classes of ribosomal ribomonucleotides 769(1997)225

Method for detection of epsilon-secondary structure in the precore region of human hepatitis B virus DNA using a fluorescence-based polymerase chain reaction–single-strand-conformation polymorphism technique with capillary electrophoresis 781(1997)307

Ropivacaine

Capillary electrochromatography with molecular imprint-based selectivity for enantiomer separation of local anaesthetics 792(1997)401

S

Sabinene

Decomposition of α -pinene and sabinene on solid sorbents, Tenax TA and Carboxen 786(1997)185

Saccharides

see also Carbohydrates, Disaccharides, Monosaccharides, Oligosaccharides, Polysaccharides and Sugars

Coupled size-exclusion chromatography–anion-exchange chromatography in the analysis of poly- and oligosaccharides 763(1997)331

Ion-pair high-performance liquid chromatography for determining disaccharide composition in heparin and heparan sulphate 765(1997)169

Separation of acyclic derivatives of sugar enantiomers using gas chromatography on chiral stationary phases of permethylated α -, β - and γ -cyclodextrin 767(1997)325

Quantitative analysis of amylopectin unit chains by means of high-performance anion-exchange chromatography with pulsed amperometric detection 800(1998)199

Saccharin

Separation and determination of four artificial sweeteners and citric acid by high-performance anion-exchange chromatography 771(1997)135

Saccharose

Quantitative evaluation of chromatograms from totally illuminated thin-layer chromatographic plates 768(1997)329

Saikosaponins

Determination of saikosaponins by micellar electrokinetic capillary chromatography 759(1997)193

Salbutamol

Rapid pharmacokinetic screening of salbutamol in plasma samples by column-switching high-performance liquid chromatography–electrospray mass spectrometry 777(1997)67

Salicin

Capillary electrophoretic analysis of salicin in *Salix* spp 781(1997)487

Salicylic acid

Quantitation of acetaminophen and salicylic acid in plasma using capillary electrophoresis without sample pretreatment. Improvement of precision 768(1997)125

Determination of hydroxyl radicals in an advanced oxidation process with salicylic acid trapping and liquid chromatography 796(1998)283

Samarium

Separation of samarium, gadolinium, terbium, dysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

Saponins

Reversed-phase high-performance liquid chromatographic determination of ginsenosides of *Panax quinquefolium* 755(1996)11

Determination of the content and the composition of the main saponins from *Solidago gigantea* AIT. using high-performance liquid chromatography 755(1996)133

Determination of saikosaponins by micellar electrokinetic capillary chromatography 759(1997)193

Identification of soyasaponins by liquid chromatography–thermospray mass spectrometry 777(1997)233

Effect of foaming power and foam stability on continuous concentration with foam counter-current chromatography 791(1997)53

Sarin

In-line respeciation: an ion-exchange ion chromatographic method applied to the separation of degradation products of chemical warfare nerve agents in soil 770(1997)253

Saxitoxins

Solid-phase extraction and high-performance liquid chromatography procedures for the analysis of paralytic shellfish toxins 798(1998)131

SCH 44342

Pharmacokinetic screening for the selection of new drug discovery candidates is greatly enhanced through the use of liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 777(1997)61

Sebutylazine

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Secobarbital

Influence of the degree of substitution of cyclodextrin sulfobutyl ether derivatives on enantioselective separations by electrokinetic chromatography 792(1997)379

Sedatives

Determination of sedatives and anesthetics in plasma by liquid chromatography–mass spectrometry with a desalting system 797(1998)227

Selenium

Investigations of the behaviour of tellurium(IV) and selenium(IV) in ion-exchange chromatography 779(1997)147

Analysis of organic and inorganic selenium anions by ion chromatography–inductively coupled plasma atomic emission spectroscopy 788(1997)105

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

High-performance liquid chromatography of selenoamino acids and organo selenium compounds. Speciation by inductively coupled plasma mass spectrometry 789(1997)349

Simultaneous speciation of aqueous selenium(IV) and selenium(VI) by high-performance liquid chromatography with ultraviolet detection 791(1997)357

Simultaneous analysis of inorganic and organic lead, mercury and selenium by capillary electrophoresis with nitrilotriacetic acid as derivatization agent 796(1998)385

Selenium compounds

Retention behavior of inorganic and organic selenium compounds on a silica-based strong-cation-exchange column with an inductively coupled plasma mass spectrometer as selenium-specific detector 789(1997)233

Selenoamino acids

Analysis of organic and inorganic selenium anions by ion chromatography–inductively coupled plasma atomic emission spectroscopy 788(1997)105

Retention behavior of inorganic and organic selenium compounds on a silica-based strong-cation-exchange column with an inductively coupled plasma mass spectrometer as selenium-specific detector 789(1997)233

High-performance liquid chromatography of selenoamino acids and organo selenium compounds. Speciation by inductively coupled plasma mass spectrometry 789(1997)349

Serotonin

Capillary zone electrophoretic separation of biogenic amines: influence of organic modifier 755(1996)142

Off-column chemiluminescence detection in capillary electrophoresis 771(1997)379

Sesquiterpene ketones

Gas chromatography–mass spectrometry coupled with pseudo-Sadtler retention indices, for the identification of components in the essential oil of *Curcuma longa* L 760(1997)303

Sesquiterpenes

Volatile secondary metabolites from *Spilanthes americana* obtained by simultaneous steam distillation–solvent extraction and supercritical fluid extraction 752(1996)223

Packed column supercritical fluid chromatography of sesquiterpene lactones with different carbon skeletons 779(1997)315

Shogaol

High-performance liquid chromatography–electrospray mass spectrometric analysis of pungent constituents of ginger 796(1998)327

Silanols

The silanol group and its role in liquid chromatography 779(1997)29

Silica hydride

Synthesis and characterization of alkyl bonded phases from a silica hydride via hydrosilation with free radical initiation 786(1997)219

Simazine

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Sobrerol

Quantification of carvone, cineole, perillaldehyde, perillyl alcohol and sobrerol by isocratic high-performance liquid chromatography 793(1998)71

Sodium

Migration behaviour of micelle counterions in micellar electrokinetic chromatography: influence on micelle mobility, efficiency and selectivity 764(1997)127

Determination of Na⁺, K⁺, Mg²⁺ and Ca²⁺ in mixtures of seawater and formation water by capillary electrophoresis 767(1997)311

Column switching for difficult cation separations 789(1997)149

Simple recognition of similar samples for the ion-chromatographic determination of the main cations 789(1997)323

Sodium chloride

Refractive-index detection by interferometric backscatter in packed-capillary high-performance liquid chromatography 762(1997)219

Sodium cholate

Migration behaviour of benzodiazepines in micellar electrokinetic chromatography 756(1996)279

Sodium deoxycholate

Migration behaviour of benzodiazepines in micellar electrokinetic chromatography 756(1996)279

Sodium dimethyldithiocarbamate

Capillary electrophoresis as a routine industrial tool for quantitative analytical testing. Determination of sodium dimethyldithiocarbamate in effluents 781(1997)541

Sodium dodecyl sulphate

Description of the retention behaviour in micellar liquid chromatography as a function of pH, surfactant and modifier concentration 769(1997)155

Quantitative analysis of sodium dodecyl sulphate by capillary electrophoresis 781(1997)67

Sodium 2-naphthalenesulfonate

²H₂O effects on the inclusion complexation of β-cyclodextrin with sodium 2-naphthalenesulfonate in capillary electrophoresis and UV spectrophotometry 800(1998)327

Sodium perchlorate

Use of sodium perchlorate at low pH for peptide separations by reversed-phase liquid chromatography. Influence of perchlorate ion on apparent hydrophilicity of positively charged amino acid side-chains 776(1997)153

Sodium phosphate glass

Capillary electrophoresis of longer-chain polyphosphates 769(1997)349

Sodium salts

Chromatographic ion-exchange and change-partner reactions. New mechanisms concerning ion–ion interactions of inorganic compounds in Sephadex G-15 columns (Review) 789(1997)21

Sodium tetra(*n*-propyl)borate

Sodium tetra(*n*-propyl)borate: a novel aqueous in situ derivatization reagent for the simultaneous determination of organomercury, -lead and -tin compounds with capillary gas chromatography–inductively coupled plasma mass spectrometry 793(1998)99

Somatotropin

Immobilized metal affinity chromatography of human growth hormone. Effect of ligand density 764(1997)27

Sorbic acid

Determination of preservatives in food products by cyclodextrin-modified capillary electrophoresis with multiwavelength detection 768(1997)334

Sotalol

Chiral bioanalysis by normal phase high-performance liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 759(1997)65

Direct liquid chromatographic enantioseparation of sotalol and other β-blockers using an α₁-acid glycoprotein-based chiral stationary phase 760(1997)193

Spermidine

High-performance liquid chromatographic determination of polyamines in milk as their 9-fluorenylmethoxycarbonyl derivatives using a column-switching technique 791(1997)79

Spermine

Electroosmotic flow mobilities in open and packed capillaries with spermine and other amine flow modifiers 768(1997)3

High-performance liquid chromatographic determination of polyamines in milk as their 9-fluorenylmethoxycarbonyl derivatives using a column-switching technique 791(1997)79

Sphingolipids

Cell lipids of the *Candida lipolytica* yeast grown on methanol 755(1996)75

Sphingomyelin

Reversed-phase separations of nitrogenous phospholipids on an octadecanoyl poly(vinyl alcohol) phase 773(1997)93

Spiramycin

Liquid chromatography of spiramycin on poly(styrene–divinylbenzene) 764(1997)43

Spirolactones

High-performance liquid chromatography methods for the separation and quantitation of spironolactone and its degradation products in aqueous formulations and of its metabolites in rat serum 797(1998)271

Spironolactone

High-performance liquid chromatography methods for the separation and quantitation of spironolactone and its degradation products in aqueous formulations and of its metabolites in rat serum 797(1998)271

SR 27417

Drug quantitation on a benchtop liquid chromatography–tandem mass spectrometry system 771(1997)119

Staphylococcal enterotoxin B

see also Toxins

Determination of Staphylococcal Enterotoxin B by on-line (micro) liquid chromatography–electrospray mass spectrometry 757(1997)51

Starch

Quantitative analysis of amylopectin unit chains by means of high-performance anion-exchange chromatography with pulsed amperometric detection 800(1998)199

Starch branching enzymes

Isolation of starch branching enzyme I from potato using γ -cyclodextrin affinity chromatography 800(1998)382

Steroid hormones

see Hormones

Steroids

Tapers and restrictors for capillary electrochromatography and capillary electrochromatography–mass spectrometry 768(1997)9

Drug quantitation on a benchtop liquid chromatography–tandem mass spectrometry system 771(1997)119

Retention behavior of synthetic corticosteroids in packed-column supercritical fluid chromatography 773(1997)277

Application of micellar electrokinetic capillary chromatography to the analysis of illicit drug seizures 780(1997)265

Anionic–zwitterionic mixed micelles in the micellar electrokinetic separation of clinically relevant steroids on a fused-silica capillary 781(1997)467

Theoretical concepts of on-line liquid chromatographic–biochemical detection systems. I. Detection systems based on labelled ligands 787(1997)27

Comparison of retention models for the dependence of retention factors on mobile phase composition in reversed-phase high-performance liquid chromatography 797(1998)23

Simplification of capillary electrochromatography procedures 798(1998)243

On-line concentration of neutral analytes for micellar electrokinetic chromatography. IV. Field-enhanced sample injection 798(1998)251

Capillary electrochromatography of steroids. Increased sensitivity by on-line concentration and comparison with high-performance liquid chromatography 798(1998)259

Sterols

Separation of indenestrol A and B isomers and enantiomers by high-performance liquid chromatography 799(1998)117

Determination of organic molecular markers in marine aerosols and sediments: one-step flash chromatography compound class fractionation and capillary gas chromatographic analysis 799(1998)215

Streptavidin

Implementation of affinity solid-phases in continuous-flow biochemical detection 776(1997)169

Theoretical concepts of on-line liquid chromatographic–biochemical detection systems. II. Detection systems based on labelled affinity proteins 787(1997)37

Streptodornase

Purification of a Streptococcal deoxyribonuclease by affinity chromatography based on a DNA–cellulose matrix 788(1997)75

Strontium

Determination of barium and strontium by capillary zone electrophoresis using an electrolyte containing sulfonazo III 767(1997)303

Strychnine

Identification of pesticide poisoning in wildlife 754(1996)463

Styrene

Qualitative and quantitative analysis of styrene and its various oligomers by liquid chromatography 771(1997)111

Evaluation of styrene in air by thermal desorption–gas chromatography 778(1997)255

Styrene–acrylonitrile

Polymer characterization using on-line coupling of thermal field flow fractionation and hydrodynamic chromatography 765(1997)135

Styrene–maleic anhydride copolymer

Composition and structure analysis of styrene–maleic anhydride copolymer by pyrolysis–gas chromatography 765(1997)279

Substance P

Analysis of bioactive peptides by liquid chromatography–high-resolution electrospray mass spectrometry 767(1997)77

Substance P and related peptides in porcine cortex: whole tissue and nuclear localization 800(1998)21

Subtilisin

Study of correlations between fine structure of bonded layer and affinity properties of silicas with attached cyclic oligopeptides 777(1997)249

Succinic acid

Gas chromatographic determination of organic acids from fruit juices by combined resin mediated methylation and extraction in supercritical carbon dioxide 785(1997)251

Sucrose

Analysis of selected alkaloids and sugars in tobacco extract 775(1997)101

Sugar alcohols

Simultaneous determination of soluble sugars and organic acids as their trimethylsilyl derivatives in apricot fruits by gas–liquid chromatography 758(1997)99

Separation of alditols of interest in food products by high-performance anion-exchange chromatography with pulsed amperometric detection 791(1997)343

Sugars

see also Carbohydrates and Saccharides

Surface plasmon resonance-based detection. An alternative to refractive index detection in high-performance liquid chromatography 759(1997)27

Determination of the absolute configuration of sugar residues using gas chromatography. Method with potential for elimination of references 769(1997)253

Microemulsion electrokinetic chromatography of diphenylhydrazones of dicarbonyl sugars 772(1997)297

Anion-exchange chromatography with electrochemical detection of alditols and sugars at a Cu_2O –carbon composite electrode 773(1997)115

Quantitative analysis of neutral and acidic sugars in whole bacterial cell hydrolysates using high-performance anion-exchange liquid chromatography–electrospray ionization tandem mass spectrometry 776(1997)205

Effect of polystyrene–divinylbenzene resin sulfonation on solute retention in high-performance liquid chromatography 797(1998)139

Separation and direct UV detection of sugars by capillary electrophoresis using chelation of copper(II) 799(1998)283

Introduction of a new scale into reversed-phase high-performance liquid chromatography of pyridylamino sugar chains for structural assignment 800(1998)187

Sulfanilamides

Separation and identification of sulfanilamides by capillary supercritical fluid chromatography–Fourier transform infrared spectroscopy 785(1997)111

Sulfates

Ion chromatography methods for the simultaneous determination of mineral anions in plant sap 752(1996)298

Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67

Use of factorial experimental design for the rapid evaluation of main and interactive factors affecting linearity in calibration curves for sulfate analysis by ion chromatography 770(1997)29

Quantitation of sulfate and thiosulfate in clinical samples by ion chromatography (Review) 789(1997)221

Analysis of difficult samples by capillary zone electrophoresis.

I. Simultaneous separation of detrimental anions in concrete digested with concentrated nitric acid 793(1998)187

Sulfhydryl groups

Removal of the fluorescent 4-(aminosulfonyl)-2,1,3-benzoxadiazole label from cysteine-containing peptides 798(1998)47

Sulfobetaines

Capillary isoelectric focusing: the problem of protein solubility 757(1997)237

Sulfonamides

Migration behavior and separation of sulfonamides in capillary zone electrophoresis. I. Influence of buffer pH and electrolyte modifier 755(1996)261

Migration behavior and separation of sulfonamides in capillary zone electrophoresis. II. Positively charged species at low pH 759(1997)203

Optimisation of a simultaneous separation of sulphonamides, dihydrofolate reductase inhibitors and β -lactam antibiotics by capillary electrophoresis 768(1997)97

Migration behavior and separation of sulfonamides in capillary zone electrophoresis. III. Citrate buffer as a background electrolyte 768(1997)105

Sulfur-selective chemiluminescence detection with packed column supercritical fluid chromatography 779(1997)307

Salt effects in capillary zone electrophoresis. I. Dependence of electrophoretic mobilities upon the hydrodynamic radius 786(1997)171

Migration behavior and selectivity of sulfonamides in capillary electrophoresis 792(1997)37

Simultaneous determination of seven sulfonamide residues in swine wastewater by high-performance liquid chromatography 793(1998)378

Sulfonates

Divalent pairing ion for ion-interaction chromatography of sulfonates and carboxylates 770(1997)51

Sulfonazo III

Determination of barium and strontium by capillary zone electrophoresis using an electrolyte containing sulfonazo III 767(1997)303

Sulfonic acids

Determination of small anions by capillary electrophoresis using indirect UV detection with sulphonated nitrosonaphthol dyes 767(1997)293

Effects of the working electrolyte (cyclodextrin type and pH) on the separation of aromatic sulphonic acids by capillary zone electrophoresis 772(1997)385

Sulfonic acids, aromatic

Analysis of polar organic pollutants in the Elbe river by flow injection analysis and high-performance liquid chromatography with tandem mass spectrometry 777(1997)127

Sulfonylureas

Analysis of primisulfuron and triasulfuron in water and soil samples by micellar electrokinetic capillary chromatography 766(1997)225

Determination of chlorsulfuron and tribenuron-methyl residues in agricultural soils 778(1997)119

Application of liquid chromatography–mass spectrometryⁿ analyses to the characterization of novel glyburide metabolites formed in vitro 794(1998)15

Microwave-assisted solvent extraction and reversed-phase liquid chromatography–UV detection for screening soils for sulfonylurea herbicides 798(1998)179

Sulfoxide

Chiral resolution of pantoprazole sodium and related sulfoxides by complex formation with bovine serum albumin in capillary electrophoresis 759(1997)185

Sulfur

Sulfur speciation by capillary electrophoresis with indirect spectrophotometric detection: In search of a suitable carrier electrolyte to maximize sensitivity 770(1997)311

Sulfur compounds

Detection of traces of oxidized and reduced sulfur compounds in small samples by combination of different high-performance liquid chromatography methods 760(1997)295

Molecular analysis of the sulphur components in a light cycle oil of a catalytic cracking unit by gas chromatography with mass spectrometric and atomic emission detection 761(1997)231

Quantitation of volatile sulphur compounds in polluted waters 778(1997)329

Gas chromatographic–mass spectrometric analysis of mercaptan odorants in liquefied petroleum gas and liquefied natural gas 791(1997)367

Sulfur compounds, volatile

Chromatographic analysis of volatile sulphur compounds in wines using the static headspace technique with flame photometric detection 773(1997)261

Problems with the determination of environmental sulphur compounds by gas chromatography (Review) 793(1998)1

Sulfur-containing compounds

Ultrafiltration: a technique for determining the molecular-mass distribution of group parameters of organic pollutants in natural waters 763(1997)315

Sulfur oxyanions

Determination of sulfur oxyanions by ion chromatography on a silica ODS column with tetrapropylammonium salt as an ion-pairing reagent 770(1997)157

Surfactants

see also Detergents and Subject Index

High-performance liquid chromatographic–fluorescence determination of aliphatic alcohol polyethoxylates and poly(ethylene glycol)s in aqueous samples 756(1996)159

Migration behaviour of benzodiazepines in micellar electrokinetic chromatography 756(1996)279

Capillary electrophoresis of herbicides. II. Evaluation of alkylglucoside chiral surfactants in the enantiomeric separation of phenoxy acid herbicides 757(1997)263

Analysis of cationic surfactants by capillary electrophoresis 757(1997)275

Micellar electrokinetic chromatography of charged and neutral drugs in acidic running buffers containing a zwitterionic surfactant, sulfonic acids or sodium dodecyl sulphate. Separation of heroin, basic by-products and adulterants 760(1997)245

High-performance liquid chromatographic analysis of a linear alkylbenzenesulfonate and its environmental biodegradation metabolites 760(1997)285

Surfactant-sensitized post-column reaction with xylenol orange for the determination of lanthanides by ion chromatography 770(1997)75

Modified electroosmotic flow by cationic surfactant additives in capillary electrophoresis. Evaluation of electrolyte systems for anion analysis 772(1997)171

Separation and quantitative determination of non-ionic surfactants used as pesticide additives (Review) 774(1997)265

Analysis of benzalkonium chloride in the effluent from European hospitals by solid-phase extraction and high-performance liquid chromatography with post-column ion-pairing and fluorescence detection 774(1997)281

Determination of alkylphenols, alkylphenoxyethoxylates and linear alkylbenzenesulfonates in sediments by accelerated solvent extraction and supercritical fluid extraction 775(1997)187

Use of different surfactants (sodium dodecyl sulfate, bile salts and ionic polymers) in micellar electrokinetic capillary chromatography. Application to the separation of organophosphorus pesticides 778(1997)201

Effect of the polar groups of anionic surfactant on migration behavior in micellar electrokinetic chromatography 781(1997)11

Determination of cationic surfactants by capillary electrophoresis with indirect photometric detection 781(1997)17

Capillary electrophoresis of cationic and anionic surfactants with indirect conductivity detection 781(1997)533

Single step method for the accurate concentration determination of polysorbate 80 786(1997)99

Selectivity of polymeric and polymer-supported pseudo-stationary phases in micellar electrokinetic chromatography 792(1997)105

Characterization of surfactants and their biointermediates by liquid chromatography–mass spectrometry 794(1998)165

On-line micellar electrokinetic chromatography–mass spectrometry with a high-molecular-mass surfactant 794(1998)317

Analysis of ethoxylated polymers by capillary electrophoresis in UV-transparent polymer networks and by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry 800(1998)13

Surfactants, multi-chain

Separation of naphthalene and flavone derivatives by micellar electrokinetic chromatography with double- and triple-chain surfactants 755(1996)147

Surfactants, non-ionic

Comparison of the retention behavior of polyethoxylated alcohols on porous graphitic carbon and polar as well as apolar bonded-silica phases 797(1998)83

Symetryn

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Syringaldehyde

Analysis of volatile components of oak wood by solvent extraction and direct thermal desorption–gas chromatography–mass spectrometry 778(1997)427

T

2,4,5-T

Determination of chlorophenoxy herbicides in waters by capillary gas chromatography with ion trap detection 758(1997)324

Tannins

Analysis of old brandy and oak extracts by capillary electrophoresis 768(1997)143

Tanshinone IIA

Extraction of Tanshinone IIA from *Salvia miltiorrhiza bunge* using supercritical fluid extraction and a new extraction technique, phytosol solvent extraction 799(1998)343

Tartaric acid

New approach to determining each component of a two-component overlapping peak by single-column anion ion chromatography 769(1997)333

Gas chromatographic determination of organic acids from fruit juices by combined resin mediated methylation and extraction in supercritical carbon dioxide 785(1997)251

Taurochenodeoxycholic acid

Bile salt micellar electrokinetic chromatography of bilirubin and related compounds 779(1997)205

Taurocholic acid

Bile salt micellar electrokinetic chromatography of bilirubin and related compounds 779(1997)205

Taurodeoxycholic acid

Bile salt micellar electrokinetic chromatography of bilirubin and related compounds 779(1997)205

Taxanes

Determination of a new polymer-bound paclitaxel derivative (PNU 166945), free paclitaxel and 7-epipaclitaxel in dog plasma and urine by reversed-phase high-performance liquid chromatography with UV detection 797(1998)295

Technetium

Technetium speciation: non-size effects in size-exclusion chromatography 755(1996)219

Teichoic acids

Quantitative analysis of neutral and acidic sugars in whole bacterial cell hydrolysates using high-performance anion-exchange liquid chromatography–electrospray ionization tandem mass spectrometry 776(1997)205

Teichuronic acids

Quantitative analysis of neutral and acidic sugars in whole bacterial cell hydrolysates using high-performance anion-exchange liquid chromatography–electrospray ionization tandem mass spectrometry 776(1997)205

Teicoplanin

Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309

Enantiomeric separation of small chiral peptides by capillary electrophoresis 792(1997)393

High-performance liquid chromatographic separation of enantiomers of unusual amino acids on a teicoplanin chiral stationary phase 793(1998)283

Tellurium

Investigations of the behaviour of tellurium(IV) and selenium(IV) in ion-exchange chromatography 779(1997)147

On the use of inductively coupled plasma mass spectrometry as an element specific detector for liquid chromatography: optimization of an industrial tellurium removal process 794(1998)219

Temephos

Identification of fenthion and temephos and their transformation products in water by high-performance liquid chromatography with diode array detection and atmospheric pressure chemical ionization mass spectrometric detection 777(1997)99

Terazosin

Enantioselective determination of terazosin in human plasma by normal phase high-performance liquid chromatography–electrospray mass spectrometry 794(1998)45

Terbium

Separation of samarium, gadolinium, terbium, dysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

Terbutaline

Ruggedness of enantiomeric separation by capillary electrophoresis and high-performance liquid chromatography with methylated cyclodextrins as chiral selectors 763(1997)139

Terbutylazine

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Terpenes

Changes in chemical composition of catalytically hydrogenated orange oil (*Citrus sinensis*) 752(1996)217

Reversed-phase high-performance liquid chromatographic determination of ginsenosides of *Panax quinquefolium* 755(1996)11

Chiral separation of monoterpenes using mixtures of sulfated β -cyclodextrins and α -cyclodextrin as chiral additives in the reversed-polarity capillary electrophoresis mode 759(1997)149

Decomposition of α -pinene and sabinene on solid sorbents, Tenax TA and Carboxen 786(1997)185

Terpenoids

Comparison of chromatographic systems for triterpenoids from Neem (*Azadirachta indica*) seeds 761(1997)53

Gas chromatographic–mass spectroscopic analysis of the acidic triterpenic fraction of mastic gum 769(1997)263

Supercritical fluid chromatography–mass spectrometry of thyme extracts (*Thymus vulgaris* L.) 773(1997)377

Tetraalkylammonium ions

Separation of hydrophobic compounds by electrokinetic chromatography with tetraalkylammonium ions 764(1997)323

Effects of tetraalkylammonium salts on the micellar electrokinetic chromatography of aniline and substituted anilines 799(1998)289

Tetrabutylammonium

Studies on elution conditions for the determination of anions by suppressed ion-interaction chromatography using a graphitized carbon column 800(1998)239

Tetracyclines

Use of metal complexation in non-aqueous capillary electrophoresis systems for the separation and improved detection of tetracyclines 779(1997)235

Tetradecyltrimethylammonium bromide

Large-volume stacking for quantitative analysis of anions in capillary electrophoresis. II. Large-volume stacking without polarity switching 757(1997)291

Tetrafluoroethane

Comparison of 1,1,1,2-tetrafluoroethane and carbon dioxide-based mobile phases for packed column supercritical fluid chromatography 782(1997)237

Tetrahydrocarbolinecarboxylic acids

Presence of tetrahydro- β -carboline-3-carboxylic acids in foods by gas chromatography–mass spectrometry as their *N*-methoxycarbonyl methyl ester derivatives 765(1997)265

Tetrahydrocarbolines

High-performance liquid chromatography–electrospray ionisation–tandem mass spectrometry for the analysis of 1,2,3,4-tetrahydro- β -carboline derivatives 767(1997)101

Tetrahydrothiophene

Gas chromatographic–mass spectrometric analysis of mercaptan odorants in liquefied petroleum gas and liquefied natural gas 791(1997)367

Tetralin derivatives

Chiral separations of indan, tetralin and benzosuberone derivatives by capillary electrophoresis 793(1998)135

Tetramethylammonium cation

Analysis of tetramine in sea snails by capillary electrophoresis–tandem mass spectrometry 781(1997)555

Tetramethylammonium chloride

Purification of insulin-like growth factor-I and related proteins using underivatized silica 753(1996)73

2,2,6,6-Tetramethyl-1-piperidinyloxy free radical

Separation of cyclic nitroxide free radicals and their redox forms with dual microelectrochemical detection 771(1997)89

Tetramethylpirrolinocarboxamidopropylphthalimide hydrochloride

Validated liquid chromatographic method for the determination of *N*-3-(2,2,5,5-tetramethyl-3-pyrrolin-3-carboxamidopropylphthalimide hydrochloride), a novel antiarrhythmic agent in human plasma 797(1998)265

Tetramethylrhodamine

Study of the enzymatic transformation of fluorescently labeled oligosaccharides in human epidermoid cells using capillary electrophoresis with laser-induced fluorescence detection 781(1997)515

Tetramine

Analysis of tetramine in sea snails by capillary electrophoresis–tandem mass spectrometry 781(1997)555

Tetrandrine

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

(*R*)-(–)-2-(2,4,5,7-Tetranitro-9-fluorenylideneaminoxy) propionic acid

Enantiomer separation of [60]fullerene derivatives by micro-column high-performance liquid chromatography using (*R*)-(–)-2-(2,4,5,7-tetranitro-9-fluorenylideneaminoxy) propionic acid as chiral stationary phase 791(1997)65

Tetraphenylporphine

Liquid chromatographic separation of metal–tetraphenylporphine complexes based on coordination interaction with immobilized neutral ligand in stationary phase 786(1997)269

Tetrazol-5-thiones

Chiral resolution of diols by capillary electrophoresis using borate–cyclodextrin complexation 761(1997)269

Thalrugosine

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography 799(1998)337

Theanine

Simultaneous determination of qualitatively important components in green tea infusions using capillary electrophoresis 758(1997)332

Theophylline

Chromatographic characterization of a molecularly imprinted polymer binding theophylline in aqueous buffers 786(1997)23

Theophylline derivatives

Enantiomeric separation of chiral theophylline derivatives by liquid chromatography on cellulose-based sorbents 760(1997)278

Thiabendazole

Complementation of direct-injection high-performance liquid chromatography and enzyme-linked immunosorbent assay for the analysis of thiabendazole in fruit juices and concentrates 754(1996)431

Simplified reversed-phase conditions for the determination of benzimidazole fungicides in fruits by high-performance liquid chromatography with UV detection 778(1997)95

Thiazide diuretics

Gradient capillary electrochromatography of drug mixtures with UV and electrospray ionisation mass spectrometric detection 768(1997)89

Thiazopyr

Determination of thiazopyr residues in soil and plants by gas chromatography with nitrogen–phosphorus detection and confirmation by gas chromatography–mass spectrometry 778(1997)193

Thimerosal

Quantification of urinary uric acid in the presence of thymol and thimerosal by high-performance liquid chromatography 763(1997)187

Thiocyanate

Thin-layer chromatographic separation, colorimetric determination and recovery of thiocyanate from photogenic waste, river and sea waters 774(1997)373

Thiodiglycol compounds

Application of liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry, and tandem mass spectrometry, to the analysis and identification of degradation products of chemical warfare agents 759(1997)79

Thiol ester proteins

Affinity chromatography of thiol ester-containing proteins 799(1998)139

Thiols

Determination of glutathione, cysteine and N-acetylcysteine in rabbit eye tissues using high-performance liquid chromatography and post-column derivatization with 5,5'-dithiobis(2-nitrobenzoic acid) 778(1997)347

Thioneins

Application of solid-phase extraction to the preconcentration of metallothionein and metals from physiological fluids 766(1997)121

Thiophenes

Molecular analysis of the sulphur components in a light cycle oil of a catalytic cracking unit by gas chromatography with mass spectrometric and atomic emission detection 761(1997)231

Thiophosphoryl nucleotides

Iodine–azide reagent in detection of thiophosphoryl nucleotides in thin-layer chromatography systems 773(1997)285

Thiosulfate

Determination of sulfur oxyanions by ion chromatography on a silica ODS column with tetrapropylammonium salt as an ion-pairing reagent 770(1997)157

Quantitation of sulfate and thiosulfate in clinical samples by ion chromatography (Review) 789(1997)221

Thrombin

Characterisation of silica-based heparin affinity sorbents from equilibrium binding studies on plasma fractions containing thrombin 762(1997)113

Study of correlations between fine structure of bonded layer and affinity properties of silicas with attached cyclic oligopeptides 777(1997)249

Thrombin inhibitors

Quantitation of an orally available thrombin inhibitor in rat, monkey and human plasma and in human urine by high-performance liquid chromatography and fluorescent post-column derivatization of arginine 762(1997)299

Thromboxanes

Analysis of prostaglandins by micellar electrokinetic capillary chromatography 761(1997)277

Thymol

Quantification of urinary uric acid in the presence of thymol and thimerosal by high-performance liquid chromatography 763(1997)187

Supercritical fluid chromatography–mass spectrometry of thyme extracts (*Thymus vulgaris* L.) 773(1997)377

Thyroxine

Optimization of post-column chemiluminescent detection for low-molecular-mass conjugates of acridinium esters 765(1997)145

Tin

Determination and speciation of metals by liquid chromatography (Review) 789(1997)301

Tocainide

Analysis of basic pharmaceuticals by capillary electrophoresis in coated capillaries and on-line mass spectrometric detection 752(1996)271

Tocopherols

Reversed-phase high-performance liquid chromatographic separations of tocopherols 782(1997)25

Determination of E vitamers in microalgae using high-performance liquid chromatography with fluorescence detection 782(1997)63

Concurrent liquid chromatographic separation and photodiode array detection of retinol, tocopherols, all-*trans*- α -carotene, all-*trans*- β -carotene and the mono-*cis* isomers of β -carotene in extracts of human plasma 787(1997)111

Tolperisone

Separation of basic drug enantiomers by capillary zone electrophoresis using ovoglycoprotein as a chiral selector 782(1997)281

Toluene

Headspace gas chromatographic determination of the plant cuticle–air partition coefficients for monocyclic aromatic hydrocarbons as environmental compartment 774(1997)213

Evaluation of environmental levels of aromatic hydrocarbons in gasoline service stations by gas chromatography 778(1997)263

Determination of volatile organic compounds in marine biota 799(1998)233

Toluene-2,4-diisocyanate

Determination of toluene-2,4-diisocyanate in environmental and workplace air by sampling onto Tenax-TA followed by thermal desorption and capillary gas chromatography using flame ionisation and ion-trap detection 771(1997)233

Toluenesulfonate

Ideal model of chromatography applied to charged solutes in reversed-phase liquid chromatography 761(1997)3

Method for determination of the chromatographic surface area of reversed-phase surfaces 761(1997)13

TO-PRO-3

Characterization of TO-PRO-3 as an intercalator for double-stranded DNA analysis with red diode laser-induced fluorescence detection 781(1997)357

Toxins

see also Mycotoxins, Neurotoxins and Staphylococcal enterotoxin B

Development of a manganese dioxide solid-phase reactor for oxidation of toxins associated with paralytic shellfish poisoning 755(1996)227

Novel methods for identification and quantification of the mushroom nephrotoxin orellanine. Thin-layer chromatography and electrophoresis screening of mushrooms with electron spin resonance determination of the toxin 758(1997)145

Optimization of sample cleanup procedure for determination of diarrhetic shellfish poisoning toxins by use of experimental design 764(1997)223

Liquid chromatography with fluorimetric, mass spectrometric and tandem mass spectrometric detection for the investigation of the seafood-toxin-producing phytoplankton, *Dinophysis acuta* 777(1997)213

Determination of mycotoxins produced by *Fusarium* isolates from banana fruits by capillary gas chromatography and high-performance liquid chromatography 778(1997)363

Detection of isoforms and isomers of rattlesnake myotoxins by capillary electrophoresis and matrix-assisted laser desorption time-of-flight mass spectrometry 781(1997)429

Analysis of tetramine in sea snails by capillary electrophoresis–tandem mass spectrometry 781(1997)555

Improvement on sample clean-up for high-performance liquid chromatographic–fluorimetric determination of diarrhetic shellfish toxins using 1-bromoacetylpyrene 793(1998)63

Solid-phase extraction and high-performance liquid chromatography procedures for the analysis of paralytic shellfish toxins 798(1998)131

Isolation of a new okadaic acid analogue from phytoplankton implicated in diarrhetic shellfish poisoning 798(1998)137

Sensitive determination of anatoxin-a, homoanatoxin-a and their degradation products by liquid chromatography with fluorimetric detection 798(1998)147

Determination of some physicochemical parameters of microcystins (cyanobacterial toxins) and trace level analysis in environmental samples using liquid chromatography 799(1998)155

Tramadol

Laboratory-developed simulated moving bed for chiral drug separations. Design of the system and separation of Tramadol enantiomers 769(1997)49

trans-1,2-Diaminocyclohexane-N,N,N',N'-tetraacetic acid

Laser induced resonance energy transfer – a novel approach towards achieving high sensitivity in capillary electrophoresis. Part B: Applications for environmental testing 767(1997)217

Transition metals

see also Metals

Use of ion chromatography for the determination of transition metals in the control of sewage-treatment-plant and related waters 770(1997)185

Tretinoin

Comparison of the separation of *cis/trans* isomers of tretinoin with different stationary phases by liquid chromatography–nuclear magnetic resonance coupling 765(1997)207

Triacylglycerols

Stereospecific analysis of the triacylglycerol fraction and linear discriminant analysis in a climatic differentiation of Umbrian extra-virgin olive oils 758(1997)109

Triacylglycerol mixture for testing capillary columns for high-temperature gas chromatography 758(1997)235

Normal-phase separation effects with lipids on a silver ion high-performance liquid chromatography column 764(1997)337

Determination of primary oxidation products of linoleic acid and triacylglycerols 767(1997)63

Non-aqueous reversed-phase high-performance liquid chromatography of synthetic triacylglycerols and diacylglycerols 782(1997)41

Analysis of cyanolipids and triacylglycerols from sapindaceae seed oils with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry 787(1997)181

Triallate

Extraction of triallate from soil with supercritical carbon dioxide and determination by gas chromatography–atomic emission detection. Comparison with a solvent extraction procedure 754(1996)257

Triasulfuron

Analysis of primisulfuron and triasulfuron in water and soil samples by micellar electrokinetic capillary chromatography 766(1997)225

Triazilam

Use of simulated liquid chromatography–diode array detection data for the definition of a guide curve in peak purity assessment by spectral comparison 786(1997)1

Triazines

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography 752(1996)147

Influence of template basicity and hydrophobicity on the molecular recognition properties of molecularly imprinted polymers 753(1996)191

High-performance separations in the determination of triazine herbicides and their residues (Review) 754(1996)17

Identification of herbicides in river water using on-line trace enrichment combined with column liquid chromatography–Fourier-transform infrared spectrometry 756(1996)145

Chemiluminescence nitrogen detection for packed-column supercritical fluid chromatography with methanol modified carbon dioxide 757(1997)183

Preconcentration of triazine herbicides from water by an ion chromatography column and determination by gas chromatography–mass spectrometry 760(1997)314

Ultra-trace-level determination of the antifouling agent Irgarol 1051 by gas chromatography with tandem mass spectrometric detection 766(1997)153

Determination of pesticides in environmental waters by solid-phase extraction and gas chromatography with electron-capture and mass spectrometry detection 771(1997)221

Rapid target analysis of microcontaminants in water by on-line single-short-column liquid chromatography combined with atmospheric pressure chemical ionization tandem mass spectrometry 777(1997)81

New method for the rapid determination of triazine herbicides and some of their main metabolites in water by using coupled-column liquid chromatography and large volume injection 778(1997)171

Optimizing recoveries of two chlorotriazine herbicide metabolites and 11 pesticides from aqueous samples using solid-phase extraction and gas chromatography–mass spectrometry 790(1997)161

Determination of hexazinone and its metabolites in groundwater by capillary electrophoresis 793(1998)349

Rapid target analysis of microcontaminants in water by on-line single-short-column liquid chromatography combined with atmospheric pressure chemical ionization ion-trap mass spectrometry 794(1998)201

Dissipation of some citrus selective residual herbicides in an irrigation well 795(1998)125

Inter-laboratory validation of solid-phase microextraction for the determination of triazine herbicides and their degradation products at ng/l level in water samples 795(1998)371

Triazolam

Simulations of liquid chromatography–diode array detector data including instrumental artefacts for the evaluation of mixture analysis techniques 758(1997)1

Determination of triazolam in a drug tablet by thermal desorption gas chromatography 761(1997)332

Triazophos

Resolution of folpet, procymidone and triazophos in high-performance liquid chromatography–diode array detection by using partial least squares calibration to cross-sections of spectrochromatograms 778(1997)183

Tribenuron-methyl

Determination of chlorsulfuron and tribenuron-methyl residues in agricultural soils 778(1997)119

Tributylbenzene

Absolute quantitation of lignin pyrolysis products using an internal standard 773(1997)227

Tributyl phosphate

Novel calibration technique for headspace analysis of semivolatiles compounds 793(1998)397

Tributyltin

Reversed-phase ion-pair chromatography with inductively coupled plasma–mass spectrometry detection for the determination of organo-tin compounds in waters 758(1997)85

Speciation of organo-tin compounds using liquid chromatography–atmospheric pressure ionisation mass spectrometry and liquid chromatography–inductively coupled plasma mass spectrometry as complementary techniques 794(1998)211

Trichloroanisole

Analysis of 2,4,6-trichloroanisole in wines using solid-phase microextraction coupled to gas chromatography–mass spectrometry 786(1997)293

Trichlorophenol

Monitoring of selected metabolites and biotransformation products from fermentation broths by high-performance liquid chromatography 763(1997)165

Simplified determination of combined residues of prochloraz and its metabolites in vegetable, fruit and wheat samples by gas chromatography 765(1997)127

Trichloropyridinol

Sub- and supercritical fluid extraction of trichloropyridinol from soil prior to immunoassay 785(1997)329

Trichothecenes

Determination of mycotoxins produced by *Fusarium* isolates from banana fruits by capillary gas chromatography and high-performance liquid chromatography 778(1997)363

Determination of trichothecene mycotoxins in wheat by use of supercritical fluid extraction and high-performance liquid chromatography with diode array detection or gas chromatography with electron capture detection 795(1998)297

Triclosan

Identification of an anti-bacterial agent in toothpaste via liquid chromatography–Fourier transform infrared spectrometry mobile phase elimination 755(1996)211

Trifluoroethanol

Modification of capillary electrophoresis selectivity in hydro-organic solutions. Dissociation constants and Stokes radius measurements of peptides in water–2,2,2-trifluoroethanol mixtures 792(1997)57

Triglycerides

Retention behavior of triglycerides in subcritical fluid chromatography with carbon dioxide mobile phase 764(1997)301

Quantitative aspects of triglyceride analysis by gas–liquid chromatography using a short metal capillary column 773(1997)233

Analysis of plant oils by subcritical fluid chromatography using pattern fitting 800(1998)317

Trihydroxybenzo[*a*]pyrene–deoxyguanosylphosphate

On-line preconcentration and high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbon–DNA adducts using copper phthalocyanine trisulfonic acid as a group-specific adsorbent 755(1996)205

Trimethylgermanium chloride

Direct determination of trimethylgermanium in water by on-column capillary gas chromatography with flame photometric detection using quartz surface-induced germanium emission 759(1997)119

Trimethylsilyldiazomethane

Determination of phenoxy acid herbicides in vegetation, utilising high-resolution gel permeation chromatographic clean-up and methylation with trimethylsilyldiazomethane prior to gas chromatographic analysis with mass-selective detection 755(1996)245

Trimethylsulfonium hydroxide

Reaction of fatty acid amides and ethanolamides with trimethylsulfonium hydroxide 773(1997)239

Trinitrotoluene

Rapid on-line chromatographic determination of trace-level munitions in aqueous samples 753(1996)81

Trace analysis of military high explosives (2,4,6-trinitrotoluene and hexahydro-1,3,5-trinitro-1,3,5-triazine) in agricultural crops 775(1997)117

Triphenylmagnesium bromide

Selective determination of Sb(III) by gas chromatography–quartz furnace atomic absorption spectrometry after derivatization with triphenylmagnesium bromide 764(1997)169

Triphenyltin

Reversed-phase ion-pair chromatography with inductively coupled plasma–mass spectrometry detection for the determination of organo-tin compounds in waters 758(1997)85

Speciation of organo-tin compounds using liquid chromatography–atmospheric pressure ionisation mass spectrometry and liquid chromatography–inductively coupled plasma mass spectrometry as complementary techniques 794(1998)211

Tris(cysteinesulfonato(-N,S)cobaltate(III))

Chromatographic separation and characterization of the photoproduct of tris(L-cysteinesulfonato-N,S)cobaltate(III) 789(1997)273

Tris(hydroxymethyl)aminomethane

Migration behaviour of micelle counterions in micellar electrokinetic chromatography: influence on micelle mobility, efficiency and selectivity 764(1997)127

Triterpene glycosides

Liquid chromatography coupled to mass spectrometry and nuclear magnetic resonance spectroscopy for the screening of plant constituents 794(1998)299

Triterpenes

Isolation of three triterpenes and several aliphatic alcohols from *Crataegus monogyna* Jacq 767(1997)340

Triterpenoid acids

Gas chromatographic–mass spectroscopic analysis of the acidic triterpenic fraction of mastic gum 769(1997)263

Triterpenoids

Comparison of chromatographic systems for triterpenoids from Neem (*Azadirachta indica*) seeds 761(1997)53

Reversed-phase high-performance liquid chromatographic separation and quantification of lantadenes using isocratic systems 786(1997)181

Effect of foaming power and foam stability on continuous concentration with foam counter-current chromatography 791(1997)53

Triton X-100

Composition analysis of detergents of the polyoxyethylene type: comparison of thin-layer chromatography, reversed-phase chromatography and matrix-assisted laser desorption/ionization mass spectrometry 791(1997)245

Triton X-114

Composition analysis of detergents of the polyoxyethylene type: comparison of thin-layer chromatography, reversed-phase chromatography and matrix-assisted laser desorption/ionization mass spectrometry 791(1997)245

Tropane alkaloids

Lesser alkaloids of cocaine-bearing plants. III. 2-Carbomethoxy-3-oxo substituted tropane esters: detection and gas chromatographic–mass spectrometric characterization of new minor alkaloids found in South American *Erythroxylum coca* var. *coca* 756(1996)185

Truxilline

Comparative determination of total isomeric truxillines in illicit, refined, South American cocaine hydrochloride using capillary gas chromatography–electron capture detection 756(1996)193

Trypsin

Study of correlations between fine structure of bonded layer and affinity properties of silicas with attached cyclic oligopeptides 777(1997)249

Tryptamine

Capillary zone electrophoretic separation of biogenic amines: influence of organic modifier 755(1996)142

Tryptophan

Enantiomeric separations of primary amino compounds by non-aqueous capillary zone electrophoresis with a chiral crown ether 757(1997)328

Partitioning of hydrophobic amino acids and oligopeptides in aqueous two-phase system containing self-aggregating block copolymer. Effects of temperature, salts and surfactants 761(1997)91

Tryptophan analysis in peptides and proteins, mainly by liquid chromatography 763(1997)1

Effect of mobile phase composition on the binding kinetics of chiral solutes on a protein-based high-performance liquid chromatography column. Interactions of D- and L-tryptophan with immobilized human serum albumin 766(1997)15

Interaction between tryptophan residues and hydrophobically modified dextran. Effect on partitioning of peptides and proteins in aqueous two-phase systems 766(1997)99

Enantiomeric properties of human albumin immobilized on porous silica supports coated with polymethacryloyl chloride 776(1997)37

Determination of γ -radiation induced products in aqueous solutions of tryptophan and synthesis of 4-, 6- and 7-hydroxytryptophan 786(1997)57

Tuberculin

High-performance liquid chromatographic assay of 8-hydroxyquinoline sulfate and its stability in immunobiological preparations 798(1998)173

Tumour necrosis factor

Role of polyethyleneimine in the purification of recombinant human tumour necrosis factor beta 760(1997)165

Tungstate

New approach to determining each component of a two-component overlapping peak by single-column anion ion chromatography 769(1997)333

Tween 20

Composition analysis of detergents of the polyoxyethylene type: comparison of thin-layer chromatography, reversed-phase chromatography and matrix-assisted laser desorption/ionization mass spectrometry 791(1997)245

Tyramine

Capillary zone electrophoretic separation of biogenic amines: influence of organic modifier 755(1996)142

Determination of biogenic amines in wines by high-performance liquid chromatography with on-column fluorescence derivatization 757(1997)311

Determination of histamine and some other amines by high-performance capillary electrophoresis with on-line mode in-capillary derivatization 790(1997)177

U

Uracil nucleosides

Resolution of enantiomers of uridine analogs, potential antiviral agents 752(1996)131

Uranium

Investigation of adducts of lanthanide and uranium β -diketonates with organophosphorus Lewis bases by supercritical fluid chromatography 793(1998)107

Urea

Simultaneous use of urea and acetonitrile as organic modifiers for optimization of resolution in micellar electrokinetic chromatography 792(1997)157

Ureidodeoxyriboside

New hydrolytic products detected in aqueous solutions of bromodeoxyuridine 797(1998)283

Uric acid

Quantification of urinary uric acid in the presence of thymol and thimerosal by high-performance liquid chromatography 763(1997)187

Uridine

Resolution of enantiomers of uridine analogs, potential antiviral agents 752(1996)131

Uridine diphosphate glucuronyl transferase

In vitro glucuronidation of 7-hydroxycoumarin and determination of 7-hydroxycoumarin and 7-hydroxycoumarin glucuronide by capillary electrophoresis 772(1997)321

V

Valine

Computerized optimization of selectivity for direct capillary gas chromatographic multicomponent separations of enantiomers 779(1997)253

Vanadium

Capillary gas chromatographic determination of vanadium in crude petroleum oil using fluorinated ketoamines as derivatizing reagents 766(1997)159

Vancomycin

Vacancy affinity capillary electrophoresis, a new method for measuring association constants 775(1997)313

Enantiomeric separation of acidic herbicides by capillary electrophoresis using vancomycin as chiral selector 781(1997)503

Enantioselectivity in capillary electrophoresis using the macrocyclic antibiotics 792(1997)309

Enantiomeric separation of small chiral peptides by capillary electrophoresis 792(1997)393

Chiral capillary electrophoresis–electrospray mass spectrometry coupling using vancomycin as chiral selector 800(1998)69

Vanillic acid

Micellar electrokinetic chromatography as an alternative to high-performance liquid chromatography for separation and determination of phenolic compounds in Japanese spirituous liquor 793(1998)409

Vanillin

Analysis of volatile components of oak wood by solvent extraction and direct thermal desorption–gas chromatography–mass spectrometry 778(1997)427

Micellar electrokinetic chromatography as an alternative to high-performance liquid chromatography for separation and determination of phenolic compounds in Japanese spirituous liquor 793(1998)409

Varidase

Purification of a Streptococcal deoxyribonuclease by affinity chromatography based on a DNA–cellulose matrix 788(1997)75

Verapamil

Chiral bioanalysis by normal phase high-performance liquid chromatography–atmospheric pressure ionization tandem mass spectrometry 759(1997)65

On-line column-switching high-performance liquid chromatography analysis of cardiovascular drugs in serum with automated sample clean-up and zone-cutting technique to perform chiral separation 762(1997)235

Separation of basic drug enantiomers by capillary zone electrophoresis using ovoglycoprotein as a chiral selector 782(1997)281

Vinblastine

Quantification of vincristine and vinblastine in *Catharanthus roseus* plants by capillary zone electrophoresis 755(1996)281

Vinclozolin

Determination of the fungicide vinclozolin in honey and bee larvae by solid-phase and solvent extraction with gas chromatography and electron-capture and mass spectrometric detection 754(1996)507

Vincristine

Quantification of vincristine and vinblastine in *Catharanthus roseus* plants by capillary zone electrophoresis 755(1996)281

Vinylguaiacol

Micellar electrokinetic chromatography as an alternative to high-performance liquid chromatography for separation and determination of phenolic compounds in Japanese spirituous liquor 793(1998)409

Vinylic organosol

Identification of potential migrants from a vinylic organosol varnish by gas chromatography–mass spectrometry and liquid chromatography–mass spectrometry 771(1997)366

Visnadin

Determination of furanochromones and pyranocoumarins in drugs and *Ammi visnaga* fruits by combined solid-phase extraction–high-performance liquid chromatography and thin-layer chromatography–high-performance liquid chromatography 797(1998)305

Visnagin

Determination of furanochromones and pyranocoumarins in drugs and *Ammi visnaga* fruits by combined solid-phase extraction–high-performance liquid chromatography and thin-layer chromatography–high-performance liquid chromatography 797(1998)305

Vitamin A palmitate

Determination of vitamin A palmitate in cereal products using supercritical fluid extraction and liquid chromatography with electrochemical detection 765(1997)215

Vitamins

see also Ascorbic acid and Retinol

Separation of retinoids by micellar electrokinetic capillary chromatography 761(1997)307

Separation and identification of vitamin A acetate isomers by supercritical fluid chromatography–¹H NMR coupling 761(1997)336

Supercritical fluid extraction and reversed-phase liquid chromatography methods for vitamin A and β -carotene. Heterogeneous distribution of vitamin A in the liver 762(1997)201

Determination of vitamin A palmitate in cereal products using supercritical fluid extraction and liquid chromatography with electrochemical detection 765(1997)215

Determination of vitamins A and E in infant milk formulae by high-performance liquid chromatography 778(1997)243

Determination of water-soluble vitamins in infant milk by high-performance liquid chromatography 778(1997)247

Simultaneous determination of vitamins A and E in rat tissues by high-performance liquid chromatography 778(1997)415

Synthesis and characterization of a strong cation exchanger based on polymer-coated silica for high-performance liquid chromatography 779(1997)113

Determination of E vitamers in microalgae using high-performance liquid chromatography with fluorescence detection 782(1997)63

Analysis of ascorbate and dehydroascorbate in plant extracts by high-resolution selected ion monitoring gas chromatography–mass spectrometry 782(1997)95

Comparison of the separation of fat-soluble vitamins using β -cyclodextrins in high-performance liquid chromatography and micellar electrokinetic chromatography 782(1997)227

Supercritical fluid chromatography–proton nuclear magnetic resonance spectroscopy coupling (Review) 785(1997)65

Simple high-performance liquid chromatographic method for the determination of all seven vitamin B₆-related compounds 790(1997)83

Separation of neutral substances by non-aqueous capillary electrophoresis through interactions with cationic additives 792(1997)475

Development of a method for quantitation of retinol and retinyl palmitate in human serum using high-performance liquid chromatography–atmospheric pressure chemical ionization–mass spectrometry 794(1998)245

Overview of the applications of liquid chromatography–mass spectrometry interfacing systems in food analysis: naturally occurring substances in food 794(1998)263

Determination of vitamin B₆ (pyridoxamine, pyridoxal and pyridoxine) in pork meat and pork meat products by liquid chromatography 795(1998)383

Rapid determination of vitamins A and E in serum with surfactant as a diluent by column-switching high-performance liquid chromatography 798(1998)125

Volatile fatty acids

Determination of anions in landfill leachates by ion chromatography 770(1997)203

Volatile organic compounds

see also Hydrocarbons

Solid-phase microextraction of volatile organic compounds. Estimation of the sorption equilibrium from the Kováts index, effect of salinity and humic acids and the study of the kinetics by the development of an "agitated/static layer" model 761(1997)205

Improvement in breakthrough volume evaluation methods for light adsorbent traps employed for volatile organic compounds determination at atmospheric concentration levels 767(1997)153

Determination of volatile organic compounds in water by purge-and-trap gas chromatography coupled to atomic emission detection 768(1997)259

Developments in the analysis of petroleum hydrocarbons in oils, petroleum products and oil-spill-related environmental samples by gas chromatography (Review) 774(1997)51

Headspace gas chromatographic determination of the plant cuticle-air partition coefficients for monocyclic aromatic hydrocarbons as environmental compartment 774(1997)213

Occurrence and determination of organic pollutants in aerosol, precipitation, and sediment samples collected at Lake Balaton 774(1997)349

Purge-and-trap gas chromatography-mass spectrometry in the analysis of volatile organochlorine compounds in water 775(1997)267

Simple laboratory-made system for the determination of C₂-C₇ hydrocarbons relevant to photochemical smog pollution 777(1997)267

Determination of volatile compounds in Txakoli wine from Biscay by gas chromatography-mass spectrometry 778(1997)225

Evaluation of environmental levels of aromatic hydrocarbons in gasoline service stations by gas chromatography 778(1997)263

Refrigerated multibed adsorption in sampling and analysis of atmospheric light hydrocarbons at ppb (v/v) and sub-ppb (v/v) concentrations 778(1997)269

Hyphenated pervaporation-solid-phase preconcentration-gas chromatography for the determination of volatile organic compounds in solid samples 779(1997)352

Analysis of volatile organic compounds in air using retention indices together with a simple thermal desorption and cold trap method 787(1997)205

Effect of addition of a graphitized carbon black trap to a glass beads trap on the cryoconcentration of some non-methane hydrocarbons in ambient air 790(1997)117

Analysis of odour and taste problems in high-density polyethylene 791(1997)213

Determination of volatile organic compounds in marine biota 799(1998)233

Volatile sulfur compounds

Quantitation of volatile sulphur compounds in polluted waters 778(1997)329

W**Warfarin**

Enantiomeric properties of human albumin immobilized on porous silica supports coated with polymethacryloyl chloride 776(1997)37

Comparison of five methods for the study of drug-protein binding in affinity capillary electrophoresis 777(1997)311

Bonded cellulose-derived high-performance liquid chromatography chiral stationary phases. III. Effect of the reticulation of the cellulose derivative on performance 791(1997)37

Solvent versatility of bonded cellulose-derived chiral stationary phases for high-performance liquid chromatography and its consequences in column loadability 793(1998)239

Water, sub-critical

Sub- and supercritical fluid extraction of trichloropyridinol from soil prior to immunoassay 785(1997)329

Water, ultra-pure

Ultra-pure water for ion chromatography 770(1997)115

Weak acids

Calculation of the composition of sample zones in capillary zone electrophoresis. IV. Weak acids and system characteristics 764(1997)111

Weak acids, monovalent

Migration behaviour of monovalent weak acids in micellar electrokinetic chromatography. Mobility model versus retention model 765(1997)295

Whey proteins

Purification by reflux electrophoresis of whey proteins and of a recombinant protein expressed in *Dictyostelium discoideum* 773(1997)299

X**Xanthenes**

pH-Zone-refining countercurrent chromatography 753(1996)1

Effect of foaming power and foam stability on continuous concentration with foam counter-current chromatography 791(1997)53

Xanthommatin

Detection of the oxidative products of 3-hydroxykynurenine using high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–electron spin resonance spectrometry and high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–mass spectrometry 773(1997)23

Xanthenes

Liquid chromatography coupled to mass spectrometry and nuclear magnetic resonance spectroscopy for the screening of plant constituents 794(1998)299

Xylenes

Headspace gas chromatographic determination of the plant cuticle–air partition coefficients for monocyclic aromatic hydrocarbons as environmental compartment 774(1997)213

Evaluation of environmental levels of aromatic hydrocarbons in gasoline service stations by gas chromatography 778(1997)263

Effect of addition of a graphitized carbon black trap to a glass beads trap on the cryoconcentration of some non-methane hydrocarbons in ambient air 790(1997)117

Determination of volatile organic compounds in marine biota 799(1998)233

Xylenol orange

Surfactant-sensitized post-column reaction with xylenol orange for the determination of lanthanides by ion chromatography 770(1997)75

Xylose

Analysis of selected alkaloids and sugars in tobacco extract 775(1997)101

Y**Ytterbium**

Separation of samarium, gadolinium, terbium, dysprosium, erbium and ytterbium by high-speed countercurrent chromatography with organophosphate ester 766(1997)282

Yttrium

Simultaneous separation of yttrium and lanthanide ions by isotachopheresis 772(1997)357

Enrichment separation of rare earth elements by high-speed countercurrent chromatography in a multilayer coiled column 789(1997)381

Z**Zearalenols**

Determination of mycotoxins produced by *Fusarium* isolates from banana fruits by capillary gas chromatography and high-performance liquid chromatography 778(1997)363

Zearalenone

Determination of mycotoxins produced by *Fusarium* isolates from banana fruits by capillary gas chromatography and high-performance liquid chromatography 778(1997)363

Ziduvodine

New procedure for the use of high-performance liquid chromatography–electrospray ionization mass spectrometry for the analysis of nucleotides and oligonucleotides 777(1997)3

Zinc

Determination of kynurenic acid by capillary electrophoresis with laser-induced fluorescence detection 781(1997)81

Zn²⁺-selective purification of recombinant proteins from the milk of transgenic animals 799(1998)125

Zinc dithiophosphates

Lability of zinc dialkyldithiophosphates under reversed-phase hplc conditions 765(1997)181

Zirconia

Factors influencing polybutadiene deposition within porous chromatographic zirconia 791(1997)45