



ELSEVIER

Journal of Chromatography A, 848 (1999) 557–566

JOURNAL OF
CHROMATOGRAPHY A

Subject Index Vol. 848

Acids, aromatic

Macrocyclic polyamine as a selective modifier in a bonded-phase capillary column for the electrophoretic separation of aromatic acids 848(1999)401

Affinity supports

Selective adsorption of poly-His tagged glutaryl acylase on tailor-made metal chelate supports 848(1999)61

Air analysis

Diffusion technique for the production of gas standards for atmospheric measurements 848(1999)239

Analysis of brominated compounds in background air by gas chromatography–high-resolution mass spectrometry 848(1999)261

Evaluation of solid-phase microextraction for sampling of volatile organic sulfur compounds in air for subsequent gas chromatographic analysis with atomic emission detection 848(1999)305

Alkaline-earth cations

Study of sugar acids separation by high-performance anion-exchange chromatography–pulsed amperometric detection using alkaline eluents spiked with Ba^{2+} , Sr^{2+} , or Ca^{2+} as acetate or nitrate salts 848(1999)71

Alkaloids

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Alkanols

Alkanol gas–liquid partition coefficients in squalane measured with packed columns. A revision of measurement methods 848(1999)203

Amines

Continuous solid-phase extraction and dansylation of low-molecular-mass amines coupled on-line with liquid chromatography and peroxyoxalate chemiluminescence-based detection 848(1999)105

Amino acids

Fluorescence properties of carbazole-*N*-(2-methyl)acetyl chloride and determination of amino compounds via high-performance liquid chromatography with pre-column fluorescence derivatization 848(1999)117

Amphetamine-like compounds

Analysis and confirmation of synthetic anorexics in adulterated traditional Chinese medicines by high-performance capillary electrophoresis 848(1999)537

Antibiotics

Enantiomeric separations by capillary electrochromatography using a macrocyclic antibiotic chiral stationary phase 848(1999)375

Antioxidants

Gas chromatographic flow method for the preconcentration and simultaneous determination of antioxidant and preservative additives in fatty foods 848(1999)529

Argentation

Positional and configurational separation of fatty acid isomers by micro reversed-phase liquid chromatography with an Ag^+ -containing mobile phase 848(1999)83

Atropine

Enantiomeric separation of atropine in *Scopolia* extract and *Scopolia Rhizome* by capillary electrophoresis using cyclodextrins as chiral selectors 848(1999)465

Benzaldehydes

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Benzoic acids

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Benzyl alcohols

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Bilirubins

Accurate and sensitive high-performance liquid chromatographic method for geometrical and structural photoisomers of bilirubin IX α using the relative molar absorptivity values 848(1999)169

Biogenic amines

Fluorescence properties of carbazole-*N*-(2-methyl)acetyl chloride and determination of amino compounds via high-performance liquid chromatography with pre-column fluorescence derivatization 848(1999)117

Buffer composition

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Macrocyclic polyamine as a selective modifier in a bonded-phase capillary column for the electrophoretic separation of aromatic acids 848(1999)401

Capillary zone electrophoresis of rigid submicron-sized particles in polyacrylamide solution. Selectivity, peak spreading and resolution 848(1999)443

Caffeine

Rapid analysis of methylated xanthenes in teas by an improved high-performance liquid chromatographic method using a polyvinylpyrrolidone pre-column 848(1999)523

Carbazolemethylacetyl chloride

Fluorescence properties of carbazole-*N*-(2-methyl)acetyl chloride and determination of amino compounds via high-performance liquid chromatography with pre-column fluorescence derivatization 848(1999)117

Carbohydrates

Study of sugar acids separation by high-performance anion-exchange chromatography–pulsed amperometric detection using alkaline eluents spiked with Ba²⁺, Sr²⁺, or Ca²⁺ as acetate or nitrate salts 848(1999)71

Cheese

Gas chromatographic flow method for the preconcentration and simultaneous determination of antioxidant and preservative additives in fatty foods 848(1999)529

Chemiluminescence detection

Continuous solid-phase extraction and dansylation of low-molecular-mass amines coupled on-line with liquid chromatography and peroxyoxalate chemiluminescence-based detection 848(1999)105

Chemometrics

Systematic optimization of peak capacity for pesticide separation in micellar electrokinetic chromatography 848(1999)347

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Chiral stationary phases, electrochromatography

Enantiomeric separations by capillary electrochromatography using a macrocyclic antibiotic chiral stationary phase 848(1999)375

Chiral stationary phases, SFC

Thermodynamic study and separation mechanism of diltiazem optical isomers in packed-column supercritical fluid chromatography 848(1999)337

Colloids

Size separation of colloidal nanoparticles using a miniscale isoelectric focusing technique 848(1999)485

Column switching

On-line trace enrichment for the simultaneous determination of microcystins in aqueous samples using high-performance liquid chromatography with diode-array detection 848(1999)179

Complexation

Analysis of metal complexes in the presence of mixed ion pairing additives in capillary electrophoresis 848(1999)473

Determination of iron(II) in natural waters by capillary zone electrophoresis using on-capillary complexation with 2,4,6-tri(2'-pyridyl)-1,3,5-triazine 848(1999)491

Computer-assisted selection of mobile phases

New approach to the computer-assisted selection of mobile phases for high-performance liquid chromatography on the basis of the Snyder theory 848(1999)1

Computer simulation

Automatic prediction of retention times in programmed-pressure isothermal gas chromatography 848(1999)229

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Coumarin

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Cyclodextrins

Enantiomeric separation of atropine in *Scopolia* extract and *Scopolia Rhizome* by capillary electrophoresis using cyclodextrins as chiral selectors 848(1999)465

Cytochromes

P450 monooxygenase in biotechnology. I. Single-step, large-scale purification method for cytochrome P450 BM-3 by anion-exchange chromatography 848(1999)149

Derivatization, electrophoresis

Size separation of colloidal nanoparticles using a miniscale isoelectric focusing technique 848(1999)485

Derivatization, LC

Continuous solid-phase extraction and dansylation of low-molecular-mass amines coupled on-line with liquid chromatography and peroxyoxalate chemiluminescence-based detection 848(1999)105

Fluorescence properties of carbazole-*N*-(2-methyl)acetyl chloride and determination of amino compounds via high-performance liquid chromatography with pre-column fluorescence derivatization 848(1999)117

Detection, LC

Comparison of the responses of triacylglycerols with an evaporative visible light scattering detector used in conventional, micro and capillary liquid chromatography 848(1999)95

Continuous solid-phase extraction and dansylation of low-molecular-mass amines coupled on-line with liquid chromatography and peroxyoxalate chemiluminescence-based detection 848(1999)105

Diltiazem

Thermodynamic study and separation mechanism of diltiazem optical isomers in packed-column supercritical fluid chromatography 848(1999)337

Dinitrobenzoyl leucine

Enantiomeric separations by capillary electrochromatography using a macrocyclic antibiotic chiral stationary phase 848(1999)375

DNA

Heparin elution of transcription factors from DNA-Sepharose columns 848(1999)131

Electrochemical detection

Study of sugar acids separation by high-performance anion-exchange chromatography-pulsed amperometric detection using alkaline eluents spiked with Ba²⁺, Sr²⁺, or Ca²⁺ as acetate or nitrate salts 848(1999)71

Electrochromatography

Enantiomeric separations by capillary electrochromatography using a macrocyclic antibiotic chiral stationary phase 848(1999)375

Elution window

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Enantiomer separation

Chromatographic separation of α -pinene isomerization products 848(1999)297

Thermodynamic study and separation mechanism of diltiazem optical isomers in packed-column supercritical fluid chromatography 848(1999)337

Enantiomeric separations by capillary electrochromatography using a macrocyclic antibiotic chiral stationary phase 848(1999)375

Enantiomeric separation of atropine in *Scopolia* extract and *Scopolia Rhizome* by capillary electrophoresis using cyclodextrins as chiral selectors 848(1999)465

Enthalpy

Thermodynamic properties for the solute transfer from the mobile to the stationary phase in reversed phase liquid chromatography obtained by squalane-impregnated C₁₈ bonded phase 848(1999)9

Entropy

Thermodynamic properties for the solute transfer from the mobile to the stationary phase in reversed phase liquid chromatography obtained by squalane-impregnated C₁₈ bonded phase 848(1999)9

Environmental analysis

Continuous solid-phase extraction and dansylation of low-molecular-mass amines coupled on-line with liquid chromatography and peroxyoxalate chemiluminescence-based detection 848(1999)105

Diffusion technique for the production of gas standards for atmospheric measurements 848(1999)239

Analysis of brominated compounds in background air by gas chromatography-high-resolution mass spectrometry 848(1999)261

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Evaluation of dynamic headspace and purge-and-trap techniques for the high-resolution gas chromatography analysis of nitrous oxide in seawater 848(1999)327

Determination of iron(II) in natural waters by capillary zone electrophoresis using on-capillary complexation with 2,4,6-tri(2'-pyridyl)-1,3,5-triazine 848(1999)491

Quantitative determination of sulfur-containing anions in complex matrices with capillary electrophoresis and conductivity detection 848(1999)503

Capillary electrophoretic determination of inorganic anions in the drainage and surface water samples 848(1999)545

Enzymes

Selective adsorption of poly-His tagged glutaryl acylase on tailor-made metal chelate supports 848(1999)61

Europium

Poly(7-oxanorborn-2-ene-5,6-dicarboxylate)-coated silica prepared by ring-opening metathesis polymerization for the selective enrichment of radioactive lanthanides 848(1999)193

Evaporative light scattering detection

Comparison of the responses of triacylglycerols with an evaporative visible light scattering detector used in conventional, micro and capillary liquid chromatography 848(1999)95

Experimental design

Systematic optimization of peak capacity for pesticide separation in micellar electrokinetic chromatography 848(1999)347

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Extraction methods

Poly(7-oxanorborn-2-ene-5,6-dicarboxylate)-coated silica prepared by ring-opening metathesis polymerization for the selective enrichment of radioactive lanthanides 848(1999)193

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Evaluation of solid-phase microextraction for sampling of volatile organic sulfur compounds in air for subsequent gas chromatographic analysis with atomic emission detection 848(1999)305

Fats

Positional and configurational separation of fatty acid isomers by micro reversed-phase liquid chromatography with an Ag⁺-containing mobile phase 848(1999)83

Fatty acids

Positional and configurational separation of fatty acid isomers by micro reversed-phase liquid chromatography with an Ag⁺-containing mobile phase 848(1999)83

Flash chromatography

Purification of closely eluting hydrophobic microcystins (peptide cyanotoxins) by normal-phase and reversed-phase flash chromatography 848(1999)515

Food analysis

Determination of the mycotoxin moniliformin in cultures of *Fusarium subglutinans* and in naturally contaminated maize by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 848(1999)185

Quantitative determination of β -ionone in red wines and grapes of Bordeaux using a stable isotope dilution assay 848(1999)317

Macrocyclic polyamine as a selective modifier in a bonded-phase capillary column for the electrophoretic separation of aromatic acids 848(1999)401

Rapid analysis of methylated xanthenes in teas by an improved high-performance liquid chromatographic method using a polyvinylpyrrolidone pre-column 848(1999)523

Gas chromatographic flow method for the preconcentration and simultaneous determination of antioxidant and preservative additives in fatty foods 848(1999)529

Fruit

Quantitative determination of β -ionone in red wines and grapes of Bordeaux using a stable isotope dilution assay 848(1999)317

Fusarium subglutinans

Determination of the mycotoxin moniliformin in cultures of *Fusarium subglutinans* and in naturally contaminated maize by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 848(1999)185

Galanthamine

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Gas standards

Diffusion technique for the production of gas standards for atmospheric measurements 848(1999)239

Geometrical isomers

Accurate and sensitive high-performance liquid chromatographic method for geometrical and structural photoisomers of bilirubin IX α using the relative molar absorptivity values 848(1999)169

Glutaryl acylase

Selective adsorption of poly-His tagged glutaryl acylase on tailor-made metal chelate supports 848(1999)61

Glycidyl methacrylate–ethylenedimethacrylate monoliths

Isocratic separations on thin glycidyl methacrylate–ethylenedimethacrylate monoliths 848(1999)51

Gold

Size separation of colloidal nanoparticles using a miniscale isoelectric focusing technique 848(1999)485

Grapes

Quantitative determination of β -ionone in red wines and grapes of Bordeaux using a stable isotope dilution assay 848(1999)317

Halocarbons

Analysis of brominated compounds in background air by gas chromatography–high-resolution mass spectrometry 848(1999)261

Headspace analysis

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Evaluation of dynamic headspace and purge-and-trap techniques for the high-resolution gas chromatography analysis of nitrous oxide in seawater 848(1999)327

Heparin

Heparin elution of transcription factors from DNA-Sepharose columns 848(1999)131

Hollow-fiber membranes

Ionic crosslinking of SO₃H-group-containing graft chains helps to capture lysozyme in a permeation mode 848(1999)161

Hydrocarbons

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Hyoscyamine

Enantiomeric separation of atropine in *Scopolia* extract and *Scopolia Rhizome* by capillary electrophoresis using cyclodextrins as chiral selectors 848(1999)465

Immobilized metal-ion affinity chromatography

Selective adsorption of poly-His tagged glutaryl acylase on tailor-made metal chelate supports 848(1999)61

Immunoassays

Noncompetitive immunoassays using protein G affinity capillary chromatography and capillary electrophoresis with laser-induced fluorescence detection 848(1999)139

Inorganic anions

Quantitative determination of sulfur-containing anions in complex matrices with capillary electrophoresis and conductivity detection 848(1999)503

Capillary electrophoretic determination of inorganic anions in the drainage and surface water samples 848(1999)545

Instrumentation

Diffusion technique for the production of gas standards for atmospheric measurements 848(1999)239

Ionones

Quantitative determination of β-ionone in red wines and grapes of Bordeaux using a stable isotope dilution assay 848(1999)317

Iron

Determination of iron(II) in natural waters by capillary zone electrophoresis using on-capillary complexation with 2,4,6-tri(2'-pyridyl)-1,3,5-triazine 848(1999)491

Isoelectric focusing

Size separation of colloidal nanoparticles using a miniscale isoelectric focusing technique 848(1999)485

Isomers, configurational

Positional and configurational separation of fatty acid isomers by micro reversed-phase liquid chromatography with an Ag⁺-containing mobile phase 848(1999)83

Isomers, positional

Positional and configurational separation of fatty acid isomers by micro reversed-phase liquid chromatography with an Ag⁺-containing mobile phase 848(1999)83

Kavalactones

Qualitative and quantitative micellar electrokinetic chromatography of kavalactones from dry extracts of *Piper methysticum* Forst. and commercial drugs 848(1999)457

Lanthanides

Poly(7-oxanorborn-2-ene-5,6-dicarboxylate)-coated silica prepared by ring-opening metathesis polymerization for the selective enrichment of radioactive lanthanides 848(1999)193

Linear solvation energy relationships

Global linear solvation energy relationships for retention prediction in reversed-phase liquid chromatography 848(1999)21

Linear solvent strength theory

Global linear solvation energy relationships for retention prediction in reversed-phase liquid chromatography 848(1999)21

Lysozyme

Ionic crosslinking of SO₃H-group-containing graft chains helps to capture lysozyme in a permeation mode 848(1999)161

Macrocyclic antibiotics

Enantiomeric separations by capillary electrochromatography using a macrocyclic antibiotic chiral stationary phase 848(1999)375

Margarines

Positional and configurational separation of fatty acid isomers by micro reversed-phase liquid chromatography with an Ag⁺-containing mobile phase 848(1999)83

Mathematical modeling

Systematic optimization of peak capacity for pesticide separation in micellar electrokinetic chromatography 848(1999)347

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Peptide mobility and peptide mapping in capillary zone electrophoresis. Experimental determination and theoretical simulation 848(1999)417

Membranes

Ionic crosslinking of SO₃H-group-containing graft chains helps to capture lysozyme in a permeation mode 848(1999)161

Isocratic separations on thin glycidyl methacrylate–ethylenedimethacrylate monoliths 848(1999)51

Metal cations

Analysis of metal complexes in the presence of mixed ion pairing additives in capillary electrophoresis 848(1999)473

Determination of iron(II) in natural waters by capillary zone electrophoresis using on-capillary complexation with 2,4,6-tri(2'-pyridyl)-1,3,5-triazine 848(1999)491

Metal complexes

Analysis of metal complexes in the presence of mixed ion pairing additives in capillary electrophoresis 848(1999)473

Metals

Size separation of colloidal nanoparticles using a miniscale isoelectric focusing technique 848(1999)485

Micellar electrokinetic chromatography

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Microcystins

On-line trace enrichment for the simultaneous determination of microcystins in aqueous samples using high-performance liquid chromatography with diode-array detection 848(1999)179

Purification of closely eluting hydrophobic microcystins (peptide cyanotoxins) by normal-phase and reversed-phase flash chromatography 848(1999)515

Mobile phase composition

Study of sugar acids separation by high-performance anion-exchange chromatography–pulsed amperometric detection using alkaline eluents spiked with Ba²⁺, Sr²⁺, or Ca²⁺ as acetate or nitrate salts 848(1999)71

Mobile phase selection

New approach to the computer-assisted selection of mobile phases for high-performance liquid chromatography on the basis of the Snyder theory 848(1999)1

Molecularly imprinted polymers

Study of the nature of recognition in molecularly imprinted polymers, II [1]. Influence of monomer–template ratio and sample load on retention and selectivity 848(1999)39

Moniliformin

Determination of the mycotoxin moniliformin in cultures of *Fusarium subglutinans* and in naturally contaminated maize by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 848(1999)185

Monoliths

Isocratic separations on thin glycidyl methacrylate–ethylenedimethacrylate monoliths 848(1999)51

Monomer–template ratio

Study of the nature of recognition in molecularly imprinted polymers, II [1]. Influence of monomer–template ratio and sample load on retention and selectivity 848(1999)39

Monoterpenes

Chromatographic separation of α -pinene isomerization products 848(1999)297

Mycotoxins

Determination of the mycotoxin moniliformin in cultures of *Fusarium subglutinans* and in naturally contaminated maize by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 848(1999)185

Neural networks, artificial

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Nicotine

Study of the nature of recognition in molecularly imprinted polymers, II [1]. Influence of monomer–template ratio and sample load on retention and selectivity 848(1999)39

Nitrotoluenes

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Nitrous oxide

Evaluation of dynamic headspace and purge-and-trap techniques for the high-resolution gas chromatography analysis of nitrous oxide in seawater 848(1999)327

Nitroux

Evaluation of dynamic headspace and purge-and-trap techniques for the high-resolution gas chromatography analysis of nitroux oxide in seawater 848(1999)327

Non-ionic organic compounds

New approach to the computer-assisted selection of mobile phases for high-performance liquid chromatography on the basis of the Snyder theory 848(1999)1

Nuclear magnetic resonance spectrometry

Characterization of stationary phases for gas chromatography by ²⁹Si NMR spectroscopy 848(1999)251

Oils

Gas chromatographic flow method for the preconcentration and simultaneous determination of antioxidant and preservative additives in fatty foods 848(1999)529

Oligonucleotides

Isocratic separations on thin glycidyl methacrylate–ethylenedimethacrylate monoliths 848(1999)51

Composition dependent separation of oligonucleotides by capillary electrophoresis in acidic buffers with application to the quality control of synthetic oligonucleotides 848(1999)435

On-capillary derivatization

Determination of iron(II) in natural waters by capillary zone electrophoresis using on-capillary complexation with 2,4,6-tri(2'-pyridyl)-1,3,5-triazine 848(1999)491

Optimization

Systematic optimization of peak capacity for pesticide separation in micellar electrokinetic chromatography 848(1999)347

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Organic acids

Study of sugar acids separation by high-performance anion-exchange chromatography–pulsed amperometric detection using alkaline eluents spiked with Ba^{2+} , Sr^{2+} , or Ca^{2+} as acetate or nitrate salts 848(1999)71

Macrocyclic polyamine as a selective modifier in a bonded-phase capillary column for the electrophoretic separation of aromatic acids 848(1999)401

Organobromine compounds

Analysis of brominated compounds in background air by gas chromatography–high-resolution mass spectrometry 848(1999)261

Organosulfur compounds

Evaluation of solid-phase microextraction for sampling of volatile organic sulfur compounds in air for subsequent gas chromatographic analysis with atomic emission detection 848(1999)305

Particles, submicron-sized

Capillary zone electrophoresis of rigid submicron-sized particles in polyacrylamide solution. Selectivity, peak spreading and resolution 848(1999)443

Partition coefficients

Alkanol gas–liquid partition coefficients in squalane measured with packed columns. A revision of measurement methods 848(1999)203

Peak shape

Use of artificial neural networks in capillary zone electrophoresis 848(1999)365

Peptides

Peptide mobility and peptide mapping in capillary zone electrophoresis. Experimental determination and theoretical simulation 848(1999)417

Perfumery compounds

Temperature dependence of the retention index for perfumery compounds on a SE-30 glass capillary column. II. The hyperbolic equation 848(1999)215

Pesticides

Systematic optimization of peak capacity for pesticide separation in micellar electrokinetic chromatography 848(1999)347

Petroleum products

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Pharmaceutical analysis

Macrocyclic polyamine as a selective modifier in a bonded-phase capillary column for the electrophoretic separation of aromatic acids 848(1999)401

Qualitative and quantitative micellar electrokinetic chromatography of kavalactones from dry extracts of *Piper methysticum* Forst. and commercial drugs 848(1999)457

Enantiomeric separation of atropine in *Scopolia* extract and *Scopolia Rhizome* by capillary electrophoresis using cyclodextrins as chiral selectors 848(1999)465

Analysis and confirmation of synthetic anorexics in adulterated traditional Chinese medicines by high-performance capillary electrophoresis 848(1999)537

Pinenes

Chromatographic separation of α -pinene isomerization products 848(1999)297

Piper methysticum

Qualitative and quantitative micellar electrokinetic chromatography of kavalactones from dry extracts of *Piper methysticum* Forst. and commercial drugs 848(1999)457

Piperonal

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Plant materials

Qualitative and quantitative micellar electrokinetic chromatography of kavalactones from dry extracts of *Piper methysticum* Forst. and commercial drugs 848(1999)457

Polyacrylamide

Capillary zone electrophoresis of rigid submicron-sized particles in polyacrylamide solution. Selectivity, peak spreading and resolution 848(1999)443

Polyamines

Macrocyclic polyamine as a selective modifier in a bonded-phase capillary column for the electrophoretic separation of aromatic acids 848(1999)401

Polynuclear aromatic hydrocarbons

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Polysiloxanes

Characterization of stationary phases for gas chromatography by ^{29}Si NMR spectroscopy 848(1999)251

Polyvinylpyrrolidone columns

Rapid analysis of methylated xanthenes in teas by an improved high-performance liquid chromatographic method using a polyvinylpyrrolidone pre-column 848(1999)523

Preparative chromatography

P450 monooxygenase in biotechnology. I. Single-step, large-scale purification method for cytochrome P450 BM-3 by anion-exchange chromatography 848(1999)149

Pressurized liquid extraction

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Programmed-pressure gas chromatography

Automatic prediction of retention times in programmed-pressure isothermal gas chromatography 848(1999)229

Purge-and-trap methods

Evaluation of dynamic headspace and purge-and-trap techniques for the high-resolution gas chromatography analysis of nitrous oxide in seawater 848(1999)327

Polystyrene

Capillary zone electrophoresis of rigid submicron-sized particles in polyacrylamide solution. Selectivity, peak spreading and resolution 848(1999)443

Preservatives

Gas chromatographic flow method for the preconcentration and simultaneous determination of antioxidant and preservative additives in fatty foods 848(1999)529

Promethium

Poly(7-oxanorborn-2-ene-5,6-dicarboxylate)-coated silica prepared by ring-opening metathesis polymerization for the selective enrichment of radioactive lanthanides 848(1999)193

Proteins

Heparin elution of transcription factors from DNA-Sepharose columns 848(1999)131

Noncompetitive immunoassays using protein G affinity capillary chromatography and capillary electrophoresis with laser-induced fluorescence detection 848(1999)139

P450 monooxygenase in biotechnology. I. Single-step, large-scale purification method for cytochrome P450 BM-3 by anion-exchange chromatography 848(1999)149

Ionic crosslinking of SO_3H -group-containing graft chains helps to capture lysozyme in a permeation mode 848(1999)161

Response surface modelling

Systematic optimization of peak capacity for pesticide separation in micellar electrokinetic chromatography 848(1999)347

Retention behaviour

Study of the nature of recognition in molecularly imprinted polymers, II [1]. Influence of monomer-template ratio and sample load on retention and selectivity 848(1999)39

Retention indices

Temperature dependence of the retention index for perfumery compounds on a SE-30 glass capillary column. II. The hyperbolic equation 848(1999)215

Retention mechanism

Thermodynamic study and separation mechanism of diltiazem optical isomers in packed-column supercritical fluid chromatography 848(1999)337

Retention models

Global linear solvation energy relationships for retention prediction in reversed-phase liquid chromatography 848(1999)21

Retention prediction

New approach to the computer-assisted selection of mobile phases for high-performance liquid chromatography on the basis of the Snyder theory 848(1999)1

Global linear solvation energy relationships for retention prediction in reversed-phase liquid chromatography 848(1999)21

Automatic prediction of retention times in programmed-pressure isothermal gas chromatography 848(1999)229

Retention times

Automatic prediction of retention times in programmed-pressure isothermal gas chromatography 848(1999)229

Ring-opening metathesis polymerization

Poly(7-oxanorborn-2-ene-5,6-dicarboxylate)-coated silica prepared by ring-opening metathesis polymerization for the selective enrichment of radioactive lanthanides 848(1999)193

Sample handling

Continuous solid-phase extraction and dansylation of low-molecular-mass amines coupled on-line with liquid chromatography and peroxyoxalate chemiluminescence-based detection 848(1999)105

Evaluation of dynamic headspace and purge-and-trap techniques for the high-resolution gas chromatography analysis of nitrous oxide in seawater 848(1999)327

Sample load

Study of the nature of recognition in molecularly imprinted polymers, II [1]. Influence of monomer–template ratio and sample load on retention and selectivity 848(1999)39

Scopolamine

Enantiomeric separation of atropine in *Scopolia* extract and *Scopolia Rhizome* by capillary electrophoresis using cyclodextrins as chiral selectors 848(1999)465

Selectivity

Study of the nature of recognition in molecularly imprinted polymers, II [1]. Influence of monomer–template ratio and sample load on retention and selectivity 848(1999)39

Silica, coating

Poly(7-oxanorborn-2-ene-5,6-dicarboxylate)-coated silica prepared by ring-opening metathesis polymerization for the selective enrichment of radioactive lanthanides 848(1999)193

Silver ion complexation

Positional and configurational separation of fatty acid isomers by micro reversed-phase liquid chromatography with an Ag⁺-containing mobile phase 848(1999)83

Snyder theory

New approach to the computer-assisted selection of mobile phases for high-performance liquid chromatography on the basis of the Snyder theory 848(1999)1

Soil

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Solid-phase microextraction

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Solute transfer

Thermodynamic properties for the solute transfer from the mobile to the stationary phase in reversed phase liquid chromatography obtained by squalane-impregnated C₁₈ bonded phase 848(1999)9

Squalane

Thermodynamic properties for the solute transfer from the mobile to the stationary phase in reversed phase liquid chromatography obtained by squalane-impregnated C₁₈ bonded phase 848(1999)9

Alkanol gas–liquid partition coefficients in squalane measured with packed columns. A revision of measurement methods 848(1999)203

Stationary phases, GC

Characterization of stationary phases for gas chromatography by ²⁹Si NMR spectroscopy 848(1999)251

Stationary phases, LC

Thermodynamic properties for the solute transfer from the mobile to the stationary phase in reversed phase liquid chromatography obtained by squalane-impregnated C₁₈ bonded phase 848(1999)9

Study of the nature of recognition in molecularly imprinted polymers, II [1]. Influence of monomer–template ratio and sample load on retention and selectivity 848(1999)39

Isocratic separations on thin glycidyl methacrylate–ethylenedimethacrylate monoliths 848(1999)51

Ionic crosslinking of SO₃H-group-containing graft chains helps to capture lysozyme in a permeation mode 848(1999)161

Rapid analysis of methylated xanthenes in teas by an improved high-performance liquid chromatographic method using a polyvinylpyrrolidone pre-column 848(1999)523

Structural isomers

Accurate and sensitive high-performance liquid chromatographic method for geometrical and structural photoisomers of bilirubin IX α using the relative molar absorptivity values 848(1999)169

Styrylpyrones

Qualitative and quantitative micellar electrokinetic chromatography of kavalactones from dry extracts of *Piper methysticum* Forst. and commercial drugs 848(1999)457

Sugar acids

Study of sugar acids separation by high-performance anion-exchange chromatography–pulsed amperometric detection using alkaline eluents spiked with Ba²⁺, Sr²⁺, or Ca²⁺ as acetate or nitrate salts 848(1999)71

Sulfur compounds

Quantitative determination of sulfur-containing anions in complex matrices with capillary electrophoresis and conductivity detection 848(1999)503

Tars

Chemical characterization and screening of hydrocarbon pollution in industrial soils by headspace solid-phase microextraction 848(1999)279

Tea

Rapid analysis of methylated xanthenes in teas by an improved high-performance liquid chromatographic method using a polyvinylpyrrolidone pre-column 848(1999)523

Terpenes

Chromatographic separation of α -pinene isomerization products 848(1999)297

Theobromine

Rapid analysis of methylated xanthenes in teas by an improved high-performance liquid chromatographic method using a polyvinylpyrrolidone pre-column 848(1999)523

Thermal redistribution

Characterization of stationary phases for gas chromatography by ^{29}Si NMR spectroscopy 848(1999)215

Thermodynamic parameters

Temperature dependence of the retention index for perfumery compounds on a SE-30 glass capillary column. II. The hyperbolic equation 848(1999)215

Thermodynamics

Thermodynamic study and separation mechanism of diltiazem optical isomers in packed-column supercritical fluid chromatography 848(1999)337

Toxins

On-line trace enrichment for the simultaneous determination of microcystins in aqueous samples using high-performance liquid chromatography with diode-array detection 848(1999)179

Purification of closely eluting hydrophobic microcystins (peptide cyanotoxins) by normal-phase and reversed-phase flash chromatography 848(1999)515

Triacylglycerols

Comparison of the responses of triacylglycerols with an evaporative visible light scattering detector used in conventional, micro and capillary liquid chromatography 848(1999)95

Triazines

Systematic optimization of peak capacity for pesticide separation in micellar electrokinetic chromatography 848(1999)347

Tripyridyltriazine

Determination of iron(II) in natural waters by capillary zone electrophoresis using on-capillary complexation with 2,4,6-tri(2'-pyridyl)-1,3,5-triazine 848(1999)491

Tryptophan

Enantiomeric separations by capillary electrochromatography using a macrocyclic antibiotic chiral stationary phase 848(1999)375

Vanillic acid

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Vanillin

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. I. Selection of surfactant, variation of pH and addition of organic modifiers or inorganic metal salts 848(1999)387

Volatile organic compounds

Diffusion technique for the production of gas standards for atmospheric measurements 848(1999)239

Evaluation of solid-phase microextraction for sampling of volatile organic sulfur compounds in air for subsequent gas chromatographic analysis with atomic emission detection 848(1999)305

Volatile sulfur compounds

Evaluation of solid-phase microextraction for sampling of volatile organic sulfur compounds in air for subsequent gas chromatographic analysis with atomic emission detection 848(1999)305

Water analysis

Continuous solid-phase extraction and dansylation of low-molecular-mass amines coupled on-line with liquid chromatography and peroxyoxalate chemiluminescence-based detection 848(1999)105

On-line trace enrichment for the simultaneous determination of microcystins in aqueous samples using high-performance liquid chromatography with diode-array detection 848(1999)179

Evaluation of dynamic headspace and purge-and-trap techniques for the high-resolution gas chromatography analysis of nitrous oxide in seawater 848(1999)327

Determination of iron(II) in natural waters by capillary zone electrophoresis using on-capillary complexation with 2,4,6-tri(2'-pyridyl)-1,3,5-triazine 848(1999)491

Quantitative determination of sulfur-containing anions in complex matrices with capillary electrophoresis and conductivity detection 848(1999)503

Capillary electrophoretic determination of inorganic anions in the drainage and surface water samples 848(1999)545

Wine

Quantitative determination of β -ionone in red wines and grapes of Bordeaux using a stable isotope dilution assay 848(1999)317

Xanthenes

Rapid analysis of methylated xanthenes in teas by an improved high-performance liquid chromatographic method using a polyvinylpyrrolidone pre-column 848(1999)523