



ELSEVIER

Journal of Chromatography A, 852 (1999) 607

---

---

JOURNAL OF  
CHROMATOGRAPHY A

---

---

### Erratum

## Erratum to “Prediction of peptide retention time in normal-phase liquid chromatography”

[J. Chromatogr. A, Vol. 811 (1998) 61–67]<sup>☆</sup>

Tatsunari Yoshida

*Tosoh Corporation, Nanyo Research Center Analytical Laboratory, 4560 Kaisei-chou, Shinnanyo-shi, Yamaguchi 764-8501, Japan*

Received 7 November 1997; received in revised form 16 March 1998; accepted 16 March 1998

*Keywords:* Retention prediction; Peptides

---

On p. 62 in Eq. (5), the first  $t_D$  should read  $t_0$ ; the correct equation is:

$$t_g i = 1/bi \{ [bi \cdot (S + 1) \cdot (t_0 \cdot k'(0) - t_D) + a^{S+1}]^{1/(S+1)} - a \} + t_0 + t_D \quad (5)$$

On p. 62 in Eq. (8), the plus sign before “(S)” should be a multiplication sign; the correct equation is:

$$k'(0) = \{ [a + b1(t_g 1 - t_0 - t_D)]^{1(S+1)} / [b1 \cdot (S + 1)] + t_D \} / t_0 \quad (8)$$

---

<sup>☆</sup>PII of original article: S0021-9673(98)00216-7