



ELSEVIER

Journal of Chromatography A, 852 (1999) 608

JOURNAL OF
CHROMATOGRAPHY A

Erratum

Erratum to “Prediction of peptide retention times in normal-phase liquid chromatography with only a single gradient run”

[J. Chromatogr. A, 841 (1999) 19–32][☆]

Tatsunari Yoshida*, Tadashi Okada

Tosoh Corporation, Nanyo Research Center Analytical Laboratory, 4560 Kaisei-chou, Shinnanyo-shi, Yamaguchi 764-8501, Japan

Received 13 November 1998; received in revised form 19 February 1999; accepted 26 February 1999

Keywords: Retention parameters; Gradient elution; Peptides

On p. 20 in Eq. (8) a closing parenthesis is missing after the log term in the denominator. The correct equation is:

$$k(0)_{\text{NPLC}} = \left(\frac{t_0 \cdot [a + b1 \cdot (\text{tg}1 - t_0 - t_D)]^{\frac{1}{p + q^e \log k(0)_{\text{NPLC}} + 1}}}{[b1 \cdot (p + q \cdot e \log k(0)_{\text{NPLC}}) + t_D]} \right) \quad (8)$$

[☆]PII of original article: S0021-9673(99)00284-8.

*Corresponding author.