

Journal of Chromatography A, 852 (1999) 613-622

JOURNAL OF CHROMATOGRAPHY A

# Subject Index Vol. 852

#### Adsorbents

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

## Adsorbers

Large-scale membrane adsorbers 852(1999)73

#### Adsorption isotherms

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

### Affinity chromatography

Affinity purification of *Schistosoma japonicum* glutathione-Stransferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

Affinity chromatography of human estrogen receptor- $\alpha$  expressed in *Saccharomyces cerevisiae*. Combination of heparinand 17 $\beta$ -estradiol-affinity chromatography 852(1999)161

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

Experimental studies on affinity chromatography in an electric field 852(1999)319

#### Affinity sorbents

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105

# Air analysis

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

## Albumin

Experimental studies on affinity chromatography in an electric field 852(1999)319

#### Alkaloids

Optimization study for the reversed-phase ion-pair liquid chromatographic determination of nicotine in commercial tobacco products 852(1999)451

#### Alkanes

Revision of a theoretical expression for gas-liquid chromatographic retention 852(1999)583

# Alkyl nitrates

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a  $\beta$ -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

#### Amberlite XAD-7 resin

Removal of detergent and solvent from solvent-detergenttreated immunoglobulins 852(1999)87

#### Amino acids

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

## Amphetamines

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

#### Angiotensin

New polymer-based prepacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

#### Anilines

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

#### Antibodies

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

0021-9673/99/\$ – see front matter © 1999 Elsevier Science B.V. All right reserved PII: S0021-9673(99)00792-X

Quantitative fast fractionation of a pool of polyclonal antibodies by immunoaffinity membrane chromatography 852(1999)129

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

# Antigens

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

# Argininealkyamides

Qualitive and quantitative analysis of new alkyl amide arginine surfactants by high-performance liquid chromatography and capillary electrophoresis 852(1999)499

## Bacillus spp

Isolation and characterization of cyclic  $\alpha$ -(1 $\rightarrow$ 4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

# **Benzoic** acids

pH effects on micelle-water partitioning determined by micellar electrokinetic chromatography 852(1999)559

# **Binding studies**

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

## **β-Blockers**

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

## **Buffer composition**

Separation of proteins and peptides by capillary electrophoresis in acid buffers containing high concentrations of surfactants 852(1999)325

## Carbohydrates

Isolation and characterization of cyclic  $\alpha$ -(1 $\rightarrow$ 4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

#### Carbon dioxide

Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383

## Carbon monoxide

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum-rhodium alloy catalysts 852(1999)507

## Catalysts

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum-rhodium alloy catalysts 852(1999)507

## Centrifugation

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

## Chemiluminescence detection

Chemiluminescence detection of heme proteins separated by capillary isoelectric focusing 852(1999)597

# **Chiral recognition**

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a  $\beta$ -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

# Chiral selectivity

Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383

# Chiral stationary phases, LC

Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383

## Chymotrypsin

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

## Cisplatin

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

#### Collagen

Binding of lead to collagen type I and V and  $\alpha_2(I)$  CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

Separation of proteins and peptides by capillary electrophoresis in acid buffers containing high concentrations of surfactants 852(1999)325

## Columns

On the need to retain unregulated choice of high-performance liquid chromatography columns 852(1999)603

# Crude oils

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

#### Cyclodextrin

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a  $\beta$ -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

#### Decanophenone

pH effects on micelle-water partitioning determined by micellar electrokinetic chromatography 852(1999)559

## **Detection**, electrophoresis

Chemiluminescence detection of heme proteins separated by capillary isoelectric focusing 852(1999)597

## Detection, GC

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

# Detection, LC

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

## Diffusion

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

# Diuretics

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

## Elastase

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

#### **Electron-capture detection**

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

## **Enantiomer separation**

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a  $\beta$ -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

## **End-capping**

Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375

## Endorphin

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

#### **Environmental analysis**

Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

# Escherichia coli

High-yield purification of biosynthetic human growth hormone secreted in *Escherichia coli* periplasmic space 852(1999)441

## **Enzyme inhibitors**

Purification of two low-molecular-mass serine proteinase inhibitors from chicken liver 852(1999)207

#### Enzymes

Affinity purification of *Schistosoma japonicum* glutathione-Stransferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

Isolation and characterization of cyclic  $\alpha$ -(1 $\rightarrow$ 4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

## Epoxides

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

#### Ethoxylates

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

#### Factor VIII

Use of surface plasmon resonance for studies of protein-protein and protein-phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Use of high-resolution techniques for the characterization of clotting factor VIII 852(1999)175

#### Fibrinogenases

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

## Fluorinated silica gel columns

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475 616

## Fuels

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

#### Generalised H-point standard addition method

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

## Gluatathione-S-transferase

Affinity purification of *Schistosoma japonicum* glutathione-Stransferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

#### Glucagon

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

## **Glucagon-like peptide-1**

Proteolytic cleavage of glucagon-like peptide-1 by pancreatic  $\beta$  cells and by fetal calf serum analyzed by mass spectrometry 852(1999)285

#### Glucans

Isolation and characterization of cyclic  $\alpha$ -(1 $\rightarrow$ 4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

## Glut1

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

# Glycoproteins

Use of high-resolution techniques for the characterization of clotting factor VIII 852(1999)175

## **Gradient elution**

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments  $^{\star}$  852(1999)43

Prediction of protein retention at gradient elution conditions in ion-exchange chromatography 852(1999)351

#### **Growth factors**

Strategy for identifying circulating fragments of insulin-like growth factor binding proteins in a hemofiltrate peptide bank 852(1999)197

#### Growth hormones

High-yield purification of biosynthetic human growth hormone secreted in *Escherichia coli* periplasmic space 852(1999)441

### Hepatitis B virus

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

# Hepatitis C virus

Introduction of the gene amplification technique to decrease the risk of hepatitis C virus transmission by plasma products 852(1999)305

#### Histidine

Histidines in affinity tags and surface clusters for immobilized metal-ion affinity chromatography of trimeric tumor necrosis factor  $\alpha = 852(1999)117$ 

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

#### Hummel-Dreyer method

Binding of lead to collagen type I and V and  $\alpha_2(I)$  CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

# Hyaluronic acid

Molecular weight determination of hyaluronic acid by gel filtration chromatography coupled to matrix-assisted laser desorption ionization mass spectrometry 852(1999)573

# Hydroperoxides

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

# $17 \hbox{-Hydroxy-16-hydroxymethyl-3-methoxyestra-1,3,5(10)-triene}$

High-performance liquid chromatographic methods for monitoring of isomers of 17-hydroxy-16-hydroxymethyl-3methoxyestra-1,3,5(10)-triene 852(1999)433

## Immobilised metal affinity chromatography

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

# Immobilization

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

#### Immobilized biomembranes

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

### Immobilized chymotrypsin

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

High concentrations of sodium chloride facilitate the use of immobilized chymotrypsin for separating virgin forms of specific trypsin inhibitors 852(1999)227

## Immobilized metal affinity chromatography

Histidines in affinity tags and surface clusters for immobilized metal-ion affinity chromatography of trimeric tumor necrosis factor  $\alpha = 852(1999)117$ 

Affinity purification of *Schistosoma japonicum* glutathione-Stransferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

#### Immunoadsorbents

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

## Immunoaffinity chromatography

Sandwich-type immunoassay with an affinity column coupled to

anti-peptide antibodies 852(1999)97 Quantitative fast fractionation of a pool of polyclonal antibodies by immunoaffinity membrane chromatography 852(1999)129

#### Immunoassays

Strategy for identifying circulating fragments of insulin-like growth factor binding proteins in a hemofiltrate peptide bank 852(1999)197

#### Immunoglobulins

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent-detergent treatment 852(1999)83

Removal of detergent and solvent from solvent-detergenttreated immunoglobulins 852(1999)87

## Immunosorbents

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

# **Inorganic anions**

Analysis of anions in aqueous samples by ion chromatography and capillary electrophoresis. A comparative study of peak modeling and validation criteria 852(1999)487

# Instrumentation

Large-scale membrane adsorbers 852(1999)73

#### Insulin

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

Proteolytic cleavage of glucagon-like peptide-1 by pancreatic  $\beta$  cells and by fetal calf serum analyzed by mass spectrometry 852(1999)285

#### Intraparticle convective flow

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

## **Isoelectric focusing**

Chemiluminescence detection of heme proteins separated by capillary isoelectric focusing 852(1999)597

# **Isoelectric points**

Ion-exchange chromatography of proteins near the isoelectric points 852(1999)31

#### **Kinetic studies**

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum-rhodium alloy catalysts 852(1999)507

#### Lachesis stenophrys

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

# Lactoglobulins

Ion-exchange chromatography of proteins near the isoelectric points 852(1999)31

#### Lead

Binding of lead to collagen type I and V and  $\alpha_2(I)$  CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

## Lead-protein interactions

Binding of lead to collagen type I and V and  $\alpha_2$ (I) CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

## Lipophilicity

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

# Mathematical modeling

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

#### Membrane chromatography

Quantitative fast fractionation of a pool of polyclonal antibodies by immunoaffinity membrane chromatography 852(1999)129 618

# Membrane dryers

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

## Membrane extraction

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

# Membranes

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

## Large-scale membrane adsorbers 852(1999)73

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

#### Metalloproteinases

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

# Methyl bromide

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

# Methyl iodide

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

#### Micelles

pH effects on micelle-water partitioning determined by micellar electrokinetic chromatography 852(1999)559

## Migration inhibitory factors

Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor 852(1999)189

# Mobile phase composition

Characterization of unstable ion-exchange chromatographic separation of proteins 852(1999)37

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

#### Mobile phase modifier

Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383

# Monolith stationary phases

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

#### Monolithic columns

Separation of oligonucleotides on novel monolithic columns with ion-exchange functional surfaces 852(1999)297

# Nicotine

Optimization study for the reversed-phase ion-pair liquid chromatographic determination of nicotine in commercial tobacco products 852(1999)451

#### NoXon-S

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

## Nucleoside monophosphates

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

#### Nucleotides

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

# **Octylphenol ethoxylates**

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

#### Oligonucleotides

Separation of oligonucleotides on novel monolithic columns with ion-exchange functional surfaces 852(1999)297

# Organochlorine compounds

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

#### Oxidation

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum-rhodium alloy catalysts 852(1999)507

# Ozone scrubbing material

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

#### Partitioning

pH effects on micelle-water partitioning determined by micellar electrokinetic chromatography 852(1999)559

# Peak shape

Analysis of anions in aqueous samples by ion chromatography and capillary electrophoresis. A comparative study of peak modeling and validation criteria 852(1999)487

## Peptides

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

New polymer-based prepacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

Application of a peptide bank from porcine brain in isolation of regulatory peptides 852(1999)273

Proteolytic cleavage of glucagon-like peptide-1 by pancreatic  $\beta$  cells and by fetal calf serum analyzed by mass spectrometry 852(1999)285

Separation of proteins and peptides by capillary electrophoresis in acid buffers containing high concentrations of surfactants 852(1999)325

#### Pesticides

Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

# pH effects

pH effects on micelle-water partitioning determined by micellar electrokinetic chromatography 852(1999)559

## Pharmaceutical analysis

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

# Phenols

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

pH effects on micelle-water partitioning determined by micellar electrokinetic chromatography 852(1999)559

#### Phenothiazines

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

#### Phenylureas

Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465

## Phospholipids

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

## Platinum-rhodium catalysts

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum-rhodium alloy catalysts 852(1999)507

# **Polychlorinated biphenyls**

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

# Polynuclear aromatic hydrocarbons

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

# Polyphenylene sulfide

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

## Polysaccharides

Molecular weight determination of hyaluronic acid by gel filtration chromatography coupled to matrix-assisted laser desorption ionization mass spectrometry 852(1999)573

#### Pore connectivity

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

# Pore network model

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

# Pore size distribution

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

## **Porous particles**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

## **Protein synthesis**

Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor 852(1999)189

## **Protein-lead interactions**

Binding of lead to collagen type I and V and  $\alpha_2(I)$  CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

## Proteins

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

Ion-exchange chromatography of proteins near the isoelectric points 852(1999)31

Characterization of unstable ion-exchange chromatographic separation of proteins 852(1999)37

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Large-scale membrane adsorbers 852(1999)73

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105

Histidines in affinity tags and surface clusters for immobilized metal-ion affinity chromatography of trimeric tumor necrosis factor  $\alpha = 852(1999)117$ 

Affinity chromatography of human estrogen receptor- $\alpha$  expressed in *Saccharomyces cerevisiae*. Combination of heparinand 17 $\beta$ -estradiol-affinity chromatography 852(1999)161

Use of high-resolution techniques for the characterization of clotting factor VIII 852(1999)175

Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor 852(1999)189

Strategy for identifying circulating fragments of insulin-like growth factor binding proteins in a hemofiltrate peptide bank 852(1999)197

Purification of two low-molecular-mass serine proteinase inhibitors from chicken liver 852(1999)207

High concentrations of sodium chloride facilitate the use of immobilized chymotrypsin for separating virgin forms of specific trypsin inhibitors 852(1999)227

Binding of lead to collagen type I and V and  $\alpha_2(I)$  CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

Proteolytic cleavage of glucagon-like peptide-1 by pancreatic  $\beta$  cells and by fetal calf serum analyzed by mass spectrometry 852(1999)285

Experimental studies on affinity chromatography in an electric field 852(1999)319

Separation of proteins and peptides by capillary electrophoresis in acid buffers containing high concentrations of surfactants 852(1999)325

Prediction of protein retention at gradient elution conditions in ion-exchange chromatography 852(1999)351

High-yield purification of biosynthetic human growth hormone secreted in *Escherichia coli* periplasmic space 852(1999)441

Chemiluminescence detection of heme proteins separated by capillary isoelectric focusing 852(1999)597

## Pyridylmethylaminoacetate ligands

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105

#### **Retention behaviour**

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

# **Retention models**

Characterization of unstable ion-exchange chromatographic separation of proteins 852(1999)37

#### **Retention prediction**

Prediction of protein retention at gradient elution conditions in ion-exchange chromatography 852(1999)351

#### 620

Revision of a theoretical expression for gas–liquid chromatographic retention 852(1999)583

#### **Reversed-flow gas chromatography**

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum-rhodium alloy catalysts 852(1999)507

#### Reviews

Use of high-resolution techniques for the characterization of clotting factor VIII (Review) 852(1999)175

## **Rhodium-platinum catalysts**

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum-rhodium alloy catalysts 852(1999)507

## RNA

Introduction of the gene amplification technique to decrease the risk of hepatitis C virus transmission by plasma products 852(1999)305

#### Saccharomyces cerevisiae

Affinity chromatography of human estrogen receptor- $\alpha$  expressed in *Saccharomyces cerevisiae*. Combination of heparinand 17 $\beta$ -estradiol-affinity chromatography 852(1999)161

## Schistosoma japonicum

Affinity purification of *Schistosoma japonicum* glutathione-Stransferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

# Sewage sludge

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

### Sodium chloride

High concentrations of sodium chloride facilitate the use of immobilized chymotrypsin for separating virgin forms of specific trypsin inhibitors 852(1999)227

#### Solid-phase peptide synthesis

Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor 852(1999)189

## Solvent-detergent method

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent-detergent treatment 852(1999)83

Removal of detergent and solvent from solvent-detergenttreated immunoglobulins 852(1999)87

# Source 5RPC columns

New polymer-based prepacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

#### Stationary phases, LC

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

New polymer-based prepacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

Separation of oligonucleotides on novel monolithic columns with ion-exchange functional surfaces 852(1999)297

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

#### Steroid hormone receptors

Affinity chromatography of human estrogen receptor- $\alpha$  expressed in *Saccharomyces cerevisiae*. Combination of heparinand 17 $\beta$ -estradiol-affinity chromatography 852(1999)161

#### Steroids

High-performance liquid chromatographic methods for monitoring of isomers of 17-hydroxy-16-hydroxymethyl-3methoxyestra-1,3,5(10)-triene 852(1999)433

## Structure-retention relationships

Revision of a theoretical expression for gas–liquid chromatographic retention 852(1999)583

## Sulfur compounds

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

## Supercritical fluid extraction

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

#### Surface coverage

Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375

# Surfactants

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

Qualitative and quantitative analysis of new alkyl amide arginine surfactants by high-performance liquid chromatography and capillary electrophoresis 852(1999)499

# Surface plasmon resonance detection

Use of surface plasmon resonance for studies of protein-protein and protein-phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

## Thermodynamic parameters

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

Revision of a theoretical expression for gas–liquid chromatographic retention 852(1999)583

# Tobacco

Optimization study for the reversed-phase ion-pair liquid chromatographic determination of nicotine in commercial tobacco products 852(1999)451

# Triacylglycerols

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

# **Tributyl phosphate**

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent-detergent treatment 852(1999)83

Removal of detergent and solvent from solvent-detergenttreated immunoglobulins 852(1999)87

# Trimethylsilyl groups

Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375

#### Triolein

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

# Triton X-100

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent-detergent treatment 852(1999)83 Removal of detergent and solvent from solvent-detergenttreated immunoglobulins 852(1999)87

## Trypsin

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

## **Trypsin inhibitors**

High concentrations of sodium chloride facilitate the use of immobilized chymotrypsin for separating virgin forms of specific trypsin inhibitors 852(1999)227

# Tumor necrosis factor $\alpha$

Histidines in affinity tags and surface clusters for immobilized metal-ion affinity chromatography of trimeric tumor necrosis factor  $\alpha = 852(1999)117$ 

#### Validation

Analysis of anions in aqueous samples by ion chromatography and capillary electrophoresis. A comparative study of peak modeling and validation criteria 852(1999)487

## Vegetable oils

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

#### Venoms

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

#### Viruses

Introduction of the gene amplification technique to decrease the risk of hepatitis C virus transmission by plasma products 852(1999)305

# Von Willebrand factor

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Use of high-resolution techniques for the characterization of clotting factor VIII 852(1999)175

## Water analysis

Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465

## 622