



ELSEVIER

Journal of Chromatography A, 867 (2000) 293–297

JOURNAL OF
CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Subject Index Vol. 867

Affinity membranes

New affinity nylon membrane used for adsorption of γ -globulin 867(2000)161

Affinity adsorbents

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Albumin

Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

Amines

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Amino acids, Dns derivatives

Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271

Analysis time

Simultaneous optimization of the analysis time and the concentration detectability in open-tubular liquid chromatography 867(2000)23

Benzene

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Capillary electrophoresis, microchip

Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271

Carboxylic acids

Effect of stationary phase hydrophobicity and mobile phase composition on the separation of carboxylic acids in ion chromatography 867(2000)131

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Cereals

Erratum to: "Liquid chromatographic method for determining added folic acid in fortified cereal products" [J. Chromatogr. A, 826 (1998) 235–240] 867(2000)289

Charge coupled device detection

High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93

Chemiluminescence detection

Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271

Cigarette smoke

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

Concentration detectability

Simultaneous optimization of the analysis time and the concentration detectability in open-tubular liquid chromatography 867(2000)23

Concentration waves, thermo-induced

Ion exchange on resins with temperature-responsive selectivity. IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57

Copper

Ion exchange on resins with temperature-responsive selectivity. IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57

Detection, electrophoresis

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271

Detection, GC

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

Detection, LC

High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93

Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

Dichlorohydrin

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Dissociation constants

Dissociation constants of phenols in methanol–water mixtures 867(2000)45

Disulfide linkages

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

Dust

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

Efficiency

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Electrochemical detection

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Enantiomer separation

Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1

Environmental analysis

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Pesticide trace analysis using solid-phase extraction and gas chromatography with electron-capture and tandem mass spectrometric detection in water samples 867(2000)235

Enzymes

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Epichlorohydrin

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Equilibrium sorptive enrichment

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Ethoxyethanol

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Expanded bed chromatography

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Extraction methods

Determination of phenols using simultaneous steam distillation–extraction 867(2000)281

Flavours

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

Fluorescence detection

High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93

Fluoxamine

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Fluoxketone

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Folic acid

Erratum to: “Liquid chromatographic method for determining added folic acid in fortified cereal products” [J. Chromatogr. A, 826 (1998) 235–240] 867(2000)289

Food analysis

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

Erratum to: “Liquid chromatographic method for determining added folic acid in fortified cereal products” [J. Chromatogr. A, 826 (1998) 235–240] 867(2000)289

Fullerene

Synthesis and characteristics of [60]fullerene polysiloxane stationary phase for capillary gas chromatography 867(2000)177

Globulin

New affinity nylon membrane used for adsorption of γ -globulin 867(2000)161

Immobilized enzymes

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

Instrumentation

Visualization of solute migration in chromatographic columns. Quantitation of the concentration in a migrating zone 867(2000)71

Iodine

Visualization of solute migration in chromatographic columns. Quantitation of the concentration in a migrating zone 867(2000)71

Ketoacids

Comparative methodology in the determination of α -oxocarboxylates in aqueous solution. Ion chromatography versus gas chromatography after oximation, extraction and esterification 867(2000)143

Ketocarboxylates

Comparative methodology in the determination of α -oxocarboxylates in aqueous solution. Ion chromatography versus gas chromatography after oximation, extraction and esterification 867(2000)143

Lactate dehydrogenase

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Lactoglobulin

Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

Mathematical modelling

Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Melengestrol

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

Membranes

New affinity nylon membrane used for adsorption of γ -globulin 867(2000)161

Methanol–water mixtures

Dissociation constants of phenols in methanol–water mixtures 867(2000)45

Micellar electrokinetic chromatography

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Microchip capillary electrophoresis

Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271

Mobile phase composition

Effect of stationary phase hydrophobicity and mobile phase composition on the separation of carboxylic acids in ion chromatography 867(2000)131

Multiangle laser light scattering detection

Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

***Neu* differentiation factor**

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

Optimization

Simultaneous optimization of the analysis time and the concentration detectability in open-tubular liquid chromatography 867(2000)23

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Oxocarboxylates

Comparative methodology in the determination of α -oxocarboxylates in aqueous solution. Ion chromatography versus gas chromatography after oximation, extraction and esterification 867(2000)143

Peptide mapping

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

Peptides

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

Pesticides

Pesticide trace analysis using solid-phase extraction and gas chromatography with electron-capture and tandem mass spectrometric detection in water samples 867(2000)235

Petroleum

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

Pharmaceutical analysis

Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1

Phenols

Dissociation constants of phenols in methanol–water mixtures 867(2000)45

Determination of phenols using simultaneous steam distillation–extraction 867(2000)281

Phenoxyalkanoic acids

Determination of phenoxyalkanoic acids and other herbicides at the ng/ml level in water by solid-phase extraction with poly(divinylbenzene-co-*N*-vinylpyrrolidone) sorbent and high-performance liquid chromatography–diode-array detection 867(2000)169

Polycarboxylic acids

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Poly(divinylbenzene-co-*N*-vinylpyrrolidone)

Determination of phenoxyalkanoic acids and other herbicides at the ng/ml level in water by solid-phase extraction with poly(divinylbenzene-co-*N*-vinylpyrrolidone) sorbent and high-performance liquid chromatography–diode-array detection 867(2000)169

Polynuclear aromatic hydrocarbons

High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93

Polysiloxane

Synthesis and characteristics of [60]fullerene polysiloxane stationary phase for capillary gas chromatography 867(2000)177

Preferential solvation

Dissociation constants of phenols in methanol–water mixtures 867(2000)45

Proteins

Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

New affinity nylon membrane used for adsorption of γ -globulin 867(2000)161

Resolution

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Sample preparation

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Sensitivity

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

Soil

Determination of phenols using simultaneous steam distillation–extraction 867(2000)281

Solute migration

Visualization of solute migration in chromatographic columns. Quantitation of the concentration in a migrating zone 867(2000)71

Stacking

New affinity nylon membrane used for adsorption of γ -globulin 867(2000)161

Stationary phases, GC

Synthesis and characteristics of [60]fullerene polysiloxane stationary phase for capillary gas chromatography 867(2000)177

Stationary phases, LC

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Effect of stationary phase hydrophobicity and mobile phase composition on the separation of carboxylic acids in ion chromatography 867(2000)131

Steady-state recycling

Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1

Steam distillation–extraction

Determination of phenols using simultaneous steam distillation–extraction 867(2000)281

Steroids

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

Thermostripping

Ion exchange on resins with temperature-responsive selectivity. IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57

Toluene

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Trenbolone

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

UV detection

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

Visualization of solute migration

Visualization of solute migration in chromatographic columns.
Quantitation of the concentration in a migrating zone
867(2000)71

Water analysis

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Pesticide trace analysis using solid-phase extraction and gas chromatography with electron-capture and tandem mass spectrometric detection in water samples 867(2000)235

Xylenes

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Zinc

Ion exchange on resins with temperature-responsive selectivity.
IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57