



ELSEVIER

Journal of Chromatography A, 868 (2000) 321

JOURNAL OF  
CHROMATOGRAPHY A

[www.elsevier.com/locate/chroma](http://www.elsevier.com/locate/chroma)

## Erratum

### Erratum to: “Optimization of conductivity of low-molecular-mass anions in capillary zone electrophoresis” [J. Chromatogr. A, 850 (1999) 355–362]<sup>☆</sup>

Martin U. Katzmayer, Christian W. Klampfl\*, Wolfgang Buchberger

Department of Analytical Chemistry, Johannes-Kepler-University Linz, Altenbergersstrasse 69, A-4040 Linz, Austria

Received 2 November 1999

**Keywords:** Conductivity detection: Detection, electrophoresis; Background electrolyte composition; Organic acids; Inorganic anions

Page 361, Table 2 should read as follows:

Table 2  
Peak areas obtained for a 10 mg/l formate standard solution as a function of the electrolyte counter-ion mobility

Counter-ion	$\mu$ (counter-ion) ( $\text{cm}^2 \text{kV}^{-1} \text{s}^{-1}$ )	Peak area (arbitrary units)
K <sup>+</sup>	0.693	146 518
Na <sup>+</sup>	0.474	143 382
Li <sup>+</sup>	0.363	132 369
TEA <sup>+</sup>	0.297	123 032
Tris <sup>+</sup>	0.259	120 203
Arg <sup>+</sup>	0.229	117 865

\*PII of original article: S0021-9673(99)00168-5

\*Corresponding author. +43-732-2468-722; +61-732-2468-679.

E-mail address: christian.klampfl@jk.uni-linz.ac.at (C.W. Klampfl)