

Subject Index Vols. 869–871

Abamectin

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

AccQ.Fluor reagent

Effect of solvents on the fumonisins analysis by high-performance liquid chromatography with AccQ.Fluor as the derivatizing reagent 870(2000)469

Acetylcysteine

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Acetylsalicylic acid

Validation of the removal of acetylsalicylic acid. Recovery and determination of residues on various surfaces by high performance liquid chromatographic 870(2000)69

Acetylserine

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Activity coefficients

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Acyclovir

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Adsorbents

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Adsorption/partition model

Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C_8 and reversed-phase C_{18} types 869(2000)57

Adsorption isotherms

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Agarose–polymer matrices

Ultra-thin-layer agarose gel electrophoresis. II. Separation of DNA fragments on composite agarose–linear polymer matrices 871(2000)289

Air analysis

Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497

Air monitoring

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Albendazole

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Albumin

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis 871(2000)311

Alcohols

2-(4-Carboxyphenyl)-6-*N,N*-diethylaminobenzofuran: a useful reagent for the sensitive determination of alcohols by high-performance liquid chromatography with fluorimetric detection 869(2000)243

Aldehydes

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Alkaloids

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

Alkyl-diol silica columns

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Alkylphenols

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

Alumina support

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Amines

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Determination of heterocyclic aromatic amines in meat extracts by liquid chromatography–ion-trap atmospheric pressure chemical ionization mass spectrometry 869(2000)307

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Amino acids

Novel cinchona alkaloid carbamate C₉-dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255

Retention/quantitation properties of the *o*-phthaldialdehyde–3-mercaptopropionic acid and the *o*-phthaldialdehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

High-performance liquid chromatographic separation of the enantiomers of unusual α -amino acid analogues 871(2000)105

Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Amino acids, PTH derivatives

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255

Aminocyclohexyldichlorobenzamide

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

Aminophenols

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Aminophenylmethylbenzothiazole

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Aminothiols

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Angiotensins

Simultaneous separation of angiotensin and endothelin peptide families by high-performance liquid chromatography: application to the specific radioimmunoassay measurement of angiotensin II or endothelin-1 from tissue 870(2000)349

Annona cherimola

High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483

Anthocyanins

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Antibacterial agents

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Antibiotics

Separation of erythromycin and related substances on base-deactivated reversed-phase silica gel columns 870(2000)227

Liquid chromatography of polymyxin B sulphate 870(2000)237

Determination of quinolones in animal tissues and eggs by high-performance liquid chromatography with photodiode-array detection 871(2000)31

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Antibodies

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Antidepressants, tricyclic

Optimized stationary phases for the high-performance liquid chromatography–electrospray ionization mass spectrometric analysis of basic pharmaceuticals 869(2000)159

Antigens

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Antioxidants

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Apigenin

Determination of apigenin in rat plasma by high-performance liquid chromatography 870(2000)463

Apples

Retention/quantitation properties of the *o*-phthalaldehyde–3-mercaptopropionic acid and the *o*-phthalaldehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

Artemisia capillaris

Fractionation and analysis of *Artemisia capillaris* Thunb. by affinity chromatography with human serum albumin as stationary phase 870(2000)501

Artifact peaks

Investigations of artifact peaks in sensitive high-performance liquid chromatography methods 869(2000)85

Aryloxyphenoxypropanoic acid

Stability-indicating high-performance liquid chromatographic assay and analysis of decomposition products of 2-[4-[(7-chloro-2-quinoxalinyloxy]phenoxy]propanoic acid 870(2000)179

Ascorbic acid

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Avermectins

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Avidin

Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151

Azaspiracid

Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13

Azinphos-methyl

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Azo dyes

Determination of sulphonated dyes in water by ion-interaction high-performance liquid chromatography 871(2000)227

Band profiles

Investigations of artifact peaks in sensitive high-performance liquid chromatography methods 869(2000)85

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C_{18} (2) columns 869(2000)181

Barbiturates

Micellar liquid chromatography for prediction of drug transport 870(2000)1

Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39

Benzenediamines

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Benzenediols

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Benzenes

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Benzimidazoles

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Benzodiazepines

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

 β -Blockers

Optimized stationary phases for the high-performance liquid chromatography–electrospray ionization mass spectrometric analysis of basic pharmaceuticals 869(2000)159

Micellar liquid chromatography for prediction of drug transport 870(2000)1

Beverages

Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Biogenic amines

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75

Biosensors

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Bis(2-ethylhexyl)sodium sulfosuccinate

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Buffer composition

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Caffeine

Determination of theophylline and its metabolites in biological samples by liquid chromatography–mass spectrometry 870(2000)87

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Camptothecin

Measurement and pharmacokinetics of unbound 20(S)-camptothecin in rat blood and brain by microdialysis coupled to microbore liquid chromatography with fluorescence detection 870(2000)221

Capillarisin

Fractionation and analysis of *Artemisia capillaris* Thunb. by affinity chromatography with human serum albumin as stationary phase 870(2000)501

Capillary array scanner

Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299

Capillary columns

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Capillary electrochromatography

Separation of peptides by strong cation-exchange capillary electrochromatography 869(2000)385

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Capillary electrophoresis

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Carbamates

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Carbolines

Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Carboxyphenyldiethylaminobenzofuran

2-(4-Carboxyphenyl)-6-*N,N*-diethylaminobenzofuran: a useful reagent for the sensitive determination of alcohols by high-performance liquid chromatography with fluorimetric detection 869(2000)243

Casein

Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371

Catecholamines

Determination of catecholamines in pheochromocytoma cell (PC-12) culture medium by microdialysis–microbore liquid chromatography 870(2000)405

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Centrifugal partition chromatography

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

Cephalosporins

Characterization of cephalosporin transfer between aqueous and colloidal phases by micellar electrokinetic chromatography 871(2000)439

Chemometrics

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Chiral stationary phases, LC

Novel cinchona alkaloid carbamate C_9 -dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Chlorogenic acid

Determination of chlorogenic acid in rat blood by microdialysis coupled with microbore liquid chromatography and its application to pharmacokinetic studies 870(2000)443

6-Chloronicotinic acid

Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497

Chlorophylls

High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483

Chloroquinoxalinyloxyphenoxypropanoic acid

Stability-indicating high-performance liquid chromatographic assay and analysis of decomposition products of 2-{4-[(7-chloro-2-quinoxalinyloxy)phenoxy]propanoic acid 870(2000)179

Chlorotheophylline

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Chlorpheniramine

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Simultaneous determination of paracetamol and chlorpheniramine in human plasma by liquid chromatography–tandem mass spectrometry 870(2000)77

Cinchona carbamates

Novel cinchona alkaloid carbamate C_9 -dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Circular dichroism detection

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

Clay

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Clenbuterol

Enantiomeric separation of clenbuterol by transient isotachopheresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachopheresis 871(2000)173

Closantel

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Clotrimazole

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Coated columns

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Codeine

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Column geometry

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

Column stability

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

Column switching

Determination of theophylline and its metabolites in biological samples by liquid chromatography–mass spectrometry 870(2000)87

Determination of the *N*-methyl-D-aspartate receptor NR2B subunit blocker Ro 63-1908 in rat, cynomolgus monkey and human plasma by high-performance liquid chromatography with column switching and fluorescence detection 870(2000)189

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Column temperature

Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3

Control of column temperature in reversed-phase liquid chromatography 869(2000)211

Computer simulation

Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Coriolis force

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

Cosmetics

Determination of sunscreen agents in cosmetic products by supercritical fluid extraction and high-performance liquid chromatography 870(2000)199

Cotinine

Application of liquid separation techniques to the determination of the main urinary nicotine metabolites 870(2000)29

Coumarin-3-carbonyl chloride

Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473

Coupled columns

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Cyclodextrins

Analysis of *cis-trans* isomers and enantiomers of sertraline by cyclodextrin-modified micellar electrokinetic chromatography 871(2000)207

Cysteine

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Derivatization, electrophoresis

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

Derivatization, LC

2-(4-Carboxyphenyl)-6-*N,N*-diethylaminobenzofuran: a useful reagent for the sensitive determination of alcohols by high-performance liquid chromatography with fluorimetric detection 869(2000)243

Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39

Retention/quantitation properties of the *o*-phthaldialdehyde–3-mercaptopropionic acid and the *o*-phthaldialdehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473

New fluorimetric method of liquid chromatography for the determination of the neurotoxin domoic acid in seafood and marine phytoplankton 871(2000)1

Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75

Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115

Detection, electrophoresis

Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299

Detection, LC

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

Diacetoxynitrophenylpropylisothiocyanate

Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115

Dialysis

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Diazabicycloundecene

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

Dibenzofulvene

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

Dichloromethylaminocyclohexylbenzamide

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

Dihydrolipoic acid

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Dihydropyridines

High-performance liquid chromatography with amperometric detection applied to the screening of 1,4-dihydropyridines in human plasma 870(2000)105

Dimenhydrinate

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Dinitrophenylhydrazines

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Dinitrophenylhydrazones

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Diols

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Dioxolanes

Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127

Diphenhydramine

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Direct injection

Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151

Dissociation constants

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Determination of sulphonated dyes in water by ion-interaction high-performance liquid chromatography 871(2000)227

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Diuretics

Modelling retention in liquid chromatography as a function of solvent composition and pH of the mobile phase 869(2000)27

DNA

Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425

Ultra-thin-layer agarose gel electrophoresis. II. Separation of DNA fragments on composite agarose–linear polymer matrices 871(2000)289

Capillary electrophoresis of DNA damage after irradiation: apoptosis and necrosis 871(2000)321

DNA fragments

Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299

Domoic acid

New fluorimetric method of liquid chromatography for the determination of the neurotoxin domoic acid in seafood and marine phytoplankton 871(2000)1

Dopamine

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Drug adsorption

Micellar liquid chromatography for prediction of drug transport 870(2000)1

Dyes

Determination of sulphonated dyes in water by ion-interaction high-performance liquid chromatography 871(2000)227

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Econazole

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Efficiency

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Electrochromatography

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Some considerations concerning the composition of the mobile phase in capillary electrochromatography 869(2000)363

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Electroosmotic flow

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Electrophoretic mobility

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Enalapril

Study of the *cis*–*trans* isomerization of enalapril by reversed-phase liquid chromatography 871(2000)189

Enantiomer analysis

Enantioselective analysis of methadone in saliva by liquid chromatography–mass spectrometry 871(2000)163

Enantiomer separation

Novel cinchona alkaloid carbamate C_9 -dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

High-performance liquid chromatographic separation of the enantiomers of unusual α -amino acid analogues 871(2000)105

Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115

Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Enantiomeric separation of clenbuterol by transient isotachopheresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachopheresis 871(2000)173

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

Analysis of *cis*–*trans* isomers and enantiomers of sertraline by cyclodextrin-modified micellar electrokinetic chromatography 871(2000)207

Endothelins

Simultaneous separation of angiotensin and endothelin peptide families by high-performance liquid chromatography: application to the specific radioimmunoassay measurement of angiotensin II or endothelin-1 from tissue 870(2000)349

Environmental analysis

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Evaluation of surface- and ground-water pollution due to herbicides in agricultural areas of Zamora and Salamanca (Spain) 869(2000)471

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Enzyme inhibition

Study of the *cis*–*trans* isomerization of enalapril by reversed-phase liquid chromatography 871(2000)189

Ephedrine

Comparison of various reversed-phase columns for the simultaneous determination of ephedrine in urine by high-performance liquid chromatography 870(2000)23

Erythromycins

Separation of erythromycin and related substances on base-deactivated reversed-phase silica gel columns 870(2000)227

Ethylbenzene

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Ethylene oxide

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Evernia prunastri

Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425

Evernic acid

Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425

Experimental design

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Explosives

Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411

Extraction methods

Microwave-assisted extraction by fast sample preparation for the systematic analysis of additives in polyolefins by high-performance liquid chromatography 869(2000)285

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515

Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

Factorial design(s)

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Field-flow fractionation

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

Flavanoids

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Flavones

Isolation and characterization of methoxylated flavones in the flowers of *Primula veris* by liquid chromatography and mass spectrometry 870(2000)453

Flavonoids

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

Determination of apigenin in rat plasma by high-performance liquid chromatography 870(2000)463

Flow patterns

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

Fluorenylmethoxycarbonyl compounds

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

Fluorescence detection

Quantitative determination of sulfonamide residues in foods of animal origin by high-performance liquid chromatography with fluorescence detection 871(2000)37

4-Fluoro-7-nitro-2,1,3-benzoxadiazole

New fluorimetric method of liquid chromatography for the determination of the neurotoxin domoic acid in seafood and marine phytoplankton 871(2000)1

Food analysis

Determination of heterocyclic aromatic amines in meat extracts by liquid chromatography–ion-trap atmospheric pressure chemical ionization mass spectrometry 869(2000)307

Retention/quantitation properties of the *o*-phthalaldehyde–3-mercaptopropionic acid and the *o*-phthalaldehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483

Quantitation of anabolic hormones and their metabolites in bovine serum and urine by liquid chromatography–tandem mass spectrometry 870(2000)511

Anion-exchange high-performance liquid chromatography with conductivity detection for the analysis of phytic acid in food 871(2000)7

Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13

Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75

Determination of phenols in wines by liquid chromatography with photodiode array and fluorescence detection 871(2000)85

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Forensic analysis

Profiling of impurities in illicit methamphetamine by high-performance liquid chromatography and capillary electrochromatography 870(2000)53

Formic acid

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Fruits

Retention/quantitation properties of the *o*-phthaldialdehyde–3-mercaptopropionic acid and the *o*-phthaldialdehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Fucans

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

Fullerene adsorbents

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Fullerenes

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Fumonisin

Effect of solvents on the fumonisins analysis by high-performance liquid chromatography with AccQ.Fluor as the derivatizing reagent 870(2000)469

Furans

Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49

Gamma radiation

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

Ganciclovir

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Gerbils

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Glutathione

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Glycoalkaloids

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

Gradient elution

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Separation of polyesters by gradient reversed-phase high-performance liquid chromatography on a 1.5 μm non-porous column 871(2000)243

Gradient polymer elution chromatography

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Heterocyclic aromatic compounds

Determination of heterocyclic aromatic amines in meat extracts by liquid chromatography–ion-trap atmospheric pressure chemical ionization mass spectrometry 869(2000)307

Hexafluoroisopropanol

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Homocysteine

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Homovanillic acid

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Honey

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

Hormones

Separation and characterization of multicomponent peptide mixtures by liquid chromatography–electrospray ionization mass spectrometry. Application to crude products of the synthesis of leuprolide 870(2000)315

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Quantitation of anabolic hormones and their metabolites in bovine serum and urine by liquid chromatography–tandem mass spectrometry 870(2000)511

Hydrocarbons

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Hydrocortisone

Direct identification and quantitation of prednisone in the presence of overlapping hydrocortisone by liquid chromatography with electrospray and atmospheric-pressure chemical-ionisation mass spectrometry 870(2000)135

Hydroxycotinine

Application of liquid separation techniques to the determination of the main urinary nicotine metabolites 870(2000)29

Hydroxyindoleacetic acid

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Hydroxyquinolines

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Ibuprofen

Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143

Ice crystals

Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391

Imidacloprid

Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497

Imidazolic compounds

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Injection techniques

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

Injector dispersion

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

Injector performance

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

Inorganic anions

Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391

Inorganic cations

Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391

Inorganic ions

Coupling of chromatographic ion-exchange reaction with change-partner reaction. Presumed mechanism on the ion–ion interaction of inorganic ions in Sephadex G-15 columns 869(2000)91

Instrumentation

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

Ion–ion interactions

Coupling of chromatographic ion-exchange reaction with change-partner reaction. Presumed mechanism on the ion–ion interaction of inorganic ions in Sephadex G-15 columns 869(2000)91

Iridium

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

ISIS 2922

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis 871(2000)311

Isotachophoresis–capillary zone electrophoresis

Enantiomeric separation of clenbuterol by transient isotachophoresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachophoresis 871(2000)173

Kerosene

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

Ketoconazole

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Ketones

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Ketoprofen

Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Lactate

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Lactones

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Least-squares analysis

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Leuprolide

Separation and characterization of multicomponent peptide mixtures by liquid chromatography–electrospray ionization mass spectrometry. Application to crude products of the synthesis of leuprolide 870(2000)315

Libraries, MS

Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411

Ligand fishing

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Linear solvation energy relationships

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Linuron

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Lipids

Analysis of pesticide residues in matrices with high lipid contents by membrane separation coupled on-line to a high-performance liquid chromatography system 869(2000)427

Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363

Lipoic acid

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Lipophilicity

Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49

Lorazepam

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

Luna C₁₈ (2)

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C₁₈ (2) columns 869(2000)181

Marine sediments

Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515

Mathematical modelling

Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C₈ and reversed-phase C₁₈ types 869(2000)57

Matrix solid-phase dispersion

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Membranes

Analysis of pesticide residues in matrices with high lipid contents by membrane separation coupled on-line to a high-performance liquid chromatography system 869(2000)427

Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151

Metal chelates

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Methadone

Enantioselective analysis of methadone in saliva by liquid chromatography–mass spectrometry 871(2000)163

Methamphetamine

Profiling of impurities in illicit methamphetamine by high-performance liquid chromatography and capillary electrochromatography 870(2000)53

Method development

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Methyl aspartate receptor blockers

Determination of the *N*-methyl-D-aspartate receptor NR2B subunit blocker Ro 63-1908 in rat, cynomolgus monkey and human plasma by high-performance liquid chromatography with column switching and fluorescence detection 870(2000)189

Micellar liquid chromatography

Micellar liquid chromatography for prediction of drug transport 870(2000)1

Microdialysis

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Determination of chlorogenic acid in rat blood by microdialysis coupled with microbore liquid chromatography and its application to pharmacokinetic studies 870(2000)443

Microporous membrane liquid–liquid extraction

Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Microwave-assisted extraction

Microwave-assisted extraction by fast sample preparation for the systematic analysis of additives in polyolefins by high-performance liquid chromatography 869(2000)285

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515

Migration time correction

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

Milk

Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371

Mobile phase composition

Modelling retention in liquid chromatography as a function of solvent composition and pH of the mobile phase 869(2000)27

Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Some considerations concerning the composition of the mobile phase in capillary electrochromatography 869(2000)363

Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127

Separation of isomeric naphthalenesulphonic acids by micro high-performance liquid chromatography with mobile phases containing cyclodextrin 871(2000)139

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Mobile phase optimisation

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Mobile phase strength

Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3

Monobromobimane

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Multivariate statistical methods

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Mycotoxins

Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473

Nanoparticles

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis 871(2000)311

Naphthalenes

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Naphthalenesulphonic acids

Separation of isomeric naphthalenesulphonic acids by micro high-performance liquid chromatography with mobile phases containing cyclodextrin 871(2000)139

Naphthoquinonesulfonate

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Nicotine

Application of liquid separation techniques to the determination of the main urinary nicotine metabolites 870(2000)29

Nifedipine

Quantitation of nifedipine in human plasma by on-line solid-phase extraction and high-performance liquid chromatography 870(2000)115

Nitrendipine

Quantitation of nifedipine in human plasma by on-line solid-phase extraction and high-performance liquid chromatography 870(2000)115

Non-aqueous capillary electrophoresis

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Nonsteroidal anti-inflammatory drugs

Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Nucleoside analogues

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Nylon

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Oligofucans

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

Oligonucleotides

Sheath liquid effects in capillary high-performance liquid chromatography–electrospray mass spectrometry of oligonucleotides 870(2000)413

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis 871(2000)311

Oligosaccharides

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

Open-tubular electrochromatography

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Optimization

Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75

Enantiomeric separation of clenbuterol by transient isotachopheresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachopheresis 871(2000)173

Optimization of the separation of phenolic compounds by micellar electrokinetic capillary chromatography 871(2000)427

Organic acids

Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391

Organometallic compounds

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Organophosphorus compounds

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Organosilane compounds

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Palladium

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Paracetamol

Simultaneous determination of paracetamol and chlorpheniramine in human plasma by liquid chromatography–tandem mass spectrometry 870(2000)77

Particle size distribution

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

Partition/adsorption model

Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C_{18} and reversed-phase C_{18} types 869(2000)57

Partition coefficients

Characterization of cephalosporin transfer between aqueous and colloidal phases by micellar electrokinetic chromatography 871(2000)439

Peptides

Separation of peptides by strong cation-exchange capillary electrochromatography 869(2000)385

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption/ionization time-of-flight mass spectrometry 870(2000)255

Signature-peptide approach to detecting proteins in complex mixtures 870(2000)295

Separation and characterization of multicomponent peptide mixtures by liquid chromatography–electrospray ionization mass spectrometry. Application to crude products of the synthesis of leuprolide 870(2000)315

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Simultaneous separation of angiotensin and endothelin peptide families by high-performance liquid chromatography: application to the specific radioimmunoassay measurement of angiotensin II or endothelin-1 from tissue 870(2000)349

Perfluorocarboxylic acids

Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245

Pesticides

Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411

Analysis of pesticide residues in matrices with high lipid contents by membrane separation coupled on-line to a high-performance liquid chromatography system 869(2000)427

On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463

Evaluation of surface- and ground-water pollution due to herbicides in agricultural areas of Zamora and Salamanca (Spain) 869(2000)471

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

Pharmaceutical analysis

Micellar liquid chromatography for prediction of drug transport 870(2000)1

Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Validation of the removal of acetylsalicylic acid. Recovery and determination of residues on various surfaces by high-performance liquid chromatographic 870(2000)69

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143

Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Stability-indicating high-performance liquid chromatographic assay and analysis of decomposition products of 2-[4-[(7-chloro-2-quinoxalinyloxy)phenoxy]propanoic acid 870(2000)179

Determination of eight water- and fat-soluble vitamins in multi-vitamin pharmaceutical formulations by high-performance liquid chromatography 870(2000)207

Determination of vitamin B₁₂ in multivitamin tablets by multimode high-performance liquid chromatography 870(2000)217

Separation of erythromycin and related substances on base-deactivated reversed-phase silica gel columns 870(2000)227

Liquid chromatography of polymyxin B sulphate 870(2000)237

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

pH effects

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by γ -radiation on titanium-grafted silica 869(2000)137

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Pheochromocytoma cells

Determination of catecholamines in pheochromocytoma cell (PC-12) culture medium by microdialysis–microbore liquid chromatography 870(2000)405

Phenolic compounds

Determination of chlorogenic acid in rat blood by microdialysis coupled with microbore liquid chromatography and its application to pharmacokinetic studies 870(2000)443

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Optimization of the separation of phenolic compounds by micellar electrokinetic capillary chromatography 871(2000)427

Phenols

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Determination of phenols in wines by liquid chromatography with photodiode array and fluorescence detection 871(2000)85

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Phenylureas

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

***o*-Phthaldialdehyde**

Effect of solvents on the fumonisins analysis by high-performance liquid chromatography with AccQ.Fluor as the derivatizing reagent 870(2000)469

Photochemical derivatization

Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39

Phytic acid

Anion-exchange high-performance liquid chromatography with conductivity detection for the analysis of phytic acid in food 871(2000)7

Plant material(s)

Isolation and characterization of methoxylated flavones in the flowers of *Primula veris* by liquid chromatography and mass spectrometry 870(2000)453

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Platinum

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

PNU-83892

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

PNU-83894

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

Polyamides

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Polyamines

Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425

Polydimethylsiloxane

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Polyesters

Separation of polyesters by gradient reversed-phase high-performance liquid chromatography on a 1.5 μm non-porous column 871(2000)243

Polyglycols

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Polymer–agarose matrices

Ultra-thin-layer agarose gel electrophoresis. II. Separation of DNA fragments on composite agarose–linear polymer matrices 871(2000)289

Polymers

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Poly(methyloctylsiloxane)

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

Poly(methyloctylsiloxane) stationary phases

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by γ -radiation on titanium-grafted silica 869(2000)137

Polymyxins

Liquid chromatography of polymyxin B sulphate 870(2000)237

Polynuclear aromatic hydrocarbons

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Polyolefins

Microwave-assisted extraction by fast sample preparation for the systematic analysis of additives in polyolefins by high-performance liquid chromatography 869(2000)285

Polyphenols

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

Polysaccharides

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

Poly(sodium 11-acrylamidoundecanoate)

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Polystyrene resins

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

Porous graphitic carbon

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245

Positional isomer separation

Study of the *cis-trans* isomerization of enalapril by reversed-phase liquid chromatography 871(2000)189

Prednisone

Direct identification and quantitation of prednisone in the presence of overlapping hydrocortisone by liquid chromatography with electrospray and atmospheric-pressure chemical-ionisation mass spectrometry 870(2000)135

Primulae veris

Isolation and characterization of methoxylated flavones in the flowers of *Primula veris* by liquid chromatography and mass spectrometry 870(2000)453

Principal component analysis

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Profens

Novel cinchona alkaloid carbamate C₉-dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Propylene oxide

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Protein sequencing

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255

Proteins

Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255

Signature-peptide approach to detecting proteins in complex mixtures 870(2000)295

Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363

Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371

Pseudoephedrine

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Pseudo-stationary phases

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Pulsed diode laser

Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299

Pyrimethamine

Simultaneous determination of sulfaquinoxaline, sulfamethazine and pyrimethamine by liquid chromatography 870(2000)169

Pyruvate

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Quaternary ammonium compounds

On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441

Quinine carbamates

Novel cinchona alkaloid carbamate C₉-dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Quinolones

Determination of quinolones in animal tissues and eggs by high-performance liquid chromatography with photodiode-array detection 871(2000)31

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Random walk model

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

Residue determination

Quantitative determination of sulfonamide residues in foods of animal origin by high-performance liquid chromatography with fluorescence detection 871(2000)37

Resolution prediction

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Retention data

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C₁₈ (2) columns 869(2000)181

Retention mechanism

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Retention modelling

Modelling retention in liquid chromatography as a function of solvent composition and pH of the mobile phase 869(2000)27

Retention parameters

Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49

Rhodium

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Robustness testing

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Ropivacaine

Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151

Rosemary

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Rotenone

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

Ro 63-1908

Determination of the *N*-methyl-D-aspartate receptor NR2B subunit blocker Ro 63-1908 in rat, cynomolgus monkey and human plasma by high-performance liquid chromatography with column switching and fluorescence detection 870(2000)189

Sample preparation

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Sample volume

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

Scoparone

Fractionation and analysis of *Artemisia capillaris* Thunb. by affinity chromatography with human serum albumin as stationary phase 870(2000)501

Selectivity

Development of a new cyano-bonded column for high-performance liquid chromatography 869(2000)143

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Separation mechanism

Separation of peptides by strong cation-exchange capillary electrochromatography 869(2000)385

Sephadex G-15

Coupling of chromatographic ion-exchange reaction with change-partner reaction. Presumed mechanism on the ion–ion interaction of inorganic ions in Sephadex G-15 columns 869(2000)91

Serotonin

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Sertraline

Analysis of *cis*–*trans* isomers and enantiomers of sertraline by cyclodextrin-modified micellar electrokinetic chromatography 871(2000)207

Sheath liquid effects

Sheath liquid effects in capillary high-performance liquid chromatography–electrospray mass spectrometry of oligonucleotides 870(2000)413

Shellfish poisoning

Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13

Signature-peptide approach

Signature-peptide approach to detecting proteins in complex mixtures 870(2000)295

Silica, titanium grafted

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by γ -radiation on titanium-grafted silica 869(2000)137

Soil

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Solanum brevidens

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

Solid-phase extraction

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

Solid-phase microextraction

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Split-flow thin cell fractionation

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

Stability studies

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Starch

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

Stationary phases, LC

Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C₈ and reversed-phase C₁₈ types 869(2000)57

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by γ -radiation on titanium-grafted silica 869(2000)137

Development of a new cyano-bonded column for high-performance liquid chromatography 869(2000)143

Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151

Optimized stationary phases for the high-performance liquid chromatography–electrospray ionization mass spectrometric analysis of basic pharmaceuticals 869(2000)159

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C₁₈ (2) columns 869(2000)181

Control of column temperature in reversed-phase liquid chromatography 869(2000)211

Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245

Separation of polyesters by gradient reversed-phase high-performance liquid chromatography on a 1.5 µm non-porous column 871(2000)243

Strong cation exchange

Separation of peptides by strong cation-exchange capillary electrochromatography 869(2000)385

Structure–retention relationships

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Sulfamethazine

Simultaneous determination of sulfaquinoxaline, sulfamethazine and pyrimethamine by liquid chromatography 870(2000)169

Sulfaquinoxaline

Simultaneous determination of sulfaquinoxaline, sulfamethazine and pyrimethamine by liquid chromatography 870(2000)169

Sulfonamide(s)

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Quantitative determination of sulfonamide residues in foods of animal origin by high-performance liquid chromatography with fluorescence detection 871(2000)37

Sunscreen agents

Determination of sunscreen agents in cosmetic products by supercritical fluid extraction and high-performance liquid chromatography 870(2000)199

Surfactant proteins

Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363

Surfactants

Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Terpenes

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Tetrahydrocarboline

Analysis of tetrahydro-β-carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Tetrahydrocarbolinecarboxylic acids

Analysis of tetrahydro-β-carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Theophyllines

Determination of theophylline and its metabolites in biological samples by liquid chromatography–mass spectrometry 870(2000)87

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Thermodynamic parameters

Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127

Thiamine

Thiamine analysis in biological media by capillary zone electrophoresis with a high-sensitivity cell 871(2000)351

Thiols

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Titanium-grafted silica

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by γ -radiation on titanium-grafted silica 869(2000)137

Toxins

Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13

Triazines

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463

Trichothecenes

Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473

Troubleshooting

Investigations of artifact peaks in sensitive high-performance liquid chromatography methods 869(2000)85

Tryptophan

Screening for defects in tryptophan metabolism 870(2000)289

Tylosin

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Vancomycin

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Vanillylmandelic acid

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Validation

Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Validation of the removal of acetylsalicylic acid. Recovery and determination of residues on various surfaces by high performance liquid chromatographic 870(2000)69

Simultaneous determination of paracetamol and chlorpheniramine in human plasma by liquid chromatography–tandem mass spectrometry 870(2000)77

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Quantitation of nifedipine in human plasma by on-line solid-phase extraction and high-performance liquid chromatography 870(2000)115

Vegetables

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Vesicles

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Vitamins

Determination of eight water- and fat-soluble vitamins in multi-vitamin pharmaceutical formulations by high-performance liquid chromatography 870(2000)207

Determination of vitamin B₁₂ in multivitamin tablets by multimode high-performance liquid chromatography 870(2000)217

Volatile organic compounds

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Water analysis

On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463

Evaluation of surface- and ground-water pollution due to herbicides in agricultural areas of Zamora and Salamanca (Spain) 869(2000)471

Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Wheat

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

Whey proteins

Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371

Wine

Optimization of the separation of phenolic compounds by micellar electrokinetic capillary chromatography 871(2000)427

Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75

Determination of phenols in wines by liquid chromatography with photodiode array and fluorescence detection 871(2000)85

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Wood

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Zirconized silica

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129