



ELSEVIER

Journal of Chromatography A, 877 (2000) 243–244

JOURNAL OF  
CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

## Author Index Vol. 877

- Bashir, W., see Paull, B. 877(2000)123
- Björklund, E., see Järemo, M. 877(2000)167
- Brinkman, U.A.Th., see Catalina, M.I. 877(2000)153
- Catalina, M.I., Dallüge, J., Vreuls, R.J.J. and Brinkman, U.A.Th.  
Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153
- Cavazzini, A., see Quiñones, I. 877(2000)1
- Chang, S.Y., see Chiang, M.-T. 877(2000)233
- Cherrak, D.E., see Khattabi, S. 877(2000)95
- Cherrak, D.E., Khattabi, S. and Guiochon, G.  
Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109
- Chiang, M.-T., Chang, S.Y. and Whang, C.-W.  
Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233
- Chowdhury, M.A.J., Ihara, H., Sagawa, T. and Hirayama, C.  
Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71
- Colmsjö, A., see Ericsson, M. 877(2000)141
- Conte, E.D., see Crick, E.W. 877(2000)87
- Crick, E.W. and Conte, E.D.  
Alkyltrimethylammonium surfactant-mediated extractions: characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87
- Dallüge, J., see Catalina, M.I. 877(2000)153
- Dorsey, J.G., see Doyle, C.A. 877(2000)25
- Dorsey, J.G., see Doyle, C.A. 877(2000)41
- Doyle, C.A., Vickers, T.J., Mann, C.K. and Dorsey, J.G.  
Characterization of C<sub>18</sub>-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of mobile phase composition 877(2000)25
- Doyle, C.A., Vickers, T.J., Mann, C.K. and Dorsey, J.G.  
Characterization of C<sub>18</sub>-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41
- Ericsson, M. and Colmsjö, A.  
Dynamic microwave-assisted extraction 877(2000)141
- Farnan, D., see Hallgren, E. 877(2000)13
- Fischer, J., see Khattabi, S. 877(2000)95
- Flanigan, V., see Gehrke, M. 877(2000)133
- Gehrke, M., Kapila, S. and Flanigan, V.  
Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133
- Grimes, B.A., see Liapis, A.I. 877(2000)181
- Guiochon, G., see Cherrak, D.E. 877(2000)109
- Guiochon, G., see Khattabi, S. 877(2000)95
- Guiochon, G., see Quiñones, I. 877(2000)1
- Hallgren, E., Kálmán, F., Farnan, D., Horváth, C. and Ståhlberg, J.  
Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13
- Hiratani, K., see Shinbo, T. 877(2000)61
- Hirayama, C., see Chowdhury, M.A.J. 877(2000)71
- Horváth, C., see Hallgren, E. 877(2000)13
- Ide, H., see Suzuki, T. 877(2000)225
- Ihara, H., see Chowdhury, M.A.J. 877(2000)71
- Iwatsubo, T., see Shinbo, T. 877(2000)61
- Jandera, P., see Khattabi, S. 877(2000)95
- Järemo, M., Björklund, E., Nilsson, N., Karlsson, L. and Mathiasson, L.  
Utilization of fat retainers in supercritical fluid extraction for the selective extraction of polychlorinated biphenyls from a model fat sample 877(2000)167
- Kálmán, F., see Hallgren, E. 877(2000)13
- Kanamori, T., see Shinbo, T. 877(2000)61
- Kanaori, K., see Suzuki, T. 877(2000)225
- Kapila, S., see Gehrke, M. 877(2000)133
- Karlsson, L., see Järemo, M. 877(2000)167
- Khattabi, S., Cherrak, D.E., Fischer, J., Jandera, P. and Guiochon, G.  
Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95
- Khattabi, S., see Cherrak, D.E. 877(2000)109
- Kim, H.-J., see Kim, I.-J. 877(2000)217
- Kim, I.-J., Park, S.-J. and Kim, H.-J.  
Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using  $\gamma$ -cyclodextrin 877(2000)217
- Liapis, A.I. and Grimes, B.A.  
Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with

- charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181
- Makino, K., see Suzuki, T. 877(2000)225
- Mann, C.K., see Doyle, C.A. 877(2000)25
- Mann, C.K., see Doyle, C.A. 877(2000)41
- Mathiasson, L., see Järemo, M. 877(2000)167
- Morii, T., see Suzuki, T. 877(2000)225
- Nagawa, Y., see Shinbo, T. 877(2000)61
- Nilsson, N., see Järemo, M. 877(2000)167
- Park, S.-J., see Kim, I.-J. 877(2000)217
- Paull, B., Twohill, E. and Bashir, W.  
Determination of trace cadmium in environmental water samples using ion-interaction reversed-phase liquid chromatography with fluorescence detection 877(2000)123
- Quiñones, I., Cavazzini, A. and Guiochon, G.  
Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1
- Sagawa, T., see Chowdhury, M.A.J. 877(2000)71
- Shimabukuro, Y., see Shinbo, T. 877(2000)61
- Shinbo, T., Shimabukuro, Y., Kanamori, T., Iwatsubo, T., Nagawa, Y. and Hiratani, K.  
Separation of aromatic isomers on cyclophane-bonded stationary phases 877(2000)61
- Ståhlberg, J., see Hallgren, E. 877(2000)13
- Suzuki, T., Yamada, M., Ide, H., Kanaori, K., Tajima, K., Morii, T. and Makino, K.  
Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225
- Tajima, K., see Suzuki, T. 877(2000)225
- Twohill, E., see Paull, B. 877(2000)123
- Vickers, T.J., see Doyle, C.A. 877(2000)25
- Vickers, T.J., see Doyle, C.A. 877(2000)41
- Vreuls, R.J.J., see Catalina, M.I. 877(2000)153
- Whang, C.-W., see Chiang, M.-T. 877(2000)233
- Yamada, M., see Suzuki, T. 877(2000)225