



ELSEVIER

Journal of Chromatography A, 877 (2000) 245–248

JOURNAL OF
CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Subject Index Vol. 877

Adsorption isotherms

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Alkyltrimethylammonium surfactants

Alkyltrimethylammonium surfactant-mediated extractions: characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Amino acids

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Aminochlorophenylbutyric acid

Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233

Aminosulfonylfluorobenzoxadiazole

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Antibiotics

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Aromatic isomers

Separation of aromatic isomers on cyclophane-bonded stationary phases 877(2000)61

Baclofen

Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233

Band profiles

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Buspirone

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Cadmium

Determination of trace cadmium in environmental water samples using ion-interaction reversed-phase liquid chromatography with fluorescence detection 877(2000)123

Capillary columns

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Cetrimide

Alkyltrimethylammonium surfactant-mediated extractions: characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Charge distribution

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Charged porous particles

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Chiral stationary phases, LC

Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Chlorobenzenes

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Chlorophenoxy acids

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

3-Chloro-1-phenyl-1-propanol

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Comb-shaped polymer

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

Crystalline phases

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

Cyclodextrins

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Cyclophane

Separation of aromatic isomers on cyclophane-bonded stationary phases 877(2000)61

Derivatization, electrophoresis

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Derivatization, GC

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Detection, GC

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Diltiazem

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Dimethyl sulphate

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Doxepin

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Electrochromatography

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Electron-capture detection

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Electroosmotic flow

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Enantiomer separation

Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Environmental analysis

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Enzymes

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Explosives

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Extraction methods

Alkyltrimethylammonium surfactant-mediated extractions: characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Dynamic microwave-assisted extraction 877(2000)141

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid-liquid extraction combined with large-volume on-column injection and gas chromatography-mass spectrometry 877(2000)153

Factorial design

Dynamic microwave-assisted extraction 877(2000)141

Fat retainers

Utilization of fat retainers in supercritical fluid extraction for the selective extraction of polychlorinated biphenyls from a model fat sample 877(2000)167

Halogenated compounds

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Homocysteine

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

8-Hydroxyquinoline 5-sulphonic acid

Determination of trace cadmium in environmental water samples using ion-interaction reversed-phase liquid chromatography with fluorescence detection 877(2000)123

Instrumentation

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Dynamic microwave-assisted extraction 877(2000)141

Intraparticle convective flow

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Intraparticle electroosmotic flow

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Intraparticle Peclet number

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Large-volume injection

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid-liquid extraction combined with large-volume on-column injection and gas chromatography-mass spectrometry 877(2000)153

Mass transfer

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Mathematical modeling

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Microwave-assisted extraction

Dynamic microwave-assisted extraction 877(2000)141

Mobile phase composition

Characterization of C_{18} -bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of mobile phase composition 877(2000)25

Characterization of C_{18} -bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41

Naphthalenedicarboxaldehyde

Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233

Net charge

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Nitroaromatic compounds

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Nitrobenzenes

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Nitrotoluenes

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Non-crystalline phases

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

Nuclease

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Nucleobase

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Oxanine

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Peak broadening

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Pesticides

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Pharmaceutical analysis

Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233

Phase transition

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

Phenoxy acids

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

1-Phenyl-1-propanol

Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95

Polychlorinated biphenyls

Utilization of fat retainers in supercritical fluid extraction for the selective extraction of polychlorinated biphenyls from a model fat sample 877(2000)167

Polynuclear aromatic hydrocarbons

Dynamic microwave-assisted extraction 877(2000)141

Protein retention

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Raman spectroscopy

Characterization of C₁₈-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of mobile phase composition 877(2000)25

Characterization of C₁₈-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41

Retention models

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Ring opening–closure equilibrium

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Shape selectivity

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

Staphylococcal nuclease

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Stationary phases, LC

Characterization of C₁₈-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of mobile phase composition 877(2000)25

Characterization of C₁₈-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41

Separation of aromatic isomers on cyclophane-bonded stationary phases 877(2000)61

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

Surfactant-mediated extraction

Alkyltrimethylammonium surfactant-mediated extractions: characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Surfactants

Alkyltrimethylammonium surfactant-mediated extractions: characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Temperature effects

Characterization of C₁₈-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41

Water analysis

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153