

Journal of Chromatography A, 888 (2000) 345-350

JOURNAL OF CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Subject Index Vol. 888

Aldicarb

Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography-atmospheric pressure chemical ionization mass spectrometry 888(2000)113

Allergens

Identification of atranorin and related potential allergens in oakmoss absolute by high-performance liquid chromatography—tandem mass spectrometry using negative ion atmospheric pressure chemical ionization 888(2000)103

Alumina

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Amino acids

Comparison of the thin-layer chromatographic properties of sulfur-containing amino acids and their aminophosphonic analogues 888(2000)335

Aminoglutethimide

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

Aminophosphonic acids

Comparison of the thin-layer chromatographic properties of sulfur-containing amino acids and their aminophosphonic analogues 888(2000)335

Anilines

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Anion exchange

Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267

Atmospheric particulate matter

Analysis of heavy metals in atmospheric particulate by ion chromatography 888(2000)145

Atranorin

Identification of atranorin and related potential allergens in oakmoss absolute by high-performance liquid chromatography—tandem mass spectrometry using negative ion atmospheric pressure chemical ionization 888(2000)103

Beer

Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321

Bile acids

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Buffer composition

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309

Chemometrics

Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197

Chiral selectors

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

Chlorophenols

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

2-Chlorovinylarsonous acid

Simultaneous extraction and derivatization of 2chlorovinylarsonous acid from soils using supercritical and pressurized fluids 888(2000)327

0021-9673/00/\$ – see front matter $\,$ © 2000 Elsevier Science B.V. All right reserved

PII: S0021-9673(00)00650-6

Counter-current chromatography

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N*,*N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Cross-linking

Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63

Cyclodextrins

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

Simultaneous determination of eleven ingredients in ophthalmic solutions by cyclodextrin-modified micellar electrokinetic chromatography with tetrabutylammonium salt 888(2000)299

Deprotonation

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Derivatization, SFE

Simultaneous extraction and derivatization of 2-chlorovinylarsonous acid from soils using supercritical and pressurized fluids 888(2000)327

Detection, electrophoresis

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309

Dibutylamine acetate

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Dihexyldiethylcarbamoyl methylenephosphonate stationary phases

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N*,*N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Distribution constants

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N*,*N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Ebrotidine

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Electrochromatography

Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267

Electrophoretic mobility

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Enantiomer separation

Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63

On-line monitoring of enantiomer concentration in chiral simulated moving bed chromatography 888(2000)73

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

Enzyme inhibitors

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Fatty acids

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Feasibility

Affinity screening by packed capillary high-performance liquid chromatography using molecular imprinted sorbents. I. Demonstration of feasibility 888(2000)23

Fibers

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

Flavonoids

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Food analysis

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography-atmospheric pressure chemical ionization mass spectrometry 888(2000)113

High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321

Fruits

Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography-atmospheric pressure chemical ionization mass spectrometry 888(2000)113

Gas chromatography

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Germanium oxide

High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43

Glassy carbon

Assessment of polycrystalline graphites as sorbents for solidphase microextraction of nonionic surfactants 888(2000)35

Glutamyl peptides

Analysis of isomeric glutamyl peptides by capillary electrophoresis. Application to stability studies 888(2000)275

Graphite

Assessment of polycrystalline graphites as sorbents for solidphase microextraction of nonionic surfactants 888(2000)35

Group-correlation method

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Group method

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Heavy metals

Analysis of heavy metals in atmospheric particulate by ion chromatography 888(2000)145

Heteroconjugation

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Hollow-fiber membranes

High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43

Hypericum peforatum

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Immobilized antibodies

Antibody immobilization to high-performance liquid chromatography supports. Characterization of maximum loading capacity for intact immunoglobulin G and Fab fragments 888(2000)13

Immunoaffinity chromatography

Antibody immobilization to high-performance liquid chromatography supports. Characterization of maximum loading capacity for intact immunoglobulin G and Fab fragments 888(2000)13

Immunoaffinity precolumns

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Inorganic anions

Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309

Inverse gas chromatography

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

Ion-pairing reagents

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Lanthanides

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N*,*N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Linear solvation energy relationships

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

Metal cations

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N*,*N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Metalloporphyrins

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Micellar electrokinetic chromatography

Open access capillary electrophoresis. A walk up capillary electrophoresis service for the synthetic chemist 888(2000)219

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

Mixed-mode retention

Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267

Mobile phase composition

Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63

Molecular imprinted sorbents

Affinity screening by packed capillary high-performance liquid chromatography using molecular imprinted sorbents. I. Demonstration of feasibility 888(2000)23

Molecular imprinting

Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63

Molecular mass

Number-average molecular mass determination of polymeric material by pyrolysis-gas chromatography 888(2000)209

Molecularly imprinted polymers

Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63

Mycotoxins

Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321

Naphthodianthrones

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Nitrophenols

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Nonaqueous capillary electrophoresis

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Oakmoss

Identification of atranorin and related potential allergens in oakmoss absolute by high-performance liquid chromatography—tandem mass spectrometry using negative ion atmospheric pressure chemical ionization 888(2000)103

Ochratoxin A

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Ochratoxins

Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321

Open access capillary electrophoresis

Open access capillary electrophoresis. A walk up capillary electrophoresis service for the synthetic chemist 888(2000)219

Ophthalmic solutions

Simultaneous determination of eleven ingredients in ophthalmic solutions by cyclodextrin-modified micellar electrokinetic chromatography with tetrabutylammonium salt 888(2000)299

Optimization

Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197

Orthogonal array design

Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197

Pencil lead

Assessment of polycrystalline graphites as sorbents for solidphase microextraction of nonionic surfactants 888(2000)35

Peptides

Analysis of isomeric glutamyl peptides by capillary electrophoresis. Application to stability studies 888(2000)275

Perchlorate

Improved method for the determination of trace perchlorate in ground and drinking waters by ion chromatography 888(2000)151

Pesticides

Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography—atmospheric pressure chemical ionization mass spectrometry 888(2000)113

Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197

Pharmaceutical analysis

Simultaneous determination of eleven ingredients in ophthalmic solutions by cyclodextrin-modified micellar electrokinetic chromatography with tetrabutylammonium salt 888(2000)299

Phenols

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Phloroglucinols

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Polybutadiene

Number-average molecular mass determination of polymeric material by pyrolysis—gas chromatography 888(2000)209

Polymers

Number-average molecular mass determination of polymeric material by pyrolysis—gas chromatography 888(2000)209

Porous membranes

High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43

Preparative chromatography

On-line monitoring of enantiomer concentration in chiral simulated moving bed chromatography 888(2000)73

Protoporphyrins

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Pyridines

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Pyrolysis

Number-average molecular mass determination of polymeric material by pyrolysis—gas chromatography 888(2000)209

Rare earth ions

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N*,*N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Response factors

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Riboflavins

High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

Sample preparation

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309

Selectivity

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Silica

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Simulated moving bed chromatography

On-line monitoring of enantiomer concentration in chiral simulated moving bed chromatography 888(2000)73

Solid-phase extraction

Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197

Sorbents

Assessment of polycrystalline graphites as sorbents for solidphase microextraction of nonionic surfactants 888(2000)35

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

St. John's Wort

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Stability studies

Analysis of isomeric glutamyl peptides by capillary electrophoresis. Application to stability studies 888(2000)275

Stationary phases, CEC

Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267

Stationary phases, LC

Antibody immobilization to high-performance liquid chromatography supports. Characterization of maximum loading capacity for intact immunoglobulin G and Fab fragments 888(2000)13

High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N*,*N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Sulfur compounds

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Comparison of the thin-layer chromatographic properties of sulfur-containing amino acids and their aminophosphonic analogues 888(2000)335

Supercritical fluid extraction

Simultaneous extraction and derivatization of 2chlorovinylarsonous acid from soils using supercritical and pressurized fluids 888(2000)327

Surface characteristics

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

Surfactants

Assessment of polycrystalline graphites as sorbents for solidphase microextraction of nonionic surfactants 888(2000)35

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

Tenax

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

Tetrabutylammonium phosphate

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Tioconazole

Open access capillary electrophoresis. A walk up capillary electrophoresis service for the synthetic chemist 888(2000)219

Titania

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Toxins

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Transient state

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

Tryptylines

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Twaron

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

Vegetables

Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography—atmospheric pressure chemical ionization mass spectrometry 888(2000)113

Visualisation of wall effects

Physical evidence of two wall effects in liquid chromatography 888(2000)1

Vitamins

High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Wall effects

Physical evidence of two wall effects in liquid chromatography 888(2000)1

Water analysis

Improved method for the determination of trace perchlorate in ground and drinking waters by ion chromatography 888(2000)151

Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197

Wine

High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Zirconia

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51