



ELSEVIER

Journal of Chromatography A, 898 (2000) 279–283

JOURNAL OF
CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Subject Index Vol. 898

Adsorbents

DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167

Affinity chromatography

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

Alcohols

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

Aldehydes

5-Amino-4-sulfanylpthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1

Alkaloids

Gas-liquid chromatographic determination of major tobacco alkaloids 898(2000)235

Alkyl aryl ketones

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

Amines

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Aminosulfanylpthalhydrazide

5-Amino-4-sulfanylpthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1

Aromatic amines

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

Aromatic hydrocarbons

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

Association constants

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

Atomic emission detection

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

Bacterial signaling molecules

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

Benzoic acids

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Biofilms

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

Buffer composition

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Building materials

Determination of ergosterol on mouldy building materials using isotope dilution and gas chromatography–tandem mass spectrometry 898(2000)227

Calibration, compound-independent

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

Capillary columns

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Catechins

Supercritical fluid extraction of catechins from *Cratoxylum prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

Catecholamines

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Cellobiohydrolase

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Cellobiose

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Cellulase

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Chemiluminescence detection

5-Amino-4-sulfanylphthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1

Chiral stationary phases, LC

Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate β -cyclodextrin bonded chiral stationary phase 898(2000)53

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Cigarette smoke

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

Clean-up methods

Application of ion-exchange cartridge clean-up in food analysis. V. Simultaneous determination of sulphonamide antibacterials in animal liver and kidney using high-performance liquid chromatography with ultraviolet and mass spectrometric detection 898(2000)95

Counter-current chromatography

Supercritical fluid extraction of catechins from *Cratoxylum prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

Cratoxylum prunifolium

Supercritical fluid extraction of catechins from *Cratoxylum prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

Derivatization, LC

5-Amino-4-sulfanylphthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1

Detection, GC

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

Detection, LC

5-Amino-4-sulfanylphthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1

Detection, TLC

Fibre optical scanning with high resolution in thin-layer chromatography 898(2000)265

Deuterium

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

Displacement chromatography

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

DNA

DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167

Dyes

Determination of dyes in foodstuffs by capillary zone electrophoresis 898(2000)271

Electroosmotic flow

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Empirical formula

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

Enantiomer separation

Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate β -cyclodextrin bonded chiral stationary phase 898(2000)53

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Ergosterol

Determination of ergosterol on mouldy building materials using isotope dilution and gas chromatography–tandem mass spectrometry 898(2000)227

Fibre optical scanning

Fibre optical scanning with high resolution in thin-layer chromatography 898(2000)265

Fluofix

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

Fluorinated bonded phases

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

Food analysis

Supercritical fluid extraction of catechins from *Cratoxylum prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

Application of ion-exchange cartridge clean-up in food analysis. V. Simultaneous determination of sulphonamide antibacterials in animal liver and kidney using high-performance liquid chromatography with ultraviolet and mass spectrometric detection 898(2000)95

Determination of dyes in foodstuffs by capillary zone electrophoresis 898(2000)271

Frontal chromatography

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

Halogen antimicrobials

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

Height equivalent to a theoretical plate

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

Homoserine lactone

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

Hydrogen

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

Helicobacter pylori

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

Immobilized metal affinity chromatography

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

Isotope analysis

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

Lactose

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Linear solvation energy relationships

Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23

Methylene selectivity

New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35

Milk

Improved method for rapid determination of vitamin A in small samples of breast milk by high-performance liquid chromatography 898(2000)179

Mobile phase composition

Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23

New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35

Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate β -cyclodextrin bonded chiral stationary phase 898(2000)53

Mould

Determination of ergosterol on mouldy building materials using isotope dilution and gas chromatography–tandem mass spectrometry 898(2000)227

Nitrogen

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

Organochlorine pesticides

Solid-phase microextraction for organochlorine pesticide residues analysis in Chinese herbal formulations 898(2000)245

Peptides

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

Perchlorate

Determination of perchlorate at parts-per-billion levels in plants by ion chromatography 898(2000)193

Pesticides

Solid-phase microextraction for organochlorine pesticide residues analysis in Chinese herbal formulations 898(2000)245

Phenols

Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Phenylaminopropyl silica

Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23

Phenyl selectivity

New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35

Polar functional groups

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

Poly(*p*-chloromethylstyrene)

DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167

Poly(ethylene glycol)

Selectivity equivalence of poly(ethylene glycol) stationary phases for gas chromatography 898(2000)211

Polyethylenimine

DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167

Polynuclear aromatic hydrocarbons

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

Polystyrene nanosphere

Separation of unmodified polystyrene nanosphere standards by capillary zone electrophoresis 898(2000)257

Porous copolymers

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

Propranolol

Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate β -cyclodextrin bonded chiral stationary phase 898(2000)53

Proteins

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

Quorum sensing

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

Response factors

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

Retention index scale

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

Retention properties

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

Retinol

Improved method for rapid determination of vitamin A in small samples of breast milk by high-performance liquid chromatography 898(2000)179

Selectivity

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

Selectivity equivalence

Selectivity equivalence of poly(ethylene glycol) stationary phases for gas chromatography 898(2000)211

Simultaneous distillation and extraction

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

Solid-phase microextraction

Solid-phase microextraction for organochlorine pesticide residues analysis in Chinese herbal formulations 898(2000)245

Stationary phases, GC

Selectivity equivalence of poly(ethylene glycol) stationary phases for gas chromatography 898(2000)211

Stationary phases, LC

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23

New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35

Sulphadimethoxine

Simplified determining procedure for routine residue monitoring of sulphamethazine and sulphadimethoxine in milk 898(2000)185

Sulphamethazine

Simplified determining procedure for routine residue monitoring of sulphamethazine and sulphadimethoxine in milk 898(2000)185

Sulphonamides

Application of ion-exchange cartridge clean-up in food analysis. V. Simultaneous determination of sulphonamide antibacterials in animal liver and kidney using high-performance liquid chromatography with ultraviolet and mass spectrometric detection 898(2000)95

Tea

Supercritical fluid extraction of catechins from *Cratoxylum prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

Thermodynamic parameters

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

Tobaccos

Gas-liquid chromatographic determination of major tobacco alkaloids 898(2000)235

Trioctylphosphine oxide

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

Vitamins

Improved method for rapid determination of vitamin A in small samples of breast milk by high-performance liquid chromatography 898(2000)179