



ELSEVIER

Journal of Chromatography A, 901 (2001) 225–339

JOURNAL OF
CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

CUMULATIVE COMPOUND INDEX

J. CHROMATOGR. A VOLS. 852–900

For all compound groups: see also sub-groups and individual compounds

For all compound sub-groups: see also main groups and individual compounds

For all individual compound: see also compound groups and sub-groups

A

A-40,926

Evaluation of the macrocyclic glycopeptide A-40,926 as a high-performance liquid chromatographic chiral selector and comparison with teicoplanin chiral stationary phase 897(2000)113

Abamectin

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Liquid chromatographic determination of emamectin, milbemectin, ivermectin and abamectin in crops and confirmation by liquid chromatography–mass spectrometry 896(2000)75

Acamprosate

Determination of homotaurine as impurity in calcium acamprosate by capillary zone electrophoresis 853(1999)421

AccQ.Fluor reagent

Effect of solvents on the fumonisins analysis by high-performance liquid chromatography with AccQ.Fluor as the derivatizing reagent 870(2000)469

Acenaphthene

Simultaneous determination of ethylbenzene, indan, indene and acenaphthene in air by capillary gas chromatography 891(2000)361

Acetaldehyde

Poly(allylamine) beads as selective sorbent for preconcentration of formaldehyde and acetaldehyde in high-performance liquid chromatographic analysis 886(2000)83

Acetals

Selection of methylation procedures for quantitation of short-chain glycerol-bound compounds formed during thermo-oxidation 863(1999)171

Acetaminophen

see Paracetamol

Acetic acid

Determination of acetic acid in aqueous samples, by water-phase derivatisation, solid-phase microextraction and gas chromatography 874(2000)225

Acetone

Determination of trace anions in organic solvents 858(1999)55

Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

Acetonitrile

Study of dead volume measurement in packed subcritical fluid chromatography with ODS columns and carbon dioxide–modifier mobile phases 866(2000)241

Acetylacetone

Liquid chromatography–mass spectrometry method for the determination of aldehydes derivatized by the Hantzsch reaction 864(1999)191

Acetylbenzocaine

Liquid chromatographic determination of benzocaine and *N*-acetylbenzocaine in the edible fillet tissue from rainbow trout 855(1999)255

Acetyl-L-carnitine

Enantiomeric purity determination of acetyl-L-carnitine by reversed-phase high-performance liquid chromatography using chiral derivatization 857(1999)127

Acetylcholinesterase inhibitors

High-performance liquid chromatography with on-line coupled UV, mass spectrometric and biochemical detection for identification of acetylcholinesterase inhibitors from natural products 872(2000)61

Acetylcysteine

Capillary electrophoretic separation of *N*-acetylcysteine and its impurities as a method for quality control of pharmaceuticals 853(1999)479

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191

Acetylsalicylic acid

Validation of the removal of acetylsalicylic acid. Recovery and determination of residues on various surfaces by high performance liquid chromatographic 870(2000)69

Acetyls erine

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Acridinone

Sample cleanup and reversed-phase high-performance liquid chromatographic analysis of polar aromatic compounds in groundwater samples from a former gas plant 862(1999)137

Acrinathrine

Gas chromatographic determination of acrinathrine and 3-phenoxybenzaldehyde residues in honey 882(2000)239

Acrylamide–allylcyclodextrin copolymer

Determination of cyclodextrin content using periodate oxidation by capillary electrophoresis 891(2000)201

Actin

Analysis of leptin gene expression in chickens using reverse transcription polymerase chain reaction and capillary electrophoresis with laser-induced fluorescence detection 853(1999)321

Actinides

Determination of U(IV) and U(VI) by ion chromatography–inductively coupled plasma mass spectrometry and its application to kinetic studies 884(2000)131

Acyclovir

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Acylcarnitines

Direct chromatographic resolution of carnitine and *O*-acylcarnitine enantiomers on a teicoplanin-bonded chiral stationary phase 857(1999)145

Analysis of carnitine and acylcarnitines in urine by capillary electrophoresis 895(2000)309

Acylglycerols

Preparative thin-layer chromatographic separation and subsequent gas chromatographic–mass spectrometric analysis of monoacylglycerols derived from butter oil by fungal degradation 855(1999)617

Analytical monitoring of the production of biodiesel by high-performance liquid chromatography with various detection methods 858(1999)13

Adrenaline

see Epinephrine

Aerosol MA 80-I

High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253

Aflatoxins

see also Toxins

Application of the assay of aflatoxins by liquid chromatography with fluorescence detection in food analysis 882(2000)1

Determination of aflatoxins in food by overpressured-layer chromatography 882(2000)11

Aglycones

Reversed-phase liquid chromatographic separation and simultaneous profiling of steroidal glycoalkaloids and their aglycones 863(1999)25

Analysis of enzymatically glucosylated flavonoids by capillary electrophoresis 895(2000)279

Alachlor

Optimisation of alachlor solid-phase microextraction from water samples using experimental design 896(2000)373

L-Alanine- β -naphthylamide

Enantiomeric purity determination of acetyl-L-carnitine by reversed-phase high-performance liquid chromatography using chiral derivatization 857(1999)127

Albendazole

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Albumins

Experimental studies on affinity chromatography in an electric field 852(1999)319

Macroporous poly(glycidyl methacrylate–triallyl isocyanurate–divinylbenzene) matrix as an anion-exchange resin for protein adsorption 855(1999)129

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Study of hydrophobic interaction adsorption of bovine serum albumin under overloaded conditions using flow microcalorimetry 865(1999)111

Kinetic study of the mass transfer of bovine serum albumin in anion-exchange chromatography 866(2000)147

Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis 871(2000)311

Erratum to: “Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles” [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

Stereoselective binding of 2,3-substituted 3-hydroxypropionic acids on an immobilised human serum albumin chiral stationary phase: stereochemical characterisation and quantitative structure–retention relationship study 876(2000)75

New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169

Scale-down of continuous protein purification by annular chromatography. Design parameters for the smallest unit 890(2000)53

Capillary electrophoresis of seed 2S albumins from *Lupinus* species 894(2000)297

Sodium dodecylsulfate capillary gel electrophoretic measurement of the concentration ratios of albumin and α_2 -macroglobulin in cerebrospinal fluid and serum of patients with neurological disorders 895(2000)339

Alcohol ethoxylates

Analysis of aliphatic alcohol ethoxylates in terms of alkyl and ethylene oxide chain lengths by reversed-phase liquid chromatography with evaporative light scattering detection 874(2000)305

Alcohols

see also Volatile organic compounds

Gas chromatographic determination of primary alcohols as their formylbenzoic esters by gas-phase luminescence detection 855(1999)583

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

Modeling chromatographic parameters by a novel graph theoretical sub-structural approach 858(1999)187

Determination of halogenated mono-alcohols and diols in water by gas chromatography with electron-capture detection 866(2000)65

2-(4-Carboxyphenyl)-6-*N,N*-diethylaminobenzofuran: a useful reagent for the sensitive determination of alcohols by high-performance liquid chromatography with fluorimetric detection 869(2000)243

Utilization of fluorescein sodium salt for the indirect fluorimetric detection in micellar electrokinetic chromatography 872(2000)247

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Electrophoretic methods in the analysis of beverages 880(2000)243

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

Aldehydes

see also Carbonyl compounds and Volatile organic compounds

Liquid chromatography–mass spectrometry method for the determination of aldehydes derivatized by the Hantzsch reaction 864(1999)191

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Determination of aldehydes in food by high-performance liquid chromatography with biosensor coupling and micromembrane suppressors 880(2000)233

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

5-Amino-4-sulfanylphthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1

Aldehydic acids

Selection of methylation procedures for quantitation of short-chain glycerol-bound compounds formed during thermoxidation 863(1999)171

Aldicarb

Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 888(2000)113

Alditols

Separation and determination of alditols and sugars by high-pH anion-exchange chromatography with pulsed amperometric detection 897(2000)195

Alkali metals

Ion-chromatographic behavior of alkali metal cations and ammonium ion on zirconium-adsorbing silica gel 884(2000)123

Alkaline phosphatase

Enzyme amplification as detection tool in continuous-flow systems I. Development of an enzyme-amplified biochemical detection system coupled on-line to flow-injection analysis 855(1999)383

Enzyme amplification as detection tool in continuous-flow systems II. On-line coupling of liquid chromatography to enzyme-amplified biochemical detection after pre-column derivatization with biotin 855(1999)397

Alkaloids

Optimization study for the reversed-phase ion-pair liquid chromatographic determination of nicotine in commercial tobacco products 852(1999)451

Application of non-aqueous capillary electrophoresis with electrochemical detection to the determination of nicotine in tobacco 853(1999)27

Capillary electrophoresis separation and permanganate chemiluminescence on-line detection of some alkaloids with β -cyclodextrin as an additive 855(1999)329

Determination of quinolizidine alkaloids in traditional Chinese herbal drugs by nonaqueous capillary electrophoresis 857(1999)303

Pre-column derivatization and gas chromatographic determination of alkaloids in bulbs of *Fritillaria* 859(1999)183

Reversed-phase liquid chromatographic separation and simultaneous profiling of steroidal glycoalkaloids and their aglycones 863(1999)25

Strategies for supercritical fluid extraction of hyoscyamine and scopolamine salts using basified modifiers 863(1999)47

Separation, identification and determination of sanguinarine in argemone and other adulterated edible oils by reversed-phase high-performance liquid chromatography 863(1999)123

Analysis of coptisine, berberine and palmatine in adulterated Chinese medicine by capillary electrophoresis–electrospray ion trap mass spectrometry 866(2000)273

Capillary electrophoretic determination of sanguinarine and chelerythrine in plant extracts and pharmaceutical preparations 866(2000)293

Enantioseparation of atropine by capillary electrophoresis using sulfated β -cyclodextrin: application to a plant extract 868(2000)285

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

Simultaneous determination of seven major isosteroidal alkaloids in bulbs of *Fritillaria* by gas chromatography 873(2000)221

Non-aqueous capillary electrophoretic enantioseparation of *N*-derivatized amino acids using cinchona alkaloids and derivatives as chiral counter-ions 875(2000)353

Analysis of tetramethylpyrazine in *Ephedrae herba* by gas chromatography–mass spectrometry and high-performance liquid chromatography 878(2000)147

On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic–diode array detection–mass spectrometric techniques 880(2000)203

Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425

Analysis of the bioactive alkaloids tetrahydro- β -carboline and β -carboline in food 881(2000)483

Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography (II) 891(2000)189

Separation of the four pairs of enantiomers of vincamine alkaloids by enantioselective high-performance liquid chromatography 893(2000)47

Gas-liquid chromatographic determination of major tobacco alkaloids 898(2000)235

Alkanals

Dynamic system for the calibration of semi-volatile carbonyl compounds in air 883(2000)171

Alkanes

Revision of a theoretical expression for gas–liquid chromatographic retention 852(1999)583

Modeling chromatographic parameters by a novel graph theoretical sub-structural approach 858(1999)187

Analysis of the equations for the temperature dependence of the retention index. II. Physico-chemical meaning of the parameters 859(1999)49

Lattice–fluid model for gas–liquid chromatography 862(1999)49

Interpreting the gas chromatographic retention of *n*-alkanes 873(2000)209

Hold-up time in gas chromatography. V. Dependence of the retention of *n*-alkanes on the chromatographic variables in isothermal gas chromatography 878(2000)125

Evaluation of time-of-flight mass spectrometric detection for fast gas chromatography 878(2000)205

Distribution coefficients of *n*-alkanes measured on wall-coated capillary columns 879(2000)157

Comparison of comprehensive two-dimensional gas chromatography and gas chromatography – mass spectrometry for the characterization of complex hydrocarbon mixtures 892(2000)29

Alkenes

Comparison of comprehensive two-dimensional gas chromatography and gas chromatography – mass spectrometry for the characterization of complex hydrocarbon mixtures 892(2000)29

Alkenylbenzenes

Solid phase microextraction of alkenylbenzenes and other flavor-related compounds from tobacco for analysis by selected ion monitoring gas chromatography–mass spectrometry 858(1999)79

S-Alk(en)ylcysteine sulfoxide

Gas chromatographic determination of *S*-alk(en)ylcysteine sulfoxides 862(1999)85

Alkylamines

Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285

Alkyl aryl ketones

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

Alkylbenzenes

Preparation of uniformly sized polymeric separation media potentially suitable for small-scale high-performance liquid chromatography and/or capillary electrochromatography 853(1999)11

Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

Life-time studies with capillary electrochromatography columns operated under different conditions 894(2000)35

Alkyl benzenesulfonates linear

Analysis of linear alkylbenzenesulfonates in water samples by large-volume injection-port derivatization and gas chromatography–mass spectrometry 857(1999)359

Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99

Assessment of the pollutant elimination efficiency by gas chromatography/mass spectrometry, liquid chromatography–mass spectrometry and –tandem mass spectrometry. Comparison of conventional and membrane-assisted biological wastewater treatment processes 889(2000)155

Inter-laboratory comparison of liquid chromatographic techniques and enzyme-linked immunosorbent assay for the determination of surfactants in wastewaters 889(2000)195

Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211

Determination of linear alkylbenzenesulfonates in wastewater treatment plants and coastal waters by automated solid-phase extraction followed by capillary electrophoresis–UV detection and confirmation by capillary electrophoresis–mass spectrometry 889(2000)221

2-Alkylcyclobutanones

2-Alkylcyclobutanones as markers for irradiated foodstuffs. III. Improvement of the field of application on the EN 1785 method by using silver ion chromatography 858(1999)109

Supercritical fluid extraction of hydrocarbons and 2-alkylcyclobutanones for the detection of irradiated foodstuffs 897(2000)259

Alkyl diphenyloxide disulfonates

High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253

Alkylmethoxypyrazines

Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxypyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93

Alkyl methylphosphonic acids

Efficiency of pretreatment of aqueous samples using a macroporous strong anion-exchange resin on the determination of nerve gas hydrolysis products by gas chromatography–mass spectrometry after *tert*.-butyldimethylsilylation 891(2000)295

Alkyl nitrates

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a β -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

Alkylphenols

Quantification of the xenoestrogens 4-*tert*.-octylphenol and bisphenol A in water and in fish tissue based on microwave assisted extraction, solid-phase extraction and liquid chromatography–mass spectrometry 864(1999)17

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

Separation of bisphenol A and three alkylphenols by micellar electrokinetic chromatography 895(2000)213

Separation of two groups of oestrogen mimicking compounds using micellar electrokinetic chromatography 895(2000)237

Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279

Alkylphosphonic acids

Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169

Alkyl polyglucosides

Investigations on the metabolism of alkyl polyglucosides and their determination in waste water by means of liquid chromatography–electrospray mass spectrometry 854(1999)221

Alkyl polyglycol ethers

Assessment of the pollutant elimination efficiency by gas chromatography/mass spectrometry, liquid chromatography–mass spectrometry and –tandem mass spectrometry. Comparison of conventional and membrane-assisted biological wastewater treatment processes 889(2000)155

Alkylsulfonic acids

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

Alkyltrimethylammonium surfactants

Alkyltrimethylammonium surfactant-mediated extractions: Characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Alkyl xanthates

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

Allergens

Identification of atranorin and related potential allergens in oakmoss absolute by high-performance liquid chromatography–tandem mass spectrometry using negative ion atmospheric pressure chemical ionization 888(2000)103

Characterisation of the rubber elongation factor from ammoniated latex by electrophoresis and mass spectrometry 890(2000)145

Allyltropane

Analysis of 2 β -carbomethoxy-3 β -(4-fluorophenyl)-*N*-(3-iodo-*E*-allyl)nortropine in rat plasma. I. Method development and validation by capillary electrophoresis 895(2000)87

Allyl carbamoylated β -cyclodextrin derivatives

Enantiomeric separations of cationic and neutral compounds by capillary electrochromatography with monolithic chiral stationary phases of β -cyclodextrin-bonded negatively charged polyacrylamide gels 893(2000)177

Altropane

Analysis of 2 β -carbomethoxy-3 β -(4-fluorophenyl)-*N*-(3-iodo-*E*-allyl)nortropane in rat plasma. I. Method development and validation by capillary electrophoresis 895(2000)87

Amides

Erratum to: "Determination of fatty acid amides as trimethylsilyl derivatives by gas chromatography with mass spectrometric detection" [J. Chromatogr. 849 (1999) 541–552] 859(1999)119

Large degree of racemization observed in the amide bond forming reaction on silica gel 878(2000)165

Amines

see also Biogenic amines and Catecholamines

Use of volatile amines as ion-pairing agents for the high-performance liquid chromatographic–tandem mass spectrometric determination of aromatic sulfonates in industrial wastewater 854(1999)175

Comparative study of divalent metals and amines as silanol-blocking agents in reversed-phase liquid chromatography 855(1999)121

Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1

Separation of enantiomers on anion exchangers modified with heparin in liquid chromatography 864(1999)25

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Protein adsorption to the bare silica wall in capillary electrophoresis. Quantitative study on the chemical composition of the background electrolyte for minimising the phenomenon 868(2000)85

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Determination of heterocyclic aromatic amines in meat extracts by liquid chromatography–ion-trap atmospheric pressure chemical ionization mass spectrometry 869(2000)307

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Capillary electrochromatography of basic compounds using octadecyl-silica stationary phases with an amine-containing mobile phase 872(2000)315

Electrophoretic methods in the analysis of beverages 880(2000)243

Combined ion-pair extraction and gas chromatography–mass spectrometry for the simultaneous determination of diamines, polyamines and aromatic amines in Port wine and grape juice 886(2000)183

Novel, trifunctional diamine for silica coating in capillary zone electrophoresis 894(2000)53

Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79

Effects of aliphatic amines on capillary electrochromatographic performance of tricyclic antidepressants on octadecylsilica 895(2000)17

Simultaneous determination of inorganic anions and cations by capillary electrophoresis with indirect UV detection 895(2000)269

Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285

Surface-ionization methods and devices of indication and identification of nitrogen-containing base molecules 896(2000)3

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Amines, aromatic

Determination of aromatic amines in water samples by capillary electrophoresis with electrochemical and fluorescence detection 895(2000)197

Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

Amines, heterocyclic

Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295

Analysis of the bioactive alkaloids tetrahydro- β -carboline and β -carboline in food 881(2000)483

Amines, heterocyclic aromatic

Determination of heterocyclic aromatic amines in meat extracts by liquid chromatography–ion-trap atmospheric pressure chemical ionization mass spectrometry 869(2000)307

Comparison of different commercial solid-phase extraction cartridges used to extract heterocyclic amines from a lyophilised meat extract 880(2000)101

Quantitative analysis of mutagenic heterocyclic aromatic amines in cooked meat using liquid chromatography–atmospheric pressure chemical ionisation tandem mass spectrometry 883(2000)89

Amino acids

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

Capillary electrochromatography with thermo-optical absorbance detection for the analysis of phenylthiohydantoin-amino acids 853(1999)131

Determination of 20 underivatized proteinic amino acids by ion-pairing chromatography and pneumatically assisted electrospray mass spectrometry 855(1999)191

Ultra-short columns for low-pressure ion chromatography 857(1999)175

Analysis of β -*N*-oxalyl-L- α , β -diaminopropionic acid and homoarginine in *Lathyrus sativus* by capillary zone electrophoresis 857(1999)295

Gas chromatographic quantification of amino acid enantiomers in food matrices by their *N*(*O*,*S*)-ethoxycarbonyl heptafluorobutyl ester derivatives 866(2000)221

Chiral and electrokinetic separation of amino acids using polypyrrole-coated adsorbents 868(2000)189

Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic β -substituted α -amino acids on a teicoplanin chiral stationary phase 868(2000)197

Novel cinchona alkaloid carbamate C₉-dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255

Retention/quantitation properties of the *o*-phthalaldehyde–3-mercaptopropionic acid and the *o*-phthalaldehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

High-performance liquid chromatographic separation of the enantiomers of unusual α -amino acid analogues 871(2000)105

Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Isolation of pure (2*S*,1'*S*,2'*S*)-2-(2'-carboxycyclopropyl)glycine from *Blighia sapida* (Akee) 873(2000)283

Chromatographic resolution of tryptophan enantiomers with L-Leu–L-Leu–L-Leu peptide. Effects of mobile phase composition and chromatographic support 874(2000)21

Chiral separation of unmodified amino acids with non-aqueous capillary electrophoresis based on the ligand-exchange principle 874(2000)285

Chiral separation of amino acids and peptides by capillary electrophoresis 875(2000)43

Enantiomer separation of drugs by micellar electrokinetic chromatography using chiral surfactants 875(2000)163

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Investigation of an enantioselective non-aqueous capillary electrochromatography system applied to the separation of chiral acids 875(2000)341

Non-aqueous capillary electrophoretic enantioseparation of *N*-derivatized amino acids using cinchona alkaloids and derivatives as chiral counter-ions 875(2000)353

Micellar electrokinetic capillary chromatographic separation and fluorescent detection of amino acids derivatized with 4-fluoro-7-nitro-2,1,3-benzoxadiazole 876(2000)183

- Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217
- Amperometric determination of underivatized amino acids at a nickel-modified gold electrode by anion-exchange chromatography 878(2000)57
- Electrophoretic methods in the analysis of beverages 880(2000)243
- Effect of the activity levels of the added proteolytic enzyme mixture on free amino acids in ripening Ossau-Iraty cheese 881(2000)69
- Chromatographic determination of amino acid enantiomers in beers and raw materials used for their manufacture 881(2000)81
- Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317
- Use of nucleic acids in the mobile phase for the determination of ascorbic acid in foods by high-performance liquid chromatography with electrochemical detection 881(2000)327
- Determination of organic acids in food samples by capillary zone electrophoresis 881(2000)357
- Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425
- Tetrahydro- β -carboline-3-carboxylic acids and contaminants of L-tryptophan 881(2000)501
- Monitoring and control of industrial downstream processing of sugar beet molasses 882(2000)329
- Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs 883(2000)249
- Low-capacity cation-exchange chromatography of ultraviolet-absorbing urinary basic metabolites using a reversed-phase column coated with hexadecylsulfonate 886(2000)297
- Comparison of the thin-layer chromatographic properties of sulfur-containing amino acids and their aminophosphonic analogues 888(2000)335
- Column-switching system for selenium speciation by coupling reversed-phase and ion-exchange high-performance liquid chromatography with microwave-assisted digestion–hydride generation–atomic fluorescence spectrometry 889(2000)33
- Role of chromatography in the analysis of sugars, carboxylic acids and amino acids in food (Review) 891(2000)1
- Use of amino acid analysis for estimating the individual concentrations of proteins in mixtures 891(2000)355
- tert*-Butylcarbamoylquinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique 892(2000)509
- Enantioseparation of anionic analytes by non-aqueous capillary electrophoresis using quinine and quinidine derivatives as chiral counter-ions 894(2000)63
- Allylamine- β -cyclodextrin copolymer. A novel chiral selector for capillary electrophoresis 894(2000)95
- Direct determination of amino acids and carbohydrates by high-performance capillary electrophoresis with refractometric detection 894(2000)253
- Chiral determination of amino acids by capillary electrophoresis and laser-induced fluorescence at picomolar concentrations 894(2000)259
- Pressurized gradient capillary electrochromatographic separation of eighteen amino acid derivatives 894(2000)337
- Determination of biologically active low-molecular-mass thiols in human blood. I. Fast qualitative and quantitative, gradient and isocratic reversed-phase high-performance liquid chromatography with photometric and fluorescence detection 895(2000)156
- Determination of biologically active low-molecular-mass thiols in human blood. II. High-performance capillary electrophoresis with photometric detection 895(2000)167
- Assays for total homocysteine and other thiols by capillary electrophoresis–laser-induced fluorescence detection. I. Preanalytical condition studies 895(2000)173
- Determination of optical purity of phosphonic acid analogues of aromatic amino acids by capillary electrophoresis with α -cyclodextrin 895(2000)301
- Screening of homocysteine from newborn blood spots by high-performance liquid chromatography with coulometric array detection 896(2000)183
- Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191
- Parameter optimization for the analysis of underivatized protein amino acids by liquid chromatography and ionspray tandem mass spectrometry 896(2000)253
- Analysis of the enzymatic racemization of D-aspartic acid to L-aspartic acid by the on-line coupling of a solid-phase extraction column and a ligand-exchange high-performance liquid chromatography column 896(2000)299

Determination of 1-aminocyclopropane-1-carboxylic acid and its structural analogue by liquid chromatography and ion spray tandem mass spectrometry 896(2000)335

Evaluation of the macrocyclic glycopeptide A-40,926 as a high-performance liquid chromatographic chiral selector and comparison with teicoplanin chiral stationary phase 897(2000)113

Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic β -substituted α -amino acids on a copper(II)-D-penicillamine chiral stationary phase 897(2000)145

Amino acids, butyloxycarbonyl derivatives

Enantioseparation of selected *N*-*tert*-butyloxycarbonyl amino acids in high-performance liquid chromatography and capillary electrophoresis with a teicoplanin chiral selector 879(2000)147

Amino acids, DNB derivatives

Quinine versus carbamoylated quinine-based chiral anion exchangers. A comparison regarding enantioselectivity for *N*-protected amino acids and other chiral acids 858(1999)1

Amino acids, DNP derivatives

Quinine versus carbamoylated quinine-based chiral anion exchangers. A comparison regarding enantioselectivity for *N*-protected amino acids and other chiral acids 858(1999)1

Amino acids, Dns derivatives

Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271

New amphiphilic aminosaccharide derivatives as chiral selectors in capillary electrophoresis 875(2000)295

Amino acids, non-protein

Determination of neurotoxin 3-*N*-oxalyl-2,3-diaminopropionic acid and non-protein amino acids in *Lathyrus sativus* by precolumn derivatization with 1-fluoro-2,4-dinitrobenzene 883(2000)113

Amino acids, PTH derivatives

Capillary electrochromatography with thermo-optical absorbance detection for the analysis of phenylthiohydantoin-amino acids 853(1999)131

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255

Amino alcohols

Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1

Enantiomeric separation of amino alcohols on protein phases using statistical experimental design. A comparative study 866(2000)15

Determination of the enantiomers of 3-*tert*-butylamino-1,2-propanediol by high-performance liquid chromatography using mass spectrometric detection 896(2000)201

Aminobenzamides

3-Aminobenzamide and 3-aminobenzoic acid, tags for capillary electrophoresis of complex carbohydrates with laser-induced fluorescent detection 863(1999)205

2-Aminobenzimidazole

Determination of thiophanate-methyl and its metabolites at trace level in spiked natural water using the supported liquid membrane extraction and the microporous membrane liquid-liquid extraction techniques combined on-line with high-performance liquid chromatography 893(2000)123

Aminobenzoic acids

3-Aminobenzamide and 3-aminobenzoic acid, tags for capillary electrophoresis of complex carbohydrates with laser-induced fluorescent detection 863(1999)205

Aminobiphenyl

Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291

Aminocaproic acid

Standardization of capillary zone electropherograms obtained by using field enhanced sample stacking 894(2000)3

Aminocaprolactam

Separation of enantiomers by capillary electrophoresis–mass spectrometry employing a partial filling technique with a chiral crown ether 875(2000)323

Aminochlorophenylbutyric acid

Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233

Amino compounds, primary

Enhancement of chiral recognition by formation of a sandwiched complex in capillary electrophoresis 875(2000)361

Aminocyclohexyldichlorobenzamide

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

Aminocyclopropanecarboxylic acid

Determination of 1-aminocyclopropane-1-carboxylic acid and its structural analogue by liquid chromatography and ion spray tandem mass spectrometry 896(2000)335

Aminoglutethimide

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

4-Amino-3-hydroxybutyric acid

Validation of an analysis method for 4-amino-3-hydroxybutyric acid by reversed-phase liquid chromatography 890(2000)273

1-Amino-2-hydroxycyclohexanecarboxylic acids

Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic β -substituted α -amino acids on a teicoplanin chiral stationary phase 868(2000)197

Aminoindanol

Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated β -cyclodextrin 875(2000)389

Aminoisobutyric acid

Detection of new sequences of peptaibol antibiotics trichothoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179

Chiral determination of amino acids by capillary electrophoresis and laser-induced fluorescence at picomolar concentrations 894(2000)259

Aminolevulinic acid

Validation of a capillary electrophoresis method for determination of 5-aminolevulinic acid and degradation products 883(2000)285

1-Amino-2-methylcyclohexanecarboxylic acids

Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic β -substituted α -amino acids on a teicoplanin chiral stationary phase 868(2000)197

Aminomethylphosphonic acid

New methods for determination of glyphosate and (aminomethyl)phosphonic acid in water and soil 886(2000)207

Amino(methylsulfanyl)phenyldiazenylphenylsulfonylethanol

Determination of gas phase peroxyacetic acid using pre-column derivatization with organic sulfide reagents and liquid chromatography 855(1999)551

Aminonaphthalenedisulfonic acid

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Aminonaphthalene trisulfonate

Analysis of 8-aminonaphthalene-1,3,6-trisulfonate-derivatized oligosaccharides by capillary electrophoresis–electrospray ionization quadrupole ion trap mass spectrometry 858(1999)229

Aminophenols

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Aminophenylmethylbenzothiazole

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Aminophosphonic acids

Comparison of the thin-layer chromatographic properties of sulfur-containing amino acids and their aminophosphonic analogues 888(2000)335

Aminopyridines

Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277

Aminopyrrolidine

Separation of enantiomers by capillary electrophoresis–mass spectrometry employing a partial filling technique with a chiral crown ether 875(2000)323

Aminosaccharides

New amphiphilic aminosaccharide derivatives as chiral selectors in capillary electrophoresis 875(2000)295

Aminosalicylic acid

Apparatus and method for interfacing capillary electrophoresis and chemiluminescence nitrogen detection for the analysis of nitrogen-containing compounds 868(2000)121

Aminosulfanylphthalhydrazide

5-Amino-4-sulfanylphthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1

Aminosulfonylfluorobenzoxadiazole

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Aminothiols

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Determination of biologically active low-molecular-mass thiols in human blood. I. Fast qualitative and quantitative, gradient and isocratic reversed-phase high-performance liquid chromatography with photometric and fluorescence detection 895(2000)156

Determination of biologically active low-molecular-mass thiols in human blood. II. High-performance capillary electrophoresis with photometric detection 895(2000)167

Ammonium

Determination of ammonium in seawater by column-switching ion chromatography 868(2000)209

Ion-chromatographic behavior of alkali metal cations and ammonium ion on zirconium-adsorbing silica gel 884(2000)123

Determination of ammonium in milk and dairy products by ion chromatography 893(2000)133

Amphetamines

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

Development of a simple in-vial liquid-phase microextraction device for drug analysis compatible with capillary gas chromatography, capillary electrophoresis and high-performance liquid chromatography 873(2000)3

Simultaneous analysis of some amphetamine derivatives in urine by nonaqueous capillary electrophoresis coupled to electrospray ionization mass spectrometry 895(2000)111

Determination of amphetamine and methamphetamine in serum via headspace derivatization solid-phase microextraction–gas chromatography–mass spectrometry 896(2000)265

Ampicillin

Stability of penicillin antibiotic residues in meat during storage. Ampicillin 882(2000)135

Amygdalin

Micellar electrokinetic chromatography for the analysis of *D*-amygdalin and its epimer in apricot kernel 866(2000)253

Amyloids

Congophilicity (Congo red affinity) of different β_2 -microglobulin conformations characterized by dye affinity capillary electrophoresis 894(2000)319

Anaesthetics

Highly sensitive bioassay of lidocaine in human plasma by high-performance liquid chromatography–tandem mass spectrometry 854(1999)3

Anatoxins

Analysis of anatoxin-a in freshwaters by automated on-line derivatization–liquid chromatography–electrospray mass spectrometry 862(1999)191

Angiotensin-converting enzyme

Analysis of angiotensin-converting enzyme by capillary electrophoresis 853(1999)185

Angiotensin-converting enzyme inhibitors

Optimization of capillary electrophoretic separation of several inhibitors of the angiotensin-converting enzyme 895(2000)33

Angiotensins

New polymer-based prepacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

Simultaneous separation of angiotensin and endothelin peptide families by high-performance liquid chromatography: application to the specific radioimmunoassay measurement of angiotensin II or endothelin-1 from tissue 870(2000)349

Anhydrides, aliphatic

Determination of aliphatic anhydrides and acids by reversed-phase liquid chromatography 868(2000)41

Anilines

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

Capillary electrochromatography with a silica column with a dynamically modified cationic surfactant 855(1999)137

Liquid chromatographic determination of aniline in table-top sweeteners based on pre-column derivatization with 1,2-naphthoquinone-4-sulfonate 859(1999)227

Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Determination of aromatic amines in water samples by capillary electrophoresis with electrochemical and fluorescence detection 895(2000)197

Anions

see also Inorganic anions and Organic acids

Determination of bromide, bromate and other anions with ion chromatography and an inductively coupled plasma mass spectrometer as element-specific detector 862(1999)39

Ion exchange-based preconcentration for the determination of anions by capillary electrophoresis 868(2000)135

Simultaneous determination of inorganic disinfection by-products and the seven standard anions by ion chromatography 890(2000)295

Anions, aromatic

Equilibrium analysis of reactions between aromatic anions and nonionic surfactant micelles by capillary zone electrophoresis 853(1999)55

Anthocyanins

Analysis of pigmented high-molecular-mass grape phenolics using ion-pair, normal-phase high-performance liquid chromatography 866(2000)25

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Electrophoretic methods in the analysis of beverages 880(2000)243

Analysis of anthocyanins in foods by liquid chromatography, liquid chromatography–mass spectrometry and capillary electrophoresis 881(2000)403

Anthracenes

Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

Anthracyclines

Anthracycline analysis by capillary electrophoresis. Application to the analysis of daunorubicin in Kaposi sarcoma tumor 853(1999)449

Anthraquinones

Preparative isolation and purification of hydroxyanthraquinones from *Rheum officinale* Baill by high-speed counter-current chromatography using pH-modulated stepwise elution 858(1999)103

High-performance liquid chromatographic determination of some anthraquinone and naphthoquinone dyes occurring in historical textiles 863(1999)235

Analysis of natural food pigments by capillary electrophoresis 880(2000)311

Antibacterial agents

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Antibiotics

Application of ion-exchange cartridge clean-up in food analysis II. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in meat using liquid chromatography with ultraviolet detection 855(1999)247

Micellar electrokinetic capillary chromatography of macrolide antibiotics. Separation of tylosin, erythromycin and their related substances 857(1999)313

Detection of new sequences of peptaibol antibiotics trichothoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179

High-performance thin-layer chromatography method for monitoring norfloxacin residues on pharmaceutical equipment surfaces 862(1999)209

Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69

Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants 868(2000)313

Reply to ‘Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants’ 868(2000)317

Separation of erythromycin and related substances on base-deactivated reversed-phase silica gel columns 870(2000)227

Liquid chromatography of polymyxin B sulphate 870(2000)237

Determination of quinolones in animal tissues and eggs by high-performance liquid chromatography with photodiode-array detection 871(2000)31

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Characterization of partition and thermodynamic properties of cephalosporins using micellar electrokinetic chromatography in glycodeoxycholic acid solution 873(2000)237

Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Optimization of separation and migration behavior of cephalosporins in capillary zone electrophoresis 879(2000)197

Analysis of polymyxin B sulfate by capillary zone electrophoresis with cyclodextrin as additive. Method development and validation 879(2000)211

Application of ion-exchange cartridge clean-up in food analysis. III. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in bovine liver and kidney by liquid chromatography with ultraviolet detection 880(2000)85

Electrophoretic methods in the analysis of beverages 880(2000)243

Quantitation of nine quinolones in chicken tissues by high-performance liquid chromatography with fluorescence detection 882(2000)63

Chromatographic analysis of tetracycline antibiotics in foods 882(2000)109

Stability of penicillin antibiotic residues in meat during storage. Ampicillin 882(2000)135

Determination of cephalosporins in raw bovine milk by high-performance liquid chromatography 882(2000)145

New configurations and applications of molecularly imprinted polymers 889(2000)15

Adaptation of capillary electrophoresis to the determination of selected cephalosporins for injection 895(2000)27

Analysis of doxycycline by capillary electrophoresis. Method development and validation 895(2000)43

Capillary Electrophoresis analysis of gentamicin sulphate with UV detection after pre-capillary derivatization with 1,2-phthalic dicarboxaldehyde and mercaptoacetic acid 895(2000)67

Evaluation of the macrocyclic glycopeptide A-40,926 as a high-performance liquid chromatographic chiral selector and comparison with teicoplanin chiral stationary phase 897(2000)113

Ion chromatographic analysis of tetracyclines using polymeric column and acidic eluent 897(2000)205

Antibodies

see also Monoclonal antibodies

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

Quantitative fast fractionation of a pool of polyclonal antibodies by immunoaffinity membrane chromatography 852(1999)129

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Antidepressants, tricyclic

Optimized stationary phases for the high-performance liquid chromatography–electrospray ionization mass spectrometric analysis of basic pharmaceuticals 869(2000)159

Effects of aliphatic amines on capillary electrochromatographic performance of tricyclic antidepressants on octadecylsilica 895(2000)17

Antifouling compounds

Trace determination of antifouling compounds by on-line solid-phase extraction–gas chromatography–mass spectrometry 885(2000)361

Antigens

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Antimicrobials

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

Antioxidants

Determination of low levels of an antioxidant in polyolefins by large-volume injection temperature-programmed packed capillary liquid chromatography 864(1999)103

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

Micellar electrokinetic chromatography with bis-(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Chromatographic analysis of tocol-derived lipid antioxidants 881(2000)197

Simultaneous determination of α -tocopherol and β -carotene in olive oil by reversed-phase high-performance liquid chromatography 881(2000)255

Analysis of several phenolic compounds with potential antioxidant properties in grape extracts and wines by high-performance liquid chromatography–photodiode array detection without sample preparation 881(2000)461

Analysis of resveratrol in wine by capillary electrophoresis 881(2000)471

Contamination in food from packaging material 882(2000)255

Determination of synthetic phenolic antioxidants in food by high-performance liquid chromatography 882(2000)271

Polymer additive analysis by pyrolysis–gas chromatography. IV. Antioxidants 891(2000)325

Apigenin

Determination of apigenin in rat plasma by high-performance liquid chromatography 870(2000)463

Apolipoproteins

Capillary electrophoretic analysis of recombinant human apolipoprotein E. Calibration mode of a protein reference material 853(1999)237

Arabinoxylans

Determination of reducing end sugar residues in oligo- and polysaccharides by gas–liquid chromatography 866(2000)97

Arginine

Analysis of β -*N*-oxalyl-L- α , β -diaminopropionic acid and homoarginine in *Lathyrus sativus* by capillary zone electrophoresis 857(1999)295

Argininealkylamides

Qualitative and quantitative analysis of new alkyl amide arginine surfactants by high-performance liquid chromatography and capillary electrophoresis 852(1999)499

Aroclor 1254

Application of solid-phase microextraction–gas chromatography–mass spectrometry to characterize intermediates in a joint solar–microbial process for total mineralization of Aroclor 1254 873(2000)53

Aroma compounds

Semipreparative reversed-phase liquid chromatographic fractionation of aroma extracts from wine and other alcoholic beverages 864(1999)77

Use of solid-phase microextraction for measuring oil–water partition coefficients and correlation with high-performance liquid chromatographic methods for lipophilicity 864(1999)183

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Erratum to: “Trace-level determination of polar flavour compounds in butter by solid-phase extraction and gas chromatography–mass spectrometry” [J. Chromatogr. A, 844 (1999) 295–305] 873(2000)299

Sample preparation for the analysis of flavors and off-flavors in foods 880(2000)3

Applications of solid-phase microextraction in food analysis 880(2000)35

Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxypyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93

Analysis of organic sulfur compounds in wine aroma 881(2000)569

Analysis of low-volatility organic sulphur compounds in wines by solid-phase microextraction and gas chromatography 881(2000)583

Characterisation of whiskeys using solid-phase microextraction with gas chromatography–mass spectrometry 896(2000)351

Characterization of odorant compounds of mussels (*Mytilus edulis*) according to their origin using gas chromatography–olfactometry and gas chromatography–mass spectrometry 896(2000)361

Arsenic

Understanding of peak deterioration in hyphenated speciation systems due to gas–liquid separation in the hydride generation interface 855(1999)291

Arsenic compounds

Speciation of dimethylarsinic acid and monomethylarsonic acid by solid-phase microextraction–gas chromatography–ion trap mass spectrometry 873(2000)129

Determination of inorganic and organic anionic arsenic species in water by ion chromatography coupled to hydride generation–inductively coupled plasma atomic emission spectrometry 884(2000)211

Evaluation of extraction procedures for the ion chromatographic determination of arsenic species in plant materials 889(2000)83

Artificial sweeteners

Liquid chromatographic determination of aniline in table-top sweeteners based on pre-column derivatization with 1,2-naphthoquinone-4-sulfonate 859(1999)227

Development of a capillary electrophoresis method for the simultaneous analysis of artificial sweeteners, preservatives and colours in soft drinks 876(2000)213

Electrophoretic methods in the analysis of beverages 880(2000)243

New configurations and applications of molecularly imprinted polymers 889(2000)15

Aryloxyphenoxypropanoic acid

Stability-indicating high-performance liquid chromatographic assay and analysis of decomposition products of 2-[4-[(7-chloro-2-quinoxalinyloxy]phenoxy]propanoic acid 870(2000)179

Ascomycin

Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69

Purification of an ascomycin derivative with simulated moving bed chromatography. A case study 874(2000)155

Ascorbic acid

see also Vitamins

Indirect detection of organic acids in non-aqueous capillary electrophoresis 853(1999)171

Direct measurement of ascorbic acid biosynthesis in *Arabidopsis* cell suspension culture using capillary electrophoresis 853(1999)381

Simultaneous measurement of volatile sulfur compounds using ascorbic acid for oxidant removal and gas chromatography–flame photometric detection 864(1999)111

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235

Dehydroascorbic acid 881(2000)299

Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables 881(2000)309

Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317

Use of nucleic acids in the mobile phase for the determination of ascorbic acid in foods by high-performance liquid chromatography with electrochemical detection 881(2000)327

Aspartame

New configurations and applications of molecularly imprinted polymers 889(2000)15

Aspartic acid

Analysis of the enzymatic racemization of D-aspartic acid to L-aspartic acid by the on-line coupling of a solid-phase extraction column and a ligand-exchange high-performance liquid chromatography column 896(2000)299

Astaxanthin

Reproducibility of the separation of astaxanthin stereoisomers on Pirkle covalent L-leucine and D-phenylglycine columns 855(1999)157

Atranorin

Identification of atranorin and related potential allergens in oakmoss absolute by high-performance liquid chromatography–tandem mass spectrometry using negative ion atmospheric pressure chemical ionization 888(2000)103

Atrazines

Tandem solid-phase extraction of atrazine ozonation products in water 866(2000)195

Determination of interfering triazine degradation products by gas chromatography–ion trap mass spectrometry 868(2000)115

Graphitized carbons for solid-phase extraction 885(2000)73

Atropine

Enantioseparation of atropine by capillary electrophoresis using sulfated β -cyclodextrin: application to a plant extract 868(2000)285

Aucubin

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Avermectins

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Avidin

Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151

Azaarenes

Determination of basic nitrogen-containing polynuclear aromatic hydrocarbons formed during thermal degradation of polymers by high-performance liquid chromatography–fluorescence detection 878(2000)171

Azadirachtin

Rapid and sensitive analysis of azadirachtin and related triterpenoids from Neem (*Azadirachta indica*) by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 886(2000)89

Azaspiracid

Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13

Azinphos-methyl

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Azirdine

Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487

Azo dyes

Determination of sulphonated dyes in water by ion-interaction high-performance liquid chromatography 871(2000)227

Determination of sulfonated azo dyes in river water samples by capillary zone electrophoresis 883(2000)277

Azure B

Investigation on the iodination reaction of methylene blue by liquid chromatography–mass spectrometry with ionspray ionisation 854(1999)57

B**Baclofen**

Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233

Barbiturates

Micellar liquid chromatography for prediction of drug transport 870(2000)1

Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39

Barium

Comparative study of divalent metals and amines as silanol-blocking agents in reversed-phase liquid chromatography 855(1999)121

Basic compounds

Capillary electrochromatography of basic compounds using octadecyl-silica stationary phases with an amine-containing mobile phase 872(2000)315

New stationary phase for liquid chromatography with chemically bonded pinane ligand: synthesis and characterization by nuclear magnetic resonance and high-performance liquid chromatographic investigations 883(2000)27

Separation of basic solutes by reversed-phase capillary electrochromatography 887(2000)209

Basic drugs

see also Drugs

Chiral resolution of basic drugs by capillary electrophoresis with new glycosaminoglycans 864(1999)163

Enantiomer separations by nonaqueous capillary electrophoresis using octakis(2,3-diacetyl-6-sulfato)- γ -cyclodextrin 892(2000)499

Optimisation and characterisation of silica-based reversed-phase liquid chromatographic systems for the analysis of basic pharmaceuticals 897(2000)1

Beauvericin

Determination of the *Fusarium* mycotoxins, fusaproliferin and beauvericin by high-performance liquid chromatography–electrospray ionization mass spectrometry 858(1999)175

Bentazone

Semi-permeable surface analytical reversed-phase column for the improved trace analysis of acidic pesticides in water with coupled-column reversed-phase liquid chromatography with UV detection. Determination of bromoxynil and bentazone in surface water 858(1999)45

Benzalkonium chloride

Determination of cationic surfactants by capillary zone electrophoresis and micellar electrokinetic chromatography with deoxycholate micelles in the presence of large organic solvent concentrations 895(2000)227

Benzenediamines

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Benzenediols

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Benzenes

Capillary electrochromatography with a silica column with a dynamically modified cationic surfactant 855(1999)137

Trapping efficiency of aqueous pollutants in multichannel thick-film silicone-rubber traps for capillary gas chromatography 863(1999)57

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Development of membrane extraction with a sorbent interface–micro gas chromatography system for field analysis 873(2000)13

Evaluation of time-of-flight mass spectrometric detection for fast gas chromatography 878(2000)205

Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121

Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

Benzenesulfonates

Stability of sulfonated derivatives of benzene and naphthalene on disposable solid-phase extraction pre-columns and in an aqueous matrix 889(2000)231

Benzenesulfonic acids

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

Benzidine

Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291

Benzilic acid

Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169

Benzimidazoles

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Benzocaine

Liquid chromatographic determination of benzocaine and N-acetylbenzocaine in the edible fillet tissue from rainbow trout 855(1999)255

Benzodiazepines

On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279

Coupling device for desorption of drugs from solid-phase extraction-pipette tips and on-line gas chromatographic analysis 858(1999)117

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

Development of a simple in-vial liquid-phase microextraction device for drug analysis compatible with capillary gas chromatography, capillary electrophoresis and high-performance liquid chromatography 873(2000)3

Comparison of separation behavior of benzodiazepines in packed capillary electrochromatography and open-tubular capillary electrochromatography 887(2000)479

Use of chiral liquid chromatography–tandem mass spectrometry to investigate the metabolism of racemic cholecystokinin-B antagonists 896(2000)217

Selective and sensitive assay for the determination of benzodiazepines by high-performance liquid chromatography with simultaneous ultraviolet and reductive electrochemical detection at the hanging mercury drop electrode 897(2000)215

Benzoic acids

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

Application of solid-phase microextraction–gas chromatography–mass spectrometry to characterize intermediates in a joint solar–microbial process for total mineralization of Aroclor 1254 873(2000)53

Split-flow injector for capillary zone electrophoresis 883(2000)223

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Benzoin

Enantiomeric separation by capillary electrophoresis with an electroosmotic flow-controlled capillary 875(2000)315

Benzophenanthridine

Separation, identification and determination of sanguinarine in argemone and other adulterated edible oils by reversed-phase high-performance liquid chromatography 863(1999)123

Benzo[a]pyrene

Separation and identification of polar degradation products of benzo[a]pyrene with ozone by atmospheric pressure chemical ionization–mass spectrometry after optimized column chromatographic clean-up 855(1999)501

Benzyl alcohol

Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudostationary phase. Influence of methanol and acetonitrile on separation selectivity 853(1999)121

Benzylamine

Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277

Benzylpenicillin

Application of ion-exchange cartridge clean-up in food analysis II. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in meat using liquid chromatography with ultraviolet detection 855(1999)247

Berberine

Analysis of coptisine, berberine and palmatine in adulterated Chinese medicine by capillary electrophoresis–electrospray ion trap mass spectrometry 866(2000)273

Beryllium

Determination of beryllium in a stream sediment by high-performance chelation ion chromatography 876(2000)127

 β -Agonists

β_2 -Agonist extraction procedures for chromatographic analysis 880(2000)69

 β -Blockers

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

Rapid development of the enantiomeric separation of β -blockers by capillary electrophoresis using an experimental design approach 855(1999)681

Discrimination between enantioselective and non-selective binding sites on cellobiohydrolase-based stationary phases by site specific competing ligands 857(1999)107

Optimized stationary phases for the high-performance liquid chromatography–electrospray ionization mass spectrometric analysis of basic pharmaceuticals 869(2000)159

Micellar liquid chromatography for prediction of drug transport 870(2000)1

Application of ligand-exchange capillary electrophoresis to the chiral separation of α -hydroxy acids and β -blockers 875(2000)307

Optimisation of procedures for the extraction of structural analogues of propranolol with molecular imprinted polymers for sample preparation 889(2000)143

Betaine

Determination of betaine in *Lycium chinense* fruits by liquid chromatography–electrospray ionization mass spectrometry 857(1999)331

Design of a simulated moving bed unit for sucrose–betaine separations 872(2000)23

Bile acids

Retention and separation studies of cholesterol and bile acids using thermostated thin-layer chromatography 857(1999)255

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Electron–capture mass spectrometry: recent advances 892(2000)329

Bile salts

Determination of cationic surfactants by capillary zone electrophoresis and micellar electrokinetic chromatography with deoxycholate micelles in the presence of large organic solvent concentrations 895(2000)227

Binaphthol

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Binaphthyldiamine

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Biochanin A

Quantitative determination of isoflavones and coumestrol in soybean by column liquid chromatography 886(2000)55

Biogenic amides

Erratum to: “Determination of fatty acid amides as trimethylsilyl derivatives by gas chromatography with mass spectrometric detection” [J. Chromatogr. 849 (1999) 541–552] 859(1999)119

Biogenic amines

see also Amines and Catecholamines

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75

Liquid chromatographic determination of biogenic amines in fermented foods after derivatization with 3,5-dinitrobenzoyl chloride 881(2000)517

Biomimetic dyes

Biomimetic dyes as affinity chromatography tools in enzyme purification (Review) 891(2000)33

Biopterin

Reversed-phase ion-pair chromatography coupled to electrospray ionisation mass spectrometry by on-line removal of the counter-ions 854(1999)155

Biotin

Enzyme amplification as detection tool in continuous-flow systems I. Development of an enzyme-amplified biochemical detection system coupled on-line to flow-injection analysis 855(1999)383

Enzyme amplification as detection tool in continuous-flow systems II. On-line coupling of liquid chromatography to enzyme-amplified biochemical detection after pre-column derivatization with biotin 855(1999)397

Analytical techniques for determining biotin 881(2000)331

Biphenyl

Application of solid-phase microextraction–gas chromatography–mass spectrometry to characterize intermediates in a joint solar–microbial process for total mineralization of Aroclor 1254 873(2000)53

Bisbenzylisoquinoline alkaloids

Determination of bisbenzylisoquinoline alkaloids by high-performance liquid chromatography (II) 891(2000)189

Bis(2-ethylhexyl)sodium sulfosuccinate

Micellar electrokinetic chromatography with bis-(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Bisethylisooctanolactones

Unknown bisethylisooctanolactone isomers in industrial waste water. Isolation, identification and occurrence in surface water 889(2000)245

Bis(hydroxybenzyl)ethylenediaminediacetic acid

Potential of ethylenediaminedi(*o*-hydroxyphenylacetic acid) and *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid for the determination of metal ions by capillary electrophoresis 890(2000)363

Separation selectivity of anionic metal complexes of *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid in ion and ion electrokinetic chromatography 895(2000)255

Bisphenol A

Quantification of the xenoestrogens 4-*tert*.-octylphenol and bisphenol A in water and in fish tissue based on microwave assisted extraction, solid-phase extraction and liquid chromatography–mass spectrometry 864(1999)17

Preparation of 4,4'-(1-[²H₆]methylene)bis-[2,3,5,6-²H₄]-phenol and its application to the measurement of bisphenol A in beverages by stable isotope dilution mass spectrometry 883(2000)163

High-performance liquid chromatography with multi-wavelength detection of the bisphenol A impurities 883(2000)291

Qualitative and quantitative analysis of a thermoset polymer, poly(benzoxazine), by pyrolysis–gas chromatography 886(2000)217

Separation of bisphenol A and three alkylphenols by micellar electrokinetic chromatography 895(2000)213

Bisphenol diglycidyl ethers

Use of solid-phase microextraction for the analysis of bisphenol A and bisphenol A diglycidyl ether in food simulants 864(1999)137

Bisphosphonates

Chromatographic analysis of bisphosphonates 868(2000)141

Analysis of bisphosphonates by capillary electrophoresis–electrospray ionization mass spectrometry 872(2000)289

Development and validation of a capillary zone electrophoretic method for the determination of bisphosphonate and phosphonate impurities in clodronate 893(2000)411

Borate

Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285

Bromate

Determination of bromide, bromate and other anions with ion chromatography and an inductively coupled plasma mass spectrometer as element-specific detector 862(1999)39

Ion chromatography determination of trace level bromate by large volume injection with conductivity and spectrophotometric detection after post column derivatisation 864(1999)263

Eliminating the chlorite interference in US Environmental Protection Agency Method 317.0 permits analysis of trace bromate levels in all drinking water matrices 882(2000)309

Comparison of on-line coupling of ion-chromatography with atmospheric pressure ionization mass spectrometry and with inductively coupled plasma mass spectrometry as tools for the ultra-trace analysis of bromate in surface water samples 884(2000)191

Performance evaluation of a method for the determination of bromate in drinking water by ion chromatography (EPA Method 317.0) and validation of EPA Method 324.0 884(2000)201

Simultaneous determination of inorganic disinfection by-products and the seven standard anions by ion chromatography 890(2000)295

Bromide

Determination of bromide, bromate and other anions with ion chromatography and an inductively coupled plasma mass spectrometer as element-specific detector 862(1999)39

Bromophenacyl trifluoromethanesulfonate

Analysis of carnitine and acylcarnitines in urine by capillary electrophoresis 895(2000)309

Bromopropanediols

Determination of halogenated mono-alcohols and diols in water by gas chromatography with electron-capture detection 866(2000)65

Bromopyridylazodiethylaminophenol

Use of ion chromatography for the determination of heavy and transition metals in biochemical samples 857(1999)343

Bromoxynil

Semi-permeable surface analytical reversed-phase column for the improved trace analysis of acidic pesticides in water with coupled-column reversed-phase liquid chromatography with UV detection. Determination of bromoxynil and bentazone in surface water 858(1999)45

Brompheniramine

Separation of brompheniramine enantiomers by capillary electrophoresis and study of chiral recognition mechanisms of cyclodextrins using NMR-spectroscopy, UV spectrometry, electrospray ionization mass spectrometry and X-ray crystallography 875(2000)471

Bupivacaine

Highly sensitive bioassay of lidocaine in human plasma by high-performance liquid chromatography–tandem mass spectrometry 854(1999)3

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Buspirone

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Butanedione monoxime

Identification of 2,3-butanedione monoxime hydrogenation products by gas chromatography–mass spectrometry in an ion trap mass spectrometer 855(1999)341

Butanol

Comparison of the performance of butanol and pentanol as modifiers in the micellar chromatographic determination of some phenethylamines 866(2000)35

Butylaminopropanediol

Determination of the enantiomers of 3-*tert.*-butylamino-1,2-propanediol by high-performance liquid chromatography coupled to evaporative light scattering detection 890(2000)239

Determination of the enantiomers of 3-*tert.*-butylamino-1,2-propanediol by high-performance liquid chromatography using mass spectrometric detection 896(2000)201

Butylcarbamoylquinine

tert.-Butylcarbamoylquinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique 892(2000)509

Butylmethyldihydrofuranones

Determination of oak lactones in barrel-aged wines and in oak extracts by stable isotope dilution analysis 857(1999)239

Butylphenol

Qualitative and quantitative analysis of a thermoset polymer, poly(benzoxazine), by pyrolysis–gas chromatography 886(2000)217

3-Butyn-2-one

Characterization of a chemical artifact in the liquid chromatographic determination of 3-butyn-2-one using the 2,4-dinitrophenylhydrazine method 886(2000)303

C**Cadmium**

Determination of trace cadmium in environmental water samples using ion-interaction reversed-phase liquid chromatography with fluorescence detection 877(2000)123

Caffeic acid

Determination of phenolic acids in fruit juices by isocratic column liquid chromatography 891(2000)183

Caffeine

Determination of theophylline and its metabolites in biological samples by liquid chromatography–mass spectrometry 870(2000)87

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175

Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235

Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425

Calcium

Comparative study of divalent metals and amines as silanol-blocking agents in reversed-phase liquid chromatography 855(1999)121

Calcium acamprostate

Determination of homotaurine as impurity in calcium acamprostate by capillary zone electrophoresis 853(1999)421

Camphor

Comparative study on camphor enantiomers behavior under the conditions of gas–liquid chromatography and reversed-phase high-performance liquid chromatography systems modified with α - and β -cyclodextrins 874(2000)73

Camptothecin

Measurement and pharmacokinetics of unbound 20(S)-camptothecin in rat blood and brain by microdialysis coupled to microbore liquid chromatography with fluorescence detection 870(2000)221

Capillarisin

Fractionation and analysis of *Artemisia capillaris* Thunb. by affinity chromatography with human serum albumin as stationary phase 870(2000)501

Carbamates

Capillary electrophoretic determination of ferric dimethyldithiocarbamate as iron(III) chelate of EDTA 857(1999)365

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Automated in-tube solid-phase microextraction–high-performance liquid chromatography for carbamate pesticide analysis 873(2000)137

Carbendazim

Determination of thiophanate-methyl and its metabolites at trace level in spiked natural water using the supported liquid membrane extraction and the microporous membrane liquid–liquid extraction techniques combined on-line with high-performance liquid chromatography 893(2000)123

CarbiDOPA

Enantioseparation of 3,4-dihydroxyphenylalanine and 2-hydrazino-2-methyl-3-(3,4-dihydroxyphenyl)propanoic acid by capillary electrophoresis using cyclodextrins 875(2000)379

Carbohydrates

see also Cyclodextrins, Disaccharides, Monosaccharides, Oligosaccharides, Polysaccharides, Saccharides and Sugars

Isolation and characterization of cyclic α -(1 \rightarrow 4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

3-Aminobenzamide and 3-aminobenzoic acid, tags for capillary electrophoresis of complex carbohydrates with laser-induced fluorescent detection 863(1999)205

Determination of reducing end sugar residues in oligo- and polysaccharides by gas–liquid chromatography 866(2000)97

Preparation of various silica-based columns for capillary electrochromatography by in-column derivatization 873(2000)247

Screening for limonoid glucosides in *Citrus tangerina* (Tanaka) Tseng by high-performance liquid chromatography–electrospray ionization mass spectrometry 874(2000)13

Electrophoretic methods in the analysis of beverages 880(2000)243

Changes in free and conjugated polyamines during starvation of sugarcane juices as analyzed by high-performance liquid chromatography 881(2000)531

Determination of inulin in meat products by high-performance liquid chromatography with refractive index detection 881(2000)591

Specific applications of capillary electrochromatography to biopolymers, including proteins, nucleic acids, peptide mapping, antibodies, and so forth (Review) 887(2000)137

Determination of glucosinolates in rapeseed and *Thlaspi caerulescens* plants by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 889(2000)75

Analysis of carbohydrates in wood and pulps employing enzymatic hydrolysis and subsequent capillary zone electrophoresis 891(2000)157

Determination of cyclodextrin content using periodate oxidation by capillary electrophoresis 891(2000)201

Direct determination of amino acids and carbohydrates by high-performance capillary electrophoresis with refractometric detection 894(2000)253

Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285

Capillary electrophoresis and off-line capillary electrophoresis–electrospray ionization quadrupole time-of-flight tandem mass spectrometry of carbohydrates 895(2000)291

Carbolines

Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Analysis of the bioactive alkaloids tetrahydro- β -carboline and β -carboline in food 881(2000)483

Carbomethoxy(4-fluorophenyl)iodoallylnortropine

Analysis of 2 β -carbomethoxy-3 β -(4-fluorophenyl)-*N*-(3-iodo-*E*-allyl)nortropine in rat plasma. I. Method development and validation by capillary electrophoresis 895(2000)87

Carbon dioxide

Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383

Study of dead volume measurement in packed subcritical fluid chromatography with ODS columns and carbon dioxide–modifier mobile phases 866(2000)241

Carbonic anhydrase

Multiple-step ligand injection affinity capillary electrophoresis for determining binding constants of ligands to receptors 897(2000)339

Carbon monoxide

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum–rhodium alloy catalysts 852(1999)507

Carbonyl compounds

see also Aldehydes, Ketones and Volatile organic compounds

Separation of carbonyl 2,4-dinitrophenylhydrazones by capillary electrochromatography with diode array detection 853(1999)487

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Dynamic system for the calibration of semi-volatile carbonyl compounds in air 883(2000)171

Solid-phase microextraction with on-fibre derivatisation applied to the analysis of volatile carbonyl compounds 886(2000)175

Microwave-assisted derivatization of volatile carbonyl compounds with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine 891(2000)267

Carboxamides

Erratum to: "Determination of fatty acid amides as trimethylsilyl derivatives by gas chromatography with mass spectrometric detection" [J. Chromatogr. 849 (1999) 541–552] 859(1999)119

Carboxyalkylphenol ethoxycarboxylates

Analysis of nonylphenol polyethoxycarboxylates and their related metabolites by on-line derivatization and ion-trap gas chromatography–mass spectrometry 862(1999)113

Carboxycyclopropylglycine

Isolation of pure (2*S*,1'*S*,2'*S*)-2-(2'-carboxycyclopropyl)glycine from *Blighia sapida* (Akee) 873(2000)283

2-Carboxyibuprofen

Uniform-sized molecularly imprinted polymer for (*S*)-ibuprofen. Retention properties in aqueous mobile phases 857(1999)117

Carboxylic acids

see also Fatty acids, Hydroxy acids and Organic acids

Equilibrium analysis of reactions between aromatic anions and nonionic surfactant micelles by capillary zone electrophoresis 853(1999)55

Non-suppressed conductivity and indirect UV detection of carboxylic acids in environmental samples by ion-exclusion chromatography using 2,6-pyridinedicarboxylic acidic eluent 859(1999)173

Effect of stationary phase hydrophobicity and mobile phase composition on the separation of carboxylic acids in ion chromatography 867(2000)131

Comparative methodology in the determination of α -oxocarboxylates in aqueous solution. Ion chromatography versus gas chromatography after oximation, extraction and esterification 867(2000)143

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97

Herbicide and plant growth regulator analysis by capillary electrophoresis 880(2000)281

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Influence of organic solvents in the mobile phase on the determination of carboxylic acids and inorganic anions in grape juice by ion chromatography 881(2000)387

Tetrahydro- β -carboline-3-carboxylic acids and contaminants of L-tryptophan 881(2000)501

Novel way of separating polyfluorocarboxylic acids by ion-exclusion chromatography 884(2000)93

Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99

Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211

Role of chromatography in the analysis of sugars, carboxylic acids and amino acids in food (Review) 891(2000)1

Application of gas chromatography coupled to chemical ionisation mass spectrometry following headspace solid-phase microextraction for the determination of free volatile fatty acids in aqueous samples 891(2000)287

Determination of 1-aminocyclopropane-1-carboxylic acid and its structural analogue by liquid chromatography and ion spray tandem mass spectrometry 896(2000)335

Carboxylic acids, perfluorinated

Determination of 20 underivatized proteinic amino acids by ion-pairing chromatography and pneumatically assisted electrospray mass spectrometry 855(1999)191

Carboxyphenyldiethylaminobenzofuran

2-(4-Carboxyphenyl)-6-*N,N*-diethylaminobenzofuran: a useful reagent for the sensitive determination of alcohols by high-performance liquid chromatography with fluorimetric detection 869(2000)243

Carnitine

Direct chromatographic resolution of carnitine and *O*-acylcarnitine enantiomers on a teicoplanin-bonded chiral stationary phase 857(1999)145

Analysis of carnitine and acylcarnitines in urine by capillary electrophoresis 895(2000)309

Carotenes

see also Vitamins

Liquid chromatography–electrospray mass spectrometry of β -carotene and xanthophylls. Validation of the analytical method 854(1999)233

Improved liquid chromatographic method for the analysis of photosynthetic pigments of higher plants 876(2000)111

C₃₀ Stationary phases for the analysis of food by liquid chromatography 880(2000)189

Simultaneous determination of α -tocopherol and β -carotene in olive oil by reversed-phase high-performance liquid chromatography 881(2000)255

Fatty acids and β -carotene in Australian purslane (*Portulaca oleracea*) varieties 893(2000)207

Carotenoids

Separation and identification of various carotenoids by C₃₀ reversed-phase high-performance liquid chromatography coupled to UV and atmospheric pressure chemical ionization mass spectrometric detection 854(1999)37

Liquid chromatography–electrospray mass spectrometry of β -carotene and xanthophylls. Validation of the analytical method 854(1999)233

C₃₀ Stationary phases for the analysis of food by liquid chromatography 880(2000)189

Chromatographic determination of carotenoids in foods 881(2000)543

Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annum* L.) powders 889(2000)41

Caseinomacropeptide

Simultaneous separation and quantitation of the major bovine whey proteins including proteose peptone and caseinomacropeptide by reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene 878(2000)183

Caseins

Analysis of recombinant and modified proteins by capillary zone electrophoresis coupled with electrospray ionization tandem mass spectrometry 855(1999)695

Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371

Evaluation of the effect of rennet type on casein proteolysis in an ovine milk cheese by means of capillary electrophoresis 881(2000)59

Catalpol

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Catechins

Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235

Electrophoretic methods in the analysis of beverages 880(2000)243

Determination of tea catechins 881(2000)411

Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425

Comparative analysis of tea catechins and theaflavins by high-performance liquid chromatography and capillary electrophoresis 881(2000)439

Supercritical fluid extraction of catechins from *Cratogeomys pruifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

Catecholamines

see also Amines and Biogenic amines

Study of catecholamines in patient urine samples by capillary electrophoresis 853(1999)527

Determination of catecholamines in pheochromocytoma cell (PC-12) culture medium by microdialysis–microbore liquid chromatography 870(2000)405

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Miniaturized thin-layer radial flow cell with interdigitated ring-shaped microarray electrode used as amperometric detector for capillary electrophoresis 891(2000)149

Determination of urinary catecholamines with capillary electrophoresis after solid-phase extraction 895(2000)317

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

Cellobiohydrolase

Discrimination between enantioselective and non-selective binding sites on cellobiohydrolase-based stationary phases by site specific competing ligands 857(1999)107

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Cellobiose

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Cellulases

Studies on the chromatographic fractionation of *Trichoderma reesei* cellulases by hydrophobic interaction 865(1999)123

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Cephalosporins

Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants 868(2000)313

Reply to 'Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants' 868(2000)317

Characterization of cephalosporin transfer between aqueous and colloidal phases by micellar electrokinetic chromatography 871(2000)439

Characterization of partition and thermodynamic properties of cephalosporins using micellar electrokinetic chromatography in glycodeoxycholic acid solution 873(2000)237

Optimization of separation and migration behavior of cephalosporins in capillary zone electrophoresis 879(2000)197

Determination of cephalosporins in raw bovine milk by high-performance liquid chromatography 882(2000)145

Adaptation of capillary electrophoresis to the determination of selected cephalosporins for injection 895(2000)27

Ceramides

Postcolumn fluorescence as an alternative to evaporative light scattering detection for ceramide analysis with gradient elution in non-aqueous reversed-phase liquid chromatography 859(1999)99

Ceramide analysis utilizing gas chromatography–mass spectrometry 876(2000)229

Retention behaviour of ceramides in sub-critical fluid chromatography in comparison with non-aqueous reversed-phase liquid chromatography 883(2000)211

Cetrimide

Alkyltrimethylammonium surfactant-mediated extractions: Characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Cetylpyridinium chloride

Determination of cationic surfactants by capillary zone electrophoresis and micellar electrokinetic chromatography with deoxycholate micelles in the presence of large organic solvent concentrations 895(2000)227

Cetyltrimethylammonium coating

Determination of inorganic anions by ion chromatography using a graphitized carbon column dynamically coated with cetyltrimethylammonium ions 855(1999)261

Chelerythrine

Capillary electrophoretic determination of sanguinarine and chelerythrine in plant extracts and pharmaceutical preparations 866(2000)293

Chelidonium

Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69

Chlorate

Simultaneous determination of inorganic disinfection by-products and the seven standard anions by ion chromatography 890(2000)295

Chloride

Determination of chloride, sulfate and nitrate in groundwater samples by ion chromatography 884(2000)185

Chlorine trifluoride

Gas chromatographic analysis of trace impurities in chlorine trifluoride 873(2000)229

Chlorite

Eliminating the chlorite interference in US Environmental Protection Agency Method 317.0 permits analysis of trace bromate levels in all drinking water matrices 882(2000)309

Simultaneous determination of inorganic disinfection by-products and the seven standard anions by ion chromatography 890(2000)295

Chlormequat

Determination of the plant growth regulator chlormequat in food by liquid chromatography–electrospray ionisation tandem mass spectrometry 878(2000)77

Improved sensitivity in detection of chlormequat by liquid chromatography–mass spectrometry 897(2000)399

Chloroanilines

Optimization of the separation of mono- and dichloroanilines in ion interaction high-performance liquid chromatography 863(1999)1

On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175

Chlorobenzenes

Trace analysis of ten chlorinated benzenes in water by headspace solid-phase microextraction 874(2000)149

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Chlorodimethylaziridine

Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487

Chlorogenic acid

Determination of chlorogenic acid in rat blood by microdialysis coupled with microbore liquid chromatography and its application to pharmacokinetic studies 870(2000)443

Determination of phenolic acids in fruit juices by isocratic column liquid chromatography 891(2000)183

4-Chloro-3-methylphenol

Trace determination of antifouling compounds by on-line solid-phase extraction–gas chromatography–mass spectrometry 885(2000)361

6-Chloronicotinic acid

Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497

Chlorophacinone

Determination of chlorophacinone and diphacinone in commercial rodenticides by liquid chromatography–UV detection and liquid chromatography–electrospray ionization mass spectrometry 891(2000)249

Chlorophenols

Nonaqueous capillary electrophoresis equipped with amperometric detection for analysis of chlorinated phenolic compounds 864(1999)323

On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175

New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Highly selective and efficient determination of US Environmental Protection Agency priority phenols employing solid-phase extraction and non-aqueous capillary electrophoresis 896(2000)95

Capillary electrophoresis–electrospray ion-trap mass spectrometry for the separation of chlorophenols 896(2000)125

Comparison of different sorbent materials for *on-line* solid-phase extraction with liquid chromatography–atmospheric pressure chemical ionization mass spectrometry of phenols 896(2000)159

Cloud point preconcentration of rather polar compounds: application to the high-performance liquid chromatographic determination of priority pollutant chlorophenols 897(2000)171

Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279

Chlorophenoxy acids

Determination of chlorophenoxy acid herbicides in water by *in situ* esterification followed by *in-vial* liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111

3-Chloro-1-phenyl-1-propanol

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Chlorophylls

High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483

Chloropropanediols

Determination of halogenated mono-alcohols and diols in water by gas chromatography with electron-capture detection 866(2000)65

Chloroquine

Separation of enantiomers on anion exchangers modified with heparin in liquid chromatography 864(1999)25

Chloroquinoxalinyloxyphenoxypropanoic acid

Stability-indicating high-performance liquid chromatographic assay and analysis of decomposition products of 2-{4-[(7-chloro-2-quinoxalinyloxy)phenoxy]propanoic acid 870(2000)179

Chlorothalonil

Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261

Determination of aqueous chlorothalonil with solid-phase microextraction and gas chromatography 896(2000)105

Chlorotheophylline

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Chlorotoluidine

Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291

Chlorotriazines

Capillary zone electrophoretic separation of neutral species of chloro-*s*-triazines in the presence of cationic surfactant monomers 878(2000)137

Identification of reactive dyes in spent dyebaths and wastewater by capillary electrophoresis–mass spectrometry 886(2000)271

2-Chlorovinylarsonous acid

Simultaneous extraction and derivatization of 2-chlorovinylarsonous acid from soils using supercritical and pressurized fluids 888(2000)327

Chlorpheniramine

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Simultaneous determination of paracetamol and chlorpheniramine in human plasma by liquid chromatography–tandem mass spectrometry 870(2000)77

Chiral capillary electrophoresis as predictor for separation of drug enantiomers in continuous flow zone electrophoresis 895(2000)51

Chlorpyrifos

Supercritical fluid extraction of chlorpyrifos and 3,5,6-trichloro-2-pyridinol from garden compost 862(1999)105

Chlorsulfuron

Determination of sulfonyleurea degradation products in soil by liquid chromatography–ultraviolet detection followed by confirmatory liquid chromatography–tandem mass spectrometry 855(1999)575

Chlozolinate

Inclusion of the fungicide chlozolinate in a multiresidue method 858(1999)123

Cholecystokinins

Use of chiral liquid chromatography–tandem mass spectrometry to investigate the metabolism of racemic cholecystokinin-B antagonists 896(2000)217

Cholesterol

Retention and separation studies of cholesterol and bile acids using thermostated thin-layer chromatography 857(1999)255

Cholesterol oxides

Determination of cholesterol oxides in processed food using high-performance liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 896(2000)321

Chromate

Ion-chromatographic assay of chromium(VI)-containing semiconductor etchants. Statistical comparison of two columns 884(2000)151

Chromium

Interface for capillary electrophoresis coupled with inductively coupled plasma atomic emission spectrometry 853(1999)141

Quantitative analysis of manganese, chromium and molybdenum by ion-pair reversed-phase high-performance liquid chromatography with pre-column derivatization and UV-visible detection 857(1999)351

Investigation of Cr(III) hydrolytic polymerisation products by capillary electrophoresis–inductively coupled plasma-mass spectrometry 872(2000)227

Chrysanthemic acid

High-performance liquid chromatography study of the enantiomer separation of chrysanthemic acid and its analogous compounds on a terguride-based stationary phase 859(1999)133

Chymotrypsin

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

Cinchona carbamates

Novel cinchona alkaloid carbamate C₉-dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Cisplatin

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

Citrinin

Reversed-phase liquid chromatographic behavior of the mycotoxins, citrinin and ochratoxin A 862(1999)147

Clenbuterol

On-line coupling of solid-phase extraction with mass spectrometry for the analysis of biological samples. I. Determination of clenbuterol in urine 863(1999)115

Enantiomeric separation of clenbuterol by transient isotachopheresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachopheresis 871(2000)173

Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography 873(2000)293

Use of molecularly imprinted solid-phase extraction for the selective clean-up of clenbuterol from calf urine 889(2000)105

Clevidipine

Packed-column supercritical fluid chromatography for the purity analysis of clevidipine, a new dihydropyridine drug 862(1999)95

Clodronate

Analysis of bisphosphonates by capillary electrophoresis–electrospray ionization mass spectrometry 872(2000)289

Development and validation of a capillary zone electrophoretic method for the determination of bisphosphonate and phosphonate impurities in clodronate 893(2000)411

Clofibric acid

Determination of rice herbicides, their transformation products and clofibric acid using on-line solid-phase extraction followed by liquid chromatography with diode array and atmospheric pressure chemical ionization mass spectrometric detection 879(2000)3

Clonixin

Determination of clonixin in plasma and urine by reversed-phase high-performance liquid chromatography 889(2000)135

Clopidol

Determination of nicarbazin and clopidol in poultry feeds by liquid chromatography 882(2000)79

Determination of clopidol residues in chicken tissues by high-performance liquid chromatography–mass spectrometry 882(2000)85

Cloransulam-methyl

Extraction of cloransulam-methyl from soil with subcritical water and supercritical CO₂ 897(2000)405

Closantel

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Clotrimazole

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Cloxacillin

Application of ion-exchange cartridge clean-up in food analysis II. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in meat using liquid chromatography with ultraviolet detection 855(1999)247

Clozapine

Separation of antipsychotic drugs (clozapine, loxapine) and their metabolites by capillary zone electrophoresis 853(1999)461

Cobalt

Natural poly-histidine affinity tag for purification of recombinant proteins on cobalt(II)-carboxymethylaspartate crosslinked agarose 864(1999)247

Integration of a microextraction system. Solvent extraction of a Co–2-nitroso-5-dimethylaminophenol complex on a microchip 894(2000)19

Cocciidiostat

Determination of nicarbazin and clopidol in poultry feeds by liquid chromatography 882(2000)79

Codeine

Method for quantification of morphine and its 3- and 6-glucuronides, codeine, codeine glucuronide and 6-monoacetylmorphine in human blood by liquid chromatography–electrospray mass spectrometry for routine analysis in forensic toxicology 854(1999)109

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Analysis of codeine, dihydrocodeine and their glucuronides in human urine by electrokinetic capillary immunoassays and capillary electrophoresis–ion trap mass spectrometry 895(2000)133

Collagen

Binding of lead to collagen type I and V and $\alpha_2(I)$ CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

Separation of proteins and peptides by capillary electrophoresis in acid buffers containing high concentrations of surfactants 852(1999)325

Use of Edman degradation sequence analysis and matrix-assisted laser desorption/ionization mass spectrometry in designing substrates for matrix metalloproteinases 890(2000)117

Color additives

Rapid quantification of hexachlorobenzene in the color additives D&C Red Nos. 27 and 28 (phloxine B) using solid-phase microextraction and gas chromatography–mass spectrometry 863(1999)37

Congo red

Congophilicity (Congo red affinity) of different β_2 -microglobulin conformations characterized by dye affinity capillary electrophoresis 894(2000)319

Copper

Ion exchange on resins with temperature-responsive selectivity. IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57

Ion-chromatographic study of the possible absorption of copper and zinc by the skin of *Rana pipiens* 884(2000)143

Copper(II)-D-penicillamine

Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic β -substituted α -amino acids on a copper(II)-D-penicillamine chiral stationary phase 897(2000)145

Coptisine

Analysis of coptisine, berberine and palmatine in adulterated Chinese medicine by capillary electrophoresis–electrospray ion trap mass spectrometry 866(2000)273

Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69

Corticosteroids

Comparison of different liquid chromatography methods for the determination of corticosteroids in biological matrices 893(2000)55

Cotinine

Application of liquid separation techniques to the determination of the main urinary nicotine metabolites 870(2000)29

Coumarin-3-carbonyl chloride

Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473

Coumarins

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669

Solid phase microextraction of alkenylbenzenes and other flavor-related compounds from tobacco for analysis by selected ion monitoring gas chromatography–mass spectrometry 858(1999)79

Preparative isolation and purification of notopterin and isoimperatorin from *Notopterygium forbesii* boisii (Chinese traditional medicinal herb) by high-speed counter-current chromatography 883(2000)67

Separation of coumarins from *Archangelica officinalis* in high-performance liquid chromatography and thin-layer chromatography systems 886(2000)75

Coumestrol

Quantitative determination of isoflavones and coumestrol in soybean by column liquid chromatography 886(2000)55

Creatinine

On-line coupling of solid-phase extraction with mass spectrometry for the analysis of biological samples. I. Determination of clenbuterol in urine 863(1999)115

Crown ethers

Separation of enantiomers by capillary electrophoresis–mass spectrometry employing a partial filling technique with a chiral crown ether 875(2000)323

Enhancement of chiral recognition by formation of a sandwiched complex in capillary electrophoresis 875(2000)361

Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113

Crystal Violet

On-line isotachophoretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219

Cupferron

Separation and direct UV detection of lanthanides complexed with cupferron by capillary electrophoresis 895(2000)263

Cyanate

Ion chromatographic determination of cyanate in saline gold processing samples 855(1999)267

Cyanide

Enhanced conductometric detection of cyanide in suppressed ion chromatography 884(2000)75

Cyanogen bromide

Simultaneous determination of cyanogen chloride and cyanogen bromide in treated water at sub- $\mu\text{g/L}$ levels by a new solid-phase microextraction–gas chromatographic–electron-capture detection method 897(2000)307

Cyanogen chloride

Simultaneous determination of cyanogen chloride and cyanogen bromide in treated water at sub- $\mu\text{g/L}$ levels by a new solid-phase microextraction–gas chromatographic–electron-capture detection method 897(2000)307

Cyanohydrins

Determination of enantiomeric excess of α -hydroxy-3-phenoxybenzeneacetonitrile and its *n*-butyl ester by chiral high-performance liquid chromatography 893(2000)189

Cyclamate

Liquid chromatographic determination of aniline in table-top sweeteners based on pre-column derivatization with 1,2-naphthoquinone-4-sulfonate 859(1999)227

Cyclodepsipeptides

Liquid chromatography–electrospray time-of-flight mass spectrometry for on-line accurate mass determination and identification of cyclodepsipeptides in a crude extract of the fungus *Metarrhizium anisopliae* 872(2000)85

Cyclodextrins

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a β -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

Electrospray mass spectrometry and supercritical fluid chromatography of methylated β -cyclodextrins 855(1999)645

Nonaqueous capillary electrophoresis equipped with amperometric detection for analysis of chlorinated phenolic compounds 864(1999)323

Enantiomer separation of α -ionone using gas chromatography with cyclodextrin derivatives as chiral stationary phases 865(1999)201

Enantioselective determination by capillary electrophoresis with cyclodextrins as chiral selectors 875(2000)89

Enantioseparation of atropine by capillary electrophoresis using sulfated β -cyclodextrin: application to a plant extract 868(2000)285

Simultaneous stereoselective analysis of tramadol and its main phase I metabolites by on-line capillary zone electrophoresis–electrospray ionization mass spectrometry 868(2000)295

Analysis of *cis*–*trans* isomers and enantiomers of sertraline by cyclodextrin-modified micellar electrokinetic chromatography 871(2000)207

Investigation of the effects of cyclodextrins and organic solvents on the separation of cationic surfactants in capillary electrophoresis 872(2000)269

Enantiomeric separation of enzymatic hydrolysis products of dihydropyrimidinone methyl ester with cationic cyclodextrin by capillary electrophoresis 872(2000)279

Comparative study on camphor enantiomers behavior under the conditions of gas–liquid chromatography and reversed-phase high-performance liquid chromatography systems modified with α - and β -cyclodextrins 874(2000)73

Enantiomeric separations of drugs using mixtures of charged and neutral cyclodextrins 875(2000)123

Separation of drug enantiomers by capillary electrophoresis in the presence of neutral cyclodextrins 875(2000)135

Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated β -cyclodextrin 875(2000)389

Fast chiral separations using sulfated β -cyclodextrin and short-end injection in capillary electrophoresis 875(2000)403

Optimization of the cyclodextrin-assisted capillary electrophoresis separation of the enantiomers of phenoxyacid herbicides 875(2000)423

Mechanistic study of opposite migration order of dimethindene enantiomers in capillary electrophoresis in the presence of native β -cyclodextrin and heptakis(2,3,6-tri-*O*-methyl)- β -cyclodextrin 875(2000)455

Separation of brompheniramine enantiomers by capillary electrophoresis and study of chiral recognition mechanisms of cyclodextrins using NMR-spectroscopy, UV spectrometry, electrospray ionization mass spectrometry and X-ray crystallography 875(2000)471

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Analysis of polymyxin B sulfate by capillary zone electrophoresis with cyclodextrin as additive. Method development and validation 879(2000)211

Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs 883(2000)249

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

Simultaneous determination of eleven ingredients in ophthalmic solutions by cyclodextrin-modified micellar electrokinetic chromatography with tetrabutylammonium salt 888(2000)299

Determination of cyclodextrin content using periodate oxidation by capillary electrophoresis 891(2000)201

Enantiomer separations by nonaqueous capillary electrophoresis using octakis(2,3-diacetyl-6-sulfato)- γ -cyclodextrin 892(2000)499

Allylamine- β -cyclodextrin copolymer. A novel chiral selector for capillary electrophoresis 894(2000)95

Methyl- β -cyclodextrin modified micellar electrokinetic capillary chromatography with laser-induced fluorescence for separation and detection of phospholipids 894(2000)129

Influence of the structure of cyclodextrins and amino acid sequence of dipeptides and tripeptides on the pH-dependent reversal of the migration order in capillary electrophoresis 894(2000)267

The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry 894(2000)345

Capillary electrophoretic separation of the enantiomers of organophosphates with a phosphorus stereogenic center using the sodium salt of octakis(2,3-diacetyl-6-sulfo)- γ -cyclodextrin as resolving agent 895(2000)247

Highly reproducible capillary zone electrophoresis of humic acids in cyclodextrin- or oligosaccharide-modified background electrolytes 895(2000)345

Characterization of sulfobutyl ether- β -cyclodextrins mixtures by anion-exchange chromatography using evaporative light scattering detection 897(2000)185

Cyclohexane

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

Cyclopentane

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

Cyclophane

Separation of aromatic isomers on cyclophane-bonded stationary phases 877(2000)61

Cyclophilin

Interaction of cyclophilin and cyclosporins monitored by affinity capillary electrophoresis 853(1999)469

Cyclopropaneamino acids

Isolation of pure (2*S*,1'*S*,2'*S*)-2-(2'-carboxycyclopropyl)glycine from *Blighia sapida* (Akee) 873(2000)283

Cyclopropanedicarboxylic acid

Determination of 1-aminocyclopropane-1-carboxylic acid and its structural analogue by liquid chromatography and ion spray tandem mass spectrometry 896(2000)335

Cycloserine

Separation of enantiomers by capillary electrophoresis–mass spectrometry employing a partial filling technique with a chiral crown ether 875(2000)323

Cyclosporins

Interaction of cyclophilin and cyclosporins monitored by affinity capillary electrophoresis 853(1999)469

Capillary electrophoresis coupled with laser-induced fluorescence polarization as a hybrid approach to ultrasensitive immunoassays 853(1999)555

Cysteine

Capillary electrophoretic separation of *N*-acetylcysteine and its impurities as a method for quality control of pharmaceuticals 853(1999)479

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Picomolar quantitation of free sulfite in foods by means of [⁵⁷Co]hydroxocobalamin and radiometric chromatography of [⁵⁷Co]sulfitocobalamin. Method, applications and significance of coexisting sulfides 881(2000)345

Determination of biologically active low-molecular-mass thiols in human blood. I. Fast qualitative and quantitative, gradient and isocratic reversed-phase high-performance liquid chromatography with photometric and fluorescence detection 895(2000)156

Determination of biologically active low-molecular-mass thiols in human blood. II. High-performance capillary electrophoresis with photometric detection 895(2000)167

Assays for total homocysteine and other thiols by capillary electrophoresis–laser-induced fluorescence detection. I. Preanalytical condition studies 895(2000)173

Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191

Cystine

Capillary electrophoretic separation of *N*-acetylcysteine and its impurities as a method for quality control of pharmaceuticals 853(1999)479

Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191

Cytochromes

Cytochrome *c* unfolding on an anionic surface 863(1999)137

Cytokinins

Determination of dissociation constants of cytokinins by capillary zone electrophoresis 878(2000)249

D**2,4-D**

Determination of some pesticides and intermediates by ion chromatography 884(2000)243

Daidzein

Quantitative determination of isoflavones and coumestrol in soybean by column liquid chromatography 886(2000)55

Dauxorubicin

Anthracycline analysis by capillary electrophoresis. Application to the analysis of daunorubicin in Kaposi sarcoma tumor 853(1999)449

D&C Red No. 27

Rapid quantification of hexachlorobenzene in the color additives D&C Red Nos. 27 and 28 (phloxine B) using solid-phase microextraction and gas chromatography–mass spectrometry 863(1999)37

D&C Red No. 28

Rapid quantification of hexachlorobenzene in the color additives D&C Red Nos. 27 and 28 (phloxine B) using solid-phase microextraction and gas chromatography–mass spectrometry 863(1999)37

DDD

Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl) ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane]) from aged contaminated soil 874(2000)257

DDE

Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl) ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane]) from aged contaminated soil 874(2000)257

DDT

Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl) ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane]) from aged contaminated soil 874(2000)257

Decanoic acid

Esterification of decanoic acid during supercritical fluid extraction employing either methanol-modified carbon dioxide or a methanol trap 858(1999)201

Decanophenone

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

Deethylatrazine

Determination of interfering triazine degradation products by gas chromatography–ion trap mass spectrometry 868(2000)115

Dehydroascorbic acid

Dehydroascorbic acid 881(2000)299

Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables 881(2000)309

Deisopropylatrazine

Determination of interfering triazine degradation products by gas chromatography–ion trap mass spectrometry 868(2000)115

Demethoxydeaminoaziridinylmethylsulfonyldaunorubicin

Determination of 4-demethoxy-3'-deamino-3'-aziridinyl-4'-methylsulphonyldaunorubicin and its 13-hydroxy metabolite by direct injection of human plasma into a column-switching liquid chromatography system with mass spectrometric detection 854(1999)81

Deoxyadenosine

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159

Deoxynivalenol

Determination of nivalenol and deoxynivalenol in wheat using liquid chromatography–mass spectrometry with negative ion atmospheric pressure chemical ionisation 854(1999)45

Quantification of deoxynivalenol in wheat using an immunoaffinity column and liquid chromatography 859(1999)23

Deramciclanc

Determination of deramciclanc and *N*-desmethylderamciclanc in human plasma by liquid chromatography–tandem mass spectrometry using off-line robotic sample pretreatment 896(2000)279

Dermatan sulfate

Analysis of glycosaminoglycan monosaccharides by capillary electrophoresis using indirect laser-induced fluorescence detection 873(2000)269

Design of experiments for capillary electrophoretic enantioresolution of salbutamol using dermatan sulfate 875(2000)411

Desmethylderamciclanc

Determination of deramciclanc and *N*-desmethylderamciclanc in human plasma by liquid chromatography–tandem mass spectrometry using off-line robotic sample pretreatment 896(2000)279

Destruxins

Liquid chromatography–electrospray time-of-flight mass spectrometry for on-line accurate mass determination and identification of cyclodepsipeptides in a crude extract of the fungus *Metarrhizium anisopliae* 872(2000)85

Deuterium

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

Dextran

Analysis of 8-aminonaphthalene-1,3,6-trisulfonate-derivatized oligosaccharides by capillary electrophoresis–electrospray ionization quadrupole ion trap mass spectrometry 858(1999)229

Molecular mass distribution of sodium alginate by high-performance size-exclusion chromatography 864(1999)199

Diacetoxynitrophenylpropylisothiocyanate

Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115

Diacetylcysteine

Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191

Diacetylcystine

Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191

Diacylglycerols

High-performance liquid chromatographic resolution of reverse isomers of 1,2-diacyl-*rac*-glycerols as 3,5-dinitrophenylurethanes 893(2000)261

Diaminobenzidine tetrahydrochloride

Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45

Diazabicycloundecene

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

Diazepam

Coupling device for desorption of drugs from solid-phase extraction-pipette tips and on-line gas chromatographic analysis 858(1999)117

Dibenzodioxin

Highly-sensitive micellar electrokinetic chromatographic analysis of dioxin-related compounds using on-line concentration 853(1999)413

Dibenzofulvene

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

Dibenzofuran

Highly-sensitive micellar electrokinetic chromatographic analysis of dioxin-related compounds using on-line concentration 853(1999)413

Dibenzopyrene diol epoxide

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159

Dibenzoyltartaric acid

Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1

Dibutylamine acetate

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Dibutylmelamine

Solid-phase extraction with a dibutylmelamine-imprinted polymer as triazine herbicide-selective sorbent 889(2000)25

Dibutylmethylphenol

Size-exclusion chromatography with on-line ultraviolet, proton nuclear magnetic resonance and mass spectrometric detection and on-line collection for off-line Fourier transform infrared spectroscopy 857(1999)89

Dicamba

Determination of some pesticides and intermediates by ion chromatography 884(2000)243

Dicarboximides

Inclusion of the fungicide chlozolinate in a multiresidue method 858(1999)123

Dicarboxyethylpyrazines

Validation of a capillary electrophoresis method for determination of 5-aminolevulinic acid and degradation products 883(2000)285

Dichlofluanid

Trace determination of antifouling compounds by on-line solid-phase extraction–gas chromatography–mass spectrometry 885(2000)361

Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261

Dichlorohydrin

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Dichloromethylaminocyclohexylbenzamide

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

Diclofenac sodium

Determination of diclofenac sodium by capillary zone electrophoresis with electrochemical detection 868(2000)101

Dicloxacillin

Application of ion-exchange cartridge clean-up in food analysis II. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in meat using liquid chromatography with ultraviolet detection 855(1999)247

Didanosine

Determination at ppb level of an anti-human immunodeficiency virus nucleoside drug by capillary electrophoresis–electrospray ionization tandem mass spectrometry 895(2000)101

Didexoyadenosine-5'-triphosphate

Determination at ppb level of an anti-human immunodeficiency virus nucleoside drug by capillary electrophoresis–electrospray ionization tandem mass spectrometry 895(2000)101

Diethylpropylvalerolactone

Unknown bisethylisooctanolactone isomers in industrial waste water. Isolation, identification and occurrence in surface water 889(2000)245

Difluorophenylacetic acids

Chromatographic separation of 3,4-difluorophenylacetic acid and its positional isomers using five different techniques 866(2000)281

Dihexylsulfosuccinate

High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253

Dihydrocodeine

Analysis of codeine, dihydrocodeine and their glucuronides in human urine by electrokinetic capillary immunoassays and capillary electrophoresis–ion trap mass spectrometry 895(2000)133

Dihydrolipoic acid

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Dihydropyridines

Resolution of newly synthesized racemic dihydropyridines with different chiral selectors by means of capillary electrophoresis 853(1999)455

High-performance liquid chromatography with amperometric detection applied to the screening of 1,4-dihydropyridines in human plasma 870(2000)105

Dihydropyrimidinone methyl ester

Enantiomeric separation of enzymatic hydrolysis products of dihydropyrimidinone methyl ester with cationic cyclodextrin by capillary electrophoresis 872(2000)279

Dihydroxyadenine

Determination of purines including 2,8-dihydroxyadenine in urine using capillary electrophoresis 894(2000)157

Dihydroxymethoxybenzoxazinone

Simultaneous determination of phenolic acids and 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one in wheat (*Triticum aestivum* L.) by gas chromatography–tandem mass spectrometry 864(1999)315

Dihydroxyphenylalanine

Enantioseparation of 3,4-dihydroxyphenylalanine and 2-hydrazino-2-methyl-3-(3,4-dihydroxyphenyl)propanoic acid by capillary electrophoresis using cyclodextrins 875(2000)379

Diisooctyl phthalate

Size-exclusion chromatography with on-line ultraviolet, proton nuclear magnetic resonance and mass spectrometric detection and on-line collection for off-line Fourier transform infrared spectroscopy 857(1999)89

Diisopropylaminoethanol

Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169

Diltiazem

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Dimedone

Liquid chromatography–mass spectrometry method for the determination of aldehydes derivatized by the Hantzsch reaction 864(1999)191

Dimenhydrinate

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Dimethindene

Mechanistic study of opposite migration order of dimethindene enantiomers in capillary electrophoresis in the presence of native β -cyclodextrin and heptakis(2,3,6-tri-*O*-methyl)- β -cyclodextrin 875(2000)455

Dimethoxyhydrochalone

Migration behavior and separation of active components in *Glycyrrhiza uralensis* Fisch and its commercial extract by micellar electrokinetic capillary chromatography 863(1999)105

Dimethylarsinic acid

Speciation of dimethylarsinic acid and monomethylarsonic acid by solid-phase microextraction–gas chromatography–ion trap mass spectrometry 873(2000)129

Dimethyldiphenylacetamide

Gas chromatography–mass spectrometry–Fourier transform infrared spectrometry and high-performance liquid chromatographic characterization of mononitro- and dinitro-isomers from the nitration of *N,N*-dimethyldiphenylacetamide 878(2000)235

Dimethyl sulfate

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Dimethyl sulfoxide

Separation of purines and pyrimidines by normal-phase high-performance liquid chromatography using dimethyl sulfoxide in binary and ternary eluents 874(2000)45

3,5-Dinitrobenzoylamides

Convenient synthesis of π -acceptor chiral stationary phases for high-performance liquid chromatography from halogen-substituted 3,5-dinitrobenzoylamides 859(1999)143

***N*-(3,5-Dinitrobenzoyl)- α -arylalkylamines**

Experimental support differentiating two proposed chiral recognition models for the resolution of *N*-(3,5-Dinitrobenzoyl)- α -arylalkylamines on high-performance liquid chromatography chiral stationary phases 886(2000)47

Dinitrobenzoyl chloride

Liquid chromatographic determination of biogenic amines in fermented foods after derivatization with 3,5-dinitrobenzoyl chloride 881(2000)517

Dinitrophenylhydrazines

High-performance liquid chromatography of methanol released from pectins after its oxidation to formaldehyde and condensation with 2,4-dinitrophenylhydrazine 863(1999)227

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Characterization of a chemical artifact in the liquid chromatographic determination of 3-butyn-2-one using the 2,4-dinitrophenylhydrazine method 886(2000)303

Dinitrophenylhydrazones

Separation of carbonyl 2,4-dinitrophenylhydrazones by capillary electrochromatography with diode array detection 853(1999)487

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Identification of the oxidation products of the reaction between α -pinene and hydroxyl radicals by gas and high-performance liquid chromatography with mass spectrometric detection 896(2000)135

3,5-Dinitrophenylurethanes

High-performance liquid chromatographic resolution of reverse isomers of 1,2-diacyl-*rac*-glycerols as 3,5-dinitrophenylurethanes 893(2000)261

Dinophysistoxin-1

Liquid chromatography–electrospray ionization mass spectrometry of the diarrhetic shellfish-poisoning toxins okadaic acid, dinophysistoxin-1 and pectenotoxin-6 in bivalves 874(2000)199

Diols

Determination of halogenated mono-alcohols and diols in water by gas chromatography with electron-capture detection 866(2000)65

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Dioxins

Evaluation of a new automated cleanup system for the analysis of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in environmental samples 893(2000)383

Dioxolanes

Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127

Diphacinone

Determination of chlorophacinone and diphacinone in commercial rodenticides by liquid chromatography–UV detection and liquid chromatography–electrospray ionization mass spectrometry 891(2000)249

Diphenhydramine

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Diphenylhexatriene

Postcolumn fluorescence as an alternative to evaporative light scattering detection for ceramide analysis with gradient elution in non-aqueous reversed-phase liquid chromatography 859(1999)99

Diphenylpyrenyl phosphine

Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159

2,2'-Dipyridyldisulfide

Specific thiol determination by micellar electrokinetic chromatography and on-column detection reaction with 2,2'-dipyridyldisulfide 895(2000)179

Disaccharides

see also Saccharides

Isocratic separations of closely-related mono- and disaccharides by high-performance anion-exchange chromatography with pulsed amperometric detection using dilute alkaline spiked with barium acetate 855(1999)539

Separation and determination of alditols and sugars by high-pH anion-exchange chromatography with pulsed amperometric detection 897(2000)195

Diterpenes

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

Dithiocarbamates

Capillary electrophoretic determination of ferric dimethyldithiocarbamate as iron(III) chelate of EDTA 857(1999)365

Dithiopyr

New gas chromatographic method for residue determination of dithiopyr in soil, wheat grain and straw 864(1999)351

Dithiothreitol

Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables 881(2000)309

Dimethyltartaric acid

Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1

Diuretics

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

Modelling retention in liquid chromatography as a function of solvent composition and pH of the mobile phase 869(2000)27

Use of a three-factor interpretive optimisation strategy in the development of an isocratic chromatographic procedure for the screening of diuretics in urine samples using micellar mobile phases 893(2000)321

Diuron

Investigation into the effects of temperature and stirring rate on the solid-phase extraction of diuron from water using a C₁₈ extraction disk 885(2000)41

DNA

see also Nucleic acids

Nanoliter-scale sample preparation methods directly coupled to polymethylmethacrylate-based microchips and gel-filled capillaries for the analysis of oligonucleotides 853(1999)107

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159

Detection of duck hepatitis B virus DNA fragments using on-column intercalating dye labeling with capillary electrophoresis–laser-induced fluorescence 853(1999)309

Analysis of leptin gene expression in chickens using reverse transcription polymerase chain reaction and capillary electrophoresis with laser-induced fluorescence detection 853(1999)321

Stepwise capillary electrophoretic separation of DNA fragments using poly(ethylene oxide) solutions in the presence of electroosmotic flow 853(1999)337

- Capillary electrophoresis of double-stranded DNA in an untreated capillary 853(1999)349
- Side-entry excitation and detection of square capillary array electrophoresis for DNA sequencing 853(1999)359
- Highly alkaline electrolyte for single-stranded DNA separations by electrophoresis in bare silica capillaries 853(1999)371
- Effect of linear polymer additives on the electroosmotic characteristics of agarose gels in ultrathin-layer electrophoresis 853(1999)511
- High-throughput genotyping of factor V Leiden mutation by ultrathin-layer agarose gel electrophoresis 853(1999)519
- Modified apparatus for voltage gradient gel electrophoresis 855(1999)723
- Separation of double-stranded DNA in conventional and isoelectric buffers: studies on stability and separation performance 859(1999)87
- DNA-binding affinities of MyoD and E47 homo- and heterodimers by capillary electrophoresis mobility shift assay 862(1999)231
- Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425
- Ultra-thin-layer agarose gel electrophoresis. II. Separation of DNA fragments on composite agarose–linear polymer matrices 871(2000)289
- Capillary electrophoresis of DNA damage after irradiation: apoptosis and necrosis 871(2000)321
- Modified capillary electrophoresis system for peptide, protein and double-stranded DNA analysis 879(2000)189
- Conductivity detection of polymerase chain reaction products separated by micro-reversed-phase liquid chromatography 883(2000)1
- Free solution mobility of DNA molecules containing variable numbers of cationic phosphoramidate internucleoside linkages 883(2000)267
- Mobile-phase-viscosity dependence on DNA separation in slalom chromatography 886(2000)1
- Application of magnetite and silica–magnetite composites to the isolation of genomic DNA 890(2000)159
- Purification of crude DNA oligonucleotides by solid-phase extraction and reversed-phase high-performance liquid chromatography 890(2000)167
- Large-scale purification of antisense oligonucleotides by high-performance membrane adsorber chromatography 890(2000)179
- Use of voltage gradient gel electrophoresis in apoptotic DNA analysis 890(2000)371
- Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23
- Effects of DNA topology, temperature and solvent viscosity on DNA retardation in slalom chromatography 893(2000)115
- Single-strand conformation polymorphism analysis to detect the p53 mutation in colon tumor samples by capillary electrophoresis 894(2000)171
- Integrated electroosmotically-driven on-line sample purification system for nanoliter DNA sequencing by capillary electrophoresis 894(2000)179
- Rapid, parallel separations of D1S80 alleles in a plastic microchannel chip 894(2000)203
- Effect of ionic strength, pH and polymer concentration on the separation of DNA fragments in the presence of electroosmotic flow 894(2000)219
- Mobility and activation energy of single-stranded DNA in denaturing cross-linked polyacrylamide slab gels 894(2000)231
- Single-strand conformation polymorphism analysis to detect the p53 mutation in colon tumor samples by capillary electrophoresis 895(2000)171
- DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167
- DNA adducts**
Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry 856(1999)515
- Capillary electrochromatography–mass spectrometry for the separation and identification of isomeric polyaromatic hydrocarbon dna adducts derived from in vitro reactions 887(2000)103
- DNA fragments**
Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299
- DNase**
Use of acidic and basic pH and calcium ion addition in the capillary zone electrophoretic characterization of recombinant human deoxyribonuclease, a complex phosphoglycoprotein 853(1999)295

Domoic acid

New fluorimetric method of liquid chromatography for the determination of the neurotoxin domoic acid in seafood and marine phytoplankton 871(2000)1

DOPA

Enantioseparation of 3,4-dihydroxyphenylalanine and 2-hydrazino-2-methyl-3-(3,4-dihydroxyphenyl)propanoic acid by capillary electrophoresis using cyclodextrins 875(2000)379

Dopamine

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Dowfax 8390

High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253

Doxepin

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Doxorubicin

Anthracycline analysis by capillary electrophoresis. Application to the analysis of daunorubicin in Kaposi sarcoma tumor 853(1999)449

Doxycycline

Analysis of doxycycline by capillary electrophoresis. Method development and validation 895(2000)43

Drugs

see also Basic drugs

Solid-phase microextraction–gas chromatography–mass spectrometry of biologically active substances in water samples 873(2000)95

Enantiomer separation of drugs by capillary electromigration techniques 875(2000)3

Enantioselective determination of drugs in body fluids by capillary electrophoresis 875(2000)27

Enantiomeric separations of drugs using mixtures of charged and neutral cyclodextrins 875(2000)123

Separation of drug enantiomers by capillary electrophoresis in the presence of neutral cyclodextrins 875(2000)135

Enantiomer separation of drugs by micellar electrokinetic chromatography using chiral surfactants 875(2000)163

Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors 875(2000)235

Applications of matrix solid-phase dispersion in food analysis 880(2000)63

Regulatory problems caused by contamination, a frequently overlooked cause of veterinary drug residues 882(2000)37

Solid-phase micro-extraction of drugs from biological matrices 885(2000)445

D1S80 alleles

Rapid, parallel separations of D1S80 alleles in a plastic microchannel chip 894(2000)203

Dyes

see also Azo dyes

Detection of duck hepatitis B virus DNA fragments using on-column intercalating dye labeling with capillary electrophoresis–laser-induced fluorescence 853(1999)309

Analysis of dyestuff degradation products by capillary electrophoresis 853(1999)503

Investigation on the iodination reaction of methylene blue by liquid chromatography–mass spectrometry with ionspray ionisation 854(1999)57

Application of electrospray mass spectrometry in the detection and determination of Remazol textile dyes 854(1999)259

High-performance liquid chromatographic determination of some anthraquinone and naphthoquinone dyes occurring in historical textiles 863(1999)235

Determination of sulphonated dyes in water by ion-interaction high-performance liquid chromatography 871(2000)227

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Pressurised liquid extraction of pentachlorophenol from dyestuffs and cotton 873(2000)287

Development of a capillary electrophoresis method for the simultaneous analysis of artificial sweeteners, preservatives and colours in soft drinks 876(2000)213

Analysis of anionic metallized azo and formazan dyes by capillary electrophoresis–mass spectrometry 886(2000)259

Identification of reactive dyes in spent dyebaths and wastewater by capillary electrophoresis–mass spectrometry 886(2000)271

Electrochemical treatment of textile dyes and their analysis by high-performance liquid chromatography with diode array detection 889(2000)253

Biomimetic dyes as affinity chromatography tools in enzyme purification (Review) 891(2000)33

Congophilicity (Congo red affinity) of different β_2 -microglobulin conformations characterized by dye affinity capillary electrophoresis 894(2000)319

Determination of dyes in foodstuffs by capillary zone electrophoresis 898(2000)271

E

Ebrotidine

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Econazole

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Ecstasy

Simultaneous analysis of some amphetamine derivatives in urine by nonaqueous capillary electrophoresis coupled to electrospray ionization mass spectrometry 895(2000)111

Elastase

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

Ellagic acid

Analysis of oak tannins by liquid chromatography-electrospray ionisation mass spectrometry 891(2000)75

Determination of phenolic acids in fruit juices by isocratic column liquid chromatography 891(2000)183

High-performance liquid chromatographic determination with photodiode array detection of ellagic acid in fresh and processed fruits 896(2000)87

Ellagitannins

Characterisation of hydrolysable tannins from leaves of *Betula pubescens* by high-performance liquid chromatography–mass spectrometry 864(1999)283

Emamectin

Liquid chromatographic determination of emamectin, milbemectin, ivermectin and abamectin in crops and confirmation by liquid chromatography–mass spectrometry 896(2000)75

EMD 122 347

Determination of the impurity profile of ethyl-2-(4-((5*R*)-3-[4-(methoxycarboxamidoiminomethyl)-phenyl]-2-oxo-5-oxazolidinylmethyl)-1-piperazinyl) acetate, citrate, a highly active platelet aggregation inhibitor, by liquid chromatography–mass spectrometry 854(1999)13

Enalapril

Study of the *cis*–*trans* isomerization of enalapril by reversed-phase liquid chromatography 871(2000)189

Endocrine disruptors

Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97

Endorphin

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

Endothelins

Simultaneous separation of angiotensin and endothelin peptide families by high-performance liquid chromatography: application to the specific radioimmunoassay measurement of angiotensin II or endothelin-1 from tissue 870(2000)349

Endoxylanases

Determination of reducing end sugar residues in oligo- and polysaccharides by gas–liquid chromatography 866(2000)97

Enterotoxins

Competitive immunoassay for staphylococcal enterotoxin A using capillary electrophoresis with laser-induced fluorescence detection 853(1999)545

Enzyme inhibitors

see also Proteins

Purification of two low-molecular-mass serine proteinase inhibitors from chicken liver 852(1999)207

Determination of human immunodeficiency virus-1 protease inhibitors in patient serum using free solution capillary zone electrophoresis 853(1999)431

Study of the *cis*–*trans* isomerization of enalapril by reversed-phase liquid chromatography 871(2000)189

High-performance liquid chromatography with on-line coupled UV, mass spectrometric and biochemical detection for identification of acetylcholinesterase inhibitors from natural products 872(2000)61

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Fast quantitation of recombinant plasminogen activator inhibitor type 1 in bacterial lysates by micropellicular reversed-phase liquid chromatography 891(2000)85

Enzymes

see also Proteins

Affinity purification of *Schistosoma japonicum* glutathione-S-transferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

Isolation and characterization of cyclic α -(1 \rightarrow 4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

Analysis of angiotension-converting enzyme by capillary electrophoresis 853(1999)185

Determination of enzymatic activity and properties of secretory phospholipase A₂ by capillary electrophoresis 853(1999)285

Use of acidic and basic pH and calcium ion addition in the capillary zone electrophoretic characterization of recombinant human deoxyribonuclease, a complex phosphoglycoprotein 853(1999)295

Direct measurement of ascorbic acid biosynthesis in *Arabidopsis* cell suspension culture using capillary electrophoresis 853(1999)381

Interaction of cyclophilin and cyclosporins monitored by affinity capillary electrophoresis 853(1999)469

A simple assay for formate dehydrogenase activity by gas chromatography–mass spectrometry 855(1999)337

Rapid reversed-phase high-performance liquid chromatographic determination of the regiospecificity of lipoxygenase products on linoleic acid 859(1999)107

Eliminating disulfide exchange during glutamyl endopeptidase digestion of native protein 864(1999)237

Studies on the chromatographic fractionation of *Trichoderma reesei* cellulases by hydrophobic interaction 865(1999)123

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Immobilised metal affinity chromatography of β -galactosidase from unclarified *Escherichia coli* homogenates using expanded bed adsorption 874(2000)27

Separation of the enzyme cofactor pyrroloquinoline quinone and three isomeric analogues by capillary electrophoresis with ion-pairing media 876(2000)193

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295

Effect of the activity levels of the added proteolytic enzyme mixture on free amino acids in ripening Ossau-Iraty cheese 881(2000)69

Use of Edman degradation sequence analysis and matrix-assisted laser desorption/ionization mass spectrometry in designing substrates for matrix metalloproteinases 890(2000)117

Biomimetic dyes as affinity chromatography tools in enzyme purification (Review) 891(2000)33

Analysis of carbohydrates in wood and pulps employing enzymatic hydrolysis and subsequent capillary zone electrophoresis 891(2000)157

Purification of Substance P endopeptidase activity in the rat ventral tegemental area with the Äkta-Purifier chromatographic system 893(2000)107

Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45

Determination of haloalkane dehalogenase activity by capillary zone electrophoresis 895(2000)219

On-line automated high-performance liquid chromatographic determination of total riboflavin phosphates using immobilized acid phosphatase as a pre-column reactor 896(2000)171

Ephedrines

Comparison of various reversed-phase columns for the simultaneous determination of ephedrines in urine by high-performance liquid chromatography 870(2000)23

Derivatization of ephedrine with *o*-phthaldialdehyde for liquid chromatography after treatment with sodium hypochlorite 893(2000)69

Epibatidine

Determination of the in vitro metabolism of (+)- and (–)-epibatidine 896(2000)229

Epichlorohydrin

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Determination of epichlorohydrin by ion chromatography 884(2000)251

Epinephrine

Analysis of epinephrine from fifteen different dental anesthetic formulations by capillary electrophoresis 853(1999)535

Epoxides

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

Ergosterol

Determination of ergosterol on mouldy building materials using isotope dilution and gas chromatography–tandem mass spectrometry 898(2000)227

Eriochrome Black T

Eriochrome Black T as a post-column reagent for the ion chromatographic determination of rare earths 855(1999)111

Erythromycins

Micellar electrokinetic capillary chromatography of macrolide antibiotics. Separation of tylosin, erythromycin and their related substances 857(1999)313

Separation of erythromycin and related substances on base-deactivated reversed-phase silica gel columns 870(2000)227

Erythropoietin

Comparative studies on the analysis of glycosylation heterogeneity of sialic acid-containing glycoproteins using capillary electrophoresis 866(2000)261

Esters

Modeling chromatographic parameters by a novel graph theoretical sub-structural approach 858(1999)187

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Estrogens

Determination of steroid sex hormones and related synthetic compounds considered as endocrine disrupters in water by liquid chromatography–diode array detection–mass spectrometry 892(2000)391

Separation of two groups of oestrogen mimicking compounds using micellar electrokinetic chromatography 895(2000)237

Ethanolamines

Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169

Ethidium bromide

Stepwise capillary electrophoretic separation of DNA fragments using poly(ethylene oxide) solutions in the presence of electroosmotic flow 853(1999)337

***N(O,S)*-Ethoxycarbonyl heptafluorobutyl ester**

Gas chromatographic quantification of amino acid enantiomers in food matrices by their *N(O,S)*-ethoxycarbonyl heptafluorobutyl ester derivatives 866(2000)221

Ethoxyethanol

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Ethoxylates

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

Characterization of nonyl phenol ethoxylates in sewage treatment plants by combined precursor ion scanning and multiple reaction monitoring 854(1999)245

Analysis of nonylphenol polyethoxycarboxylates and their related metabolites by on-line derivatization and ion-trap gas chromatography–mass spectrometry 862(1999)113

Assessment of the pollutant elimination efficiency by gas chromatography–mass spectrometry, liquid chromatography–mass spectrometry and –tandem mass spectrometry. Comparison of conventional and membrane-assisted biological wastewater treatment processes 889(2000)155

Inter-laboratory comparison of liquid chromatographic techniques and enzyme-linked immunosorbent assay for the determination of surfactants in wastewaters 889(2000)195

Concentration of nonylphenol and its polyethoxylated derivatives by polymer-mediated extraction 891(2000)69

Ethylbenzene

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Development of membrane extraction with a sorbent interface–micro gas chromatography system for field analysis 873(2000)13

Simultaneous determination of ethylbenzene, indan, indene and acenaphthene in air by capillary gas chromatography 891(2000)361

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

Ethylene

Automated capillary gas chromatographic system to monitor ethylene emitted from biological materials 868(2000)249

Ethylenediaminedi(*o*-hydroxyphenylacetic acid)

Potential of ethylenediaminedi(*o*-hydroxyphenylacetic acid) and *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid for the determination of metal ions by capillary electrophoresis 890(2000)363

Ethylene oxide

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

4-Ethylguaiaicol

Quantitative analysis of 4-ethylphenol and 4-ethylguaiaicol in red wine 874(2000)101

Ethylidenebis(tryptophan)

Tetrahydro- β -carboline-3-carboxylic acids and contaminants of L-tryptophan 881(2000)501

Ethyl-2-(4-((5*R*)-3-[4-(methoxycarboxamidoiminomethyl)-phenyl]-2-oxo-5-oxazolidinylmethyl)-1-piperazinyl) acetate

Determination of the impurity profile of ethyl-2-(4-((5*R*)-3-[4-(methoxycarboxamidoiminomethyl)-phenyl]-2-oxo-5-oxazolidinylmethyl)-1-piperazinyl) acetate, citrate, a highly active platelet aggregation inhibitor, by liquid chromatography–mass spectrometry 854(1999)13

4-Ethylphenol

Quantitative analysis of 4-ethylphenol and 4-ethylguaiaicol in red wine 874(2000)101

Etodolac

Separation and identification of etodolac and its urinary phase I metabolites using capillary electrochromatography and on-line capillary electrochromatography–electrospray ionisation mass spectrometry coupling 887(2000)393

Eudragit polymers

Ion-exclusion controlled size-exclusion chromatography of methacrylic acid–methyl methacrylate copolymers 872(2000)91

Evernic acid

Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425

Explosives

n-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133

Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Application of solid-phase microextraction to the recovery of explosives and ignitable liquid residues from forensic specimens 885(2000)419

F

Factor V Leiden mutation

High-throughput genotyping of factor V Leiden mutation by ultrathin-layer agarose gel electrophoresis 853(1999)519

Factor VIII

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Use of high-resolution techniques for the characterization of clotting factor VIII 852(1999)175

Erratum to: “Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes”. [J. Chromatogr. A, 852 (1999) 59-71] 872(2000)323

Fats

Automatic microgravimetric determination of fats in milk products by use of supercritical fluid extraction with on-line piezoelectric detection 874(2000)265

Fatty acid amides

Erratum to: “Determination of fatty acid amides as trimethylsilyl derivatives by gas chromatography with mass spectrometric detection” [J. Chromatogr. 849 (1999) 541–552] 859(1999)119

Fatty acid esters

Combined liquid and gas chromatographic characterisation of polyglycerol fatty acid esters 896(2000)239

Fatty acid hydroperoxides

Simultaneous micellar electrokinetic chromatographic determination of isomeric fatty acid hydroperoxides and corresponding hydroxy fatty acids 894(2000)145

Fatty acid methyl esters

Analytical monitoring of the production of biodiesel by high-performance liquid chromatography with various detection methods 858(1999)13

Introduction of large volumes of water-containing samples into a gas chromatograph. Improved retention of volatile solutes through the swing system 876(2000)135

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Fatty acids

see also Carboxylic acids and Organic acids

Indirect detection of organic acids in non-aqueous capillary electrophoresis 853(1999)171

New sheathless interface for coupling capillary electrophoresis to electrospray mass spectrometry evaluated by the analysis of fatty acids and prostaglandins 854(1999)141

Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

Esterification of decanoic acid during supercritical fluid extraction employing either methanol-modified carbon dioxide or a methanol trap 858(1999)201

Rapid reversed-phase high-performance liquid chromatographic determination of the regioselectivity of lipoxygenase products on linoleic acid 859(1999)107

New hydrolysis method for extremely small amount of lipids and capillary gas chromatographic analysis as *N(O)-tert.-butyldimethylsilyl* fatty acid derivatives compared with methyl ester derivatives 862(1999)199

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

Selection of methylation procedures for quantitation of short-chain glycerol-bound compounds formed during thermoxidation 863(1999)171

Development of a headspace solid-phase microextraction procedure for the determination of free volatile fatty acids in waste waters 873(2000)107

Ceramide analysis utilizing gas chromatography–mass spectrometry 876(2000)229

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Determination of organic acids in food samples by capillary zone electrophoresis 881(2000)357

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Application of gas chromatography coupled to chemical ionisation mass spectrometry following headspace solid-phase microextraction for the determination of free volatile fatty acids in aqueous samples 891(2000)287

Polymer additive analysis by pyrolysis–gas chromatography. III. Lubricants 891(2000)313

Gas chromatographic determination of the fatty-acid content of heat-treated green beans 891(2000)367

Fatty acids and β -carotene in Australian purslane (*Portulaca oleracea*) varieties 893(2000)207

Development and validation of a capillary electrophoresis method for the measurement of short-chain organic acids in natural rubber latex 894(2000)135

Simultaneous micellar electrokinetic chromatographic determination of isomeric fatty acid hydroperoxides and corresponding hydroxy fatty acids 894(2000)145

Fenoterol

On-line isotachophoretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219

Fenpyroximate

Determination of fenpyroximate in apples by supercritical fluid extraction and packed capillary liquid chromatography with UV detection 880(2000)121

Ferbam

Capillary electrophoretic determination of ferric dimethyldithiocarbamate as iron(III) chelate of EDTA 857(1999)365

Fermononetin

Migration behavior and separation of active components in *Glycyrrhiza uralensis* Fisch and its commercial extract by micellar electrokinetic capillary chromatography 863(1999)105

Ferulic acid

Determination of phenolic acids in fruit juices by isocratic column liquid chromatography 891(2000)183

Fibrinogenases

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

FK-506

Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69

Flavanoids

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Flavins

Chromatographic determination of riboflavin and its derivatives in food 881(2000)285

Flavones

Isolation and characterization of methoxylated flavones in the flowers of *Primula veris* by liquid chromatography and mass spectrometry 870(2000)453

Flavonoid aglycones

Liquid chromatographic method for the separation and quantification of prominent flavonoid aglycones 897(2000)177

Flavonoids

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

Determination of apigenin in rat plasma by high-performance liquid chromatography 870(2000)463

Liquid chromatography–electrospray ionization mass spectrometry study of the flavonoids of the roots of *Astragalus mongholicus* and *A. membranaceus* 876(2000)87

On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic–diode array detection–mass spectrometric techniques 880(2000)203

Comparative analysis of tea catechins and theaflavins by high-performance liquid chromatography and capillary electrophoresis 881(2000)439

Spectrophotometric and coulometric detection in the high-performance liquid chromatography of flavonoids and optimization of sample treatment for the determination of quercetin in orange juice 881(2000)449

High-speed counter-current chromatography of apple procyanidins 886(2000)65

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Analysis of enzymatically glucosylated flavonoids by capillary electrophoresis 895(2000)279

Flufenacet

Analysis of flufenacet in soil, wheat grain and straw by gas chromatography 886(2000)319

Fluorenone

Sample cleanup and reversed-phase high-performance liquid chromatographic analysis of polar aromatic compounds in groundwater samples from a former gas plant 862(1999)137

Fluorenylmethoxycarbonyl compounds

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

Fluorescein

Utilization of fluorescein sodium salt for the indirect fluorimetric detection in micellar electrokinetic chromatography 872(2000)247

Fluorocinnamic acid

Analytical techniques used for monitoring the biodegradation of fluorinated compounds in waste streams from pharmaceutical production 889(2000)177

1-Fluoro-2,4-dinitrobenzene

Determination of neurotoxin 3-*N*-oxalyl-2,3-diaminopropionic acid and non-protein amino acids in *Lathyrus sativus* by precolumn derivatization with 1-fluoro-2,4-dinitrobenzene 883(2000)113

Fluoronitrobenzoxadiazole

New fluorimetric method of liquid chromatography for the determination of the neurotoxin domoic acid in seafood and marine phytoplankton 871(2000)1

Micellar electrokinetic capillary chromatographic separation and fluorescent detection of amino acids derivatized with 4-fluoro-7-nitro-2,1,3-benzoxadiazole 876(2000)183

Fluoroquinolones

High-performance thin-layer chromatography method for monitoring norfloxacin residues on pharmaceutical equipment surfaces 862(1999)209

Direct liquid chromatographic enantiomer separation of new fluoroquinolones including gemifloxacin 879(2000)113

Fluvoxamine

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Capillary electrochromatography of basic compounds using octadecyl-silica stationary phases with an amine-containing mobile phase 872(2000)315

Fluvoxetone

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Folate vitamers

Determination of folate vitamers in food and in Italian reference diet by high-performance liquid chromatography 855(1999)237

Folic acid

Analysis of some folate monoglutamates by high-performance liquid chromatography–mass spectrometry. I 864(1999)59

Erratum to: ‘‘Liquid chromatographic method for determining added folic acid in fortified cereal products’’ [J. Chromatogr. A, 826 (1998) 235–240] 867(2000)289

Folinic acid

Analysis of some folate monoglutamates by high-performance liquid chromatography–mass spectrometry. I 864(1999)59

Formaldehyde

High-performance liquid chromatography of methanol released from pectins after its oxidation to formaldehyde and condensation with 2,4-dinitrophenylhydrazine 863(1999)227

Poly(allylamine) beads as selective sorbent for preconcentration of formaldehyde and acetaldehyde in high-performance liquid chromatographic analysis 886(2000)83

Simultaneous supercritical fluid derivatization and extraction of formaldehyde by the Hantzsch reaction 896(2000)51

Formate dehydrogenase

A simple assay for formate dehydrogenase activity by gas chromatography–mass spectrometry 855(1999)337

Formic acid

A simple assay for formate dehydrogenase activity by gas chromatography–mass spectrometry 855(1999)337

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Formononetin

Quantitative determination of isoflavones and coumestrol in soybean by column liquid chromatography 886(2000)55

Fritillaria alkaloids

Pre-column derivatization and gas chromatographic determination of alkaloids in bulbs of *Fritillaria* 859(1999)183

Fructans

Determination of inulin in meat products by high-performance liquid chromatography with refractive index detection 881(2000)591

Fructose

Optimization of simulated moving bed plants with low efficient stationary phases: separation of fructose and glucose 868(2000)169

Determination of inulin in meat products by high-performance liquid chromatography with refractive index detection 881(2000)591

Fucans

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

Fullerenes

Synthesis and characteristics of [60]fullerene polysiloxane stationary phase for capillary gas chromatography 867(2000)177

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Novel approach to the analysis and use of fullerenes in capillary electrophoresis 873(2000)257

Fullerenes separation with monomeric type C₃₀ stationary phase in high-performance liquid chromatography 883(2000)55

Fumaric acid

Determination of fumaric acid in apple juice by on-line coupled capillary isotachopheresis–capillary zone electrophoresis with UV detection 891(2000)175

Fumonisin

Effect of solvents on the fumonisins analysis by high-performance liquid chromatography with AccQ.Fluor as the derivatizing reagent 870(2000)469

Furaldehyde

Determination of aldehydes in food by high-performance liquid chromatography with biosensor coupling and micromembrane suppressors 880(2000)233

Furans

Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49

Evaluation of a new automated cleanup system for the analysis of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in environmental samples 893(2000)383

Furfurals

High-performance liquid chromatographic determination of Maillard compounds in store-brand and name-brand ultra-high-temperature-treated cows' milk 881(2000)599

Furosine

High-performance liquid chromatographic determination of Maillard compounds in store-brand and name-brand ultra-high-temperature-treated cows' milk 881(2000)599

Furoylquinolinecarboxaldehyde

Labeling effects on the isoelectric point of green fluorescent protein 853(1999)21

Methyl- β -cyclodextrin modified micellar electrokinetic capillary chromatography with laser-induced fluorescence for separation and detection of phospholipids 894(2000)129

Fusaproliferin

Determination of the *Fusarium* mycotoxins, fusaproliferin and beauvericin by high-performance liquid chromatography–electrospray ionization mass spectrometry 858(1999)175

G **β -Galactosidase**

Immobilised metal affinity chromatography of β -galactosidase from unclarified *Escherichia coli* homogenates using expanded bed adsorption 874(2000)27

Galectins

Reinforcement of frontal affinity chromatography for effective analysis of lectin–oligosaccharide interactions 890(2000)261

Gallic acid

Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235

Separation of gallic acid from *Cornus officinalis* Sieb. et Zucc by high-speed counter-current chromatography 886(2000)309

Analysis of oak tannins by liquid chromatography–electrospray ionisation mass spectrometry 891(2000)75

Determination of phenolic acids in fruit juices by isocratic column liquid chromatography 891(2000)183

Gallotannins

Characterisation of hydrolysable tannins from leaves of *Betula pubescens* by high-performance liquid chromatography–mass spectrometry 864(1999)283

Ganciclovir

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Gases

Gas chromatographic analysis of trace impurities in chlorine trifluoride 873(2000)229

Gemifloxacin

Direct liquid chromatographic enantiomer separation of new fluoroquinolones including gemifloxacin 879(2000)113

Genistein

Quantitative determination of isoflavones and coumestrol in soybean by column liquid chromatography 886(2000)55

Gentamicin sulfate

Capillary Electrophoresis analysis of gentamicin sulphate with UV detection after pre-capillary derivatization with 1,2-phthalic dicarboxaldehyde and mercaptoacetic acid 895(2000)67

Germanium oxide

High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43

Ginsenosides

Liquid chromatography–electrospray mass spectrometric identification of ginsenosides in *Panax ginseng* roots 854(1999)69

High-performance thin-layer chromatographic determination of six major ginsenosides in *Panax ginseng* 868(2000)269

Globulins

New affinity nylon membrane used for adsorption of γ -globulin 867(2000)161

Congophilicity (Congo red affinity) of different β_2 -microglobulin conformations characterized by dye affinity capillary electrophoresis 894(2000)319

Sodium dodecylsulfate capillary gel electrophoretic measurement of the concentration ratios of albumin and α_2 -macroglobulin in cerebrospinal fluid and serum of patients with neurological disorders 895(2000)339

Glutathione-S-transferase

Affinity purification of *Schistosoma japonicum* glutathione-S-transferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

Glucagon

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

Glucagon-like peptide-1

Proteolytic cleavage of glucagon-like peptide-1 by pancreatic β cells and by fetal calf serum analyzed by mass spectrometry 852(1999)285

Glucans

Isolation and characterization of cyclic α -(1 \rightarrow 4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

Glucobrassicin

Optimization of glucosinolate separation by micellar electrokinetic capillary chromatography using a Doehlert's experimental design 864(1999)155

Glucose

Optimization of simulated moving bed plants with low efficient stationary phases: separation of fructose and glucose 868(2000)169

Glucose ester

Analysis of oak tannins by liquid chromatography-electrospray ionisation mass spectrometry 891(2000)75

Glucosides

Screening for limonoid glucosides in *Citrus tangerina* (Tanaka) Tseng by high-performance liquid chromatography–electrospray ionization mass spectrometry 874(2000)13

Glucosinolates

Optimization of glucosinolate separation by micellar electrokinetic capillary chromatography using a Doehlert's experimental design 864(1999)155

Determination of glucosinolates in rapeseed and *Thlaspi caerulescens* plants by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 889(2000)75

Glucotropaeolin

Optimization of glucosinolate separation by micellar electrokinetic capillary chromatography using a Doehlert's experimental design 864(1999)155

Glucuronides

Method for quantification of morphine and its 3- and 6-glucuronides, codeine, codeine glucuronide and 6-monoacetylmorphine in human blood by liquid chromatography–electrospray mass spectrometry for routine analysis in forensic toxicology 854(1999)109

Analysis of codeine, dihydrocodeine and their glucuronides in human urine by electrokinetic capillary immunoassays and capillary electrophoresis–ion trap mass spectrometry 895(2000)133

Glut1

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

Glutamates

Analysis of some folate monoglutamates by high-performance liquid chromatography–mass spectrometry. I 864(1999)59

Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317

Glutamyl endopeptidase

Eliminating disulfide exchange during glutamyl endopeptidase digestion of native protein 864(1999)237

Glutamyl peptides

Analysis of isomeric glutamyl peptides by capillary electrophoresis. Application to stability studies 888(2000)275

Glutathione

Quantitation of oxidized and reduced glutathione in plasma by micellar electrokinetic capillary electrophoresis 853(1999)215

Glutathione oxidation in real time by thermospray liquid chromatography–mass spectrometry 862(1999)161

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Determination of biologically active low-molecular-mass thiols in human blood. I. Fast qualitative and quantitative, gradient and isocratic reversed-phase high-performance liquid chromatography with photometric and fluorescence detection 895(2000)156

Determination of biologically active low-molecular-mass thiols in human blood. II. High-performance capillary electrophoresis with photometric detection 895(2000)167

Assays for total homocysteine and other thiols by capillary electrophoresis–laser-induced fluorescence detection. I. Preanalytical condition studies 895(2000)173

Specific thiol determination by micellar electrokinetic chromatography and on-column detection reaction with 2,2'-dipyridyldisulfide 895(2000)179

Glycerides

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Glycerols

Preparative thin-layer chromatographic separation and subsequent gas chromatographic–mass spectrometric analysis of monoacylglycerols derived from butter oil by fungal degradation 855(1999)617

Analytical monitoring of the production of biodiesel by high-performance liquid chromatography with various detection methods 858(1999)13

Combined liquid and gas chromatographic characterisation of polyglycerol fatty acid esters 896(2000)239

Glycoalkaloids

Reversed-phase liquid chromatographic separation and simultaneous profiling of steroidal glycoalkaloids and their aglycones 863(1999)25

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

Glycoamylase

Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295

Glycodeoxycholic acid

Characterization of partition and thermodynamic properties of cephalosporins using micellar electrokinetic chromatography in glycodeoxycholic acid solution 873(2000)237

Glycopeptides

Microscale characterization of the structure–activity relationship of a heparin-binding glycopeptide using affinity capillary electrophoresis and immobilized enzymes 853(1999)189

Glycoproteins

Use of high-resolution techniques for the characterization of clotting factor VIII 852(1999)175

Use of acidic and basic pH and calcium ion addition in the capillary zone electrophoretic characterization of recombinant human deoxyribonuclease, a complex phosphoglycoprotein 853(1999)295

Liquid chromatography–electrospray mass spectrometry as a tool for the analysis of sulfated oligosaccharides from mucin glycoproteins 854(1999)131

Integrated approach to the multidimensional analysis of complex biological samples by microseparation techniques. Analysis of glycoprotein factor associated with cancer cachexia 857(1999)183

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VII. Comparison of chiral recognition ability of ovoglycoprotein from chicken and Japanese quail egg whites 858(1999)155

One-step capillary isoelectric focusing for the separation of the recombinant human immunodeficiency virus envelope glycoprotein glycoforms 866(2000)121

Comparative studies on the analysis of glycosylation heterogeneity of sialic acid-containing glycoproteins using capillary electrophoresis 866(2000)261

Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors 875(2000)235

Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255

Glycosaminoglycans

Chiral resolution of basic drugs by capillary electrophoresis with new glycosaminoglycans 864(1999)163

Analysis of glycosaminoglycan monosaccharides by capillary electrophoresis using indirect laser-induced fluorescence detection 873(2000)269

Glycosides

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Chromatographic determination of riboflavin and its derivatives in food 881(2000)285

Monoterpenes in grape juice and wines 881(2000)557

Analysis of enzymatically glucosylated flavonoids by capillary electrophoresis 895(2000)279

Glycosidic surfactants

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Glycosphingolipids

Capillary electrochromatography with novel stationary phases. IV. Retention behavior of glycosphingolipids on porous and non-porous octadecyl sulfonated silica 887(2000)287

Glycyrrhetic acid

Migration behavior and separation of active components in *Glycyrrhiza uralensis* Fisch and its commercial extract by micellar electrokinetic capillary chromatography 863(1999)105

Glycyrrhizic acid

Migration behavior and separation of active components in *Glycyrrhiza uralensis* Fisch and its commercial extract by micellar electrokinetic capillary chromatography 863(1999)105

Glyphosate

New methods for determination of glyphosate and (aminomethyl)phosphonic acid in water and soil 886(2000)207

Extraction of glyphosate by a supported liquid membrane technique 889(2000)93

Gold clusters

Synthesis and study of the cluster $[\text{Pd}(\text{AuPPh}_3)_6(\text{HgNO}_3)](\text{NO}_3)$ by reversed-phase high-performance liquid chromatography, thermoanalysis, infrared, nuclear magnetic resonance, and fast atom bombardment mass spectrometry 862(1999)29

Gonyautoxin

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatroscan (mark 5) and flame thermionic detection 855(1999)657

Goserelin

Liquid chromatography–electrospray mass spectrometry of multicomponent peptide mixtures. Characterization of a mixture from the synthesis of the hormone goserelin 889(2000)119

Green fluorescent protein

Labeling effects on the isoelectric point of green fluorescent protein 853(1999)21

Green leaf volatile compounds

Retention index database for identification of general green leaf volatiles in plants by coupled capillary gas chromatography–mass spectrometry 890(2000)313

Growth hormones

see also Hormones

High-yield purification of biosynthetic human growth hormone secreted in *Escherichia coli* periplasmic space 852(1999)441

Separation of oxidized and deamidated human growth hormone variants by isocratic reversed-phase high-performance liquid chromatography 855(1999)147

Development of analytical and preparative chromatographic separations of novel growth hormone secretagogue compounds 872(2000)75

Determination of trenbolone and zeranol in bovine muscle and liver by liquid chromatography–electrospray mass spectrometry 882(2000)53

Use of molecularly imprinted solid-phase extraction for the selective clean-up of clenbuterol from calf urine 889(2000)105

Guanidine

Reversed-phase ion-pair chromatography coupled to electrospray ionisation mass spectrometry by on-line removal of the counterions 854(1999)155

H**Halides**

Simultaneous determination of cyanogen chloride and cyanogen bromide in treated water at sub- $\mu\text{g/L}$ levels by a new solid-phase microextraction–gas chromatographic–electron-capture detection method 897(2000)307

Haloacetic acids

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

Solid-phase microextraction coupled with gas chromatography–ion trap mass spectrometry for the analysis of haloacetic acids in water 859(1999)159

Haloacetonitriles

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

Haloalkane dehalogenase

Determination of haloalkane dehalogenase activity by capillary zone electrophoresis 895(2000)219

Haloalkanes

Determination of haloalkane dehalogenase activity by capillary zone electrophoresis 895(2000)219

Halogenated compounds

see also Organochlorine compounds, Organohalogen compounds, Polychlorinated biphenyls, Polychlorinated dibenzodioxins, Polychlorinated dibenzofurans, and Polychlorinated dibenzothiophenes

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

Determination of halogenated mono-alcohols and diols in water by gas chromatography with electron-capture detection 866(2000)65

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Halogens

Determination of bromide, bromate and other anions with ion chromatography and an inductively coupled plasma mass spectrometer as element-specific detector 862(1999)39

Halohydrins

Determination of halogenated mono-alcohols and diols in water by gas chromatography with electron-capture detection 866(2000)65

Heavy metals

see also Metals

Use of ion chromatography for the determination of heavy and transition metals in biochemical samples 857(1999)343

Analysis of heavy metals in atmospheric particulate by ion chromatography 888(2000)145

Hemoglobins

Kinetics of appearance of hemorphins from bovine hemoglobin peptic hydrolysates by a direct coupling of reversed-phase high-performance liquid chromatography and electrospray ionization mass spectrometry 873(2000)185

High-performance liquid chromatography coupled to enzyme-amplified biochemical detection for the analysis of hemoglobin after pre-column biotinylation 886(2000)19

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. II. Separation of proteins 890(2000)37

Hemorphins

Kinetics of appearance of hemorphins from bovine hemoglobin peptic hydrolysates by a direct coupling of reversed-phase high-performance liquid chromatography and electrospray ionization mass spectrometry 873(2000)185

Heparin

Microscale characterization of the structure–activity relationship of a heparin-binding glycopeptide using affinity capillary electrophoresis and immobilized enzymes 853(1999)189

Analysis of glycosaminoglycan monosaccharides by capillary electrophoresis using indirect laser-induced fluorescence detection 873(2000)269

Hexachlorobenzenes

Rapid quantification of hexachlorobenzene in the color additives D&C Red Nos. 27 and 28 (phloxine B) using solid-phase microextraction and gas chromatography–mass spectrometry 863(1999)37

Hexadecylsulfonate

Low-capacity cation-exchange chromatography of ultraviolet-absorbing urinary basic metabolites using a reversed-phase column coated with hexadecylsulfonate 886(2000)297

Hexafluoroisopropanol

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Hexenuronic acid

Analysis of carbohydrates in wood and pulps employing enzymatic hydrolysis and subsequent capillary zone electrophoresis 891(2000)157

Hexobarbital

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VII. Comparison of chiral recognition ability of ovoglycoprotein from chicken and Japanese quail egg whites 858(1999)155

Hexoses

Isocratic separations of closely-related mono- and disaccharides by high-performance anion-exchange chromatography with pulsed amperometric detection using dilute alkaline spiked with barium acetate 855(1999)539

Hippuric acid

Field amplified injection in the presence of salts for capillary electrophoresis 853(1999)3

Histamine

Use of capillary electrophoresis with UV detection as a screening method to determine histamine in fish samples 853(1999)541

Histidine

Histidines in affinity tags and surface clusters for immobilized metal-ion affinity chromatography of trimeric tumor necrosis factor α 852(1999)117

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

HIV-p24

Preliminary comparison of precursor scans and liquid chromatography–tandem mass spectrometry on a hybrid quadrupole time-of-flight mass spectrometer 854(1999)119

Homoarginine

Analysis of β -*N*-oxalyl-L- α , β -diaminopropionic acid and homoarginine in *Lathyrus sativus* by capillary zone electrophoresis 857(1999)295

Homocysteine

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using γ -cyclodextrin 877(2000)217

Determination of biologically active low-molecular-mass thiols in human blood. I. Fast qualitative and quantitative, gradient and isocratic reversed-phase high-performance liquid chromatography with photometric and fluorescence detection 895(2000)156

Determination of biologically active low-molecular-mass thiols in human blood. II. High-performance capillary electrophoresis with photometric detection 895(2000)167

Assays for total homocysteine and other thiols by capillary electrophoresis–laser-induced fluorescence detection. I. Preanalytical condition studies 895(2000)173

Screening of homocysteine from newborn blood spots by high-performance liquid chromatography with coulometric array detection 896(2000)183

Homoserine lactone

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

Homotaurine

Determination of homotaurine as impurity in calcium acamprostate by capillary zone electrophoresis 853(1999)421

Homovanillic acid

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Hormones

see also Growth hormones and Plant hormones

Analysis of leptin gene expression in chickens using reverse transcription polymerase chain reaction and capillary electrophoresis with laser-induced fluorescence detection 853(1999)321

Erratum to: "Determination of fatty acid amides as trimethylsilyl derivatives by gas chromatography with mass spectrometric detection" [J. Chromatogr. 849 (1999) 541–552] 859(1999)119

Separation and characterization of multicomponent peptide mixtures by liquid chromatography–electrospray ionization mass spectrometry. Application to crude products of the synthesis of leuprolide 870(2000)315

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Quantitation of anabolic hormones and their metabolites in bovine serum and urine by liquid chromatography–tandem mass spectrometry 870(2000)511

Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97

Recognition of oxytocin and oxytocin-related peptides in aqueous media using a molecularly imprinted polymer synthesized by the epitope approach 889(2000)111

Liquid chromatography–electrospray mass spectrometry of multicomponent peptide mixtures. Characterization of a mixture from the synthesis of the hormone goserelin 889(2000)119

Analysis of human parathyroid hormone (1–84) products. Separation of a major impurity in synthetic products by ion-pairing reversed-phase high-performance liquid chromatography 890(2000)127

Determination of steroid sex hormones and related synthetic compounds considered as endocrine disruptors in water by liquid chromatography–diode array detection–mass spectrometry 892(2000)391

Analysis of a peptide hormone mixture of therapeutic interest by liquid chromatography coupled to high-flow pneumatically assisted electrospray mass spectrometry 893(2000)95

Use of chiral liquid chromatography–tandem mass spectrometry to investigate the metabolism of racemic cholecystokinin-B antagonists 896(2000)217

Humic acids

Highly reproducible capillary zone electrophoresis of humic acids in cyclodextrin- or oligosaccharide-modified background electrolytes 895(2000)345

Humic substances

Capillary electrophoretic behaviour of humic substances in physical gels 863(1999)195

Hyaluronic acid

Molecular weight determination of hyaluronic acid by gel filtration chromatography coupled to matrix-assisted laser desorption ionization mass spectrometry 852(1999)573

Hydrazinomethyl(dihydroxyphenyl)propanoic acid monohydrate

Enantioseparation of 3,4-dihydroxyphenylalanine and 2-hydrazino-2-methyl-3-(3,4-dihydroxyphenyl)propanoic acid by capillary electrophoresis using cyclodextrins 875(2000)379

Hydrazoic acid

Gas chromatographic study of complexing in the system hydrazoic acid–tributyl phosphate–nitric acid–uranyl nitrate 859(1999)41

Hydrazones

Separation of carbonyl 2,4-dinitrophenylhydrazones by capillary electrochromatography with diode array detection 853(1999)487

Hydrocarbons

see also Volatile organic compounds

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

Selective extraction of hydrocarbons, phosphonates and phosphonic acids from soils by successive supercritical fluid and pressurized liquid extractions 866(2000)231

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Extraction of hydrocarbon contamination from soils using accelerated solvent extraction 874(2000)217

Characterization of a commercial gas chromatography–flame ionization detection system for the in situ determination of C₅–C₁₀ hydrocarbons in ambient air 878(2000)215

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

Comparison of comprehensive two-dimensional gas chromatography and gas chromatography – mass spectrometry for the characterization of complex hydrocarbon mixtures 892(2000)29

Supercritical fluid extraction of hydrocarbons and 2-alkylcyclobutanones for the detection of irradiated foodstuffs 897(2000)259

Hydrocarbons, aromatic

see also Polynuclear aromatic hydrocarbons

New operational modes for multidimensional and comprehensive gas chromatography by using cryogenic modulation 866(2000)203

Time-resolved cryogenic modulation for targeted multidimensional capillary gas chromatography analysis 892(2000)15

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

Hydrochloric acid

Experimental study on moving neutralization reaction boundary created with the strong reactive electrolytes of HCl and NaOH in agarose gel 891(2000)337

Hydrocortisone

Direct identification and quantitation of prednisone in the presence of overlapping hydrocortisone by liquid chromatography with electrospray and atmospheric-pressure chemical-ionisation mass spectrometry 870(2000)135

Hydrogen

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

Hydrogen peroxide

Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45

Hydroperoxides

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

Hydroquinone

Analysis of hydroquinone and some of its ethers by using capillary electrochromatography 887(2000)489

Hydroxocobalamin

Picomolar quantitation of free sulfite in foods by means of [⁵⁷Co]hydroxocobalamin and radiometric chromatography of [⁵⁷Co]sulfitecobalamin. Method, applications and significance of coexisting sulfides 881(2000)345

Hydroxy acids

Liquid chromatographic resolution of 2-hydroxycarboxylic acids on a new chiral stationary phase derived from (*S*)-leucine 868(2000)31

Enantiomeric separation and discrimination of 2-hydroxy acids as *O*-trifluoroacetylated (*S*)-(+)-3-methyl-2-butyl esters by achiral dual-capillary column gas chromatography 874(2000)91

Application of ligand-exchange capillary electrophoresis to the chiral separation of α-hydroxy acids and β-blockers 875(2000)307

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Configurational analysis of chiral acids as *O*-trifluoroacetylated (–)-menthyl esters by achiral dual-capillary column gas chromatography 891(2000)257

Hydroxyanthraquinones

Computer-aided development of a high-performance liquid chromatographic method for the determination of hydroxyanthraquinone derivatives in Chinese herb medicine rhubarb 857(1999)167

Preparative isolation and purification of hydroxyanthraquinones from *Rheum officinale* Baill by high-speed counter-current chromatography using pH-modulated stepwise elution 858(1999)103

Hydroxybiphenyls

Application of solid-phase microextraction–gas chromatography–mass spectrometry to characterize intermediates in a joint solar–microbial process for total mineralization of Aroclor 1254 873(2000)53

Hydroxycotinine

Application of liquid separation techniques to the determination of the main urinary nicotine metabolites 870(2000)29

Hydroxyethylcellulose

Quantitation of protein binding to the capillary wall in acidic, isoelectric buffers and means for minimizing the phenomenon 894(2000)273

Hydroxy fatty acids

Simultaneous micellar electrokinetic chromatographic determination of isomeric fatty acid hydroperoxides and corresponding hydroxy fatty acids 894(2000)145

17-Hydroxy-16-hydroxymethyl-3-methoxyestra-1,3,5(10)-triene

High-performance liquid chromatographic methods for monitoring of isomers of 17-hydroxy-16-hydroxymethyl-3-methoxyestra-1,3,5(10)-triene 852(1999)433

2-Hydroxyibuprofen

Uniform-sized molecularly imprinted polymer for (*S*)-ibuprofen. Retention properties in aqueous mobile phases 857(1999)117

Hydroxyindoleacetic acid

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Hydroxymethylfuraldehyde

Determination of aldehydes in food by high-performance liquid chromatography with biosensor coupling and micromembrane suppressors 880(2000)233

 α -Hydroxy-3-phenoxybenzeneacetonitrile

Determination of enantiomeric excess of α -hydroxy-3-phenoxybenzeneacetonitrile and its *n*-butyl ester by chiral high-performance liquid chromatography 893(2000)189

3-Hydroxypropionic acid

Stereoselective binding of 2,3-substituted 3-hydroxypropionic acids on an immobilised human serum albumin chiral stationary phase: stereochemical characterisation and quantitative structure–retention relationship study 876(2000)75

Hydroxypropylcellulose

Separation of proteins by sodium dodecylsulfate capillary electrophoresis in hydroxypropylcellulose sieving matrix with laser-induced fluorescence detection 894(2000)291

Hydroxyquinolines

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

8-Hydroxyquinoline 5-sulfonic acid

Determination of trace cadmium in environmental water samples using ion-interaction reversed-phase liquid chromatography with fluorescence detection 877(2000)123

Hyoscyamine

Strategies for supercritical fluid extraction of hyoscyamine and scopolamine salts using basified modifiers 863(1999)47

Enantioseparation of atropine by capillary electrophoresis using sulfated β -cyclodextrin: application to a plant extract 868(2000)285

I**Ibuprofen**

see also Nonsteroidal anti-inflammatory drugs and Profens

Uniform-sized molecularly imprinted polymer for (*S*)-ibuprofen. Retention properties in aqueous mobile phases 857(1999)117

Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143

Solid-phase microextraction–gas chromatography–mass spectrometry of biologically active substances in water samples 873(2000)95

Imazosulfuron

Liquid chromatographic determination of imazosulfuron in drinking water and in soil using ultraviolet detection 857(1999)327

Imidacloprid

Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497

Imidazoles

Determination of four nitroimidazole residues in poultry meat by liquid chromatography–mass spectrometry 882(2000)89

Imidazolic compounds

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Imides

Gas chromatographic–mass spectrometric separation, identification and determination of C₆–C₁₂ cyclic imides in thermally treated epoxy and alkyd resins to locate hot spots inside large electricity generators 862(1999)17

Immunoglobulins

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent–detergent treatment 852(1999)83

Removal of detergent and solvent from solvent–detergent-treated immunoglobulins 852(1999)87

Use of spray-dried zirconia microspheres in the separation of immunoglobulins from cell culture supernatant 890(2000)15

Scale-down of continuous protein purification by annular chromatography. Design parameters for the smallest unit 890(2000)53

Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy 893(2000)235

Indan

Simultaneous determination of ethylbenzene, indan, indene and acenaphthene in air by capillary gas chromatography 891(2000)361

Indanediones

Determination of chlorofacinone and diphacinone in commercial rodenticides by liquid chromatography–UV detection and liquid chromatography–electrospray ionization mass spectrometry 891(2000)249

Indene

Simultaneous determination of ethylbenzene, indan, indene and acenaphthene in air by capillary gas chromatography 891(2000)361

Indole

Determination of indole-3-acetic acid, tryptophan and other indoles in must and wine by high-performance liquid chromatography with fluorescence detection 855(1999)227

Indole-3-acetic acid

Determination of indole-3-acetic acid, tryptophan and other indoles in must and wine by high-performance liquid chromatography with fluorescence detection 855(1999)227

Ingenol

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

Inorganic anions

see also Anions, Inorganic ions and Special Issue Index

Analysis of anions in aqueous samples by ion chromatography and capillary electrophoresis. A comparative study of peak modeling and validation criteria 852(1999)487

Electrophoretic mobility study on ion–ion interactions in an aqueous solution 853(1999)63

Quantification of anions and cations in environmental water samples. Measurements with capillary electrophoresis and indirect-UV detection 853(1999)403

Analysis of low-molecular-mass inorganic and organic anions by ion chromatography–atmospheric pressure ionization mass spectrometry 854(1999)275

Determination of inorganic anions by ion chromatography using a graphitized carbon column dynamically coated with cetyltrimethylammonium ions 855(1999)261

Ultra-short columns for low-pressure ion chromatography 857(1999)175

Potential for chlorate interference in ion chromatographic determination of total nitrogen in natural waters following alkaline persulfate digestion 857(1999)337

Determination of trace anions in organic solvents 858(1999)55

Effect of drying on the degradation of cationic surfactants and separation performance in capillary zone electrophoresis of inorganic anions 863(1999)81

Ion chromatography determination of trace level bromate by large volume injection with conductivity and spectrophotometric detection after post column derivatisation 864(1999)263

Erratum to: “Optimization of conductivity of low-molecular-mass anions in capillary zone electrophoresis” [J. Chromatogr. A, 850 (1999) 355–362] 868(2000)321

Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391

Capillary electrophoretic determination of thiosulfate and its oxidation products 879(2000)235

Separation of anions by ion chromatography–capillary electrophoresis 879(2000)245

Electrophoretic methods in the analysis of beverages 880(2000)243

Influence of organic solvents in the mobile phase on the determination of carboxylic acids and inorganic anions in grape juice by ion chromatography 881(2000)387

Advances in the determination of inorganic anions by ion chromatography 881(2000)607

Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629

Eliminating the chlorite interference in US Environmental Protection Agency Method 317.0 permits analysis of trace bromate levels in all drinking water matrices 882(2000)309

- On-line microdialysis–ion chromatographic determination of inorganic anions in olive-oil mill wastewater 882(2000)321
- Detection techniques in ion analysis: what are our choices? 884(2000)3
- Application of neutral hydrophobic hypercrosslinked polystyrene to the separation of inorganic anions by ion chromatography 884(2000)41
- Simplified column-switching technology for the determination of traces of anions in the presence of high concentrations of other anions 884(2000)53
- Retention behaviour of strong acid anions in ion-exclusion chromatography on sulfonate and carboxylate stationary phases 884(2000)61
- Enhanced conductometric detection of cyanide in suppressed ion chromatography 884(2000)75
- Determination of thiosulfate, thiocyanate and polythionates in a mixture by ion-pair chromatography with ultraviolet absorbance detection 884(2000)81
- Suppressed electrostatic ion chromatography with tetraborate as eluent and its application to the determination of inorganic anions in snow and rainwater 884(2000)161
- Simultaneous ion-exclusion/cation-exchange chromatography of anions and cations in acid rain waters on a weakly acidic cation-exchange resin by elution with sulfosalicylic acid 884(2000)167
- Determination of inorganic anions in environmental waters with a hydroxide-selective column 884(2000)175
- Determination of chloride, sulfate and nitrate in groundwater samples by ion chromatography 884(2000)185
- Comparison of on-line coupling of ion-chromatography with atmospheric pressure ionization mass spectrometry and with inductively coupled plasma mass spectrometry as tools for the ultra-trace analysis of bromate in surface water samples 884(2000)191
- Determination of inorganic and organic anionic arsenic species in water by ion chromatography coupled to hydride generation–inductively coupled plasma atomic emission spectrometry 884(2000)211
- Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261
- Elution mechanism in electrostatic ion chromatography with histidine as an isoelectric ampholytic mobile phase 884(2000)287
- Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297
- Ultra-rapid analysis of nitrate and nitrite by capillary electrophoresis 884(2000)311
- Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309
- New preconditioning strategy for the determination of inorganic anions with capillary zone electrophoresis using indirect UV detection 890(2000)355
- Simple interface for capillary electrophoresis–inductively coupled plasma atomic emission spectrometry 891(2000)139
- Peak shapes in open tubular ion-exchange capillary electrochromatography of inorganic anions 892(2000)303
- Capillary zone electrophoresis in phosphate buffer – known or unknown? 894(2000)89
- Determination of soluble anions and cations from waters of pulp and paper mills with on-line coupled capillary electrophoresis 895(2000)189
- Simultaneous determination of inorganic anions and cations by capillary electrophoresis with indirect UV detection 895(2000)269
- Inorganic cations**
see also Inorganic ions, Metal cations and Special Issue Index
- Interface for capillary electrophoresis coupled with inductively coupled plasma atomic emission spectrometry 853(1999)141
- Quantification of anions and cations in environmental water samples. Measurements with capillary electrophoresis and indirect-UV detection 853(1999)403
- Direct determination of small cations in proteinaceous samples using a flow injection–capillary electrophoresis system 857(1999)321
- Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391
- Electrophoretic methods in the analysis of beverages 880(2000)243
- Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425
- Eliminating the chlorite interference in US Environmental Protection Agency Method 317.0 permits analysis of trace bromate levels in all drinking water matrices 882(2000)309

Detection techniques in ion analysis: what are our choices?
884(2000)3

Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113

Simultaneous ion-exclusion/cation-exchange chromatography of anions and cations in acid rain waters on a weakly acidic cation-exchange resin by elution with sulfosalicylic acid 884(2000)167

Determination of inorganic and organic anionic arsenic species in water by ion chromatography coupled to hydride generation–inductively coupled plasma atomic emission spectrometry 884(2000)211

Determination of inorganic cations in brine solutions by ion chromatography 884(2000)223

Trace analysis of heavy metals in groundwater samples by ion chromatography with post-column reaction and ultraviolet–visible detection 884(2000)229

Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261

Elution mechanism in electrostatic ion chromatography with histidine as an isoelectric ampholytic mobile phase 884(2000)287

On-line preconcentration of niobium(V) and tantalum(V) as 4-(2-pyridylazo) resorcinol–citrate ternary complexes in geological samples by ion interaction high-performance liquid chromatography 885(2000)369

Capillary zone electrophoresis in phosphate buffer – known or unknown? 894(2000)89

Determination of soluble anions and cations from waters of pulp and paper mills with on-line coupled capillary electrophoresis 895(2000)189

Simultaneous determination of inorganic anions and cations by capillary electrophoresis with indirect UV detection 895(2000)269

Inorganic ions

see also Inorganic anions and Inorganic cations

Developments in sample preparation and separation techniques for the determination of inorganic ions by ion chromatography and capillary electrophoresis 856(1999)145

Coupling of chromatographic ion-exchange reaction with change-partner reaction. Presumed mechanism on the ion–ion interaction of inorganic ions in Sephadex G-15 columns 869(2000)91

Inositol

Ion chromatographic determination of inositol in infant formulae and clinical products for enteral feeding 879(2000)129

Insulin

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

Proteolytic cleavage of glucagon-like peptide-1 by pancreatic β cells and by fetal calf serum analyzed by mass spectrometry 852(1999)285

Eliminating disulfide exchange during glutamyl endopeptidase digestion of native protein 864(1999)237

Novel simulated moving-bed method for reduced solvent consumption 873(2000)149

Quantitative analysis of urinary C-peptide by liquid chromatography–tandem mass spectrometry with a stable isotopically labelled internal standard 896(2000)275

Interleukins

Analysis of recombinant and modified proteins by capillary zone electrophoresis coupled with electrospray ionization tandem mass spectrometry 855(1999)695

Inulin

Determination of inulin in meat products by high-performance liquid chromatography with refractive index detection 881(2000)591

Iodine

Visualization of solute migration in chromatographic columns. Quantitation of the concentration in a migrating zone 867(2000)71

Iodoacetamidofluorescein

Assays for total homocysteine and other thiols by capillary electrophoresis–laser-induced fluorescence detection. I. Preanalytical condition studies 895(2000)173

Ionenes

Ion chromatographic selectivity of polyelectrolyte sorbents based on some aliphatic and aromatic ionenes 884(2000)31

Ionogenic compounds

Reversed-phase high-performance liquid chromatography of ionogenic compounds: comparison of retention models 859(1999)1

Ionones

Enantiomer separation of α -ionone using gas chromatography with cyclodextrin derivatives as chiral stationary phases 865(1999)201

Ions

Retention models for ions in chromatography 855(1999)3

Migration of ions in capillary electrochromatography
892(2000)291

Irganox

Determination of low levels of an antioxidant in polyolefins by large-volume injection temperature-programmed packed capillary liquid chromatography 864(1999)103

Irgarols

Monitoring of priority pesticides and other organic pollutants in river water from Portugal by gas chromatography–mass spectrometry and liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)13

Trace determination of antifouling compounds by on-line solid-phase extraction–gas chromatography–mass spectrometry
885(2000)361

Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261

Iridium

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Iridoid glycosides

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Iron

Eliminating the chlorite interference in US Environmental Protection Agency Method 317.0 permits analysis of trace bromate levels in all drinking water matrices 882(2000)309

ISIS 2922

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis
871(2000)311

Isoascorbic acid

Indirect detection of organic acids in non-aqueous capillary electrophoresis 853(1999)171

Isocyanates

Determination of 1-(2-methoxyphenyl)piperazine derivatives of isocyanates at low concentrations by temperature-programmed miniaturized liquid chromatography 892(2000)67

Isoflavones

Separation and purification of isoflavones from *Pueraria lobata* by high-speed counter-current chromatography 855(1999)709

Quantitative determination of isoflavones and coumestrol in soybean by column liquid chromatography 886(2000)55

Isoimperatorin

Preparative isolation and purification of notopterol and isoimperatorin from *Notopterygium forbesii* boiss (Chinese traditional medicinal herb) by high-speed counter-current chromatography 883(2000)67

Isoliquiritigenin

Migration behavior and separation of active components in *Glycyrrhiza uralensis* Fisch and its commercial extract by micellar electrokinetic capillary chromatography 863(1999)105

Isopropanol

Determination of trace anions in organic solvents 858(1999)55

Headspace gas chromatography method for the analysis of volatile impurities in a hormone replacement transdermal patch
859(1999)113

N-Isopropylacrylamide

Concentration of nonylphenol and its polyethoxylated derivatives by polymer-mediated extraction 891(2000)69

Isovaline

Detection of new sequences of peptaibol antibiotics trichotoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179

Chiral determination of amino acids by capillary electrophoresis and laser-induced fluorescence at picomolar concentrations
894(2000)259

Ivermectin

Liquid chromatographic determination of emamectin, milbemectin, ivermectin and abamectin in crops and confirmation by liquid chromatography–mass spectrometry
896(2000)75

K

Ketoacids

Comparative methodology in the determination of α -oxocarboxylates in aqueous solution. Ion chromatography versus gas chromatography after oximation, extraction and esterification
867(2000)143

Ketocarboxylates

Comparative methodology in the determination of α -oxocarboxylates in aqueous solution. Ion chromatography versus gas chromatography after oximation, extraction and esterification
867(2000)143

Ketoconazole

Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399

Ketones

see also Carbonyl compounds and Volatile organic compounds

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

Modeling chromatographic parameters by a novel graph theoretical sub-structural approach 858(1999)187

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Pressurised liquid extraction of ketones of polycyclic aromatic hydrocarbons from soil 893(2000)201

Ketoprofen

see also Nonsteroidal anti-inflammatory drugs and Profens

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VII. Comparison of chiral recognition ability of ovoglycoprotein from chicken and Japanese quail egg whites 858(1999)155

Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Kitols

Liquid chromatography–nuclear magnetic resonance spectroscopy 856(1999)199

Krypton

Development and characterization of chlorine-selective pulsed discharge emission detector for gas chromatography 872(2000)141

Applications using the chlorine-selective pulsed discharge emission detector 872(2000)167

Characterization of chlorinated compounds using a dual chlorine-selective pulsed discharge emission detector–helium-pulsed discharge photoionization detector system 872(2000)179

L**Lactalbumin**

Analysis of bovine whey proteins in soybean dairy-like products by capillary electrophoresis 859(1999)77

Lactams

Stability of penicillin antibiotic residues in meat during storage. Ampicillin 882(2000)135

Determination of cephalosporins in raw bovine milk by high-performance liquid chromatography 882(2000)145

New configurations and applications of molecularly imprinted polymers 889(2000)15

Lactate

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Lactate dehydrogenase

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Lactic acid

Direct chiral resolution of lactic acid in food products by capillary electrophoresis 875(2000)371

Lactic acid oligomers

Capillary zone electrophoretic analysis of positively charged poly(ethylene oxide) macromolecules using non-covalent polycation-coated fused-silica capillary and indirect UV detection 864(1999)335

Lactoglobulins

Ion-exchange chromatography of proteins near the isoelectric points 852(1999)31

Analysis of bovine whey proteins in soybean dairy-like products by capillary electrophoresis 859(1999)77

Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

Lactones

Determination of oak lactones in barrel-aged wines and in oak extracts by stable isotope dilution analysis 857(1999)239

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Unknown bisethylisooctanolactone isomers in industrial waste water. Isolation, identification and occurrence in surface water 889(2000)245

Lactose

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

Lactuloses

Isocratic separations of closely-related mono- and disaccharides by high-performance anion-exchange chromatography with pulsed amperometric detection using dilute alkaline spiked with barium acetate 855(1999)539

Lanthanides

see also Rare earth ions

Eriochrome Black T as a post-column reagent for the ion chromatographic determination of rare earths 855(1999)111

Time-resolved high-performance liquid chromatography fluorescence detector using a nanosecond pulsed light source for detecting lanthanide-chelated compounds 859(1999)13

Improved capillary electrophoresis method for measuring rare-earth elements in synthetic geochemical standards 884(2000)317

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N,N*-diethylcarbamoil methylene phosphonate as stationary phase 888(2000)137

Separation and direct UV detection of lanthanides complexed with cupferron by capillary electrophoresis 895(2000)263

Laser-induced fluorescence

Anthracycline analysis by capillary electrophoresis. Application to the analysis of daunorubicine in Kaposi sarcoma tumor 853(1999)449

Lead

Binding of lead to collagen type I and V and $\alpha_2(I)$ CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

Lecithin

Ion chromatographic determination of inositol in infant formulae and clinical products for enteral feeding 879(2000)129

Lectins

Reinforcement of frontal affinity chromatography for effective analysis of lectin–oligosaccharide interactions 890(2000)261

Leptin

Analysis of leptin gene expression in chickens using reverse transcription polymerase chain reaction and capillary electrophoresis with laser-induced fluorescence detection 853(1999)321

Leucine

Liquid chromatographic resolution of 2-hydroxycarboxylic acids on a new chiral stationary phase derived from (*S*)-leucine 868(2000)31

Leuprolide

Separation and characterization of multicomponent peptide mixtures by liquid chromatography–electrospray ionization mass spectrometry. Application to crude products of the synthesis of leuprolide 870(2000)315

Lidocaine

Highly sensitive bioassay of lidocaine in human plasma by high-performance liquid chromatography–tandem mass spectrometry 854(1999)3

Coupling device for desorption of drugs from solid-phase extraction-pipette tips and on-line gas chromatographic analysis 858(1999)117

Multiple solid-phase microextraction 878(2000)27

Limonoid glucosides

Screening for limonoid glucosides in *Citrus tangerina* (*Tanaka*) *Tseng* by high-performance liquid chromatography–electrospray ionization mass spectrometry 874(2000)13

Linoleic acid

Rapid reversed-phase high-performance liquid chromatographic determination of the regioselectivity of lipoxygenase products on linoleic acid 859(1999)107

Linolenic acid

Fatty acids and β -carotene in Australian purslane (*Portulaca oleracea*) varieties 893(2000)207

Linuron

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Lipid hydroperoxides

Determination of peroxides by capillary zone electrophoresis with amperometric detection 868(2000)127

Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159

Lipids

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry I. High-temperature gas chromatography 855(1999)601

Preparative thin-layer chromatographic separation and subsequent gas chromatographic–mass spectrometric analysis of monoacylglycerols derived from butter oil by fungal degradation 855(1999)617

Erratum to: "Determination of fatty acid amides as trimethylsilyl derivatives by gas chromatography with mass spectrometric detection" [J. Chromatogr. 849 (1999) 541–552] 859(1999)119

Application of polydivinylbenzene liquid chromatography columns to remove lipid material from fish tissue extracts for the analysis of semivolatiles 859(1999)235

New hydrolysis method for extremely small amount of lipids and capillary gas chromatographic analysis as *N(O)-tert.-butyldimethylsilyl* fatty acid derivatives compared with methyl ester derivatives 862(1999)199

Efficient method for (lyso)phospholipid class separation by high-performance liquid chromatography using an evaporative light-scattering detector 864(1999)179

High-performance liquid chromatography method for the characterization of rhamnolipid mixtures produced by *Pseudomonas aeruginosa* UG2 on corn oil 864(1999)211

Analysis of pesticide residues in matrices with high lipid contents by membrane separation coupled on-line to a high-performance liquid chromatography system 869(2000)427

Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363

Quantitation and characterization of phospholipids in pharmaceutical formulations by liquid chromatography–mass spectrometry 876(2000)97

Chromatographic analysis of unsaponifiable compounds of olive oils and fat-containing foods 881(2000)105

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Chromatographic analysis of tocol-derived lipid antioxidants 881(2000)197

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. II: High-temperature gas chromatography–chemical ionization mass spectrometry 883(2000)75

Update on solid-phase extraction for the analysis of lipid classes and related compounds 885(2000)321

Capillary electrochromatography with novel stationary phases. IV. Retention behavior of glycosphingolipids on porous and non-porous octadecyl sulfonated silica 887(2000)287

Separation of the molecular species of intact phosphatidylethanolamines and their *N*-monomethyl and *N,N*-dimethyl derivatives by high-performance liquid chromatography on a C_8 column 891(2000)349

Methyl- β -cyclodextrin modified micellar electrokinetic capillary chromatography with laser-induced fluorescence for separation and detection of phospholipids 894(2000)129

Lipoic acid

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Lipoproteins

Capillary electrophoretic analysis of recombinant human apolipoprotein E. Calibration mode of a protein reference material 853(1999)237

Enantioselective binding analysis of verapamil to plasma lipoproteins by capillary electrophoresis–frontal analysis 875(2000)447

Lipoxygenases

Rapid reversed-phase high-performance liquid chromatographic determination of the regioselectivity of lipoxygenase products on linoleic acid 859(1999)107

Lithium

Determination of lithium in wines by ion chromatography 881(2000)645

Lorazepam

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

Loxapine

Separation of antipsychotic drugs (clozapine, loxapine) and their metabolites by capillary zone electrophoresis 853(1999)461

Lubricants

Polymer additive analysis by pyrolysis–gas chromatography. III. Lubricants 891(2000)313

Lysophospholipids

Efficient method for (lyso)phospholipid class separation by high-performance liquid chromatography using an evaporative light-scattering detector 864(1999)179

Lysozyme

Lysozyme refolding with immobilized GroEL column chromatography 878(2000)197

Pressure-induced retention of the lysozyme on reversed-phase liquid chromatography 891(2000)207

M

Macroglobulin

Sodium dodecylsulfate capillary gel electrophoretic measurement of the concentration ratios of albumin and α_2 -macroglobulin in cerebrospinal fluid and serum of patients with neurological disorders 895(2000)339

Macrolide antibiotics

Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69

Magnesium

Comparative study of divalent metals and amines as silanol-blocking agents in reversed-phase liquid chromatography 855(1999)121

Malic acid

Determination of fumaric acid in apple juice by on-line coupled capillary isotachopheresis–capillary zone electrophoresis with UV detection 891(2000)175

Malonates

Liquid chromatography–electrospray ionization mass spectrometry study of the flavonoids of the roots of *Astragalus mongholicus* and *A. membranaceus* 876(2000)87

Mandelic acid

Investigation of mandelic acid bonding on Pirkle type chromatographic stationary phases by Raman spectroscopy 893(2000)37

Manganese

Quantitative analysis of manganese, chromium and molybdenum by ion-pair reversed-phase high-performance liquid chromatography with pre-column derivatization and UV-visible detection 857(1999)351

Matrix metalloproteinases

Use of Edman degradation sequence analysis and matrix-assisted laser desorption/ionization mass spectrometry in designing substrates for matrix metalloproteinases 890(2000)117

Melengestrol

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

Metal cations

see also Inorganic cations

How to achieve higher repeatability and reproducibility in capillary electrophoresis 853(1999)83

Ultra-short columns for low-pressure ion chromatography 857(1999)175

Use of ion chromatography for the determination of heavy and transition metals in biochemical samples 857(1999)343

Investigation of Cr(III) hydrolytic polymerisation products by capillary electrophoresis–inductively coupled plasma-mass spectrometry 872(2000)227

Determination of lithium in wines by ion chromatography 881(2000)645

Ion-chromatographic behavior of alkali metal cations and ammonium ion on zirconium-adsorbing silica gel 884(2000)123

Improved capillary electrophoresis method for measuring rare-earth elements in synthetic geochemical standards 884(2000)317

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N,N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69

Potential of ethylenediaminedi(*o*-hydroxyphenylacetic acid) and *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid for the determination of metal ions by capillary electrophoresis 890(2000)363

Discoloration of ceramic hydroxyapatite used for protein chromatography 891(2000)93

Simple interface for capillary electrophoresis–inductively coupled plasma atomic emission spectrometry 891(2000)139

Standardization of capillary zone electropherograms obtained by using field enhanced sample stacking 894(2000)3

Simultaneous determination of alkali, alkaline earth and transition metal ions by capillary electrophoresis with indirect UV detection 895(2000)205

Metal chelates

Quantitative analysis of manganese, chromium and molybdenum by ion-pair reversed-phase high-performance liquid chromatography with pre-column derivatization and UV-visible detection 857(1999)351

Capillary electrophoretic determination of ferric dimethyldithiocarbamate as iron(III) chelate of EDTA 857(1999)365

Time-resolved high-performance liquid chromatography fluorescence detector using a nanosecond pulsed light source for detecting lanthanide-chelated compounds 859(1999)13

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Adsorption and selectivity characteristics of several human serum proteins with immobilised hard Lewis metal ion–chelate adsorbents 890(2000)95

Potential of ethylenediaminedi(*o*-hydroxyphenylacetic acid) and *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid for the determination of metal ions by capillary electrophoresis 890(2000)363

Separation selectivity of anionic metal complexes of *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid in ion and ion electrokinetic chromatography 895(2000)255

Metal complexes

On-line preconcentration of niobium(V) and tantalum(V) as 4-(2-pyridylazo) resorcinol–citrate ternary complexes in geological samples by ion interaction high-performance liquid chromatography 885(2000)369

Analysis of anionic metallized azo and formazan dyes by capillary electrophoresis–mass spectrometry 886(2000)259

Simple interface for capillary electrophoresis–inductively coupled plasma atomic emission spectrometry 891(2000)139

Integration of a microextraction system. Solvent extraction of a Co–2-nitroso-5-dimethylaminophenol complex on a microchip 894(2000)19

Separation and direct UV detection of lanthanides complexed with cupferron by capillary electrophoresis 895(2000)263

Metallomesogenic polymer

Synthesis and analytical properties of a novel columnar metallomesogenic polymer 862(1999)65

Metalloporphyrins

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Metalloproteinases

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

Use of Edman degradation sequence analysis and matrix-assisted laser desorption/ionization mass spectrometry in designing substrates for matrix metalloproteinases 890(2000)117

Metals

see also Heavy metals and Transition metals

Understanding of peak deterioration in hyphenated speciation systems due to gas–liquid separation in the hydride generation interface 855(1999)291

Ion-chromatographic study of the possible absorption of copper and zinc by the skin of *Rana pipiens* 884(2000)143

Trace analysis of heavy metals in groundwater samples by ion chromatography with post-column reaction and ultraviolet–visible detection 884(2000)229

Capillary electrophoresis–inductively coupled plasma mass spectrometry interface with minimised dead volume for high separation efficiency 891(2000)129

Metamitron

Determination of some pesticides and intermediates by ion chromatography 884(2000)243

Methacrylic acid

Determination of methacrylic acid in the drain of a biotrickling filter using isotachopheresis and capillary zone electrophoresis 862(1999)237

Methacrylic acid–methyl methacrylate polymers

Ion-exclusion controlled size-exclusion chromatography of methacrylic acid–methyl methacrylate copolymers 872(2000)91

Methadone

Enantioselective analysis of methadone in saliva by liquid chromatography–mass spectrometry 871(2000)163

Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography 873(2000)293

Chiral capillary electrophoresis as predictor for separation of drug enantiomers in continuous flow zone electrophoresis 895(2000)51

Methamphetamine

Profiling of impurities in illicit methamphetamine by high-performance liquid chromatography and capillary electrochromatography 870(2000)53

Determination of amphetamine and methamphetamine in serum via headspace derivatization solid-phase microextraction–gas chromatography–mass spectrometry 896(2000)265

Methanol

High-performance liquid chromatography of methanol released from pectins after its oxidation to formaldehyde and condensation with 2,4-dinitrophenylhydrazine 863(1999)227

Study of dead volume measurement in packed subcritical fluid chromatography with ODS columns and carbon dioxide–modifier mobile phases 866(2000)241

Influence of pH*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63

Methoxyglucobrassicin

Optimization of glucosinolate separation by micellar electrokinetic capillary chromatography using a Doehler's experimental design 864(1999)155

Methoxyphenylpiperazine

Determination of 1-(2-methoxyphenyl)piperazine derivatives of isocyanates at low concentrations by temperature-programmed miniaturized liquid chromatography 892(2000)67

Methyl aspartate receptor blockers

Determination of the *N*-methyl-D-aspartate receptor NR2B subunit blocker Ro 63-1908 in rat, cynomolgus monkey and human plasma by high-performance liquid chromatography with column switching and fluorescence detection 870(2000)189

Methyl bromide

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

Methylbutyldimethyltrifluoroacetamide

Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97

Methyl *tert*-butyl ether

Analysis of methyl *tert*-butyl ether and its degradation products by direct aqueous injection onto gas chromatography with mass spectrometry or flame ionization detection systems 857(1999)205

3-Methyl-1-(2',4'-dinitrophenyl)pyrazol

Characterization of a chemical artifact in the liquid chromatographic determination of 3-butyn-2-one using the 2,4-dinitrophenylhydrazine method 886(2000)303

Methylene blue

Investigation on the iodination reaction of methylene blue by liquid chromatography–mass spectrometry with ionspray ionisation 854(1999)57

Sensitive determination of dissolved sulfide in estuarine water by solid-phase extraction and high-performance liquid chromatography of methylene blue 883(2000)305

Methylformamide

Highly selective and efficient determination of US Environmental Protection Agency priority phenols employing solid-phase extraction and non-aqueous capillary electrophoresis 896(2000)95

Methyl iodide

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

(Methylidobutyl)methylpiperazine

Novel, trifunctional diamine for silica coating in capillary zone electrophoresis 894(2000)53

Quantitation of protein binding to the capillary wall in acidic, isoelectric buffers and means for minimizing the phenomenon 894(2000)273

Methylmercury

Experimental design of a microwave-assisted extraction–derivatization method for the analysis of methylmercury 889(2000)185

Methyl oleate

Novel use of a dual-zone restricted access sorbent: normal-phase solid-phase extraction separation of methyl oleate from polynuclear aromatic hydrocarbons stemming from semi-permeable membrane devices 883(2000)143

Methylphosphonic acid

Efficiency of pretreatment of aqueous samples using a macroporous strong anion-exchange resin on the determination of nerve gas hydrolysis products by gas chromatography–mass spectrometry after *tert*-butyldimethylsilylation 891(2000)295

Methylpyrrolidone

Determination of trace anions in organic solvents 858(1999)55

Methyltolyl sulfide

Determination of gas phase peroxyacetic acid using pre-column derivatization with organic sulfide reagents and liquid chromatography 855(1999)551

Metsulfuron-methyl

Determination of sulfonyleurea degradation products in soil by liquid chromatography–ultraviolet detection followed by confirmatory liquid chromatography–tandem mass spectrometry 855(1999)575

Microcystins

Isolation and preparative purification of microcystin variants 883(2000)103

Microglobulins

Congophilicity (Congo red affinity) of different β_2 -microglobulin conformations characterized by dye affinity capillary electrophoresis 894(2000)319

Milbemectin

Liquid chromatographic determination of emamectin, milbemectin, ivermectin and abamectin in crops and confirmation by liquid chromatography–mass spectrometry 896(2000)75

Mildronate

Analysis of carnitine and acylcarnitines in urine by capillary electrophoresis 895(2000)309

Molybdenum

Quantitative analysis of manganese, chromium and molybdenum by ion-pair reversed-phase high-performance liquid chromatography with pre-column derivatization and UV-visible detection 857(1999)351

Monoacetylmorphine

Method for quantification of morphine and its 3- and 6-glucuronides, codeine, codeine glucuronide and 6-monoacetylmorphine in human blood by liquid chromatography–electrospray mass spectrometry for routine analysis in forensic toxicology 854(1999)109

Monoacylglycerols

Preparative thin-layer chromatographic separation and subsequent gas chromatographic–mass spectrometric analysis of monoacylglycerols derived from butter oil by fungal degradation 855(1999)617

Monobromobimane

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Determination of biologically active low-molecular-mass thiols in human blood. I. Fast qualitative and quantitative, gradient and isocratic reversed-phase high-performance liquid chromatography with photometric and fluorescence detection 895(2000)156

Determination of biologically active low-molecular-mass thiols in human blood. II. High-performance capillary electrophoresis with photometric detection 895(2000)167

Monoclonal antibodies

see also Antibodies

Capillary electrophoresis coupled with laser-induced fluorescence polarization as a hybrid approach to ultrasensitive immunoassays 853(1999)555

Use of spray-dried zirconia microspheres in the separation of immunoglobulins from cell culture supernatant 890(2000)15

Monoglutamates

Analysis of some folate monoglutamates by high-performance liquid chromatography–mass spectrometry. I 864(1999)59

Monomethylarsonic acid

Speciation of dimethylarsinic acid and monomethylarsonic acid by solid-phase microextraction–gas chromatography–ion trap mass spectrometry 873(2000)129

Monosaccharides

see also Saccharides

Isocratic separations of closely-related mono- and disaccharides by high-performance anion-exchange chromatography with pulsed amperometric detection using dilute alkaline spiked with barium acetate 855(1999)539

Analysis of glycosaminoglycan monosaccharides by capillary electrophoresis using indirect laser-induced fluorescence detection 873(2000)269

Analysis of carbohydrates in wood and pulps employing enzymatic hydrolysis and subsequent capillary zone electrophoresis 891(2000)157

Separation and determination of alditols and sugars by high-pH anion-exchange chromatography with pulsed amperometric detection 897(2000)195

Monoterpenes

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Monoterpenes in grape juice and wines 881(2000)557

Morphine

Routine determination of morphine, morphine 3- β -D-glucuronide and morphine 6- β -D-glucuronide in human serum by liquid chromatography coupled to electrospray mass spectrometry 854(1999)93

Method for quantification of morphine and its 3- and 6-glucuronides, codeine, codeine glucuronide and 6-monoacetylmorphine in human blood by liquid chromatography–electrospray mass spectrometry for routine analysis in forensic toxicology 854(1999)109

Determination of morphine by capillary electrophoresis immunoassay in thermally reversible hydrogel-modified buffer and laser-induced fluorescence detection 895(2000)1

Micellar electrokinetic and high-performance liquid chromatographic determination of potential manufacturing impurities in pholcodine 895(2000)9

Morphine glucuronides

Routine determination of morphine, morphine 3- β -D-glucuronide and morphine 6- β -D-glucuronide in human serum by liquid chromatography coupled to electrospray mass spectrometry 854(1999)93

Morpholines

Application and comparison of derivatized cellulose and amylose chiral stationary phases for the separation of enantiomers of pharmaceutical compounds by high-performance liquid chromatography 855(1999)411

Mycotoxins

see also Toxins

Determination of nivalenol and deoxynivalenol in wheat using liquid chromatography–mass spectrometry with negative ion atmospheric pressure chemical ionisation 854(1999)45

Determination of the *Fusarium* mycotoxins, fusaproliferin and beauvericin by high-performance liquid chromatography–electrospray ionization mass spectrometry 858(1999)175

Determination of zearalenone in grains by high-performance liquid chromatography–tandem mass spectrometry after solid-phase extraction with RP-18 columns or immunoaffinity columns 858(1999)167

Quantification of deoxynivalenol in wheat using an immunoaffinity column and liquid chromatography 859(1999)23

Reversed-phase liquid chromatographic behavior of the mycotoxins, citrinin and ochratoxin A 862(1999)147

Determination of ochratoxin A in wine by means of immunoaffinity column clean-up and high-performance liquid chromatography 864(1999)89

Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473

Chromatographic determination of the mycotoxin patulin in fruit and fruit juices 882(2000)17

Simultaneous determination of trichothecene mycotoxins and zearalenone in cereals by gas chromatography–mass spectrometry 882(2000)23

Solid-phase extraction applied to the determination of ochratoxin A in wines by reversed-phase high-performance liquid chromatography 882(2000)29

Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321

Determination of patulin in apple juice by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 897(2000)365

MyoD-E47 protein

DNA-binding affinities of MyoD and E47 homo- and heterodimers by capillary electrophoresis mobility shift assay 862(1999)231

Myoglobin

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. II. Separation of proteins 890(2000)37

N**Nafcillin**

Application of ion-exchange cartridge clean-up in food analysis II. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in meat using liquid chromatography with ultraviolet detection 855(1999)247

Naphthalenedicarboxaldehyde

Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233

Naphthalenes

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Naphthalenesulfonates

Equilibrium analysis of reactions between aromatic anions and nonionic surfactant micelles by capillary zone electrophoresis 853(1999)55

On-line solid-phase extraction–ion-pair liquid chromatography–electrospray mass spectrometry for the trace determination of naphthalene monosulphonates in water 854(1999)187

Solid-phase extraction of polycyclic aromatic compounds 885(2000)273

Stability of sulfonated derivatives of benzene and naphthalene on disposable solid-phase extraction pre-columns and in an aqueous matrix 889(2000)231

Determination of naphthalenesulfonates in water by on-line ion-pair solid-phase extraction and ion-pair liquid chromatography with fast-scanning fluorescence detection 890(2000)289

Naphthalenesulfonic acids

Use of volatile amines as ion-pairing agents for the high-performance liquid chromatographic–tandem mass spectrometric determination of aromatic sulfonates in industrial wastewater 854(1999)175

Separation of isomeric naphthalenesulphonic acids by micro high-performance liquid chromatography with mobile phases containing cyclodextrin 871(2000)139

Naphthodianthrones

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Naphthoquinones

High-performance liquid chromatographic determination of some anthraquinone and naphthoquinone dyes occurring in historical textiles 863(1999)235

Naphthoquinonesulfonate

Liquid chromatographic determination of aniline in table-top sweeteners based on pre-column derivatization with 1,2-naphthoquinone-4-sulfonate 859(1999)227

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Naphthylamine

Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291

Neosaxitoxin

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatrosan (mark 5) and flame thermionic detection 855(1999)657

Neostigmine

On-line isotachophoretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219

Neuropeptides

Analysis of neuropeptide Y and its metabolites by high-performance liquid chromatography–electrospray ionization mass spectrometry and integrated sample clean-up with a novel restricted-access sulphonic acid cation exchanger 890(2000)135

Neurotoxins

see also Toxins

Analysis of β -*N*-oxalyl-L- α , β -diaminopropionic acid and homoarginine in *Lathyrus sativus* by capillary zone electrophoresis 857(1999)295

Nicarbazin

Determination of nicarbazin and clopidol in poultry feeds by liquid chromatography 882(2000)79

Nicotinamide

Analysis of 2 β -carbomethoxy-3 β -(4-fluorophenyl)-*N*-(3-iodo-*E*-allyl)nortropine in rat plasma. I. Method development and validation by capillary electrophoresis 895(2000)87

Nicotine

Optimization study for the reversed-phase ion-pair liquid chromatographic determination of nicotine in commercial tobacco products 852(1999)451

Application of non-aqueous capillary electrophoresis with electrochemical detection to the determination of nicotine in tobacco 853(1999)27

Application of liquid separation techniques to the determination of the main urinary nicotine metabolites 870(2000)29

Nifedipine

Quantitation of nifedipine in human plasma by on-line solid-phase extraction and high-performance liquid chromatography 870(2000)115

Niobium

On-line preconcentration of niobium(V) and tantalum(V) as 4-(2-pyridylazo) resorcinol–citrate ternary complexes in geological samples by ion interaction high-performance liquid chromatography 885(2000)369

Nitrate

Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629

Determination of chloride, sulfate and nitrate in groundwater samples by ion chromatography 884(2000)185

Ultra-rapid analysis of nitrate and nitrite by capillary electrophoresis 884(2000)311

Nitrated polynuclear aromatic hydrocarbons

Determination of 1,3-, 1,6-, 1,8-dinitropyrene and 1-nitropyrene in airborne particulate by column liquid chromatography with electrochemical detection 897(2000)393

Nitrendipine

Quantitation of nifedipine in human plasma by on-line solid-phase extraction and high-performance liquid chromatography 870(2000)115

Nitric acid

Gas chromatographic study of complexing in the system hydrazoic acid–tributyl phosphate–nitric acid–uranyl nitrate 859(1999)41

Nitrite

Ultra-rapid analysis of nitrate and nitrite by capillary electrophoresis 884(2000)311

Nitrobenzenes

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Nitrobenzoyl chloride

Determination of sugar alcohols in confectioneries by high-performance liquid chromatography after nitrobenzoylation 893(2000)195

Nitro compounds

Gas chromatography–mass spectrometry–Fourier transform infrared spectrometry and high-performance liquid chromatographic characterization of mononitro- and dinitro-isomers from the nitration of *N,N*-dimethyldiphenylacetamide 878(2000)235

Nitro explosives

n-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Nitrogen

Potential for chlorate interference in ion chromatographic determination of total nitrogen in natural waters following alkaline persulfate digestion 857(1999)337

Gas chromatographic analysis of trace impurities in chlorine trifluoride 873(2000)229

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

Nitroimidazoles

Determination of four nitroimidazole residues in poultry meat by liquid chromatography–mass spectrometry 882(2000)89

Nitrophenols

Pressure-controlled high-performance liquid chromatographic study on the influence of rim chemistry on partial molar volume differences between free and complexed cyclodextrins 863(1999)127

New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169

Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235

Split-flow injector for capillary zone electrophoresis 883(2000)223

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Highly selective and efficient determination of US Environmental Protection Agency priority phenols employing solid-phase extraction and non-aqueous capillary electrophoresis 896(2000)95

Comparison of different sorbent materials for *on-line* solid-phase extraction with liquid chromatography–atmospheric pressure chemical ionization mass spectrometry of phenols 896(2000)159

2-Nitrophenylhydrazine hydrochloride

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Nitropyrenes

Determination of 1-nitropyrene in herbs after selective enrichment by a sol–gel-generated immunoaffinity column 880(2000)113

Determination of 1,3-, 1,6-, 1,8-dinitropyrene and 1-nitropyrene in airborne particulate by column liquid chromatography with electrochemical detection 897(2000)393

Nitrosodiethanolamine

Screening method for determining the presence of N-nitrosodiethanolamine in cosmetics by open-tubular capillary electrochromatography 887(2000)497

Nitrosodimethylaminophenol

Integration of a microextraction system. Solvent extraction of a Co–2-nitroso-5-dimethylaminophenol complex on a microchip 894(2000)19

Nitrotoluenes

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669

On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175

Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133

Nivalenol

Determination of nivalenol and deoxynivalenol in wheat using liquid chromatography–mass spectrometry with negative ion atmospheric pressure chemical ionisation 854(1999)45

Nomilin glucoside

Screening for limonoid glucosides in *Citrus tangerina* (Tanaka) Tseng by high-performance liquid chromatography–electrospray ionization mass spectrometry 874(2000)13

Nonsteroidal anti-inflammatory drugs

see also Profens

Determination of diclofenac sodium by capillary zone electrophoresis with electrochemical detection 868(2000)101

Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Development of a simple in-vial liquid-phase microextraction device for drug analysis compatible with capillary gas chromatography, capillary electrophoresis and high-performance liquid chromatography 873(2000)3

Solid-phase microextraction–gas chromatography–mass spectrometry of biologically active substances in water samples 873(2000)95

Development and robustness testing of a nonaqueous capillary electrophoresis method for the analysis of nonsteroidal anti-inflammatory drugs 874(2000)121

Determination of clonixin in plasma and urine by reversed-phase high-performance liquid chromatography 889(2000)135

Capillary electrochromatography and capillary electrochromatography–electrospray mass spectrometry for the separation of non-steroidal anti-inflammatory drugs 895(2000)123

Nonylphenol

Application of pressurized liquid extraction followed by gas chromatography–mass spectrometry to determine 4-nonylphenols in sediments 866(2000)79

Concentration of nonylphenol and its polyethoxylated derivatives by polymer-mediated extraction 891(2000)69

Nonylphenol ethoxylates

Characterization of nonyl phenol ethoxylates in sewage treatment plants by combined precursor ion scanning and multiple reaction monitoring 854(1999)245

Analysis of nonylphenol polyethoxycarboxylates and their related metabolites by on-line derivatization and ion-trap gas chromatography–mass spectrometry 862(1999)113

Assessment of the pollutant elimination efficiency by gas chromatography/mass spectrometry, liquid chromatography–mass spectrometry and –tandem mass spectrometry. Comparison of conventional and membrane-assisted biological wastewater treatment processes 889(2000)155

Norflloxacin

High-performance thin-layer chromatography method for monitoring norflloxacin residues on pharmaceutical equipment surfaces 862(1999)209

Notopterol

Preparative isolation and purification of notopterol and isoimperatorin from *Notopterygium forbesii* boiss (Chinese traditional medicinal herb) by high-speed counter-current chromatography 883(2000)67

NoXon-S

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

Nuclease

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Nucleic acids

see also DNA and RNA

Use of nucleic acids in the mobile phase for the determination of ascorbic acid in foods by high-performance liquid chromatography with electrochemical detection 881(2000)327

Specific applications of capillary electrochromatography to biopolymers, including proteins, nucleic acids, peptide mapping, antibodies, and so forth (Review) 887(2000)137

Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3

Nucleobase

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Nucleoside analogues

Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159

Nucleoside monophosphates

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

Nucleosides

Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry 856(1999)515

Determination at ppb level of an anti-human immunodeficiency virus nucleoside drug by capillary electrophoresis–electrospray ionization tandem mass spectrometry 895(2000)101

Nucleotides

see also Oligonucleotides and Polynucleotides

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry 856(1999)515

Chromatographic determination of riboflavin and its derivatives in food 881(2000)285

Development of a method for the analysis of nucleotides from the mantle tissue of the mussel *Mytilus galloprovincialis* 891(2000)99

Nylon

see Polyamides

Nystatin

Novel simulated moving-bed method for reduced solvent consumption 873(2000)149

O

Obacunone glucoside

Screening for limonoid glucosides in *Citrus tangerina* (Tanaka) Tseng by high-performance liquid chromatography–electrospray ionization mass spectrometry 874(2000)13

Ochratoxin A

Reversed-phase liquid chromatographic behavior of the mycotoxins, citrinin and ochratoxin A 862(1999)147

Determination of ochratoxin A in wine by means of immunoaffinity column clean-up and high-performance liquid chromatography 864(1999)89

Solid-phase extraction applied to the determination of ochratoxin A in wines by reversed-phase high-performance liquid chromatography 882(2000)29

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321

Octadecyl(dibutylhydroxyphenyl) propionate

Size-exclusion chromatography with on-line ultraviolet, proton nuclear magnetic resonance and mass spectrometric detection and on-line collection for off-line Fourier transform infrared spectroscopy 857(1999)89

***N*-Octyl bicycloheptene dicarboximide**

Chiral high-performance liquid chromatography of *N*-octyl bicycloheptene dicarboximide and confirmatory studies using liquid chromatography–tandem mass spectrometry and two-dimensional nuclear magnetic resonance spectroscopy 864(1999)271

4-*tert*-Octylphenol

Quantification of the xenoestrogens 4-*tert*-octylphenol and bisphenol A in water and in fish tissue based on microwave assisted extraction, solid-phase extraction and liquid chromatography–mass spectrometry 864(1999)17

Octylphenol ethoxylates

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

Octylthioglucoside

Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121

Okadaic acid

Liquid chromatography–electrospray ionization mass spectrometry of the diarrhetic shellfish-poisoning toxins okadaic acid, dinophysistoxin-1 and pectenotoxin-6 in bivalves 874(2000)199

Development and validation of a high-performance liquid chromatographic method using fluorimetric detection for the determination of the diarrhetic shellfish poisoning toxin okadaic acid without chlorinated solvents 876(2000)117

Oleuropein

Liquid chromatography with electrospray ionisation mass spectrometric detection of phenolic compounds from *Olea europaea* 855(1999)529

Oligofucans

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

Oligoguluronic acids

Analysis of oligomannuronic acids and oligoguluronic acids by high-performance anion-exchange chromatography and electrospray ionization mass spectrometry 884(2000)105

Oligomannuronic acids

Analysis of oligomannuronic acids and oligoguluronic acids by high-performance anion-exchange chromatography and electrospray ionization mass spectrometry 884(2000)105

Oligonucleotides

see also Nucleotides

Separation of oligonucleotides on novel monolithic columns with ion-exchange functional surfaces 852(1999)297

Nanoliter-scale sample preparation methods directly coupled to polymethylmethacrylate-based microchips and gel-filled capillaries for the analysis of oligonucleotides 853(1999)107

Application of capillary electrophoresis at low pH to oligonucleotides quality control 853(1999)355

Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry 856(1999)515

High-performance liquid chromatography separation methods for the analysis of peptide nucleic acids 864(1999)49

Sheath liquid effects in capillary high-performance liquid chromatography–electrospray mass spectrometry of oligonucleotides 870(2000)413

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis 871(2000)311

Specific applications of capillary electrochromatography to biopolymers, including proteins, nucleic acids, peptide mapping, antibodies, and so forth (Review) 887(2000)137

Purification of crude DNA oligonucleotides by solid-phase extraction and reversed-phase high-performance liquid chromatography 890(2000)167

Large-scale purification of antisense oligonucleotides by high-performance membrane adsorber chromatography 890(2000)179

Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23

Oligosaccharides

see also Saccharides

Liquid chromatography–electrospray mass spectrometry as a tool for the analysis of sulfated oligosaccharides from mucin glycoproteins 854(1999)131

Analysis of 8-aminonaphthalene-1,3,6-trisulfonate-derivatized oligosaccharides by capillary electrophoresis–electrospray ionization quadrupole ion trap mass spectrometry 858(1999)229

Efficiency studies in nonaqueous capillary electrophoresis 859(1999)203

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

Reinforcement of frontal affinity chromatography for effective analysis of lectin–oligosaccharide interactions 890(2000)261

Highly reproducible capillary zone electrophoresis of humic acids in cyclodextrin- or oligosaccharide-modified background electrolytes 895(2000)345

Oligourethanes

Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79

Opiates

Comparison of phenyl-type columns in the development of a fast liquid chromatographic system for eighteen opiates commonly found in forensic toxicology 896(2000)311

Opioids

Head-column field-amplified sample stacking in binary system capillary electrophoresis. Preparation of extracts for determination of opioids in microliter amounts of body fluids 853(1999)95

Kinetics of appearance of hemorphins from bovine hemoglobin peptic hydrolysates by a direct coupling of reversed-phase high-performance liquid chromatography and electrospray ionization mass spectrometry 873(2000)185

Analysis of codeine, dihydrocodeine and their glucuronides in human urine by electrokinetic capillary immunoassays and capillary electrophoresis–ion trap mass spectrometry 895(2000)133

Surface-ionization methods and devices of indication and identification of nitrogen-containing base molecules 896(2000)3

Organic acids

see also Carboxylic acids, Hydroxy acids and Fatty acids

Electrophoretic mobility study on ion–ion interactions in an aqueous solution 853(1999)63

Indirect detection of organic acids in non-aqueous capillary electrophoresis 853(1999)171

Determination of tartaric acid in solid wine residues by capillary electrophoresis and indirect UV detection 853(1999)181

- Use of volatile amines as ion-pairing agents for the high-performance liquid chromatographic–tandem mass spectrometric determination of aromatic sulfonates in industrial wastewater 854(1999)175
- Analysis of low-molecular-mass inorganic and organic anions by ion chromatography–atmospheric pressure ionization mass spectrometry 854(1999)275
- Liquid chromatographic–mass spectrometric determination of phenolic compounds using a capillary-scale particle beam interface 855(1999)515
- Ultra-short columns for low-pressure ion chromatography 857(1999)175
- Reversed-phase high-performance liquid chromatography of ionogenic compounds: comparison of retention models 859(1999)1
- Statistics aided optimization for high-performance liquid chromatographic analysis of organic acids in tobacco 859(1999)153
- Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1
- Chromatographic separation of 3,4-difluorophenylacetic acid and its positional isomers using five different techniques 866(2000)281
- Influence of pH*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63
- Erratum to: “Optimization of conductivity of low-molecular-mass anions in capillary zone electrophoresis” [J. Chromatogr. A, 850 (1999) 355–362] 868(2000)261
- Determination of aliphatic anhydrides and acids by reversed-phase liquid chromatography 868(2000)41
- Calculation of electrophoretic mobilities in water–organic modifier mixtures in capillary electrophoresis 868(2000)277
- Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23
- Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341
- Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391
- Gas chromatographic–mass spectrometric analysis of acids and phenols in distilled alcohol beverages. Application of anion-exchange disk extraction combined with in-vial elution and silylation 873(2000)29
- Separation of anions by ion chromatography–capillary electrophoresis 879(2000)245
- Electrophoretic methods in the analysis of beverages 880(2000)243
- Use of nucleic acids in the mobile phase for the determination of ascorbic acid in foods by high-performance liquid chromatography with electrochemical detection 881(2000)327
- Determination of organic acids in food samples by capillary zone electrophoresis 881(2000)357
- Development and validation of a capillary electrophoresis method for direct measurement of isocitric, citric, tartaric and malic acids as adulteration markers in orange juice 881(2000)395
- Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425
- Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629
- Detection techniques in ion analysis: what are our choices? 884(2000)3
- Retention behaviour of strong acid anions in ion-exclusion chromatography on sulfonate and carboxylate stationary phases 884(2000)61
- Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261
- Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297
- Determination of fumaric acid in apple juice by on-line coupled capillary isotachopheresis–capillary zone electrophoresis with UV detection 891(2000)175
- Novel, trifunctional diamine for silica coating in capillary zone electrophoresis 894(2000)53
- Capillary zone electrophoresis in phosphate buffer – known or unknown? 894(2000)89
- Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117
- Development and validation of a capillary electrophoresis method for the measurement of short-chain organic acids in natural rubber latex 894(2000)135

Simultaneous determination of inorganic anions and cations by capillary electrophoresis with indirect UV detection 895(2000)269

Characterization of sulfobutyl ether- β -cyclodextrins mixtures by anion-exchange chromatography using evaporative light scattering detection 897(2000)185

Preparation and evaluation of an imidazole-coated capillary column for the electrophoretic separation of aromatic acids 897(2000)383

Organic bases

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

Organoarsenic compounds

Speciation of dimethylarsinic acid and monomethylarsonic acid by solid-phase microextraction–gas chromatography–ion trap mass spectrometry 873(2000)129

Determination of inorganic and organic anionic arsenic species in water by ion chromatography coupled to hydride generation–inductively coupled plasma atomic emission spectrometry 884(2000)211

Evaluation of extraction procedures for the ion chromatographic determination of arsenic species in plant materials 889(2000)83

Organochlorine compounds

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

Multiresidue determination of persistent organochlorine and organophosphorus compounds in whale tissues using automated liquid chromatographic clean up and gas chromatographic–mass spectrometric detection 855(1999)633

On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

Separation and identification of perchlorinated polycyclic aromatic hydrocarbons by high-performance liquid chromatography and ultraviolet absorption spectroscopy 864(1999)173

Prediction of gas chromatographic retention indices of polychlorinated dibenzothiophenes on non-polar columns 866(2000)105

Development and characterization of chlorine-selective pulsed discharge emission detector for gas chromatography 872(2000)141

Applications using the chlorine-selective pulsed discharge emission detector 872(2000)167

Characterization of chlorinated compounds using a dual chlorine-selective pulsed discharge emission detector–helium-pulsed discharge photoionization detector system 872(2000)179

Extraction of DDT [1,1,1-trichloro-2,2-bis(*p*-chlorophenyl) ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane] from aged contaminated soil 874(2000)257

Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83

Simplified method for the determination of chlorinated fungicides and insecticides in fruits by gas chromatography 882(2000)193

Liquid chromatographic profiles of individual compounds of technical toxaphene 886(2000)123

Multi-residue methodology for pesticide screening in wines 889(2000)59

Monitoring organochlorine pesticides from landfill leachates by gas chromatography–electron-capture detection after solid-phase microextraction 891(2000)305

Solid-phase microextraction for organochlorine pesticide residues analysis in Chinese herbal formulations 898(2000)245

Organofluorine compounds

Analytical techniques used for monitoring the biodegradation of fluorinated compounds in waste streams from pharmaceutical production 889(2000)177

Organohalogen compounds

see also Halogenated compounds, Organochlorine compounds, Polychlorinated biphenyls, Polychlorinated dibenzodioxins, Polychlorinated dibenzofurans and Polychlorinated dibenzothiophenes

Pulsed discharge emission detector: an element-selective detector for gas chromatography 872(2000)119

Organomercury compounds

Gas chromatographic determination of organomercury following aqueous derivatization with sodium tetraethylborate and sodium tetraphenylborate. Comparative study of gas chromatography coupled with atomic fluorescence spectrometry, atomic emission spectrometry and mass spectrometry 876(2000)147

Application of chromatographic and electrophoretic methodology to the speciation of organomercury compounds in food analysis 882(2000)283

Experimental design of a microwave-assisted extraction–derivatization method for the analysis of methylmercury 889(2000)185

Organometallic compounds

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Organophosphorus compounds

Multiresidue determination of persistent organochlorine and organophosphorus compounds in whale tissues using automated liquid chromatographic clean up and gas chromatographic–mass spectrometric detection 855(1999)633

On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279

Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215

Stability of organophosphorus insecticides on graphitized carbon black extraction cartridges used for large volumes of surface water 879(2000)73

Herbicide and plant growth regulator analysis by capillary electrophoresis 880(2000)281

Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159

Multiresidue method for the rapid determination of organophosphorus insecticides in grapes, must and wine 882(2000)213

Multiple-technique analytical characterization of a mixture containing chemical-weapons simulant from a munition 883(2000)185

Multi-residue methodology for pesticide screening in wines 889(2000)59

Capillary electrophoretic separation of the enantiomers of organophosphates with a phosphorus stereogenic center using the sodium salt of octakis(2,3-diacetyl-6-sulfo)- γ -cyclodextrin as resolving agent 895(2000)247

Organoselenium compounds

High-performance liquid chromatography of selenium compounds utilizing perfluorinated carboxylic acid ion-pairing agents and inductively coupled plasma and electrospray ionization mass spectrometric detection 866(2000)51

Column-switching system for selenium speciation by coupling reversed-phase and ion-exchange high-performance liquid chromatography with microwave-assisted digestion–hydride generation–atomic fluorescence spectrometry 889(2000)33

Organosilane compounds

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Organosilicons, cyclic

Low-carbon silica sorbents for solid-phase extraction 868(2000)1

Organosulfur compounds

see also Sulfur compounds

On the performance and inertness of different materials used for the enrichment of sulfur compounds from air and gaseous samples 864(1999)345

Prediction of gas chromatographic retention indices of polychlorinated dibenzothiophenes on non-polar columns 866(2000)105

Analysis of organic sulfur compounds in wine aroma 881(2000)569

Organotin compounds

Analysis of organotin compounds by gas chromatography–reactive-flow detection 858(1999)245

Determination of butyltin species in water and sediment by solid-phase microextraction–gas chromatography–flame ionization detection 873(2000)63

Determination of butyltin and phenyltin species by reversed-phase liquid chromatography and fluorimetric detection 878(2000)69

Speciation of butyl- and phenyltin compounds in sediments using pressurized liquid extraction and liquid chromatography–inductively coupled plasma mass spectrometry 879(2000)137

Direct Grignard pentylation of organotin-contaminated lard samples followed by capillary gas chromatography with flame photometric detection 886(2000)197

Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149

Ormeloxifene

Fast chiral separations using sulfated β -cyclodextrin and short-end injection in capillary electrophoresis 875(2000)403

Ovalbumin

Suspended bed chromatography, a new approach in downstream processing 890(2000)45

Ovoglycoprotein

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VII. Comparison of chiral recognition ability of ovoglycoprotein from chicken and Japanese quail egg whites 858(1999)155

Oxacillin

Application of ion-exchange cartridge clean-up in food analysis II. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in meat using liquid chromatography with ultraviolet detection 855(1999)247

Oxalate

Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629

Oxalyldiaminopropionic acid

Analysis of β -*N*-oxalyl-L- α , β -diaminopropionic acid and homoarginine in *Lathyrus sativus* by capillary zone electrophoresis 857(1999)295

Determination of neurotoxin 3-*N*-oxalyl-2,3-diaminopropionic acid and non-protein amino acids in *Lathyrus sativus* by precolumn derivatization with 1-fluoro-2,4-dinitrobenzene 883(2000)113

Oxanine

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Oxocarboxylates

Comparative methodology in the determination of α -oxocarboxylates in aqueous solution. Ion chromatography versus gas chromatography after oximation, extraction and esterification 867(2000)143

Oxygen

Gas chromatographic analysis of trace impurities in chlorine trifluoride 873(2000)229

Oxyhalides

Analysis of low-molecular-mass inorganic and organic anions by ion chromatography–atmospheric pressure ionization mass spectrometry 854(1999)275

Oxyhalogens

Determination of bromide, bromate and other anions with ion chromatography and an inductively coupled plasma mass spectrometer as element-specific detector 862(1999)39

Oxytocin

Recognition of oxytocin and oxytocin-related peptides in aqueous media using a molecularly imprinted polymer synthesized by the epitope approach 889(2000)111

Ozone

Separation and identification of polar degradation products of benzo[*a*]pyrene with ozone by atmospheric pressure chemical ionization–mass spectrometry after optimized column chromatographic clean-up 855(1999)501

Application of gas chromatography–cryocondensation–Fourier transform infrared spectroscopy and gas chromatography–mass spectrometry to the identification of gas phase reaction products from the α -pinene/ozone reaction 864(1999)299

P

Paclitaxel

Optimization of throughput and desorbent consumption in simulated moving-bed chromatography for paclitaxel purification 855(1999)71

Palladium

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Palladium–gold

Synthesis and study of the cluster $[\text{Pd}(\text{AuPPh}_3)_6(\text{HgNO}_3)](\text{NO}_3)$ by reversed-phase high-performance liquid chromatography, thermoanalysis, infrared, nuclear magnetic resonance, and fast atom bombardment mass spectrometry 862(1999)29

Palmatine

Analysis of coptisine, berberine and palmatine in adulterated Chinese medicine by capillary electrophoresis–electrospray ion trap mass spectrometry 866(2000)273

Paracetamol

Simultaneous determination of paracetamol and chlorpheniramine in human plasma by liquid chromatography–tandem mass spectrometry 870(2000)77

Parathyroid hormone

Analysis of human parathyroid hormone (1–84) products. Separation of a major impurity in synthetic products by ion-pairing reversed-phase high-performance liquid chromatography 890(2000)127

Patulin

Chromatographic determination of the mycotoxin patulin in fruit and fruit juices 882(2000)17

Determination of patulin in apple juice by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 897(2000)365

Pectenotoxin-6

Liquid chromatography–electrospray ionization mass spectrometry of the diarrhetic shellfish-poisoning toxins okadaic acid, dinophysistoxin-1 and pectenotoxin-6 in bivalves 874(2000)199

Pectins

Chiral selectors from fruit: application of citrus pectins to enantiomer separations in capillary electrophoresis 857(1999)285

High-performance liquid chromatography of methanol released from pectins after its oxidation to formaldehyde and condensation with 2,4-dinitrophenylhydrazine 863(1999)227

Penicillins

Application of ion-exchange cartridge clean-up in food analysis. III. Determination of benzylpenicillin, phenoxymethyl-penicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in bovine liver and kidney by liquid chromatography with ultraviolet detection 880(2000)85

Stability of penicillin antibiotic residues in meat during storage. Ampicillin 882(2000)135

New configurations and applications of molecularly imprinted polymers 889(2000)15

Pentachlorophenol

Pressurised liquid extraction of pentachlorophenol from dyestuffs and cotton 873(2000)287

Pentafluorobenzylhydroxylamine

Microwave-assisted derivatization of volatile carbonyl compounds with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine 891(2000)267

Pentanol

Comparison of the performance of butanol and pentanol as modifiers in the micellar chromatographic determination of some phenethylamines 866(2000)35

Peptaibiotics

Detection of new sequences of peptaibol antibiotics trichothoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179

C-Peptide

Quantitative analysis of urinary C-peptide by liquid chromatography–tandem mass spectrometry with a stable isotopically labelled internal standard 896(2000)275

Peptidases

Analysis of angiotension-converting enzyme by capillary electrophoresis 853(1999)185

Peptide nucleic acids

High-performance liquid chromatography separation methods for the analysis of peptide nucleic acids 864(1999)49

Peptides

see also Special Issue Index

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

New polymer-based preppacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

Application of a peptide bank from porcine brain in isolation of regulatory peptides 852(1999)273

Proteolytic cleavage of glucagon-like peptide-1 by pancreatic β cells and by fetal calf serum analyzed by mass spectrometry 852(1999)285

Separation of proteins and peptides by capillary electrophoresis in acid buffers containing high concentrations of surfactants 852(1999)325

General experimental aspects of the use of isoelectric buffers in capillary electrophoresis 853(1999)71

Microscale characterization of the structure–activity relationship of a heparin-binding glycopeptide using affinity capillary electrophoresis and immobilized enzymes 853(1999)189

Development and validation of a capillary electrophoresis method for the characterization of protegrin IB-367 853(1999)197

Phosphopeptide analysis by on-line immobilized metal-ion affinity chromatography–capillary electrophoresis–electrospray ionization mass spectrometry 853(1999)225

Capillary electrochromatography of peptides in a microfabricated system 853(1999)257

- Separation of selected peptides by capillary electroosmotic chromatography using 3 μm reversed-phase bonded silica and mixed-mode phases 853(1999)263
- Analysis of leptin gene expression in chickens using reverse transcription polymerase chain reaction and capillary electrophoresis with laser-induced fluorescence detection 853(1999)321
- Interaction of cyclophilin and cyclosporins monitored by affinity capillary electrophoresis 853(1999)469
- Preliminary comparison of precursor scans and liquid chromatography–tandem mass spectrometry on a hybrid quadrupole time-of-flight mass spectrometer 854(1999)119
- Capillary electrochromatography with a silica column with a dynamically modified cationic surfactant 855(1999)137
- Capillary columns with in situ formed porous monolithic packing for micro high-performance liquid chromatography and capillary electrochromatography 855(1999)273
- Analysis of recombinant and modified proteins by capillary zone electrophoresis coupled with electrospray ionization tandem mass spectrometry 855(1999)695
- Recent developments in microcolumn liquid chromatography 856(1999)117
- Studies on the relationship between structure and electrophoretic mobility of α -helical and β -sheet peptides using capillary zone electrophoresis 857(1999)263
- Capillary electrochromatography of proteins and peptides with porous-layer open-tubular columns 858(1999)91
- Chiral separations using polymeric dipeptide surfactants: effect of number of chiral centers and steric factors 858(1999)219
- Glutathione oxidation in real time by thermospray liquid chromatography–mass spectrometry 862(1999)161
- Detection of new sequences of peptaibol antibiotics trichothoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179
- Natural poly-histidine affinity tag for purification of recombinant proteins on cobalt(II)-carboxymethylaspartate crosslinked agarose 864(1999)247
- Rapid reversed-phase separation of proteins and peptides using optimized ‘moulded’ monolithic poly(styrene-*co*-divinylbenzene) columns 865(1999)169
- Reversed-phase liquid chromatography of proteins and peptides using multimodal copolymer-encapsulated silica 866(2000)1
- Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151
- Separation of peptides by strong cation-exchange capillary electrochromatography 869(2000)385
- Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255
- Signature-peptide approach to detecting proteins in complex mixtures 870(2000)295
- Separation and characterization of multicomponent peptide mixtures by liquid chromatography–electrospray ionization mass spectrometry. Application to crude products of the synthesis of leuprolide 870(2000)315
- Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335
- Simultaneous separation of angiotensin and endothelin peptide families by high-performance liquid chromatography: application to the specific radioimmunoassay measurement of angiotensin II or endothelin-1 from tissue 870(2000)349
- Liquid chromatography–electrospray time-of-flight mass spectrometry for on-line accurate mass determination and identification of cyclodepsipeptides in a crude extract of the fungus *Metarrhizium anisopliae* 872(2000)85
- Kinetics of appearance of hemorphins from bovine hemoglobin peptic hydrolysates by a direct coupling of reversed-phase high-performance liquid chromatography and electrospray ionization mass spectrometry 873(2000)185
- Chiral separation of amino acids and peptides by capillary electrophoresis 875(2000)43
- Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255
- Modified capillary electrophoresis system for peptide, protein and double-stranded DNA analysis 879(2000)189
- Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides 881(2000)1
- Isolation and preparative purification of microcystin variants 883(2000)103
- Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

- Specific applications of capillary electrochromatography to biopolymers, including proteins, nucleic acids, peptide mapping, antibodies, and so forth 887(2000)137
- Capillary electroosmotic chromatography of peptides 887(2000)165
- Capillary electrochromatography of proteins and peptides with a cationic acrylic monolith 887(2000)465
- Analysis of isomeric glutamyl peptides by capillary electrophoresis. Application to stability studies 888(2000)275
- Recognition of oxytocin and oxytocin-related peptides in aqueous media using a molecularly imprinted polymer synthesized by the epitope approach 889(2000)111
- Liquid chromatography–electrospray mass spectrometry of multicomponent peptide mixtures. Characterization of a mixture from the synthesis of the hormone goserelin 889(2000)119
- Trapping the monomeric α -helical state during unfolding of coiled-coils by reversed-phase liquid chromatography 890(2000)81
- Use of Edman degradation sequence analysis and matrix-assisted laser desorption/ionization mass spectrometry in designing substrates for matrix metalloproteinases 890(2000)117
- Analysis of human parathyroid hormone (1–84) products. Separation of a major impurity in synthetic products by ion-pairing reversed-phase high-performance liquid chromatography 890(2000)127
- Analysis of neuropeptide Y and its metabolites by high-performance liquid chromatography–electrospray ionization mass spectrometry and integrated sample clean-up with a novel restricted-access sulphonic acid cation exchanger 890(2000)135
- Characterisation of the rubber elongation factor from ammoniated latex by electrophoresis and mass spectrometry 890(2000)145
- Determination of enantiomers in a synthetic argininal peptide using capillary zone electrophoresis and high-performance liquid chromatography 891(2000)115
- Approaches towards the quantitative analysis of peptides and proteins by reversed-phase high-performance liquid chromatography in the absence of a pure reference sample 891(2000)235
- Novel microfabricated device for electrokinetically induced pressure flow and electrospray ionization mass spectrometry 892(2000)195
- Applications of mass spectrometry to food proteins and peptides 893(2000)1
- Development of simultaneous purification methodology for multiple synthetic peptides by reversed-phase sample displacement chromatography 893(2000)81
- Analysis of a peptide hormone mixture of therapeutic interest by liquid chromatography coupled to high-flow pneumatically assisted electrospray mass spectrometry 893(2000)95
- Liquid chromatographic fractionation of small peptides from wine 893(2000)281
- Influence of temperature on the behaviour of small linear peptides in capillary electrochromatography 893(2000)401
- Comparative study of capillary electroosmotic chromatography and electrically assisted gradient nano-liquid chromatography for the separation of peptides 894(2000)241
- Influence of the structure of cyclodextrins and amino acid sequence of dipeptides and tripeptides on the pH-dependent reversal of the migration order in capillary electrophoresis 894(2000)267
- The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry 894(2000)345
- Capillary electrophoresis and off-line capillary electrophoresis–electrospray ionization quadrupole time-of-flight tandem mass spectrometry of carbohydrates 895(2000)291
- Use of chiral liquid chromatography–tandem mass spectrometry to investigate the metabolism of racemic cholecystokinin-B antagonists 896(2000)217
- Quantitative analysis of urinary C-peptide by liquid chromatography–tandem mass spectrometry with a stable isotopically labelled internal standard 896(2000)275
- Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83
- Perchlorate**
Improved method for the determination of trace perchlorate in ground and drinking waters by ion chromatography 888(2000)151
- Determination of perchlorate at parts-per-billion levels in plants by ion chromatography 898(2000)193
- Perfluorocarboxylic acids**
Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245
- Perfumery compounds**
Analysis of the equations for the temperature dependence of the retention index. I. Relation between equations 858(1999)65

Periodate

Determination of cyclodextrin content using periodate oxidation by capillary electrophoresis 891(2000)201

Permanganate

Capillary electrophoresis separation and permanganate chemiluminescence on-line detection of some alkaloids with β -cyclodextrin as an additive 855(1999)329

Peroxidase

Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45

Peroxides

Determination of gas phase peroxyacetic acid using pre-column derivatization with organic sulfide reagents and liquid chromatography 855(1999)551

Determination of peroxides by capillary zone electrophoresis with amperometric detection 868(2000)127

Peroxyacetic acid

Determination of gas phase peroxyacetic acid using pre-column derivatization with organic sulfide reagents and liquid chromatography 855(1999)551

Post-column oxidative derivatization for the liquid chromatographic determination of phenothiazines 890(2000)281

Persulfates

Determination of peroxides by capillary zone electrophoresis with amperometric detection 868(2000)127

Perylene

Video-taped sample evaporation in hot chambers simulating gas chromatography split/splitless injectors. I. Thermospray injection 897(2000)237

Video-taped sample evaporation in hot chambers simulating gas chromatography split and splitless injectors. II. Injection with band formation 897(2000)247

Pesticides

see also Special Issue Index

Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

Simultaneous determination of antifouling herbicides in marina water samples by on-line solid-phase extraction followed by liquid chromatography–mass spectrometry 854(1999)197

Atmospheric pressure ionisation multiple mass spectrometric analysis of pesticides 854(1999)207

Multiresidue determination of persistent organochlorine and organophosphorus compounds in whale tissues using automated liquid chromatographic clean up and gas chromatographic–mass spectrometric detection 855(1999)633

On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279

Ultra-trace-level determination of polar pesticides and their transformation products in surface and estuarine water samples using column liquid chromatography–electrospray tandem mass spectrometry 857(1999)157

Liquid chromatographic determination of imazosulfuron in drinking water and in soil using ultraviolet detection 857(1999)327

Capillary electrophoretic determination of ferric dimethyldithiocarbamate as iron(III) chelate of EDTA 857(1999)365

Semi-permeable surface analytical reversed-phase column for the improved trace analysis of acidic pesticides in water with coupled-column reversed-phase liquid chromatography with UV detection. Determination of bromoxynil and bentazone in surface water 858(1999)45

Inclusion of the fungicide chlozolinate in a multiresidue method 858(1999)123

Analysis of some pesticides in water samples using solid-phase microextraction–gas chromatography with different mass spectrometric techniques 859(1999)193

Supercritical fluid extraction of chlorpyrifos and 3,5,6-trichloro-2-pyridinol from garden compost 862(1999)105

Determination of polar pesticides with atmospheric pressure chemical ionisation mass spectrometry–mass spectrometry using methanol and/or acetonitrile for solid-phase desorption and gradient liquid chromatography 863(1999)147

Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121

Soil thin-layer chromatography and pesticide mobility through soil microstructures. New technical approach 864(1999)145

New gas chromatographic method for residue determination of dithiopyr in soil, wheat grain and straw 864(1999)351

Determination of pesticides in waters by automatic on-line solid-phase extraction–capillary electrophoresis 866(2000)137

Tandem solid-phase extraction of atrazine ozonation products in water 866(2000)195

- Pesticide trace analysis using solid-phase extraction and gas chromatography with electron-capture and tandem mass spectrometric detection in water samples 867(2000)235
- Low-carbon silica sorbents for solid-phase extraction 868(2000)1
- Does further clean-up reduce the matrix enhancement effect in gas chromatographic analysis of pesticide residues in food? 868(2000)51
- Determination of interfering triazine degradation products by gas chromatography–ion trap mass spectrometry 868(2000)115
- Comparison of gas chromatography–pulsed flame photometric detection–mass spectrometry, automated mass spectral deconvolution and identification system and gas chromatography–tandem mass spectrometry as tools for trace level detection and identification 868(2000)229
- Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411
- Analysis of pesticide residues in matrices with high lipid contents by membrane separation coupled on-line to a high-performance liquid chromatography system 869(2000)427
- On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441
- Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451
- Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463
- Evaluation of surface- and ground-water pollution due to herbicides in agricultural areas of Zamora and Salamanca (Spain) 869(2000)471
- Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481
- Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487
- Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497
- Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43
- Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57
- Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67
- Applications using the chlorine-selective pulsed discharge emission detector 872(2000)167
- Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215
- Trace-level determination of triazines and several degradation products in environmental waters by disk solid-phase extraction and micellar electrokinetic chromatography 872(2000)299
- Automated in-tube solid-phase microextraction–high-performance liquid chromatography for carbamate pesticide analysis 873(2000)137
- Trace analysis of pesticide residues in water by high-speed narrow-bore capillary gas chromatography–mass spectrometry with programmable temperature vaporizer 874(2000)81
- Solid-phase extraction followed by high-performance liquid chromatography–ionspray interface–mass spectrometry for monitoring of herbicides in environmental water 874(2000)187
- Determination of triazines in soil leachates by solid-phase microextraction coupled to gas chromatography–mass spectrometry 874(2000)247
- Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl) ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane] from aged contaminated soil 874(2000)257
- Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207
- Optimization of the cyclodextrin-assisted capillary electrophoresis separation of the enantiomers of phenoxyacid herbicides 875(2000)423
- Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133
- Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

- Determination of the plant growth regulator chlormequat in food by liquid chromatography–electrospray ionisation tandem mass spectrometry 878(2000)77
- Simultaneous determination of acidic and non-acidic pesticides in natural waters by liquid chromatography–mass spectrometry 878(2000)87
- Capillary zone electrophoretic separation of neutral species of chloro-*s*-triazines in the presence of cationic surfactant monomers 878(2000)137
- Gas chromatography–mass spectrometry–Fourier transform infrared spectrometry and high-performance liquid chromatographic characterization of mononitro- and dinitro-isomers from the nitration of *N,N*-dimethyldiphenylacetamide 878(2000)235
- Determination of rice herbicides, their transformation products and clofibric acid using on-line solid-phase extraction followed by liquid chromatography with diode array and atmospheric pressure chemical ionization mass spectrometric detection 879(2000)3
- Monitoring of priority pesticides and other organic pollutants in river water from Portugal by gas chromatography–mass spectrometry and liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)13
- Part-per-trillion level determination of antifouling pesticides and their byproducts in seawater samples by off-line solid-phase extraction followed by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)27
- Comparison of various liquid chromatographic methods involving UV and atmospheric pressure chemical ionization mass spectrometric detection for the efficient trace analysis of phenylurea herbicides in various types of water samples 879(2000)39
- Application of liquid chromatography with mass spectrometry combined with photodiode array detection and tandem mass spectrometry for monitoring pesticides in surface waters 879(2000)51
- Stability of organophosphorus insecticides on graphitized carbon black extraction cartridges used for large volumes of surface water 879(2000)73
- Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83
- Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121
- Applications of solid-phase microextraction in food analysis 880(2000)35
- Applications of matrix solid-phase dispersion in food analysis 880(2000)63
- Determination of fenpyroximate in apples by supercritical fluid extraction and packed capillary liquid chromatography with UV detection 880(2000)121
- Thin-layer chromatography in food and agricultural analysis 880(2000)129
- Herbicide and plant growth regulator analysis by capillary electrophoresis 880(2000)281
- Pesticide residue determination in fruit and vegetables by liquid chromatography–mass spectrometry 882(2000)153
- Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175
- Simplified method for the determination of chlorinated fungicides and insecticides in fruits by gas chromatography 882(2000)193
- Multiresidue analysis of seventeen pesticides in wine by gas chromatography with mass-selective detection 882(2000)205
- Multiresidue method for the rapid determination of organophosphorus insecticides in grapes, must and wine 882(2000)213
- Multiresidue method for the rapid determination – in grape, must and wine – of fungicides frequently used on vineyards 882(2000)221
- Interlaboratory study of identification and quantitation of multiresidue pyrethroids in agricultural products by gas chromatography–mass spectrometry 882(2000)231
- Gas chromatographic determination of acrinathrine and 3-phenoxybenzaldehyde residues in honey 882(2000)239
- Chromatographic characterization of molecularly imprinted polymers binding the herbicide 2,4,5-trichlorophenoxyacetic acid 883(2000)119
- Use of ion chromatography–electrospray mass spectrometry for the determination of ionic compounds in agricultural chemicals 884(2000)23
- Determination of some pesticides and intermediates by ion chromatography 884(2000)243
- Investigation into the effects of temperature and stirring rate on the solid-phase extraction of diuron from water using a C₁₈ extraction disk 885(2000)41
- Graphitized carbons for solid-phase extraction 885(2000)73

- Multiresidue methods using solid-phase extraction techniques for monitoring priority pesticides, including triazines and degradation products, in ground and surface waters 885(2000)217
- Solid-phase extraction of acidic herbicides 885(2000)237
- Solid-phase extraction of quaternary ammonium herbicides 885(2000)251
- On-line dual-precolum-based trace enrichment for the determination of polar and acidic microcontaminants in river water by liquid chromatography with diode-array UV and tandem mass spectrometric detection 885(2000)377
- Solid-phase microextraction in pesticide residue analysis 885(2000)389
- Liquid chromatographic profiles of individual compounds of technical toxaphene 886(2000)123
- New methods for determination of glyphosate and (aminomethyl)phosphonic acid in water and soil 886(2000)207
- Analysis of flufenacet in soil, wheat grain and straw by gas chromatography 886(2000)319
- Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 888(2000)113
- Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197
- Solid-phase extraction with a dibutylmelamine-imprinted polymer as triazine herbicide-selective sorbent 889(2000)25
- Multi-residue methodology for pesticide screening in wines 889(2000)59
- Extraction of glyphosate by a supported liquid membrane technique 889(2000)93
- Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261
- Indirect analysis of urea herbicides from environmental water using solid-phase microextraction 890(2000)303
- Analysis of agrochemicals by capillary electrophoresis (Review) 891(2000)45
- Determination of chlorophacinone and diphacinone in commercial rodenticides by liquid chromatography–UV detection and liquid chromatography–electrospray ionization mass spectrometry 891(2000)249
- Monitoring organochlorine pesticides from landfill leachates by gas chromatography–electron-capture detection after solid-phase microextraction 891(2000)305
- Reversed migration micellar electrokinetic chromatography with off-line and on-line concentration analysis of phenylurea herbicides 892(2000)187
- Pesticide residue analysis in foodstuffs applying capillary gas chromatography with mass spectrometric detection. State-of-the-art use of modified DFG-multimethod S19 and automated data evaluation 892(2000)347
- Determination of pesticides in vegetables using large-volume injection column liquid chromatography–electrospray tandem mass spectrometry 892(2000)379
- Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435
- Determination of fungicides in natural waters using solid-phase microextraction and gas chromatography coupled with electron-capture and mass spectrometric detection 893(2000)143
- High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157
- Solid-phase microextraction for the determination of systemic and non-volatile pesticides in river water using gas chromatography with nitrogen–phosphorous and electron-capture detection 893(2000)347
- On-line coupling of solid-phase extraction to gas chromatography with fast solvent vaporization and concentration in an open injector liner. Analysis of pesticides in aqueous samples 896(2000)41
- Liquid chromatographic determination of emamectin, milbemectin, ivermectin and abamectin in crops and confirmation by liquid chromatography–mass spectrometry 896(2000)75
- Determination of aqueous chlorothalonil with solid-phase microextraction and gas chromatography 896(2000)105
- Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111
- Optimisation of alachlor solid-phase microextraction from water samples using experimental design 896(2000)373
- Solid-phase microextraction for organochlorine pesticide residues analysis in Chinese herbal formulations 898(2000)245
- Phenanthridinone**
Sample cleanup and reversed-phase high-performance liquid chromatographic analysis of polar aromatic compounds in groundwater samples from a former gas plant 862(1999)137

Phenethylamines

Comparison of the performance of butanol and pentanol as modifiers in the micellar chromatographic determination of some phenethylamines 866(2000)35

Phenolic acids

Simultaneous determination of phenolic acids and 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one in wheat (*Triticum aestivum* L.) by gas chromatography–tandem mass spectrometry 864(1999)315

Determination of phenolic acids in fruit juices by isocratic column liquid chromatography 891(2000)183

Phenolic compounds

Electrophoretic mobility study on ion–ion interactions in an aqueous solution 853(1999)63

Liquid chromatographic–mass spectrometric determination of phenolic compounds using a capillary-scale particle beam interface 855(1999)515

Liquid chromatography with electrospray ionisation mass spectrometric detection of phenolic compounds from *Olea europaea* 855(1999)529

Analysis of pigmented high-molecular-mass grape phenolics using ion-pair, normal-phase high-performance liquid chromatography 866(2000)25

Determination of chlorogenic acid in rat blood by microdialysis coupled with microbore liquid chromatography and its application to pharmacokinetic studies 870(2000)443

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Optimization of the separation of phenolic compounds by micellar electrokinetic capillary chromatography 871(2000)427

On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic–diode array detection–mass spectrometric techniques 880(2000)203

Electrophoretic methods in the analysis of beverages 880(2000)243

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Determination of organic acids in food samples by capillary zone electrophoresis 881(2000)357

Determination of tea catechins 881(2000)411

Analysis of several phenolic compounds with potential antioxidant properties in grape extracts and wines by high-performance liquid chromatography–photodiode array detection without sample preparation 881(2000)461

Determination of synthetic phenolic antioxidants in food by high-performance liquid chromatography 882(2000)271

Optimization strategy and validation of one chromatographic method as approach to determine the phenolic compounds from different sources 897(2000)161

Phenols

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudostationary phase. Influence of methanol and acetonitrile on separation selectivity 853(1999)121

Efficient solid-phase extraction procedures for trace enrichment of priority phenols from industrial effluents with high total organic carbon content 857(1999)97

Synthesis and analytical properties of a novel columnar metallomesogenic polymer 862(1999)65

Trapping efficiency of aqueous pollutants in multichannel thick-film silicone-rubber traps for capillary gas chromatography 863(1999)57

Pressure-controlled high-performance liquid chromatographic study on the influence of rim chemistry on partial molar volume differences between free and complexed cyclodextrins 863(1999)127

Gas chromatographic profiling and screening for phenols as isobutoxycarbonyl derivatives in aqueous samples 866(2000)87

Dissociation constants of phenols in methanol–water mixtures 867(2000)45

Determination of phenols using simultaneous steam distillation–extraction 867(2000)281

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Determination of phenols in wines by liquid chromatography with photodiode array and fluorescence detection 871(2000)85

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

- Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357
- On-column amperometric detection in capillary electrophoresis with an improved high-voltage electric field decoupler 872(2000)259
- Gas chromatographic–mass spectrometric analysis of acids and phenols in distilled alcohol beverages. Application of anion-exchange disk extraction combined with in-vial elution and silylation 873(2000)29
- New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169
- Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235
- Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97
- Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121
- Preparation of 4,4'-(1-[²H₆]methylene)bis-[2,3,5,6-²H₄]-phenol and its application to the measurement of bisphenol A in beverages by stable isotope dilution mass spectrometry 883(2000)163
- Solid-phase extraction of phenols 885(2000)291
- Interpretive optimisation strategy applied to the isocratic separation of phenols by reversed-phase liquid chromatography with acetonitrile–water and methanol–water mobile phases 886(2000)31
- Qualitative and quantitative analysis of a thermoset polymer, poly(benzoxazine), by pyrolysis–gas chromatography 886(2000)217
- Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251
- Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3
- High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157
- Separation of bisphenol A and three alkylphenols by micellar electrokinetic chromatography 895(2000)213
- Separation of two groups of oestrogen mimicking compounds using micellar electrokinetic chromatography 895(2000)237
- Highly selective and efficient determination of US Environmental Protection Agency priority phenols employing solid-phase extraction and non-aqueous capillary electrophoresis 896(2000)95
- Capillary electrophoresis–electrospray ion-trap mass spectrometry for the separation of chlorophenols 896(2000)125
- Comparison of different sorbent materials for *on-line* solid-phase extraction with liquid chromatography–atmospheric pressure chemical ionization mass spectrometry of phenols 896(2000)159
- Analysis of phenols in pyrolysis oils by gel permeation chromatography and multidimensional liquid chromatography 896(2000)343
- Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279
- Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23
- Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111
- Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133
- Phenothiazines**
Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313
- Post-column oxidative derivatization for the liquid chromatographic determination of phenothiazines 890(2000)281
- Phenoxy acids**
Determination of polar pesticides with atmospheric pressure chemical ionisation mass spectrometry–mass spectrometry using methanol and/or acetonitrile for solid-phase desorption and gradient liquid chromatography 863(1999)147
- Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207
- Optimization of the cyclodextrin-assisted capillary electrophoresis separation of the enantiomers of phenoxyacid herbicides 875(2000)423

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111

Phenoxyalkanoic acids

Determination of phenoxyalkanoic acids and other herbicides at the ng/ml level in water by solid-phase extraction with poly(divinylbenzene-co-*N*-vinylpyrrolidone) sorbent and high-performance liquid chromatography–diode-array detection 867(2000)169

Phenoxybenzaldehyde

Gas chromatographic determination of acrinathrine and 3-phenoxybenzaldehyde residues in honey 882(2000)239

Phenoxymethylpenicillin

Application of ion-exchange cartridge clean-up in food analysis II. Determination of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in meat using liquid chromatography with ultraviolet detection 855(1999)247

Phenylacetic acid

Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117

Phenylalanine

Apparatus and method for interfacing capillary electrophoresis and chemiluminescence nitrogen detection for the analysis of nitrogen-containing compounds 868(2000)121

Electron–capture mass spectrometry: recent advances 892(2000)329

Phenylalanine anilide

Application of molecularly imprinted polymers in supercritical fluid chromatography 897(2000)317

Phenylamines

Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

(Phenylamino)alanine

Tetrahydro- β -carboline-3-carboxylic acids and contaminants of L-tryptophan 881(2000)501

Phenylborate

Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285

2-Phenylethanol

Calculation of chromatographic band profiles with an implicit isotherm 862(1999)1

α -Phenylethylamine

Convenient synthesis of π -acceptor chiral stationary phases for high-performance liquid chromatography from halogen-substituted 3,5-dinitrobenzoylamides 859(1999)143

1-Phenyl-1-propanol

Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95

Enantioseparation of 1-phenyl-1-propanol by simulated moving bed under linear and nonlinear conditions 893(2000)307

3-Phenylpropanol

Calculation of chromatographic band profiles with an implicit isotherm 862(1999)1

Phenylureas

Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465

Determination of polar pesticides with atmospheric pressure chemical ionisation mass spectrometry–mass spectrometry using methanol and/or acetonitrile for solid-phase desorption and gradient liquid chromatography 863(1999)147

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Comparison of various liquid chromatographic methods involving UV and atmospheric pressure chemical ionization mass spectrometric detection for the efficient trace analysis of phenylurea herbicides in various types of water samples 879(2000)39

Reversed migration micellar electrokinetic chromatography with off-line and on-line concentration analysis of phenylurea herbicides 892(2000)187

Pheophytin a

Improved liquid chromatographic method for the analysis of photosynthetic pigments of higher plants 876(2000)111

Pheromones

Use of solid-phase microextraction in the investigation of chemical communication in social wasps 873(2000)73

Role of solid-phase microextraction in the identification of highly volatile pheromones of two Rhinoceros beetles *Scapanes australis* and *Strategus aloeus* (Coleoptera, Scarabaeidae, Dynastinae) 885(2000)433

Phloroglucinols

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

Phloxines

Rapid quantification of hexachlorobenzene in the color additives D&C Red Nos. 27 and 28 (phloxine B) using solid-phase microextraction and gas chromatography–mass spectrometry 863(1999)37

Pholcodine

Micellar electrokinetic and high-performance liquid chromatographic determination of potential manufacturing impurities in pholcodine 895(2000)9

Phorbol

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

Phosphatases

On-line automated high-performance liquid chromatographic determination of total riboflavin phosphates using immobilized acid phosphatase as a pre-column reactor 896(2000)171

Phosphates

Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121

Analysis of condensed phosphates in food products by ion chromatography with an on-line hydroxide eluent generator 881(2000)639

Capillary electrophoretic separation of the enantiomers of organophosphates with a phosphorus stereogenic center using the sodium salt of octakis(2,3-diacetyl-6-sulfo)- γ -cyclodextrin as resolving agent 895(2000)247

On-line automated high-performance liquid chromatographic determination of total riboflavin phosphates using immobilized acid phosphatase as a pre-column reactor 896(2000)171

Phosphatidylethanolamines

Separation of the molecular species of intact phosphatidylethanolamines and their *N*-monomethyl and *N,N*-dimethyl derivatives by high-performance liquid chromatography on a C_8 column 891(2000)349

Phosphines

Synthesis and study of the cluster $[Pd(AuPPh_3)_6(HgNO_3)](NO_3)$ by reversed-phase high-performance liquid chromatography, thermoanalysis, infrared, nuclear magnetic resonance, and fast atom bombardment mass spectrometry 862(1999)29

Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159

Phosphinic acids

Determination of optical purity of phosphonic acid analogues of aromatic amino acids by capillary electrophoresis with α -cyclodextrin 895(2000)301

Phosphoglycoproteins

Use of acidic and basic pH and calcium ion addition in the capillary zone electrophoretic characterization of recombinant human deoxyribonuclease, a complex phosphoglycoprotein 853(1999)295

Phospholipases

Determination of enzymatic activity and properties of secretory phospholipase A_2 by capillary electrophoresis 853(1999)285

Phospholipids

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

Efficient method for (lyso)phospholipid class separation by high-performance liquid chromatography using an evaporative light-scattering detector 864(1999)179

Erratum to: “Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes”. [J. Chromatogr. A, 852 (1999) 59-71] 872(2000)323

Quantitation and characterization of phospholipids in pharmaceutical formulations by liquid chromatography–mass spectrometry 876(2000)97

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Methyl- β -cyclodextrin modified micellar electrokinetic capillary chromatography with laser-induced fluorescence for separation and detection of phospholipids 894(2000)129

Phosphonates

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

Selective extraction of hydrocarbons, phosphonates and phosphonic acids from soils by successive supercritical fluid and pressurized liquid extractions 866(2000)231

Development and validation of a capillary zone electrophoretic method for the determination of bisphosphonate and phosphonate impurities in clodronate 893(2000)411

Phosphonic acids

Selective extraction of hydrocarbons, phosphonates and phosphonic acids from soils by successive supercritical fluid and pressurized liquid extractions 866(2000)231

New methods for determination of glyphosate and (aminomethyl)phosphonic acid in water and soil 886(2000)207

Determination of optical purity of phosphonic acid analogues of aromatic amino acids by capillary electrophoresis with α -cyclodextrin 895(2000)301

Phosphopeptides

Phosphopeptide analysis by on-line immobilized metal-ion affinity chromatography–capillary electrophoresis–electrospray ionization mass spectrometry 853(1999)225

Phosphoproteins

Phosphopeptide analysis by on-line immobilized metal-ion affinity chromatography–capillary electrophoresis–electrospray ionization mass spectrometry 853(1999)225

Phosphoramidates

Free solution mobility of DNA molecules containing variable numbers of cationic phosphoramidate internucleoside linkages 883(2000)267

Photosynthetic pigments

Improved liquid chromatographic method for the analysis of photosynthetic pigments of higher plants 876(2000)111

Phthalates

Fractionation and toxicity evaluation of waste waters 889(2000)149

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

o-Phthaldialdehyde

Effect of solvents on the fumonisins analysis by high-performance liquid chromatography with AccQ.Fluor as the derivatizing reagent 870(2000)469

Derivatization of ephedrine with *o*-phthaldialdehyde for liquid chromatography after treatment with sodium hypochlorite 893(2000)69

Phthalic esters

Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121

Phytic acid

Anion-exchange high-performance liquid chromatography with conductivity detection for the analysis of phytic acid in food 871(2000)7

Phytol

Separation of stereoisomers in a simulated moving bed-supercritical fluid chromatography plant 865(1999)175

Pigments

Analysis of natural food pigments by capillary electrophoresis 880(2000)311

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annuum* L.) powders 889(2000)41

Solid-phase extraction and high-performance liquid chromatographic separation of pigments of red wines 889(2000)51

Colour pigments of *Trichoderma harzianum*. Preliminary investigations with thin-layer chromatography–Fourier transform infrared spectroscopy and high-performance liquid chromatography with diode array and mass spectrometric detection 896(2000)61

Separation and quantitation of colour pigments of chili powder (*Capsicum frutescens*) by high-performance liquid chromatography–diode array detection 896(2000)69

Pinanes

New stationary phase for liquid chromatography with chemically bonded pinane ligand: synthesis and characterization by nuclear magnetic resonance and high-performance liquid chromatographic investigations 883(2000)27

Pinenes

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

Application of gas chromatography–cryocondensation–Fourier transform infrared spectroscopy and gas chromatography–mass spectrometry to the identification of gas phase reaction products from the α -pinene/ozone reaction 864(1999)299

Identification of the oxidation products of the reaction between α -pinene and hydroxyl radicals by gas and high-performance liquid chromatography with mass spectrometric detection 896(2000)135

Piperazines

Novel, trifunctional diamine for silica coating in capillary zone electrophoresis 894(2000)53

Piperonal

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669

Solid phase microextraction of alkenylbenzenes and other flavor-related compounds from tobacco for analysis by selected ion monitoring gas chromatography–mass spectrometry 858(1999)79

Pirimcarb

Comparison of capillary electrochromatography with high-performance liquid chromatography for the analysis of pirimicarb and related compounds 855(1999)215

Plant hormones

Herbicide and plant growth regulator analysis by capillary electrophoresis 880(2000)281

Plasticizers

Polymer additive analysis by pyrolysis–gas chromatography. I. Plasticizers 883(2000)199

Platelet aggregation inhibitors

Determination of the impurity profile of ethyl-2-(4-((5*R*)-3-[4-(methoxycarboxamidoiminomethyl)-phenyl]-2-oxo-5-oxazolidinylmethyl)-1-piperazinyl) acetate, citrate, a highly active platelet aggregation inhibitor, by liquid chromatography–mass spectrometry 854(1999)13

Platinum

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

PNU-83892

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

PNU-83894

Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201

PNU-159548

Determination of 4-demethoxy-3'-deamino-3'-aziridinyl-4'-methylsulphonyldaunorubicin and its 13-hydroxy metabolite by direct injection of human plasma into a column-switching liquid chromatography system with mass spectrometric detection 854(1999)81

PNU-169884

Determination of 4-demethoxy-3'-deamino-3'-aziridinyl-4'-methylsulphonyldaunorubicin and its 13-hydroxy metabolite by direct injection of human plasma into a column-switching liquid chromatography system with mass spectrometric detection 854(1999)81

Poly(allylamine)

Poly(allylamine) beads as selective sorbent for preconcentration of formaldehyde and acetaldehyde in high-performance liquid chromatographic analysis 886(2000)83

Polyamides

Supercritical fluid extraction of nylon 6,6 oligomers and their characterization via liquid chromatography coupled with mass spectrometry 855(1999)715

Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. I. Injection procedure and column stability 876(2000)37

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. II. Methods of detection and quantification and overall long-term performance 878(2000)45

Polyamines

Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425

Changes in free and conjugated polyamines during starvation of sugarcane juices as analyzed by high-performance liquid chromatography 881(2000)531

Determination of hair polyamines as *N*-ethoxycarbonyl-*N*-pentafluoropropionyl derivatives by gas chromatography–mass spectrometry 897(2000)295

Poly(benzoxazine)

Qualitative and quantitative analysis of a thermoset polymer, poly(benzoxazine), by pyrolysis–gas chromatography 886(2000)217

Polybetaines

Characterization of synthetic polyelectrolytes by capillary electrophoresis 894(2000)105

Polybutadiene

Number-average molecular mass determination of polymeric material by pyrolysis–gas chromatography 888(2000)209

Polycarboxylic acids

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Polychlorinated biphenyl atropisomers

Enantiomer separation of polychlorinated biphenyl atropisomers and polychlorinated biphenyl retention behavior on modified cyclodextrin capillary gas chromatography columns 866(2000)213

Polychlorinated biphenyls

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

Low-carbon silica sorbents for solid-phase extraction 868(2000)1

Applications using the chlorine-selective pulsed discharge emission detector 872(2000)167

On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175

Utilization of fat retainers in supercritical fluid extraction for the selective extraction of polychlorinated biphenyls from a model fat sample 877(2000)167

Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83

Liquid chromatographic profiles of individual compounds of technical toxaphene 886(2000)123

Validation of an analytical procedure for polychlorinated biphenyls, coplanar polychlorinated biphenyls and polycyclic aromatic hydrocarbons in environmental samples 886(2000)153

Comparison of two extraction methods independently developed on two conceptually different automated supercritical fluid extraction systems for the determination of polychlorinated biphenyls in sediments 891(2000)195

Polychlorinated dibenzodioxins

Highly-sensitive micellar electrokinetic chromatographic analysis of dioxin-related compounds using on-line concentration 853(1999)413

Polychlorinated dibenzothiophenes

Prediction of gas chromatographic retention indices of polychlorinated dibenzothiophenes on non-polar columns 866(2000)105

Poly(*p*-chloromethylstyrene)

DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167

Polycyclic aromatic hydrocarbons

see Polynuclear aromatic hydrocarbons

Poly(diallyldimethylammonium chloride)

Separation of anions by ion chromatography–capillary electrophoresis 879(2000)245

Poly(dimethylsiloxane)

Lattice–fluid model for gas–liquid chromatography 862(1999)49

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Poly(divinylbenzene-co-*N*-vinylpyrrolidone)

Determination of phenoxyalkanoic acids and other herbicides at the ng/ml level in water by solid-phase extraction with poly(divinylbenzene-co-*N*-vinylpyrrolidone) sorbent and high-performance liquid chromatography–diode-array detection 867(2000)169

Poly(divinylbenzene)

Application of polydivinylbenzene liquid chromatography columns to remove lipid material from fish tissue extracts for the analysis of semivolatiles organics 859(1999)235

Polyesters

Separation of polyesters by gradient reversed-phase high-performance liquid chromatography on a 1.5 μm non-porous column 871(2000)243

Polyethylene

Comparison of methods for quantitative analysis of additives in low-density polyethylene using supercritical fluid and enhanced solvent extraction 858(1999)209

Polymer additive analysis by pyrolysis–gas chromatography. III. Lubricants 891(2000)313

Poly(ethylene glycol)

Capillary electrophoretic behaviour of humic substances in physical gels 863(1999)195

Influence of pH*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63

Liquid adsorption chromatography of polyethers: experiments and simulation 890(2000)195

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Selectivity equivalence of poly(ethylene glycol) stationary phases for gas chromatography 898(2000)211

Poly(ethylene oxide)

Stepwise capillary electrophoretic separation of DNA fragments using poly(ethylene oxide) solutions in the presence of electroosmotic flow 853(1999)337

Capillary zone electrophoretic analysis of positively charged poly(ethylene oxide) macromolecules using non-covalent polycation-coated fused-silica capillary and indirect UV detection 864(1999)335

Simultaneous quantitative trace analysis of anionic and nonionic surfactant mixtures by reversed-phase liquid chromatography 878(2000)99

Effect of ionic strength, pH and polymer concentration on the separation of DNA fragments in the presence of electroosmotic flow 894(2000)219

Polyethylenimine

DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167

Polyfluorocarboxylic acids

Novel way of separating polyfluorocarboxylic acids by ion-exclusion chromatography 884(2000)93

Polyglycerols

Combined liquid and gas chromatographic characterisation of polyglycerol fatty acid esters 896(2000)239

Poly(glycidyl methacrylate)

Modified capillary electrophoresis system for peptide, protein and double-stranded DNA analysis 879(2000)189

Polyglycols

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Polymers

Purification of protein and recycling of polymers in a new aqueous two-phase system using two thermoseparating polymers 864(1999)31

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

Ion-exclusion controlled size-exclusion chromatography of methacrylic acid–methyl methacrylate copolymers 872(2000)91

Investigation of Cr(III) hydrolytic polymerisation products by capillary electrophoresis–inductively coupled plasma–mass spectrometry 872(2000)227

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. I. Injection procedure and column stability 876(2000)37

Determination of basic nitrogen-containing polynuclear aromatic hydrocarbons formed during thermal degradation of polymers by high-performance liquid chromatography–fluorescence detection 878(2000)171

Contamination in food from packaging material 882(2000)255

Polymer additive analysis by pyrolysis–gas chromatography. I. Plasticizers 883(2000)199

Qualitative and quantitative analysis of a thermoset polymer, poly(benzoxazine), by pyrolysis–gas chromatography 886(2000)217

Polymer additive analysis by pyrolysis–gas chromatography. II. Flame retardants 886(2000)225

Number-average molecular mass determination of polymeric material by pyrolysis–gas chromatography 888(2000)209

Liquid adsorption chromatography of polyethers: experiments and simulation 890(2000)195

Determination of cyclodextrin content using periodate oxidation by capillary electrophoresis 891(2000)201

Polymer additive analysis by pyrolysis–gas chromatography. III. Lubricants 891(2000)313

Polymer additive analysis by pyrolysis–gas chromatography. IV. Antioxidants 891(2000)325

Characterization of synthetic polyelectrolytes by capillary electrophoresis 894(2000)105

Off-line size-exclusion chromatographic fractionation–matrix-assisted laser desorption ionization time-of-flight mass spectrometry for polymer characterization. Theoretical and experimental study 896(2000)19

Polymers, monolithic

Design of the monolithic polymers used in capillary electrochromatography columns 887(2000)3

Poly(methyloctylsiloxane)

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

Polymyxins

Liquid chromatography of polymyxin B sulphate 870(2000)237

Polymyxin B sulfate

Analysis of polymyxin B sulfate by capillary zone electrophoresis with cyclodextrin as additive. Method development and validation 879(2000)211

Polynuclear aromatic compounds

Laser-induced dispersed fluorescence detection of polycyclic aromatic compounds in soil extracts separated by capillary electrochromatography 883(2000)231

Polynuclear aromatic hydrocarbons

see also Hydrocarbons, aromatic

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

Capillary electrochromatography with macroporous particles 853(1999)45

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159

- Separation and identification of polar degradation products of benzo[a]pyrene with ozone by atmospheric pressure chemical ionization–mass spectrometry after optimized column chromatographic clean-up 855(1999)501
- On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279
- n*-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133
- Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29
- Shape-selective separation of polycyclic aromatic hydrocarbons on protoporphyrin-silica phases. Effect of surface porphyrin distribution on column efficiency 859(1999)121
- Application of polydivinylbenzene liquid chromatography columns to remove lipid material from fish tissue extracts for the analysis of semivolatiles organics 859(1999)235
- Analysis of high-molecular-mass polycyclic aromatic hydrocarbons in environmental samples using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 863(1999)13
- Separation and identification of perchlorinated polycyclic aromatic hydrocarbons by high-performance liquid chromatography and ultraviolet absorption spectroscopy 864(1999)173
- High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93
- Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101
- Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329
- Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505
- Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515
- Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415
- Novel approach to the analysis and use of fullerenes in capillary electrophoresis 873(2000)257
- Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography 873(2000)293
- Dynamic microwave-assisted extraction 877(2000)141
- Determination of basic nitrogen-containing polynuclear aromatic hydrocarbons formed during thermal degradation of polymers by high-performance liquid chromatography–fluorescence detection 878(2000)171
- Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83
- Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121
- Solid-phase microextraction for determining the distribution of sixteen US Environmental Protection Agency polycyclic aromatic hydrocarbons in water samples 879(2000)177
- Chromatographic analysis of minor constituents in vegetable oils 881(2000)131
- Polycyclic aromatic hydrocarbons in edible fats and oils: occurrence and analytical methods 882(2000)245
- Novel use of a dual-zone restricted access sorbent: normal-phase solid-phase extraction separation of methyl oleate from polynuclear aromatic hydrocarbons stemming from semi-permeable membrane devices 883(2000)143
- Pressurised liquid extraction of polycyclic aromatic hydrocarbons from contaminated soils 883(2000)151
- Solid-phase extraction of polycyclic aromatic compounds 885(2000)273
- Validation of an analytical procedure for polychlorinated biphenyls, coplanar polychlorinated biphenyls and polycyclic aromatic hydrocarbons in environmental samples 886(2000)153
- Static subcritical water extraction with simultaneous solid-phase extraction for determining polycyclic aromatic hydrocarbons on environmental solids 886(2000)237
- Capillary electrochromatography–mass spectrometry for the separation and identification of isomeric polyaromatic hydrocarbon dna adducts derived from *in vitro* reactions 887(2000)103
- Preparation of stationary phases for open-tubular capillary electrochromatography using the sol–gel method 887(2000)253
- Solid-phase extraction of sugar cane soot extract for analysis by gas chromatography with flame ionisation and mass spectrometric detection 889(2000)281

Miniaturised pressurised liquid extraction of polycyclic aromatic hydrocarbons from soil and sediment with subsequent large-volume injection–gas chromatography 891(2000)275

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421

Pressurised liquid extraction of ketones of polycyclic aromatic hydrocarbons from soil 893(2000)201

Investigation of the nematic–isotropic transition of a liquid crystalline polymer and determination of molecular diffusion coefficients using gas chromatography 893(2000)359

Laterally attached liquid crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. II. Optimization of the molecular parameter of the polymer 897(2000)131

Determination of polycyclic aromatic hydrocarbons in waste water by off-line coupling of solid-phase microextraction with column liquid chromatography 897(2000)153

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

Polynucleotides

see Nucleotides and Special Issue Index

Polyolefins

Determination of low levels of an antioxidant in polyolefins by large-volume injection temperature-programmed packed capillary liquid chromatography 864(1999)103

Microwave-assisted extraction by fast sample preparation for the systematic analysis of additives in polyolefins by high-performance liquid chromatography 869(2000)285

Polyols

Separation and determination of alditols and sugars by high-pH anion-exchange chromatography with pulsed amperometric detection 897(2000)195

Polyphenols

Determination of polyphenols by high-performance liquid chromatography with inhibited chemiluminescence detection 855(1999)171

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

Separation of polyphenols in Canary Islands wine by capillary zone electrophoresis without preconcentration 874(2000)111

Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425

Comparative analysis of tea catechins and theaflavins by high-performance liquid chromatography and capillary electrophoresis 881(2000)439

High-speed counter-current chromatography of apple procyanidins 886(2000)65

Polyphenylene sulfide

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

Polyphosphates

Analysis of condensed phosphates in food products by ion chromatography with an on-line hydroxide eluent generator 881(2000)639

Polysaccharides

see also Saccharides

Molecular weight determination of hyaluronic acid by gel filtration chromatography coupled to matrix-assisted laser desorption ionization mass spectrometry 852(1999)573

Molecular mass distribution of sodium alginate by high-performance size-exclusion chromatography 864(1999)199

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

Analysis of oligomannuronic acids and oligoguluronic acids by high-performance anion-exchange chromatography and electrospray ionization mass spectrometry 884(2000)105

Enantioseparations in non-aqueous capillary electrochromatography using polysaccharide type chiral stationary phases 887(2000)439

Determination of free carrier protein in protein–polysaccharide conjugate vaccines by micellar electrokinetic chromatography 894(2000)311

Polysiloxane

Synthesis and characteristics of [60]fullerene polysiloxane stationary phase for capillary gas chromatography 867(2000)177

Poly(sodium 11-acrylamidoundecanoate)

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Polystyrene

Preparation of uniformly sized polymeric separation media potentially suitable for small-scale high-performance liquid chromatography and/or capillary electrochromatography 853(1999)11

Lattice–fluid model for gas–liquid chromatography 862(1999)49

Modelling of the pore flow in capillary electrochromatography 887(2000)199

Separation of unmodified polystyrene nanosphere standards by capillary zone electrophoresis 898(2000)257

Poly(styrene–divinylbenzene)

Capillary columns with in situ formed porous monolithic packing for micro high-performance liquid chromatography and capillary electrochromatography 855(1999)273

Polythionates

Determination of thiosulfate, thiocyanate and polythionates in a mixture by ion-pair chromatography with ultraviolet absorbance detection 884(2000)81

Poly(vinyl alcohol)

Capillary electrophoretic behaviour of humic substances in physical gels 863(1999)195

Influence of pH*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63

Polyvinylpyridines

Characterization of synthetic polyelectrolytes by capillary electrophoresis 894(2000)105

Potassium sorbate

Rapid high-performance liquid chromatography method for the analysis of sodium benzoate and potassium sorbate in foods 883(2000)299

Prednisone

Direct identification and quantitation of prednisone in the presence of overlapping hydrocortisone by liquid chromatography with electrospray and atmospheric-pressure chemical-ionisation mass spectrometry 870(2000)135

Preservatives

Development of a capillary electrophoresis method for the simultaneous analysis of artificial sweeteners, preservatives and colours in soft drinks 876(2000)213

Rapid high-performance liquid chromatography method for the analysis of sodium benzoate and potassium sorbate in foods 883(2000)299

Prion proteins

Use of capillary electrophoresis and fluorescent labeled peptides to detect the abnormal prion protein in the blood of animals that are infected with a transmissible spongiform encephalopathy 853(1999)207

Proanthocyanidins

Analysis of pigmented high-molecular-mass grape phenolics using ion-pair, normal-phase high-performance liquid chromatography 866(2000)25

Procyanidins

Fractionation of apple procyanidins by size-exclusion chromatography 855(1999)181

High-speed counter-current chromatography of apple procyanidins 886(2000)65

Fractionation of apple procyanidins according to their degree of polymerization by normal-phase high-performance liquid chromatography 890(2000)251

Profens

see also Ibuprofen, Ketoprofen and Nonsteroidal anti-inflammatory drugs

Novel cinchona alkaloid carbamate C₉-dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Investigation of an enantioselective non-aqueous capillary electrochromatography system applied to the separation of chiral acids 875(2000)341

Progestogens

Determination of steroid sex hormones and related synthetic compounds considered as endocrine disrupters in water by liquid chromatography–diode array detection–mass spectrometry 892(2000)391

Propranolol

Direct, preparative enantioselective chromatography of propranolol hydrochloride and thioridazine hydrochloride using carbon dioxide-based mobile phases 865(1999)227

Optimisation of procedures for the extraction of structural analogues of propranolol with molecular imprinted polymers for sample preparation 889(2000)143

Application of molecularly imprinted polymers in supercritical fluid chromatography 897(2000)317

Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate β -cyclodextrin bonded chiral stationary phase 898(2000)53

Propylene oxide

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Prostaglandins

New sheathless interface for coupling capillary electrophoresis to electrospray mass spectrometry evaluated by the analysis of fatty acids and prostaglandins 854(1999)141

Solid- and liquid-phase extraction for the gas chromatographic–tandem mass spectrometric quantification of 2,3-dinor-thromboxane B₂ and 2,3-dinor-6-oxo-prostaglandin F_{1α} in human urine 885(2000)351

Protease inhibitors

Determination of human immunodeficiency virus-1 protease inhibitors in patient serum using free solution capillary zone electrophoresis 853(1999)431

Protegrins

Development and validation of a capillary electrophoresis method for the characterization of protegrin IB-367 853(1999)197

Proteins

see also Antibodies, Enzyme inhibitors, Enzymes, Immunoglobulins, Monoclonal antibodies and Special Issue Index

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

Ion-exchange chromatography of proteins near the isoelectric points 852(1999)31

Characterization of unstable ion-exchange chromatographic separation of proteins 852(1999)37

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Large-scale membrane adsorbers 852(1999)73

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105

Labeling effects on the isoelectric point of green fluorescent protein 853(1999)21

General experimental aspects of the use of isoelectric buffers in capillary electrophoresis 853(1999)71

Use of capillary electrophoresis and fluorescent labeled peptides to detect the abnormal prion protein in the blood of animals that are infected with a transmissible spongiform encephalopathy 853(1999)207

Phosphopeptide analysis by on-line immobilized metal-ion affinity chromatography–capillary electrophoresis–electrospray ionization mass spectrometry 853(1999)225

Electrokinetically-driven cation-exchange chromatography of proteins and its comparison with pressure-driven high-performance liquid chromatography 853(1999)243

Microemulsion electrokinetic chromatography of proteins 853(1999)277

Competitive immunoassay for staphylococcal enterotoxin A using capillary electrophoresis with laser-induced fluorescence detection 853(1999)545

Liquid chromatography–electrospray mass spectrometry as a tool for the analysis of sulfated oligosaccharides from mucin glycoproteins 854(1999)131

Macroporous poly(glycidyl methacrylate–triallyl isocyanurate–divinylbenzene) matrix as an anion-exchange resin for protein adsorption 855(1999)129

Separation of oxidized and deamidated human growth hormone variants by isocratic reversed-phase high-performance liquid chromatography 855(1999)147

Renaturation of heterodimeric platelet-derived growth factor from inclusion bodies of recombinant *Escherichia coli* using size-exclusion chromatography 855(1999)203

Capillary columns with in situ formed porous monolithic packing for micro high-performance liquid chromatography and capillary electrochromatography 855(1999)273

Analysis of recombinant and modified proteins by capillary zone electrophoresis coupled with electrospray ionization tandem mass spectrometry 855(1999)695

Recent developments in microcolumn liquid chromatography 856(1999)117

Continuous-bed chromatography for the analysis and purification of recombinant human basic fibroblast growth factor 857(1999)137

Integrated approach to the multidimensional analysis of complex biological samples by microseparation techniques. Analysis of glycoprotein factor associated with cancer cachexia 857(1999)183

- Measurement of electroosmotic flow in plastic imprinted microfluid devices and the effect of protein adsorption on flow rate 857(1999)275
- Direct determination of small cations in proteinaceous samples using a flow injection–capillary electrophoresis system 857(1999)321
- Capillary electrochromatography of proteins and peptides with porous-layer open-tubular columns 858(1999)91
- Analysis of bovine whey proteins in soybean dairy-like products by capillary electrophoresis 859(1999)77
- Improved liquid chromatographic separation of different proteins by designing functional surfaces of cattle bone-originated apatite 862(1999)217
- Cytochrome *c* unfolding on an anionic surface 863(1999)137
- Purification of protein and recycling of polymers in a new aqueous two-phase system using two thermoseparating polymers 864(1999)31
- Considerations of sample application and elution during size-exclusion chromatography-based protein refolding 864(1999)229
- Eliminating disulfide exchange during glutamyl endopeptidase digestion of native protein 864(1999)237
- Natural poly-histidine affinity tag for purification of recombinant proteins on cobalt(II)-carboxymethylaspartate crosslinked agarose 864(1999)247
- Influence of column design on process-scale ion-exchange chromatography 865(1999)3
- Polycations as displacer in high-performance bioseparation 865(1999)27
- Study of hydrophobic interaction adsorption of bovine serum albumin under overloaded conditions using flow microcalorimetry 865(1999)111
- Expanded-bed chromatography in primary protein purification 865(1999)129
- Electrophoretic transfer of proteins across polyacrylamide membranes 865(1999)145
- Protein diffusion in charged polyacrylamide gels. Visualization and analysis 865(1999)155
- Rapid reversed-phase separation of proteins and peptides using optimized ‘moulded’ monolithic poly(styrene-*co*-divinylbenzene) columns 865(1999)169
- Reversed-phase liquid chromatography of proteins and peptides using multimodal copolymer-encapsulated silica 866(2000)1
- One-step capillary isoelectric focusing for the separation of the recombinant human immunodeficiency virus envelope glycoprotein glycoforms 866(2000)121
- Comparative studies on the analysis of glycosylation heterogeneity of sialic acid-containing glycoproteins using capillary electrophoresis 866(2000)261
- Size-exclusion chromatography–multiangle laser light scattering analysis of β -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105
- New affinity nylon membrane used for adsorption of γ -globulin 867(2000)161
- Protein adsorption to the bare silica wall in capillary electrophoresis. Quantitative study on the chemical composition of the background electrolyte for minimising the phenomenon 868(2000)85
- Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151
- Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255
- Signature-peptide approach to detecting proteins in complex mixtures 870(2000)295
- Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363
- Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371
- Microcalorimetric studies of interactions between proteins and hydrophobic ligands in hydrophobic interaction chromatography: effects of ligand chain length, density and the amount of bound protein 872(2000)37
- Erratum to: “Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes”. [J. Chromatogr. A, 852 (1999) 59-71] 872(2000)323
- The influence of cell adsorbent interactions on protein adsorption in expanded beds 873(2000)195
- Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. II. Effects of adsorbed, neutral polymers on quenching the interaction 874(2000)293

- Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors 875(2000)235
- Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255
- Enantioselective binding analysis of verapamil to plasma lipoproteins by capillary electrophoresis–frontal analysis 875(2000)447
- Simultaneous separation and quantitation of the major bovine whey proteins including proteose peptone and caseinomacropptide by reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene 878(2000)183
- Lysozyme refolding with immobilized GroEL column chromatography 878(2000)197
- Determination of cow's milk in non-bovine and mixed cheeses by capillary electrophoresis of whey proteins in acidic isoelectric buffers 878(2000)261
- Modified capillary electrophoresis system for peptide, protein and double-stranded DNA analysis 879(2000)189
- Perfusion chromatography: an emergent technique for the analysis of food proteins 880(2000)169
- Electrophoretic methods in the analysis of beverages 880(2000)243
- Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides 881(2000)1
- Electrophoresis of cereal storage proteins 881(2000)23
- Determination by perfusion reversed-phase high-performance liquid chromatography of the soybean protein content of commercial soybean products prepared directly from whole soybeans 881(2000)37
- Characterization of commercial soybean products by conventional and perfusion reversed-phase high-performance liquid chromatography and multivariate analysis 881(2000)47
- Evaluation of the effect of rennet type on casein proteolysis in an ovine milk cheese by means of capillary electrophoresis 881(2000)59
- High-performance liquid chromatography coupled to enzyme-amplified biochemical detection for the analysis of hemoglobin after pre-column biotinylation 886(2000)19
- Protein identification platform utilizing micro dispensing technology interfaced to matrix-assisted laser desorption ionization time-of-flight mass spectrometry 886(2000)99
- Resolution and identification of the protein components of the photosystem II antenna system of higher plants by reversed-phase liquid chromatography with electrospray-mass spectrometric detection 886(2000)111
- Specific applications of capillary electrochromatography to biopolymers, including proteins, nucleic acids, peptide mapping, antibodies, and so forth 887(2000)137
- Capillary electrochromatography of proteins and peptides with a cationic acrylic monolith 887(2000)465
- Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3
- Use of spray-dried zirconia microspheres in the separation of immunoglobulins from cell culture supernatant 890(2000)15
- High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. II. Separation of proteins 890(2000)37
- Suspended bed chromatography, a new approach in downstream processing 890(2000)45
- Scale-down of continuous protein purification by annular chromatography. Design parameters for the smallest unit 890(2000)53
- Protein losses in ion-exchange and hydrophobic interaction high-performance liquid chromatography 890(2000)73
- Trapping the monomeric α -helical state during unfolding of coiled-coils by reversed-phase liquid chromatography 890(2000)81
- Adsorption and selectivity characteristics of several human serum proteins with immobilised hard Lewis metal ion–chelate adsorbents 890(2000)95
- Characterisation of the rubber elongation factor from ammoniated latex by electrophoresis and mass spectrometry 890(2000)145
- Fast quantitation of recombinant plasminogen activator inhibitor type 1 in bacterial lysates by micropellicular reversed-phase liquid chromatography 891(2000)85
- Discoloration of ceramic hydroxyapatite used for protein chromatography 891(2000)93
- Pressure-induced retention of the lysozyme on reversed-phase liquid chromatography 891(2000)207
- Approaches towards the quantitative analysis of peptides and proteins by reversed-phase high-performance liquid chromatography in the absence of a pure reference sample 891(2000)235

Use of amino acid analysis for estimating the individual concentrations of proteins in mixtures 891(2000)355

Applications of mass spectrometry to food proteins and peptides 893(2000)1

Protein mapping by two-dimensional high performance liquid chromatography 893(2000)293

Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79

Quantitation of protein binding to the capillary wall in acidic, isoelectric buffers and means for minimizing the phenomenon 894(2000)273

Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. III: Effects of adsorbed surfactants on quenching the interaction 894(2000)281

Separation of proteins by sodium dodecylsulfate capillary electrophoresis in hydroxypropylcellulose sieving matrix with laser-induced fluorescence detection 894(2000)291

Capillary electrophoresis of seed 2S albumins from *Lupinus* species 894(2000)297

Determination of free carrier protein in protein–polysaccharide conjugate vaccines by micellar electrokinetic chromatography 894(2000)311

Congophilicity (Congo red affinity) of different β_2 -microglobulin conformations characterized by dye affinity capillary electrophoresis 894(2000)319

Rapid analysis of covalently and non-covalently fluorophore-labeled proteins using ultra-thin-layer sodium dodecylsulfate gel electrophoresis 894(2000)329

Sodium dodecylsulfate capillary gel electrophoretic measurement of the concentration ratios of albumin and α_2 -macroglobulin in cerebrospinal fluid and serum of patients with neurological disorders 895(2000)339

Protein adsorption on novel acrylamido-based polymeric ion-exchangers. I. morphology and equilibrium adsorption 897(2000)65

Protein adsorption on novel acrylamido-based polymeric ion exchangers. II. Adsorption rates and column behavior 897(2000)81

Comparison of chromatographic ion-exchange resins. I. Strong anion-exchange resins 897(2000)99

Multiple-step ligand injection affinity capillary electrophoresis for determining binding constants of ligands to receptors 897(2000)339

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

Proteose peptone

Simultaneous separation and quantitation of the major bovine whey proteins including proteose peptone and caseinomacropptide by reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene 878(2000)183

Protoporphyrins

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Provitamins

Simultaneous determination of fat-soluble vitamins and provitamins in milk by microcolumn liquid chromatography 891(2000)109

Pseudoephedrine

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Puerarins

Separation and purification of isoflavones from *Pueraria lobata* by high-speed counter-current chromatography 855(1999)709

Pulegone

Solid phase microextraction of alkenylbenzenes and other flavor-related compounds from tobacco for analysis by selected ion monitoring gas chromatography–mass spectrometry 858(1999)79

Pullulan

Molecular mass distribution of sodium alginate by high-performance size-exclusion chromatography 864(1999)199

Purines

Separation of purines and pyrimidines by normal-phase high-performance liquid chromatography using dimethyl sulfoxide in binary and ternary eluents 874(2000)45

Determination of purines including 2,8-dihydroxyadenine in urine using capillary electrophoresis 894(2000)157

Pyrazines

Identification of 2,3-butanedione monoxime hydrogenation products by gas chromatography–mass spectrometry in an ion trap mass spectrometer 855(1999)341

Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxypyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93

Pyrazoles

Identification of 2,3-butanedione monoxime hydrogenation products by gas chromatography–mass spectrometry in an ion trap mass spectrometer 855(1999)341

Pyrethrin esters

Separation of natural pyrethrum extracts using micellar electrokinetic chromatography 863(1999)89

Pyrethroids

High-performance liquid chromatography study of the enantiomer separation of chrysanthemic acid and its analogous compounds on a terguride-based stationary phase 859(1999)133

Interlaboratory study of identification and quantitation of multiresidue pyrethroids in agricultural products by gas chromatography–mass spectrometry 882(2000)231

2,6-Pyridinedicarboxylic acid

Non-suppressed conductivity and indirect UV detection of carboxylic acids in environmental samples by ion-exclusion chromatography using 2,6-pyridinedicarboxylic acidic eluent 859(1999)173

Pyridines

Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Pyrimethamine

Simultaneous determination of sulfaquinoxaline, sulfamethazine and pyrimethamine by liquid chromatography 870(2000)169

Pyrimidines

Comparison of capillary electrochromatography with high-performance liquid chromatography for the analysis of pirimicarb and related compounds 855(1999)215

Separation of purines and pyrimidines by normal-phase high-performance liquid chromatography using dimethyl sulfoxide in binary and ternary eluents 874(2000)45

Pyrroles

Identification of 2,3-butanedione monoxime hydrogenation products by gas chromatography–mass spectrometry in an ion trap mass spectrometer 855(1999)341

Pyrroloquinoline quinone

Separation of the enzyme cofactor pyrroloquinoline quinone and three isomeric analogues by capillary electrophoresis with ion-pairing media 876(2000)193

Pyruvate

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Q**Quaternary ammonium compounds**

On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441

Investigation of the effects of cyclodextrins and organic solvents on the separation of cationic surfactants in capillary electrophoresis 872(2000)269

Solid-phase extraction of quaternary ammonium herbicides 885(2000)251

Quercetin

Spectrophotometric and coulometric detection in the high-performance liquid chromatography of flavonoids and optimization of sample treatment for the determination of quercetin in orange juice 881(2000)449

Quinine

Quinine versus carbamoylated quinine-based chiral anion exchangers. A comparison regarding enantioselectivity for N-protected amino acids and other chiral acids 858(1999)1

Quinine carbamates

Novel cinchona alkaloid carbamate C₉-dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111

Quinolinone

Sample cleanup and reversed-phase high-performance liquid chromatographic analysis of polar aromatic compounds in groundwater samples from a former gas plant 862(1999)137

Quinolizidines

Determination of quinolizidine alkaloids in traditional Chinese herbal drugs by nonaqueous capillary electrophoresis 857(1999)303

Quinolones

Determination of quinolones in animal tissues and eggs by high-performance liquid chromatography with photodiode-array detection 871(2000)31

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Influence of pH and pK_a values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

New amphiphilic aminosaccharide derivatives as chiral selectors in capillary electrophoresis 875(2000)295

Quantitation of nine quinolones in chicken tissues by high-performance liquid chromatography with fluorescence detection 882(2000)63

Quinones

Preparative isolation and purification of hydroxyanthraquinones from *Rheum officinale* Baill by high-speed counter-current chromatography using pH-modulated stepwise elution 858(1999)103

Quinuclidinol

Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169

R

Racemorphan

Cyclodextrin-mediated micellar electrokinetic chromatography and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry for the enantiomer separation of racemorphan in human urine 853(1999)439

Rare earth ions

see also Lanthanides

Eriochrome Black T as a post-column reagent for the ion chromatographic determination of rare earths 855(1999)111

Separation and determination of rare earth elements by Dowex 2-X8 resin using sodium trimetaphosphate as elution agent 874(2000)311

Improved capillary electrophoresis method for measuring rare-earth elements in synthetic geochemical standards 884(2000)317

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N,N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Capillary electrophoresis–inductively coupled plasma mass spectrometry interface with minimised dead volume for high separation efficiency 891(2000)129

Separation and direct UV detection of lanthanides complexed with cupferron by capillary electrophoresis 895(2000)263

Recombinant proteins

see also Proteins

Capillary electrophoretic analysis of recombinant human apolipoprotein E. Calibration mode of a protein reference material 853(1999)237

Remazol dyes

Application of electrospray mass spectrometry in the detection and determination of Remazol textile dyes 854(1999)259

Resveratrol

Analysis of resveratrol in wine by capillary electrophoresis 881(2000)471

Retinal isomers

A reversed-phase high-performance liquid chromatographic method to analyze retinal isomers 864(1999)257

Retinoids

C_{30} Stationary phases for the analysis of food by liquid chromatography 880(2000)189

High-performance liquid chromatography of non-polar retinoid isomers 881(2000)183

Retinol

see also Vitamins

Improved method for rapid determination of vitamin A in small samples of breast milk by high-performance liquid chromatography 898(2000)179

Retinyl esters

Nonaqueous electrochromatography on continuous bed columns of sol–gel bonded large-pore C_{18} material: separation of retinyl esters 890(2000)347

Rhamnolipids

High-performance liquid chromatography method for the characterization of rhamnolipid mixtures produced by *Pseudomonas aeruginosa* UG2 on corn oil 864(1999)211

Rhodium

Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217

Riboflavin phosphates

On-line automated high-performance liquid chromatographic determination of total riboflavin phosphates using immobilized acid phosphatase as a pre-column reactor 896(2000)171

Riboflavins

Chromatographic determination of riboflavin and its derivatives in food 881(2000)285

High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

Ribozymes

Use of high-performance liquid chromatographic fractionation of large RNA molecules in the assay of group I intron ribozyme activity 868(2000)109

RNA

see also Nucleic acids

Introduction of the gene amplification technique to decrease the risk of hepatitis C virus transmission by plasma products 852(1999)305

Analysis of leptin gene expression in chickens using reverse transcription polymerase chain reaction and capillary electrophoresis with laser-induced fluorescence detection 853(1999)321

Use of high-performance liquid chromatographic fractionation of large RNA molecules in the assay of group I intron ribozyme activity 868(2000)109

Ro 63-1908

Determination of the *N*-methyl-D-aspartate receptor NR2B subunit blocker Ro 63-1908 in rat, cynomolgus monkey and human plasma by high-performance liquid chromatography with column switching and fluorescence detection 870(2000)189

Ropivacaine

Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151

Rotenone

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

S**Saccharides**

see also Carbohydrates, Disaccharides, Monosaccharides, Oligosaccharides, Polysaccharides and Sugars

Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255

New amphiphilic aminosaccharide derivatives as chiral selectors in capillary electrophoresis 875(2000)295

Salbutamol

Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography 873(2000)293

Design of experiments for capillary electrophoretic enantioresolution of salbutamol using dermatan sulfate 875(2000)411

On-line isotachophoretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219

Sanguinarine

Separation, identification and determination of sanguinarine in argemone and other adulterated edible oils by reversed-phase high-performance liquid chromatography 863(1999)123

Capillary electrophoretic determination of sanguinarine and chelerythrine in plant extracts and pharmaceutical preparations 866(2000)293

Saponins

On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic–diode array detection–mass spectrometric techniques 880(2000)203

Saxitoxin

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatroscan (mark 5) and flame thermionic detection 855(1999)657

Scoparone

Fractionation and analysis of *Artemisia capillaris* Thunb. by affinity chromatography with human serum albumin as stationary phase 870(2000)501

Scopolamine

Strategies for supercritical fluid extraction of hyoscyamine and scopolamine salts using basified modifiers 863(1999)47

On-line isotachophoretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219

Sea-Nine 211

Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261

Selenium

Column-switching system for selenium speciation by coupling reversed-phase and ion-exchange high-performance liquid chromatography with microwave-assisted digestion–hydride generation–atomic fluorescence spectrometry 889(2000)33

Selenium compounds

High-performance liquid chromatography of selenium compounds utilizing perfluorinated carboxylic acid ion-pairing agents and inductively coupled plasma and electrospray ionization mass spectrometric detection 866(2000)51

Speciation of selenium compounds with ion-pair reversed-phase liquid chromatography using inductively coupled plasma mass spectrometry as element-specific detection 874(2000)55

Semicarbazide

Determination of enantiomers in a synthetic argininal peptide using capillary zone electrophoresis and high-performance liquid chromatography 891(2000)115

Serotonin

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Sertraline

Analysis of *cis*–*trans* isomers and enantiomers of sertraline by cyclodextrin-modified micellar electrokinetic chromatography 871(2000)207

Sialic acid

Comparative studies on the analysis of glycosylation heterogeneity of sialic acid-containing glycoproteins using capillary electrophoresis 866(2000)261

Silicate

Determination of silicate in water by ion exclusion chromatography with conductivity detection 874(2000)143

Sinigrin

Optimization of glucosinolate separation by micellar electrokinetic capillary chromatography using a Doehlert's experimental design 864(1999)155

Skatole

Determination of indole-3-acetic acid, tryptophan and other indoles in must and wine by high-performance liquid chromatography with fluorescence detection 855(1999)227

Sodium

Determination of ammonium in seawater by column-switching ion chromatography 868(2000)209

Sodium alginate

Molecular mass distribution of sodium alginate by high-performance size-exclusion chromatography 864(1999)199

Sodium benzoate

Rapid high-performance liquid chromatography method for the analysis of sodium benzoate and potassium sorbate in foods 883(2000)299

Sodium chloride

High concentrations of sodium chloride facilitate the use of immobilized chymotrypsin for separating virgin forms of specific trypsin inhibitors 852(1999)227

Sodium dodecylsulfate

Comparison of the performance of butanol and pentanol as modifiers in the micellar chromatographic determination of some phenethylamines 866(2000)35

New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169

Simultaneous quantitative trace analysis of anionic and nonionic surfactant mixtures by reversed-phase liquid chromatography 878(2000)99

Separation of proteins by sodium dodecylsulfate capillary electrophoresis in hydroxypropylcellulose sieving matrix with laser-induced fluorescence detection 894(2000)291

Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285

Sodium hydroxide

Experimental study on moving neutralization reaction boundary created with the strong reactive electrolytes of HCl and NaOH in agarose gel 891(2000)337

Sodium hypochlorite

Derivatization of ephedrine with *o*-phthaldialdehyde for liquid chromatography after treatment with sodium hypochlorite 893(2000)69

Sodium tetraethylborate

Gas chromatographic determination of organomercury following aqueous derivatization with sodium tetraethylborate and sodium tetraphenylborate. Comparative study of gas chromatography coupled with atomic fluorescence spectrometry, atomic emission spectrometry and mass spectrometry 876(2000)147

Sodium tetraphenylborate

Gas chromatographic determination of organomercury following aqueous derivatization with sodium tetraethylborate and sodium tetraphenylborate. Comparative study of gas chromatography coupled with atomic fluorescence spectrometry, atomic emission spectrometry and mass spectrometry 876(2000)147

Sodium trimetaphosphate

Separation and determination of rare earth elements by Dowex 2-X8 resin using sodium trimetaphosphate as elution agent 874(2000)311

Sotalol

Temperature-induced inversion of elution order in the enantioseparation of sotalol on a cellobiohydrolase I-based stationary phase 858(1999)33

Spiro[2H-benzofuro(2,3-*a*)quinolizine-2,4'-(1'H)pyrimidin]-2'-(3'H)-one, 1,3,4,5',6,6',7,12b-octahydro-1',3'-dimethyl-, (2*S*-*trans*)-

Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated β -cyclodextrin 875(2000)389

Staphylococcal nuclease

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Starch

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

Steroidal glycoalkaloids

Reversed-phase liquid chromatographic separation and simultaneous profiling of steroidal glycoalkaloids and their aglycones 863(1999)25

Steroid hormone receptors

Affinity chromatography of human estrogen receptor- α expressed in *Saccharomyces cerevisiae*. Combination of heparin- and 17β -estradiol-affinity chromatography 852(1999)161

Steroid hormones

Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121

Steroids

High-performance liquid chromatographic methods for monitoring of isomers of 17-hydroxy-16-hydroxymethyl-3-methoxyestra-1,3,5(10)-triene 852(1999)433

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

Analysis of steroids in environmental water samples using solid-phase extraction and ion-trap gas chromatography–mass spectrometry and gas chromatography–tandem mass spectrometry 872(2000)309

Steroid profiles determined by capillary electrochromatography, laser-induced fluorescence detection and electrospray–mass spectrometry 887(2000)379

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Comparison of different liquid chromatography methods for the determination of corticosteroids in biological matrices 893(2000)55

Sterol esters

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

Sterols

Determination of tocopherols and sterols in vegetable oils by solid-phase extraction and subsequent capillary gas chromatographic analysis 857(1999)231

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Analysis of olive and hazelnut oil mixtures by high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry of triacylglycerols and gas–liquid chromatography of non-saponifiable compounds (tocopherols and sterols) 881(2000)149

Strong acids

Retention behaviour of strong acid anions in ion-exclusion chromatography on sulfonate and carboxylate stationary phases 884(2000)61

Styrene phosphonate

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

Substance P

Preliminary comparison of precursor scans and liquid chromatography–tandem mass spectrometry on a hybrid quadrupole time-of-flight mass spectrometer 854(1999)119

Substance P endopeptidase

Purification of Substance P endopeptidase activity in the rat ventral tegmental area with the Äkta-Purifier chromatographic system 893(2000)107

Sucrose

Design of a simulated moving bed unit for sucrose–betaine separations 872(2000)23

Sugar alcohols

Determination of sugar alcohols in confectioneries by high-performance liquid chromatography after nitrobenzoylation 893(2000)195

Sugars

see also Carbohydrates and Saccharides

Design of a simulated moving bed unit for sucrose–betaine separations 872(2000)23

Role of chromatography in the analysis of sugars, carboxylic acids and amino acids in food (Review) 891(2000)1

Separation and determination of alditols and sugars by high-pH anion-exchange chromatography with pulsed amperometric detection 897(2000)195

Sulfadimethoxine

Simplified determining procedure for routine residue monitoring of sulphamethazine and sulphadimethoxine in milk 898(2000)185

Sulfa drugs

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

Sulfamethazine

Simultaneous determination of sulfaquinoxaline, sulfamethazine and pyrimethamine by liquid chromatography 870(2000)169

Simplified determining procedure for routine residue monitoring of sulphamethazine and sulphadimethoxine in milk 898(2000)185

Sulfaquinoxaline

Simultaneous determination of sulfaquinoxaline, sulfamethazine and pyrimethamine by liquid chromatography 870(2000)169

Sulfates

Liquid chromatography–electrospray mass spectrometry as a tool for the analysis of sulfated oligosaccharides from mucin glycoproteins 854(1999)131

Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629

Determination of chloride, sulfate and nitrate in groundwater samples by ion chromatography 884(2000)185

Sulfides

Picomolar quantitation of free sulfite in foods by means of [⁵⁷Co]hydroxocobalamin and radiometric chromatography of [⁵⁷Co]sulfitocobalamin. Method, applications and significance of coexisting sulfides 881(2000)345

Sensitive determination of dissolved sulfide in estuarine water by solid-phase extraction and high-performance liquid chromatography of methylene blue 883(2000)305

Sulfite

Picomolar quantitation of free sulfite in foods by means of [⁵⁷Co]hydroxocobalamin and radiometric chromatography of [⁵⁷Co]sulfitocobalamin. Method, applications and significance of coexisting sulfides 881(2000)345

Sulfolane

Gas chromatographic–mass spectrometric determination of sulfolane in wetland vegetation exposed to sour gas-contaminated groundwater 859(1999)69

Sulfonamides

Detection and identification of sulphonamide drugs in municipal waste water by liquid chromatography coupled with electrospray ionisation tandem mass spectrometry 854(1999)163

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Quantitative determination of sulfonamide residues in foods of animal origin by high-performance liquid chromatography with fluorescence detection 871(2000)37

Selective deuterium exchange during superheated heavy water chromatography–nuclear magnetic resonance spectroscopy–mass spectrometry of sulfonamides 886(2000)289

Application of ion-exchange cartridge clean-up in food analysis. V. Simultaneous determination of sulphonamide antibacterials in animal liver and kidney using high-performance liquid chromatography with ultraviolet and mass spectrometric detection 898(2000)95

Sulfonates

On-line solid-phase extraction–ion-pair liquid chromatography–electrospray mass spectrometry for the trace determination of naphthalene monosulphonates in water 854(1999)187

Stability of sulfonated derivatives of benzene and naphthalene on disposable solid-phase extraction pre-columns and in an aqueous matrix 889(2000)231

Sulfonates, aromatic

Solid-phase extraction of polar hydrophilic aromatic sulfonates followed by capillary zone electrophoresis–UV absorbance detection and ion-pair liquid chromatography–diode array UV detection and electrospray mass spectrometry 890(2000)225

Sulfonic acids

Use of volatile amines as ion-pairing agents for the high-performance liquid chromatographic–tandem mass spectrometric determination of aromatic sulfonates in industrial wastewater 854(1999)175

Sulfonylureas

Determination of sulfonylurea degradation products in soil by liquid chromatography–ultraviolet detection followed by confirmatory liquid chromatography–tandem mass spectrometry 855(1999)575

Herbicide and plant growth regulator analysis by capillary electrophoresis 880(2000)281

Sulfophenylcarboxylic acids

Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99

Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211

Sulfosalicylic acid

Simultaneous ion-exclusion/cation-exchange chromatography of anions and cations in acid rain waters on a weakly acidic cation-exchange resin by elution with sulfosalicylic acid 884(2000)167

Sulfur compounds

see also Organosulfur compounds

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

Simultaneous measurement of volatile sulfur compounds using ascorbic acid for oxidant removal and gas chromatography–flame photometric detection 864(1999)111

On the performance and inertness of different materials used for the enrichment of sulfur compounds from air and gaseous samples 864(1999)345

Prediction of gas chromatographic retention indices of polychlorinated dibenzothiophenes on non-polar columns 866(2000)105

Pulsed discharge emission detector: an element-selective detector for gas chromatography 872(2000)119

Determination of sulphur compounds in beer using headspace solid-phase microextraction and gas chromatographic analysis with pulsed flame photometric detection 872(2000)203

Capillary electrophoretic determination of thiosulfate and its oxidation products 879(2000)235

Analysis of organic sulfur compounds in wine aroma 881(2000)569

Analysis of low-volatility organic sulphur compounds in wines by solid-phase microextraction and gas chromatography 881(2000)583

Determination of thiosulfate, thiocyanate and polythionates in a mixture by ion-pair chromatography with ultraviolet absorbance detection 884(2000)81

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Comparison of the thin-layer chromatographic properties of sulfur-containing amino acids and their aminophosphonic analogues 888(2000)335

Determination of glucosinolates in rapeseed and *Thlaspi caerulescens* plants by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 889(2000)75

Specific thiol determination by micellar electrokinetic chromatography and on-column detection reaction with 2,2'-dipyridyldisulfide 895(2000)179

Sunscreen agents

Determination of sunscreen agents in cosmetic products by supercritical fluid extraction and high-performance liquid chromatography 870(2000)199

Determination of sunscreen agents in cosmetic products using microwave-assisted extraction and liquid chromatography 876(2000)243

Surfactant proteins

Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363

Surfactants

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

Qualitative and quantitative analysis of new alkyl amide arginine surfactants by high-performance liquid chromatography and capillary electrophoresis 852(1999)499

Equilibrium analysis of reactions between aromatic anions and nonionic surfactant micelles by capillary zone electrophoresis 853(1999)55

Investigations on the metabolism of alkyl polyglucosides and their determination in waste water by means of liquid chromatography–electrospray mass spectrometry 854(1999)221

Characterization of nonyl phenol ethoxylates in sewage treatment plants by combined precursor ion scanning and multiple reaction monitoring 854(1999)245

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669

Analysis of linear alkylbenzenesulfonates in water samples by large-volume injection-port derivatization and gas chromatography–mass spectrometry 857(1999)359

Chiral separations using polymeric dipeptide surfactants: effect of number of chiral centers and steric factors 858(1999)219

Effect of drying on the degradation of cationic surfactants and separation performance in capillary zone electrophoresis of inorganic anions 863(1999)81

Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants 868(2000)313

Reply to 'Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants' 868(2000)317

Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Micellar electrokinetic chromatography with bis-(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Investigation of the effects of cyclodextrins and organic solvents on the separation of cationic surfactants in capillary electrophoresis 872(2000)269

Determination of anionic surfactants during wastewater recycling process by ion pair chromatography with suppressed conductivity detection 874(2000)207

Analysis of aliphatic alcohol ethoxylates in terms of alkyl and ethylene oxide chain lengths by reversed-phase liquid chromatography with evaporative light scattering detection 874(2000)305

Enantiomer separation of drugs by micellar electrokinetic chromatography using chiral surfactants 875(2000)163

Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography 875(2000)179

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Alkyltrimethylammonium surfactant-mediated extractions: Characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

Simultaneous quantitative trace analysis of anionic and nonionic surfactant mixtures by reversed-phase liquid chromatography 878(2000)99

Capillary zone electrophoretic separation of neutral species of chloro-*s*-triazines in the presence of cationic surfactant monomers 878(2000)137

Statistical evaluation of linear solvation energy relationship models used to characterize chemical selectivity in micellar electrokinetic chromatography 886(2000)245

Assessment of polycrystalline graphites as sorbents for solid-phase microextraction of nonionic surfactants 888(2000)35

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99

Assessment of the pollutant elimination efficiency by gas chromatography/mass spectrometry, liquid chromatography–mass spectrometry and –tandem mass spectrometry. Comparison of conventional and membrane-assisted biological wastewater treatment processes 889(2000)155

Inter-laboratory comparison of liquid chromatographic techniques and enzyme-linked immunosorbent assay for the determination of surfactants in wastewaters 889(2000)195

Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211

Determination of linear alkylbenzenesulfonates in wastewater treatment plants and coastal waters by automated solid-phase extraction followed by capillary electrophoresis–UV detection and confirmation by capillary electrophoresis–mass spectrometry 889(2000)221

High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253

Chiral determination of amino acids by capillary electrophoresis and laser-induced fluorescence at picomolar concentrations 894(2000)259

Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. III: Effects of adsorbed surfactants on quenching the interaction 894(2000)281

The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry 894(2000)345

Selectivity in microemulsion electrokinetic chromatography 897(2000)375

T

Tamoxifen

Non-aqueous capillary electrophoresis of tamoxifen and its acid hydrolysis products 895(2000)81

Tannins

Fractionation of apple procyanidins by size-exclusion chromatography 855(1999)181

Characterisation of hydrolysable tannins from leaves of *Betula pubescens* by high-performance liquid chromatography– mass spectrometry 864(1999)283

Analysis of pigmented high-molecular-mass grape phenolics using ion-pair, normal-phase high-performance liquid chromatography 866(2000)25

High-speed counter-current chromatography of apple procyanidins 886(2000)65

Fractionation of apple procyanidins according to their degree of polymerization by normal-phase high-performance liquid chromatography 890(2000)251

Analysis of oak tannins by liquid chromatography-electrospray ionisation mass spectrometry 891(2000)75

Tantalum

On-line preconcentration of niobium(V) and tantalum(V) as 4-(2-pyridylazo) resorcinol–citrate ternary complexes in geological samples by ion interaction high-performance liquid chromatography 885(2000)369

Tartaric acid

Determination of tartaric acid in solid wine residues by capillary electrophoresis and indirect UV detection 853(1999)181

Ion-chromatographic behavior of alkali metal cations and ammonium ion on zirconium-adsorbing silica gel 884(2000)123

Determination of fumaric acid in apple juice by on-line coupled capillary isotachopheresis–capillary zone electrophoresis with UV detection 891(2000)175

Taurine

Determination of homotaurine as impurity in calcium acamprostate by capillary zone electrophoresis 853(1999)421

Taxoids

Analysis of taxol and major taxoids in Himalayan yew, *Taxus wallichiana* 858(1999)239

Taxol

Analysis of taxol and major taxoids in Himalayan yew, *Taxus wallichiana* 858(1999)239

Teicoplanin

Direct chromatographic resolution of carnitine and *O*-acylcarnitine enantiomers on a teicoplanin-bonded chiral stationary phase 857(1999)145

Enantioseparation of selected *N*-*tert*-butyloxycarbonyl amino acids in high-performance liquid chromatography and capillary electrophoresis with a teicoplanin chiral selector 879(2000)147

Evaluation of the macrocyclic glycopeptide A-40,926 as a high-performance liquid chromatographic chiral selector and comparison with teicoplanin chiral stationary phase 897(2000)113

Enantioselective reversed-phase and non-aqueous capillary electrochromatography using a teicoplanin chiral stationary phase 897(2000)349

Terguride

High-performance liquid chromatography study of the enantiomer separation of chrysanthemic acid and its analogous compounds on a terguride-based stationary phase 859(1999)133

Terpenes

Separation and identification of various carotenoids by C₃₀ reversed-phase high-performance liquid chromatography coupled to UV and atmospheric pressure chemical ionization mass spectrometric detection 854(1999)37

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Monoterpenes in grape juice and wines 881(2000)557

Terpenoids

Determination of octanol–water partition coefficient for terpenoids using reversed-phase high-performance liquid chromatography 864(1999)221

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Tetraalkylammonium bromide

Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants 868(2000)313

Reply to ‘Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants’ 868(2000)317

Tetraalkylammonium salts

Analysis of nonylphenol polyethoxycarboxylates and their related metabolites by on-line derivatization and ion-trap gas chromatography–mass spectrometry 862(1999)113

Tetraborate

Suppressed electrostatic ion chromatography with tetraborate as eluent and its application to the determination of inorganic anions in snow and rainwater 884(2000)161

Tetrabutylammonium phosphate

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Tetracyclines

Chromatographic analysis of tetracycline antibiotics in foods 882(2000)109

Ion chromatographic analysis of tetracyclines using polymeric column and acidic eluent 897(2000)205

Tetradecyldimethylamine

Effect of drying on the degradation of cationic surfactants and separation performance in capillary zone electrophoresis of inorganic anions 863(1999)81

Tetradecyltrimethylammonium bromide

Effect of drying on the degradation of cationic surfactants and separation performance in capillary zone electrophoresis of inorganic anions 863(1999)81

Tetraethylenepentamine

Quantitation of protein binding to the capillary wall in acidic, isoelectric buffers and means for minimizing the phenomenon 894(2000)273

Tetrahydrocannabinol

High-performance liquid chromatographic determination of Δ^9 -tetrahydrocannabinol and the corresponding acid in hemp containing foods with special regard to the fluorescence properties of Δ^9 -tetrahydrocannabinol 872(2000)101

Tetrahydrocannabinolic acid

High-performance liquid chromatographic determination of Δ^9 -tetrahydrocannabinol and the corresponding acid in hemp containing foods with special regard to the fluorescence properties of Δ^9 -tetrahydrocannabinol 872(2000)101

Tetrahydrocarbolinecarboxylic acids

Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Tetrahydro- β -carboline-3-carboxylic acids and contaminants of L-tryptophan 881(2000)501

Tetrahydrocarbolines

Analysis of tetrahydro- β -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Analysis of the bioactive alkaloids tetrahydro- β -carboline and β -carboline in food 881(2000)483

Tetramethylammonium hydroxide

New hydrolysis method for extremely small amount of lipids and capillary gas chromatographic analysis as *N(O)-tert*-butyldimethylsilyl fatty acid derivatives compared with methyl ester derivatives 862(1999)199

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

Tetramethylpyrazine

Analysis of tetramethylpyrazine in *Ephedrae herba* by gas chromatography–mass spectrometry and high-performance liquid chromatography 878(2000)147

Tetramethylrhodamine

Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285

Thalidomide

Simultaneous separation and enantioseparation of thalidomide and its hydroxylated metabolites using high-performance liquid chromatography in common-size columns, capillary liquid chromatography and nonaqueous capillary electrochromatography 876(2000)157

Theaflavins

Comparative analysis of tea catechins and theaflavins by high-performance liquid chromatography and capillary electrophoresis 881(2000)439

Theanine

Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235

Theophyllines

Determination of theophylline and its metabolites in biological samples by liquid chromatography–mass spectrometry 870(2000)87

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Thiamine

Thiamine analysis in biological media by capillary zone electrophoresis with a high-sensitivity cell 871(2000)351

Determination of thiamine by high-performance liquid chromatography 881(2000)267

Thiazides

Liquid chromatographic separation of the stereoisomers of thiazide diuretics 876(2000)221

Thifensulfuron-methyl

Determination of sulfonylurea degradation products in soil by liquid chromatography–ultraviolet detection followed by confirmatory liquid chromatography–tandem mass spectrometry 855(1999)575

Thiocyanate

Determination of thiosulfate, thiocyanate and polythionates in a mixture by ion-pair chromatography with ultraviolet absorbance detection 884(2000)81

Thiocyanomethylthiobenzothiazole

Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261

Thiodiglycol compounds

Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169

Thioglycol methylate

Speciation of dimethylarsinic acid and monomethylarsonic acid by solid-phase microextraction–gas chromatography–ion trap mass spectrometry 873(2000)129

Thiols

Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381

Qualitative and quantitative determination of biologically active low-molecular-mass thiols in human blood by reversed-phase high-performance liquid chromatography with photometry and fluorescence detection 870(2000)433

Determination of biologically active low-molecular-mass thiols in human blood. I. Fast qualitative and quantitative, gradient and isocratic reversed-phase high-performance liquid chromatography with photometric and fluorescence detection 895(2000)156

Determination of biologically active low-molecular-mass thiols in human blood. II. High-performance capillary electrophoresis with photometric detection 895(2000)167

Assays for total homocysteine and other thiols by capillary electrophoresis–laser-induced fluorescence detection. I. Preanalytical condition studies 895(2000)173

Specific thiol determination by micellar electrokinetic chromatography and on-column detection reaction with 2,2'-dipyridyldisulfide 895(2000)179

Thiophanate-methyl

Determination of thiophanate-methyl and its metabolites at trace level in spiked natural water using the supported liquid membrane extraction and the microporous membrane liquid–liquid extraction techniques combined on-line with high-performance liquid chromatography 893(2000)123

Thiopropional-S-oxide

Possible interest of various sample transfer techniques for fast gas chromatography–mass spectrometric analysis of true onion volatiles 896(2000)117

Thioridazine

Direct, preparative enantioselective chromatography of propranolol hydrochloride and thioridazine hydrochloride using carbon dioxide-based mobile phases 865(1999)227

Thiosulfate

Capillary electrophoretic determination of thiosulfate and its oxidation products 879(2000)235

Determination of thiosulfate, thiocyanate and polythionates in a mixture by ion-pair chromatography with ultraviolet absorbance detection 884(2000)81

Thiosulfates

Possible interest of various sample transfer techniques for fast gas chromatography–mass spectrometric analysis of true onion volatiles 896(2000)117

Thorium

Separation and determination of rare earth elements by Dowex 2-X8 resin using sodium trimetaphosphate as elution agent 874(2000)311

Thromboxanes

Solid- and liquid-phase extraction for the gas chromatographic–tandem mass spectrometric quantification of 2,3-dinor-thromboxane B₂ and 2,3-dinor-6-oxo-prostaglandin F_{1α} in human urine 885(2000)351

Tioconazole

Open access capillary electrophoresis. A walk up capillary electrophoresis service for the synthetic chemist 888(2000)219

Tocopherol acetate

Determination of tocopherol acetate in emulsified nutritional supplements by solid-phase extraction and high-performance liquid chromatography with fluorescence detection 881(2000)243

Tocopherols

see also Vitamins

Determination of tocopherols and sterols in vegetable oils by solid-phase extraction and subsequent capillary gas chromatographic analysis 857(1999)231

C₃₀ Stationary phases for the analysis of food by liquid chromatography 880(2000)189

Analysis of olive and hazelnut oil mixtures by high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry of triacylglycerols and gas–liquid chromatography of non-saponifiable compounds (tocopherols and sterols) 881(2000)149

Chromatographic analysis of tocol-derived lipid antioxidants 881(2000)197

Normal-phase high-performance liquid chromatography of tocopherols and tocotrienols. Comparison of different chromatographic columns 881(2000)217

Automated analysis of vitamin E isomers in vegetable oils by continuous membrane extraction and liquid chromatography–electrochemical detection 881(2000)229

Rapid determination of vitamin E in vegetable oils by reversed-phase high-performance liquid chromatography 881(2000)251

Simultaneous determination of α -tocopherol and β -carotene in olive oil by reversed-phase high-performance liquid chromatography 881(2000)255

Quantitative determination of α -, β -, γ - and δ -tocopherols in human serum by high-performance liquid chromatography and gas chromatography–mass spectrometry as trimethylsilyl derivatives with a two-step sample preparation 896(2000)209

Tocotrienols

Chromatographic analysis of tocol-derived lipid antioxidants 881(2000)197

Normal-phase high-performance liquid chromatography of tocopherols and tocotrienols. Comparison of different chromatographic columns 881(2000)217

Tolperisone

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VII. Comparison of chiral recognition ability of ovoglycoprotein from chicken and Japanese quail egg whites 858(1999)155

Toluene

Headspace gas chromatography method for the analysis of volatile impurities in a hormone replacement transdermal patch 859(1999)113

Separation of stereoisomers in a simulated moving bed-supercritical fluid chromatography plant 865(1999)175

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Development of membrane extraction with a sorbent interface–micro gas chromatography system for field analysis 873(2000)13

Evaluation of time-of-flight mass spectrometric detection for fast gas chromatography 878(2000)205

Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

Toluidines

Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277

Toxaphene

Liquid chromatographic profiles of individual compounds of technical toxaphene 886(2000)123

Toxins

see also Aflatoxins, Mycotoxins and Neurotoxins

Competitive immunoassay for staphylococcal enterotoxin A using capillary electrophoresis with laser-induced fluorescence detection 853(1999)545

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatroscan (mark 5) and flame thermionic detection 855(1999)657

Analysis of anatoxin-a in freshwaters by automated on-line derivatization–liquid chromatography–electrospray mass spectrometry 862(1999)191

Reversed-phase liquid chromatographic separation and simultaneous profiling of steroidal glycoalkaloids and their aglycones 863(1999)25

Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13

Isolation and preparative purification of microcystin variants 883(2000)103

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Tramadol

Simultaneous stereoselective analysis of tramadol and its main phase I metabolites by on-line capillary zone electrophoresis–electrospray ionization mass spectrometry 868(2000)295

Capillary electrophoresis with laser-induced fluorescence in clinical drug development. Routine application and future aspects 895(2000)147

Transition metals

see also Metals

Use of ion chromatography for the determination of heavy and transition metals in biochemical samples 857(1999)343

Improved scheme of chelation ion chromatography with a mixed eluent for the simultaneous analysis of transition metals at $\mu\text{g l}^{-1}$ levels 883(2000)127

Trenbolone

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

Determination of trenbolone and zeranol in bovine muscle and liver by liquid chromatography–electrospray mass spectrometry 882(2000)53

Triacylglycerols

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

Analysis of olive and hazelnut oil mixtures by high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry of triacylglycerols and gas–liquid chromatography of non-saponifiable compounds (tocopherols and sterols) 881(2000)149

Triazines

On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279

Determination of polar pesticides with atmospheric pressure chemical ionisation mass spectrometry–mass spectrometry using methanol and/or acetonitrile for solid-phase desorption and gradient liquid chromatography 863(1999)147

Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121

Determination of interfering triazine degradation products by gas chromatography–ion trap mass spectrometry 868(2000)115

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463

Trace-level determination of triazines and several degradation products in environmental waters by disk solid-phase extraction and micellar electrokinetic chromatography 872(2000)299

Determination of triazines in soil leachates by solid-phase microextraction coupled to gas chromatography–mass spectrometry 874(2000)247

Capillary zone electrophoretic separation of neutral species of chloro-*s*-triazines in the presence of cationic surfactant monomers 878(2000)137

Herbicide and plant growth regulator analysis by capillary electrophoresis 880(2000)281

Multiresidue methods using solid-phase extraction techniques for monitoring priority pesticides, including triazines and degradation products, in ground and surface waters 885(2000)217

Identification of reactive dyes in spent dyebaths and wastewater by capillary electrophoresis–mass spectrometry 886(2000)271

Solid-phase extraction with a dibutylmelamine-imprinted polymer as triazine herbicide-selective sorbent 889(2000)25

Tribenuron-methyl

Determination of sulfonylurea degradation products in soil by liquid chromatography–ultraviolet detection followed by confirmatory liquid chromatography–tandem mass spectrometry 855(1999)575

Tributyl phosphate

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent–detergent treatment 852(1999)83

Removal of detergent and solvent from solvent–detergent-treated immunoglobulins 852(1999)87

Gas chromatographic study of complexing in the system hydrazoic acid–tributyl phosphate–nitric acid–uranyl nitrate 859(1999)41

Trichloroethene

High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253

2,4,5-Trichlorophenoxyacetic acid

Chromatographic characterization of molecularly imprinted polymers binding the herbicide 2,4,5-trichlorophenoxyacetic acid 883(2000)119

Trichloropyridinol

Supercritical fluid extraction of chlorpyrifos and 3,5,6-trichloro-2-pyridinol from garden compost 862(1999)105

Trichosanthin

Capillary electrophoretic determination of apoptosis of HeLa cells induced by trichosanthin 894(2000)165

Trichothecenes

Determination of nivalenol and deoxynivalenol in wheat using liquid chromatography–mass spectrometry with negative ion atmospheric pressure chemical ionisation 854(1999)45

Quantification of deoxynivalenol in wheat using an immunoaffinity column and liquid chromatography 859(1999)23

Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473

Simultaneous determination of trichothecene mycotoxins and zearalenone in cereals by gas chromatography–mass spectrometry 882(2000)23

Trichothoxin

Detection of new sequences of peptaibol antibiotics trichothoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179

Triclabendazole

Simultaneous determination of triclabendazole and its sulphoxide and sulphone metabolites in bovine milk by high-performance liquid chromatography 882(2000)99

O-Trifluoroacetylated (–)-menthyl esters

Configurational analysis of chiral acids as O-trifluoroacetylated (–)-menthyl esters by achiral dual-capillary column gas chromatography 891(2000)257

Triglycerides

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Trihalomethanes

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

On-line monitoring trihalomethanes in chlorinated water by membrane introduction–fast gas chromatography mass–spectrometry 893(2000)169

Trimethylhydrazinium propionate

Analysis of carnitine and acylcarnitines in urine by capillary electrophoresis 895(2000)309

Trioctylphosphine oxide

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

Triolein

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

Triple-helical peptides

Use of Edman degradation sequence analysis and matrix-assisted laser desorption/ionization mass spectrometry in designing substrates for matrix metalloproteinases 890(2000)117

Tripolyphosphate

Determination of tripolyphosphate in frozen cod and scallop adductor by ion chromatography 884(2000)89

Triterpenes

Screening for limonoid glucosides in *Citrus tangerina* (Tanaka) Tseng by high-performance liquid chromatography–electrospray ionization mass spectrometry 874(2000)13

Triterpenic acids

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Triterpenoids

High-performance thin-layer chromatographic determination of six major ginsenosides in *Panax ginseng* 868(2000)269

Rapid and sensitive analysis of azadirachtin and related triterpenoids from Neem (*Azadirachta indica*) by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 886(2000)89

Triton X-100

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent–detergent treatment 852(1999)83

Removal of detergent and solvent from solvent–detergent-treated immunoglobulins 852(1999)87

Triton X-114

Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3

Tropane alkaloids

Strategies for supercritical fluid extraction of hyoscyamine and scopolamine salts using basified modifiers 863(1999)47

Enantioseparation of atropine by capillary electrophoresis using sulfated β -cyclodextrin: application to a plant extract 868(2000)285

Trypsin

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

Trypsin inhibitors

High concentrations of sodium chloride facilitate the use of immobilized chymotrypsin for separating virgin forms of specific trypsin inhibitors 852(1999)227

Tryptophan

Determination of indole-3-acetic acid, tryptophan and other indoles in must and wine by high-performance liquid chromatography with fluorescence detection 855(1999)227

Screening for defects in tryptophan metabolism 870(2000)289

Chromatographic resolution of tryptophan enantiomers with L-Leu–L-Leu–L-Leu peptide. Effects of mobile phase composition and chromatographic support 874(2000)21

Tetrahydro- β -carboline-3-carboxylic acids and contaminants of L-tryptophan 881(2000)501

Tryptophol

Determination of indole-3-acetic acid, tryptophan and other indoles in must and wine by high-performance liquid chromatography with fluorescence detection 855(1999)227

Tryptylines

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Tumor necrosis factor α

Histidines in affinity tags and surface clusters for immobilized metal-ion affinity chromatography of trimeric tumor necrosis factor α 852(1999)117

Twaron

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

Tylosins

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Micellar electrokinetic capillary chromatography of macrolide antibiotics. Separation of tylosin, erythromycin and their related substances 857(1999)313

Tyramine

Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113

U**Uranium**

Determination of U(IV) and U(VI) by ion chromatography–inductively coupled plasma mass spectrometry and its application to kinetic studies 884(2000)131

Uranyl nitrate

Gas chromatographic study of complexing in the system hydrazoic acid–tributyl phosphate–nitric acid–uranyl nitrate 859(1999)41

Uridine

Comparision of the enantioseparation of racemic uridine analogs on Whelk-O 1 and ChiralPak-AD columns 866(2000)183

Uronic acid

Analysis of carbohydrates in wood and pulps employing enzymatic hydrolysis and subsequent capillary zone electrophoresis 891(2000)157

V**Vancomycin**

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Multiple-step ligand injection affinity capillary electrophoresis for determining binding constants of ligands to receptors 897(2000)339

Vanillin

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669

Vanillylmandelic acid

Direct measurement of homovanillic, vanillylmandelic and 5-hydroxyindoleacetic acids in urine by capillary electrophoresis 871(2000)341

Verapamil

Enantioselective binding analysis of verapamil to plasma lipoproteins by capillary electrophoresis–frontal analysis 875(2000)447

Verbascoside

Liquid chromatography with electrospray ionisation mass spectrometric detection of phenolic compounds from *Olea europaea* 855(1999)529

Vincamine alkaloids

Separation of the four pairs of enantiomers of vincamine alkaloids by enantioselective high-performance liquid chromatography 893(2000)47

Vinylsulfones

Identification of reactive dyes in spent dyebaths and wastewater by capillary electrophoresis–mass spectrometry 886(2000)271

Vitamins

see also Ascorbic acid, Carotenes, Tocopherols and Retinol

Direct measurement of ascorbic acid biosynthesis in *Arabidopsis* cell suspension culture using capillary electrophoresis 853(1999)381

Elimination of adsorption effects in the analysis of water-soluble vitamins in pharmaceutical formulations by capillary electrophoresis 853(1999)391

Liquid chromatography–nuclear magnetic resonance spectroscopy 856(1999)199

Direct chromatographic resolution of carnitine and *O*-acylcarnitine enantiomers on a teicoplanin-bonded chiral stationary phase 857(1999)145

Analysis of some folate monoglutamates by high-performance liquid chromatography–mass spectrometry. I 864(1999)59

A reversed-phase high-performance liquid chromatographic method to analyze retinal isomers 864(1999)257

Determination of eight water- and fat-soluble vitamins in multi-vitamin pharmaceutical formulations by high-performance liquid chromatography 870(2000)207

Determination of vitamin B₁₂ in multivitamin tablets by multimode high-performance liquid chromatography 870(2000)217

Determination of vitamins A and E in milk powder using supercritical fluid extraction for sample clean-up 874(2000)275

Microemulsion electrokinetic chromatography in suppressed electroosmotic flow environment. Separation of fat-soluble vitamins 876(2000)201

C₃₀ Stationary phases for the analysis of food by liquid chromatography 880(2000)189

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Analysis of olive and hazelnut oil mixtures by high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry of triacylglycerols and gas–liquid chromatography of non-saponifiable compounds (tocopherols and sterols) 881(2000)149

Determination of vitamins in food-matrix Standard Reference Materials 881(2000)171

High-performance liquid chromatography of non-polar retinoid isomers 881(2000)183

Determination of vitamin D₂ in emulsified nutritional supplements by solid-phase extraction and column-switching high-performance liquid chromatography with UV detection 881(2000)189

Chromatographic analysis of tocol-derived lipid antioxidants 881(2000)197

Normal-phase high-performance liquid chromatography of tocopherols and tocotrienols. Comparison of different chromatographic columns 881(2000)217

Automated analysis of vitamin E isomers in vegetable oils by continuous membrane extraction and liquid chromatography–electrochemical detection 881(2000)229

Determination of tocopherol acetate in emulsified nutritional supplements by solid-phase extraction and high-performance liquid chromatography with fluorescence detection 881(2000)243

Rapid determination of vitamin E in vegetable oils by reversed-phase high-performance liquid chromatography 881(2000)251

Simultaneous determination of α -tocopherol and β -carotene in olive oil by reversed-phase high-performance liquid chromatography 881(2000)255

Determination of vitamin K₁ in emulsified nutritional supplements by solid-phase extraction and high-performance liquid chromatography with postcolumn reduction on a platinum catalyst and fluorescence detection 881(2000)261

Determination of thiamine by high-performance liquid chromatography 881(2000)267

Chromatographic determination of riboflavin and its derivatives in food 881(2000)285

Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables 881(2000)309

Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317

Analytical techniques for determining biotin 881(2000)331

Picomolar quantitation of free sulfite in foods by means of [⁵⁷Co]hydroxocobalamin and radiometric chromatography of [⁵⁷Co]sulfitecobalamin. Method, applications and significance of coexisting sulfides 881(2000)345

Determination of organic acids in food samples by capillary zone electrophoresis 881(2000)357

Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425

High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Simultaneous determination of fat-soluble vitamins and provitamins in milk by microcolumn liquid chromatography 891(2000)109

Determination of vitamin B₁₂ in multivitamin tablets and fermentation medium by high-performance liquid chromatography with fluorescence detection 891(2000)243

Quantitative determination of α -, β -, γ - and δ -tocopherols in human serum by high-performance liquid chromatography and gas chromatography–mass spectrometry as trimethylsilyl derivatives with a two-step sample preparation 896(2000)209

Improved method for rapid determination of vitamin A in small samples of breast milk by high-performance liquid chromatography 898(2000)179

Volatile organic compounds

see also Alcohols, Carbonyl compounds and Hydrocarbons

Evaluation of a dual-sorbent trap for monitoring organic compounds in air 855(1999)593

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

Automated gas chromatography with cryogenic/sorbent trap for the measurement of volatile organic compounds in the atmosphere 863(1999)183

Application of gas chromatography–cryocondensation–Fourier transform infrared spectroscopy and gas chromatography–mass spectrometry to the identification of gas phase reaction products from the α -pinene/ozone reaction 864(1999)299

On the performance and inertness of different materials used for the enrichment of sulfur compounds from air and gaseous samples 864(1999)345

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Analysis of volatile contaminants in vegetable oils by headspace solid-phase microextraction with Carboxen-based fibres 873(2000)79

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Characterization of a commercial gas chromatography–flame ionization detection system for the in situ determination of C₅–C₁₀ hydrocarbons in ambient air 878(2000)215

Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

Sorbent trapping of volatile organic compounds from air 885(2000)129

Application of solid-phase microextraction for determination of organic vapours in gaseous matrices 885(2000)405

Role of solid-phase microextraction in the identification of highly volatile pheromones of two Rhinoceros beetles *Scapanes australis* and *Strategus aloeus* (Coleoptera, Scarabaeidae, Dynastinae) 885(2000)433

Evidence for selectivity of absorption of volatile organic compounds by a polydimethylsiloxane solid-phase microextraction fibre 885(2000)457

Fullerenes-extracted soot: a new adsorbent for collecting volatile organic compounds in ambient air 886(2000)313

Importance of handling organic atmospheric pollutants for assessing air quality 889(2000)271

Retention index database for identification of general green leaf volatiles in plants by coupled capillary gas chromatography–mass spectrometry 890(2000)313

Microwave-assisted derivatization of volatile carbonyl compounds with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine 891(2000)267

Application of gas chromatography coupled to chemical ionisation mass spectrometry following headspace solid-phase microextraction for the determination of free volatile fatty acids in aqueous samples 891(2000)287

Time-weighted average sampling of volatile and semi-volatile airborne organic compounds by the solid-phase microextraction device 892(2000)455

Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the ng l⁻¹ concentration level 893(2000)367

Cryogen free automated gas chromatography for the measurement of ambient volatile organic compounds 896(2000)31

Characterisation of whiskeys using solid-phase microextraction with gas chromatography–mass spectrometry 896(2000)351

Characterization of odorant compounds of mussels (*Mytilus edulis*) according to their origin using gas chromatography–olfactometry and gas chromatography–mass spectrometry 896(2000)361

VX

Multiple-technique analytical characterization of a mixture containing chemical-weapons simulant from a munition 883(2000)185

Methanol chemical ionization quadrupole ion trap mass spectrometry of *O*-ethyl *S*-[2-(diisopropylamino)ethyl] methylphosphonothiolate (VX) and its degradation products 893(2000)393

W

Warfarin

Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1

Waxes

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Polymer additive analysis by pyrolysis–gas chromatography. III. Lubricants 891(2000)313

Whey proteins

Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371

Determination of cow's milk in non-bovine and mixed cheeses by capillary electrophoresis of whey proteins in acidic isoelectric buffers 878(2000)261

X

Xanthates

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

Xanthophylls

Liquid chromatography–electrospray mass spectrometry of β -carotene and xanthophylls. Validation of the analytical method 854(1999)233

Xenoestrogens

Quantification of the xenoestrogens 4-*tert*-octylphenol and bisphenol A in water and in fish tissue based on microwave assisted extraction, solid-phase extraction and liquid chromatography–mass spectrometry 864(1999)17

Xylenes

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Development of membrane extraction with a sorbent interface–micro gas chromatography system for field analysis 873(2000)13

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

Z

Zearalanone

Determination of zearalanone in grains by high-performance liquid chromatography–tandem mass spectrometry after solid-phase extraction with RP-18 columns or immunoaffinity columns 858(1999)167

Zearalenone

Determination of zearalenone in grains by high-performance liquid chromatography–tandem mass spectrometry after solid-phase extraction with RP-18 columns or immunoaffinity columns 858(1999)167

Simultaneous determination of trichothecene mycotoxins and zearalenone in cereals by gas chromatography–mass spectrometry 882(2000)23

Zeranol

Determination of trenbolone and zeranol in bovine muscle and liver by liquid chromatography–electrospray mass spectrometry 882(2000)53

Zinc

Ion exchange on resins with temperature-responsive selectivity. IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57

Ion-chromatographic study of the possible absorption of copper and zinc by the skin of *Rana pipiens* 884(2000)143