

## CUMULATIVE SUBJECT INDEX

### J. CHROMATOGR. A VOLS. 852–900

# A

#### **Accelerated solvent extraction**

see Pressurized liquid extraction

#### **Acid–base equilibria**

Dissociation of polyvalent electrolytes 853(1999)35

#### **$\alpha_1$ -Acid glycoprotein**

Comparative studies on the analysis of glycosylation heterogeneity of sialic acid-containing glycoproteins using capillary electrophoresis 866(2000)261

#### **Activation energy**

Mobility and activation energy of single-stranded DNA in denaturing cross-linked polyacrylamide slab gels 894(2000)231

#### **Activity coefficients**

Influence of pH and  $pK_a$  values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

#### **Adenovirus**

Capillary zone electrophoresis of a recombinant adenovirus 895(2000)329

#### **Adsorbents**

see also Affinity adsorbents, Sorbents and Stationary phases

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Large-scale membrane adsorbents 852(1999)73

Evaluation of a dual-sorbent trap for monitoring organic compounds in air 855(1999)593

Heats of adsorption of some organic compounds on  $\beta$ -cyclodextrin determined by gas–solid chromatography 864(1999)293

Low-carbon silica sorbents for solid-phase extraction 868(2000)1

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

Pellicular expanded bed matrix suitable for high flow rates 878(2000)17

Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

Effect of surface heterogeneity on adsorption on solid surfaces. Application of inverse gas chromatography in the studies of energetic heterogeneity of adsorbents 886(2000)133

Fullerenes-extracted soot: a new adsorbent for collecting volatile organic compounds in ambient air 886(2000)313

Adsorption and selectivity characteristics of several human serum proteins with immobilised hard Lewis metal ion–chelate adsorbents 890(2000)95

Solid-phase extraction of polar hydrophilic aromatic sulfonates followed by capillary zone electrophoresis–UV absorbance detection and ion-pair liquid chromatography–diode array UV detection and electrospray mass spectrometry 890(2000)225

DNA adsorption onto polyethylenimine-attached poly(*p*-chloromethylstyrene) beads 898(2000)167

#### Adsorption

Elimination of adsorption effects in the analysis of water-soluble vitamins in pharmaceutical formulations by capillary electrophoresis 853(1999)391

The influence of cell adsorbent interactions on protein adsorption in expanded beds 873(2000)195

Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. II. Effects of adsorbed, neutral polymers on quenching the interaction 874(2000)293

Modified capillary electrophoresis system for peptide, protein and double-stranded DNA analysis 879(2000)189

Suspended bed chromatography, a new approach in downstream processing 890(2000)45

Liquid adsorption chromatography of polyethers: experiments and simulation 890(2000)195

Quantitation of protein binding to the capillary wall in acidic, isoelectric buffers and means for minimizing the phenomenon 894(2000)273

Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. III: Effects of adsorbed surfactants on quenching the interaction 894(2000)281

Protein adsorption on novel acrylamido-based polymeric ion-exchangers. I. morphology and equilibrium adsorption 897(2000)65

Protein adsorption on novel acrylamido-based polymeric ion exchangers. II. Adsorption rates and column behavior 897(2000)81

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

#### Adsorption enthalpy

Microcalorimetric studies of interactions between proteins and hydrophobic ligands in hydrophobic interaction chromatography: effects of ligand chain length, density and the amount of bound protein 872(2000)37

#### Adsorption isotherms

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

Calculation of chromatographic band profiles with an implicit isotherm 862(1999)1

Quantification of single solute and competitive adsorption isotherms using a closed-loop perturbation method 865(1999)51

Determination of the thermodynamic contribution to peak asymmetry of basic solutes in reversed-phase liquid chromatography 865(1999)97

Separation of stereoisomers in a simulated moving bed-supercritical fluid chromatography plant 865(1999)175

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Study on the accuracy of the elution by characteristic point method for the determination of single component isotherms 872(2000)1

Kinetics of solid-phase extraction and solid-phase microextraction in thin adsorbent layer with saturation sorption isotherm 873(2000)39

Determination of mobile phase effect on single-component adsorption isotherm by use of numerical estimation 874(2000)1

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Effect of surface heterogeneity on adsorption on solid surfaces. Application of inverse gas chromatography in the studies of energetic heterogeneity of adsorbents 886(2000)133

#### Adsorption kinetics

Enantiomers separation by simulated moving bed chromatography. Non-instantaneous equilibrium at the solid–fluid interface 865(1999)187

#### Adsorption–partition model

See Partition–adsorption systems

#### Affinity adsorbents

see also Immunoabsorbents and Stationary phases

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105

Reinforcement of frontal affinity chromatography for effective analysis of lectin–oligosaccharide interactions 890(2000)261

Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy 893(2000)235

#### **Affinity capillary electrochromatography**

Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors 875(2000)235

#### **Affinity capillary electrophoresis**

Microscale characterization of the structure–activity relationship of a heparin-binding glycopeptide using affinity capillary electrophoresis and immobilized enzymes 853(1999)189

Interaction of cyclophilin and cyclosporins monitored by affinity capillary electrophoresis 853(1999)469

Congophilicity (Congo red affinity) of different  $\beta_2$ -microglobulin conformations characterized by dye affinity capillary electrophoresis 894(2000)319

#### **Affinity chromatography**

see also Immobilized metal affinity chromatography and Immunoaffinity chromatography

Affinity purification of *Schistosoma japonicum* glutathione-S-transferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

Affinity chromatography of human estrogen receptor- $\alpha$  expressed in *Saccharomyces cerevisiae*. Combination of heparin- and 17 $\beta$ -estradiol-affinity chromatography 852(1999)161

Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken 852(1999)217

Experimental studies on affinity chromatography in an electric field 852(1999)319

Biomimetic dyes as affinity chromatography tools in enzyme purification (Review) 891(2000)33

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

#### **Affinity membranes**

New affinity nylon membrane used for adsorption of  $\gamma$ -globulin 867(2000)161

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

#### **Agarose beads**

Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy 893(2000)235

#### **Agarose gels**

Effect of linear polymer additives on the electroosmotic characteristics of agarose gels in ultrathin-layer electrophoresis 853(1999)511

#### **Agarose–polymer matrices**

Ultra-thin-layer agarose gel electrophoresis. II. Separation of DNA fragments on composite agarose–linear polymer matrices 871(2000)289

#### **Agarose-stainless steel composite**

Pellicular expanded bed matrix suitable for high flow rates 878(2000)17

#### **Air analysis**

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

Separation of carbonyl 2,4-dinitrophenylhydrazones by capillary electrochromatography with diode array detection 853(1999)487

Evaluation of a dual-sorbent trap for monitoring organic compounds in air 855(1999)593

Automated gas chromatography with cryogenic/sorbent trap for the measurement of volatile organic compounds in the atmosphere 863(1999)183

Simultaneous measurement of volatile sulfur compounds using ascorbic acid for oxidant removal and gas chromatography–flame photometric detection 864(1999)111

Application of gas chromatography–cryocondensation–Fourier transform infrared spectroscopy and gas chromatography–mass spectrometry to the identification of gas phase reaction products from the  $\alpha$ -pinene/ozone reaction 864(1999)299

On the performance and inertness of different materials used for the enrichment of sulfur compounds from air and gaseous samples 864(1999)345

2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

Determination of imidacloprid and its metabolite 6-chloronicotinic acid in greenhouse air by high-performance liquid chromatography with diode-array detection 869(2000)497

Characterization of a commercial gas chromatography–flame ionization detection system for the in situ determination of  $C_5$ – $C_{10}$  hydrocarbons in ambient air 878(2000)215

Application of solid-phase microextraction for determination of organic vapours in gaseous matrices 885(2000)405

Importance of handling organic atmospheric pollutants for assessing air quality 889(2000)271

Simultaneous determination of ethylbenzene, indan, indene and acenaphthene in air by capillary gas chromatography 891(2000)361

Determination of 1-(2-methoxyphenyl)piperazine derivatives of isocyanates at low concentrations by temperature-programmed miniaturized liquid chromatography 892(2000)67

Time-weighted average sampling of volatile and semi-volatile airborne organic compounds by the solid-phase microextraction device 892(2000)455

Cryogen free automated gas chromatography for the measurement of ambient volatile organic compounds 896(2000)31

Identification of the oxidation products of the reaction between  $\alpha$ -pinene and hydroxyl radicals by gas and high-performance liquid chromatography with mass spectrometric detection 896(2000)135

#### **Airborne particulate**

Analysis of heavy metals in atmospheric particulate by ion chromatography 888(2000)145

Determination of 1,3-, 1,6-, 1,8-dinitropyrene and 1-nitropyrene in airborne particulate by column liquid chromatography with electrochemical detection 897(2000)393

#### **Albumin stationary phase**

Synthesis of a silica-bonded bovine serum albumin *s*-triazine chiral stationary phase for high-performance liquid chromatographic resolution of enantiomers 866(2000)173

#### **Alkyl fluorenyl stationary phases**

*n*-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133

#### **Alkyl-diol silica columns**

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

#### **Allelochemicals**

Simultaneous determination of phenolic acids and 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one in wheat (*Triticum aestivum* L.) by gas chromatography–tandem mass spectrometry 864(1999)315

#### **Allium spp.**

Possible interest of various sample transfer techniques for fast gas chromatography–mass spectrometric analysis of true onion volatiles 896(2000)117

#### **Alloys**

Quantitative analysis of manganese, chromium and molybdenum by ion-pair reversed-phase high-performance liquid chromatography with pre-column derivatization and UV-visible detection 857(1999)351

#### **Alumina**

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

#### **Amberlite XAD-7 resin**

Removal of detergent and solvent from solvent–detergent-treated immunoglobulins 852(1999)87

#### **Aminophosphonate-bonded silica**

Determination of beryllium in a stream sediment by high-performance chelation ion chromatography 876(2000)127

#### **Amperometric detection**

see also Electrochemical detection

Determination of peroxides by capillary zone electrophoresis with amperometric detection 868(2000)127

Amperometric determination of underivatized amino acids at a nickel-modified gold electrode by anion-exchange chromatography 878(2000)57

#### **Ampholytes**

Elution mechanism in electrostatic ion chromatography with histidine as an isoelectric ampholytic mobile phase 884(2000)287

#### **Amphoteric buffers**

General experimental aspects of the use of isoelectric buffers in capillary electrophoresis 853(1999)71

#### **Amylose stationary phases**

Application and comparison of derivatized cellulose and amylose chiral stationary phases for the separation of enantiomers of pharmaceutical compounds by high-performance liquid chromatography 855(1999)411

#### **Analysis time**

High-speed gas chromatography: an overview of various concepts 856(1999)315

Simultaneous optimization of the analysis time and the concentration detectability in open-tubular liquid chromatography 867(2000)23

**Anion exchangers**

see also Ion exchangers

Application of experimental design for the characterisation of a novel elution system for high-capacity anion chromatography with suppressed conductivity detection 855(1999)91

Quinine versus carbamoylated quinine-based chiral anion exchangers. A comparison regarding enantioselectivity for N-protected amino acids and other chiral acids 858(1999)1

Kinetic study of the mass transfer of bovine serum albumin in anion-exchange chromatography 866(2000)147

Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267

Pressurized-flow anion-exchange capillary electrochromatography using a polymeric ion-exchange stationary phase 890(2000)337

Comparison of chromatographic ion-exchange resins. I. Strong anion-exchange resins 897(2000)99

***Annona cherimola***

High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483

**Annular chromatography**

Scale-down of continuous protein purification by annular chromatography. Design parameters for the smallest unit 890(2000)53

***Apis* spp.**

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. I. High-temperature gas chromatography 855(1999)601

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. II: High-temperature gas chromatography–chemical ionization mass spectrometry 883(2000)75

**Apples**

see also Fruits

Fractionation of apple procyanidins by size-exclusion chromatography 855(1999)181

Retention/quantitation properties of the *o*-phthalaldehyde–3-mercaptopropionic acid and the *o*-phthalaldehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

Determination of fenpyroximate in apples by supercritical fluid extraction and packed capillary liquid chromatography with UV detection 880(2000)121

High-speed counter-current chromatography of apple procyanidins 886(2000)65

Fractionation of apple procyanidins according to their degree of polymerization by normal-phase high-performance liquid chromatography 890(2000)251

Determination of 1-aminocyclopropane-1-carboxylic acid and its structural analogue by liquid chromatography and ion spray tandem mass spectrometry 896(2000)335

**Aqueous two-phase systems**

Purification of protein and recycling of polymers in a new aqueous two-phase system using two thermoseparating polymers 864(1999)31

***Arabidopsis thaliana***

Direct measurement of ascorbic acid biosynthesis in *Arabidopsis* cell suspension culture using capillary electrophoresis 853(1999)381

***Archangelica officinalis***

Separation of coumarins from *Archangelica officinalis* in high-performance liquid chromatography and thin-layer chromatography systems 886(2000)75

**Argemone oil**

Separation, identification and determination of sanguinarine in argemone and other adulterated edible oils by reversed-phase high-performance liquid chromatography 863(1999)123

***Artemisia capillaris***

Fractionation and analysis of *Artemisia capillaris* Thunb. by affinity chromatography with human serum albumin as stationary phase 870(2000)501

**Artifact peaks**

Investigations of artifact peaks in sensitive high-performance liquid chromatography methods 869(2000)85

**Association constants**

Electrophoretic mobility study on ion–ion interactions in an aqueous solution 853(1999)63

Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants 868(2000)313

Reply to ‘Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants’ 868(2000)317

Calculation of stability constants for the chiral selector–enantiomer interactions from electrophoretic mobilities 875(2000)431

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

***Astragalus membranaceus***

Liquid chromatography–electrospray ionization mass spectrometry study of the flavonoids of the roots of *Astragalus mongholicus* and *A. membranaceus* 876(2000)87

***Astragalus mongholicus***

Liquid chromatography–electrospray ionization mass spectrometry study of the flavonoids of the roots of *Astragalus mongholicus* and *A. membranaceus* 876(2000)87

**Atmospheric pressure ionization**

see also Mass spectrometry

Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

State-of-the-art in liquid chromatography–mass spectrometry 856(1999)179

**Atomic emission detection**

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

**Atomic fluorescence detection**

Column-switching system for selenium speciation by coupling reversed-phase and ion-exchange high-performance liquid chromatography with microwave-assisted digestion–hydride generation–atomic fluorescence spectrometry 889(2000)33

**Automation**

Determination of pesticides in waters by automatic on-line solid-phase extraction–capillary electrophoresis 866(2000)137

Automated capillary gas chromatographic system to monitor ethylene emitted from biological materials 868(2000)249

Automated in-tube solid-phase microextraction–high-performance liquid chromatography for carbamate pesticide analysis 873(2000)137

Automated analysis of vitamin E isomers in vegetable oils by continuous membrane extraction and liquid chromatography–electrochemical detection 881(2000)229

Automating solid-phase extraction: current aspects and future prospects 885(2000)97

Cryogen free automated gas chromatography for the measurement of ambient volatile organic compounds 896(2000)31

On-line coupling of solid-phase extraction to gas chromatography with fast solvent vaporization and concentration in an open injector liner. Analysis of pesticides in aqueous samples 896(2000)41

On-line automated high-performance liquid chromatographic determination of total riboflavin phosphates using immobilized acid phosphatase as a pre-column reactor 896(2000)171

***Azadirachta indica***

Rapid and sensitive analysis of azadirachtin and related triterpenoids from Neem (*Azadirachta indica*) by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 886(2000)89

**Azo dyes**

Determination of sulphonated dyes in water by ion-interaction high-performance liquid chromatography 871(2000)227

**B*****Bacillus* spp.**

Isolation and characterization of cyclic  $\alpha$ -(1→4)-glucans having degrees of polymerization 9–31 and their quantitative analysis by high-performance anion-exchange chromatography with pulsed amperometric detection 852(1999)407

**Background electrolyte composition**

see also Buffer composition, Chiral selectors and Organic modifiers

Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudostationary phase. Influence of methanol and acetonitrile on separation selectivity 853(1999)121

Highly alkaline electrolyte for single-stranded DNA separations by electrophoresis in bare silica capillaries 853(1999)371

Protein adsorption to the bare silica wall in capillary electrophoresis. Quantitative study on the chemical composition of the background electrolyte for minimising the phenomenon 868(2000)85

Calculation of electrophoretic mobilities in water–organic modifier mixtures in capillary electrophoresis 868(2000)277

Erratum to: “Optimization of conductivity of low-molecular-mass anions in capillary zone electrophoresis” [J. Chromatogr. A, 850 (1999) 355–362] 868(2000)321

Development and robustness testing of a nonaqueous capillary electrophoresis method for the analysis of nonsteroidal anti-inflammatory drugs 874(2000)121

- Enantiomeric separations of drugs using mixtures of charged and neutral cyclodextrins 875(2000)123
- New amphiphilic aminosaccharide derivatives as chiral selectors in capillary electrophoresis 875(2000)295
- New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169
- Separation of anions by ion chromatography–capillary electrophoresis 879(2000)245
- Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277
- Determination of enantiomers in a synthetic argininal peptide using capillary zone electrophoresis and high-performance liquid chromatography 891(2000)115
- Enantiomer separations by nonaqueous capillary electrophoresis using octakis(2,3-diacetyl-6-sulfato)- $\gamma$ -cyclodextrin 892(2000)499
- Comparison of separation selectivity in capillary electrokinetic chromatography using a cationic linear polymeric pseudo-stationary phase or monomeric additives of similar structure 894(2000)25
- Capillary zone electrophoresis in phosphate buffer – known or unknown? 894(2000)89
- Capillary zone electrophoresis in phosphate buffer – known or unknown? 894(2000)89
- Allylamine– $\beta$ -cyclodextrin copolymer. A novel chiral selector for capillary electrophoresis 894(2000)95
- Capillary electrochromatography and capillary electrochromatography–electrospray mass spectrometry for the separation of non-steroidal anti-inflammatory drugs 895(2000)123
- Simultaneous determination of alkali, alkaline earth and transition metal ions by capillary electrophoresis with indirect UV detection 895(2000)205
- Determination of cationic surfactants by capillary zone electrophoresis and micellar electrokinetic chromatography with deoxycholate micelles in the presence of large organic solvent concentrations 895(2000)227
- Capillary electrophoretic separation of the enantiomers of organophosphates with a phosphorus stereogenic center using the sodium salt of octakis(2,3-diacetyl-6-sulfo)- $\gamma$ -cyclodextrin as resolving agent 895(2000)247
- Simultaneous determination of inorganic anions and cations by capillary electrophoresis with indirect UV detection 895(2000)269
- Highly reproducible capillary zone electrophoresis of humic acids in cyclodextrin- or oligosaccharide-modified background electrolytes 895(2000)345
- Bacteria**  
Pyrolysis–gas–liquid chromatography with atomic emission detection for the identification of *Corynebacterium* species 863(1999)243
- Bacterial signaling molecules**  
Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153
- Band broadening**  
see also Band profiles and Peak shape  
Recent developments in microcolumn liquid chromatography 856(1999)117  
Use of equations for the description of experimental dependence of the height equivalent to a theoretical plate on carrier gas velocity in capillary gas–liquid chromatography 872(2000)111  
Characteristics of particle-loaded monolithic sol–gel columns for capillary electrochromatography. I. Structural, electrical and band-broadening properties 887(2000)277
- Band formation**  
Video-taped sample evaporation in hot chambers simulating gas chromatography split and splitless injectors. II. Injection with band formation 897(2000)247
- Band profiles**  
see also Band broadening and Peak shape  
Dynamics of capillary electrochromatography. II. Comparison of column efficiency parameters in microscale high-performance liquid chromatography and capillary electrochromatography 855(1999)349  
Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. III. Results obtained with Kromasil C<sub>18</sub> columns 855(1999)423  
Calculation of chromatographic band profiles with an implicit isotherm 862(1999)1  
Visualization of sample introduction in liquid chromatographic columns. Contribution of a flow distributor on the sample band shape 865(1999)83  
Investigations of artifact peaks in sensitive high-performance liquid chromatography methods 869(2000)85

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C<sub>18</sub> (2) columns 869(2000)181

Determination of mobile phase effect on single-component adsorption isotherm by use of numerical estimation 874(2000)1

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Control of dispersion in capillary electrochromatography coupled to UV and mass spectrometric detection 887(2000)409

#### **Bed consolidation**

Evaluation of the secondary consolidation of columns for liquid chromatography by ultrasonic irradiation 878(2000)153

#### **Bed porosity**

Hydrodynamic aspects of slurry packing processes in microcolumn liquid chromatography 883(2000)11

#### **Beer**

Determination of sulphur compounds in beer using headspace solid-phase microextraction and gas chromatographic analysis with pulsed flame photometric detection 872(2000)203

Electrophoretic methods in the analysis of beverages 880(2000)243

Chromatographic determination of amino acid enantiomers in beers and raw materials used for their manufacture 881(2000)81

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321

#### **Beeswax**

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. I: High-temperature gas chromatography 855(1999)601

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. II: High-temperature gas chromatography–chemical ionization mass spectrometry 883(2000)75

#### ***Betula pubescens***

Characterisation of hydrolysable tannins from leaves of *Betula pubescens* by high-performance liquid chromatography–mass spectrometry 864(1999)283

#### **Beverages**

see also Beer, Soft drinks, Tea and Wine

Semipreparative reversed-phase liquid chromatographic fractionation of aroma extracts from wine and other alcoholic beverages 864(1999)77

Analysis of tetrahydro-β-carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23

Gas chromatographic–mass spectrometric analysis of acids and phenols in distilled alcohol beverages. Application of anion-exchange disk extraction combined with in-vial elution and silylation 873(2000)29

Determination of aldehydes in food by high-performance liquid chromatography with biosensor coupling and micromembrane suppressors 880(2000)233

Preparation of 4,4'-(1-[<sup>2</sup>H<sub>6</sub>]methylene)bis-[2,3,5,6-<sup>2</sup>H<sub>4</sub>]-phenol and its application to the measurement of bisphenol A in beverages by stable isotope dilution mass spectrometry 883(2000)163

Characterisation of whiskeys using solid-phase microextraction with gas chromatography–mass spectrometry 896(2000)351

#### **Binding constants**

Calculation of stability constants for the chiral selector–enantiomer interactions from electrophoretic mobilities 875(2000)431

Multiple-step ligand injection affinity capillary electrophoresis for determining binding constants of ligands to receptors 897(2000)339

#### **Binding studies**

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

Enantioselective binding analysis of verapamil to plasma lipoproteins by capillary electrophoresis–frontal analysis 875(2000)447

#### **Biochemical detection**

High-performance liquid chromatography with on-line coupled UV, mass spectrometric and biochemical detection for identification of acetylcholinesterase inhibitors from natural products 872(2000)61

High-performance liquid chromatography coupled to enzyme-amplified biochemical detection for the analysis of hemoglobin after pre-column biotinylation 886(2000)19



**Biodiesel**

Analytical monitoring of the production of biodiesel by high-performance liquid chromatography with various detection methods 858(1999)13

**Biofilms**

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

**Biosensors**

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

Determination of aldehydes in food by high-performance liquid chromatography with biosensor coupling and micromembrane suppressors 880(2000)233

Monitoring and control of industrial downstream processing of sugar beet molasses 882(2000)329

Capillary electrophoresis coupled to biosensor detection 892(2000)143

**Biotrickling filter**

Determination of methacrylic acid in the drain of a biotrickling filter using isotachopheresis and capillary zone electrophoresis 862(1999)237

***Blighia sapida***

Isolation of pure (2*S*,1'*S*,2'*S*)-2-(2'-carboxycyclopropyl)glycine from *Blighia sapida* (Akee) 873(2000)283

**Blue-green alga**

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

**Book Reviews**

Size exclusion chromatography (by S. Mori and H.G. Barth) 877(2000)239

Analytical pyrolysis of natural organic polymers (by S.C. Moldoreanu) 877(2000)241

**Breakthrough volume**

Contributions of theory to method development in solid-phase extraction 885(2000)17

**Brine**

Determination of inorganic cations in brine solutions by ion chromatography 884(2000)223

**Buffer composition**

see also Background electrolyte composition, Chiral selectors and Organic modifiers

Separation of proteins and peptides by capillary electrophoresis in acid buffers containing high concentrations of surfactants 852(1999)325

General experimental aspects of the use of isoelectric buffers in capillary electrophoresis 853(1999)71

Use of acidic and basic pH and calcium ion addition in the capillary zone electrophoretic characterization of recombinant human deoxyribonuclease, a complex phosphoglycoprotein 853(1999)295

Stepwise capillary electrophoretic separation of DNA fragments using poly(ethylene oxide) solutions in the presence of electroosmotic flow 853(1999)337

Capillary electrophoresis of double-stranded DNA in an untreated capillary 853(1999)349

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669

Separation of double-stranded DNA in conventional and isoelectric buffers: studies on stability and separation performance 859(1999)87

Efficiency studies in nonaqueous capillary electrophoresis 859(1999)203

Effects of organic modifiers on retention mechanism and selectivity in micellar electrokinetic capillary chromatography studied by linear solvation energy relationships 863(1999)69

Capillary electrophoretic behaviour of humic substances in physical gels 863(1999)195

Nonaqueous capillary electrophoresis equipped with amperometric detection for analysis of chlorinated phenolic compounds 864(1999)323

Simultaneous stereoselective analysis of tramadol and its main phase I metabolites by on-line capillary zone electrophoresis–electrospray ionization mass spectrometry 868(2000)295

Migration behavior and separation of benzenediamines, aminophenols and benzenediols by capillary zone electrophoresis 871(2000)357

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. II. Effects of adsorbed, neutral polymers on quenching the interaction 874(2000)293

Enantiomeric separations by nonaqueous capillary electrophoresis 875(2000)277

- Non-aqueous capillary electrophoretic enantioseparation of *N*-derivatized amino acids using cinchona alkaloids and derivatives as chiral counter-ions 875(2000)353
- Microemulsion electrokinetic chromatography in suppressed electroosmotic flow environment. Separation of fat-soluble vitamins 876(2000)201
- Optimization of separation and migration behavior of cephalosporins in capillary zone electrophoresis 879(2000)197
- Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs 883(2000)249
- Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297
- Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241
- Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309
- High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. I. Effect of buffer composition on the gradient shape 890(2000)25
- High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. II. Separation of proteins 890(2000)37
- Non-aqueous capillary electrophoresis 892(2000)155
- Background theory and applications of microemulsion electrokinetic chromatography 892(2000)171
- Characterization of synthetic polyelectrolytes by capillary electrophoresis 894(2000)105
- Effect of ionic strength, pH and polymer concentration on the separation of DNA fragments in the presence of electroosmotic flow 894(2000)219
- Chiral determination of amino acids by capillary electrophoresis and laser-induced fluorescence at picomolar concentrations 894(2000)259
- Influence of the structure of cyclodextrins and amino acid sequence of dipeptides and tripeptides on the pH-dependent reversal of the migration order in capillary electrophoresis 894(2000)267
- Quantitation of protein binding to the capillary wall in acidic, isoelectric buffers and means for minimizing the phenomenon 894(2000)273
- Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. III: Effects of adsorbed surfactants on quenching the interaction 894(2000)281
- Capillary electrophoresis of seed 2S albumins from *Lupinus* species 894(2000)297
- The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry 894(2000)345
- Chiral capillary electrophoresis as predictor for separation of drug enantiomers in continuous flow zone electrophoresis 895(2000)51
- Separation of bisphenol A and three alkylphenols by micellar electrokinetic chromatography 895(2000)213
- Analysis of enzymatically glucosylated flavonoids by capillary electrophoresis 895(2000)279
- Effects of amine modifiers on the separation of tetramethylrhodamine-labeled mono- and oligosaccharides by capillary zone electrophoresis 895(2000)285
- Capillary zone electrophoresis of a recombinant adenovirus 895(2000)329
- Highly selective and efficient determination of US Environmental Protection Agency priority phenols employing solid-phase extraction and non-aqueous capillary electrophoresis 896(2000)95
- Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133
- Building materials**  
Determination of ergosterol on mouldy building materials using isotope dilution and gas chromatography–tandem mass spectrometry 898(2000)227
- Butter**  
Erratum to: “Trace-level determination of polar flavour compounds in butter by solid-phase extraction and gas chromatography–mass spectrometry” [J. Chromatogr. A, 844 (1999) 295–305] 873(2000)299
- Determination of cholesterol oxides in processed food using high-performance liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 896(2000)321

# C

## Calibration curves

Ion-chromatographic study of the possible absorption of copper and zinc by the skin of *Rana pipiens* 884(2000)143

Ion-chromatographic assay of chromium(VI)-containing semiconductor etchants. Statistical comparison of two columns 884(2000)151

## Calibration, compound-independent

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

## Capillary array electrophoresis

Sub-microliter DNA sequencing for capillary array electrophoresis 894(2000)191

## Capillary array scanner

Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299

## Capillary columns

see also Coated capillaries and Packed capillary columns

Chiral separations by open tubular capillary electrokinetic chromatography 853(1999)151

Capillary electrochromatography of peptides in a microfabricated system 853(1999)257

High-speed gas chromatography: an overview of various concepts 856(1999)315

Capillary electrochromatography of proteins and peptides with porous-layer open-tubular columns 858(1999)91

Capillary zone electrophoretic analysis of positively charged poly(ethylene oxide) macromolecules using non-covalent polycation-coated fused-silica capillary and indirect UV detection 864(1999)335

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Preparation of various silica-based columns for capillary electrochromatography by in-column derivatization 873(2000)247

Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography 873(2000)293

Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. II. Effects of adsorbed, neutral polymers on quenching the interaction 874(2000)293

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Distribution coefficients of *n*-alkanes measured on wall-coated capillary columns 879(2000)157

Modified capillary electrophoresis system for peptide, protein and double-stranded DNA analysis 879(2000)189

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

Open tubular capillary electrokinetic chromatography in etched fused-silica tubes (Review) 887(2000)31

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

Advances in column technology and instrumentation in capillary electrochromatography 892(2000)257

Capillary electrochromatography on silica columns: factors influencing performance 892(2000)279

Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23

Investigation of the nematic–isotropic transition of a liquid crystalline polymer and determination of molecular diffusion coefficients using gas chromatography 893(2000)359

Life-time studies with capillary electrochromatography columns operated under different conditions 894(2000)35

Unexpected surface chemistry in capillaries for electrophoresis 894(2000)73

Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79

Quantitation of protein binding to the capillary wall in acidic, isoelectric buffers and means for minimizing the phenomenon 894(2000)273

Quantitative studies on the adsorption of proteins to the bare silica wall in capillary electrophoresis. III: Effects of adsorbed surfactants on quenching the interaction 894(2000)281

Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133

#### **Capillary columns, hydrophilic coated**

Influence of pH\*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63

#### **Capillary electrophoresis**

see also Affinity capillary electrophoresis, Electrokinetic chromatography, Electrophoresis, Isoelectric focusing, Isotachopheresis, Micellar electrokinetic chromatography, Non-aqueous capillary electrochromatography, Non-aqueous capillary electrophoresis, Partial-filling capillary electrophoresis and Special Issue Index

Overview of capillary electrophoresis and capillary electrochromatography 856(1999)443

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261

Alternative methods providing enhanced sensitivity and selectivity in capillary electroseparation experiments 892(2000)203

Standardization of capillary zone electropherograms obtained by using field enhanced sample stacking 894(2000)3

New method for standardization of electropherograms obtained in capillary zone electrophoresis 894(2000)11

Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117

Chiral capillary electrophoresis as predictor for separation of drug enantiomers in continuous flow zone electrophoresis 895(2000)51

Determination of soluble anions and cations from waters of pulp and paper mills with on-line coupled capillary electrophoresis 895(2000)189

Robustness testing for a capillary electrophoresis method using the "short-end injection" technique 897(2000)329

#### **Capillary electrophoresis–biosensors detection**

Capillary electrophoresis coupled to biosensor detection 892(2000)143

#### **Capillary electrophoresis–chemiluminescence nitrogen detection**

Apparatus and method for interfacing capillary electrophoresis and chemiluminescence nitrogen detection for the analysis of nitrogen-containing compounds 868(2000)121

#### **Capillary electrophoresis immunoassay**

Determination of morphine by capillary electrophoresis immunoassay in thermally reversible hydrogel-modified buffer and laser-induced fluorescence detection 895(2000)1

#### **Capillary electrophoresis–inductively coupled plasma atomic emission spectrometry**

Interface for capillary electrophoresis coupled with inductively coupled plasma atomic emission spectrometry 853(1999)141

#### **Capillary electrophoresis–ion chromatography**

Separation of anions by ion chromatography–capillary electrophoresis 879(2000)245

#### **Capillary electrophoresis–mass spectrometry**

see also Mass spectrometry and Special Issue Index

New sheathless interface for coupling capillary electrophoresis to electrospray mass spectrometry evaluated by the analysis of fatty acids and prostaglandins 854(1999)141

Investigation of Cr(III) hydrolytic polymerisation products by capillary electrophoresis–inductively coupled plasma-mass spectrometry 872(2000)227

Capillary electrophoresis–inductively coupled plasma mass spectrometry interface with minimised dead volume for high separation efficiency 891(2000)129

Capillary electrophoresis and off-line capillary electrophoresis–electrospray ionization quadrupole time-of-flight tandem mass spectrometry of carbohydrates 895(2000)291

#### **Capillary electrophoresis mobility shift assay**

DNA-binding affinities of MyoD and E47 homo- and heterodimers by capillary electrophoresis mobility shift assay 862(1999)231

#### ***Capsicum annuum***

Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annuum* L.) powders 889(2000)41

#### ***Capsicum frutescens***

Separation and quantitation of colour pigments of chili powder (*Capsicum frutescens*) by high-performance liquid chromatography–diode array detection 896(2000)69

**Cardboard**

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

***Carpinus betulus***

Retention index database for identification of general green leaf volatiles in plants by coupled capillary gas chromatography–mass spectrometry 890(2000)313

***Carthamus tinctorius***

Analysis of natural food pigments by capillary electrophoresis 880(2000)311

**Catalysts**

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum–rhodium alloy catalysts 852(1999)507

Purity testing of organometallic catalysts by packed capillary supercritical fluid chromatography 868(2000)261

**Cation exchangers**

see also Ion exchangers

Pore size distributions of cation-exchange adsorbents determined by inverse size-exclusion chromatography 883(2000)39

**Cell**

Capillary electrophoretic determination of apoptosis of HeLa cells induced by trichosanthin 894(2000)165

**Cell adhesion**

The influence of cell adsorbent interactions on protein adsorption in expanded beds 873(2000)195

**Cellobiohydrolase stationary phases**

Temperature-induced inversion of elution order in the enantioseparation of sotalol on a cellobiohydrolase I-based stationary phase 858(1999)33

Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1

**Cellulose stationary phases**

Application and comparison of derivatized cellulose and amylose chiral stationary phases for the separation of enantiomers of pharmaceutical compounds by high-performance liquid chromatography 855(1999)411

**Central composite design**

Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215

Capillary electrophoresis analysis of gentamicin sulphate with UV detection after pre-capillary derivatization with 1,2-phthalic dicarboxaldehyde and mercaptoacetic acid 895(2000)67

Robustness testing for a capillary electrophoresis method using the “short-end injection” technique 897(2000)329

**Centrifugal partition chromatography**

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

Improvement purification of sulfated oligofucan by ion-exchange displacement centrifugal partition chromatography 869(2000)353

**Centrifugation**

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

**Cereals**

Erratum to: “Liquid chromatographic method for determining added folic acid in fortified cereal products” [J. Chromatogr. A, 826 (1998) 235–240] 867(2000)289

Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides 881(2000)1

Electrophoresis of cereal storage proteins 881(2000)23

Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175

**Characteristic point method**

Study on the accuracy of the elution by characteristic point method for the determination of single component isotherms 872(2000)1

**Charge coupled device detection**

High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93

Capillary electrophoresis with laser-induced fluorescence in clinical drug development. Routine application and future aspects 895(2000)147

**Charge distribution**

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

**Charged porous particles**

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

**Cheese**

Amperometric determination of underivatized amino acids at a nickel-modified gold electrode by anion-exchange chromatography 878(2000)57

- Determination of cow's milk in non-bovine and mixed cheeses by capillary electrophoresis of whey proteins in acidic isoelectric buffers 878(2000)261
- Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides 881(2000)1
- Evaluation of the effect of rennet type on casein proteolysis in an ovine milk cheese by means of capillary electrophoresis 881(2000)59
- Effect of the activity levels of the added proteolytic enzyme mixture on free amino acids in ripening Ossau-Iraty cheese 881(2000)69
- Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79
- Chelidonium majus**  
Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69
- Chemical ionization, methanol**  
Methanol chemical ionization quadrupole ion trap mass spectrometry of *O*-ethyl *S*-[2-(diisopropylamino)ethyl] methylphosphonothiolate (VX) and its degradation products 893(2000)393
- Chemiluminescence detection**  
Chemiluminescence detection of heme proteins separated by capillary isoelectric focusing 852(1999)597
- Determination of polyphenols by high-performance liquid chromatography with inhibited chemiluminescence detection 855(1999)171
- Capillary electrophoresis separation and permanganate chemiluminescence on-line detection of some alkaloids with  $\beta$ -cyclodextrin as an additive 855(1999)329
- Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271
- 5-Amino-4-sulfanylphthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1
- Chemometrics**  
see also specific methods
- Application of several modified peak purity assays to real complex multicomponent mixtures by high-performance liquid chromatography with diode-array detection 855(1999)487
- Pyrolysis–gas–liquid chromatography with atomic emission detection for the identification of *Corynebacterium* species 863(1999)243
- Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41
- Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95
- Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331
- Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215
- Substitution of fish species detected by thin-layer isoelectric focusing and a computer-assisted method for the evaluation of gels 880(2000)303
- Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197
- Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annum* L.) powders 889(2000)41
- Optimization methods in chromatography and capillary electrophoresis 892(2000)75
- Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149
- Exploratory chemometric analysis of the classification of pharmaceutical substances based on chromatographic data 897(2000)23
- Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227
- Chip technology**  
Nanoliter-scale sample preparation methods directly coupled to polymethylmethacrylate-based microchips and gel-filled capillaries for the analysis of oligonucleotides 853(1999)107
- Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271
- Novel microfabricated device for electrokinetically induced pressure flow and electrospray ionization mass spectrometry 892(2000)195
- Integration of a microextraction system. Solvent extraction of a Co–2-nitroso-5-dimethylaminophenol complex on a microchip 894(2000)19

Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45

Rapid, parallel separations of D1S80 alleles in a plastic microchannel chip 894(2000)203

### Chiral recognition

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a  $\beta$ -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

Enhancement of chiral recognition by formation of a sandwiched complex in capillary electrophoresis 875(2000)361

Experimental support differentiating two proposed chiral recognition models for the resolution of *N*-(3,5-Dinitrobenzoyl)- $\alpha$ -arylalkylamines on high-performance liquid chromatography chiral stationary phases 886(2000)47

### Chiral selectivity

Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383

### Chiral selectors

see also Buffer composition and Mobile phase composition

Chiral selectors from fruit: application of citrus pectins to enantiomer separations in capillary electrophoresis 857(1999)285

Quinine versus carbamoylated quinine-based chiral anion exchangers. A comparison regarding enantioselectivity for *N*-protected amino acids and other chiral acids 858(1999)1

Chiral resolution of basic drugs by capillary electrophoresis with new glycosaminoglycans 864(1999)163

Chiral separation of amino acids and peptides by capillary electrophoresis 875(2000)43

Enantioselective determination by capillary electrophoresis with cyclodextrins as chiral selectors 875(2000)89

Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors 875(2000)235

Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255

Enantiomeric separations by nonaqueous capillary electrophoresis 875(2000)277

New amphiphilic aminosaccharide derivatives as chiral selectors in capillary electrophoresis 875(2000)295

Non-aqueous capillary electrophoretic enantioseparation of *N*-derivatized amino acids using cinchona alkaloids and derivatives as chiral counter-ions 875(2000)353

Calculation of stability constants for the chiral selector–enantiomer interactions from electrophoretic mobilities 875(2000)431

Mechanistic study of opposite migration order of dimethindene enantiomers in capillary electrophoresis in the presence of native  $\beta$ -cyclodextrin and heptakis(2,3,6-tri-*O*-methyl)- $\beta$ -cyclodextrin 875(2000)455

Separation of brompheniramine enantiomers by capillary electrophoresis and study of chiral recognition mechanisms of cyclodextrins using NMR-spectroscopy, UV spectrometry, electrospray ionization mass spectrometry and X-ray crystallography 875(2000)471

Enantioseparation of selected *N*-*tert*-butyloxycarbonyl amino acids in high-performance liquid chromatography and capillary electrophoresis with a teicoplanin chiral selector 879(2000)147

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

*tert*-Butylcarbamoylquinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique 892(2000)509

Enantioseparation of anionic analytes by non-aqueous capillary electrophoresis using quinine and quinidine derivatives as chiral counter-ions 894(2000)63

Allylamine- $\beta$ -cyclodextrin copolymer. A novel chiral selector for capillary electrophoresis 894(2000)95

Influence of the structure of cyclodextrins and amino acid sequence of dipeptides and tripeptides on the pH-dependent reversal of the migration order in capillary electrophoresis 894(2000)267

### Chiral stationary phases

see also Enantiomer separation

### Chiral stationary phases, electrochromatography

Separation of enantiomers by packed capillary electrochromatography on a cellulose-based stationary phase 875(2000)331

Investigation of an enantioselective non-aqueous capillary electrochromatography system applied to the separation of chiral acids 875(2000)341

Chiral anion exchangers applied to capillary electrochromatography enantioseparation of oppositely charged chiral analytes: investigation of stationary and mobile phase parameters 887(2000)421

Enantioseparations in non-aqueous capillary electrochromatography using polysaccharide type chiral stationary phases 887(2000)439

- Enantiomer separations by capillary electrochromatography using chiral stationary phases 887(2000)457
- Electrochromatography 892(2000)219
- Enantiomeric separations of cationic and neutral compounds by capillary electrochromatography with monolithic chiral stationary phases of  $\beta$ -cyclodextrin-bonded negatively charged polyacrylamide gels 893(2000)177
- Enantioselective reversed-phase and non-aqueous capillary electrochromatography using a teicoplanin chiral stationary phase 897(2000)349
- Chiral stationary phases, GC**
- Enantiomer separation of  $\alpha$ -ionone using gas chromatography with cyclodextrin derivatives as chiral stationary phases 865(1999)201
- Chiral stationary phases, LC**
- Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383
- Application and comparison of derivatized cellulose and amylose chiral stationary phases for the separation of enantiomers of pharmaceutical compounds by high-performance liquid chromatography 855(1999)411
- Discrimination between enantioselective and non-selective binding sites on cellobiohydrolase-based stationary phases by site specific competing ligands 857(1999)107
- Uniform-sized molecularly imprinted polymer for (*S*)-ibuprofen. Retention properties in aqueous mobile phases 857(1999)117
- Direct chromatographic resolution of carnitine and *O*-acylcarnitine enantiomers on a teicoplanin-bonded chiral stationary phase 857(1999)145
- Quinine versus carbamoylated quinine-based chiral anion exchangers. A comparison regarding enantioselectivity for N-protected amino acids and other chiral acids 858(1999)1
- Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VII. Comparison of chiral recognition ability of ovoglycoprotein from chicken and Japanese quail egg whites 858(1999)155
- High-performance liquid chromatography study of the enantiomer separation of chrysanthemic acid and its analogous compounds on a terguride-based stationary phase 859(1999)133
- Convenient synthesis of  $\pi$ -acceptor chiral stationary phases for high-performance liquid chromatography from halogen-substituted 3,5-dinitrobenzoylamides 859(1999)143
- Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1
- Synthesis of a silica-bonded bovine serum albumin *s*-triazine chiral stationary phase for high-performance liquid chromatographic resolution of enantiomers 866(2000)173
- Liquid chromatographic resolution of 2-hydroxycarboxylic acids on a new chiral stationary phase derived from (*S*)-leucine 868(2000)31
- Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic  $\beta$ -substituted  $\alpha$ -amino acids on a teicoplanin chiral stationary phase 868(2000)197
- Novel cinchona alkaloid carbamate  $C_9$ -dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111
- Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395
- Stereoselective binding of 2,3-substituted 3-hydroxypropionic acids on an immobilised human serum albumin chiral stationary phase: stereochemical characterisation and quantitative structure–retention relationship study 876(2000)75
- Liquid chromatographic separation of the stereoisomers of thiazide diuretics 876(2000)221
- Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95
- Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109
- Large degree of racemization observed in the amide bond forming reaction on silica gel 878(2000)165
- Experimental support differentiating two proposed chiral recognition models for the resolution of *N*-(3,5-Dinitrobenzoyl)- $\alpha$ -arylalkylamines on high-performance liquid chromatography chiral stationary phases 886(2000)47
- Investigation of mandelic acid bonding on Pirkle type chromatographic stationary phases by Raman spectroscopy 893(2000)37
- Separation of the four pairs of enantiomers of vincamine alkaloids by enantioselective high-performance liquid chromatography 893(2000)47
- Evaluation of the macrocyclic glycopeptide A-40,926 as a high-performance liquid chromatographic chiral selector and comparison with teicoplanin chiral stationary phase 897(2000)113



Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic  $\beta$ -substituted  $\alpha$ -amino acids on a copper(II)-*D*-penicillamine chiral stationary phase 897(2000)145

Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate  $\beta$ -cyclodextrin bonded chiral stationary phase 898(2000)53

Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63

#### **Chiral stationary phases, SFC**

Application of molecularly imprinted polymers in supercritical fluid chromatography 897(2000)317

#### **Chlorinated water**

On-line monitoring trihalomethanes in chlorinated water by membrane introduction–fast gas chromatography mass–spectrometry 893(2000)169

#### **Chlorine-selective detection**

Development and characterization of chlorine-selective pulsed discharge emission detector for gas chromatography 872(2000)141

Applications using the chlorine-selective pulsed discharge emission detector 872(2000)167

Characterization of chlorinated compounds using a dual chlorine-selective pulsed discharge emission detector–helium-pulsed discharge photoionization detector system 872(2000)179

#### **Chromarod**

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatrosan (mark 5) and flame thermionic detection 855(1999)657

#### **Chromatofocusing**

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. I. Effect of buffer composition on the gradient shape 890(2000)25

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. II. Separation of proteins 890(2000)37

#### **Cigarette smoke**

see also Tobacco

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection 867(2000)187

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

#### **Circular dichroism detection**

Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181

#### **Citrus tangerina**

Screening for limonoid glucosides in *Citrus tangerina* (Tanaka) Tseng by high-performance liquid chromatography–electrospray ionization mass spectrometry 874(2000)13

#### **Clays**

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279

#### **Clean-up methods**

see also Extraction methods, Preconcentration, Sample handling, Sample preparation and Special Issue Index

Recent developments in microcolumn liquid chromatography 856(1999)117

Does further clean-up reduce the matrix enhancement effect in gas chromatographic analysis of pesticide residues in food? 868(2000)51

Evaluation of a new automated cleanup system for the analysis of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in environmental samples 893(2000)383

Application of ion-exchange cartridge clean-up in food analysis. V. Simultaneous determination of sulphonamide antibacterials in animal liver and kidney using high-performance liquid chromatography with ultraviolet and mass spectrometric detection 898(2000)95

#### **Closed-loop perturbation**

Quantification of single solute and competitive adsorption isotherms using a closed-loop perturbation method 865(1999)51

#### **Closed-loop stripping analysis**

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

Retention index database for identification of general green leaf volatiles in plants by coupled capillary gas chromatography–mass spectrometry 890(2000)313

**Cloud point preconcentration**

Exploratory chemometric analysis of the classification of pharmaceutical substances based on chromatographic data 897(2000)23

Cloud point preconcentration of rather polar compounds: application to the high-performance liquid chromatographic determination of priority pollutant chlorophenols 897(2000)171

**Cluster analysis**

Substitution of fish species detected by thin-layer isoelectric focusing and a computer-assisted method for the evaluation of gels 880(2000)303

**Coated capillaries**

see also Capillary columns

Influence of pH\*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Enantiomeric separation by capillary electrophoresis with an electroosmotic flow-controlled capillary 875(2000)315

Novel, trifunctional diamine for silica coating in capillary zone electrophoresis 894(2000)53

Capillary zone electrophoresis of a recombinant adenovirus 895(2000)329

**Column design**

Influence of column design on process-scale ion-exchange chromatography 865(1999)3

Improved column preparation and performance in capillary electrochromatography 887(2000)367

**Column efficiency**

Influence of column radial heterogeneity on peak fronting in linear chromatography 857(1999)69

Comparison of the repeatability of quantitative data measured in high-performance liquid chromatography with UV and atmospheric pressure chemical ionization mass spectrometric detection 873(2000)163

**Column fabrication**

Aspects of column fabrication for packed capillary electrochromatography 887(2000)347

**Column geometry**

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

**Column headers**

Visualization of viscous fingering in high-performance liquid chromatographic columns. Influence of the header design 865(1999)73

**Column packing**

Comparison of different packing methods for capillary electrochromatography columns 887(2000)307

**Column radial heterogeneity**

Influence of column radial heterogeneity on peak fronting in linear chromatography 857(1999)69

Influence of the column radial heterogeneity on the determination of single-component isotherms by the elution by characteristic point method 891(2000)217

**Column reproducibility**

Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. II. Minimizing column reproducibility problems 897(2000)51

**Column stability**

Studies of retention and stability of a horizontally polymerized bonded phase for reversed-phase liquid chromatography 868(2000)153

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

**Column switching**

see also Coupled columns

Determination of 4-demethoxy-3'-deamino-3'-aziridinyl-4'-methylsulphonyldaunorubicin and its 13-hydroxy metabolite by direct injection of human plasma into a column-switching liquid chromatography system with mass spectrometric detection 854(1999)81

Determination of ammonium in seawater by column-switching ion chromatography 868(2000)209

Determination of theophylline and its metabolites in biological samples by liquid chromatography–mass spectrometry 870(2000)87

Determination of the *N*-methyl-D-aspartate receptor NR2B subunit blocker Ro 63-1908 in rat, cynomolgus monkey and human plasma by high-performance liquid chromatography with column switching and fluorescence detection 870(2000)189

Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153

Direct analysis of food samples by high-performance liquid chromatography 880(2000)149

Determination of vitamin D<sub>2</sub> in emulsified nutritional supplements by solid-phase extraction and column-switching high-performance liquid chromatography with UV detection 881(2000)189

Simplified column-switching technology for the determination of traces of anions in the presence of high concentrations of other anions 884(2000)53

Column-switching system for selenium speciation by coupling reversed-phase and ion-exchange high-performance liquid chromatography with microwave-assisted digestion–hydride generation–atomic fluorescence spectrometry 889(2000)33

Analysis of neuropeptide Y and its metabolites by high-performance liquid chromatography–electrospray ionization mass spectrometry and integrated sample clean-up with a novel restricted-access sulphonic acid cation exchanger 890(2000)135

Analysis of the enzymatic racemization of D-aspartic acid to L-aspartic acid by the on-line coupling of a solid-phase extraction column and a ligand-exchange high-performance liquid chromatography column 896(2000)299

Analysis of phenols in pyrolysis oils by gel permeation chromatography and multidimensional liquid chromatography 896(2000)343

#### Column temperature

Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3

Control of column temperature in reversed-phase liquid chromatography 869(2000)211

#### Columns

On the need to retain unregulated choice of high-performance liquid chromatography columns 852(1999)603

Ultra-short columns for low-pressure ion chromatography 857(1999)175

Trace analysis of pesticide residues in water by high-speed narrow-bore capillary gas chromatography–mass spectrometry with programmable temperature vaporizer 874(2000)81

#### Combinatorial chemistry

Solid-phase extraction for combinatorial libraries 885(2000)305

#### Comb-shaped polymer

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

#### Cometary nucleus analysis

Gas chromatography for in situ analysis of a cometary nucleus: characterization and optimization of diphenyl/dimethylpolysiloxane stationary phases 863(1999)157

#### Complex formation stoichiometry

Ion exchange on resins with temperature-responsive selectivity. III. Influence of complex formation stoichiometry on temperature dependence of resin selectivity 868(2000)143

#### Complexation

see also Derivatization

Gas chromatographic study of complexing in the system hydrazoic acid–tributyl phosphate–nitric acid–uranyl nitrate 859(1999)41

On-line preconcentration of niobium(V) and tantalum(V) as 4-(2-pyridylazo) resorcinol–citrate ternary complexes in geological samples by ion interaction high-performance liquid chromatography 885(2000)369

Simultaneous determination of alkali, alkaline earth and transition metal ions by capillary electrophoresis with indirect UV detection 895(2000)205

Separation selectivity of anionic metal complexes of *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid in ion and ion electrokinetic chromatography 895(2000)255

#### Complexation constants

Calculation of stability constants for the chiral selector–enantiomer interactions from electrophoretic mobilities 875(2000)431

#### Compost

Supercritical fluid extraction of chlorpyrifos and 3,5,6-trichloro-2-pyridinol from garden compost 862(1999)105

#### Computer simulation

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. II. Two-run assay procedures 857(1999)21

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. III. Improving the accuracy of computer simulation 857(1999)41

Computer-aided development of a high-performance liquid chromatographic method for the determination of hydroxyanthraquinone derivatives in Chinese herb medicine rhubarb 857(1999)167

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Use of optimization software to determine rugged analysis conditions in high-performance liquid chromatography 868(2000)13

Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Erratum to: "Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles" [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

Hydrodynamic aspects of slurry packing processes in microcolumn liquid chromatography 883(2000)11

Importance of handling organic atmospheric pollutants for assessing air quality 889(2000)271

Liquid adsorption chromatography of polyethers: experiments and simulation 890(2000)195

Simulation of electrophoretic separations by the flux-corrected transport method 890(2000)321

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487

#### **Concentration detectability**

Simultaneous optimization of the analysis time and the concentration detectability in open-tubular liquid chromatography 867(2000)23

#### **Concentration waves, thermo-induced**

Ion exchange on resins with temperature-responsive selectivity. IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57

#### **Conductivity detection**

Application of experimental design for the characterisation of a novel elution system for high-capacity anion chromatography with suppressed conductivity detection 855(1999)91

Non-suppressed conductivity and indirect UV detection of carboxylic acids in environmental samples by ion-exclusion chromatography using 2,6-pyridinedicarboxylic acid eluent 859(1999)173

Erratum to: "Optimization of conductivity of low-molecular-mass anions in capillary zone electrophoresis" [J. Chromatogr. A, 850 (1999) 355–362] 868(2000)321

Determination of silicate in water by ion exclusion chromatography with conductivity detection 874(2000)143

Conductivity detection of polymerase chain reaction products separated by micro-reversed-phase liquid chromatography 883(2000)1

Enhanced conductometric detection of cyanide in suppressed ion chromatography 884(2000)75

Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261

#### **Confectioneries**

Determination of sugar alcohols in confectioneries by high-performance liquid chromatography after nitrobenzoylation 893(2000)195

#### **Confocal micro-Raman spectroscopy**

Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy 893(2000)235

#### **Confocal scanning laser microscopy**

Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy 893(2000)235

#### **Continuous beds**

Continuous-bed chromatography for the analysis and purification of recombinant human basic fibroblast growth factor 857(1999)137

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Erratum to: "Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles" [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

Nonaqueous electrochromatography on continuous bed columns of sol-gel bonded large-pore C<sub>18</sub> material: separation of retinyl esters 890(2000)347

#### **Continuous subcritical water extraction**

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

**Coriolis force**

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

***Cornus officinalis***

Separation of gallic acid from *Cornus officinalis* Sieb. et Zucc by high-speed counter-current chromatography 886(2000)309

***Corynebacterium* spp.**

Pyrolysis–gas–liquid chromatography with atomic emission detection for the identification of *Corynebacterium* species 863(1999)243

**Cosmetics**

Determination of sunscreen agents in cosmetic products by supercritical fluid extraction and high-performance liquid chromatography 870(2000)199

Determination of sunscreen agents in cosmetic products using microwave-assisted extraction and liquid chromatography 876(2000)243

Screening method for determining the presence of N-nitrosodiethanolamine in cosmetics by open-tubular capillary electrochromatography 887(2000)497

**Cotton**

Pressurised liquid extraction of pentachlorophenol from dyestuffs and cotton 873(2000)287

**Counter-current chromatography**

see also Centrifugal partition chromatography

Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69

Preparative isolation and purification of notopterin and isoimperatorin from *Notopterygium forbesii* Boiss (Chinese traditional medicinal herb) by high-speed counter-current chromatography 883(2000)67

High-speed counter-current chromatography of apple procyanidins 886(2000)65

Relationship between retention, linear velocity and flow for counter-current chromatography 886(2000)283

Separation of gallic acid from *Cornus officinalis* Sieb. et Zucc by high-speed counter-current chromatography 886(2000)309

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N,N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

Supercritical fluid extraction of catechins from *Cratoxylum prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

**Coupled columns**

see also Column switching

Semi-permeable surface analytical reversed-phase column for the improved trace analysis of acidic pesticides in water with coupled-column reversed-phase liquid chromatography with UV detection. Determination of bromoxynil and bentazone in surface water 858(1999)45

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

New operational modes for multidimensional and comprehensive gas chromatography by using cryogenic modulation 866(2000)203

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Time-resolved cryogenic modulation for targeted multidimensional capillary gas chromatography analysis 892(2000)15

Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117

***Cratoxylum prunifolium***

Supercritical fluid extraction of catechins from *Cratoxylum prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

**Critical effects**

Robustness testing of a reversed-phase high-performance liquid chromatographic assay: comparison of fractional and asymmetrical factorial designs 874(2000)167

**Critical micellar concentration**

Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants 868(2000)313

Reply to 'Determination of critical micelle concentration and interactions between cephalosporins and charged surfactants' 868(2000)317

**Cross-linking**

Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63

***Croton* oil**

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

**Cryogenic trapping**

New operational modes for multidimensional and comprehensive gas chromatography by using cryogenic modulation 866(2000)203

Time-resolved cryogenic modulation for targeted multidimensional capillary gas chromatography analysis 892(2000)15

Possible interest of various sample transfer techniques for fast gas chromatography–mass spectrometric analysis of true onion volatiles 896(2000)117

**Crystalline phases**

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

**Curve resolution**

Application of several modified peak purity assays to real complex multicomponent mixtures by high-performance liquid chromatography with diode-array detection 855(1999)487

**Cyano bonded phase**

Dependence of cyano bonded phase hydrolytic stability on ligand structure and solution pH 893(2000)245

**Cyclodextrin adsorbents**

Heats of adsorption of some organic compounds on  $\beta$ -cyclodextrin determined by gas–solid chromatography 864(1999)293

**Cyclodextrin stationary phases**

Pressure-controlled high-performance liquid chromatographic study on the influence of rim chemistry on partial molar volume differences between free and complexed cyclodextrins 863(1999)127

Insoluble  $\beta$ -cyclodextrin polymer for capillary gas chromatographic separation of enantiomers and isomers 883(2000)137

**D****Dairy products**

Determination of ammonium in milk and dairy products by ion chromatography 893(2000)133

**Deamidation**

Determination of ammonium in milk and dairy products by ion chromatography 893(2000)133

**Deconvolution**

Laser-induced dispersed fluorescence detection of polycyclic aromatic compounds in soil extracts separated by capillary electrochromatography 883(2000)231

**Densitometry**

High-performance thin-layer chromatography method for monitoring norfloxacin residues on pharmaceutical equipment surfaces 862(1999)209

**Density**

Different elution modes and field programming in gravitational field-flow fractionation. 2. Experimental verification of the range of conditions for flow-rate and carrier liquid density programming 868(2000)217

**Deprotonation**

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

**Derivatization**

see also Complexation

**Derivatization, electrochromatography**

Preparation of various silica-based columns for capillary electrochromatography by in-column derivatization 873(2000)247

**Derivatization, electrophoresis**

3-Aminobenzamide and 3-aminobenzoic acid, tags for capillary electrophoresis of complex carbohydrates with laser-induced fluorescent detection 863(1999)205

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

Micellar electrokinetic capillary chromatographic separation and fluorescent detection of amino acids derivatized with 4-fluoro-7-nitro-2,1,3-benzoxadiazole 876(2000)183

Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using  $\gamma$ -cyclodextrin 877(2000)217

Enantioseparation of anionic analytes by non-aqueous capillary electrophoresis using quinine and quinidine derivatives as chiral counter-ions 894(2000)63

Methyl- $\beta$ -cyclodextrin modified micellar electrokinetic capillary chromatography with laser-induced fluorescence for separation and detection of phospholipids 894(2000)129

Rapid analysis of covalently and non-covalently fluorophore-labeled proteins using ultra-thin-layer sodium dodecylsulfate gel electrophoresis 894(2000)329

Capillary electrophoresis analysis of gentamicin sulphate with UV detection after pre-capillary derivatization with 1,2-phthalic dicarboxaldehyde and mercaptoacetic acid 895(2000)67

Analysis of carnitine and acylcarnitines in urine by capillary electrophoresis 895(2000)309

#### **Derivatization, GC**

Gas chromatographic determination of primary alcohols as their formylbenzoic esters by gas-phase luminescence detection 855(1999)583

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. I. High-temperature gas chromatography 855(1999)601

Analysis of linear alkylbenzenesulfonates in water samples by large-volume injection-port derivatization and gas chromatography–mass spectrometry 857(1999)359

Erratum to: “Determination of fatty acid amides as trimethylsilyl derivatives by gas chromatography with mass spectrometric detection” [J. Chromatogr. 849 (1999) 541–552] 859(1999)119

Pre-column derivatization and gas chromatographic determination of alkaloids in bulbs of *Fritillaria* 859(1999)183

Solid-phase microextraction coupled with gas chromatography–ion trap mass spectrometry for the analysis of haloacetic acids in water 859(1999)159

Analysis of nonylphenol polyethoxycarboxylates and their related metabolites by on-line derivatization and ion-trap gas chromatography–mass spectrometry 862(1999)113

New hydrolysis method for extremely small amount of lipids and capillary gas chromatographic analysis as *N(O)-tert.-butyldimethylsilyl* fatty acid derivatives compared with methyl ester derivatives 862(1999)199

Gas chromatographic profiling and screening for phenols as isobutoxycarbonyl derivatives in aqueous samples 866(2000)87

Gas chromatographic quantification of amino acid enantiomers in food matrices by their *N(O,S)-ethoxycarbonyl* heptafluorobutyl ester derivatives 866(2000)221

Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215

Enantiomeric separation and discrimination of 2-hydroxy acids as *O*-trifluoroacetylated (*S*)-(+)-3-methyl-2-butyl esters by achiral dual-capillary column gas chromatography 874(2000)91

Determination of acetic acid in aqueous samples, by water-phase derivatization, solid-phase microextraction and gas chromatography 874(2000)225

Gas chromatographic determination of organomercury following aqueous derivatization with sodium tetraethylborate and sodium tetraphenylborate. Comparative study of gas chromatography coupled with atomic fluorescence spectrometry, atomic emission spectrometry and mass spectrometry 876(2000)147

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert.*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97

Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175

Solid-phase microextraction with on-fibre derivatization applied to the analysis of volatile carbonyl compounds 886(2000)175

Experimental design of a microwave-assisted extraction–derivatization method for the analysis of methylmercury 889(2000)185

Microwave-assisted derivatization of volatile carbonyl compounds with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine 891(2000)267

Determination of hair polyamines as *N*-ethoxycarbonyl-*N*-pentafluoropropionyl derivatives by gas chromatography–mass spectrometry 897(2000)295

#### **Derivatization, LC**

Enzyme amplification as detection tool in continuous-flow systems. II. On-line coupling of liquid chromatography to enzyme-amplified biochemical detection after pre-column derivatization with biotin 855(1999)397

Determination of gas phase peroxyacetic acid using pre-column derivatization with organic sulfide reagents and liquid chromatography 855(1999)551

Enantiomeric purity determination of acetyl-L-carnitine by reversed-phase high-performance liquid chromatography using chiral derivatization 857(1999)127

Use of ion chromatography for the determination of heavy and transition metals in biochemical samples 857(1999)343

Liquid chromatographic determination of aniline in table-top sweeteners based on pre-column derivatization with 1,2-naphthoquinone-4-sulfonate 859(1999)227

- High-performance liquid chromatography of methanol released from pectins after its oxidation to formaldehyde and condensation with 2,4-dinitrophenylhydrazine 863(1999)227
- Liquid chromatography–mass spectrometry method for the determination of aldehydes derivatized by the Hantzsch reaction 864(1999)191
- 2-(4-Carboxyphenyl)-6-*N,N*-diethylaminobenzofuran: a useful reagent for the sensitive determination of alcohols by high-performance liquid chromatography with fluorimetric detection 869(2000)243
- Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39
- Retention/quantitation properties of the *o*-phthaldialdehyde–3-mercaptopropionic acid and the *o*-phthaldialdehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271
- Determination of lipoic acid and dihydrolipoic acid in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 870(2000)381
- Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection 870(2000)473
- New fluorimetric method of liquid chromatography for the determination of the neurotoxin domoic acid in seafood and marine phytoplankton 871(2000)1
- Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75
- Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115
- Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159
- Liquid chromatographic determination of biogenic amines in fermented foods after derivatization with 3,5-dinitrobenzoyl chloride 881(2000)517
- High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365
- Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175
- Determination of neurotoxin 3-*N*-oxalyl-2,3-diaminopropionic acid and non-protein amino acids in *Lathyrus sativus* by precolumn derivatization with 1-fluoro-2,4-dinitrobenzene 883(2000)113
- Characterization of a chemical artifact in the liquid chromatographic determination of 3-butyn-2-one using the 2,4-dinitrophenylhydrazine method 886(2000)303
- Post-column oxidative derivatization for the liquid chromatographic determination of phenothiazines 890(2000)281
- Determination of enantiomers in a synthetic argininal peptide using capillary zone electrophoresis and high-performance liquid chromatography 891(2000)115
- Derivatization of ephedrine with *o*-phthaldialdehyde for liquid chromatography after treatment with sodium hypochlorite 893(2000)69
- Determination of sugar alcohols in confectioneries by high-performance liquid chromatography after nitrobenzoylation 893(2000)195
- Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111
- 5-Amino-4-sulfanylphthalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1
- Derivatization, pyrolysis**  
Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235
- Derivatization, SFE**  
Esterification of decanoic acid during supercritical fluid extraction employing either methanol-modified carbon dioxide or a methanol trap 858(1999)201
- Simultaneous extraction and derivatization of 2-chlorovinylarsonous acid from soils using supercritical and pressurized fluids 888(2000)327
- Simultaneous supercritical fluid derivatization and extraction of formaldehyde by the Hantzsch reaction 896(2000)51
- Derivatization, SPE**  
Solid-phase extraction of phenols 885(2000)291
- Detection, electrochromatography**  
Capillary electrochromatography with thermo-optical absorbance detection for the analysis of phenylthiohydantoin-amino acids 853(1999)131
- Use of on-line mass spectrometric detection in capillary electrochromatography (Review) 887(2000)85



Control of dispersion in capillary electrochromatography coupled to UV and mass spectrometric detection 887(2000)409

#### Detection, electrophoresis

Chemiluminescence detection of heme proteins separated by capillary isoelectric focusing 852(1999)597

Labeling effects on the isoelectric point of green fluorescent protein 853(1999)21

Application of non-aqueous capillary electrophoresis with electrochemical detection to the determination of nicotine in tobacco 853(1999)27

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159

Indirect detection of organic acids in non-aqueous capillary electrophoresis 853(1999)171

Side-entry excitation and detection of square capillary array electrophoresis for DNA sequencing 853(1999)359

Capillary electrophoresis coupled with laser-induced fluorescence polarization as a hybrid approach to ultrasensitive immunoassays 853(1999)555

Electrokinetic chromatography 856(1999)465

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Microchip capillary electrophoresis using on-line chemiluminescence detection 867(2000)271

Determination of diclofenac sodium by capillary zone electrophoresis with electrochemical detection 868(2000)101

Apparatus and method for interfacing capillary electrophoresis and chemiluminescence nitrogen detection for the analysis of nitrogen-containing compounds 868(2000)121

Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299

Utilization of fluorescein sodium salt for the indirect fluorimetric detection in micellar electrokinetic chromatography 872(2000)247

On-column amperometric detection in capillary electrophoresis with an improved high-voltage electric field decoupler 872(2000)259

Analysis of glycosaminoglycan monosaccharides by capillary electrophoresis using indirect laser-induced fluorescence detection 873(2000)269

Detection techniques in ion analysis: what are our choices? 884(2000)3

Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261

Determination of ebrotidine and its metabolites by capillary electrophoresis with UV and mass spectrometry detection 888(2000)281

Separation of metalloporphyrins by capillary electrophoresis with UV detection and inductively coupled plasma mass spectrometric detection 888(2000)293

Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309

Simple interface for capillary electrophoresis–inductively coupled plasma atomic emission spectrometry 891(2000)139

Miniaturized thin-layer radial flow cell with interdigitated ring-shaped microarray electrode used as amperometric detector for capillary electrophoresis 891(2000)149

Non-aqueous capillary electrophoresis 892(2000)155

Direct determination of amino acids and carbohydrates by high-performance capillary electrophoresis with refractometric detection 894(2000)253

Rapid analysis of covalently and non-covalently fluorophore-labeled proteins using ultra-thin-layer sodium dodecylsulfate gel electrophoresis 894(2000)329

Non-aqueous capillary electrophoresis of tamoxifen and its acid hydrolysis products 895(2000)81

Capillary electrophoresis with laser-induced fluorescence in clinical drug development. Routine application and future aspects 895(2000)147

Specific thiol determination by micellar electrokinetic chromatography and on-column detection reaction with 2,2'-dipyridyldisulfide 895(2000)179

Determination of aromatic amines in water samples by capillary electrophoresis with electrochemical and fluorescence detection 895(2000)197

#### Detection, FIA

Enzyme amplification as detection tool in continuous-flow systems. I. Development of an enzyme-amplified biochemical detection system coupled on-line to flow-injection analysis 855(1999)383

#### Detection, GC

Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535

- Gas chromatographic determination of primary alcohols as their formylbenzoic esters by gas-phase luminescence detection 855(1999)583
- Gas chromatography with spectroscopic detectors 856(1999)349
- Analysis of organotin compounds by gas chromatography–reactive-flow detection 858(1999)245
- Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection 867(2000)187
- Comparison of gas chromatography–pulsed flame photometric detection–mass spectrometry, automated mass spectral deconvolution and identification system and gas chromatography–tandem mass spectrometry as tools for trace level detection and identification 868(2000)229
- Towards a quantitative definition of perfection in chromatographic analyses 868(2000)305
- Pulsed discharge emission detector: an element-selective detector for gas chromatography 872(2000)119
- Development and characterization of chlorine-selective pulsed discharge emission detector for gas chromatography 872(2000)141
- Applications using the chlorine-selective pulsed discharge emission detector 872(2000)167
- Characterization of chlorinated compounds using a dual chlorine-selective pulsed discharge emission detector–helium-pulsed discharge photoionization detector system 872(2000)179
- Determination of sulphur compounds in beer using headspace solid-phase microextraction and gas chromatographic analysis with pulsed flame photometric detection 872(2000)203
- Gas chromatographic determination of organomercury following aqueous derivatization with sodium tetraethylborate and sodium tetraphenylborate. Comparative study of gas chromatography coupled with atomic fluorescence spectrometry, atomic emission spectrometry and mass spectrometry 876(2000)147
- Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133
- Surface-ionization methods and devices of indication and identification of nitrogen-containing base molecules 896(2000)3
- Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149
- Characterization of odorant compounds of mussels (*Mytilus edulis*) according to their origin using gas chromatography–olfactometry and gas chromatography–mass spectrometry 896(2000)361
- Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227
- Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201
- Detection, LC**
- Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59
- Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361
- Reversed-phase ion-pair chromatography coupled to electrospray ionisation mass spectrometry by on-line removal of the counter-ions 854(1999)155
- Application of experimental design for the characterisation of a novel elution system for high-capacity anion chromatography with suppressed conductivity detection 855(1999)91
- Determination of polyphenols by high-performance liquid chromatography with inhibited chemiluminescence detection 855(1999)171
- Enzyme amplification as detection tool in continuous-flow systems. II. On-line coupling of liquid chromatography to enzyme-amplified biochemical detection after pre-column derivatization with biotin 855(1999)397
- Liquid chromatography–nuclear magnetic resonance spectroscopy 856(1999)199
- Liquid chromatography–Fourier-transform infrared spectrometry 856(1999)213
- Size-exclusion chromatography with on-line ultraviolet, proton nuclear magnetic resonance and mass spectrometric detection and on-line collection for off-line Fourier transform infrared spectroscopy 857(1999)89
- Analytical monitoring of the production of biodiesel by high-performance liquid chromatography with various detection methods 858(1999)13
- Time-resolved high-performance liquid chromatography fluorescence detector using a nanosecond pulsed light source for detecting lanthanide-chelated compounds 859(1999)13

- Postcolumn fluorescence as an alternative to evaporative light scattering detection for ceramide analysis with gradient elution in non-aqueous reversed-phase liquid chromatography 859(1999)99
- Non-suppressed conductivity and indirect UV detection of carboxylic acids in environmental samples by ion-exclusion chromatography using 2,6-pyridinedicarboxylic acidic eluent 859(1999)173
- Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169
- High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93
- Size-exclusion chromatography–multiangle laser light scattering analysis of  $\beta$ -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105
- Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261
- Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275
- Quantitative analysis of hydrophobic pulmonary surfactant proteins by high-performance liquid chromatography with light-scattering detection 870(2000)363
- Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181
- High-performance liquid chromatography with on-line coupled UV, mass spectrometric and biochemical detection for identification of acetylcholinesterase inhibitors from natural products 872(2000)61
- Ion-exclusion controlled size-exclusion chromatography of methacrylic acid–methyl methacrylate copolymers 872(2000)91
- Erratum to: “Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes”. [J. Chromatogr. A, 852 (1999) 59–71] 872(2000)323
- Speciation of selenium compounds with ion-pair reversed-phase liquid chromatography using inductively coupled plasma mass spectrometry as element-specific detection 874(2000)55
- Using a differential refractive index detector as a pressure transducer for online viscometry in exclusion chromatography 878(2000)1
- Amperometric determination of underivatized amino acids at a nickel-modified gold electrode by anion-exchange chromatography 878(2000)57
- Determination of aldehydes in food by high-performance liquid chromatography with biosensor coupling and micromembrane suppressors 880(2000)233
- Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159
- Conductivity detection of polymerase chain reaction products separated by micro-reversed-phase liquid chromatography 883(2000)1
- Detection techniques in ion analysis: what are our choices? 884(2000)3
- Enhanced conductometric detection of cyanide in suppressed ion chromatography 884(2000)75
- Determination of inorganic and organic anionic arsenic species in water by ion chromatography coupled to hydride generation–inductively coupled plasma atomic emission spectrometry 884(2000)211
- High-performance liquid chromatography coupled to enzyme-amplified biochemical detection for the analysis of hemoglobin after pre-column biotinylation 886(2000)19
- Column-switching system for selenium speciation by coupling reversed-phase and ion-exchange high-performance liquid chromatography with microwave-assisted digestion–hydride generation–atomic fluorescence spectrometry 889(2000)33
- Determination of naphthalenesulfonates in water by on-line ion-pair solid-phase extraction and ion-pair liquid chromatography with fast-scanning fluorescence detection 890(2000)289
- Determination of steroid sex hormones and related synthetic compounds considered as endocrine disrupters in water by liquid chromatography–diode array detection–mass spectrometry 892(2000)391
- Screening of homocysteine from newborn blood spots by high-performance liquid chromatography with coulometric array detection 896(2000)183
- 5-Amino-4-sulfanylphtalhydrazide as a chemiluminescence derivatization reagent for aromatic aldehydes in liquid chromatography 898(2000)1
- Detection, SFC**  
Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

**Detection, TLC**

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatroskan (mark 5) and flame thermionic detection 855(1999)657

The state of the art in thin-layer chromatography–mass spectrometry: a critical appraisal 856(1999)429

Fibre optical scanning with high resolution in thin-layer chromatography 898(2000)265

**Deuterium exchange**

Selective deuterium exchange during superheated heavy water chromatography–nuclear magnetic resonance spectroscopy–mass spectrometry of sulfonamides 886(2000)289

**Dialysis**

see also Microdialysis

Membrane-based sample preparation coupled on-line to chromatography or electrophoresis 856(1999)55

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Direct analysis of food samples by high-performance liquid chromatography 880(2000)149

**Diastereomer separation**

Chiral separation of amino acids and peptides by capillary electrophoresis 875(2000)43

Determination of enantiomers in a synthetic argininal peptide using capillary zone electrophoresis and high-performance liquid chromatography 891(2000)115

Chiral determination of amino acids by capillary electrophoresis and laser-induced fluorescence at picomolar concentrations 894(2000)259

**Diesel fuel**

Extraction of hydrocarbon contamination from soils using accelerated solvent extraction 874(2000)217

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

**Diets**

Determination of folate vitamers in food and in Italian reference diet by high-performance liquid chromatography 855(1999)237

**Differential signal detection**

Differential signal detection system for improved analysis of overlapping signals in liquid chromatography 859(1999)221

**Diffusion**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Protein diffusion in charged polyacrylamide gels. Visualization and analysis 865(1999)155

Protein adsorption on novel acrylamido-based polymeric ion exchangers. II. Adsorption rates and column behavior 897(2000)81

**Diffusion coefficients**

Kinetic study of the mass transfer of bovine serum albumin in anion-exchange chromatography 866(2000)147

Investigation of the nematic–isotropic transition of a liquid crystalline polymer and determination of molecular diffusion coefficients using gas chromatography 893(2000)359

**Diffusion tubes**

Dynamic system for the calibration of semi-volatile carbonyl compounds in air 883(2000)171

**Dihexyldiethylcarbamoyl methylenephosphonate stationary phases**

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N,N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

**Diphenyl/dimethylpolysiloxane stationary phases**

Gas chromatography for in situ analysis of a cometary nucleus: characterization and optimization of diphenyl/dimethylpolysiloxane stationary phases 863(1999)157

***Diplodus vulgaris***

Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99

**Direct immersion**

Applications of solid-phase microextraction in food analysis 880(2000)35

**Direct injection**

Analysis of methyl *tert*-butyl ether and its degradation products by direct aqueous injection onto gas chromatography with mass spectrometry or flame ionization detection systems 857(1999)205

Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151

**Disinfection by-products**

Simultaneous determination of inorganic disinfection by-products and the seven standard anions by ion chromatography 890(2000)295

**Dispersion**

see also Matrix solid-phase dispersion

Control of dispersion in capillary electrochromatography coupled to UV and mass spectrometric detection 887(2000)409

**Displacement chromatography**

Polycations as displacer in high-performance bioseparation 865(1999)27

Optimization of ion-exchange displacement separations. I. Validation of an iterative scheme and its use as a methods development tool 876(2000)51

Optimization of ion-exchange displacement separations. II. Comparison of displacement separations on various ion-exchange resins 876(2000)63

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

**Displacers**

Polycations as displacer in high-performance bioseparation 865(1999)27

**Dissociation constants**

Dissociation of polyvalent electrolytes 853(1999)35

Dissociation constants of phenols in methanol–water mixtures 867(2000)45

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Determination of sulphonated dyes in water by ion-interaction high-performance liquid chromatography 871(2000)227

Influence of pH and  $pK_a$  values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Determination of dissociation constants of cytokinins by capillary zone electrophoresis 878(2000)249

Reinforcement of frontal affinity chromatography for effective analysis of lectin–oligosaccharide interactions 890(2000)261

**Distribution coefficients**

Distribution coefficients of *n*-alkanes measured on wall-coated capillary columns 879(2000)157

Liquid adsorption chromatography of polyethers: experiments and simulation 890(2000)195

**Distribution constants**

Behavior of lanthanides in countercurrent chromatography using dihexyl-*N,N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137

**Distribution equilibrium**

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

**Disulfide linkages**

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

**DNA sequencing**

Nanoliter-scale sample preparation methods directly coupled to polymethylmethacrylate-based microchips and gel-filled capillaries for the analysis of oligonucleotides 853(1999)107

**Doehlert's experimental design**

Optimization of glucosinolate separation by micellar electrokinetic capillary chromatography using a Doehlert's experimental design 864(1999)155

**Doping analysis**

Cyclodextrin-mediated micellar electrokinetic chromatography and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry for the enantiomer separation of racemorphan in human urine 853(1999)439

**Dovetailed capillaries**

Enantiomeric separation by capillary electrophoresis with an electroosmotic flow-controlled capillary 875(2000)315

**Drug adsorption**

Micellar liquid chromatography for prediction of drug transport 870(2000)1

**Dust**

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

**Dynamic capacity**

Comparison of chromatographic ion-exchange resins. I. Strong anion-exchange resins 897(2000)99

**Dynamically modified silica capillary electrochromatography**

Capillary electrochromatography with a silica column with a dynamically modified cationic surfactant 855(1999)137

# E

**Efficiency**

Dynamics of capillary electrochromatography. II. Comparison of column efficiency parameters in microscale high-performance liquid chromatography and capillary electrochromatography 855(1999)349

Efficiency studies in nonaqueous capillary electrophoresis  
859(1999)203

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns  
869(2000)319

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

Hydrophobic-phase-modified fused-silica columns for capillary electrophoresis 871(2000)279

Capillary electrophoresis–inductively coupled plasma mass spectrometry interface with minimised dead volume for high separation efficiency 891(2000)129

Influence of temperature on the behaviour of small linear peptides in capillary electrochromatography 893(2000)401

Life-time studies with capillary electrochromatography columns operated under different conditions 894(2000)35

### Eggs

Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides  
881(2000)1

### Electrically assisted gradient elution

Comparative study of capillary electroosmotic chromatography and electrically assisted gradient nano-liquid chromatography for the separation of peptides 894(2000)241

### Electric field strength

Simulation of electrophoretic separations by the flux-corrected transport method 890(2000)321

### Electrochemical detection

see also Amperometric detection

Application of non-aqueous capillary electrophoresis with electrochemical detection to the determination of nicotine in tobacco 853(1999)27

Nonaqueous capillary electrophoresis equipped with amperometric detection for analysis of chlorinated phenolic compounds 864(1999)323

Simultaneous determination of polycarboxylic acids by capillary electrophoresis with a copper electrode 867(2000)261

Determination of diclofenac sodium by capillary zone electrophoresis with electrochemical detection 868(2000)101

On-column amperometric detection in capillary electrophoresis with an improved high-voltage electric field decoupler  
872(2000)259

Miniaturized thin-layer radial flow cell with interdigitated ring-shaped microarray electrode used as amperometric detector for capillary electrophoresis 891(2000)149

Screening of homocysteine from newborn blood spots by high-performance liquid chromatography with coulometric array detection 896(2000)183

### Electrochromatography

see also Special Issue Index

Preparation of uniformly sized polymeric separation media potentially suitable for small-scale high-performance liquid chromatography and/or capillary electrochromatography  
853(1999)11

Capillary electrochromatography with macroporous particles  
853(1999)45

Chiral separations by open tubular capillary electrokinetic chromatography 853(1999)151

Electrokinetically-driven cation-exchange chromatography of proteins and its comparison with pressure-driven high-performance liquid chromatography 853(1999)243

Capillary electrochromatography of peptides in a microfabricated system 853(1999)257

Separation of selected peptides by capillary electroosmotic chromatography using 3  $\mu\text{m}$  reversed-phase bonded silica and mixed-mode phases 853(1999)263

Separation of carbonyl 2,4-dinitrophenylhydrazones by capillary electrochromatography with diode array detection  
853(1999)487

Capillary electrochromatography with a silica column with a dynamically modified cationic surfactant 855(1999)137

Comparison of capillary electrochromatography with high-performance liquid chromatography for the analysis of pirimicarb and related compounds 855(1999)215

Capillary columns with in situ formed porous monolithic packing for micro high-performance liquid chromatography and capillary electrochromatography 855(1999)273

Dynamics of capillary electrochromatography. II. Comparison of column efficiency parameters in microscale high-performance liquid chromatography and capillary electrochromatography  
855(1999)349

Overview of capillary electrophoresis and capillary electrochromatography 856(1999)443

- Capillary electrochromatography of proteins and peptides with porous-layer open-tubular columns 858(1999)91
- Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319
- Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329
- Some considerations concerning the composition of the mobile phase in capillary electrochromatography 869(2000)363
- Separation of peptides by strong cation-exchange capillary electrochromatography 869(2000)385
- Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395
- Capillary electrochromatography of basic compounds using octadecyl-silica stationary phases with an amine-containing mobile phase 872(2000)315
- Preparation of various silica-based columns for capillary electrochromatography by in-column derivatization 873(2000)247
- Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography 873(2000)293
- Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors 875(2000)235
- Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255
- Separation of enantiomers by packed capillary electrochromatography on a cellulose-based stationary phase 875(2000)331
- Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181
- On-line isotachophoretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219
- Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277
- Design of the monolithic polymers used in capillary electrochromatography columns (Review) 887(2000)3
- Packing columns for capillary electrochromatography (Review) 887(2000)43
- Instrumentation for capillary electrochromatography (Review) 887(2000)55
- Use of on-line mass spectrometric detection in capillary electrochromatography 887(2000)85
- Capillary electrochromatography–mass spectrometry for the separation and identification of isomeric polyaromatic hydrocarbon dna adducts derived from in vitro reactions 887(2000)103
- Gradient elution techniques for capillary electrochromatography 887(2000)115
- Specific applications of capillary electrochromatography to biopolymers, including proteins, nucleic acids, peptide mapping, antibodies, and so forth (Review) 887(2000)137
- Capillary electroosmotic chromatography of peptides (Review) 887(2000)165
- Theory for migration of ions in capillary electrochromatography 887(2000)187
- Modelling of the pore flow in capillary electrochromatography 887(2000)199
- Separation of basic solutes by reversed-phase capillary electrochromatography 887(2000)209
- Separation of acidic compounds by strong anion-exchange capillary electrochromatography 887(2000)223
- Comparison of aqueous and non-aqueous capillary electrochromatography for the separation of basic solutes 887(2000)233
- High temperature and temperature programming in capillary electrochromatography 887(2000)245
- Preparation of stationary phases for open-tubular capillary electrochromatography using the sol–gel method 887(2000)253
- Capillary electrochromatography using continuous-bed columns of sol–gel bonded silica particles with mixed-mode octadecyl and propylsulfonic acid functional groups 887(2000)265
- Characteristics of particle-loaded monolithic sol–gel columns for capillary electrochromatography. I. Structural, electrical and band-broadening properties 887(2000)277
- Capillary electrochromatography with novel stationary phases. IV. Retention behavior of glycosphingolipids on porous and non-porous octadecyl sulfonated silica 887(2000)287
- Stationary phases for capillary electrochromatography (Review) 887(2000)313

- Towards the column bed stabilization of columns in capillary electroosmotic chromatography. Immobilization of microparticulate silica columns to a continuous bed 887(2000)327
- Towards the ultimate minimum particle diameter of silica packings in capillary electrochromatography 887(2000)339
- Aspects of column fabrication for packed capillary electrochromatography 887(2000)347
- Improved column preparation and performance in capillary electrochromatography 887(2000)367
- Steroid profiles determined by capillary electrochromatography, laser-induced fluorescence detection and electrospray–mass spectrometry 887(2000)379
- Separation and identification of etodolac and its urinary phase I metabolites using capillary electrochromatography and on-line capillary electrochromatography–electrospray ionisation mass spectrometry coupling 887(2000)393
- Chiral anion exchangers applied to capillary electrochromatography enantioseparation of oppositely charged chiral analytes: investigation of stationary and mobile phase parameters 887(2000)421
- Capillary electrochromatography of proteins and peptides with a cationic acrylic monolith 887(2000)465
- Analysis of hydroquinone and some of its ethers by using capillary electrochromatography 887(2000)489
- Screening method for determining the presence of N-nitrosodiethanolamine in cosmetics by open-tubular capillary electrochromatography 887(2000)497
- Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267
- Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Peclet number in capillary electrochromatography from pore network theory 890(2000)61
- Pressurized-flow anion-exchange capillary electrochromatography using a polymeric ion-exchange stationary phase 890(2000)337
- Nonaqueous electrochromatography on continuous bed columns of sol–gel bonded large-pore  $C_{18}$  material: separation of retinyl esters 890(2000)347
- Alternative methods providing enhanced sensitivity and selectivity in capillary electroseparation experiments 892(2000)203
- Electrochromatography 892(2000)219
- Advances in column technology and instrumentation in capillary electrochromatography 892(2000)257
- Capillary electrochromatography on silica columns: factors influencing performance 892(2000)279
- Migration of ions in capillary electrochromatography 892(2000)291
- Peak shapes in open tubular ion-exchange capillary electrochromatography of inorganic anions 892(2000)303
- Influence of temperature on the behaviour of small linear peptides in capillary electrochromatography 893(2000)401
- Comparison of separation selectivity in capillary electrokinetic chromatography using a cationic linear polymeric pseudo-stationary phase or monomeric additives of similar structure 894(2000)25
- Life-time studies with capillary electrochromatography columns operated under different conditions 894(2000)35
- Pressurized gradient capillary electrochromatographic separation of eighteen amino acid derivatives 894(2000)337
- Effects of aliphatic amines on capillary electrochromatographic performance of tricyclic antidepressants on octadecylsilica 895(2000)17
- Capillary electrochromatography and capillary electrochromatography–electrospray mass spectrometry for the separation of non-steroidal anti-inflammatory drugs 895(2000)123
- Enantioselective reversed-phase and non-aqueous capillary electrochromatography using a teicoplanin chiral stationary phase 897(2000)349
- Electrodes**  
Determination of diclofenac sodium by capillary zone electrophoresis with electrochemical detection 868(2000)101
- Miniaturized thin-layer radial flow cell with interdigitated ring-shaped microarray electrode used as amperometric detector for capillary electrophoresis 891(2000)149
- Electrodialysis**  
Membrane-based sample preparation coupled on-line to chromatography or electrophoresis 856(1999)55
- Electrokinetic chromatography**  
see also Micellar electrokinetic chromatography and Microemulsion electrokinetic chromatography
- Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudostationary phase. Influence of methanol and acetonitrile on separation selectivity 853(1999)121



- Electrokinetic chromatography 856(1999)465
- Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography 875(2000)179
- New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169
- Potential of ethylenediaminedi(*o*-hydroxyphenylacetic acid) and *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid for the determination of metal ions by capillary electrophoresis 890(2000)363
- Comparison of separation selectivity in capillary electrokinetic chromatography using a cationic linear polymeric pseudo-stationary phase or monomeric additives of similar structure 894(2000)25
- Comparative study of capillary electroendosmotic chromatography and electrically assisted gradient nano-liquid chromatography for the separation of peptides 894(2000)241
- Separation selectivity of anionic metal complexes of *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid in ion and ion electrokinetic chromatography 895(2000)255
- Electrokinetic injection**  
Sub-microliter DNA sequencing for capillary array electrophoresis 894(2000)191
- Electron-capture detection**  
Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535
- Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133
- Electron-capture mass spectrometry**  
Electron-capture mass spectrometry: recent advances 892(2000)329
- Electroosmotic flow**  
Effect of linear polymer additives on the electroosmotic characteristics of agarose gels in ultrathin-layer electrophoresis 853(1999)511
- Capillary columns with in situ formed porous monolithic packing for micro high-performance liquid chromatography and capillary electrochromatography 855(1999)273
- Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669
- Measurement of electroosmotic flow in plastic imprinted microfluid devices and the effect of protein adsorption on flow rate 857(1999)275
- Influence of pH\*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63
- Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329
- Enantiomeric separation by capillary electrophoresis with an electroosmotic flow-controlled capillary 875(2000)315
- Microemulsion electrokinetic chromatography in suppressed electroosmotic flow environment. Separation of fat-soluble vitamins 876(2000)201
- Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181
- Stationary phase structures enhancing electroosmotic flow in capillary electrochromatography 887(2000)299
- Comparison of different packing methods for capillary electrochromatography columns 887(2000)307
- Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Peclet number in capillary electrochromatography from pore network theory 890(2000)61
- Electrochromatography 892(2000)219
- Novel, trifunctional diamine for silica coating in capillary zone electrophoresis 894(2000)53
- Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79
- Effect of ionic strength, pH and polymer concentration on the separation of DNA fragments in the presence of electroosmotic flow 894(2000)219
- Dynamic modification of the capillary wall for electrophoretic separations of small ions 898(2000)133
- Electrophoretically mediated microanalysis**  
Interaction of cyclophilin and cyclosporins monitored by affinity capillary electrophoresis 853(1999)469
- Electrophoretic mobility**  
Electrophoretic mobility study on ion-ion interactions in an aqueous solution 853(1999)63
- Calculation of electrophoretic mobilities in water-organic modifier mixtures in capillary electrophoresis 868(2000)277
- Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile-water media 871(2000)367

Influence of pH and  $pK_a$  values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Calculation of stability constants for the chiral selector–enantiomer interactions from electrophoretic mobilities 875(2000)431

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

New method for standardization of electropherograms obtained in capillary zone electrophoresis 894(2000)11

Mobility and activation energy of single-stranded DNA in denaturing cross-linked polyacrylamide slab gels 894(2000)231

### Electrospray ionization

see also Mass spectrometry

New sheathless interface for coupling capillary electrophoresis to electrospray mass spectrometry evaluated by the analysis of fatty acids and prostaglandins 854(1999)141

Liquid chromatography–electrospray mass spectrometry of  $\beta$ -carotene and xanthophylls. Validation of the analytical method 854(1999)233

Application of electrospray mass spectrometry in the detection and determination of Remazol textile dyes 854(1999)259

Novel microfabricated device for electrokinetically induced pressure flow and electrospray ionization mass spectrometry 892(2000)195

### Electrostatic ion chromatography

Suppressed electrostatic ion chromatography with tetraborate as eluent and its application to the determination of inorganic anions in snow and rainwater 884(2000)161

Elution mechanism in electrostatic ion chromatography with histidine as an isoelectric ampholytic mobile phase 884(2000)287

### Electrostatic repulsion

Different elution modes and field programming in gravitational field-flow fractionation. 2. Experimental verification of the range of conditions for flow-rate and carrier liquid density programming 868(2000)217

### Elemental analysis

Discoloration of ceramic hydroxyapatite used for protein chromatography 891(2000)93

### Elution by characteristic point

Influence of the column radial heterogeneity on the determination of single-component isotherms by the elution by characteristic point method 891(2000)217

### Empirical formula

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

### EN 1785

2-Alkylcyclobutanones as markers for irradiated foodstuffs. III. Improvement of the field of application on the EN 1785 method by using silver ion chromatography 858(1999)109

### Enantiomer bonding

Investigation of mandelic acid bonding on Pirkle type chromatographic stationary phases by Raman spectroscopy 893(2000)37

### Enantiomeric excess

Determination of enantiomeric excess of  $\alpha$ -hydroxy-3-phenoxybenzeneacetonitrile and its *n*-butyl ester by chiral high-performance liquid chromatography 893(2000)189

### Enantiomer separation

see also Chiral stationary phases and Special Issue Index

Alkyl nitrates as achiral and chiral solute probes in gas chromatography. Novel properties of a  $\beta$ -cyclodextrin derivative and characterization of its enantioselective forces 852(1999)525

Chiral separations by open tubular capillary electrokinetic chromatography 853(1999)151

Cyclodextrin-mediated micellar electrokinetic chromatography and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry for the enantiomer separation of racemorphan in human urine 853(1999)439

Resolution of newly synthesized racemic dihydropyridines with different chiral selectors by means of capillary electrophoresis 853(1999)455

Application and comparison of derivatized cellulose and amylose chiral stationary phases for the separation of enantiomers of pharmaceutical compounds by high-performance liquid chromatography 855(1999)411

Rapid development of the enantiomeric separation of  $\beta$ -blockers by capillary electrophoresis using an experimental design approach 855(1999)681

Discrimination between enantioselective and non-selective binding sites on cellobiohydrolase-based stationary phases by site specific competing ligands 857(1999)107

Enantiomeric purity determination of acetyl-L-carnitine by reversed-phase high-performance liquid chromatography using chiral derivatization 857(1999)127

Direct chromatographic resolution of carnitine and *O*-acylcarnitine enantiomers on a teicoplanin-bonded chiral stationary phase 857(1999)145

- Chiral selectors from fruit: application of citrus pectins to enantiomer separations in capillary electrophoresis 857(1999)285
- Quinine versus carbamoylated quinine-based chiral anion exchangers. A comparison regarding enantioselectivity for N-protected amino acids and other chiral acids 858(1999)1
- Temperature-induced inversion of elution order in the enantioseparation of sotalol on a cellobiohydrolase I-based stationary phase 858(1999)33
- Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VII. Comparison of chiral recognition ability of ovoglycoprotein from chicken and Japanese quail egg whites 858(1999)155
- Chiral separations using polymeric dipeptide surfactants: effect of number of chiral centers and steric factors 858(1999)219
- High-performance liquid chromatography study of the enantiomer separation of chrysanthemic acid and its analogous compounds on a terguride-based stationary phase 859(1999)133
- Convenient synthesis of  $\pi$ -acceptor chiral stationary phases for high-performance liquid chromatography from halogen-substituted 3,5-dinitrobenzoylamides 859(1999)143
- Chromatographic evaluation of structure selective and enantioselective retention of amines and acids on cellobiohydrolase I wild type and its mutant D214N 864(1999)1
- Separation of enantiomers on anion exchangers modified with heparin in liquid chromatography 864(1999)25
- Chiral resolution of basic drugs by capillary electrophoresis with new glycosaminoglycans 864(1999)163
- Chiral high-performance liquid chromatography of *N*-octyl bicycloheptene dicarboximide and confirmatory studies using liquid chromatography–tandem mass spectrometry and two-dimensional nuclear magnetic resonance spectroscopy 864(1999)271
- Development of an high-performance liquid chromatographic simulated moving bed separation from an industrial perspective 865(1999)35
- Enantiomers separation by simulated moving bed chromatography. Non-instantaneous equilibrium at the solid–fluid interface 865(1999)187
- Enantiomer separation of  $\alpha$ -ionone using gas chromatography with cyclodextrin derivatives as chiral stationary phases 865(1999)201
- Preparative chromatographic resolution of enantiomers using polar organic solvents with polysaccharide chiral stationary phases 865(1999)211
- Direct, preparative enantioselective chromatography of propranolol hydrochloride and thioridazine hydrochloride using carbon dioxide-based mobile phases 865(1999)227
- Enantiomeric separation of amino alcohols on protein phases using statistical experimental design. A comparative study 866(2000)15
- Synthesis of a silica-bonded bovine serum albumin *s*-triazine chiral stationary phase for high-performance liquid chromatographic resolution of enantiomers 866(2000)173
- Comparison of the enantioseparation of racemic uridine analogs on Whelk-O 1 and ChiralPak-AD columns 866(2000)183
- Enantiomer separation of polychlorinated biphenyl atropisomers and polychlorinated biphenyl retention behavior on modified cyclodextrin capillary gas chromatography columns 866(2000)213
- Gas chromatographic quantification of amino acid enantiomers in food matrices by their *N*(*O,S*)-ethoxycarbonyl heptafluorobutyl ester derivatives 866(2000)221
- Micellar electrokinetic chromatography for the analysis of *D*-amygdalin and its epimer in apricot kernel 866(2000)253
- Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1
- Liquid chromatographic resolution of 2-hydroxycarboxylic acids on a new chiral stationary phase derived from (*S*)-leucine 868(2000)31
- Chiral and electrokinetic separation of amino acids using polypyrrole-coated adsorbents 868(2000)189
- Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic  $\beta$ -substituted  $\alpha$ -amino acids on a teicoplanin chiral stationary phase 868(2000)197
- Enantioseparation of atropine by capillary electrophoresis using sulfated  $\beta$ -cyclodextrin: application to a plant extract 868(2000)285
- Simultaneous stereoselective analysis of tramadol and its main phase I metabolites by on-line capillary zone electrophoresis–electrospray ionization mass spectrometry 868(2000)295
- Novel cinchona alkaloid carbamate  $C_9$ -dimers as chiral anion-exchange type selectors for high-performance liquid chromatography 869(2000)111
- High-performance liquid chromatographic separation of the enantiomers of unusual  $\alpha$ -amino acid analogues 871(2000)105

- Application of (1*S*,2*S*)- and (1*R*,2*R*)-1,3-diacetoxy-1-(4-nitrophenyl)-2-propylisothiocyanate to the indirect enantioseparation of racemic proteinogenic amino acids 871(2000)115
- Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127
- Application of the restricted-access precolumn packing material alkyl-diol silica in a column-switching system for the determination of ketoprofen enantiomers in horse plasma 871(2000)153
- Enantioselective analysis of methadone in saliva by liquid chromatography–mass spectrometry 871(2000)163
- Enantiomeric separation of clenbuterol by transient isotachopheresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachopheresis 871(2000)173
- Stereospecific analysis of lorazepam in plasma by chiral column chromatography with a circular dichroism-based detector 871(2000)181
- Enantiospecific determination of PNU-83894 and its major metabolite, PNU-83892, in plasma, and its application to the characterization of the enantioselective pharmacokinetics of PNU-83894 in the dog 871(2000)201
- Analysis of *cis*–*trans* isomers and enantiomers of sertraline by cyclodextrin-modified micellar electrokinetic chromatography 871(2000)207
- Enantiomeric separation of enzymatic hydrolysis products of dihydropyrimidinone methyl ester with cationic cyclodextrin by capillary electrophoresis 872(2000)279
- Chromatographic resolution of tryptophan enantiomers with L-Leu–L-Leu–L-Leu peptide. Effects of mobile phase composition and chromatographic support 874(2000)21
- Comparative study on camphor enantiomers behavior under the conditions of gas–liquid chromatography and reversed-phase high-performance liquid chromatography systems modified with  $\alpha$ - and  $\beta$ -cyclodextrins 874(2000)73
- Enantiomeric separation and discrimination of 2-hydroxy acids as *O*-trifluoroacetylated (*S*)-(+)-3-methyl-2-butyl esters by achiral dual-capillary column gas chromatography 874(2000)91
- Chiral separation of unmodified amino acids with non-aqueous capillary electrophoresis based on the ligand-exchange principle 874(2000)285
- Enantiomer separation of drugs by capillary electromigration techniques 875(2000)3
- Enantioselective determination of drugs in body fluids by capillary electrophoresis 875(2000)27
- Chiral separation of amino acids and peptides by capillary electrophoresis 875(2000)43
- Enantioselective determination by capillary electrophoresis with cyclodextrins as chiral selectors 875(2000)89
- Enantiomeric separations of drugs using mixtures of charged and neutral cyclodextrins 875(2000)123
- Separation of drug enantiomers by capillary electrophoresis in the presence of neutral cyclodextrins 875(2000)135
- Enantiomer separation of drugs by micellar electrokinetic chromatography using chiral surfactants 875(2000)163
- Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography 875(2000)179
- Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207
- Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors 875(2000)235
- Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255
- Enantiomeric separations by nonaqueous capillary electrophoresis 875(2000)277
- New amphiphilic aminosaccharide derivatives as chiral selectors in capillary electrophoresis 875(2000)295
- Application of ligand-exchange capillary electrophoresis to the chiral separation of  $\alpha$ -hydroxy acids and  $\beta$ -blockers 875(2000)307
- Enantiomeric separation by capillary electrophoresis with an electroosmotic flow-controlled capillary 875(2000)315
- Separation of enantiomers by capillary electrophoresis–mass spectrometry employing a partial filling technique with a chiral crown ether 875(2000)323
- Separation of enantiomers by packed capillary electrochromatography on a cellulose-based stationary phase 875(2000)331
- Investigation of an enantioselective non-aqueous capillary electrochromatography system applied to the separation of chiral acids 875(2000)341
- Non-aqueous capillary electrophoretic enantioseparation of *N*-derivatized amino acids using cinchona alkaloids and derivatives as chiral counter-ions 875(2000)353

- Direct chiral resolution of lactic acid in food products by capillary electrophoresis 875(2000)371
- Enantioseparation of 3,4-dihydroxyphenylalanine and 2-hydrazino-2-methyl-3-(3,4-dihydroxyphenyl)propanoic acid by capillary electrophoresis using cyclodextrins 875(2000)379
- Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated  $\beta$ -cyclodextrin 875(2000)389
- Fast chiral separations using sulfated  $\beta$ -cyclodextrin and short-end injection in capillary electrophoresis 875(2000)403
- Design of experiments for capillary electrophoretic enantioresolution of salbutamol using dermatan sulfate 875(2000)411
- Optimization of the cyclodextrin-assisted capillary electrophoresis separation of the enantiomers of phenoxyacid herbicides 875(2000)423
- Calculation of stability constants for the chiral selector–enantiomer interactions from electrophoretic mobilities 875(2000)431
- Enantioselective binding analysis of verapamil to plasma lipoproteins by capillary electrophoresis–frontal analysis 875(2000)447
- Mechanistic study of opposite migration order of dimethindene enantiomers in capillary electrophoresis in the presence of native  $\beta$ -cyclodextrin and heptakis(2,3,6-tri-*O*-methyl)- $\beta$ -cyclodextrin 875(2000)455
- Separation of brompheniramine enantiomers by capillary electrophoresis and study of chiral recognition mechanisms of cyclodextrins using NMR-spectroscopy, UV spectrometry, electrospray ionization mass spectrometry and X-ray crystallography 875(2000)471
- Stereoselective binding of 2,3-substituted 3-hydroxypropionic acids on an immobilised human serum albumin chiral stationary phase: stereochemical characterisation and quantitative structure–retention relationship study 876(2000)75
- Simultaneous separation and enantioseparation of thalidomide and its hydroxylated metabolites using high-performance liquid chromatography in common-size columns, capillary liquid chromatography and nonaqueous capillary electrochromatography 876(2000)157
- Liquid chromatographic separation of the stereoisomers of thiazide diuretics 876(2000)221
- Study of the adsorption behavior of the enantiomers of 1-phenyl-1-propanol on a cellulose-based chiral stationary phase 877(2000)95
- Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109
- Chiral separation of homocysteine by derivatization with 4-aminosulfonyl-7-fluoro-2,1,3-benzoxadiazole followed by capillary electrophoresis using  $\gamma$ -cyclodextrin 877(2000)217
- Case of enantiomer impurity identification by normal-phase chiral high-performance liquid chromatography with optical rotation and mass spectrometric detection 878(2000)35
- Large degree of racemization observed in the amide bond forming reaction on silica gel 878(2000)165
- Direct liquid chromatographic enantiomer separation of new fluoroquinolones including gemifloxacin 879(2000)113
- Enantioseparation of selected *N-tert*-butyloxycarbonyl amino acids in high-performance liquid chromatography and capillary electrophoresis with a teicoplanin chiral selector 879(2000)147
- Chromatographic determination of amino acid enantiomers in beers and raw materials used for their manufacture 881(2000)81
- Insoluble  $\beta$ -cyclodextrin polymer for capillary gas chromatographic separation of enantiomers and isomers 883(2000)137
- Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs 883(2000)249
- Chiral anion exchangers applied to capillary electrochromatography enantioseparation of oppositely charged chiral analytes: investigation of stationary and mobile phase parameters 887(2000)421
- Enantioseparations in non-aqueous capillary electrochromatography using polysaccharide type chiral stationary phases 887(2000)439
- Enantiomer separations by capillary electrochromatography using chiral stationary phases 887(2000)457
- Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63
- On-line monitoring of enantiomer concentration in chiral simulated moving bed chromatography 888(2000)73
- Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241
- Determination of the enantiomers of 3-*tert*-butylamino-1,2-propanediol by high-performance liquid chromatography coupled to evaporative light scattering detection 890(2000)239

- Determination of enantiomers in a synthetic argininal peptide using capillary zone electrophoresis and high-performance liquid chromatography 891(2000)115
- Configurational analysis of chiral acids as *O*-trifluoroacetylated (–)-menthyl esters by achiral dual-capillary column gas chromatography 891(2000)257
- Non-aqueous capillary electrophoresis 892(2000)155
- Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487
- Enantiomer separations by nonaqueous capillary electrophoresis using octakis(2,3-diacetyl-6-sulfato)- $\gamma$ -cyclodextrin 892(2000)499
- tert*-Butylcarbamoylquinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique 892(2000)509
- Separation of the four pairs of enantiomers of vincamine alkaloids by enantioselective high-performance liquid chromatography 893(2000)47
- Enantioseparation of anionic analytes by non-aqueous capillary electrophoresis using quinine and quinidine derivatives as chiral counter-ions 894(2000)63
- Allylamine- $\beta$ -cyclodextrin copolymer. A novel chiral selector for capillary electrophoresis 894(2000)95
- Chiral determination of amino acids by capillary electrophoresis and laser-induced fluorescence at picomolar concentrations 894(2000)259
- Influence of the structure of cyclodextrins and amino acid sequence of dipeptides and tripeptides on the pH-dependent reversal of the migration order in capillary electrophoresis 894(2000)267
- Chiral capillary electrophoresis as predictor for separation of drug enantiomers in continuous flow zone electrophoresis 895(2000)51
- Capillary electrophoretic separation of the enantiomers of organophosphates with a phosphorus stereogenic center using the sodium salt of octakis(2,3-diacetyl-6-sulfo)- $\gamma$ -cyclodextrin as resolving agent 895(2000)247
- Determination of optical purity of phosphonic acid analogues of aromatic amino acids by capillary electrophoresis with  $\alpha$ -cyclodextrin 895(2000)301
- Determination of the enantiomers of 3-*tert*-butylamino-1,2-propanediol by high-performance liquid chromatography using mass spectrometric detection 896(2000)201
- Use of chiral liquid chromatography–tandem mass spectrometry to investigate the metabolism of racemic cholecystokinin-B antagonists 896(2000)217
- Determination of the in vitro metabolism of (+)- and (–)-epibatidine 896(2000)229
- Analysis of the enzymatic racemization of *D*-aspartic acid to *L*-aspartic acid by the on-line coupling of a solid-phase extraction column and a ligand-exchange high-performance liquid chromatography column 896(2000)299
- Evaluation of the macrocyclic glycopeptide A-40,926 as a high-performance liquid chromatographic chiral selector and comparison with teicoplanin chiral stationary phase 897(2000)113
- Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic  $\beta$ -substituted  $\alpha$ -amino acids on a copper(II)-*D*-penicillamine chiral stationary phase 897(2000)145
- Enantioselective reversed-phase and non-aqueous capillary electrochromatography using a teicoplanin chiral stationary phase 897(2000)349
- Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate  $\beta$ -cyclodextrin bonded chiral stationary phase 898(2000)53
- Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations 898(2000)63
- End-capping**  
Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375
- Enhanced solvent extraction**  
Comparison of methods for quantitative analysis of additives in low-density polyethylene using supercritical fluid and enhanced solvent extraction 858(1999)209
- Entangled polymers**  
Capillary electrophoretic behaviour of humic substances in physical gels 863(1999)195
- Environmental analysis**  
see also Special Issue Index
- Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465
- Enhancement of electron-capture detection of methyl bromide in air by iodination 852(1999)535
- Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

- Quantification of anions and cations in environmental water samples. Measurements with capillary electrophoresis and indirect-UV detection 853(1999)403
- Separation of carbonyl 2,4-dinitrophenylhydrazones by capillary electrochromatography with diode array detection 853(1999)487
- Detection and identification of sulphonamide drugs in municipal waste water by liquid chromatography coupled with electrospray ionisation tandem mass spectrometry 854(1999)163
- Use of volatile amines as ion-pairing agents for the high-performance liquid chromatographic–tandem mass spectrometric determination of aromatic sulfonates in industrial wastewater 854(1999)175
- On-line solid-phase extraction–ion-pair liquid chromatography–electrospray mass spectrometry for the trace determination of naphthalene monosulphonates in water 854(1999)187
- Simultaneous determination of antifouling herbicides in marina water samples by on-line solid-phase extraction followed by liquid chromatography–mass spectrometry 854(1999)197
- Investigations on the metabolism of alkyl polyglucosides and their determination in waste water by means of liquid chromatography–electrospray mass spectrometry 854(1999)221
- On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279
- Comprehensive two-dimensional gas chromatography: a hyphenated method with strong coupling between the two dimensions 856(1999)331
- Ultra-trace-level determination of polar pesticides and their transformation products in surface and estuarine water samples using column liquid chromatography–electrospray tandem mass spectrometry 857(1999)157
- Analysis of methyl *tert.*-butyl ether and its degradation products by direct aqueous injection onto gas chromatography with mass spectrometry or flame ionization detection systems 857(1999)205
- Potential for chlorate interference in ion chromatographic determination of total nitrogen in natural waters following alkaline persulfate digestion 857(1999)337
- Analysis of linear alkylbenzenesulfonates in water samples by large-volume injection-port derivatization and gas chromatography–mass spectrometry 857(1999)359
- Semi-permeable surface analytical reversed-phase column for the improved trace analysis of acidic pesticides in water with coupled-column reversed-phase liquid chromatography with UV detection. Determination of bromoxynil and bentazone in surface water 858(1999)45
- Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29
- Gas chromatographic–mass spectrometric determination of sulfolane in wetland vegetation exposed to sour gas-contaminated groundwater 859(1999)69
- Analysis of some pesticides in water samples using solid-phase microextraction–gas chromatography with different mass spectrometric techniques 859(1999)193
- Sample cleanup and reversed-phase high-performance liquid chromatographic analysis of polar aromatic compounds in groundwater samples from a former gas plant 862(1999)137
- Analysis of high-molecular-mass polycyclic aromatic hydrocarbons in environmental samples using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 863(1999)13
- Automated gas chromatography with cryogenic/sorbent trap for the measurement of volatile organic compounds in the atmosphere 863(1999)183
- Simultaneous measurement of volatile sulfur compounds using ascorbic acid for oxidant removal and gas chromatography–flame photometric detection 864(1999)111
- Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121
- Soil thin-layer chromatography and pesticide mobility through soil microstructures. New technical approach 864(1999)145
- New gas chromatographic method for residue determination of dithiopyr in soil, wheat grain and straw 864(1999)351
- Application of pressurized liquid extraction followed by gas chromatography–mass spectrometry to determine 4-nonylphenols in sediments 866(2000)79
- Determination of pesticides in waters by automatic on-line solid-phase extraction–capillary electrophoresis 866(2000)137
- On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207
- Pesticide trace analysis using solid-phase extraction and gas chromatography with electron-capture and tandem mass spectrometric detection in water samples 867(2000)235
- 2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography–mass spectrometry 869(2000)251

- Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411
- Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451
- Evaluation of surface- and ground-water pollution due to herbicides in agricultural areas of Zamora and Salamanca (Spain) 869(2000)471
- Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487
- Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523
- Trace-level determination of triazines and several degradation products in environmental waters by disk solid-phase extraction and micellar electrokinetic chromatography 872(2000)299
- Analysis of steroids in environmental water samples using solid-phase extraction and ion-trap gas chromatography–mass spectrometry and gas chromatography–tandem mass spectrometry 872(2000)309
- Solid-phase microextraction–gas chromatography–mass spectrometry of biologically active substances in water samples 873(2000)95
- Determination of triazines in soil leachates by solid-phase microextraction coupled to gas chromatography–mass spectrometry 874(2000)247
- Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl)ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane] from aged contaminated soil 874(2000)257
- Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153
- Characterization of a commercial gas chromatography–flame ionization detection system for the in situ determination of C<sub>5</sub>–C<sub>10</sub> hydrocarbons in ambient air 878(2000)215
- Determination of rice herbicides, their transformation products and clofibric acid using on-line solid-phase extraction followed by liquid chromatography with diode array and atmospheric pressure chemical ionization mass spectrometric detection 879(2000)3
- Monitoring of priority pesticides and other organic pollutants in river water from Portugal by gas chromatography–mass spectrometry and liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)13
- Part-per-trillion level determination of antifouling pesticides and their byproducts in seawater samples by off-line solid-phase extraction followed by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)27
- Comparison of various liquid chromatographic methods involving UV and atmospheric pressure chemical ionization mass spectrometric detection for the efficient trace analysis of phenylurea herbicides in various types of water samples 879(2000)39
- Application of liquid chromatography with mass spectrometry combined with photodiode array detection and tandem mass spectrometry for monitoring pesticides in surface waters 879(2000)51
- Stability of organophosphorus insecticides on graphitized carbon black extraction cartridges used for large volumes of surface water 879(2000)73
- Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83
- Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97
- Speciation of butyl- and phenyltin compounds in sediments using pressurized liquid extraction and liquid chromatography–inductively coupled plasma mass spectrometry 879(2000)137
- Solid-phase microextraction for determining the distribution of sixteen US Environmental Protection Agency polycyclic aromatic hydrocarbons in water samples 879(2000)177
- Advances in the determination of inorganic anions by ion chromatography 881(2000)607
- On-line microdialysis–ion chromatographic determination of inorganic anions in olive-oil mill wastewater 882(2000)321
- Pressurised liquid extraction of polycyclic aromatic hydrocarbons from contaminated soils 883(2000)151
- Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113



- Simultaneous ion-exclusion/cation-exchange chromatography of anions and cations in acid rain waters on a weakly acidic cation-exchange resin by elution with sulfosalicylic acid 884(2000)167
- Determination of inorganic anions in environmental waters with a hydroxide-selective column 884(2000)175
- Determination of chloride, sulfate and nitrate in groundwater samples by ion chromatography 884(2000)185
- Comparison of on-line coupling of ion-chromatography with atmospheric pressure ionization mass spectrometry and with inductively coupled plasma mass spectrometry as tools for the ultra-trace analysis of bromate in surface water samples 884(2000)191
- Determination of inorganic and organic anionic arsenic species in water by ion chromatography coupled to hydride generation–inductively coupled plasma atomic emission spectrometry 884(2000)211
- Trace analysis of heavy metals in groundwater samples by ion chromatography with post-column reaction and ultraviolet–visible detection 884(2000)229
- Fifty years of solid-phase extraction in water analysis – historical development and overview 885(2000)3
- Solid-phase extraction for multiresidue analysis of organic contaminants in water 885(2000)195
- Multiresidue methods using solid-phase extraction techniques for monitoring priority pesticides, including triazines and degradation products, in ground and surface waters 885(2000)217
- Solid-phase extraction of quaternary ammonium herbicides 885(2000)251
- On-line dual-precursor-based trace enrichment for the determination of polar and acidic microcontaminants in river water by liquid chromatography with diode-array UV and tandem mass spectrometric detection 885(2000)377
- Application of solid-phase microextraction for determination of organic vapours in gaseous matrices 885(2000)405
- Validation of an analytical procedure for polychlorinated biphenyls, coplanar polychlorinated biphenyls and polycyclic aromatic hydrocarbons in environmental samples 886(2000)153
- New methods for determination of glyphosate and (aminomethyl)phosphonic acid in water and soil 886(2000)207
- Static subcritical water extraction with simultaneous solid-phase extraction for determining polycyclic aromatic hydrocarbons on environmental solids 886(2000)237
- Identification of reactive dyes in spent dyebaths and wastewater by capillary electrophoresis–mass spectrometry 886(2000)271
- Solid-phase microextraction: a promising technique for sample preparation in environmental analysis 889(2000)3
- Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99
- Fractionation and toxicity evaluation of waste waters 889(2000)149
- Inter-laboratory comparison of liquid chromatographic techniques and enzyme-linked immunosorbent assay for the determination of surfactants in wastewaters 889(2000)195
- Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211
- Determination of linear alkylbenzenesulfonates in wastewater treatment plants and coastal waters by automated solid-phase extraction followed by capillary electrophoresis–UV detection and confirmation by capillary electrophoresis–mass spectrometry 889(2000)221
- Unknown bisethylisooctanolactone isomers in industrial waste water. Isolation, identification and occurrence in surface water 889(2000)245
- Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261
- Importance of handling organic atmospheric pollutants for assessing air quality 889(2000)271
- Indirect analysis of urea herbicides from environmental water using solid-phase microextraction 890(2000)303
- Analysis of agrochemicals by capillary electrophoresis (Review) 891(2000)45
- Comparison of two extraction methods independently developed on two conceptually different automated supercritical fluid extraction systems for the determination of polychlorinated biphenyls in sediments 891(2000)195
- Application of gas chromatography coupled to chemical ionisation mass spectrometry following headspace solid-phase microextraction for the determination of free volatile fatty acids in aqueous samples 891(2000)287
- Monitoring organochlorine pesticides from landfill leachates by gas chromatography–electron-capture detection after solid-phase microextraction 891(2000)305

- Simultaneous determination of ethylbenzene, indan, indene and acenaphthene in air by capillary gas chromatography 891(2000)361
- Determination of steroid sex hormones and related synthetic compounds considered as endocrine disrupters in water by liquid chromatography–diode array detection–mass spectrometry 892(2000)391
- Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421
- Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435
- Determination of fungicides in natural waters using solid-phase microextraction and gas chromatography coupled with electron-capture and mass spectrometric detection 893(2000)143
- Enantiomeric separations of cationic and neutral compounds by capillary electrochromatography with monolithic chiral stationary phases of  $\beta$ -cyclodextrin-bonded negatively charged polyacrylamide gels 893(2000)177
- Pressurised liquid extraction of ketones of polycyclic aromatic hydrocarbons from soil 893(2000)201
- High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253
- High-performance liquid chromatographic resolution of reverse isomers of 1,2-diacyl-*rac*-glycerols as 3,5-dinitrophenylurethanes 893(2000)261
- Enantioseparation of 1-phenyl-1-propanol by simulated moving bed under linear and nonlinear conditions 893(2000)307
- Solid-phase microextraction for the determination of systemic and non-volatile pesticides in river water using gas chromatography with nitrogen–phosphorous and electron-capture detection 893(2000)347
- Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the  $\text{ng l}^{-1}$  concentration level 893(2000)367
- Determination of aromatic amines in water samples by capillary electrophoresis with electrochemical and fluorescence detection 895(2000)197
- Simultaneous determination of alkali, alkaline earth and transition metal ions by capillary electrophoresis with indirect UV detection 895(2000)205
- Separation of two groups of oestrogen mimicking compounds using micellar electrokinetic chromatography 895(2000)237
- Cryogen free automated gas chromatography for the measurement of ambient volatile organic compounds 896(2000)31
- On-line coupling of solid-phase extraction to gas chromatography with fast solvent vaporization and concentration in an open injector liner. Analysis of pesticides in aqueous samples 896(2000)41
- Determination of aqueous chlorothalonil with solid-phase microextraction and gas chromatography 896(2000)105
- Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111
- Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149
- Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279
- Enzyme amplification**  
Enzyme amplification as detection tool in continuous-flow systems. I. Development of an enzyme-amplified biochemical detection system coupled on-line to flow-injection analysis 855(1999)383
- Enzyme amplification as detection tool in continuous-flow systems. II. On-line coupling of liquid chromatography to enzyme-amplified biochemical detection after pre-column derivatization with biotin 855(1999)397
- High-performance liquid chromatography coupled to enzyme-amplified biochemical detection for the analysis of hemoglobin after pre-column biotinylation 886(2000)19
- Ephedrae herba***  
Analysis of tetramethylpyrazine in *Ephedrae herba* by gas chromatography–mass spectrometry and high-performance liquid chromatography 878(2000)147
- Equilibrium-dispersive model**  
Calculation of chromatographic band profiles with an implicit isotherm 862(1999)1
- Equilibrium sorptive enrichment**  
On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207
- Equilibrium theory**  
Design and optimisation of a simulated moving bed unit: role of deviations from equilibrium theory 876(2000)3
- Escherichia coli***  
High-yield purification of biosynthetic human growth hormone secreted in *Escherichia coli* periplasmic space 852(1999)441

Renaturation of heterodimeric platelet-derived growth factor from inclusion bodies of recombinant *Escherichia coli* using size-exclusion chromatography 855(1999)203

Immobilised metal affinity chromatography of  $\beta$ -galactosidase from unclarified *Escherichia coli* homogenates using expanded bed adsorption 874(2000)27

#### Essential oils

see also Oils

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

#### Etchants

Ion-chromatographic assay of chromium(VI)-containing semiconductor etchants. Statistical comparison of two columns 884(2000)151

#### Etched capillaries

Open tubular capillary electrokinetic chromatography in etched fused-silica tubes (Review) 887(2000)31

#### *Eucalyptus globulus*

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

#### *Euphorbia leuconeura*

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

#### Evaporative light scattering detection

Hindered amine stabilizers investigated by the use of packed capillary temperature-programmed liquid chromatography. I. Poly(6-(1,1,3,3-tetramethylbutyl)-amino)-1,3,5-triazine-2,4-diyl(2,2,6,6-tetramethyl-4-piperidyl)imino)-1,6-hexanedyl(2,2,6,6-tetramethyl-4-piperidyl)imino) 874(2000)65

Determination of the enantiomers of 3-*tert*-butylamino-1,2-propanediol by high-performance liquid chromatography coupled to evaporative light scattering detection 890(2000)239

Characterization of sulfobutyl ether- $\beta$ -cyclodextrins mixtures by anion-exchange chromatography using evaporative light scattering detection 897(2000)185

#### *Evernia prunastri*

Separation of polyamines, conjugated to DNA, by reversed-phase high-performance liquid chromatography 870(2000)425

#### Expanded bed chromatography

Expanded-bed chromatography in primary protein purification (Review) 865(1999)129

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

The influence of cell adsorbent interactions on protein adsorption in expanded beds 873(2000)195

Immobilised metal affinity chromatography of  $\beta$ -galactosidase from unclarified *Escherichia coli* homogenates using expanded bed adsorption 874(2000)27

Pellicular expanded bed matrix suitable for high flow rates 878(2000)17

#### Experimental design

Application of experimental design for the characterisation of a novel elution system for high-capacity anion chromatography with suppressed conductivity detection 855(1999)91

Rapid development of the enantiomeric separation of  $\beta$ -blockers by capillary electrophoresis using an experimental design approach 855(1999)681

Immobilisation and evaluation of a vancomycin chiral stationary phase for capillary electrochromatography 869(2000)395

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215

Design of experiments for capillary electrophoretic enantioresolution of salbutamol using dermatan sulfate 875(2000)411

Spectrophotometric and coulometric detection in the high-performance liquid chromatography of flavonoids and optimization of sample treatment for the determination of quercetin in orange juice 881(2000)449

Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs 883(2000)249

Experimental design of a microwave-assisted extraction–derivatization method for the analysis of methylmercury 889(2000)185

Capillary electrophoresis analysis of gentamicin sulphate with UV detection after pre-capillary derivatization with 1,2-phthalic dicarboxaldehyde and mercaptoacetic acid 895(2000)67

Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149

Optimisation of alachlor solid-phase microextraction from water samples using experimental design 896(2000)373

Enantioselective reversed-phase and non-aqueous capillary electrochromatography using a teicoplanin chiral stationary phase 897(2000)349

#### Extraction methods

see also Clean-up methods, Preconcentration, Pressurized liquid extraction, Sample handling, Sample preparation, Solid-phase extraction, Solid-phase microextraction, Supercritical fluid extraction and Special Issue Index

Evaluation of a dual-sorbent trap for monitoring organic compounds in air 855(1999)593

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

Solid-phase extraction: method development, sorbents, and coupling with liquid chromatography 856(1999)3

Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217

Study of applicability of various solid-phase extraction materials for sample handling in screening analysis of organic micropollutants in water 857(1999)247

Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29

Gas chromatographic study of complexing in the system hydrazoic acid–tributyl phosphate–nitric acid–uranyl nitrate 859(1999)41

Supercritical fluid extraction of chlorpyrifos and 3,5,6-trichloro-2-pyridinol from garden compost 862(1999)105

Strategies for supercritical fluid extraction of hyoscyamine and scopolamine salts using basified modifiers 863(1999)47

Trapping efficiency of aqueous pollutants in multichannel thick-film silicone-rubber traps for capillary gas chromatography 863(1999)57

Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121

Use of solid-phase microextraction for measuring oil–water partition coefficients and correlation with high-performance liquid chromatographic methods for lipophilicity 864(1999)183

Application of pressurized liquid extraction followed by gas chromatography–mass spectrometry to determine 4-nonylphenols in sediments 866(2000)79

Selective extraction of hydrocarbons, phosphonates and phosphonic acids from soils by successive supercritical fluid and pressurized liquid extractions 866(2000)231

Determination of phenols using simultaneous steam distillation–extraction 867(2000)281

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Microwave-assisted extraction by fast sample preparation for the systematic analysis of additives in polyolefins by high-performance liquid chromatography 869(2000)285

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515

Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151

Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

- Determination of sulphur compounds in beer using headspace solid-phase microextraction and gas chromatographic analysis with pulsed flame photometric detection 872(2000)203
- Development of a simple in-vial liquid-phase microextraction device for drug analysis compatible with capillary gas chromatography, capillary electrophoresis and high-performance liquid chromatography 873(2000)3
- Development of membrane extraction with a sorbent interface–micro gas chromatography system for field analysis 873(2000)13
- Analysis of volatile contaminants in vegetable oils by headspace solid-phase microextraction with Carboxen-based fibres 873(2000)79
- Solid-phase microextraction–gas chromatography–mass spectrometry of biologically active substances in water samples 873(2000)95
- Development of a headspace solid-phase microextraction procedure for the determination of free volatile fatty acids in waste waters 873(2000)107
- Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117
- Automated in-tube solid-phase microextraction–high-performance liquid chromatography for carbamate pesticide analysis 873(2000)137
- On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175
- Immobilised metal affinity chromatography of  $\beta$ -galactosidase from unclarified *Escherichia coli* homogenates using expanded bed adsorption 874(2000)27
- Trace analysis of ten chlorinated benzenes in water by headspace solid-phase microextraction 874(2000)149
- Extraction of hydrocarbon contamination from soils using accelerated solvent extraction 874(2000)217
- Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl)ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane] from aged contaminated soil 874(2000)257
- Alkyltrimethylammonium surfactant-mediated extractions: Characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87
- Dynamic microwave-assisted extraction 877(2000)141
- Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153
- Simultaneous quantitative trace analysis of anionic and nonionic surfactant mixtures by reversed-phase liquid chromatography 878(2000)99
- Stability of organophosphorus insecticides on graphitized carbon black extraction cartridges used for large volumes of surface water 879(2000)73
- Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121
- Applications of matrix solid-phase dispersion in food analysis 880(2000)63
- $\beta_2$ -Agonist extraction procedures for chromatographic analysis 880(2000)69
- Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxy-pyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93
- Determination of fenpyroximate in apples by supercritical fluid extraction and packed capillary liquid chromatography with UV detection 880(2000)121
- Automated analysis of vitamin E isomers in vegetable oils by continuous membrane extraction and liquid chromatography–electrochemical detection 881(2000)229
- Gas chromatographic determination of acrinathrine and 3-phenoxybenzaldehyde residues in honey 882(2000)239
- Multiresidue methods using solid-phase extraction techniques for monitoring priority pesticides, including triazines and degradation products, in ground and surface waters 885(2000)217
- Solid-phase extraction of quaternary ammonium herbicides 885(2000)251
- Role of solid-phase microextraction in the identification of highly volatile pheromones of two Rhinoceros beetles *Scapanes australis* and *Strategus aloeus* (Coleoptera, Scarabaeidae, Dynastinae) 885(2000)433
- Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annum* L.) powders 889(2000)41
- Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69
- Evaluation of extraction procedures for the ion chromatographic determination of arsenic species in plant materials 889(2000)83

- Extraction of glyphosate by a supported liquid membrane technique 889(2000)93
- Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99
- Optimisation of procedures for the extraction of structural analogues of propranolol with molecular imprinted polymers for sample preparation 889(2000)143
- Experimental design of a microwave-assisted extraction–derivatization method for the analysis of methylmercury 889(2000)185
- Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211
- Stability of sulfonated derivatives of benzene and naphthalene on disposable solid-phase extraction pre-columns and in an aqueous matrix 889(2000)231
- Solid-phase extraction of sugar cane soot extract for analysis by gas chromatography with flame ionisation and mass spectrometric detection 889(2000)281
- Comparison of two extraction methods independently developed on two conceptually different automated supercritical fluid extraction systems for the determination of polychlorinated biphenyls in sediments 891(2000)195
- Miniaturised pressurised liquid extraction of polycyclic aromatic hydrocarbons from soil and sediment with subsequent large-volume injection–gas chromatography 891(2000)275
- Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421
- Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435
- High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157
- Pressurised liquid extraction of ketones of polycyclic aromatic hydrocarbons from soil 893(2000)201
- Integration of a microextraction system. Solvent extraction of a Co–2-nitroso-5-dimethylaminophenol complex on a microchip 894(2000)19
- Simultaneous supercritical fluid derivatization and extraction of formaldehyde by the Hantzsch reaction 896(2000)51
- Possible interest of various sample transfer techniques for fast gas chromatography–mass spectrometric analysis of true onion volatiles 896(2000)117
- Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291
- Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269
- Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279

## F

### Factorial designs

- Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505
- Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45
- Robustness testing of a reversed-phase high-performance liquid chromatographic assay: comparison of fractional and asymmetrical factorial designs 874(2000)167
- Dynamic microwave-assisted extraction 877(2000)141
- Simultaneous quantitative trace analysis of anionic and nonionic surfactant mixtures by reversed-phase liquid chromatography 878(2000)99
- Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs 883(2000)249
- Experimental design of a microwave-assisted extraction–derivatization method for the analysis of methylmercury 889(2000)185
- Simultaneous supercritical fluid derivatization and extraction of formaldehyde by the Hantzsch reaction 896(2000)51
- Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291
- Optimisation of alachlor solid-phase microextraction from water samples using experimental design 896(2000)373

***Fagus sylvatica***

Retention index database for identification of general green leaf volatiles in plants by coupled capillary gas chromatography–mass spectrometry 890(2000)313

**Fat retainers**

Utilization of fat retainers in supercritical fluid extraction for the selective extraction of polychlorinated biphenyls from a model fat sample 877(2000)167

**Fats**

Chromatographic analysis of unsaponifiable compounds of olive oils and fat-containing foods 881(2000)105

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Polycyclic aromatic hydrocarbons in edible fats and oils: occurrence and analytical methods 882(2000)245

**Fibers**

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

**Fibre optical scanning**

Fibre optical scanning with high resolution in thin-layer chromatography 898(2000)265

**Field-amplified sample injection/stacking**

see Field-enhanced sample stacking

**Field-enhanced sample stacking**

Field amplified injection in the presence of salts for capillary electrophoresis 853(1999)3

Head-column field-amplified sample stacking in binary system capillary electrophoresis. Preparation of extracts for determination of opioids in microliter amounts of body fluids 853(1999)95

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

Standardization of capillary zone electropherograms obtained by using field enhanced sample stacking 894(2000)3

**Field-flow fractionation**

Different elution modes and field programming in gravitational field-flow fractionation. 2. Experimental verification of the range of conditions for flow-rate and carrier liquid density programming 868(2000)217

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

**Field programming**

Different elution modes and field programming in gravitational field-flow fractionation. 2. Experimental verification of the range of conditions for flow-rate and carrier liquid density programming 868(2000)217

**Filters**

Adaptive numerical morphological filter for identifying chromatographic signals 855(1999)305

**Filtration**

Membrane-based sample preparation coupled on-line to chromatography or electrophoresis 856(1999)55

**Fish**

Use of capillary electrophoresis with UV detection as a screening method to determine histamine in fish samples 853(1999)541

Liquid chromatographic determination of benzocaine and N-acetylbenzocaine in the edible fillet tissue from rainbow trout 855(1999)255

Substitution of fish species detected by thin-layer isoelectric focusing and a computer-assisted method for the evaluation of gels 880(2000)303

Determination of aflatoxins in food by overpressured-layer chromatography 882(2000)11

Determination of tripolyphosphate in frozen cod and scallop adductor by ion chromatography 884(2000)89

Validation of an analytical procedure for polychlorinated biphenyls, coplanar polychlorinated biphenyls and polycyclic aromatic hydrocarbons in environmental samples 886(2000)153

**Flame photometric detection**

Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149

**Flame retardants**

Polymer additive analysis by pyrolysis–gas chromatography. II. Flame retardants 886(2000)225

**Flame thermionic detection**

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatroscan (mark 5) and flame thermionic detection 855(1999)657

**Flash chromatography**

On-column refractive index detector for flash chromatography 886(2000)9

**Flavours**

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection 867(2000)187

**Flotation reagents**

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

**Flow cell**

Miniaturized thin-layer radial flow cell with interdigitated ring-shaped microarray electrode used as amperometric detector for capillary electrophoresis 891(2000)149

**Flow distributor**

Visualization of sample introduction in liquid chromatographic columns. Contribution of a flow distributor on the sample band shape 865(1999)83

**Flow injection analysis**

Enzyme amplification as detection tool in continuous-flow systems. I. Development of an enzyme-amplified biochemical detection system coupled on-line to flow-injection analysis 855(1999)383

Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159

**Flow microcalorimetry**

Study of hydrophobic interaction adsorption of bovine serum albumin under overloaded conditions using flow microcalorimetry 865(1999)111

**Flow patterns**

Influence of flow patterns on chromatographic efficiency in centrifugal partition chromatography 869(2000)339

**Flow rate**

Pellicular expanded bed matrix suitable for high flow rates 878(2000)17

Relationship between retention, linear velocity and flow for counter-current chromatography 886(2000)283

**Flow velocity**

Use of equations for the description of experimental dependence of the height equivalent to a theoretical plate on carrier gas velocity in capillary gas–liquid chromatography 872(2000)111

Relationship between retention, linear velocity and flow for counter-current chromatography 886(2000)283

**Flow velocity distribution**

Influence of column radial heterogeneity on peak fronting in linear chromatography 857(1999)69

**Fluidised bed**

Immobilised metal affinity chromatography of  $\beta$ -galactosidase from unclarified *Escherichia coli* homogenates using expanded bed adsorption 874(2000)27

Pellicular expanded bed matrix suitable for high flow rates 878(2000)17

**Fluofix**

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

**Fluorenyl stationary phases**

*n*-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133

**Fluorescence detection**

see also Laser-induced fluorescence detection

Labeling effects on the isoelectric point of green fluorescent protein 853(1999)21

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159

Time-resolved high-performance liquid chromatography fluorescence detector using a nanosecond pulsed light source for detecting lanthanide-chelated compounds 859(1999)13

High-sensitivity chromatographic detector incorporating three-dimensional charge coupled device fluorimetry 867(2000)93

Quantitative determination of sulfonamide residues in foods of animal origin by high-performance liquid chromatography with fluorescence detection 871(2000)37

Utilization of fluorescein sodium salt for the indirect fluorimetric detection in micellar electrokinetic chromatography 872(2000)247

Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159

Rapid analysis of covalently and non-covalently fluorophore-labeled proteins using ultra-thin-layer sodium dodecylsulfate gel electrophoresis 894(2000)329

**Fluorescence line narrowing spectroscopy**

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159



**Fluorescence non-line narrowing spectroscopy**

On-line identification of diastereomeric dibenzo[*a,l*]pyrene diol epoxide-derived deoxyadenosine adducts by capillary electrophoresis–fluorescence line-narrowing and non-line narrowing spectroscopy 853(1999)159

**Fluorescence polarization**

Capillary electrophoresis coupled with laser-induced fluorescence polarization as a hybrid approach to ultrasensitive immunoassays 853(1999)555

**Fluorinated bonded phases**

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

**Fluorinated silica gel columns**

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

**Flux-corrected transport**

Simulation of electrophoretic separations by the flux-corrected transport method 890(2000)321

**Food analysis**

see also Special Issue Index

Use of capillary electrophoresis with UV detection as a screening method to determine histamine in fish samples 853(1999)541

Separation and identification of various carotenoids by C<sub>30</sub> reversed-phase high-performance liquid chromatography coupled to UV and atmospheric pressure chemical ionization mass spectrometric detection 854(1999)37

Determination of nivalenol and deoxynivalenol in wheat using liquid chromatography–mass spectrometry with negative ion atmospheric pressure chemical ionisation 854(1999)45

Fractionation of apple procyanidins by size-exclusion chromatography 855(1999)181

Determination of folate vitamers in food and in Italian reference diet by high-performance liquid chromatography 855(1999)237

Liquid chromatographic determination of benzocaine and *N*-acetylbenzocaine in the edible fillet tissue from rainbow trout 855(1999)255

Liquid chromatographic–mass spectrometric determination of phenolic compounds using a capillary-scale particle beam interface 855(1999)515

Determination of oak lactones in barrel-aged wines and in oak extracts by stable isotope dilution analysis 857(1999)239

Use of ion chromatography for the determination of heavy and transition metals in biochemical samples 857(1999)343

2-Alkylcyclobutanones as markers for irradiated foodstuffs. III. Improvement of the field of application on the EN 1785 method by using silver ion chromatography 858(1999)109

Inclusion of the fungicide chlozolinate in a multiresidue method 858(1999)123

Determination of the *Fusarium* mycotoxins, fusaproliferin and beauvericin by high-performance liquid chromatography–electrospray ionization mass spectrometry 858(1999)175

Quantification of deoxynivalenol in wheat using an immunoaffinity column and liquid chromatography 859(1999)23

Analysis of bovine whey proteins in soybean dairy-like products by capillary electrophoresis 859(1999)77

Analysis of some folate monoglutamates by high-performance liquid chromatography–mass spectrometry. I 864(1999)59

Semipreparative reversed-phase liquid chromatographic fractionation of aroma extracts from wine and other alcoholic beverages 864(1999)77

Determination of ochratoxin A in wine by means of immunoaffinity column clean-up and high-performance liquid chromatography 864(1999)89

High-performance liquid chromatography of selenium compounds utilizing perfluorinated carboxylic acid ion-pairing agents and inductively coupled plasma and electrospray ionization mass spectrometric detection 866(2000)51

Ultra trace detection of a wide range of anabolic steroids in meat by gas chromatography coupled to mass spectrometry 867(2000)219

Erratum to: “Liquid chromatographic method for determining added folic acid in fortified cereal products” [J. Chromatogr. A, 826 (1998) 235–240] 867(2000)289

Does further clean-up reduce the matrix enhancement effect in gas chromatographic analysis of pesticide residues in food? 868(2000)51

Determination of heterocyclic aromatic amines in meat extracts by liquid chromatography–ion-trap atmospheric pressure chemical ionization mass spectrometry 869(2000)307

Retention/quantitation properties of the *o*-phthalaldehyde–3-mercaptopropionic acid and the *o*-phthalaldehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

- High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483
- Quantitation of anabolic hormones and their metabolites in bovine serum and urine by liquid chromatography–tandem mass spectrometry 870(2000)511
- Anion-exchange high-performance liquid chromatography with conductivity detection for the analysis of phytic acid in food 871(2000)7
- Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13
- Analysis of tetrahydro- $\beta$ -carboline-3-carboxylic acids in foods by solid-phase extraction and reversed-phase high-performance liquid chromatography combined with fluorescence detection 871(2000)23
- Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67
- Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75
- Determination of phenols in wines by liquid chromatography with photodiode array and fluorescence detection 871(2000)85
- Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95
- Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403
- High-performance liquid chromatographic determination of  $\Delta^9$ -tetrahydrocannabinol and the corresponding acid in hemp containing foods with special regard to the fluorescence properties of  $\Delta^9$ -tetrahydrocannabinol 872(2000)101
- Determination of sulphur compounds in beer using headspace solid-phase microextraction and gas chromatographic analysis with pulsed flame photometric detection 872(2000)203
- Utilization of fluorescein sodium salt for the indirect fluorimetric detection in micellar electrokinetic chromatography 872(2000)247
- Gas chromatographic–mass spectrometric analysis of acids and phenols in distilled alcohol beverages. Application of anion-exchange disk extraction combined with in-vial elution and silylation 873(2000)29
- Erratum to: “Trace-level determination of polar flavour compounds in butter by solid-phase extraction and gas chromatography–mass spectrometry” [J. Chromatogr. A, 844 (1999) 295–305] 873(2000)299
- Separation of polyphenols in Canary Islands wine by capillary zone electrophoresis without preconcentration 874(2000)111
- Automatic microgravimetric determination of fats in milk products by use of supercritical fluid extraction with on-line piezoelectric detection 874(2000)265
- Direct chiral resolution of lactic acid in food products by capillary electrophoresis 875(2000)371
- Development of a capillary electrophoresis method for the simultaneous analysis of artificial sweeteners, preservatives and colours in soft drinks 876(2000)213
- Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235
- Amperometric determination of underivatized amino acids at a nickel-modified gold electrode by anion-exchange chromatography 878(2000)57
- Determination of the plant growth regulator chlormequat in food by liquid chromatography–electrospray ionisation tandem mass spectrometry 878(2000)77
- Simultaneous separation and quantitation of the major bovine whey proteins including proteose peptone and caseinomacropeptide by reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene 878(2000)183
- Determination of cow’s milk in non-bovine and mixed cheeses by capillary electrophoresis of whey proteins in acidic isoelectric buffers 878(2000)261
- Sample preparation for the analysis of flavors and off-flavors in foods 880(2000)3
- Applications of solid-phase microextraction in food analysis 880(2000)35
- Applications of matrix solid-phase dispersion in food analysis 880(2000)63
- Application of ion-exchange cartridge clean-up in food analysis. III. Determination of benzylpenicillin, phenoxymethyl-penicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in bovine liver and kidney by liquid chromatography with ultraviolet detection 880(2000)85
- Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxypyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93

- Comparison of different commercial solid-phase extraction cartridges used to extract heterocyclic amines from a lyophilised meat extract 880(2000)101
- Determination of 1-nitropyrene in herbs after selective enrichment by a sol–gel-generated immunoaffinity column 880(2000)113
- Determination of fenpyroximate in apples by supercritical fluid extraction and packed capillary liquid chromatography with UV detection 880(2000)121
- Thin-layer chromatography in food and agricultural analysis 880(2000)129
- Direct analysis of food samples by high-performance liquid chromatography 880(2000)149
- Perfusion chromatography: an emergent technique for the analysis of food proteins 880(2000)169
- C<sub>30</sub> Stationary phases for the analysis of food by liquid chromatography 880(2000)189
- Determination of aldehydes in food by high-performance liquid chromatography with biosensor coupling and micromembrane suppressors 880(2000)233
- Electrophoretic methods in the analysis of beverages 880(2000)243
- Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295
- Substitution of fish species detected by thin-layer isoelectric focusing and a computer-assisted method for the evaluation of gels 880(2000)303
- Analysis of natural food pigments by capillary electrophoresis 880(2000)311
- Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides 881(2000)1
- Determination by perfusion reversed-phase high-performance liquid chromatography of the soybean protein content of commercial soybean products prepared directly from whole soybeans 881(2000)37
- Characterization of commercial soybean products by conventional and perfusion reversed-phase high-performance liquid chromatography and multivariate analysis 881(2000)47
- Evaluation of the effect of rennet type on casein proteolysis in an ovine milk cheese by means of capillary electrophoresis 881(2000)59
- Effect of the activity levels of the added proteolytic enzyme mixture on free amino acids in ripening Ossau-Iraty cheese 881(2000)69
- Chromatographic determination of amino acid enantiomers in beers and raw materials used for their manufacture 881(2000)81
- Chromatographic analysis of unsaponifiable compounds of olive oils and fat-containing foods 881(2000)105
- Chromatographic analysis of minor constituents in vegetable oils 881(2000)131
- Analysis of olive and hazelnut oil mixtures by high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry of triacylglycerols and gas–liquid chromatography of non-saponifiable compounds (tocopherols and sterols) 881(2000)149
- Development of phosphine reagents for the high-performance liquid chromatographic–fluorometric determination of lipid hydroperoxides 881(2000)159
- Determination of vitamins in food-matrix Standard Reference Materials 881(2000)171
- Determination of vitamin D<sub>2</sub> in emulsified nutritional supplements by solid-phase extraction and column-switching high-performance liquid chromatography with UV detection 881(2000)189
- Chromatographic analysis of tocol-derived lipid antioxidants 881(2000)197
- Normal-phase high-performance liquid chromatography of tocopherols and tocotrienols. Comparison of different chromatographic columns 881(2000)217
- Automated analysis of vitamin E isomers in vegetable oils by continuous membrane extraction and liquid chromatography–electrochemical detection 881(2000)229
- Determination of tocopherol acetate in emulsified nutritional supplements by solid-phase extraction and high-performance liquid chromatography with fluorescence detection 881(2000)243
- Rapid determination of vitamin E in vegetable oils by reversed-phase high-performance liquid chromatography 881(2000)251
- Simultaneous determination of  $\alpha$ -tocopherol and  $\beta$ -carotene in olive oil by reversed-phase high-performance liquid chromatography 881(2000)255
- Determination of vitamin K<sub>1</sub> in emulsified nutritional supplements by solid-phase extraction and high-performance liquid chromatography with postcolumn reduction on a platinum catalyst and fluorescence detection 881(2000)261

- Determination of thiamine by high-performance liquid chromatography 881(2000)267
- Chromatographic determination of riboflavin and its derivatives in food 881(2000)285
- Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables 881(2000)309
- Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317
- Use of nucleic acids in the mobile phase for the determination of ascorbic acid in foods by high-performance liquid chromatography with electrochemical detection 881(2000)327
- Analytical techniques for determining biotin 881(2000)331
- Picomolar quantitation of free sulfite in foods by means of [<sup>57</sup>Co]hydroxocobalamin and radiometric chromatography of [<sup>57</sup>Co]sulfitecobalamin. Method, applications and significance of coexisting sulfides 881(2000)345
- Determination of organic acids in food samples by capillary zone electrophoresis 881(2000)357
- High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365
- Influence of organic solvents in the mobile phase on the determination of carboxylic acids and inorganic anions in grape juice by ion chromatography 881(2000)387
- Development and validation of a capillary electrophoresis method for direct measurement of isocitric, citric, tartaric and malic acids as adulteration markers in orange juice 881(2000)395
- Analysis of anthocyanins in foods by liquid chromatography, liquid chromatography–mass spectrometry and capillary electrophoresis 881(2000)403
- Determination of tea catechins 881(2000)411
- Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425
- Comparative analysis of tea catechins and theaflavins by high-performance liquid chromatography and capillary electrophoresis 881(2000)439
- Spectrophotometric and coulometric detection in the high-performance liquid chromatography of flavonoids and optimization of sample treatment for the determination of quercetin in orange juice 881(2000)449
- Analysis of several phenolic compounds with potential antioxidant properties in grape extracts and wines by high-performance liquid chromatography–photodiode array detection without sample preparation 881(2000)461
- Analysis of resveratrol in wine by capillary electrophoresis 881(2000)471
- Analysis of the bioactive alkaloids tetrahydro- $\beta$ -carboline and  $\beta$ -carboline in food 881(2000)483
- Liquid chromatographic determination of biogenic amines in fermented foods after derivatization with 3,5-dinitrobenzoyl chloride 881(2000)517
- Changes in free and conjugated polyamines during starvation of sugarcane juices as analyzed by high-performance liquid chromatography 881(2000)531
- Chromatographic determination of carotenoids in foods 881(2000)543
- Monoterpenes in grape juice and wines 881(2000)557
- Analysis of organic sulfur compounds in wine aroma 881(2000)569
- Analysis of low-volatility organic sulphur compounds in wines by solid-phase microextraction and gas chromatography 881(2000)583
- Determination of inulin in meat products by high-performance liquid chromatography with refractive index detection 881(2000)591
- High-performance liquid chromatographic determination of Maillard compounds in store-brand and name-brand ultra-high-temperature-treated cows' milk 881(2000)599
- Advances in the determination of inorganic anions by ion chromatography 881(2000)607
- Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629
- Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629
- Analysis of condensed phosphates in food products by ion chromatography with an on-line hydroxide eluent generator 881(2000)639
- Determination of lithium in wines by ion chromatography 881(2000)645
- Application of the assay of aflatoxins by liquid chromatography with fluorescence detection in food analysis 882(2000)1
- Determination of aflatoxins in food by overpressured-layer chromatography 882(2000)11

- Chromatographic determination of the mycotoxin patulin in fruit and fruit juices 882(2000)17
- Simultaneous determination of trichothecene mycotoxins and zearalenone in cereals by gas chromatography–mass spectrometry 882(2000)23
- Solid-phase extraction applied to the determination of ochratoxin A in wines by reversed-phase high-performance liquid chromatography 882(2000)29
- Regulatory problems caused by contamination, a frequently overlooked cause of veterinary drug residues 882(2000)37
- Determination of trenbolone and zeranol in bovine muscle and liver by liquid chromatography–electrospray mass spectrometry 882(2000)53
- Quantitation of nine quinolones in chicken tissues by high-performance liquid chromatography with fluorescence detection 882(2000)63
- Determination of clopidol residues in chicken tissues by high-performance liquid chromatography–mass spectrometry 882(2000)85
- Determination of four nitroimidazole residues in poultry meat by liquid chromatography–mass spectrometry 882(2000)89
- Simultaneous determination of triclabendazole and its sulphoxide and sulphone metabolites in bovine milk by high-performance liquid chromatography 882(2000)99
- Chromatographic analysis of tetracycline antibiotics in foods 882(2000)109
- Stability of penicillin antibiotic residues in meat during storage. Ampicillin 882(2000)135
- Determination of cephalosporins in raw bovine milk by high-performance liquid chromatography 882(2000)145
- Pesticide residue determination in fruit and vegetables by liquid chromatography–mass spectrometry 882(2000)153
- Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175
- Simplified method for the determination of chlorinated fungicides and insecticides in fruits by gas chromatography 882(2000)193
- Multiresidue analysis of seventeen pesticides in wine by gas chromatography with mass-selective detection 882(2000)205
- Multiresidue method for the rapid determination of organophosphorus insecticides in grapes, must and wine 882(2000)213
- Multiresidue method for the rapid determination – in grape, must and wine – of fungicides frequently used on vineyards 882(2000)221
- Interlaboratory study of identification and quantitation of multiresidue pyrethroids in agricultural products by gas chromatography–mass spectrometry 882(2000)231
- Gas chromatographic determination of acrinathrine and 3-phenoxybenzaldehyde residues in honey 882(2000)239
- Contamination in food from packaging material 882(2000)255
- Determination of synthetic phenolic antioxidants in food by high-performance liquid chromatography 882(2000)271
- Application of chromatographic and electrophoretic methodology to the speciation of organomercury compounds in food analysis 882(2000)283
- Preparation of 4,4'-(1-[<sup>2</sup>H<sub>6</sub>]methylene)bis-[2,3,5,6-<sup>2</sup>H<sub>4</sub>]-phenol and its application to the measurement of bisphenol A in beverages by stable isotope dilution mass spectrometry 883(2000)163
- Rapid high-performance liquid chromatography method for the analysis of sodium benzoate and potassium sorbate in foods 883(2000)299
- Determination of tripolyphosphate in frozen cod and scallop adductor by ion chromatography 884(2000)89
- Solid-phase extraction of quaternary ammonium herbicides 885(2000)251
- High-speed counter-current chromatography of apple procyanidins 886(2000)65
- Solid-phase microextraction with on-fibre derivatisation applied to the analysis of volatile carbonyl compounds 886(2000)175
- Combined ion-pair extraction and gas chromatography–mass spectrometry for the simultaneous determination of diamines, polyamines and aromatic amines in Port wine and grape juice 886(2000)183
- Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85
- Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 888(2000)113
- High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

- Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129
- Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography 888(2000)321
- Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annuum* L.) powders 889(2000)41
- Solid-phase extraction and high-performance liquid chromatographic separation of pigments of red wines 889(2000)51
- Multi-residue methodology for pesticide screening in wines 889(2000)59
- Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69
- Role of chromatography in the analysis of sugars, carboxylic acids and amino acids in food (Review) 891(2000)1
- Analysis of agrochemicals by capillary electrophoresis (Review) 891(2000)45
- Determination of fumaric acid in apple juice by on-line coupled capillary isotachopheresis–capillary zone electrophoresis with UV detection 891(2000)175
- Gas chromatographic determination of the fatty-acid content of heat-treated green beans 891(2000)367
- Pesticide residue analysis in foodstuffs applying capillary gas chromatography with mass spectrometric detection. State-of-the-art use of modified DFG-multimethod S19 and automated data evaluation 892(2000)347
- Determination of pesticides in vegetables using large-volume injection column liquid chromatography–electrospray tandem mass spectrometry 892(2000)379
- Applications of mass spectrometry to food proteins and peptides 893(2000)1
- Determination of sugar alcohols in confectioneries by high-performance liquid chromatography after nitrobenzoylation 893(2000)195
- Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79
- Separation and quantitation of colour pigments of chili powder (*Capsicum frutescens*) by high-performance liquid chromatography–diode array detection 896(2000)69
- Liquid chromatographic determination of emamectin, milbemectin, ivermectin and abamectin in crops and confirmation by liquid chromatography–mass spectrometry 896(2000)75
- High-performance liquid chromatographic determination with photodiode array detection of ellagic acid in fresh and processed fruits 896(2000)87
- Determination of cholesterol oxides in processed food using high-performance liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 896(2000)321
- Determination of 1-aminocyclopropane-1-carboxylic acid and its structural analogue by liquid chromatography and ion spray tandem mass spectrometry 896(2000)335
- Characterisation of whiskeys using solid-phase microextraction with gas chromatography–mass spectrometry 896(2000)351
- Characterization of odorant compounds of mussels (*Mytilus edulis*) according to their origin using gas chromatography–olfactometry and gas chromatography–mass spectrometry 896(2000)361
- Separation and determination of alditols and sugars by high-pH anion-exchange chromatography with pulsed amperometric detection 897(2000)195
- Supercritical fluid extraction of catechins from *Cratogeomys pruifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75
- Application of ion-exchange cartridge clean-up in food analysis. V. Simultaneous determination of sulphonamide antibacterials in animal liver and kidney using high-performance liquid chromatography with ultraviolet and mass spectrometric detection 898(2000)95
- Determination of dyes in foodstuffs by capillary zone electrophoresis 898(2000)271
- Food simulants**  
Use of solid-phase microextraction for the analysis of bisphenol A and bisphenol A diglycidyl ether in food simulants 864(1999)137
- Forensic analysis**  
Method for quantification of morphine and its 3- and 6-glucuronides, codeine, codeine glucuronide and 6-monoacetylmorphine in human blood by liquid chromatography–electrospray mass spectrometry for routine analysis in forensic toxicology 854(1999)109
- Profiling of impurities in illicit methamphetamine by high-performance liquid chromatography and capillary electrochromatography 870(2000)53

Application of solid-phase microextraction to the recovery of explosives and ignitable liquid residues from forensic specimens 885(2000)419

Efficiency of pretreatment of aqueous samples using a macroporous strong anion-exchange resin on the determination of nerve gas hydrolysis products by gas chromatography–mass spectrometry after *tert*.-butyldimethylsilylation 891(2000)295

Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23

Comparison of phenyl-type columns in the development of a fast liquid chromatographic system for eighteen opiates commonly found in forensic toxicology 896(2000)311

#### **Fourier-transform infrared spectrometry**

Liquid chromatography–Fourier-transform infrared spectrometry 856(1999)213

#### **Frit diameter**

Visualization of sample introduction in liquid chromatography columns. The effect of the frit diameter 855(1999)367

#### ***Fritillaria* spp.**

Simultaneous determination of seven major isosteroidal alkaloids in bulbs of *Fritillaria* by gas chromatography 873(2000)221

#### **Fritless packed capillaries**

Improved column preparation and performance in capillary electrochromatography 887(2000)367

#### **Frontal analysis**

Enantioselective binding analysis of verapamil to plasma lipoproteins by capillary electrophoresis–frontal analysis 875(2000)447

Determination of the lumped mass transfer rate coefficient by frontal analysis 890(2000)211

#### **Frontal chromatography**

Reinforcement of frontal affinity chromatography for effective analysis of lectin–oligosaccharide interactions 890(2000)261

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

#### **Fruit juices**

Fractionation of apple procyanidins by size-exclusion chromatography 855(1999)181

Electrophoretic methods in the analysis of beverages 880(2000)243

Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Influence of organic solvents in the mobile phase on the determination of carboxylic acids and inorganic anions in grape juice by ion chromatography 881(2000)387

Development and validation of a capillary electrophoresis method for direct measurement of isocitric, citric, tartaric and malic acids as adulteration markers in orange juice 881(2000)395

Spectrophotometric and coulometric detection in the high-performance liquid chromatography of flavonoids and optimization of sample treatment for the determination of quercetin in orange juice 881(2000)449

Monoterpenes in grape juice and wines 881(2000)557

Chromatographic determination of the mycotoxin patulin in fruit and fruit juices 882(2000)17

Rapid high-performance liquid chromatography method for the analysis of sodium benzoate and potassium sorbate in foods 883(2000)299

Combined ion-pair extraction and gas chromatography–mass spectrometry for the simultaneous determination of diamines, polyamines and aromatic amines in Port wine and grape juice 886(2000)183

Determination of fumaric acid in apple juice by on-line coupled capillary isotachopheresis–capillary zone electrophoresis with UV detection 891(2000)175

Determination of patulin in apple juice by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 897(2000)365

#### **Fruits**

see also Apples and Grapes

Fractionation of apple procyanidins by size-exclusion chromatography 855(1999)181

Inclusion of the fungicide chlozolinate in a multiresidue method 858(1999)123

Retention/quantitation properties of the *o*-phthaldialdehyde–3-mercaptopropionic acid and the *o*-phthaldialdehyde–*N*-acetyl-L-cysteine amino acid derivatives in reversed-phase high-performance liquid chromatography 870(2000)271

Study of polyphenols in grape berries by reversed-phase high-performance liquid chromatography 870(2000)449

High-performance liquid chromatographic screening of chlorophyll derivatives produced during fruit storage 870(2000)483

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Determination of fenpyroximate in apples by supercritical fluid extraction and packed capillary liquid chromatography with UV detection 880(2000)121

Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables 881(2000)309

Analysis of anthocyanins in foods by liquid chromatography, liquid chromatography–mass spectrometry and capillary electrophoresis 881(2000)403

Chromatographic determination of the mycotoxin patulin in fruit and fruit juices 882(2000)17

Pesticide residue determination in fruit and vegetables by liquid chromatography–mass spectrometry 882(2000)153

Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175

Simplified method for the determination of chlorinated fungicides and insecticides in fruits by gas chromatography 882(2000)193

Interlaboratory study of identification and quantitation of multiresidue pyrethroids in agricultural products by gas chromatography–mass spectrometry 882(2000)231

High-performance liquid chromatographic determination with photodiode array detection of ellagic acid in fresh and processed fruits 896(2000)87

#### **Fuels**

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

#### **Fullerene adsorbents**

Analytical potential of fullerene as adsorbent for organic and organometallic compounds from aqueous solutions 869(2000)101

#### **Fullerenes-extracted soot**

Fullerenes-extracted soot: a new adsorbent for collecting volatile organic compounds in ambient air 886(2000)313

#### ***Fusarium subglutinans***

Determination of the *Fusarium* mycotoxins, fusaproliferin and beauvericin by high-performance liquid chromatography–electrospray ionization mass spectrometry 858(1999)175

## G

#### **Gamma radiation**

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by  $\gamma$ -radiation on titanium-grafted silica 869(2000)137

#### ***Gardenia jasminoides***

Analysis of natural food pigments by capillary electrophoresis 880(2000)311

#### **Gas chromatography**

New operational modes for multidimensional and comprehensive gas chromatography by using cryogenic modulation 866(2000)203

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

Time-resolved cryogenic modulation for targeted multidimensional capillary gas chromatography analysis 892(2000)15

Comparison of comprehensive two-dimensional gas chromatography and gas chromatography–mass spectrometry for the characterization of complex hydrocarbon mixtures 892(2000)29

#### **Gas chromatography, dynamic**

Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487



**Gas chromatography, fast**

Evaluation of time-of-flight mass spectrometric detection for fast gas chromatography 878(2000)205

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

**Gas chromatography, comprehensive two-dimensional**

Comprehensive two-dimensional gas chromatography: a hyphenated method with strong coupling between the two dimensions 856(1999)331

Comparison of comprehensive two-dimensional gas chromatography and gas chromatography–mass spectrometry for the characterization of complex hydrocarbon mixtures 892(2000)29

**Gas chromatography, high-speed**

High-speed gas chromatography: an overview of various concepts 856(1999)315

**Gas chromatography–mass spectrometry**

see also Mass spectrometry

Comparison of comprehensive two-dimensional gas chromatography and gas chromatography–mass spectrometry for the characterization of complex hydrocarbon mixtures 892(2000)29

**Gas–liquid separation interface**

Understanding of peak deterioration in hyphenated speciation systems due to gas–liquid separation in the hydride generation interface 855(1999)291

**Gasoline**

Analysis of methyl *tert.*-butyl ether and its degradation products by direct aqueous injection onto gas chromatography with mass spectrometry or flame ionization detection systems 857(1999)205

**Gas-phase luminescence detection**

Gas chromatographic determination of primary alcohols as their formylbenzoic esters by gas-phase luminescence detection 855(1999)583

**Generalised *H*-point standard addition method**

Generalised *H*-point standard addition method for the isolation of the analyte signal from the sample signal when coelution of unknown compounds occurs in liquid chromatography 852(1999)361

**Geochemistry**

Improved capillary electrophoresis method for measuring rare-earth elements in synthetic geochemical standards 884(2000)317

**Geological samples**

On-line preconcentration of niobium(V) and tantalum(V) as 4-(2-pyridylazo) resorcinol–citrate ternary complexes in geological samples by ion interaction high-performance liquid chromatography 885(2000)369

**Gerbils**

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

**Glassy carbon**

Assessment of polycrystalline graphites as sorbents for solid-phase microextraction of nonionic surfactants 888(2000)35

***Glycyrrhiza uralensis***

Migration behavior and separation of active components in *Glycyrrhiza uralensis* Fisch and its commercial extract by micellar electrokinetic capillary chromatography 863(1999)105

**Gradient elution**

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

Prediction of protein retention at gradient elution conditions in ion-exchange chromatography 852(1999)351

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. II. Two-run assay procedures 857(1999)21

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. III. Improving the accuracy of computer simulation 857(1999)41

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Separation of polyesters by gradient reversed-phase high-performance liquid chromatography on a 1.5  $\mu\text{m}$  non-porous column 871(2000)243

Novel simulated moving-bed method for reduced solvent consumption 873(2000)149

Analysis of condensed phosphates in food products by ion chromatography with an on-line hydroxide eluent generator 881(2000)639

Gradient elution techniques for capillary electrochromatography (Review) 887(2000)115

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. I. Effect of buffer composition on the gradient shape 890(2000)25

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. II. Separation of proteins 890(2000)37

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Comparative study of capillary electroendosmotic chromatography and electrically assisted gradient nano-liquid chromatography for the separation of peptides 894(2000)241

Pressurized gradient capillary electrochromatographic separation of eighteen amino acid derivatives 894(2000)337

Optimization strategy and validation of one chromatographic method as approach to determine the phenolic compounds from different sources 897(2000)161

#### **Gradient polymer elution chromatography**

Analysis of organosilicone copolymers by gradient polymer elution chromatography with evaporative light scattering detection 871(2000)269

#### **Gradient time**

Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. I. Optimizing selectivity and resolution 897(2000)37

Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. II. Minimizing column reproducibility problems 897(2000)51

#### **Granular activated carbons**

Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

#### **Grapes**

see also Fruits

Analysis of anthocyanins in foods by liquid chromatography, liquid chromatography–mass spectrometry and capillary electrophoresis 881(2000)403

Analysis of several phenolic compounds with potential antioxidant properties in grape extracts and wines by high-performance liquid chromatography–photodiode array detection without sample preparation 881(2000)461

Multiresidue method for the rapid determination of organophosphorus insecticides in grapes, must and wine 882(2000)213

Multiresidue method for the rapid determination – in grape, must and wine – of fungicides frequently used on vineyards 882(2000)221

#### **Graphite**

Assessment of polycrystalline graphites as sorbents for solid-phase microextraction of nonionic surfactants 888(2000)35

#### **Graphitized carbon black**

see also Porous graphitized carbon

Determination of inorganic anions by ion chromatography using a graphitized carbon column dynamically coated with cetyltrimethylammonium ions 855(1999)261

Stability of organophosphorus insecticides on graphitized carbon black extraction cartridges used for large volumes of surface water 879(2000)73

Graphitized carbons for solid-phase extraction 885(2000)73

#### **Gravitational field-flow fractionation**

Different elution modes and field programming in gravitational field-flow fractionation. 2. Experimental verification of the range of conditions for flow-rate and carrier liquid density programming 868(2000)217

#### **Grignard pentylation**

Direct Grignard pentylation of organotin-contaminated lard samples followed by capillary gas chromatography with flame photometric detection 886(2000)197

#### **Group-correlation method**

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

#### **Group method**

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

#### **Growth factors**

Strategy for identifying circulating fragments of insulin-like growth factor binding proteins in a hemofiltrate peptide bank 852(1999)197

Renaturation of heterodimeric platelet-derived growth factor from inclusion bodies of recombinant *Escherichia coli* using size-exclusion chromatography 855(1999)203

Continuous-bed chromatography for the analysis and purification of recombinant human basic fibroblast growth factor 857(1999)137

# H

## Hair

Determination of hair polyamines as *N*-ethoxycarbonyl-*N*-pentafluoropropionyl derivatives by gas chromatography–mass spectrometry 897(2000)295

## Half-normal plots

Robustness testing of a reversed-phase high-performance liquid chromatographic assay: comparison of fractional and asymmetrical factorial designs 874(2000)167

## Hanging mercury drop electrode

Selective and sensitive assay for the determination of benzodiazepines by high-performance liquid chromatography with simultaneous ultraviolet and reductive electrochemical detection at the hanging mercury drop electrode 897(2000)215

## Hantzsch reaction

Liquid chromatography–mass spectrometry method for the determination of aldehydes derivatized by the Hantzsch reaction 864(1999)191

Simultaneous supercritical fluid derivatization and extraction of formaldehyde by the Hantzsch reaction 896(2000)51

## Header design

Visualization of viscous fingering in high-performance liquid chromatographic columns. Influence of the header design 865(1999)73

## Headspace analysis

Determination of sulphur compounds in beer using headspace solid-phase microextraction and gas chromatographic analysis with pulsed flame photometric detection 872(2000)203

Analysis of volatile contaminants in vegetable oils by headspace solid-phase microextraction with Carboxen-based fibres 873(2000)79

Development of a headspace solid-phase microextraction procedure for the determination of free volatile fatty acids in waste waters 873(2000)107

Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117

Trace analysis of ten chlorinated benzenes in water by headspace solid-phase microextraction 874(2000)149

Applications of solid-phase microextraction in food analysis 880(2000)35

Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxypyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93

Application of solid-phase microextraction for determination of organic vapours in gaseous matrices 885(2000)405

Role of solid-phase microextraction in the identification of highly volatile pheromones of two Rhinoceros beetles *Scapanes australis* and *Strategus aloeus* (Coleoptera, Scarabaeidae, Dynastinae) 885(2000)433

Influence of fibre coating in headspace solid-phase microextraction–gas chromatographic analysis of aromatic and medicinal plants 892(2000)469

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

Simultaneous determination of cyanogen chloride and cyanogen bromide in treated water at sub- $\mu\text{g/L}$  levels by a new solid-phase microextraction–gas chromatographic–electron-capture detection method 897(2000)307

## Heat of adsorption

Heats of adsorption of some organic compounds on  $\beta$ -cyclodextrin determined by gas–solid chromatography 864(1999)293

Application of neutral hydrophobic hypercrosslinked polystyrene to the separation of inorganic anions by ion chromatography 884(2000)41

## Heat transfer

Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227

## Height equivalent to a theoretical plate

Use of equations for the description of experimental dependence of the height equivalent to a theoretical plate on carrier gas velocity in capillary gas–liquid chromatography 872(2000)111

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

## *Helicobacter pylori*

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

## Hemp

High-performance liquid chromatographic determination of  $\Delta^9$ -tetrahydrocannabinol and the corresponding acid in hemp containing foods with special regard to the fluorescence properties of  $\Delta^9$ -tetrahydrocannabinol 872(2000)101

**Heparin-modified stationary phases**

Separation of enantiomers on anion exchangers modified with heparin in liquid chromatography 864(1999)25

**Hepatitis B virus**

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

Detection of duck hepatitis B virus DNA fragments using on-column intercalating dye labeling with capillary electrophoresis–laser-induced fluorescence 853(1999)309

**Hepatitis C virus**

Introduction of the gene amplification technique to decrease the risk of hepatitis C virus transmission by plasma products 852(1999)305

**Heteroconjugation**

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

***Hevea brasiliensis***

Characterisation of the rubber elongation factor from ammoniated latex by electrophoresis and mass spectrometry 890(2000)145

**High-performance liquid chromatography, fast**

Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3

**Hindered amine stabilizers**

Hindered amine stabilizers investigated by the use of packed capillary temperature-programmed liquid chromatography. I. Poly(6-(1,1,3,3-tetramethylbutyl)-amino)-1,3,5-triazine-2,4-diyl(2,2,6,6-tetramethyl-4-piperidyl)imino)-1,6-hexanediyl (2,2,6,6-tetramethyl-4-piperidyl)imino) 874(2000)65

**Hold-up times**

Hold-up time in gas chromatography. V. Dependence of the retention of *n*-alkanes on the chromatographic variables in isothermal gas chromatography 878(2000)125

**Hold-up volume**

Low-energy interactions in high-performance liquid chromatography 872(2000)49

**Hollow fibers**

Development of a simple in-vial liquid-phase microextraction device for drug analysis compatible with capillary gas chromatography, capillary electrophoresis and high-performance liquid chromatography 873(2000)3

**Hollow-fiber membranes**

High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43

**Honey**

Determination of rotenone residues in raw honey by solid-phase extraction and high-performance liquid chromatography 871(2000)67

Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629

Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography 881(2000)629

Gas chromatographic determination of acrinathrine and 3-phenoxybenzaldehyde residues in honey 882(2000)239

**Hot spots**

Gas chromatographic–mass spectrometric separation, identification and determination of C<sub>6</sub>–C<sub>12</sub> cyclic imides in thermally treated epoxy and alkyd resins to locate hot spots inside large electricity generators 862(1999)17

**Hot water extraction**

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

**Human immunodeficiency virus**

Determination of human immunodeficiency virus-1 protease inhibitors in patient serum using free solution capillary zone electrophoresis 853(1999)431

One-step capillary isoelectric focusing for the separation of the recombinant human immunodeficiency virus envelope glycoprotein glycoforms 866(2000)121

**Hummel–Dreyer method**

Binding of lead to collagen type I and V and  $\alpha_2(I)$  CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

**Hydride generation**

Understanding of peak deterioration in hyphenated speciation systems due to gas–liquid separation in the hydride generation interface 855(1999)291

**Hydrodynamic injection**

Field amplified injection in the presence of salts for capillary electrophoresis 853(1999)3

**Hydrodynamic lift forces**

Different elution modes and field programming in gravitational field-flow fractionation. 2. Experimental verification of the range of conditions for flow-rate and carrier liquid density programming 868(2000)217

**Hydrodynamic radius**

Modelling of the pore flow in capillary electrochromatography 887(2000)199

**Hydrogels**

Determination of morphine by capillary electrophoresis immunoassay in thermally reversible hydrogel-modified buffer and laser-induced fluorescence detection 895(2000)1

**Hydrolysis**

Analysis of carbohydrates in wood and pulps employing enzymatic hydrolysis and subsequent capillary zone electrophoresis 891(2000)157

Determination of urinary catecholamines with capillary electrophoresis after solid-phase extraction 895(2000)317

**Hydrolytic stability**

Dependence of cyano bonded phase hydrolytic stability on ligand structure and solution pH 893(2000)245

**Hydrophobic interaction chromatography**

Study of hydrophobic interaction adsorption of bovine serum albumin under overloaded conditions using flow microcalorimetry 865(1999)111

Studies on the chromatographic fractionation of *Trichoderma reesei* cellulases by hydrophobic interaction 865(1999)123

Protein losses in ion-exchange and hydrophobic interaction high-performance liquid chromatography 890(2000)73

**Hydrophobicity**

see also Lipophilicity

Chromatographic properties of reversed-phase stationary phases under pressure- and electro-driven conditions 862(1999)121

Fractionation and toxicity evaluation of waste waters 889(2000)149

**Hydrophobic selectivity**

Chromatographic properties of reversed-phase stationary phases under pressure- and electro-driven conditions 862(1999)121

**Hydrosilation**

Open tubular capillary electrokinetic chromatography in etched fused-silica tubes (Review) 887(2000)31

**Hydroxyapatite**

Improved liquid chromatographic separation of different proteins by designing functional surfaces of cattle bone-originated apatite 862(1999)217

Discoloration of ceramic hydroxyapatite used for protein chromatography 891(2000)93

**Hydroxyl radicals**

Identification of the oxidation products of the reaction between  $\alpha$ -pinene and hydroxyl radicals by gas and high-performance liquid chromatography with mass spectrometric detection 896(2000)135

***Hypericum perforatum***

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

**Hyphenated techniques**

see also Special Issue Index

Size-exclusion chromatography with on-line ultraviolet, proton nuclear magnetic resonance and mass spectrometric detection and on-line collection for off-line Fourier transform infrared spectroscopy 857(1999)89

Multiple hyphenation of liquid chromatography with nuclear magnetic resonance spectroscopy, mass spectrometry and beyond 892(2000)315

**I****Ice crystals**

Analysis of size-classified ice crystals by capillary electrophoresis 871(2000)391

**Ignitable liquids**

Application of solid-phase microextraction to the recovery of explosives and ignitable liquid residues from forensic specimens 885(2000)419

**Imidazole coating**

Preparation and evaluation of an imidazole-coated capillary column for the electrophoretic separation of aromatic acids 897(2000)383

**Immobilization**

Biomembrane-affinity centrifugal analyses of solute interactions with membrane proteins 852(1999)93

Towards the column bed stabilization of columns in capillary electroendosmotic chromatography. Immobilization of microparticulate silica columns to a continuous bed 887(2000)327

**Immobilized antibodies**

Antibody immobilization to high-performance liquid chromatography supports. Characterization of maximum loading capacity for intact immunoglobulin G and Fab fragments 888(2000)13

**Immobilized biomembranes**

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

**Immobilized cellulose derivatives**

Separation of enantiomers by packed capillary electrochromatography on a cellulose-based stationary phase 875(2000)331

**Immobilized chaperonins**

Lysozyme refolding with immobilized GroEL column chromatography 878(2000)197

**Immobilized chymotrypsin**

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

High concentrations of sodium chloride facilitate the use of immobilized chymotrypsin for separating virgin forms of specific trypsin inhibitors 852(1999)227

**Immobilized enzymes**

Microscale characterization of the structure–activity relationship of a heparin-binding glycopeptide using affinity capillary electrophoresis and immobilized enzymes 853(1999)189

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

On-line automated high-performance liquid chromatographic determination of total riboflavin phosphates using immobilized acid phosphatase as a pre-column reactor 896(2000)171

**Immobilized metal affinity chromatography**

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105

Histidines in affinity tags and surface clusters for immobilized metal-ion affinity chromatography of trimeric tumor necrosis factor  $\alpha$  852(1999)117

Affinity purification of *Schistosoma japonicum* glutathione-S-transferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

Characterisation by immobilised metal ion affinity chromatographic procedures of the binding behaviour of several synthetic peptides designed to have high affinity for Cu(II) ions 852(1999)261

Phosphopeptide analysis by on-line immobilized metal-ion affinity chromatography–capillary electrophoresis–electrospray ionization mass spectrometry 853(1999)225

Natural poly-histidine affinity tag for purification of recombinant proteins on cobalt(II)-carboxymethylaspartate crosslinked agarose 864(1999)247

Adsorption and selectivity characteristics of several human serum proteins with immobilised hard Lewis metal ion–chelate adsorbents 890(2000)95

Designing new metal affinity peptides by random mutagenesis of a natural metal-binding site 898(2000)83

**Immobilized proteins**

Eliminating disulfide exchange during glutamyl endopeptidase digestion of native protein 864(1999)237

Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy 893(2000)235

**Immunoabsorbents**

see also Affinity adsorbents

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

Quantification of deoxynivalenol in wheat using an immunoaffinity column and liquid chromatography 859(1999)23

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

**Immunoaffinity chromatography**

see also Affinity chromatography

Sandwich-type immunoassay with an affinity column coupled to anti-peptide antibodies 852(1999)97

Quantitative fast fractionation of a pool of polyclonal antibodies by immunoaffinity membrane chromatography 852(1999)129

Determination of 1-nitropyrene in herbs after selective enrichment by a sol–gel-generated immunoaffinity column 880(2000)113

Antibody immobilization to high-performance liquid chromatography supports. Characterization of maximum loading capacity for intact immunoglobulin G and Fab fragments 888(2000)13

**Immunoassays**

Strategy for identifying circulating fragments of insulin-like growth factor binding proteins in a hemofiltrate peptide bank 852(1999)197

Competitive immunoassay for staphylococcal enterotoxin A using capillary electrophoresis with laser-induced fluorescence detection 853(1999)545

Capillary electrophoresis coupled with laser-induced fluorescence polarization as a hybrid approach to ultrasensitive immunoassays 853(1999)555

Determination of morphine by capillary electrophoresis immunoassay in thermally reversible hydrogel-modified buffer and laser-induced fluorescence detection 895(2000)1

Analysis of codeine, dihydrocodeine and their glucuronides in human urine by electrokinetic capillary immunoassays and capillary electrophoresis–ion trap mass spectrometry 895(2000)133

#### **Immunoextraction**

Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29

#### **Imprinted polymers**

see also Molecular imprinting

Measurement of electroosmotic flow in plastic imprinted microfluid devices and the effect of protein adsorption on flow rate 857(1999)275

#### **In-capillary reaction micellar electrokinetic chromatography**

Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295

#### **Indirect detection**

Indirect detection of organic acids in non-aqueous capillary electrophoresis 853(1999)171

Determination of tartaric acid in solid wine residues by capillary electrophoresis and indirect UV detection 853(1999)181

Utilization of fluorescein sodium salt for the indirect fluorimetric detection in micellar electrokinetic chromatography 872(2000)247

Analysis of glycosaminoglycan monosaccharides by capillary electrophoresis using indirect laser-induced fluorescence detection 873(2000)269

Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113

New preconditioning strategy for the determination of inorganic anions with capillary zone electrophoresis using indirect UV detection 890(2000)355

#### **Inductively coupled plasma mass spectrometry**

see also Mass spectrometry

Liquid chromatography–inductively coupled plasma mass spectrometry 856(1999)243

Speciation of selenium compounds with ion-pair reversed-phase liquid chromatography using inductively coupled plasma mass spectrometry as element-specific detection 874(2000)55

#### **Industrial effluents**

Efficient solid-phase extraction procedures for trace enrichment of priority phenols from industrial effluents with high total organic carbon content 857(1999)97

#### **Infant formulae**

Ion chromatographic determination of inositol in infant formulae and clinical products for enteral feeding 879(2000)129

#### **Infrared spectrometry**

Liquid chromatography–Fourier-transform infrared spectrometry 856(1999)213

Gas chromatography with spectroscopic detectors 856(1999)349

#### **Injection methods**

see also Large-volume injections, Sample introduction and Sample stacking

Field amplified injection in the presence of salts for capillary electrophoresis 853(1999)3

Head-column field-amplified sample stacking in binary system capillary electrophoresis. Preparation of extracts for determination of opioids in microliter amounts of body fluids 853(1999)95

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

Programmed temperature vaporiser-based injection in capillary gas chromatography 856(1999)259

Coupling of biological sample handling and capillary electrophoresis 856(1999)483

Analysis of methyl *tert*.-butyl ether and its degradation products by direct aqueous injection onto gas chromatography with mass spectrometry or flame ionization detection systems 857(1999)205

Coupling device for desorption of drugs from solid-phase extraction-pipette tips and on-line gas chromatographic analysis 858(1999)117

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

Fast chiral separations using sulfated  $\beta$ -cyclodextrin and short-end injection in capillary electrophoresis 875(2000)403

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. I. Injection procedure and column stability 876(2000)37

Introduction of large volumes of water-containing samples into a gas chromatograph. Improved retention of volatile solutes through the swing system 876(2000)135

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. II. Methods of detection and quantification and overall long-term performance 878(2000)45

Direct analysis of food samples by high-performance liquid chromatography 880(2000)149

Split-flow injector for capillary zone electrophoresis 883(2000)223

Efficiency through combining high-performance liquid chromatography and high resolution gas chromatography: progress 1995–1999 892(2000)407

Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111

Video-taped sample evaporation in hot chambers simulating gas chromatography split/splitless injectors. I. Thermospray injection 897(2000)237

Video-taped sample evaporation in hot chambers simulating gas chromatography split and splitless injectors. II. Injection with band formation 897(2000)247

Robustness testing for a capillary electrophoresis method using the “short-end injection” technique 897(2000)329

### **Injector dispersion**

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

### **Injector performance**

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

### **Instrumentation**

Large-scale membrane adsorbers 852(1999)73

Application of non-aqueous capillary electrophoresis with electrochemical detection to the determination of nicotine in tobacco 853(1999)27

Nanoliter-scale sample preparation methods directly coupled to polymethylmethacrylate-based microchips and gel-filled capillaries for the analysis of oligonucleotides 853(1999)107

Interface for capillary electrophoresis coupled with inductively coupled plasma atomic emission spectrometry 853(1999)141

Side-entry excitation and detection of square capillary array electrophoresis for DNA sequencing 853(1999)359

New sheathless interface for coupling capillary electrophoresis to electrospray mass spectrometry evaluated by the analysis of fatty acids and prostaglandins 854(1999)141

Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

Modified apparatus for voltage gradient gel electrophoresis 855(1999)723

Comprehensive two-dimensional gas chromatography: a hyphenated method with strong coupling between the two dimensions 856(1999)331

Gas chromatography with spectroscopic detectors 856(1999)349

Integrated approach to the multidimensional analysis of complex biological samples by microseparation techniques. Analysis of glycoprotein factor associated with cancer cachexia 857(1999)183

Measurement of electroosmotic flow in plastic imprinted microfluid devices and the effect of protein adsorption on flow rate 857(1999)275

Coupling device for desorption of drugs from solid-phase extraction-pipette tips and on-line gas chromatographic analysis 858(1999)117

Time-resolved high-performance liquid chromatography fluorescence detector using a nanosecond pulsed light source for detecting lanthanide-chelated compounds 859(1999)13

Nonaqueous capillary electrophoresis equipped with amperometric detection for analysis of chlorinated phenolic compounds 864(1999)323

Visualization of solute migration in chromatographic columns. Quantitation of the concentration in a migrating zone 867(2000)71

Apparatus and method for interfacing capillary electrophoresis and chemiluminescence nitrogen detection for the analysis of nitrogen-containing compounds 868(2000)121

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

Design of a simulated moving bed unit for sucrose–betaine separations 872(2000)23



- Investigation of Cr(III) hydrolytic polymerisation products by capillary electrophoresis–inductively coupled plasma–mass spectrometry 872(2000)227
- On-column amperometric detection in capillary electrophoresis with an improved high-voltage electric field decoupler 872(2000)259
- Introduction of large volumes of water-containing samples into a gas chromatograph. Improved retention of volatile solutes through the swing system 876(2000)135
- Application of electron-attachment reactions to enhance selectivity of electron-capture detector for nitroaromatic explosives 877(2000)133
- Dynamic microwave-assisted extraction 877(2000)141
- Conductivity detection of polymerase chain reaction products separated by micro-reversed-phase liquid chromatography 883(2000)1
- Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. II: High-temperature gas chromatography–chemical ionization mass spectrometry 883(2000)75
- Split-flow injector for capillary zone electrophoresis 883(2000)223
- Protein identification platform utilizing micro dispensing technology interfaced to matrix-assisted laser desorption ionization time-of-flight mass spectrometry 886(2000)99
- Instrumentation for capillary electrochromatography 887(2000)55
- Scale-down of continuous protein purification by annular chromatography. Design parameters for the smallest unit 890(2000)53
- Capillary electrophoresis–inductively coupled plasma mass spectrometry interface with minimised dead volume for high separation efficiency 891(2000)129
- Simple interface for capillary electrophoresis–inductively coupled plasma atomic emission spectrometry 891(2000)139
- Miniaturized thin-layer radial flow cell with interdigitated ring-shaped microarray electrode used as amperometric detector for capillary electrophoresis 891(2000)149
- Novel microfabricated device for electrokinetically induced pressure flow and electrospray ionization mass spectrometry 892(2000)195
- Advances in column technology and instrumentation in capillary electrochromatography 892(2000)257
- Electron–capture mass spectrometry: recent advances 892(2000)329
- Integration of a microextraction system. Solvent extraction of a Co–2-nitroso-5-dimethylaminophenol complex on a microchip 894(2000)19
- Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45
- Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117
- Integrated electroosmotically-driven on-line sample purification system for nanoliter DNA sequencing by capillary electrophoresis 894(2000)179
- Rapid, parallel separations of D1S80 alleles in a plastic microchannel chip 894(2000)203
- Surface-ionization methods and devices of indication and identification of nitrogen-containing base molecules 896(2000)3
- Interfaces**  
see also Mass spectrometry
- Interfaces, CE–chemiluminescence nitrogen detection**  
Apparatus and method for interfacing capillary electrophoresis and chemiluminescence nitrogen detection for the analysis of nitrogen-containing compounds 868(2000)121
- Interfaces, CEC–MS**  
Use of on-line mass spectrometric detection in capillary electrochromatography (Review) 887(2000)85
- Interfaces, CE–ICP–AES**  
Interface for capillary electrophoresis coupled with inductively coupled plasma atomic emission spectrometry 853(1999)141
- Simple interface for capillary electrophoresis–inductively coupled plasma atomic emission spectrometry 891(2000)139
- Interfaces, CE–MS**  
New sheathless interface for coupling capillary electrophoresis to electrospray mass spectrometry evaluated by the analysis of fatty acids and prostaglandins 854(1999)141
- Capillary electrophoresis–inductively coupled plasma mass spectrometry interface with minimised dead volume for high separation efficiency 891(2000)129
- Novel microfabricated device for electrokinetically induced pressure flow and electrospray ionization mass spectrometry 892(2000)195

**Interfaces, GC–MS**

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. II: High-temperature gas chromatography–chemical ionization mass spectrometry 883(2000)75

**Interfaces, LC–FT-IR**

Liquid chromatography–Fourier-transform infrared spectrometry 856(1999)213

**Interfaces, LC–GC**

Efficiency through combining high-performance liquid chromatography and high resolution gas chromatography: progress 1995–1999 892(2000)407

**Interfaces, LC–MS**

Liquid chromatographic–mass spectrometric determination of phenolic compounds using a capillary-scale particle beam interface 855(1999)515

Liquid chromatography–inductively coupled plasma mass spectrometry 856(1999)243

Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides 881(2000)1

Pesticide residue determination in fruit and vegetables by liquid chromatography–mass spectrometry 882(2000)153

Protein identification platform utilizing micro dispensing technology interfaced to matrix-assisted laser desorption ionization time-of-flight mass spectrometry 886(2000)99

**Interfaces, LC–NMR–MS–IR**

Size-exclusion chromatography with on-line ultraviolet, proton nuclear magnetic resonance and mass spectrometric detection and on-line collection for off-line Fourier transform infrared spectroscopy 857(1999)89

**Interfaces, SFC–MS**

Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

**Interfaces, solid-phase extraction-pipette tips–programmed-temperature vaporiser**

Coupling device for desorption of drugs from solid-phase extraction-pipette tips and on-line gas chromatographic analysis 858(1999)117

**Interfaces, SPE–GC**

On-line coupling of solid-phase extraction to gas chromatography with fast solvent vaporization and concentration in an open injector liner. Analysis of pesticides in aqueous samples 896(2000)41

**Interfaces, SPE–MS**

On-line coupling of solid-phase extraction with mass spectrometry for the analysis of biological samples. I. Determination of clenbuterol in urine 863(1999)115

**Interfaces, SPME–HPLC**

Automated in-tube solid-phase microextraction–high-performance liquid chromatography for carbamate pesticide analysis 873(2000)137

**Interfaces, TLC–MS**

The state of the art in thin-layer chromatography–mass spectrometry: a critical appraisal 856(1999)429

**Interlaboratory studies**

Interlaboratory study of identification and quantitation of multiresidue pyrethroids in agricultural products by gas chromatography–mass spectrometry 882(2000)231

Quantitative analysis of mutagenic heterocyclic aromatic amines in cooked meat using liquid chromatography–atmospheric pressure chemical ionisation tandem mass spectrometry 883(2000)89

Inter-laboratory comparison of liquid chromatographic techniques and enzyme-linked immunosorbent assay for the determination of surfactants in wastewaters 889(2000)195

Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the ng l<sup>-1</sup> concentration level 893(2000)367

**Intraparticle convective flow**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

**Intraparticle electroosmotic flow**

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Peclet number in capillary electrochromatography from pore network theory 890(2000)61

**Intraparticle electroosmotic velocity**

Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Peclet number in capillary electrochromatography from pore network theory 890(2000)61

**Intraparticle Peclet number**

Modeling the velocity field of the electroosmotic flow in charged capillaries and in capillary columns packed with charged particles: interstitial and intraparticle velocities in capillary electrochromatography systems 877(2000)181

**Inverse gas chromatography**

Effect of surface heterogeneity on adsorption on solid surfaces. Application of inverse gas chromatography in the studies of energetic heterogeneity of adsorbents 886(2000)133

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

Investigation of the nematic–isotropic transition of a liquid crystalline polymer and determination of molecular diffusion coefficients using gas chromatography 893(2000)359

**Inverse size-exclusion chromatography**

Pore size distributions of cation-exchange adsorbents determined by inverse size-exclusion chromatography 883(2000)39

**Iodination**

Investigation on the iodination reaction of methylene blue by liquid chromatography–mass spectrometry with ionspray ionisation 854(1999)57

**Ion chromatography**

see also Special Issue Index

Advances in the determination of inorganic anions by ion chromatography 881(2000)607

**Ion chromatography–capillary electrophoresis**

Separation of anions by ion chromatography–capillary electrophoresis 879(2000)245

Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261

**Ion electrokinetic chromatography**

Potential of ethylenediaminedi(*o*-hydroxyphenylacetic acid) and *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid for the determination of metal ions by capillary electrophoresis 890(2000)363

Separation selectivity of anionic metal complexes of *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid in ion and ion electrokinetic chromatography 895(2000)255

**Ion-exchange chromatography**

Cytochrome *c* unfolding on an anionic surface 863(1999)137

Optimization of ion-exchange displacement separations. I. Validation of an iterative scheme and its use as a methods development tool 876(2000)51

Optimization of ion-exchange displacement separations. II. Comparison of displacement separations on various ion-exchange resins 876(2000)63

**Ion-exchange electrochromatography**

Separation of acidic compounds by strong anion-exchange capillary electrochromatography 887(2000)223

**Ion-exclusion/cation-exchange chromatography**

Simultaneous ion-exclusion/cation-exchange chromatography of anions and cations in acid rain waters on a weakly acidic cation-exchange resin by elution with sulfosalicylic acid 884(2000)167

**Ion exchangers**

see also Anion exchangers, Cation exchangers and Stationary phases

Pore size distributions of cation-exchange adsorbents determined by inverse size-exclusion chromatography 883(2000)39

Protein adsorption on novel acrylamido-based polymeric ion-exchangers. I. morphology and equilibrium adsorption 897(2000)65

Protein adsorption on novel acrylamido-based polymeric ion exchangers. II. Adsorption rates and column behavior 897(2000)81

Comparison of chromatographic ion-exchange resins. I. Strong anion-exchange resins 897(2000)99

**Ionic strength**

Effect of ionic strength, pH and polymer concentration on the separation of DNA fragments in the presence of electroosmotic flow 894(2000)219

**Ion–ion interactions**

Electrophoretic mobility study on ion–ion interactions in an aqueous solution 853(1999)63

Coupling of chromatographic ion-exchange reaction with change-partner reaction. Presumed mechanism on the ion–ion interaction of inorganic ions in Sephadex G-15 columns 869(2000)91

**Ionization mechanisms**

State-of-the-art in liquid chromatography–mass spectrometry 856(1999)179

**Ion-pair chromatography**

Reversed-phase ion-pair chromatography coupled to electrospray ionisation mass spectrometry by on-line removal of the counter-ions 854(1999)155

Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23

**Ion-pair solid-phase extraction**

Ion-pair solid-phase extraction 885(2000)343

**Ion-pairing reagents**

Use of volatile amines as ion-pairing agents for the high-performance liquid chromatographic–tandem mass spectrometric determination of aromatic sulfonates in industrial wastewater 854(1999)175

Determination of 20 underivatized proteinic amino acids by ion-pairing chromatography and pneumatically assisted electrospray mass spectrometry 855(1999)191

High-performance liquid chromatography of selenium compounds utilizing perfluorinated carboxylic acid ion-pairing agents and inductively coupled plasma and electrospray ionization mass spectrometric detection 866(2000)51

Combined ion-pair extraction and gas chromatography–mass spectrometry for the simultaneous determination of diamines, polyamines and aromatic amines in Port wine and grape juice 886(2000)183

High-performance ion-pair chromatographic behaviour of conjugated bile acids with di-*n*-butylamine acetate 888(2000)93

Development of a method for the analysis of nucleotides from the mantle tissue of the mussel *Mytilus galloprovincialis* 891(2000)99

Capillary electrophoresis of seed 2S albumins from *Lupinus* species 894(2000)297

**Isobutoxycarbonyl derivatization**

Gas chromatographic profiling and screening for phenols as isobutoxycarbonyl derivatives in aqueous samples 866(2000)87

**Isoelectric buffers**

General experimental aspects of the use of isoelectric buffers in capillary electrophoresis 853(1999)71

Separation of double-stranded DNA in conventional and isoelectric buffers: studies on stability and separation performance 859(1999)87

**Isoelectric focusing**

Chemiluminescence detection of heme proteins separated by capillary isoelectric focusing 852(1999)597

Labeling effects on the isoelectric point of green fluorescent protein 853(1999)21

The unvalidity of Kohlrausch' regulating function for Svensson's isoelectric focusing and stationary electrolysis at steady state 863(1999)219

One-step capillary isoelectric focusing for the separation of the recombinant human immunodeficiency virus envelope glycoprotein glycoforms 866(2000)121

Substitution of fish species detected by thin-layer isoelectric focusing and a computer-assisted method for the evaluation of gels 880(2000)303

Experimental study on moving neutralization reaction boundary created with the strong reactive electrolytes of HCl and NaOH in agarose gel 891(2000)337

**Isoelectric points**

Ion-exchange chromatography of proteins near the isoelectric points 852(1999)31

**Isotachophoresis**

Coupling of biological sample handling and capillary electrophoresis 856(1999)483

The unvalidity of Kohlrausch' regulating function for Svensson's isoelectric focusing and stationary electrolysis at steady state 863(1999)219

On-line isotachophoretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219

Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117

**Isotachophoresis–capillary zone electrophoresis**

Enantiomeric separation of clenbuterol by transient isotachophoresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachophoresis 871(2000)173

Determination of fumaric acid in apple juice by on-line coupled capillary isotachophoresis–capillary zone electrophoresis with UV detection 891(2000)175

**Isothermal titration calorimetry**

Microcalorimetric studies of interactions between proteins and hydrophobic ligands in hydrophobic interaction chromatography: effects of ligand chain length, density and the amount of bound protein 872(2000)37

**Isotope analysis**

Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis 884(2000)261

Comparison of chromatographic methods for hydrogen isotope separation by Pd beds 898(2000)125

**K****Kerosene**

Determination of phenolic antioxidants in aviation jet fuel 871(2000)235

**Kinetic studies**

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum–rhodium alloy catalysts 852(1999)507

Kinetic study of the mass transfer of bovine serum albumin in anion-exchange chromatography 866(2000)147

Kinetics of solid-phase extraction and solid-phase microextraction in thin adsorbent layer with saturation sorption isotherm 873(2000)39

Kinetics of appearance of hemorphins from bovine hemoglobin peptic hydrolysates by a direct coupling of reversed-phase high-performance liquid chromatography and electrospray ionization mass spectrometry 873(2000)185

Enantioselective determination of drugs in body fluids by capillary electrophoresis 875(2000)27

Determination of U(IV) and U(VI) by ion chromatography–inductively coupled plasma mass spectrometry and its application to kinetic studies 884(2000)131

Determination of epichlorohydrin by ion chromatography 884(2000)251

Determination of the lumped mass transfer rate coefficient by frontal analysis 890(2000)211

Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487

**Kohlrausch function**

The invalidity of Kohlrausch' regulating function for Svensson's isoelectric focusing and stationary electrolysis at steady state 863(1999)219

Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117

**Kraft pulp**

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

**Kromasil C<sub>18</sub>**

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. III. Results obtained with Kromasil C<sub>18</sub> columns 855(1999)423

**L*****Lachesis stenophrys***

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

**Lard**

Direct Grignard pentylation of organotin-contaminated lard samples followed by capillary gas chromatography with flame photometric detection 886(2000)197

**Large-volume injections**

see also Injection methods

Programmed temperature vaporiser-based injection in capillary gas chromatography 856(1999)259

Analysis of linear alkylbenzenesulfonates in water samples by large-volume injection-port derivatization and gas chromatography–mass spectrometry 857(1999)359

Determination of low levels of an antioxidant in polyolefins by large-volume injection temperature-programmed packed capillary liquid chromatography 864(1999)103

Trace analysis of pesticide residues in water by high-speed narrow-bore capillary gas chromatography–mass spectrometry with programmable temperature vaporizer 874(2000)81

Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153

Comparison of various liquid chromatographic methods involving UV and atmospheric pressure chemical ionization mass spectrometric detection for the efficient trace analysis of phenylurea herbicides in various types of water samples 879(2000)39

Determination of 1-(2-methoxyphenyl)piperazine derivatives of isocyanates at low concentrations by temperature-programmed miniaturized liquid chromatography 892(2000)67

Determination of pesticides in vegetables using large-volume injection column liquid chromatography–electrospray tandem mass spectrometry 892(2000)379

Efficiency through combining high-performance liquid chromatography and high resolution gas chromatography: progress 1995–1999 892(2000)407

Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111

**Laser-induced dispersed fluorescence detection**

Laser-induced dispersed fluorescence detection of polycyclic aromatic compounds in soil extracts separated by capillary electrochromatography 883(2000)231

**Laser-induced fluorescence detection**

Anthracycline analysis by capillary electrophoresis. Application to the analysis of daunorubicin in Kaposi sarcoma tumor 853(1999)449

Competitive immunoassay for staphylococcal enterotoxin A using capillary electrophoresis with laser-induced fluorescence detection 853(1999)545

**Laser-induced fluorescence polarization detection**

Capillary electrophoresis coupled with laser-induced fluorescence polarization as a hybrid approach to ultrasensitive immunoassays 853(1999)555

**Latex**

Characterisation of the rubber elongation factor from ammoniated latex by electrophoresis and mass spectrometry 890(2000)145

Development and validation of a capillary electrophoresis method for the measurement of short-chain organic acids in natural rubber latex 894(2000)135

***Lathyrus sativus***

Analysis of  $\beta$ -*N*-oxalyl-L- $\alpha$ , $\beta$ -diaminopropionic acid and homoarginine in *Lathyrus sativus* by capillary zone electrophoresis 857(1999)295

Determination of neurotoxin 3-*N*-oxalyl-2,3-diaminopropionic acid and non-protein amino acids in *Lathyrus sativus* by precolumn derivatization with 1-fluoro-2,4-dinitrobenzene 883(2000)113

**Lattice–fluid models**

Lattice–fluid model for gas–liquid chromatography 862(1999)49

**Lead–protein interactions**

Binding of lead to collagen type I and V and  $\alpha_2$ (I) CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

**Least-squares analysis**

Determination of amino acids in overlapped capillary electrophoresis peaks by means of partial least-squares regression 871(2000)331

**Libraries, MS**

Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411

**Ligand fishing**

Micropreparative ligand fishing with a cuvette-based optical mirror resonance biosensor 869(2000)261

**Ligand structure**

Dependence of cyano bonded phase hydrolytic stability on ligand structure and solution pH 893(2000)245

**Ligand-exchange capillary electrophoresis**

Chiral separation of unmodified amino acids with non-aqueous capillary electrophoresis based on the ligand-exchange principle 874(2000)285

Application of ligand-exchange capillary electrophoresis to the chiral separation of  $\alpha$ -hydroxy acids and  $\beta$ -blockers 875(2000)307

**Ligand-exchange chromatography**

Analysis of the enzymatic racemization of D-aspartic acid to L-aspartic acid by the on-line coupling of a solid-phase extraction column and a ligand-exchange high-performance liquid chromatography column 896(2000)299

**Light-harvesting complex**

Resolution and identification of the protein components of the photosystem II antenna system of higher plants by reversed-phase liquid chromatography with electrospray-mass spectrometric detection 886(2000)111

**Limiting retention factors**

Reversed-phase high-performance liquid chromatography of ionogenic compounds: comparison of retention models 859(1999)1

**Linear free energy relationships**

Characterization of reversed-phase columns using the linear free energy relationship. III. Effect of the organic modifier and the mobile phase composition 893(2000)214

**Linear solvation energy relationships**

Effects of organic modifiers on retention mechanism and selectivity in micellar electrokinetic capillary chromatography studied by linear solvation energy relationships 863(1999)69

Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335

Revised linear solvation energy relationship coefficients for the 77-phase McReynolds data set based on an updated set of solute descriptors 878(2000)115

Statistical evaluation of linear solvation energy relationship models used to characterize chemical selectivity in micellar electrokinetic chromatography 886(2000)245

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

Liquid chromatography–electrospray mass spectrometry of multicomponent peptide mixtures. Characterization of a mixture from the synthesis of the hormone goserelin 889(2000)119

Analysis of a peptide hormone mixture of therapeutic interest by liquid chromatography coupled to high-flow pneumatically assisted electrospray mass spectrometry 893(2000)95

Calculation of Abraham solute descriptors from McReynolds gas chromatographic retention data 893(2000)339

Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23

#### **Liner performance**

Video-taped sample evaporation in hot chambers simulating gas chromatography split and splitless injectors. II. Injection with band formation 897(2000)247

#### **Lipophilicity**

see also Hydrophobicity

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

Use of solid-phase microextraction for measuring oil–water partition coefficients and correlation with high-performance liquid chromatographic methods for lipophilicity 864(1999)183

Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49

#### **Liquid chromatography–gas chromatography**

Efficiency through combining high-performance liquid chromatography and high resolution gas chromatography: progress 1995–1999 892(2000)407

#### **Liquid chromatography–mass spectrometry**

see also Mass spectrometry and Special Issue Index

Reversed-phase ion-pair chromatography coupled to electrospray ionisation mass spectrometry by on-line removal of the counterions 854(1999)155

State-of-the-art in liquid chromatography–mass spectrometry 856(1999)179

Application of liquid chromatography with mass spectrometry combined with photodiode array detection and tandem mass spectrometry for monitoring pesticides in surface waters 879(2000)51

Pesticide residue determination in fruit and vegetables by liquid chromatography–mass spectrometry 882(2000)153

Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435

#### **Liquid crystalline polymers**

Synthesis and properties of two new liquid crystals: an analytical and thermodynamic study 859(1999)59

Investigation of the nematic–isotropic transition of a liquid crystalline polymer and determination of molecular diffusion coefficients using gas chromatography 893(2000)359

Laterally attached liquid crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. II. Optimization of the molecular parameter of the polymer 897(2000)131

#### **Liquid-phase microextraction**

Development of a simple in-vial liquid-phase microextraction device for drug analysis compatible with capillary gas chromatography, capillary electrophoresis and high-performance liquid chromatography 873(2000)3

#### **Low-energy interactions**

Low-energy interactions in high-performance liquid chromatography 872(2000)49

#### **Luminescence detection**

Gas chromatographic determination of primary alcohols as their formylbenzoic esters by gas-phase luminescence detection 855(1999)583

#### **Luna C<sub>18</sub> (2)**

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C<sub>18</sub> (2) columns 869(2000)181

#### **Lupinus spp.**

Capillary electrophoresis of seed 2S albumins from *Lupinus* species 894(2000)297

#### **Lycium chinense**

Determination of betaine in *Lycium chinense* fruits by liquid chromatography–electrospray ionization mass spectrometry 857(1999)331

## M

#### **Magnetapheresis**

Analytical magnetapheresis of magnetically susceptible particles 874(2000)131

#### **Magnetic particles**

Analytical magnetapheresis of magnetically susceptible particles 874(2000)131

**Magnetic separation**

Magnetic split-flow thin fractionation of magnetically susceptible particles 857(1999)193

**Magnetic sorbents**

Application of magnetite and silica–magnetite composites to the isolation of genomic DNA 890(2000)159

**Magnetite sorbents**

Application of magnetite and silica–magnetite composites to the isolation of genomic DNA 890(2000)159

**Malaprade's reaction**

Determination of cyclodextrin content using periodate oxidation by capillary electrophoresis 891(2000)201

**Marine sediments**

see also Sediments

Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515

Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211

**Marjoram**

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

**Mass spectrometry**

see also Capillary electrophoresis-mass spectrometry, Gas chromatography-mass spectrometry, Inductively coupled plasma mass spectrometry, Interfaces, Liquid chromatography-mass spectrometry, Matrix-assisted laser desorption ionization mass spectrometry, Supercritical fluid chromatography-mass spectrometry, Tandem mass spectrometry, Special Issue Index and specific ionization techniques

Preliminary comparison of precursor scans and liquid chromatography–tandem mass spectrometry on a hybrid quadrupole time-of-flight mass spectrometer 854(1999)119

Atmospheric pressure ionisation multiple mass spectrometric analysis of pesticides 854(1999)207

Characterization of nonyl phenol ethoxylates in sewage treatment plants by combined precursor ion scanning and multiple reaction monitoring 854(1999)245

Application of electrospray mass spectrometry in the detection and determination of Remazol textile dyes 854(1999)259

State-of-the-art in liquid chromatography–mass spectrometry 856(1999)179

Gas chromatography with spectroscopic detectors 856(1999)349

The state of the art in thin-layer chromatography–mass spectrometry: a critical appraisal 856(1999)429

Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry 856(1999)515

Analysis of 8-aminonaphthalene-1,3,6-trisulfonate-derivatized oligosaccharides by capillary electrophoresis–electrospray ionization quadrupole ion trap mass spectrometry 858(1999)229

Detection of new sequences of peptaibol antibiotics trichothoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179

On-line coupling of solid-phase extraction with mass spectrometry for the analysis of biological samples. I. Determination of clenbuterol in urine 863(1999)115

Comparison of gas chromatography–pulsed flame photometric detection–mass spectrometry, automated mass spectral deconvolution and identification system and gas chromatography–tandem mass spectrometry as tools for trace level detection and identification 868(2000)229

Application of spectral libraries for high-performance liquid chromatography–atmospheric pressure ionisation mass spectrometry to the analysis of pesticide and explosive residues in environmental samples 869(2000)411

Evaluation of time-of-flight mass spectrometric detection for fast gas chromatography 878(2000)205

Investigation of combwax of honeybees with high-temperature gas chromatography and high-temperature gas chromatography–chemical ionization mass spectrometry. II: High-temperature gas chromatography–chemical ionization mass spectrometry 883(2000)75

Comparison of on-line coupling of ion-chromatography with atmospheric pressure ionization mass spectrometry and with inductively coupled plasma mass spectrometry as tools for the ultra-trace analysis of bromate in surface water samples 884(2000)191

Selective deuterium exchange during superheated heavy water chromatography–nuclear magnetic resonance spectroscopy–mass spectrometry of sulfonamides 886(2000)289

Use of on-line mass spectrometric detection in capillary electrochromatography 887(2000)85

Capillary electrochromatography–mass spectrometry for the separation and identification of isomeric polyaromatic hydrocarbon DNA adducts derived from in vitro reactions (Review) 887(2000)103



Separation and identification of etodolac and its urinary phase I metabolites using capillary electrochromatography and on-line capillary electrochromatography–electrospray ionisation mass spectrometry coupling 887(2000)393

Novel microfabricated device for electrokinetically induced pressure flow and electrospray ionization mass spectrometry 892(2000)195

Electron–capture mass spectrometry: recent advances 892(2000)329

Applications of mass spectrometry to food proteins and peptides 893(2000)1

Methanol chemical ionization quadrupole ion trap mass spectrometry of *O*-ethyl *S*-[2-(diisopropylamino)ethyl] methylphosphonothiolate (VX) and its degradation products 893(2000)393

The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry 894(2000)345

Simultaneous analysis of some amphetamine derivatives in urine by nonaqueous capillary electrophoresis coupled to electrospray ionization mass spectrometry 895(2000)111

Parameter optimization for the analysis of underivatized protein amino acids by liquid chromatography and ionspray tandem mass spectrometry 896(2000)253

#### Mass transfer

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Kinetic study of the mass transfer of bovine serum albumin in anion-exchange chromatography 866(2000)147

Erratum to: “Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles” [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

Adsorption behavior and prediction of the band profiles of the enantiomers of 3-chloro-1-phenyl-1-propanol. Influence of the mass transfer kinetics 877(2000)109

Determination of the lumped mass transfer rate coefficient by frontal analysis 890(2000)211

#### Mass transfer resistance

Influence of the column radial heterogeneity on the determination of single-component isotherms by the elution by characteristic point method 891(2000)217

#### Mathematical modelling

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Understanding of peak deterioration in hyphenated speciation systems due to gas–liquid separation in the hydride generation interface 855(1999)291

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Calculation of electrophoretic mobilities in water–organic modifier mixtures in capillary electrophoresis 868(2000)277

Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C<sub>8</sub> and reversed-phase C<sub>18</sub> types 869(2000)57

Study on the accuracy of the elution by characteristic point method for the determination of single component isotherms 872(2000)1

Scale-down of continuous protein purification by annular chromatography. Design parameters for the smallest unit 890(2000)53

Kinetics of solid-phase extraction and solid-phase microextraction in thin adsorbent layer with saturation sorption isotherm 873(2000)39

Erratum to: “Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles” [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

Adsorption equilibria and overloaded band profiles of basic drugs in a reversed-phase system 877(2000)1

Modelling of the pore flow in capillary electrochromatography 887(2000)199

Optimization methods in chromatography and capillary electrophoresis 892(2000)75

Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Peclet number in capillary electrochromatography from pore network theory 890(2000)61

**Matrix effects**

Does further clean-up reduce the matrix enhancement effect in gas chromatographic analysis of pesticide residues in food? 868(2000)51

Solid-phase extraction of phenols 885(2000)291

Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421

**Matrix solid-phase dispersion**

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Determination of abamectin in citrus fruits by liquid chromatography–electrospray ionization mass spectrometry 871(2000)57

Applications of matrix solid-phase dispersion in food analysis 880(2000)63

Matrix solid-phase dispersion 885(2000)115

**Matrix-assisted laser desorption ionization mass spectrometry**

see also Mass spectrometry

Protein identification platform utilizing micro dispensing technology interfaced to matrix-assisted laser desorption ionization time-of-flight mass spectrometry 886(2000)99

The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry 894(2000)345

Off-line size-exclusion chromatographic fractionation–matrix-assisted laser desorption ionization time-of-flight mass spectrometry for polymer characterization. Theoretical and experimental study 896(2000)19

**Meat**

Determination of folate vitamers in food and in Italian reference diet by high-performance liquid chromatography 855(1999)237

Comparison of different commercial solid-phase extraction cartridges used to extract heterocyclic amines from a lyophilised meat extract 880(2000)101

Determination of inulin in meat products by high-performance liquid chromatography with refractive index detection 881(2000)591

Determination of clopidol residues in chicken tissues by high-performance liquid chromatography–mass spectrometry 882(2000)85

Determination of four nitroimidazole residues in poultry meat by liquid chromatography–mass spectrometry 882(2000)89

Stability of penicillin antibiotic residues in meat during storage. Ampicillin 882(2000)135

**Membrane chromatography**

Quantitative fast fractionation of a pool of polyclonal antibodies by immunoaffinity membrane chromatography 852(1999)129

**Membrane dryers**

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

**Membrane extraction**

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

Membrane-based sample preparation coupled on-line to chromatography or electrophoresis 856(1999)55

Development of membrane extraction with a sorbent interface–micro gas chromatography system for field analysis 873(2000)13

**Membranes**

see also Hollow-fiber membranes

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Large-scale membrane adsorbers 852(1999)73

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

Membrane-based sample preparation coupled on-line to chromatography or electrophoresis 856(1999)55

Coupling of biological sample handling and capillary electrophoresis 856(1999)483

Electrophoretic transfer of proteins across polyacrylamide membranes 865(1999)145

New affinity nylon membrane used for adsorption of  $\gamma$ -globulin 867(2000)161

Analysis of pesticide residues in matrices with high lipid contents by membrane separation coupled on-line to a high-performance liquid chromatography system 869(2000)427

Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151

Erratum to: "Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes". [J. Chromatogr. A, 852 (1999) 59-71] 872(2000)323

Development of membrane extraction with a sorbent interface–micro gas chromatography system for field analysis 873(2000)13

Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83

Novel use of a dual-zone restricted access sorbent: normal-phase solid-phase extraction separation of methyl oleate from polynuclear aromatic hydrocarbons stemming from semi-permeable membrane devices 883(2000)143

High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43

Extraction of glyphosate by a supported liquid membrane technique 889(2000)93

Large-scale purification of antisense oligonucleotides by high-performance membrane adsorber chromatography 890(2000)179

#### **Metalloprotoporphyrin-silica stationary phases**

Shape-selective separation of polycyclic aromatic hydrocarbons on protoporphyrin-silica phases. Effect of surface porphyrin distribution on column efficiency 859(1999)121

#### ***Metarrhizium anisopliae***

Liquid chromatography–electrospray time-of-flight mass spectrometry for on-line accurate mass determination and identification of cyclodepsipeptides in a crude extract of the fungus *Metarrhizium anisopliae* 872(2000)85

#### **Method development**

Solid-phase extraction: method development, sorbents, and coupling with liquid chromatography 856(1999)3

Towards a quantitative definition of perfection in chromatographic analyses 868(2000)305

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215

Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs 883(2000)249

Contributions of theory to method development in solid-phase extraction 885(2000)17

Practitioner's guide to method development in thin-layer chromatography 892(2000)123

Analysis of 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-fluorophenyl)-*N*-(3-iodo-*E*-allyl)nortropine in rat plasma. I. Method development and validation by capillary electrophoresis 895(2000)87

Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. II. Minimizing column reproducibility problems 897(2000)51

#### **Methylation**

Esterification of decanoic acid during supercritical fluid extraction employing either methanol-modified carbon dioxide or a methanol trap 858(1999)201

Selection of methylation procedures for quantitation of short-chain glycerol-bound compounds formed during thermoxidation 863(1999)171

#### **Methylene increments**

Analysis of the equations for the temperature dependence of the retention index. I. Relation between equations 858(1999)65

#### **Methylene selectivity**

New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35

#### **Micellar electrokinetic chromatography**

see also Electrokinetic chromatography

Widening of the elution window in micellar electrokinetic chromatography with cationic surfactants. II. Cationic additives and modifiers of the electroosmotic flow 855(1999)669

Effects of organic modifiers on retention mechanism and selectivity in micellar electrokinetic capillary chromatography studied by linear solvation energy relationships 863(1999)69

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Enantiomer separation of drugs by micellar electrokinetic chromatography using chiral surfactants 875(2000)163

Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography 875(2000)179

Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis 875(2000)207

Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295

Statistical evaluation of linear solvation energy relationship models used to characterize chemical selectivity in micellar electrokinetic chromatography 886(2000)245

Open access capillary electrophoresis. A walk up capillary electrophoresis service for the synthetic chemist 888(2000)219

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry 894(2000)345

**Micellar electrokinetic chromatography, reversed migration**  
Reversed migration micellar electrokinetic chromatography with off-line and on-line concentration analysis of phenylurea herbicides 892(2000)187

**Micellar liquid chromatography**  
Micellar liquid chromatography for prediction of drug transport 870(2000)1

Use of a three-factor interpretive optimisation strategy in the development of an isocratic chromatographic procedure for the screening of diuretics in urine samples using micellar mobile phases 893(2000)321

**Micelles**  
pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

**Micelle-mediated extraction**  
Micelle-mediated extraction for concentrating hydrophobic organic compounds 879(2000)121

**Microchannels**  
Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45

Rapid, parallel separations of D1S80 alleles in a plastic microchannel chip 894(2000)203

**Microchips**  
see Chip technology

**Micro liquid chromatography**  
Recent developments in microcolumn liquid chromatography 856(1999)117

**Microdialysis**  
see also Dialysis

Monitoring of extracellular pyruvate, lactate, and ascorbic acid during cerebral ischemia: a microdialysis study in awake gerbils 870(2000)389

Determination of chlorogenic acid in rat blood by microdialysis coupled with microbore liquid chromatography and its application to pharmacokinetic studies 870(2000)443

On-line microdialysis–ion chromatographic determination of inorganic anions in olive-oil mill wastewater 882(2000)321

**Microemulsion electrokinetic chromatography**  
Microemulsion electrokinetic chromatography of proteins 853(1999)277

Microemulsion electrokinetic chromatography in suppressed electroosmotic flow environment. Separation of fat-soluble vitamins 876(2000)201

Background theory and applications of microemulsion electrokinetic chromatography 892(2000)171

Selectivity in microemulsion electrokinetic chromatography 897(2000)375

**Microgravimetry**  
Automatic microgravimetric determination of fats in milk products by use of supercritical fluid extraction with on-line piezoelectric detection 874(2000)265

**Microporous membrane liquid–liquid extraction**  
Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523

Determination of thiophanate-methyl and its metabolites at trace level in spiked natural water using the supported liquid membrane extraction and the microporous membrane liquid–liquid extraction techniques combined on-line with high-performance liquid chromatography 893(2000)123

**Microwave-assisted derivatization**  
Microwave-assisted derivatization of volatile carbonyl compounds with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine 891(2000)267

**Microwave-assisted extraction**  
Microwave-assisted extraction by fast sample preparation for the systematic analysis of additives in polyolefins by high-performance liquid chromatography 869(2000)285

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media 869(2000)515

Determination of sunscreen agents in cosmetic products using microwave-assisted extraction and liquid chromatography 876(2000)243

Dynamic microwave-assisted extraction 877(2000)141

Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annuum* L.) powders 889(2000)41

Experimental design of a microwave-assisted extraction–derivatization method for the analysis of methylmercury 889(2000)185

Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279

#### **Migration factor**

Capillary electrochromatography of proteins and peptides with a cationic acrylic monolith 887(2000)465

#### **Migration inhibitory factors**

Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor 852(1999)189

#### **Migration time correction**

Migration time correction for the analysis of derivatized amino acids and oligosaccharides by micellar capillary electrochromatography 869(2000)375

#### **Migration times**

How to achieve higher repeatability and reproducibility in capillary electrophoresis 853(1999)83

New method for standardization of electropherograms obtained in capillary zone electrophoresis 894(2000)11

#### **Milk**

Direct determination of small cations in proteinaceous samples using a flow injection–capillary electrophoresis system 857(1999)321

Analysis of major ovine milk proteins by reversed-phase high-performance liquid chromatography and flow injection analysis with electrospray ionization mass spectrometry 870(2000)371

Electrophoretic methods in the analysis of beverages 880(2000)243

Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides 881(2000)1

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

High-performance liquid chromatographic determination of Maillard compounds in store-brand and name-brand ultra-high-temperature-treated cows' milk 881(2000)599

Simultaneous determination of triclabendazole and its sulphoxide and sulphone metabolites in bovine milk by high-performance liquid chromatography 882(2000)99

Determination of cephalosporins in raw bovine milk by high-performance liquid chromatography 882(2000)145

Improved method for rapid determination of vitamin A in small samples of breast milk by high-performance liquid chromatography 898(2000)179

#### **Milk powder**

Determination of vitamins A and E in milk powder using supercritical fluid extraction for sample clean-up 874(2000)275

#### **Mineralization**

Application of solid-phase microextraction–gas chromatography–mass spectrometry to characterize intermediates in a joint solar–microbial process for total mineralization of Aroclor 1254 873(2000)53

#### **Mixed-mode chromatography**

Behavior of basic compounds in ion-exchange capillary electrochromatography with low-pH carrier electrolytes 884(2000)277

Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267

Comparative study of capillary electroendosmotic chromatography and electrically assisted gradient nano-liquid chromatography for the separation of peptides 894(2000)241

#### **Mobile phase composition**

see also Chiral selectors and Organic modifiers

Characterization of unstable ion-exchange chromatographic separation of proteins 852(1999)37

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

Empirical relationship between chiral selectivity and mobile phase modifier properties 852(1999)383

- Separation of selected peptides by capillary electroosmotic chromatography using 3  $\mu\text{m}$  reversed-phase bonded silica and mixed-mode phases 853(1999)263
- Application of experimental design for the characterisation of a novel elution system for high-capacity anion chromatography with suppressed conductivity detection 855(1999)91
- Comparative study of divalent metals and amines as silanol-blocking agents in reversed-phase liquid chromatography 855(1999)121
- Determination of inorganic anions by ion chromatography using a graphitized carbon column dynamically coated with cetyltrimethylammonium ions 855(1999)261
- Application and comparison of derivatized cellulose and amylose chiral stationary phases for the separation of enantiomers of pharmaceutical compounds by high-performance liquid chromatography 855(1999)411
- Isocratic separations of closely-related mono- and disaccharides by high-performance anion-exchange chromatography with pulsed amperometric detection using dilute alkaline spiked with barium acetate 855(1999)539
- Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1
- Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. III. Improving the accuracy of computer simulation 857(1999)41
- Retention and separation studies of cholesterol and bile acids using thermostated thin-layer chromatography 857(1999)255
- Quantitative analysis of manganese, chromium and molybdenum by ion-pair reversed-phase high-performance liquid chromatography with pre-column derivatization and UV-visible detection 857(1999)351
- Temperature-induced inversion of elution order in the enantioseparation of sotalol on a cellobiohydrolase I-based stationary phase 858(1999)33
- Optimization of glucosinolate separation by micellar electrokinetic capillary chromatography using a Doehlert's experimental design 864(1999)155
- Comparison of the performance of butanol and pentanol as modifiers in the micellar chromatographic determination of some phenethylamines 866(2000)35
- Effect of stationary phase hydrophobicity and mobile phase composition on the separation of carboxylic acids in ion chromatography 867(2000)131
- Modelling retention in liquid chromatography as a function of solvent composition and pH of the mobile phase 869(2000)27
- Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49
- Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329
- Some considerations concerning the composition of the mobile phase in capillary electrochromatography 869(2000)363
- Separation of potentially therapeutic peptide hormones by liquid chromatography. Optimisation of the composition and pH of the mobile phase 870(2000)335
- Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127
- Separation of isomeric naphthalenesulphonic acids by micro high-performance liquid chromatography with mobile phases containing cyclodextrin 871(2000)139
- Separation of some platinum group metal chelates with 8-hydroxyquinoline by various high-performance liquid chromatographic methods 871(2000)217
- Analysis of higher polyamide-6 oligomers on a silica-based reversed-phase column with a gradient of formic acid as compared with hexafluoroisopropanol 871(2000)259
- Capillary electrochromatography of basic compounds using octadecyl-silica stationary phases with an amine-containing mobile phase 872(2000)315
- Determination of mobile phase effect on single-component adsorption isotherm by use of numerical estimation 874(2000)1
- Chromatographic resolution of tryptophan enantiomers with L-Leu-L-Leu-L-Leu peptide. Effects of mobile phase composition and chromatographic support 874(2000)21
- Separation of purines and pyrimidines by normal-phase high-performance liquid chromatography using dimethyl sulfoxide in binary and ternary eluents 874(2000)45
- Separation of enantiomers by packed capillary electrochromatography on a cellulose-based stationary phase 875(2000)331
- Complementary mobile-phase optimisation for resolution enhancement in high-performance liquid chromatography 876(2000)17
- Characterization of C<sub>18</sub>-bonded liquid chromatographic stationary phases by Raman spectroscopy: The effect of mobile phase composition 877(2000)25

- Characterization of C<sub>18</sub>-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41
- High-performance liquid chromatography of non-polar retinoid isomers 881(2000)183
- Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317
- Use of nucleic acids in the mobile phase for the determination of ascorbic acid in foods by high-performance liquid chromatography with electrochemical detection 881(2000)327
- Influence of organic solvents in the mobile phase on the determination of carboxylic acids and inorganic anions in grape juice by ion chromatography 881(2000)387
- Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113
- Simultaneous ion-exclusion/cation-exchange chromatography of anions and cations in acid rain waters on a weakly acidic cation-exchange resin by elution with sulfosalicylic acid 884(2000)167
- Trace analysis of heavy metals in groundwater samples by ion chromatography with post-column reaction and ultraviolet-visible detection 884(2000)229
- Elution mechanism in electrostatic ion chromatography with histidine as an isoelectric ampholytic mobile phase 884(2000)287
- Interpretive optimisation strategy applied to the isocratic separation of phenols by reversed-phase liquid chromatography with acetonitrile–water and methanol–water mobile phases 886(2000)31
- Separation of acidic compounds by strong anion-exchange capillary electrochromatography 887(2000)223
- Comparison of aqueous and non-aqueous capillary electrochromatography for the separation of basic solutes 887(2000)233
- Chiral anion exchangers applied to capillary electrochromatography enantioseparation of oppositely charged chiral analytes: investigation of stationary and mobile phase parameters 887(2000)421
- Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63
- Development of a method for the analysis of nucleotides from the mantle tissue of the mussel *Mytilus galloprovincialis* 891(2000)99
- Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107
- Characterization of reversed-phase columns using the linear free energy relationship. III. Effect of the organic modifier and the mobile phase composition 893(2000)215
- Use of a three-factor interpretive optimisation strategy in the development of an isocratic chromatographic procedure for the screening of diuretics in urine samples using micellar mobile phases 893(2000)321
- Effects of aliphatic amines on capillary electrochromatographic performance of tricyclic antidepressants on octadecylsilica 895(2000)17
- Optimisation and characterisation of silica-based reversed-phase liquid chromatographic systems for the analysis of basic pharmaceuticals 897(2000)1
- Characterization of sulfobutyl ether- $\beta$ -cyclodextrins mixtures by anion-exchange chromatography using evaporative light scattering detection 897(2000)185
- Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23
- New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35
- Effect of mobile phase composition on the separation of propranolol enantiomers using a perphenylcarbamate  $\beta$ -cyclodextrin bonded chiral stationary phase 898(2000)53
- Mobile phase flow**  
On the possibility of shear-driven chromatography: a theoretical performance analysis 855(1999)57
- Mobile phase strength**  
Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3
- Molasses**  
Monitoring and control of industrial downstream processing of sugar beet molasses 882(2000)329
- Molecular descriptors**  
Modeling chromatographic parameters by a novel graph theoretical sub-structural approach 858(1999)187

**Molecular imprinting**

Uniform-sized molecularly imprinted polymer for (*S*)-ibuprofen. Retention properties in aqueous mobile phases 857(1999)117

Chiral and electrokinetic separation of amino acids using polypyrrole-coated adsorbents 868(2000)189

Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255

Chromatographic characterization of molecularly imprinted polymers binding the herbicide 2,4,5-trichlorophenoxyacetic acid 883(2000)119

Highly selective separations by capillary electrochromatography: molecular imprint polymer sorbents (Review) 887(2000)125

Affinity screening by packed capillary high-performance liquid chromatography using molecular imprinted sorbents. I. Demonstration of feasibility 888(2000)23

Influence of mobile phase composition and cross-linking density on the enantiomeric recognition properties of molecularly imprinted polymers 888(2000)63

New configurations and applications of molecularly imprinted polymers 889(2000)15

Solid-phase extraction with a dibutylmelamine-imprinted polymer as triazine herbicide-selective sorbent 889(2000)25

Use of molecularly imprinted solid-phase extraction for the selective clean-up of clenbuterol from calf urine 889(2000)105

Recognition of oxytocin and oxytocin-related peptides in aqueous media using a molecularly imprinted polymer synthesized by the epitope approach 889(2000)111

Optimisation of procedures for the extraction of structural analogues of propranolol with molecular imprinted polymers for sample preparation 889(2000)143

Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435

Application of molecularly imprinted polymers in supercritical fluid chromatography 897(2000)317

**Molecular interactions**

Characterization of reversed-phase columns using the linear free energy relationship. III. Effect of the organic modifier and the mobile phase composition 893(2000)215

**Molecular mass**

Number-average molecular mass determination of polymeric material by pyrolysis–gas chromatography 888(2000)209

Characterization of synthetic polyelectrolytes by capillary electrophoresis 894(2000)105

Off-line size-exclusion chromatographic fractionation–matrix-assisted laser desorption ionization time-of-flight mass spectrometry for polymer characterization. Theoretical and experimental study 896(2000)19

**Molecular mass distribution**

Molecular mass distribution of sodium alginate by high-performance size-exclusion chromatography 864(1999)199

**Molecular shape recognition**

Laterally attached liquid crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. II. Optimization of the molecular parameter of the polymer 897(2000)131

**Monascus**

Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295

Analysis of natural food pigments by capillary electrophoresis 880(2000)311

**Monolithic columns**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Separation of oligonucleotides on novel monolithic columns with ion-exchange functional surfaces 852(1999)297

Capillary columns with in situ formed porous monolithic packing for micro high-performance liquid chromatography and capillary electrochromatography 855(1999)273

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Erratum to: “Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles” [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

Design of the monolithic polymers used in capillary electrochromatography columns (Review) 887(2000)3

Steroid profiles determined by capillary electrochromatography, laser-induced fluorescence detection and electrospray–mass spectrometry 887(2000)379

Capillary electrochromatography of proteins and peptides with a cationic acrylic monolith 887(2000)465

**Monolithic sol–gel**

Characteristics of particle-loaded monolithic sol–gel columns for capillary electrochromatography. I. Structural, electrical and band-broadening properties 887(2000)277



**Morphological signals**

Adaptive numerical morphological filter for identifying chromatographic signals 855(1999)305

**Mould**

Determination of ergosterol on mouldy building materials using isotope dilution and gas chromatography–tandem mass spectrometry 898(2000)227

**Moving boundary system**

The invalidity of Kohlrausch' regulating function for Svensson's isoelectric focusing and stationary electrolysis at steady state 863(1999)219

**Moving chemical reaction boundary**

Experimental study on moving neutralization reaction boundary created with the strong reactive electrolytes of HCl and NaOH in agarose gel 891(2000)337

**Multiangle laser light scattering detection**

Size-exclusion chromatography–multiangle laser light scattering analysis of  $\beta$ -lactoglobulin and bovine serum albumin in aqueous solution with added salt 867(2000)105

**Multidimensional separation**

Integrated approach to the multidimensional analysis of complex biological samples by microseparation techniques. Analysis of glycoprotein factor associated with cancer cachexia 857(1999)183

**Multiple extraction**

Multiple solid-phase microextraction 878(2000)27

**Multiple headspace extraction**

Reduction of adsorption phenomena of volatile aldehydes and aromatic compounds for static headspace analysis of cellulose based packaging materials 897(2000)269

**Multiple linear regression analysis**

Calculation of Abraham solute descriptors from McReynolds gas chromatographic retention data 893(2000)339

**Multiple-step ligand injection**

Multiple-step ligand injection affinity capillary electrophoresis for determining binding constants of ligands to receptors 897(2000)339

**Multiresidue analysis**

Solid-phase extraction for multiresidue analysis of organic contaminants in water 885(2000)195

**Multivariate analysis**

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Characterization of commercial soybean products by conventional and perfusion reversed-phase high-performance liquid chromatography and multivariate analysis 881(2000)47

Robustness testing for a capillary electrophoresis method using the "short-end injection" technique 897(2000)329

**Multi-capillary columns**

High-speed gas chromatography: an overview of various concepts 856(1999)315

**Must**

Determination of indole-3-acetic acid, tryptophan and other indoles in must and wine by high-performance liquid chromatography with fluorescence detection 855(1999)227

Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxypyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93

Multiresidue method for the rapid determination of organophosphorus insecticides in grapes, must and wine 882(2000)213

Multiresidue method for the rapid determination – in grape, must and wine – of fungicides frequently used on vineyards 882(2000)221

**Mutagens**

Quantitative analysis of mutagenic heterocyclic aromatic amines in cooked meat using liquid chromatography–atmospheric pressure chemical ionisation tandem mass spectrometry 883(2000)89

***Mytilus edulis***

Characterization of odorant compounds of mussels (*Mytilus edulis*) according to their origin using gas chromatography–olfactometry and gas chromatography–mass spectrometry 896(2000)361

***Mytilus galloprovincialis***

Development of a method for the analysis of nucleotides from the mantle tissue of the mussel *Mytilus galloprovincialis* 891(2000)99

# N

**Nanoparticles**

Determination of oligonucleotide ISIS 2922 in nanoparticulate delivery systems by capillary zone electrophoresis 871(2000)311

**Nanoreactors**

Nanoliter-scale sample preparation methods directly coupled to polymethylmethacrylate-based microchips and gel-filled capillaries for the analysis of oligonucleotides 853(1999)107

**Nano liquid chromatography**

Separation of enantiomers by packed capillary electrochromatography on a cellulose-based stationary phase 875(2000)331

**Nebulizing interface**

Capillary electrophoresis–inductively coupled plasma mass spectrometry interface with minimised dead volume for high separation efficiency 891(2000)129

**Needle Search**

Application of several modified peak purity assays to real complex multicomponent mixtures by high-performance liquid chromatography with diode-array detection 855(1999)487

**Neem**

Rapid and sensitive analysis of azadirachtin and related triterpenoids from Neem (*Azadirachta indica*) by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 886(2000)89

**Net charge**

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

**Network model**

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Erratum to: “Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles” [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

**Neu differentiation factor**

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

**Neural networks, artificial**

Pressurized-flow anion-exchange capillary electrochromatography using a polymeric ion-exchange stationary phase 890(2000)337

Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227

**Non-aqueous capillary electrochromatography**

Enantiomeric separations by nonaqueous capillary electrophoresis 875(2000)277

Investigation of an enantioselective non-aqueous capillary electrochromatography system applied to the separation of chiral acids 875(2000)341

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Non-aqueous capillary electrophoresis 892(2000)155

Electrochromatography 892(2000)219

Enantiomer separations by nonaqueous capillary electrophoresis using octakis(2,3-diacetyl-6-sulfato)- $\gamma$ -cyclodextrin 892(2000)499

*tert*-Butylcarbamoylquinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique 892(2000)509

Simultaneous analysis of some amphetamine derivatives in urine by nonaqueous capillary electrophoresis coupled to electrospray ionization mass spectrometry 895(2000)111

**Non-aqueous capillary electrophoresis**

Application of non-aqueous capillary electrophoresis with electrochemical detection to the determination of nicotine in tobacco 853(1999)27

Indirect detection of organic acids in non-aqueous capillary electrophoresis 853(1999)171

Determination of quinolizidine alkaloids in traditional Chinese herbal drugs by nonaqueous capillary electrophoresis 857(1999)303

Efficiency studies in nonaqueous capillary electrophoresis 859(1999)203

Nonaqueous capillary electrophoresis equipped with amperometric detection for analysis of chlorinated phenolic compounds 864(1999)323

Influence of pH\*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63

Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

Development and robustness testing of a nonaqueous capillary electrophoresis method for the analysis of nonsteroidal anti-inflammatory drugs 874(2000)121

Chiral separation of unmodified amino acids with non-aqueous capillary electrophoresis based on the ligand-exchange principle 874(2000)285

Non-aqueous capillary electrophoretic enantioseparation of *N*-derivatized amino acids using cinchona alkaloids and derivatives as chiral counter-ions 875(2000)353

Enantiomeric separations by nonaqueous capillary electrophoresis 875(2000)277

Enantioseparation of anionic analytes by non-aqueous capillary electrophoresis using quinine and quinidine derivatives as chiral counter-ions 894(2000)63

Non-aqueous capillary electrophoresis of tamoxifen and its acid hydrolysis products 895(2000)81

Highly selective and efficient determination of US Environmental Protection Agency priority phenols employing solid-phase extraction and non-aqueous capillary electrophoresis 896(2000)95

#### **Non-crystalline phases**

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

#### **Non-equilibrium effects**

Enantiomers separation by simulated moving bed chromatography. Non-instantaneous equilibrium at the solid–fluid interface 865(1999)187

#### **Non-linear chromatography**

Enantiomers separation by simulated moving bed chromatography. Non-instantaneous equilibrium at the solid–fluid interface 865(1999)187

#### ***Nostoc***

Separation and identification of hydrocarbons and other volatile compounds from cultured blue-green alga *Nostoc* sp. by gas chromatography–mass spectrometry using serially coupled capillary columns with consecutive nonpolar and semipolar stationary phases 862(1999)221

#### ***Notopterygium forbesii***

Preparative isolation and purification of notopterol and isoimperatorin from *Notopterygium forbesii* boiss (Chinese traditional medicinal herb) by high-speed counter-current chromatography 883(2000)67

#### **Nuclear magnetic resonance spectrometry**

Liquid chromatography–nuclear magnetic resonance spectroscopy 856(1999)199

*n*-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133

New stationary phase for liquid chromatography with chemically bonded pinane ligand: synthesis and characterization by nuclear magnetic resonance and high-performance liquid chromatographic investigations 883(2000)27

Fullerenes separation with monomeric type C<sub>30</sub> stationary phase in high-performance liquid chromatography 883(2000)55

Selective deuterium exchange during superheated heavy water chromatography–nuclear magnetic resonance spectroscopy–mass spectrometry of sulfonamides 886(2000)289

## O

#### **Oakmoss**

Identification of atranorin and related potential allergens in oakmoss absolute by high-performance liquid chromatography–tandem mass spectrometry using negative ion atmospheric pressure chemical ionization 888(2000)103

#### **Octanol-water partition coefficients**

Determination of octanol–water partition coefficient for terpenoids using reversed-phase high-performance liquid chromatography 864(1999)221

#### **Off-flavor compounds**

Sample preparation for the analysis of flavors and off-flavors in foods 880(2000)3

#### **Oils**

see also Essential oils, Olive oil, Petroleum and Vegetable oils

Automated on-line membrane extraction liquid chromatographic determination of phenols in crude oils, gasolines and diesel fuels 852(1999)395

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

Preparative thin-layer chromatographic separation and subsequent gas chromatographic–mass spectrometric analysis of monoacylglycerols derived from butter oil by fungal degradation 855(1999)617

Separation, identification and determination of sanguinarine in argemone and other adulterated edible oils by reversed-phase high-performance liquid chromatography 863(1999)123

Analysis of volatile contaminants in vegetable oils by headspace solid-phase microextraction with Carboxen-based fibres 873(2000)79

Extraction of hydrocarbon contamination from soils using accelerated solvent extraction 874(2000)217

Analysis of olive and hazelnut oil mixtures by high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry of triacylglycerols and gas–liquid chromatography of non-saponifiable compounds (tocopherols and sterols) 881(2000)149

Rapid determination of vitamin E in vegetable oils by reversed-phase high-performance liquid chromatography 881(2000)251

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Polycyclic aromatic hydrocarbons in edible fats and oils: occurrence and analytical methods 882(2000)245

Determination of cholesterol oxides in processed food using high-performance liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 896(2000)321

Analysis of phenols in pyrolysis oils by gel permeation chromatography and multidimensional liquid chromatography 896(2000)343

#### **Oil–water partition coefficients**

Use of solid-phase microextraction for measuring oil–water partition coefficients and correlation with high-performance liquid chromatographic methods for lipophilicity 864(1999)183

#### **Olfactometry**

Characterization of odorant compounds of mussels (*Mytilus edulis*) according to their origin using gas chromatography–olfactometry and gas chromatography–mass spectrometry 896(2000)361

#### **Olive oil**

Chromatographic analysis of unsaponifiable compounds of olive oils and fat-containing foods 881(2000)105

Analysis of olive and hazelnut oil mixtures by high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry of triacylglycerols and gas–liquid chromatography of non-saponifiable compounds (tocopherols and sterols) 881(2000)149

Simultaneous determination of  $\alpha$ -tocopherol and  $\beta$ -carotene in olive oil by reversed-phase high-performance liquid chromatography 881(2000)255

On-line microdialysis–ion chromatographic determination of inorganic anions in olive-oil mill wastewater 882(2000)321

#### **Olives**

Liquid chromatography with electrospray ionisation mass spectrometric detection of phenolic compounds from *Olea europaea* 855(1999)529

#### **Open access capillary electrophoresis**

Open access capillary electrophoresis. A walk up capillary electrophoresis service for the synthetic chemist 888(2000)219

#### **Open tubular columns**

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

#### **Open-tubular capillary electrochromatography**

Influence of mobile phase composition on electroosmotic flow velocity, solute retention and column efficiency in open-tubular reversed-phase capillary electrochromatography 869(2000)329

Comparison of separation behavior of benzodiazepines in packed capillary electrochromatography and open-tubular capillary electrochromatography 887(2000)479

Peak shapes in open tubular ion-exchange capillary electrochromatography of inorganic anions 892(2000)303

#### **Ophthalmic solutions**

Simultaneous determination of eleven ingredients in ophthalmic solutions by cyclodextrin-modified micellar electrokinetic chromatography with tetrabutylammonium salt 888(2000)299

#### **Optical emission spectrometry**

Discoloration of ceramic hydroxyapatite used for protein chromatography 891(2000)93

#### **Optical rotation**

Case of enantiomer impurity identification by normal-phase chiral high-performance liquid chromatography with optical rotation and mass spectrometric detection 878(2000)35

#### **Optimization**

Optimization of throughput and desorbent consumption in simulated moving-bed chromatography for paclitaxel purification 855(1999)71

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. II. Two-run assay procedures 857(1999)21

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. III. Improving the accuracy of computer simulation 857(1999)41

- Computer-aided development of a high-performance liquid chromatographic method for the determination of hydroxyanthraquinone derivatives in Chinese herb medicine rhubarb 857(1999)167
- Retention and separation studies of cholesterol and bile acids using thermostated thin-layer chromatography 857(1999)255
- Optimization of the separation of mono- and dichloroanilines in ion interaction high-performance liquid chromatography 863(1999)1
- Simultaneous optimization of the analysis time and the concentration detectability in open-tubular liquid chromatography 867(2000)23
- Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247
- Use of optimization software to determine rugged analysis conditions in high-performance liquid chromatography 868(2000)13
- Computer simulation for the convenient optimization of isocratic reversed-phase liquid chromatographic separations by varying temperature and mobile phase strength 869(2000)3
- Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505
- Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75
- Enantiomeric separation of clenbuterol by transient isotachopheresis–capillary zone electrophoresis–UV detection. New optimization technique for transient isotachopheresis 871(2000)173
- Optimization of the separation of phenolic compounds by micellar electrokinetic capillary chromatography 871(2000)427
- Development of a method for the simultaneous determination of phosphoric and amino acid group containing pesticides by gas chromatography with mass-selective detection. Optimization of the derivatization procedure using an experimental design approach 872(2000)215
- Design of experiments for capillary electrophoretic enantioresolution of salbutamol using dermatan sulfate 875(2000)411
- Complementary mobile-phase optimisation for resolution enhancement in high-performance liquid chromatography 876(2000)17
- Optimization of ion-exchange displacement separations. I. Validation of an iterative scheme and its use as a methods development tool 876(2000)51
- Optimization of ion-exchange displacement separations. II. Comparison of displacement separations on various ion-exchange resins 876(2000)63
- Interpretive optimisation strategy applied to the isocratic separation of phenols by reversed-phase liquid chromatography with acetonitrile–water and methanol–water mobile phases 886(2000)31
- Capillary electroosmotic chromatography of peptides 887(2000)165
- Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197
- Optimization methods in chromatography and capillary electrophoresis 892(2000)75
- Use of a three-factor interpretive optimisation strategy in the development of an isocratic chromatographic procedure for the screening of diuretics in urine samples using micellar mobile phases 893(2000)321
- Optimization of parameters for the analysis of aromatic amines in finger-paints 896(2000)291
- Optimisation and characterisation of silica-based reversed-phase liquid chromatographic systems for the analysis of basic pharmaceuticals 897(2000)1
- Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. I. Optimizing selectivity and resolution 897(2000)37
- Optimization strategy and validation of one chromatographic method as approach to determine the phenolic compounds from different sources 897(2000)161
- Organic modifiers**  
see also Background electrolyte composition, Buffer composition and Mobile phase composition
- Effects of organic modifiers on retention mechanism and selectivity in micellar electrokinetic capillary chromatography studied by linear solvation energy relationships 863(1999)69
- Orthogonal array design**  
Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197
- Orthogonal projection**  
Application of several modified peak purity assays to real complex multicomponent mixtures by high-performance liquid chromatography with diode-array detection 855(1999)487
- Overlapping signals**  
Differential signal detection system for improved analysis of overlapping signals in liquid chromatography 859(1999)221

**Overload gradient elution**

Determination of mobile phase effect on single-component adsorption isotherm by use of numerical estimation 874(2000)1

**Overpressured layer chromatography**

Determination of aflatoxins in food by overpressured-layer chromatography 882(2000)11

**Oxidation**

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum–rhodium alloy catalysts 852(1999)507

**Ozonation**

Tandem solid-phase extraction of atrazine ozonation products in water 866(2000)195

**Ozone scrubbing material**

Investigation of membrane dryers and evaluation of a new ozone scrubbing material for the sampling of organosulphur compounds in air 852(1999)589

**P****Packed capillary columns**

see also Capillary columns

Determination of low levels of an antioxidant in polyolefins by large-volume injection temperature-programmed packed capillary liquid chromatography 864(1999)103

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

**Packed column capillary electrochromatography**

Packing columns for capillary electrochromatography (Review) 887(2000)43

Comparison of separation behavior of benzodiazepines in packed capillary electrochromatography and open-tubular capillary electrochromatography 887(2000)479

**Packed columns**

High-speed gas chromatography: an overview of various concepts 856(1999)315

**Packing methods**

Evaluation of the secondary consolidation of columns for liquid chromatography by ultrasonic irradiation 878(2000)153

Comparison of different packing methods for capillary electrochromatography columns 887(2000)307

Packing columns for capillary electrochromatography (Review) 887(2000)43

***Panax ginseng***

Liquid chromatography–electrospray mass spectrometric identification of ginsenosides in *Panax ginseng* roots 854(1999)69

High-performance thin-layer chromatographic determination of six major ginsenosides in *Panax ginseng* 868(2000)269

**Paper**

Determination of soluble anions and cations from waters of pulp and paper mills with on-line coupled capillary electrophoresis 895(2000)189

**Partial-filling capillary electrophoresis**

Simultaneous stereoselective analysis of tramadol and its main phase I metabolites by on-line capillary zone electrophoresis–electrospray ionization mass spectrometry 868(2000)295

Separation of enantiomers by capillary electrophoresis–mass spectrometry employing a partial filling technique with a chiral crown ether 875(2000)323

*tert.*-Butylcarbamoylquinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique 892(2000)509

**Partial least squares calibration**

Determination of naphthalenesulfonates in water by on-line ion-pair solid-phase extraction and ion-pair liquid chromatography with fast-scanning fluorescence detection 890(2000)289

**Particle analysis**

Capillary zone electrophoresis of a recombinant adenovirus 895(2000)329

**Particle diameter**

Towards the ultimate minimum particle diameter of silica packings in capillary electrochromatography 887(2000)339

**Particle size distribution**

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

**Partition–adsorption systems**

Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C<sub>8</sub> and reversed-phase C<sub>18</sub> types 869(2000)57

Separation of purines and pyrimidines by normal-phase high-performance liquid chromatography using dimethyl sulfoxide in binary and ternary eluents 874(2000)45

**Partitioning**

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

Purification of protein and recycling of polymers in a new aqueous two-phase system using two thermoseparating polymers 864(1999)31

Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69

Use of solid-phase microextraction for measuring oil–water partition coefficients and correlation with high-performance liquid chromatographic methods for lipophilicity 864(1999)183

Determination of octanol–water partition coefficient for terpenoids using reversed-phase high-performance liquid chromatography 864(1999)221

Characterization of cephalosporin transfer between aqueous and colloidal phases by micellar electrokinetic chromatography 871(2000)439

Characterization of partition and thermodynamic properties of cephalosporins using micellar electrokinetic chromatography in glycodeoxycholic acid solution 873(2000)237

**Peak areas**

How to achieve higher repeatability and reproducibility in capillary electrophoresis 853(1999)83

Comparison of the repeatability of quantitative data measured in high-performance liquid chromatography with UV and atmospheric pressure chemical ionization mass spectrometric detection 873(2000)163

**Peak capacity**

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

**Peak dispersion**

Liquid adsorption chromatography of polyethers: experiments and simulation 890(2000)195

**Peak overlap**

Differential signal detection system for improved analysis of overlapping signals in liquid chromatography 859(1999)221

**Peak purity**

Application of several modified peak purity assays to real complex multicomponent mixtures by high-performance liquid chromatography with diode-array detection 855(1999)487

Interpretive optimisation strategy applied to the isocratic separation of phenols by reversed-phase liquid chromatography with acetonitrile–water and methanol–water mobile phases 886(2000)31

**Peak resolution**

Complementary mobile-phase optimisation for resolution enhancement in high-performance liquid chromatography 876(2000)17

**Peak shape**

see also Band broadening and Band profiles

Analysis of anions in aqueous samples by ion chromatography and capillary electrophoresis. A comparative study of peak modeling and validation criteria 852(1999)487

Influence of column radial heterogeneity on peak fronting in linear chromatography 857(1999)69

Determination of the thermodynamic contribution to peak asymmetry of basic solutes in reversed-phase liquid chromatography 865(1999)97

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

Peak shapes in open tubular ion-exchange capillary electrochromatography of inorganic anions 892(2000)303

Influence of temperature on the behaviour of small linear peptides in capillary electrochromatography 893(2000)401

Life-time studies with capillary electrochromatography columns operated under different conditions 894(2000)35

Capillary zone electrophoresis in phosphate buffer – known or unknown? 894(2000)89

**Pecllet number**

Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Pecllet number in capillary electrochromatography from pore network theory 890(2000)61

**Pellicillar expanded bed matrix**

Pellicular expanded bed matrix suitable for high flow rates 878(2000)17

**Pencil lead**

Assessment of polycrystalline graphites as sorbents for solid-phase microextraction of nonionic surfactants 888(2000)35

***Penicillium roquefortii***

Preparative thin-layer chromatographic separation and subsequent gas chromatographic–mass spectrometric analysis of monoacylglycerols derived from butter oil by fungal degradation 855(1999)617

**Penicillium spp.**

Chromatographic determination of the mycotoxin patulin in fruit and fruit juices 882(2000)17

**Peptide mapping**

Peptide map procedure using immobilized protease cartridges in tandem for disulfide linkage identification of *neu* differentiation factor epidermal growth factor domain 867(2000)151

**Perfusion chromatography**

Perfusion chromatography: an emergent technique for the analysis of food proteins 880(2000)169

Determination by perfusion reversed-phase high-performance liquid chromatography of the soybean protein content of commercial soybean products prepared directly from whole soybeans 881(2000)37

Characterization of commercial soybean products by conventional and perfusion reversed-phase high-performance liquid chromatography and multivariate analysis 881(2000)47

**Petroleum**

see also Oils

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection 867(2000)187

Comprehensive two-dimensional gas chromatography: a hyphenated method with strong coupling between the two dimensions 856(1999)331

**Pharmaceutical analysis**

Immobilized biomembrane chromatography of highly lipophilic drugs 852(1999)313

Capillary electrophoretic study of cisplatin interaction with nucleoside monophosphates, di- and trinucleotides 852(1999)337

Elimination of adsorption effects in the analysis of water-soluble vitamins in pharmaceutical formulations by capillary electrophoresis 853(1999)391

Determination of homotaurine as impurity in calcium acamprostate by capillary zone electrophoresis 853(1999)421

Resolution of newly synthesized racemic dihydropyridines with different chiral selectors by means of capillary electrophoresis 853(1999)455

Separation of antipsychotic drugs (clozapine, loxapine) and their metabolites by capillary zone electrophoresis 853(1999)461

Capillary electrophoretic separation of *N*-acetylcysteine and its impurities as a method for quality control of pharmaceuticals 853(1999)479

Analysis of epinephrine from fifteen different dental anesthetic formulations by capillary electrophoresis 853(1999)535

Determination of the impurity profile of ethyl-2-(4-((5*R*)-3-[4-(methoxycarbonylaminomethyl)-phenyl]-2-oxo-5-oxazolidinylmethyl)-1-piperazinyl) acetate, citrate, a highly active platelet aggregation inhibitor, by liquid chromatography–mass spectrometry 854(1999)13

Comparison of turbulent-flow chromatography with automated solid-phase extraction in 96-well plates and liquid–liquid extraction used as plasma sample preparation techniques for liquid chromatography–tandem mass spectrometry 854(1999)23

Investigation on the iodination reaction of methylene blue by liquid chromatography–mass spectrometry with ionspray ionisation 854(1999)57

Application and comparison of derivatized cellulose and amylose chiral stationary phases for the separation of enantiomers of pharmaceutical compounds by high-performance liquid chromatography 855(1999)411

Separation and purification of isoflavones from *Pueraria lobata* by high-speed counter-current chromatography 855(1999)709

Determination of quinolizidine alkaloids in traditional Chinese herbal drugs by nonaqueous capillary electrophoresis 857(1999)303

Micellar electrokinetic capillary chromatography of macrolide antibiotics. Separation of tylosin, erythromycin and their related substances 857(1999)313

Headspace gas chromatography method for the analysis of volatile impurities in a hormone replacement transdermal patch 859(1999)113

High-performance thin-layer chromatography method for monitoring norfloxacin residues on pharmaceutical equipment surfaces 862(1999)209

Migration behavior and separation of active components in *Glycyrrhiza uralensis* Fisch and its commercial extract by micellar electrokinetic capillary chromatography 863(1999)105

Preparative chromatographic resolution of enantiomers using polar organic solvents with polysaccharide chiral stationary phases 865(1999)211

Direct, preparative enantioselective chromatography of propranolol hydrochloride and thioridazine hydrochloride using carbon dioxide-based mobile phases 865(1999)227

Analysis of coptisine, berberine and palmatine in adulterated Chinese medicine by capillary electrophoresis–electrospray ion trap mass spectrometry 866(2000)273



- Capillary electrophoretic determination of sanguinarine and chelerythrine in plant extracts and pharmaceutical preparations 866(2000)293
- Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1
- Micellar liquid chromatography for prediction of drug transport 870(2000)1
- Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39
- Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45
- Validation of the removal of acetylsalicylic acid. Recovery and determination of residues on various surfaces by high performance liquid chromatographic 870(2000)69
- Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97
- Determination by high-performance liquid chromatography of ketoprofen in vitro in rat skin permeation samples 870(2000)143
- Automated liquid membrane extraction for high-performance liquid chromatography of Ropivacaine metabolites in urine 870(2000)151
- Development of a sensitive method for the determination of ganciclovir by reversed-phase high-performance liquid chromatography 870(2000)159
- Stability-indicating high-performance liquid chromatographic assay and analysis of decomposition products of 2-{4-[(7-chloro-2-quinoxalinyloxy]phenoxy}propanoic acid 870(2000)179
- Determination of eight water- and fat-soluble vitamins in multi-vitamin pharmaceutical formulations by high-performance liquid chromatography 870(2000)207
- Determination of vitamin B<sub>12</sub> in multivitamin tablets by multimode high-performance liquid chromatography 870(2000)217
- Separation of erythromycin and related substances on base-deactivated reversed-phase silica gel columns 870(2000)227
- Liquid chromatography of polymyxin B sulphate 870(2000)237
- Capillary zone electrophoretic separation and determination of imidazolic antifungal drugs 871(2000)399
- Ion-exclusion controlled size-exclusion chromatography of methacrylic acid–methyl methacrylate copolymers 872(2000)91
- Development and robustness testing of a nonaqueous capillary electrophoresis method for the analysis of nonsteroidal anti-inflammatory drugs 874(2000)121
- Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography 875(2000)255
- Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated  $\beta$ -cyclodextrin 875(2000)389
- Separation of brompheniramine enantiomers by capillary electrophoresis and study of chiral recognition mechanisms of cyclodextrins using NMR-spectroscopy, UV spectrometry, electrospray ionization mass spectrometry and X-ray crystallography 875(2000)471
- Quantitation and characterization of phospholipids in pharmaceutical formulations by liquid chromatography–mass spectrometry 876(2000)97
- Analysis of baclofen by capillary electrophoresis with laser-induced fluorescence detection 877(2000)233
- Analysis of tetramethylpyrazine in *Ephedrae herba* by gas chromatography–mass spectrometry and high-performance liquid chromatography 878(2000)147
- Analysis of polymyxin B sulfate by capillary zone electrophoresis with cyclodextrin as additive. Method development and validation 879(2000)211
- Analytical techniques for determining biotin 881(2000)331
- Preparative isolation and purification of notopterol and isoimperatorin from *Notopterygium forbesii* boiss (Chinese traditional medicinal herb) by high-speed counter-current chromatography 883(2000)67
- Validation of a capillary electrophoresis method for determination of 5-aminolevulinic acid and degradation products 883(2000)285
- Simultaneous determination of eleven ingredients in ophthalmic solutions by cyclodextrin-modified micellar electrokinetic chromatography with tetrabutylammonium salt 888(2000)299
- Liquid chromatography–electrospray mass spectrometry of multicomponent peptide mixtures. Characterization of a mixture from the synthesis of the hormone goserelin 889(2000)119

Determination of vitamin B<sub>12</sub> in multivitamin tablets and fermentation medium by high-performance liquid chromatography with fluorescence detection 891(2000)243

Background theory and applications of microemulsion electrokinetic chromatography 892(2000)171

Development and validation of a capillary zone electrophoretic method for the determination of bisphosphonate and phosphonate impurities in clodronate 893(2000)411

Effects of aliphatic amines on capillary electrochromatographic performance of tricyclic antidepressants on octadecylsilica 895(2000)17

Adaptation of capillary electrophoresis to the determination of selected cephalosporins for injection 895(2000)27

Determination at ppb level of an anti-human immunodeficiency virus nucleoside drug by capillary electrophoresis–electrospray ionization tandem mass spectrometry 895(2000)101

Capillary electrophoresis with laser-induced fluorescence in clinical drug development. Routine application and future aspects 895(2000)147

Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191

Use of chiral liquid chromatography–tandem mass spectrometry to investigate the metabolism of racemic cholecystokinin-B antagonists 896(2000)217

#### **Phase transition**

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

#### **pH effects**

pH effects on micelle–water partitioning determined by micellar electrokinetic chromatography 852(1999)559

Influence of pH\*-value of methanolic electrolytes on electroosmotic flow in hydrophilic coated capillaries 868(2000)63

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by  $\gamma$ -radiation on titanium-grafted silica 869(2000)137

Influence of pH and p*K*<sub>a</sub> values on electrophoretic behaviour of quinolones in aqueous and hydro-organic media 871(2000)381

Effect of ionic strength, pH and polymer concentration on the separation of DNA fragments in the presence of electroosmotic flow 894(2000)219

Influence of the structure of cyclodextrins and amino acid sequence of dipeptides and tripeptides on the pH-dependent reversal of the migration order in capillary electrophoresis 894(2000)267

Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279

#### **Phenyl selectivity**

New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35

#### **Phenylaminopropyl silica**

Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23

#### **Pheochromocytoma cells**

Determination of catecholamines in pheochromocytoma cell (PC-12) culture medium by microdialysis–microbore liquid chromatography 870(2000)405

#### **pH gradients**

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. I. Effect of buffer composition on the gradient shape 890(2000)25

High-performance chromatofocusing using linear and concave pH gradients formed with simple buffer mixtures. II. Separation of proteins 890(2000)37

#### **Phosphate buffers**

Capillary zone electrophoresis in phosphate buffer – known or unknown? 894(2000)89

#### **Photochemical derivatization**

Analysis of barbiturates by micro-high-performance liquid chromatography with post-column photochemical derivatization 870(2000)39

#### **Photosystem II**

Resolution and identification of the protein components of the photosystem II antenna system of higher plants by reversed-phase liquid chromatography with electrospray-mass spectrometric detection 886(2000)111

#### **$\pi$ -Acceptors**

Convenient synthesis of  $\pi$ -acceptor chiral stationary phases for high-performance liquid chromatography from halogen-substituted 3,5-dinitrobenzoylamides 859(1999)143

#### **$\pi$ – $\pi$ Interactions**

*n*-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133

**Piezoelectric detection**

Automatic microgravimetric determination of fats in milk products by use of supercritical fluid extraction with on-line piezoelectric detection 874(2000)265

**Pitch**

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

**Planar chromatography**

see also Thin-layer chromatography

Planar chromatography at the turn of the century 856(1999)399

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

**Plant materials**

Liquid chromatography–electrospray mass spectrometric identification of ginsenosides in *Panax ginseng* roots 854(1999)69

Fast, sensitive and selective liquid chromatographic–tandem mass spectrometric determination of tumor-promoting diterpene esters 855(1999)563

Separation and purification of isoflavones from *Pueraria lobata* by high-speed counter-current chromatography 855(1999)709

Analysis of  $\beta$ -*N*-oxalyl-L- $\alpha$ , $\beta$ -diaminopropionic acid and homoarginine in *Lathyrus sativus* by capillary zone electrophoresis 857(1999)295

Preparative isolation and purification of hydroxyanthraquinones from *Rheum officinale* Baill by high-speed counter-current chromatography using pH-modulated stepwise elution 858(1999)103

Analysis of taxol and major taxoids in Himalayan yew, *Taxus wallichiana* 858(1999)239

Gas chromatographic–mass spectrometric determination of sulfolane in wetland vegetation exposed to sour gas-contaminated groundwater 859(1999)69

Capillary electrophoretic determination of sanguinarine and chelerythrine in plant extracts and pharmaceutical preparations 866(2000)293

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Enantioseparation of atropine by capillary electrophoresis using sulfated  $\beta$ -cyclodextrin: application to a plant extract 868(2000)285

Isolation and characterization of methoxylated flavones in the flowers of *Primula veris* by liquid chromatography and mass spectrometry 870(2000)453

Liquid chromatography–electrospray ionization mass spectrometry study of the flavonoids of the roots of *Astragalus mongholicus* and *A. membranaceus* 876(2000)87

Improved liquid chromatographic method for the analysis of photosynthetic pigments of higher plants 876(2000)111

On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic–diode array detection–mass spectrometric techniques 880(2000)203

Solid-phase extraction of quaternary ammonium herbicides 885(2000)251

Determination of glucosinolates in rapeseed and *Thlaspi caerulescens* plants by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 889(2000)75

Evaluation of extraction procedures for the ion chromatographic determination of arsenic species in plant materials 889(2000)83

Retention index database for identification of general green leaf volatiles in plants by coupled capillary gas chromatography–mass spectrometry 890(2000)313

Influence of fibre coating in headspace solid-phase microextraction–gas chromatographic analysis of aromatic and medicinal plants 892(2000)469

**Platinum–rhodium catalysts**

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum–rhodium alloy catalysts 852(1999)507

**Polar functional groups**

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

**Polyacrylamide gels**

Mobility and activation energy of single-stranded DNA in denaturing cross-linked polyacrylamide slab gels 894(2000)231

**Polyacrylamide membranes**

Electrophoretic transfer of proteins across polyacrylamide membranes 865(1999)145

**Polydimethylsiloxane fibres**

Evidence for selectivity of absorption of volatile organic compounds by a polydimethylsiloxane solid-phase microextraction fibre 885(2000)457

**Polydimethylsiloxane sorbents**

On the performance and inertness of different materials used for the enrichment of sulfur compounds from air and gaseous samples 864(1999)345

**Polydispersity**

Off-line size-exclusion chromatographic fractionation–matrix-assisted laser desorption ionization time-of-flight mass spectrometry for polymer characterization. Theoretical and experimental study 896(2000)19

**Polyelectrolyte sorbents**

Ion chromatographic selectivity of polyelectrolyte sorbents based on some aliphatic and aromatic ionenes 884(2000)31

**Polyelectrolytes**

Dissociation of polyvalent electrolytes 853(1999)35

Characterization of synthetic polyelectrolytes by capillary electrophoresis 894(2000)105

**Polyethyleneimine pseudostationary phase**

Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudostationary phase. Influence of methanol and acetonitrile on separation selectivity 853(1999)121

**Poly(glycidyl methacrylate–triallyl isocyanurate–divinylbenzene) resins**

Macroporous poly(glycidyl methacrylate–triallyl isocyanurate–divinylbenzene) matrix as an anion-exchange resin for protein adsorption 855(1999)129

**Polymer–agarose matrices**

Ultra-thin-layer agarose gel electrophoresis. II. Separation of DNA fragments on composite agarose–linear polymer matrices 871(2000)289

**Polymer-based sorbents**

Recent developments in polymer-based sorbents for solid-phase extraction 885(2000)51

**Polymer-encapsulated silica**

Reversed-phase liquid chromatography of proteins and peptides using multimodal copolymer-encapsulated silica 866(2000)1

**Polymeric adsorbents**

Solid-phase extraction of polar hydrophilic aromatic sulfonates followed by capillary zone electrophoresis–UV absorbance detection and ion-pair liquid chromatography–diode array UV detection and electrospray mass spectrometry 890(2000)225

**Polymeric gels**

Protein diffusion in charged polyacrylamide gels. Visualization and analysis 865(1999)155

Protein adsorption on novel acrylamido-based polymeric ion-exchangers. I. morphology and equilibrium adsorption 897(2000)65

Protein adsorption on novel acrylamido-based polymeric ion exchangers. II. Adsorption rates and column behavior 897(2000)81

**Polymer microfluid devices**

Measurement of electroosmotic flow in plastic imprinted microfluid devices and the effect of protein adsorption on flow rate 857(1999)275

**Polymer stationary phases**

Preparation of uniformly sized polymeric separation media potentially suitable for small-scale high-performance liquid chromatography and/or capillary electrochromatography 853(1999)11

**Poly(methyloctylsiloxane) stationary phases**

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by  $\gamma$ -radiation on titanium-grafted silica 869(2000)137

**Poly(styrene–divinylbenzene) stationary phases**

Rapid reversed-phase separation of proteins and peptides using optimized ‘moulded’ monolithic poly(styrene–*co*-divinylbenzene) columns 865(1999)169

Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23

**Polystyrene resins**

Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins 869(2000)171

**Polystyrene stationary phases**

Application of neutral hydrophobic hypercrosslinked polystyrene to the separation of inorganic anions by ion chromatography 884(2000)41

**Pore connectivity**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Erratum to: “Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles” [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

**Pore diffusion**

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Erratum to: "Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles" [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

#### **Pore diffusion coefficient**

Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Peclet number in capillary electrochromatography from pore network theory 890(2000)61

#### **Pore flow**

Modelling of the pore flow in capillary electrochromatography 887(2000)199

#### **Pore network model**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Determination of the intraparticle electroosmotic volumetric flow-rate, velocity and Peclet number in capillary electrochromatography from pore network theory 890(2000)61

#### **Pore size distribution**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Modeling and simulation of the dynamic behavior of monoliths. Effects of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles 865(1999)13

Erratum to: "Modeling and simulation of the dynamic behavior of monoliths. Effect of pore structure from pore network model analysis and comparison with columns packed with porous spherical particles" [J. Chromatogr. A, 865 (1999) 13–25] 873(2000)301

Pore size distributions of cation-exchange adsorbents determined by inverse size-exclusion chromatography 883(2000)39

#### **Poroshell packings**

Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3

#### **Porous copolymers**

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

#### **Porous graphitized carbon**

see also Graphitized carbon black

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245

Graphitized carbons for solid-phase extraction 885(2000)73

#### **Porous particles**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3

Synthesis of spherical porous silicas in the micron and submicron size range: challenges and opportunities for miniaturized high-resolution chromatographic and electrokinetic separations 892(2000)47

#### **Porous-layer open-tubular columns**

Capillary electrochromatography of proteins and peptides with porous-layer open-tubular columns 858(1999)91

#### **Porphyrin-silica stationary phases**

Shape-selective separation of polycyclic aromatic hydrocarbons on protoporphyrin-silica phases. Effect of surface porphyrin distribution on column efficiency 859(1999)121

#### ***Portulaca oleracea***

Fatty acids and  $\beta$ -carotene in Australian purslane (*Portulaca oleracea*) varieties 893(2000)207

#### **Positional isomers**

Equilibrium analysis of reactions between aromatic anions and nonionic surfactant micelles by capillary zone electrophoresis 853(1999)55

Separation of stereoisomers in a simulated moving bed-supercritical fluid chromatography plant 865(1999)175

Chromatographic separation of 3,4-difluorophenylacetic acid and its positional isomers using five different techniques 866(2000)281

Study of the *cis-trans* isomerization of enalapril by reversed-phase liquid chromatography 871(2000)189

Insoluble  $\beta$ -cyclodextrin polymer for capillary gas chromatographic separation of enantiomers and isomers 883(2000)137

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

#### **Preconcentration**

see also Clean-up methods, Extraction methods, Sample handling, Sample preparation and Special Issue Index

- Ion exchange-based preconcentration for the determination of anions by capillary electrophoresis 868(2000)135
- Poly(allylamine) beads as selective sorbent for preconcentration of formaldehyde and acetaldehyde in high-performance liquid chromatographic analysis 886(2000)83
- Preferential solvation**  
Dissociation constants of phenols in methanol–water mixtures 867(2000)45
- Preparative chromatography**  
see also Special Issue Index
- Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69
- Influence of column design on process-scale ion-exchange chromatography 865(1999)3
- Development of an high-performance liquid chromatographic simulated moving bed separation from an industrial perspective 865(1999)35
- Expanded-bed chromatography in primary protein purification (Review) 865(1999)129
- Separation of stereoisomers in a simulated moving bed-supercritical fluid chromatography plant 865(1999)175
- Enantiomer separation of  $\alpha$ -ionone using gas chromatography with cyclodextrin derivatives as chiral stationary phases 865(1999)201
- Preparative chromatographic resolution of enantiomers using polar organic solvents with polysaccharide chiral stationary phases 865(1999)211
- Direct, preparative enantioselective chromatography of propranolol hydrochloride and thioridazine hydrochloride using carbon dioxide-based mobile phases 865(1999)227
- Design of a simulated moving bed unit for sucrose–betaine separations 872(2000)23
- Development of analytical and preparative chromatographic separations of novel growth hormone secretagogue compounds 872(2000)75
- Determination of mobile phase effect on single-component adsorption isotherm by use of numerical estimation 874(2000)1
- Design and optimisation of a simulated moving bed unit: role of deviations from equilibrium theory 876(2000)3
- Isolation and preparative purification of microcystin variants 883(2000)103
- On-line monitoring of enantiomer concentration in chiral simulated moving bed chromatography 888(2000)73
- Suspended bed chromatography, a new approach in downstream processing 890(2000)45
- Discoloration of ceramic hydroxyapatite used for protein chromatography 891(2000)93
- Development of simultaneous purification methodology for multiple synthetic peptides by reversed-phase sample displacement chromatography 893(2000)81
- Enantioseparation of 1-phenyl-1-propanol by simulated moving bed under linear and nonlinear conditions 893(2000)307
- Comparison of chromatographic ion-exchange resins. I. Strong anion-exchange resins 897(2000)99
- Preparative electrophoresis**  
Electrophoretic transfer of proteins across polyacrylamide membranes 865(1999)145
- Chiral capillary electrophoresis as predictor for separation of drug enantiomers in continuous flow zone electrophoresis 895(2000)51
- Pressure dependent retention**  
Pressure-controlled high-performance liquid chromatographic study on the influence of rim chemistry on partial molar volume differences between free and complexed cyclodextrins 863(1999)127
- Pressure-induced retention**  
Pressure-induced retention of the lysozyme on reversed-phase liquid chromatography 891(2000)207
- Pressurised liquid extraction of ketones of polycyclic aromatic hydrocarbons from soil 893(2000)201
- Pressure transducer**  
Using a differential refractive index detector as a pressure transducer for online viscometry in exclusion chromatography 878(2000)1
- Pressurized flow**  
Pressurized-flow anion-exchange capillary electrochromatography using a polymeric ion-exchange stationary phase 890(2000)337
- Pressurized gradient electrochromatography**  
Pressurized gradient capillary electrochromatographic separation of eighteen amino acid derivatives 894(2000)337
- Pressurized hot water extraction**  
Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

**Pressurized liquid extraction**

Application of pressurized liquid extraction followed by gas chromatography–mass spectrometry to determine 4-nonylphenols in sediments 866(2000)79

Selective extraction of hydrocarbons, phosphonates and phosphonic acids from soils by successive supercritical fluid and pressurized liquid extractions 866(2000)231

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

Pressurised liquid extraction of pentachlorophenol from dyestuffs and cotton 873(2000)287

Extraction of hydrocarbon contamination from soils using accelerated solvent extraction 874(2000)217

Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl)ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane] from aged contaminated soil 874(2000)257

Pressurised liquid extraction of polycyclic aromatic hydrocarbons from contaminated soils 883(2000)151

Evaluation of extraction procedures for the ion chromatographic determination of arsenic species in plant materials 889(2000)83

Miniaturised pressurised liquid extraction of polycyclic aromatic hydrocarbons from soil and sediment with subsequent large-volume injection–gas chromatography 891(2000)275

Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421

***Primulae veris***

Isolation and characterization of methoxylated flavones in the flowers of *Primula veris* by liquid chromatography and mass spectrometry 870(2000)453

**Principal component analysis**

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Exploratory chemometric analysis of the classification of pharmaceutical substances based on chromatographic data 897(2000)23

**Process control**

Determination of soluble anions and cations from waters of pulp and paper mills with on-line coupled capillary electrophoresis 895(2000)189

**Process monitoring**

Monitoring and control of industrial downstream processing of sugar beet molasses 882(2000)329

Analytical techniques used for monitoring the biodegradation of fluorinated compounds in waste streams from pharmaceutical production 889(2000)177

**Programmed-temperature ...**

see also Temperature programming

**Programmed-temperature vaporizer**

Programmed temperature vaporiser-based injection in capillary gas chromatography 856(1999)259

Coupling device for desorption of drugs from solid-phase extraction-pipette tips and on-line gas chromatographic analysis 858(1999)117

Trace analysis of pesticide residues in water by high-speed narrow-bore capillary gas chromatography–mass spectrometry with programmable temperature vaporizer 874(2000)81

Introduction of large volumes of water-containing samples into a gas chromatograph. Improved retention of volatile solutes through the swing system 876(2000)135

Efficiency through combining high-performance liquid chromatography and high resolution gas chromatography: progress 1995–1999 892(2000)407

**Protein adsorption**

Protein adsorption to the bare silica wall in capillary electrophoresis. Quantitative study on the chemical composition of the background electrolyte for minimising the phenomenon 868(2000)85

Microcalorimetric studies of interactions between proteins and hydrophobic ligands in hydrophobic interaction chromatography: effects of ligand chain length, density and the amount of bound protein 872(2000)37

**Protein coiled cells**

Trapping the monomeric  $\alpha$ -helical state during unfolding of coiled-coils by reversed-phase liquid chromatography 890(2000)81

**Protein folding**

Considerations of sample application and elution during size-exclusion chromatography-based protein refolding 864(1999)229

Trapping the monomeric  $\alpha$ -helical state during unfolding of coiled-coils by reversed-phase liquid chromatography 890(2000)81

**Protein phases**

Enantiomeric separation of amino alcohols on protein phases using statistical experimental design. A comparative study 866(2000)15

**Protein–lead interactions**

Binding of lead to collagen type I and V and  $\alpha_2(I)$  CNBr (3,5) fragment by a modified Hummel–Dreyer method 852(1999)245

**Protein retention**

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

**Protein sequencing**

Identification of unusual (modified and non-encoded) amino acid residues in peptides by combinations of high-performance liquid chromatography and high-performance capillary electrophoresis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry 870(2000)255

**Protein synthesis**

Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor 852(1999)189

**Proton binding curves**

Dissociation of polyvalent electrolytes 853(1999)35

**Protoporphyrin-silica stationary phases**

Shape-selective separation of polycyclic aromatic hydrocarbons on protoporphyrin-silica phases. Effect of surface porphyrin distribution on column efficiency 859(1999)121

***Pseudomonas aeruginosa***

High-performance liquid chromatography method for the characterization of rhamnolipid mixtures produced by *Pseudomonas aeruginosa* UG2 on corn oil 864(1999)211

**Pseudo-stationary phases**

Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudostationary phase. Influence of methanol and acetonitrile on separation selectivity 853(1999)121

Electrokinetic chromatography 856(1999)465

Chiral separations using polymeric dipeptide surfactants: effect of number of chiral centers and steric factors 858(1999)219

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

Macromolecular surfactant as a pseudo-stationary phase in micellar electrokinetic capillary chromatography 871(2000)415

Novel approach to the analysis and use of fullerenes in capillary electrophoresis 873(2000)257

Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography 875(2000)179

New pseudo-stationary phases for electrokinetic capillary chromatography. Complexes between bovine serum albumin and sodium dodecyl sulfate 876(2000)169

Comparison of separation selectivity in capillary electrokinetic chromatography using a cationic linear polymeric pseudo-stationary phase or monomeric additives of similar structure 894(2000)25

Selectivity in microemulsion electrokinetic chromatography 897(2000)375

***Pueraria lobata***

Separation and purification of isoflavones from *Pueraria lobata* by high-speed counter-current chromatography 855(1999)709

**Pulsed diode laser**

Capillary array scanner for time-resolved detection and identification of fluorescently labelled DNA fragments 871(2000)299

**Pulsed discharge emission detection**

Pulsed discharge emission detector: an element-selective detector for gas chromatography 872(2000)119

Development and characterization of chlorine-selective pulsed discharge emission detector for gas chromatography 872(2000)141

Applications using the chlorine-selective pulsed discharge emission detector 872(2000)167

Characterization of chlorinated compounds using a dual chlorine-selective pulsed discharge emission detector–helium-pulsed discharge photoionization detector system 872(2000)179

**Pulsed discharge photoionization detection**

Characterization of chlorinated compounds using a dual chlorine-selective pulsed discharge emission detector–helium-pulsed discharge photoionization detector system 872(2000)179

**Pulsed flame photometric detection**

Comparison of gas chromatography–pulsed flame photometric detection–mass spectrometry, automated mass spectral deconvolution and identification system and gas chromatography–tandem mass spectrometry as tools for trace level detection and identification 868(2000)229

Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149

**Purge-and-trap methods**

Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the  $\text{ng l}^{-1}$  concentration level 893(2000)367

**Pyridylmethylaminoacetate ligands**

New ligand, *N*-(2-pyridylmethyl)aminoacetate, for use in the immobilised metal ion affinity chromatographic separation of proteins 852(1999)105



**Pyrolysis**

Pyrolysis–gas–liquid chromatography with atomic emission detection for the identification of *Corynebacterium* species 863(1999)243

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

Polymer additive analysis by pyrolysis–gas chromatography. I. Plasticizers 883(2000)199

Qualitative and quantitative analysis of a thermoset polymer, poly(benzoxazine), by pyrolysis–gas chromatography 886(2000)217

Polymer additive analysis by pyrolysis–gas chromatography. II. Flame retardants 886(2000)225

Number-average molecular mass determination of polymeric material by pyrolysis–gas chromatography 888(2000)209

Polymer additive analysis by pyrolysis–gas chromatography. III. Lubricants 891(2000)313

Polymer additive analysis by pyrolysis–gas chromatography. IV. Antioxidants 891(2000)325

**Pyrolysis oils**

Analysis of phenols in pyrolysis oils by gel permeation chromatography and multidimensional liquid chromatography 896(2000)343

**Q****Quality assurance**

Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the  $\text{ng l}^{-1}$  concentration level 893(2000)367

**Quality control**

Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the  $\text{ng l}^{-1}$  concentration level 893(2000)367

**Quantitative structure–retention relationships**

see Structure-retention relationships

**Quorum sensing**

Degradation pathway of homoserine lactone bacterial signal molecules by halogen antimicrobials identified by liquid chromatography with photodiode array and mass spectrometric detection 898(2000)153

**R****Racemization**

Large degree of racemization observed in the amide bond forming reaction on silica gel 878(2000)165

**Rainbow trout**

Quantification of the xenoestrogens 4-*tert*-octylphenol and bisphenol A in water and in fish tissue based on microwave assisted extraction, solid-phase extraction and liquid chromatography–mass spectrometry 864(1999)17

**Raman spectroscopy**

Characterization of  $\text{C}_{18}$ -bonded liquid chromatographic stationary phases by Raman spectroscopy: The effect of mobile phase composition 877(2000)25

Characterization of  $\text{C}_{18}$ -bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41

Investigation of mandelic acid bonding on Pirkle type chromatographic stationary phases by Raman spectroscopy 893(2000)37

**Random walk model**

Analysis of channel-geometry effects on separation efficiency in rectangular-capillary electrochromatography columns 869(2000)319

**Reaction-dispersive model**

Calculation of chromatographic band profiles with an implicit isotherm 862(1999)1

**Reactive-flow detection**

Analysis of organotin compounds by gas chromatography–reactive-flow detection 858(1999)245

**Receptor–ligand interactions**

Multiple-step ligand injection affinity capillary electrophoresis for determining binding constants of ligands to receptors 897(2000)339

**Refractive index detection**

Using a differential refractive index detector as a pressure transducer for online viscometry in exclusion chromatography 878(2000)1

On-column refractive index detector for flash chromatography 886(2000)9

Direct determination of amino acids and carbohydrates by high-performance capillary electrophoresis with refractometric detection 894(2000)253

**Regioisomers**

Rapid reversed-phase high-performance liquid chromatographic determination of the regioselectivity of lipoxygenase products on linoleic acid 859(1999)107

**Regression analysis**

Hold-up time in gas chromatography. V. Dependence of the retention of *n*-alkanes on the chromatographic variables in isothermal gas chromatography 878(2000)125

Calculation of Abraham solute descriptors from McReynolds gas chromatographic retention data 893(2000)339

**Regression models**

Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227

**Repeatability**

How to achieve higher repeatability and reproducibility in capillary electrophoresis 853(1999)83

Comparison of the repeatability of quantitative data measured in high-performance liquid chromatography with UV and atmospheric pressure chemical ionization mass spectrometric detection 873(2000)163

**Reproducibility**

How to achieve higher repeatability and reproducibility in capillary electrophoresis 853(1999)83

**Residence time distribution**

The influence of cell adsorbent interactions on protein adsorption in expanded beds 873(2000)195

**Resin selectivity**

Ion exchange on resins with temperature-responsive selectivity. III. Influence of complex formation stoichiometry on temperature dependence of resin selectivity 868(2000)143

**Resolution**

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. II. Two-run assay procedures 857(1999)21

Resolution optimisation in micellar electrokinetic chromatography using empirical models 867(2000)247

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Robustness testing for a capillary electrophoresis method using the “short-end injection” technique 897(2000)329

**Resolution prediction**

Prediction of electrophoretic mobilities in non-aqueous capillary electrophoresis. Optimal separation of quinolones in acetonitrile–water media 871(2000)367

**Response factors**

Group method approach to the estimation of response factors of unavailable substances in quantitative gas chromatography 888(2000)159

Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227

Optimization and evaluation of atomic emission gas chromatographic detection for nitrogen using the 388 nm molecular emission spectral band 898(2000)201

**Restricted-access media**

Direct analysis of food samples by high-performance liquid chromatography 880(2000)149

Novel use of a dual-zone restricted access sorbent: normal-phase solid-phase extraction separation of methyl oleate from polynuclear aromatic hydrocarbons stemming from semi-permeable membrane devices 883(2000)143

Analysis of neuropeptide Y and its metabolites by high-performance liquid chromatography–electrospray ionization mass spectrometry and integrated sample clean-up with a novel restricted-access sulphonic acid cation exchanger 890(2000)135

Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435

**Retention**

Relationship between retention, linear velocity and flow for counter-current chromatography 886(2000)283

**Retention, pressure-dependent**

Pressure-controlled high-performance liquid chromatographic study on the influence of rim chemistry on partial molar volume differences between free and complexed cyclodextrins 863(1999)127

**Retention behaviour**

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

Enantiomer separation of polychlorinated biphenyl atropisomers and polychlorinated biphenyl retention behavior on modified cyclodextrin capillary gas chromatography columns 866(2000)213

Retention behaviour of ceramides in sub-critical fluid chromatography in comparison with non-aqueous reversed-phase liquid chromatography 883(2000)211

Capillary electrochromatography with novel stationary phases. IV. Retention behavior of glycosphingolipids on porous and non-porous octadecyl sulfonated silica 887(2000)287

Pressure-induced retention of the lysozyme on reversed-phase liquid chromatography 891(2000)207

Influence of temperature on the behaviour of small linear peptides in capillary electrochromatography 893(2000)401

#### **Retention chart**

Retention behaviour of ceramides in sub-critical fluid chromatography in comparison with non-aqueous reversed-phase liquid chromatography 883(2000)211

#### **Retention data**

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. III. Results obtained with Kromasil C<sub>18</sub> columns 855(1999)423

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C<sub>18</sub> (2) columns 869(2000)181

#### **Retention factors**

Application of experimental design for the characterisation of a novel elution system for high-capacity anion chromatography with suppressed conductivity detection 855(1999)91

Comparison of capillary electrochromatography with high-performance liquid chromatography for the analysis of pirimicarb and related compounds 855(1999)215

Life-time studies with capillary electrochromatography columns operated under different conditions 894(2000)35

#### **Retention gap**

Efficiency through combining high-performance liquid chromatography and high resolution gas chromatography: progress 1995–1999 892(2000)407

#### **Retention indices**

Analysis of the equations for the temperature dependence of the retention index. I. Relation between equations 858(1999)65

Analysis of the equations for the temperature dependence of the retention index. II. Physico-chemical meaning of the parameters 859(1999)49

Gas chromatographic profiling and screening for phenols as isobutoxycarbonyl derivatives in aqueous samples 866(2000)87

Prediction of gas chromatographic retention indices of polychlorinated dibenzothiophenes on non-polar columns 866(2000)105

Retention index database for identification of general green leaf volatiles in plants by coupled capillary gas chromatography–mass spectrometry 890(2000)313

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

#### **Retention mechanisms**

Molecular mechanism of retention in reversed-phase high-performance liquid chromatography and classification of modern stationary phases by using quantitative structure–retention relationships 855(1999)455

Effects of organic modifiers on retention mechanism and selectivity in micellar electrokinetic capillary chromatography studied by linear solvation energy relationships 863(1999)69

Retention mechanism, isocratic and gradient-elution separation and characterization of (co)polymers in normal-phase and reversed-phase high-performance liquid chromatography 869(2000)65

Application of neutral hydrophobic hypercrosslinked polystyrene to the separation of inorganic anions by ion chromatography 884(2000)41

Retention behaviour of strong acid anions in ion-exclusion chromatography on sulfonate and carboxylate stationary phases 884(2000)61

Effects of DNA topology, temperature and solvent viscosity on DNA retardation in slalom chromatography 893(2000)115

Characterization of sulfobutyl ether- $\beta$ -cyclodextrins mixtures by anion-exchange chromatography using evaporative light scattering detection 897(2000)185

#### **Retention models**

Characterization of unstable ion-exchange chromatographic separation of proteins 852(1999)37

Retention models for ions in chromatography 855(1999)3

Reversed-phase high-performance liquid chromatography of ionogenic compounds: comparison of retention models 859(1999)1

Modelling retention in liquid chromatography as a function of solvent composition and pH of the mobile phase 869(2000)27

Protein retention in ion-exchange chromatography: effect of net charge and charge distribution 877(2000)13

Separation of basic solutes by reversed-phase capillary electrochromatography 887(2000)209

**Retention parameters**

Lipophilicity characterization by reversed-phase liquid chromatography of some furan derivatives 869(2000)49

**Retention prediction**

Prediction of protein retention at gradient elution conditions in ion-exchange chromatography 852(1999)351

Revision of a theoretical expression for gas–liquid chromatographic retention 852(1999)583

**Retention properties**

Retention properties of the fluorinated bonded phase on liquid chromatography of aromatic hydrocarbons 898(2000)141

**Retention time distribution method**

Pellicular expanded bed matrix suitable for high flow rates 878(2000)17

**Retention times**

Comparison of the repeatability of quantitative data measured in high-performance liquid chromatography with UV and atmospheric pressure chemical ionization mass spectrometric detection 873(2000)163

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. III. Improving the accuracy of computer simulation 857(1999)41

Hold-up time in gas chromatography. V. Dependence of the retention of *n*-alkanes on the chromatographic variables in isothermal gas chromatography 878(2000)125

**Retention volumes**

Lattice–fluid model for gas–liquid chromatography 862(1999)49

Investigation of the nematic–isotropic transition of a liquid crystalline polymer and determination of molecular diffusion coefficients using gas chromatography 893(2000)359

**Reversed-flow gas chromatography**

Gas chromatographic kinetic study of carbon monoxide oxidation over platinum–rhodium alloy catalysts 852(1999)507

**Reversed-phase materials**

Chromatographic properties of reversed-phase stationary phases under pressure- and electro-driven conditions 862(1999)121

**Reverse isomers**

High-performance liquid chromatographic resolution of reverse isomers of 1,2-diacyl-*rac*-glycerols as 3,5-dinitrophenylurethanes 893(2000)261

**Reviews**

Use of high-resolution techniques for the characterization of clotting factor VIII (Review) 852(1999)175

Retention models for ions in chromatography (Review) 855(1999)3

Solid-phase extraction: method development, sorbents, and coupling with liquid chromatography (Review) 856(1999)3

Membrane-based sample preparation coupled on-line to chromatography or electrophoresis (Review) 856(1999)55

Supercritical fluids in separation science – the dreams, the reality and the future (Review) 856(1999)83

Recent developments in microcolumn liquid chromatography (Review) 856(1999)117

Developments in sample preparation and separation techniques for the determination of inorganic ions by ion chromatography and capillary electrophoresis (Review) 856(1999)145

State-of-the-art in liquid chromatography–mass spectrometry (Review) 856(1999)179

Liquid chromatography–nuclear magnetic resonance spectroscopy (Review) 856(1999)199

Liquid chromatography–Fourier-transform infrared spectrometry (Review) 856(1999)213

Liquid chromatography–inductively coupled plasma mass spectrometry (Review) 856(1999)243

Programmed temperature vaporiser-based injection in capillary gas chromatography (Review) 856(1999)259

On-line combination of aqueous-sample preparation and capillary gas chromatography (Review) 856(1999)279

High-speed gas chromatography: an overview of various concepts (Review) 856(1999)315

Comprehensive two-dimensional gas chromatography: a hyphenated method with strong coupling between the two dimensions (Review) 856(1999)331

Gas chromatography with spectroscopic detectors (Review) 856(1999)349

Planar chromatography at the turn of the century (Review) 856(1999)399

The state of the art in thin-layer chromatography–mass spectrometry: a critical appraisal (Review) 856(1999)429

Overview of capillary electrophoresis and capillary electrochromatography (Review) 856(1999)443

Electrokinetic chromatography (Review) 856(1999)465

Coupling of biological sample handling and capillary electrophoresis (Review) 856(1999)483

Expanded-bed chromatography in primary protein purification (Review) 865(1999)129

- Enantiomer separation of drugs by capillary electromigration techniques (Review) 875(2000)3
- Enantioselective determination of drugs in body fluids by capillary electrophoresis (Review) 875(2000)27
- Chiral separation of amino acids and peptides by capillary electrophoresis (Review) 875(2000)43
- Enantioselective determination by capillary electrophoresis with cyclodextrins as chiral selectors (Review) 875(2000)89
- Enantiomeric separations of drugs using mixtures of charged and neutral cyclodextrins (Review) 875(2000)123
- Separation of drug enantiomers by capillary electrophoresis in the presence of neutral cyclodextrins (Review) 875(2000)135
- Enantiomer separation of drugs by micellar electrokinetic chromatography using chiral surfactants (Review) 875(2000)163
- Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography (Review) 875(2000)179
- Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis (Review) 875(2000)207
- Enantiomer separation of drugs by capillary electrophoresis using proteins as chiral selectors (Review) 875(2000)235
- Enantiomer separation of chiral pharmaceuticals by capillary electrochromatography (Review) 875(2000)255
- Enantiomeric separations by nonaqueous capillary electrophoresis (Review) 875(2000)277
- Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants (Review) 879(2000)83
- Sample preparation for the analysis of flavors and off-flavors in foods (Review) 880(2000)3
- Applications of solid-phase microextraction in food analysis (Review) 880(2000)35
- $\beta_2$ -Agonist extraction procedures for chromatographic analysis (Review) 880(2000)69
- Thin-layer chromatography in food and agricultural analysis (Review) 880(2000)129
- Direct analysis of food samples by high-performance liquid chromatography (Review) 880(2000)149
- Perfusion chromatography: an emergent technique for the analysis of food proteins (Review) 880(2000)169
- C<sub>30</sub> Stationary phases for the analysis of food by liquid chromatography (Review) 880(2000)189
- On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic–diode array detection–mass spectrometric techniques (Review) 880(2000)203
- Electrophoretic methods in the analysis of beverages (Review) 880(2000)243
- Herbicide and plant growth regulator analysis by capillary electrophoresis (Review) 880(2000)281
- Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry (Review) 880(2000)295
- Analysis of natural food pigments by capillary electrophoresis (Review) 880(2000)311
- Application of chromatography and mass spectrometry to the characterization of food proteins and derived peptides (Review) 881(2000)1
- Electrophoresis of cereal storage proteins (Review) 881(2000)23
- Authentication of vegetable oils by chromatographic techniques (Review) 881(2000)93
- Chromatographic analysis of unsaponifiable compounds of olive oils and fat-containing foods (Review) 881(2000)105
- Chromatographic analysis of minor constituents in vegetable oils (Review) 881(2000)131
- Chromatographic analysis of tocol-derived lipid antioxidants (Review) 881(2000)197
- Determination of thiamine by high-performance liquid chromatography (Review) 881(2000)267
- Dehydroascorbic acid (Review) 881(2000)299
- Analytical techniques for determining biotin (Review) 881(2000)331
- Determination of organic acids in food samples by capillary zone electrophoresis (Review) 881(2000)357
- High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides (Review) 881(2000)365
- Analysis of anthocyanins in foods by liquid chromatography, liquid chromatography–mass spectrometry and capillary electrophoresis (Review) 881(2000)403
- Determination of tea catechins (Review) 881(2000)411

- Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis (Review) 881(2000)425
- Analysis of resveratrol in wine by capillary electrophoresis (Review) 881(2000)471
- Chromatographic determination of carotenoids in foods (Review) 881(2000)543
- Monoterpenes in grape juice and wines (Review) 881(2000)557
- Analysis of organic sulfur compounds in wine aroma (Review) 881(2000)569
- Advances in the determination of inorganic anions by ion chromatography (Review) 881(2000)607
- Chromatographic determination of the mycotoxin patulin in fruit and fruit juices (Review) 882(2000)17
- Regulatory problems caused by contamination, a frequently overlooked cause of veterinary drug residues (Review) 882(2000)37
- Chromatographic analysis of tetracycline antibiotics in foods (Review) 882(2000)109
- Pesticide residue determination in fruit and vegetables by liquid chromatography–mass spectrometry (Review) 882(2000)153
- Analysis of herbicide residues in cereals, fruits and vegetables (Review) 882(2000)175
- Polycyclic aromatic hydrocarbons in edible fats and oils: occurrence and analytical methods (Review) 882(2000)245
- Contamination in food from packaging material (Review) 882(2000)255
- Determination of synthetic phenolic antioxidants in food by high-performance liquid chromatography (Review) 882(2000)271
- Application of chromatographic and electrophoretic methodology to the speciation of organomercury compounds in food analysis (Review) 882(2000)283
- Detection techniques in ion analysis: what are our choices? (Review) 884(2000)3
- Recent developments in the separation of inorganic and small organic ions by capillary electrophoresis (Review) 884(2000)261
- Fifty years of solid-phase extraction in water analysis – historical development and overview (Review) 885(2000)3
- Contributions of theory to method development in solid-phase extraction (Review) 885(2000)17
- Recent developments in polymer-based sorbents for solid-phase extraction (Review) 885(2000)51
- Graphitized carbons for solid-phase extraction (Review) 885(2000)73
- Automating solid-phase extraction: current aspects and future prospects (Review) 885(2000)97
- Matrix solid-phase dispersion (Review) 885(2000)115
- Sorbent trapping of volatile organic compounds from air (Review) 885(2000)129
- Evolution of solid-phase microextraction technology (Review) 885(2000)153
- Solid-phase extraction for multiresidue analysis of organic contaminants in water (Review) 885(2000)195
- Multiresidue methods using solid-phase extraction techniques for monitoring priority pesticides, including triazines and degradation products, in ground and surface waters (Review) 885(2000)217
- Solid-phase extraction of acidic herbicides (Review) 885(2000)237
- Solid-phase extraction of quaternary ammonium herbicides (Review) 885(2000)251
- Solid-phase extraction of polycyclic aromatic compounds (Review) 885(2000)273
- Solid-phase extraction of phenols (Review) 885(2000)291
- Solid-phase extraction for combinatorial libraries (Review) 885(2000)305
- Update on solid-phase extraction for the analysis of lipid classes and related compounds (Review) 885(2000)321
- Ion-pair solid-phase extraction (Review) 885(2000)343
- Solid-phase microextraction in pesticide residue analysis (Review) 885(2000)389
- Application of solid-phase microextraction for determination of organic vapours in gaseous matrices (Review) 885(2000)405
- Solid-phase micro-extraction of drugs from biological matrices (Review) 885(2000)445
- Effect of surface heterogeneity on adsorption on solid surfaces. Application of inverse gas chromatography in the studies of energetic heterogeneity of adsorbents (Review) 886(2000)133

- Design of the monolithic polymers used in capillary electrochromatography columns (Review) 887(2000)3
- Open tubular capillary electrokinetic chromatography in etched fused-silica tubes (Review) 887(2000)31
- Packing columns for capillary electrochromatography (Review) 887(2000)43
- Instrumentation for capillary electrochromatography (Review) 887(2000)55
- Use of on-line mass spectrometric detection in capillary electrochromatography (Review) 887(2000)85
- Capillary electrochromatography–mass spectrometry for the separation and identification of isomeric polyaromatic hydrocarbon DNA adducts derived from in vitro reactions (Review) 887(2000)103
- Gradient elution techniques for capillary electrochromatography (Review) 887(2000)115
- Highly selective separations by capillary electrochromatography: molecular imprint polymer sorbents (Review) 887(2000)125
- Specific applications of capillary electrochromatography to biopolymers, including proteins, nucleic acids, peptide mapping, antibodies, and so forth (Review) 887(2000)137
- Capillary electroosmotic chromatography of peptides (Review) 887(2000)165
- Stationary phases for capillary electrochromatography (Review) 887(2000)313
- Solid-phase microextraction: a promising technique for sample preparation in environmental analysis (Review) 889(2000)3
- New configurations and applications of molecularly imprinted polymers (Review) 889(2000)15
- Role of chromatography in the analysis of sugars, carboxylic acids and amino acids in food (Review) 891(2000)1
- Biomimetic dyes as affinity chromatography tools in enzyme purification (Review) 891(2000)33
- Analysis of agrochemicals by capillary electrophoresis (Review) 891(2000)45
- Synthesis of spherical porous silicas in the micron and submicron size range: challenges and opportunities for miniaturized high-resolution chromatographic and electrokinetic separations (Review) 892(2000)47
- Chromatographic silanol activity test procedures: the quest for a universal test (Review) 892(2000)57
- Optimization methods in chromatography and capillary electrophoresis (Review) 892(2000)75
- Practitioner's guide to method development in thin-layer chromatography (Review) 892(2000)123
- Capillary electrophoresis coupled to biosensor detection (Review) 892(2000)143
- Non-aqueous capillary electrophoresis (Review) 892(2000)155
- Background theory and applications of microemulsion electrokinetic chromatography (Review) 892(2000)171
- Alternative methods providing enhanced sensitivity and selectivity in capillary electroseparation experiments (Review) 892(2000)203
- Electrochromatography (Review) 892(2000)219
- Advances in column technology and instrumentation in capillary electrochromatography (Review) 892(2000)257
- Capillary electrochromatography on silica columns: factors influencing performance (Review) 892(2000)279
- Multiple hyphenation of liquid chromatography with nuclear magnetic resonance spectroscopy, mass spectrometry and beyond (Review) 892(2000)315
- Electron–capture mass spectrometry: recent advances (Review) 892(2000)329
- Efficiency through combining high-performance liquid chromatography and high resolution gas chromatography: progress 1995–1999 (Review) 892(2000)407
- Recent and future developments of liquid chromatography in pesticide trace analysis (Review) 892(2000)435
- Applications of mass spectrometry to food proteins and peptides (Review) 893(2000)1
- Optimisation and characterisation of silica-based reversed-phase liquid chromatographic systems for the analysis of basic pharmaceuticals (Review) 897(2000)1
- Rheum officinale***  
Preparative isolation and purification of hydroxyanthraquinones from *Rheum officinale* Baill by high-speed counter-current chromatography using pH-modulated stepwise elution 858(1999)103
- Rhodium–platinum catalysts**  
Gas chromatographic kinetic study of carbon monoxide oxidation over platinum–rhodium alloy catalysts 852(1999)507

**Rhubarb**

Computer-aided development of a high-performance liquid chromatographic method for the determination of hydroxyanthraquinone derivatives in Chinese herb medicine rhubarb 857(1999)167

**Rice**

Applications of in-capillary reaction micellar electrokinetic chromatography in the food industry 880(2000)295

**Ring opening–closure equilibrium**

Influence of ring opening–closure equilibrium of oxanine, a novel damaged nucleobase, on migration behavior in capillary electrophoresis 877(2000)225

**Robustness testing**

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Robustness testing of a reversed-phase high-performance liquid chromatographic assay: comparison of fractional and asymmetrical factorial designs 874(2000)167

Robustness testing for a capillary electrophoresis method using the “short-end injection” technique 897(2000)329

**Rosemary**

Liquid chromatographic–mass spectrometric analysis of supercritical-fluid extracts of rosemary plants 870(2000)491

**Rubber**

Development and validation of a capillary electrophoresis method for the measurement of short-chain organic acids in natural rubber latex 894(2000)135

**Rubber elongation factor**

Characterisation of the rubber elongation factor from ammoniated latex by electrophoresis and mass spectrometry 890(2000)145

**Ruditapes decussatus**

Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99

**Ruggedness**

Use of optimization software to determine rugged analysis conditions in high-performance liquid chromatography 868(2000)13

**S*****Saccharomyces cerevisiae***

Affinity chromatography of human estrogen receptor- $\alpha$  expressed in *Saccharomyces cerevisiae*. Combination of heparin- and 17 $\beta$ -estradiol-affinity chromatography 852(1999)161

**Saline gold processing samples**

Ion chromatographic determination of cyanate in saline gold processing samples 855(1999)267

***Sambucus nigra***

Analysis of natural food pigments by capillary electrophoresis 880(2000)311

**Sample capacity**

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

**Sample displacement chromatography**

Development of simultaneous purification methodology for multiple synthetic peptides by reversed-phase sample displacement chromatography 893(2000)81

**Sample evaporation**

Video-taped sample evaporation in hot chambers simulating gas chromatography split/splitless injectors. I. Thermospray injection 897(2000)237

Video-taped sample evaporation in hot chambers simulating gas chromatography split and splitless injectors. II. Injection with band formation 897(2000)247

**Sample handling**

see also Clean-up methods, Extraction methods, Preconcentration, Sample preparation and Special Issue Index

Nanoliter-scale sample preparation methods directly coupled to polymethylmethacrylate-based microchips and gel-filled capillaries for the analysis of oligonucleotides 853(1999)107

Highly-sensitive micellar electrokinetic chromatographic analysis of dioxin-related compounds using on-line concentration 853(1999)413

Quantification of deoxynivalenol in wheat using an immunoaffinity column and liquid chromatography 859(1999)23

On the performance and inertness of different materials used for the enrichment of sulfur compounds from air and gaseous samples 864(1999)345



Comparison of nutshell granular activated carbons to commercial adsorbents for the purge-and-trap gas chromatographic analysis of volatile organic compounds 879(2000)169

On-line isotachopheretic sample focusing for loadability enhancement in capillary electrochromatography–mass spectrometry 879(2000)219

Application of ion-exchange cartridge clean-up in food analysis. III. Determination of benzylpenicillin, phenoxymethyl-penicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in bovine liver and kidney by liquid chromatography with ultraviolet detection 880(2000)85

Analysis of organic sulfur compounds in wine aroma 881(2000)569

On-line microdialysis–ion chromatographic determination of inorganic anions in olive-oil mill wastewater 882(2000)321

Analysis of neuropeptide Y and its metabolites by high-performance liquid chromatography–electrospray ionization mass spectrometry and integrated sample clean-up with a novel restricted-access sulphonic acid cation exchanger 890(2000)135

Efficiency of pretreatment of aqueous samples using a macroporous strong anion-exchange resin on the determination of nerve gas hydrolysis products by gas chromatography–mass spectrometry after *tert*.-butyldimethylsilylation 891(2000)295

Time-weighted average sampling of volatile and semi-volatile airborne organic compounds by the solid-phase microextraction device 892(2000)455

Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the  $\text{ng l}^{-1}$  concentration level 893(2000)367

Integrated electroosmotically-driven on-line sample purification system for nanoliter DNA sequencing by capillary electrophoresis 894(2000)179

Determination of deramciclane and *N*-desmethylderamciclane in human plasma by liquid chromatography–tandem mass spectrometry using off-line robotic sample pretreatment 896(2000)279

### Sample introduction

see also Injection methods

Visualization of sample introduction in liquid chromatography columns. The effect of the frit diameter 855(1999)367

Visualization of sample introduction in liquid chromatographic columns. Contribution of a flow distributor on the sample band shape 865(1999)83

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. I. Injection procedure and column stability 876(2000)37

### Sample preparation

see also Clean-up methods, Extraction methods, Preconcentration, Sample handling and Special Issue Index

Comparison of turbulent-flow chromatography with automated solid-phase extraction in 96-well plates and liquid–liquid extraction used as plasma sample preparation techniques for liquid chromatography–tandem mass spectrometry 854(1999)23

Membrane-based sample preparation coupled on-line to chromatography or electrophoresis 856(1999)55

Developments in sample preparation and separation techniques for the determination of inorganic ions by ion chromatography and capillary electrophoresis 856(1999)145

Sample cleanup and reversed-phase high-performance liquid chromatographic analysis of polar aromatic compounds in groundwater samples from a former gas plant 862(1999)137

On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279

Coupling of biological sample handling and capillary electrophoresis 856(1999)483

On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207

Determination of albendazole and its main metabolites in ovine plasma by liquid chromatography with dialysis as an integrated sample preparation technique 870(2000)121

Gas chromatographic–mass spectrometric analysis of acids and phenols in distilled alcohol beverages. Application of anion-exchange disk extraction combined with in-vial elution and silylation 873(2000)29

Sample preparation for the analysis of flavors and off-flavors in foods 880(2000)3

Determination of 1-nitropyrene in herbs after selective enrichment by a sol–gel-generated immunoaffinity column 880(2000)113

Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175

Multiresidue analysis of seventeen pesticides in wine by gas chromatography with mass-selective detection 882(2000)205

Polycyclic aromatic hydrocarbons in edible fats and oils: occurrence and analytical methods 882(2000)245

Automating solid-phase extraction: current aspects and future prospects 885(2000)97

On-line preconcentration of niobium(V) and tantalum(V) as 4-(2-pyridylazo) resorcinol–citrate ternary complexes in geological samples by ion interaction high-performance liquid chromatography 885(2000)369

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Analysis of highly saline samples by capillary zone electrophoresis: enhanced direct UV detection of inorganic anions using on-capillary preconcentration and clean-up techniques 888(2000)309

Reversed migration micellar electrokinetic chromatography with off-line and on-line concentration analysis of phenylurea herbicides 892(2000)187

Purification of Substance P endopeptidase activity in the rat ventral tegemental area with the Äkta-Purifier chromatographic system 893(2000)107

Quantitative determination of  $\alpha$ -,  $\beta$ -,  $\gamma$ - and  $\delta$ -tocopherols in human serum by high-performance liquid chromatography and gas chromatography–mass spectrometry as trimethylsilyl derivatives with a two-step sample preparation 896(2000)209

Analysis of phenols in pyrolysis oils by gel permeation chromatography and multidimensional liquid chromatography 896(2000)343

### Sample stacking

see also Injection methods

Field amplified injection in the presence of salts for capillary electrophoresis 853(1999)3

Head-column field-amplified sample stacking in binary system capillary electrophoresis. Preparation of extracts for determination of opioids in microliter amounts of body fluids 853(1999)95

Capillary electrophoresis of double-stranded DNA in an untreated capillary 853(1999)349

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

Electrokinetic chromatography 856(1999)465

Standardization of capillary zone electropherograms obtained by using field enhanced sample stacking 894(2000)3

### Sample volume

Performance of experimental sample injectors for high-performance liquid chromatography microcolumns 869(2000)231

### Sandwich injection

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. I. Injection procedure and column stability 876(2000)37

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. II. Methods of detection and quantification and overall long-term performance 878(2000)45

### Scanning fluorescence detection, fast

Determination of naphthalenesulfonates in water by on-line ion-pair solid-phase extraction and ion-pair liquid chromatography with fast-scanning fluorescence detection 890(2000)289

### Scapanes australis

Role of solid-phase microextraction in the identification of highly volatile pheromones of two Rhinoceros beetles *Scapanes australis* and *Strategus aloeus* (Coleoptera, Scarabaeidae, Dynastinae) 885(2000)433

### Schistosoma japonicum

Affinity purification of *Schistosoma japonicum* glutathione-S-transferase and its site-directed mutants with glutathione affinity chromatography and immobilized metal affinity chromatography 852(1999)151

### Scopolia japonica

Strategies for supercritical fluid extraction of hyoscyamine and scopolamine salts using basified modifiers 863(1999)47

### Secondary consolidation

Evaluation of the secondary consolidation of columns for liquid chromatography by ultrasonic irradiation 878(2000)153

### Sediments

see also Marine sediments

Application of pressurized liquid extraction followed by gas chromatography–mass spectrometry to determine 4-nonylphenols in sediments 866(2000)79

Determination of butyltin species in water and sediment by solid-phase microextraction–gas chromatography–flame ionization detection 873(2000)63

Speciation of butyl- and phenyltin compounds in sediments using pressurized liquid extraction and liquid chromatography–inductively coupled plasma mass spectrometry 879(2000)137

Validation of an analytical procedure for polychlorinated biphenyls, coplanar polychlorinated biphenyls and polycyclic aromatic hydrocarbons in environmental samples 886(2000)153

Comparison of two extraction methods independently developed on two conceptually different automated supercritical fluid extraction systems for the determination of polychlorinated biphenyls in sediments 891(2000)195

Miniaturised pressurised liquid extraction of polycyclic aromatic hydrocarbons from soil and sediment with subsequent large-volume injection–gas chromatography 891(2000)275

### Selectivity

see also Special Issue Index

Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudostationary phase. Influence of methanol and acetonitrile on separation selectivity 853(1999)121

Comparison of capillary electrochromatography with high-performance liquid chromatography for the analysis of pirimicarb and related compounds 855(1999)215

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1

Effects of organic modifiers on retention mechanism and selectivity in micellar electrokinetic capillary chromatography studied by linear solvation energy relationships 863(1999)69

Development of a new cyano-bonded column for high-performance liquid chromatography 869(2000)143

Increasing the reliability of the identification by high-performance liquid chromatography by means of selective and/or sensitive detection 869(2000)275

Evidence for selectivity of absorption of volatile organic compounds by a polydimethylsiloxane solid-phase microextraction fibre 885(2000)457

Characterization of chemical selectivity in micellar electrokinetic chromatography. VI. Effects of surfactant counter-ion 888(2000)229

Separation of acidic solutes by nonaqueous capillary electrophoresis in acetonitrile-based media. Combined effects of deprotonation and heteroconjugation 888(2000)251

Alternative methods providing enhanced sensitivity and selectivity in capillary electroseparation experiments 892(2000)203

Characterization of reversed-phase columns using the linear free energy relationship. III. Effect of the organic modifier and the mobile phase composition 893(2000)215

Comparison of separation selectivity in capillary electrokinetic chromatography using a cationic linear polymeric pseudo-stationary phase or monomeric additives of similar structure 894(2000)25

Characterization of synthetic polyelectrolytes by capillary electrophoresis 894(2000)105

Separation selectivity of anionic metal complexes of *N,N'*-bis(hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid in ion and ion electrokinetic chromatography 895(2000)255

Selectivity differences for  $C_{18}$  and  $C_8$  reversed-phase columns as a function of temperature and gradient steepness. I. Optimizing selectivity and resolution 897(2000)37

Selectivity differences for  $C_{18}$  and  $C_8$  reversed-phase columns as a function of temperature and gradient steepness. II. Minimizing column reproducibility problems 897(2000)51

Selectivity in microemulsion electrokinetic chromatography 897(2000)375

Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13

### Selectivity equivalence

Selectivity equivalence of poly(ethylene glycol) stationary phases for gas chromatography 898(2000)211

### Semipermeable membrane devices

Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83

### Sensitivity

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection 867(2000)187

Alternative methods providing enhanced sensitivity and selectivity in capillary electroseparation experiments 892(2000)203

### Sephadex G-15

Coupling of chromatographic ion-exchange reaction with change-partner reaction. Presumed mechanism on the ion–ion interaction of inorganic ions in Sephadex G-15 columns 869(2000)91

### Sequential projection pursuit

Exploratory chemometric analysis of the classification of pharmaceutical substances based on chromatographic data 897(2000)23

### Sewage sludge

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29

**Sewage treatment plants**

Characterization of nonyl phenol ethoxylates in sewage treatment plants by combined precursor ion scanning and multiple reaction monitoring 854(1999)245

**Shape selectivity**

Studies of retention and stability of a horizontally polymerized bonded phase for reversed-phase liquid chromatography 868(2000)153

Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71

**Shear-driven chromatography**

On the possibility of shear-driven chromatography: a theoretical performance analysis 855(1999)57

**Sheath liquid effects**

Sheath liquid effects in capillary high-performance liquid chromatography–electrospray mass spectrometry of oligonucleotides 870(2000)413

**Shellfish poisoning**

Separation of paralytic shellfish poisoning toxins on Chromarods-SIII by thin-layer chromatography with the Iatroscan (mark 5) and flame thermionic detection 855(1999)657

Development of a method for the identification of azaspiracid in shellfish by liquid chromatography–tandem mass spectrometry 871(2000)13

Liquid chromatography–electrospray ionization mass spectrometry of the diarrhetic shellfish-poisoning toxins okadaic acid, dinophysistoxin-1 and pectenotoxin-6 in bivalves 874(2000)199

Development and validation of a high-performance liquid chromatographic method using fluorimetric detection for the determination of the diarrhetic shellfish poisoning toxin okadaic acid without chlorinated solvents 876(2000)117

**Short tandem repeats sequences**

Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23

**Short-end injection**

Fast chiral separations using sulfated  $\beta$ -cyclodextrin and short-end injection in capillary electrophoresis 875(2000)403

Robustness testing for a capillary electrophoresis method using the “short-end injection” technique 897(2000)329

**Signal averaging**

Towards a quantitative definition of perfection in chromatographic analyses 868(2000)305

**Signal processing**

Adaptive numerical morphological filter for identifying chromatographic signals 855(1999)305

**Signature-peptide approach**

Signature-peptide approach to detecting proteins in complex mixtures 870(2000)295

**Silanization**

Open tubular capillary electrokinetic chromatography in etched fused-silica tubes (Review) 887(2000)31

**Silanol activity**

Chromatographic silanol activity test procedures: the quest for a universal test 892(2000)57

**Silanol groups, residual**

Comparison of aqueous and non-aqueous capillary electrochromatography for the separation of basic solutes 887(2000)233

**Silanols**

Chromatographic properties of reversed-phase stationary phases under pressure- and electro-driven conditions 862(1999)121

**Silanol-blocking agents**

Comparative study of divalent metals and amines as silanol-blocking agents in reversed-phase liquid chromatography 855(1999)121

**Silanophilicity**

Studies of retention and stability of a horizontally polymerized bonded phase for reversed-phase liquid chromatography 868(2000)153

**Silica**

Low-carbon silica sorbents for solid-phase extraction 868(2000)1

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3

Synthesis of spherical porous silicas in the micron and submicron size range: challenges and opportunities for miniaturized high-resolution chromatographic and electrokinetic separations 892(2000)47

Chromatographic silanol activity test procedures: the quest for a universal test 892(2000)57

Novel trifunctional diamine for silica coating in capillary zone electrophoresis 894(2000)53

**Silica, aluminium adsorbing**

Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113

**Silica, microparticulate**

Towards the column bed stabilization of columns in capillary electroendosmotic chromatography. Immobilization of microparticulate silica columns to a continuous bed 887(2000)327

**Silica, octadecyl sulfonated**

Capillary electrochromatography with novel stationary phases. IV. Retention behavior of glycosphingolipids on porous and non-porous octadecyl sulfonated silica 887(2000)287

**Silica, titanium grafted**

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by  $\gamma$ -radiation on titanium-grafted silica 869(2000)137

**Silica, zirconized**

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

**Silica–magnetite sorbents**

Application of magnetite and silica–magnetite composites to the isolation of genomic DNA 890(2000)159

**Silica packings**

Towards the ultimate minimum particle diameter of silica packings in capillary electrochromatography 887(2000)339

Capillary electrochromatography on silica columns: factors influencing performance 892(2000)279

**SIMPLISMA**

Application of several modified peak purity assays to real complex multicomponent mixtures by high-performance liquid chromatography with diode-array detection 855(1999)487

**Simulated moving bed chromatography**

Optimization of throughput and desorbent consumption in simulated moving-bed chromatography for paclitaxel purification 855(1999)71

Development of an high-performance liquid chromatographic simulated moving bed separation from an industrial perspective 865(1999)35

Separation of stereoisomers in a simulated moving bed-supercritical fluid chromatography plant 865(1999)175

Enantiomers separation by simulated moving bed chromatography. Non-instantaneous equilibrium at the solid–fluid interface 865(1999)187

Optimization of simulated moving bed plants with low efficient stationary phases: separation of fructose and glucose 868(2000)169

Design of a simulated moving bed unit for sucrose–betaine separations 872(2000)23

Novel simulated moving-bed method for reduced solvent consumption 873(2000)149

Purification of an ascomycin derivative with simulated moving bed chromatography. A case study 874(2000)155

Design and optimisation of a simulated moving bed unit: role of deviations from equilibrium theory 876(2000)3

On-line monitoring of enantiomer concentration in chiral simulated moving bed chromatography 888(2000)73

Enantioseparation of 1-phenyl-1-propanol by simulated moving bed under linear and nonlinear conditions 893(2000)307

**Simultaneous distillation and extraction**

Analysis of polycyclic aromatic hydrocarbons, phenols and aromatic amines in particulate phase cigarette smoke using simultaneous distillation and extraction as a sole sample clean-up step 898(2000)111

**Single-strand conformation polymorphism analysis**

Single-strand conformation polymorphism analysis to detect the p53 mutation in colon tumor samples by capillary electrophoresis 894(2000)171

**Size-exclusion chromatography**

Off-line size-exclusion chromatographic fractionation–matrix-assisted laser desorption ionization time-of-flight mass spectrometry for polymer characterization. Theoretical and experimental study 896(2000)19

**Slalom chromatography**

Mobile-phase-viscosity dependence on DNA separation in slalom chromatography 886(2000)1

Effects of DNA topology, temperature and solvent viscosity on DNA retardation in slalom chromatography 893(2000)115

**Slurry packing**

Hydrodynamic aspects of slurry packing processes in microcolumn liquid chromatography 883(2000)11

**Soft drinks**

Development of a capillary electrophoresis method for the simultaneous analysis of artificial sweeteners, preservatives and colours in soft drinks 876(2000)213

Electrophoretic methods in the analysis of beverages 880(2000)243

**Software, integration**

How to achieve higher repeatability and reproducibility in capillary electrophoresis 853(1999)83

**Soil**

Determination of sulfonylurea degradation products in soil by liquid chromatography–ultraviolet detection followed by confirmatory liquid chromatography–tandem mass spectrometry 855(1999)575

Liquid chromatographic determination of imazosulfuron in drinking water and in soil using ultraviolet detection 857(1999)327

Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29

Soil thin-layer chromatography and pesticide mobility through soil microstructures. New technical approach 864(1999)145

New gas chromatographic method for residue determination of dithiopyr in soil, wheat grain and straw 864(1999)351

Selective extraction of hydrocarbons, phosphonates and phosphonic acids from soils by successive supercritical fluid and pressurized liquid extractions 866(2000)231

Determination of phenols using simultaneous steam distillation–extraction 867(2000)281

Comparison of gas chromatography–pulsed flame photometric detection–mass spectrometry, automated mass spectral deconvolution and identification system and gas chromatography–tandem mass spectrometry as tools for trace level detection and identification 868(2000)229

Study of the behaviour of azinphos-methyl in a clay mineral by batch and column leaching 869(2000)481

Determination of linuron and related compounds in soil by microwave-assisted solvent extraction and reversed-phase liquid chromatography with UV detection 869(2000)487

Extraction of hydrocarbon contamination from soils using accelerated solvent extraction 874(2000)217

Determination of triazines in soil leachates by solid-phase microextraction coupled to gas chromatography–mass spectrometry 874(2000)247

Extraction of DDT [1,1,1,-trichloro-2,2-bis(*p*-chlorophenyl)ethane] and its metabolites DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethylene] and DDD [1,1-dichloro-2,2-bis(*p*-chlorophenyl)- ethane] from aged contaminated soil 874(2000)257

Pressurized liquid extraction of polycyclic aromatic hydrocarbons from contaminated soils 883(2000)151

Laser-induced dispersed fluorescence detection of polycyclic aromatic compounds in soil extracts separated by capillary electrochromatography 883(2000)231

Solid-phase extraction of quaternary ammonium herbicides 885(2000)251

New methods for determination of glyphosate and (aminomethyl)phosphonic acid in water and soil 886(2000)207

Analysis of flufenacet in soil, wheat grain and straw by gas chromatography 886(2000)319

Analysis of agrochemicals by capillary electrophoresis (Review) 891(2000)45

Miniaturised pressurised liquid extraction of polycyclic aromatic hydrocarbons from soil and sediment with subsequent large-volume injection–gas chromatography 891(2000)275

Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421

Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435

Pressurised liquid extraction of ketones of polycyclic aromatic hydrocarbons from soil 893(2000)201

Continuous microwave-assisted extraction, solvent changeover and preconcentration of monophenols in agricultural soils 897(2000)279

Extraction of cloransulam-methyl from soil with subcritical water and supercritical CO<sub>2</sub> 897(2000)405

***Solanum brevidens***

Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301

**Solid-phase extraction**

see also Extraction methods

Solid-phase extraction: method development, sorbents, and coupling with liquid chromatography 856(1999)3

Coupling of biological sample handling and capillary electrophoresis 856(1999)483

Efficient solid-phase extraction procedures for trace enrichment of priority phenols from industrial effluents with high total organic carbon content 857(1999)97

Study of applicability of various solid-phase extraction materials for sample handling in screening analysis of organic micropollutants in water 857(1999)247

- Determination of zearalenone in grains by high-performance liquid chromatography–tandem mass spectrometry after solid-phase extraction with RP-18 columns or immunoaffinity columns 858(1999)167
- Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29
- Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121
- Low-carbon silica sorbents for solid-phase extraction 868(2000)1
- Does further clean-up reduce the matrix enhancement effect in gas chromatographic analysis of pesticide residues in food? 868(2000)51
- Comparison of commercial solid-phase extraction sorbents for the sample preparation of potato glycoalkaloids 869(2000)301
- Kinetics of solid-phase extraction and solid-phase microextraction in thin adsorbent layer with saturation sorption isotherm 873(2000)39
- Part-per-trillion level determination of antifouling pesticides and their byproducts in seawater samples by off-line solid-phase extraction followed by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)27
- Comparison of different commercial solid-phase extraction cartridges used to extract heterocyclic amines from a lyophilised meat extract 880(2000)101
- Quantitative analysis of mutagenic heterocyclic aromatic amines in cooked meat using liquid chromatography–atmospheric pressure chemical ionisation tandem mass spectrometry 883(2000)89
- Novel use of a dual-zone restricted access sorbent: normal-phase solid-phase extraction separation of methyl oleate from polynuclear aromatic hydrocarbons stemming from semi-permeable membrane devices 883(2000)143
- Fifty years of solid-phase extraction in water analysis – historical development and overview 885(2000)3
- Contributions of theory to method development in solid-phase extraction 885(2000)17
- Investigation into the effects of temperature and stirring rate on the solid-phase extraction of diuron from water using a C<sub>18</sub> extraction disk 885(2000)41
- Recent developments in polymer-based sorbents for solid-phase extraction 885(2000)51
- Graphitized carbons for solid-phase extraction 885(2000)73
- Automating solid-phase extraction: current aspects and future prospects 885(2000)97
- Matrix solid-phase dispersion 885(2000)115
- Multiresidue methods using solid-phase extraction techniques for monitoring priority pesticides, including triazines and degradation products, in ground and surface waters 885(2000)217
- Solid-phase extraction of acidic herbicides 885(2000)237
- Solid-phase extraction of quaternary ammonium herbicides 885(2000)251
- Solid-phase extraction of polycyclic aromatic compounds 885(2000)273
- Solid-phase extraction of phenols 885(2000)291
- Solid-phase extraction for combinatorial libraries 885(2000)305
- Update on solid-phase extraction for the analysis of lipid classes and related compounds 885(2000)321
- Ion-pair solid-phase extraction 885(2000)343
- On-line dual-precursor-based trace enrichment for the determination of polar and acidic microcontaminants in river water by liquid chromatography with diode-array UV and tandem mass spectrometric detection 885(2000)377
- Static subcritical water extraction with simultaneous solid-phase extraction for determining polycyclic aromatic hydrocarbons on environmental solids 886(2000)237
- Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197
- Solid-phase extraction with a dibutylmelamine-imprinted polymer as triazine herbicide-selective sorbent 889(2000)25
- Solid-phase extraction and high-performance liquid chromatographic separation of pigments of red wines 889(2000)51
- Use of molecularly imprinted solid-phase extraction for the selective clean-up of clenbuterol from calf urine 889(2000)105
- Solid-phase extraction of polar hydrophilic aromatic sulfonates followed by capillary zone electrophoresis–UV absorbance detection and ion-pair liquid chromatography–diode array UV detection and electrospray mass spectrometry 890(2000)225

- Determination of naphthalenesulfonates in water by on-line ion-pair solid-phase extraction and ion-pair liquid chromatography with fast-scanning fluorescence detection 890(2000)289
- Determination of phenolic acids in fruit juices by isocratic column liquid chromatography 891(2000)183
- Comparison of different liquid chromatography methods for the determination of corticosteroids in biological matrices 893(2000)55
- Comparison of different sorbent materials for *on-line* solid-phase extraction with liquid chromatography–atmospheric pressure chemical ionization mass spectrometry of phenols 896(2000)159
- Analysis of the enzymatic racemization of D-aspartic acid to L-aspartic acid by the on-line coupling of a solid-phase extraction column and a ligand-exchange high-performance liquid chromatography column 896(2000)299
- Solid-phase extraction–mass spectrometry**  
On-line coupling of solid-phase extraction with mass spectrometry for the analysis of biological samples.  
I. Determination of clenbuterol in urine 863(1999)115
- Solid-phase microextraction**  
see also Extraction methods
- Solid-phase microextraction coupled with gas chromatography–ion trap mass spectrometry for the analysis of haloacetic acids in water 859(1999)159
- Analysis of biogenic amines by solid-phase microextraction and high-performance liquid chromatography with electrochemical detection 870(2000)395
- Kinetics of solid-phase extraction and solid-phase microextraction in thin adsorbent layer with saturation sorption isotherm 873(2000)39
- Analysis of volatile contaminants in vegetable oils by headspace solid-phase microextraction with Carboxen-based fibres 873(2000)79
- Development of a headspace solid-phase microextraction procedure for the determination of free volatile fatty acids in waste waters 873(2000)107
- Screening of Brazilian fruit aromas using solid-phase microextraction–gas chromatography–mass spectrometry 873(2000)117
- Multiple solid-phase microextraction 878(2000)27
- Solid-phase microextraction for determining the distribution of sixteen US Environmental Protection Agency polycyclic aromatic hydrocarbons in water samples 879(2000)177
- Applications of solid-phase microextraction in food analysis 880(2000)35
- Evolution of solid-phase microextraction technology 885(2000)153
- Solid-phase microextraction in pesticide residue analysis 885(2000)389
- Application of solid-phase microextraction for determination of organic vapours in gaseous matrices 885(2000)405
- Application of solid-phase microextraction to the recovery of explosives and ignitable liquid residues from forensic specimens 885(2000)419
- Role of solid-phase microextraction in the identification of highly volatile pheromones of two Rhinoceros beetles *Scapanes australis* and *Strategus aloeus* (Coleoptera, Scarabaeidae, Dynastinae) 885(2000)433
- Solid-phase micro-extraction of drugs from biological matrices 885(2000)445
- Evidence for selectivity of absorption of volatile organic compounds by a polydimethylsiloxane solid-phase microextraction fibre 885(2000)457
- Solid-phase microextraction: a promising technique for sample preparation in environmental analysis 889(2000)3
- Indirect analysis of urea herbicides from environmental water using solid-phase microextraction 890(2000)303
- Influence of fibre coating in headspace solid-phase microextraction–gas chromatographic analysis of aromatic and medicinal plants 892(2000)469
- Determination of fungicides in natural waters using solid-phase microextraction and gas chromatography coupled with electron-capture and mass spectrometric detection 893(2000)143
- High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157
- Solid-phase microextraction for the determination of systemic and non-volatile pesticides in river water using gas chromatography with nitrogen–phosphorous and electron-capture detection 893(2000)347
- Characterisation of whiskeys using solid-phase microextraction with gas chromatography–mass spectrometry 896(2000)351
- Determination of polycyclic aromatic hydrocarbons in waste water by off-line coupling of solid-phase microextraction with column liquid chromatography 897(2000)153



Simultaneous determination of cyanogen chloride and cyanogen bromide in treated water at sub- $\mu\text{g/L}$  levels by a new solid-phase microextraction–gas chromatographic–electron-capture detection method 897(2000)307

Solid-phase microextraction for organochlorine pesticide residues analysis in Chinese herbal formulations 898(2000)245

#### **Solid-phase microextraction, in-tube**

Automated in-tube solid-phase microextraction–high-performance liquid chromatography for carbamate pesticide analysis 873(2000)137

#### **Solid-phase peptide synthesis**

Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor 852(1999)189

#### **Solid-phase trapping**

On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175

#### **Solid-state NMR spectroscopy**

Fullerenes separation with monomeric type  $\text{C}_{30}$  stationary phase in high-performance liquid chromatography 883(2000)55

#### **Solubility**

Application of centrifugal counter-current chromatography to the separation of macrolide antibiotic analogues. I. Selection of solvent systems based on solubility and partition coefficient investigations 864(1999)69

#### **Solute descriptors**

Revised linear solvation energy relationship coefficients for the 77-phase McReynolds data set based on an updated set of solute descriptors 878(2000)115

Calculation of Abraham solute descriptors from McReynolds gas chromatographic retention data 893(2000)339

#### **Solute migration**

Visualization of solute migration in chromatographic columns. Quantitation of the concentration in a migrating zone 867(2000)71

#### **Solvating gas chromatography**

Fast gas chromatography: packed column solvating gas chromatography versus open tubular column gas chromatography 892(2000)3

#### **Solvent–detergent method**

Ion-exchange chromatography separation of the detergent and the solvent from immunoglobulins after solvent–detergent treatment 852(1999)83

Removal of detergent and solvent from solvent–detergent-treated immunoglobulins 852(1999)87

#### **Solvent elimination**

Liquid chromatography–Fourier-transform infrared spectrometry 856(1999)213

#### **Solvent strength**

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

#### **Sol–gel**

Determination of 1-nitropyrene in herbs after selective enrichment by a sol–gel-generated immunoaffinity column 880(2000)113

Capillary electrochromatography using continuous-bed columns of sol–gel bonded silica particles with mixed-mode octadecyl and propylsulfonic acid functional groups 887(2000)265

Preparation of stationary phases for open-tubular capillary electrochromatography using the sol–gel method 887(2000)253

High-performance polyethylene glycol-coated solid-phase microextraction fibers using sol–gel technology 893(2000)157

#### **Sophora spp.**

Determination of quinolizidine alkaloids in traditional Chinese herbal drugs by nonaqueous capillary electrophoresis 857(1999)303

#### **Sorbent retention**

Contributions of theory to method development in solid-phase extraction 885(2000)17

Recent developments in polymer-based sorbents for solid-phase extraction 885(2000)51

#### **Sorbent saturation**

Kinetics of solid-phase extraction and solid-phase microextraction in thin adsorbent layer with saturation sorption isotherm 873(2000)39

#### **Sorbent trapping**

Sorbent trapping of volatile organic compounds from air 885(2000)129

#### **Sorbents**

see also Adsorbents

Solid-phase extraction: method development, sorbents, and coupling with liquid chromatography 856(1999)3

On the performance and inertness of different materials used for the enrichment of sulfur compounds from air and gaseous samples 864(1999)345

Novel use of a dual-zone restricted access sorbent: normal-phase solid-phase extraction separation of methyl oleate from polynuclear aromatic hydrocarbons stemming from semi-permeable membrane devices 883(2000)143

Solid-phase extraction of phenols 885(2000)291

Assessment of polycrystalline graphites as sorbents for solid-phase microextraction of nonionic surfactants 888(2000)35

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Solid-phase extraction with a dibutylmelamine-imprinted polymer as triazine herbicide-selective sorbent 889(2000)25

Comparison of different sorbent materials for *on-line* solid-phase extraction with liquid chromatography–atmospheric pressure chemical ionization mass spectrometry of phenols 896(2000)159

#### Source 5RPC columns

New polymer-based prepacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

#### Soxhlet extraction

Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421

#### Soybean

Analysis of bovine whey proteins in soybean dairy-like products by capillary electrophoresis 859(1999)77

Determination by perfusion reversed-phase high-performance liquid chromatography of the soybean protein content of commercial soybean products prepared directly from whole soybeans 881(2000)37

Characterization of commercial soybean products by conventional and perfusion reversed-phase high-performance liquid chromatography and multivariate analysis 881(2000)47

Quantitative determination of isoflavones and coumestrol in soybean by column liquid chromatography 886(2000)55

#### *Sparus aurata*

Extraction and isolation of linear alkylbenzene sulfonates and their intermediate metabolites from various marine organisms 889(2000)99

#### Speciation

Understanding of peak deterioration in hyphenated speciation systems due to gas–liquid separation in the hydride generation interface 855(1999)291

Speciation of selenium compounds with ion-pair reversed-phase liquid chromatography using inductively coupled plasma mass spectrometry as element-specific detection 874(2000)55

#### Speed optimisation

High-speed gas chromatography: an overview of various concepts 856(1999)315

#### Split/splitless injection

Video-taped sample evaporation in hot chambers simulating gas chromatography split/splitless injectors. I. Thermospray injection 897(2000)237

Video-taped sample evaporation in hot chambers simulating gas chromatography split and splitless injectors. II. Injection with band formation 897(2000)247

#### Split-flow injector

Split-flow injector for capillary zone electrophoresis 883(2000)223

#### Split-flow thin fractionation

Magnetic split-flow thin fractionation of magnetically susceptible particles 857(1999)193

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

#### St. John's Wort

Evaluation of major active components in St. John's Wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation 888(2000)85

#### Stability constants

Comparative study on camphor enantiomers behavior under the conditions of gas–liquid chromatography and reversed-phase high-performance liquid chromatography systems modified with  $\alpha$ - and  $\beta$ -cyclodextrins 874(2000)73

Calculation of stability constants for the chiral selector–enantiomer interactions from electrophoretic mobilities 875(2000)431

#### Stability studies

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Stability of organophosphorus insecticides on graphitized carbon black extraction cartridges used for large volumes of surface water 879(2000)73

Use of an amino acid in the mobile phase for the determination of ascorbic acid in food by high-performance liquid chromatography with electrochemical detection 881(2000)317

Stability of penicillin antibiotic residues in meat during storage. Ampicillin 882(2000)135

Analysis of isomeric glutamyl peptides by capillary electrophoresis. Application to stability studies 888(2000)275

Stability of sulfonated derivatives of benzene and naphthalene on disposable solid-phase extraction pre-columns and in an aqueous matrix 889(2000)231

**Stable isotope dilution**

Preparation of 4,4'-(1-[<sup>2</sup>H<sub>6</sub>]methylethylidene)bis-[2,3,5,6-<sup>2</sup>H<sub>4</sub>]-phenol and its application to the measurement of bisphenol A in beverages by stable isotope dilution mass spectrometry 883(2000)163

**Stacking**

New affinity nylon membrane used for adsorption of  $\gamma$ -globulin 867(2000)161

**Standard Reference Materials**

Determination of vitamins in food-matrix Standard Reference Materials 881(2000)171

**Standardization**

Standardization of capillary zone electropherograms obtained by using field enhanced sample stacking 894(2000)3

**Stationary phases**

see also Adsorbents, Affinity adsorbents, Chiral stationary phases, Ion exchangers and Sorbents

**Stationary phases, electrochromatography**

Preparation of uniformly sized polymeric separation media potentially suitable for small-scale high-performance liquid chromatography and/or capillary electrochromatography 853(1999)11

Capillary electrochromatography with macroporous particles 853(1999)45

Chromatographic properties of reversed-phase stationary phases under pressure- and electro-driven conditions 862(1999)121

Capillary electrochromatography of basic compounds using octadecyl-silica stationary phases with an amine-containing mobile phase 872(2000)315

Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography 873(2000)293

Packing columns for capillary electrochromatography 887(2000)43

Highly selective separations by capillary electrochromatography: molecular imprint polymer sorbents (Review) 887(2000)125

Comparison of aqueous and non-aqueous capillary electrochromatography for the separation of basic solutes 887(2000)233

Preparation of stationary phases for open-tubular capillary electrochromatography using the sol-gel method 887(2000)253

Capillary electrochromatography using continuous-bed columns of sol-gel bonded silica particles with mixed-mode octadecyl and propylsulfonic acid functional groups 887(2000)265

Characteristics of particle-loaded monolithic sol-gel columns for capillary electrochromatography. I. Structural, electrical and band-broadening properties 887(2000)277

Capillary electrochromatography with novel stationary phases. IV. Retention behavior of glycosphingolipids on porous and non-porous octadecyl sulfonated silica 887(2000)287

Stationary phase structures enhancing electroosmotic flow in capillary electrochromatography 887(2000)299

Stationary phases for capillary electrochromatography (Review) 887(2000)313

Towards the column bed stabilization of columns in capillary electroosmotic chromatography. Immobilization of microparticulate silica columns to a continuous bed 887(2000)327

Aspects of column fabrication for packed capillary electrochromatography 887(2000)347

Pressurized-flow anion-exchange capillary electrochromatography using a polymeric ion-exchange stationary phase 890(2000)337

Nonaqueous electrochromatography on continuous bed columns of sol-gel bonded large-pore C<sub>18</sub> material: separation of retinyl esters 890(2000)347

Advances in column technology and instrumentation in capillary electrochromatography 892(2000)257

Capillary electrochromatography on silica columns: factors influencing performance 892(2000)279

**Stationary phases, GC**

Synthesis and properties of two new liquid crystals: an analytical and thermodynamic study 859(1999)59

Synthesis and analytical properties of a novel columnar metallomesogenic polymer 862(1999)65

Gas chromatography for in situ analysis of a cometary nucleus: characterization and optimization of diphenyl/dimethylpolysiloxane stationary phases 863(1999)157

Synthesis and characteristics of [60]fullerene polysiloxane stationary phase for capillary gas chromatography 867(2000)177

Insoluble  $\beta$ -cyclodextrin polymer for capillary gas chromatographic separation of enantiomers and isomers 883(2000)137

Selectivity equivalence of poly(ethylene glycol) stationary phases for gas chromatography 898(2000)211

**Stationary phases, LC**

Network modeling of the convective flow and diffusion of molecules adsorbing in monoliths and in porous particles packed in a chromatographic column 852(1999)3

Retention behavior of very large biomolecules in ion-exchange chromatography 852(1999)25

Oriented immobilization of chymotrypsin by use of suitable antibodies coupled to a nonporous solid support 852(1999)141

New polymer-based prepacked column for the reversed-phase liquid chromatographic separation of peptides over the pH range 2–12 852(1999)255

Separation of oligonucleotides on novel monolithic columns with ion-exchange functional surfaces 852(1999)297

Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375

Separation and characterization of octylphenol ethoxylate surfactants used by reversed-phase high-performance liquid chromatography on branched fluorinated silica gel columns 852(1999)475

Preparation of uniformly sized polymeric separation media potentially suitable for small-scale high-performance liquid chromatography and/or capillary electrochromatography 853(1999)11

Capillary electrochromatography with macroporous particles 853(1999)45

Chiral separations by open tubular capillary electrokinetic chromatography 853(1999)151

Electrokinetically-driven cation-exchange chromatography of proteins and its comparison with pressure-driven high-performance liquid chromatography 853(1999)243

Separation of selected peptides by capillary electroosmotic chromatography using 3  $\mu\text{m}$  reversed-phase bonded silica and mixed-mode phases 853(1999)263

Macroporous poly(glycidyl methacrylate–triallyl isocyanurate–divinylbenzene) matrix as an anion-exchange resin for protein adsorption 855(1999)129

Capillary electrochromatography with a silica column with a dynamically modified cationic surfactant 855(1999)137

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. III. Results obtained with Kromasil C<sub>18</sub> columns 855(1999)423

Molecular mechanism of retention in reversed-phase high-performance liquid chromatography and classification of modern stationary phases by using quantitative structure–retention relationships 855(1999)455

Determination of inorganic anions by ion chromatography using a graphitized carbon column dynamically coated with cetyltrimethylammonium ions 855(1999)261

*n*-Alkyl fluorenyl phases in chromatography. II. Dynamic behavior and high-performance liquid chromatography applications 858(1999)133

Shape-selective separation of polycyclic aromatic hydrocarbons on protoporphyrin-silica phases. Effect of surface porphyrin distribution on column efficiency 859(1999)121

Chromatographic properties of reversed-phase stationary phases under pressure- and electro-driven conditions 862(1999)121

Reversed-phase liquid chromatographic behavior of the mycotoxins citrinin and ochratoxin A 862(1999)147

Pressure-controlled high-performance liquid chromatographic study on the influence of rim chemistry on partial molar volume differences between free and complexed cyclodextrins 863(1999)127

Determination of the thermodynamic contribution to peak asymmetry of basic solutes in reversed-phase liquid chromatography 865(1999)97

Rapid reversed-phase separation of proteins and peptides using optimized 'moulded' monolithic poly(styrene–*co*-divinylbenzene) columns 865(1999)169

Reversed-phase liquid chromatography of proteins and peptides using multimodal copolymer-encapsulated silica 866(2000)1

Enhanced performance of expanded bed chromatography on rigid superporous adsorbent matrix 867(2000)113

Effect of stationary phase hydrophobicity and mobile phase composition on the separation of carboxylic acids in ion chromatography 867(2000)131

Studies of retention and stability of a horizontally polymerized bonded phase for reversed-phase liquid chromatography 868(2000)153

Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C<sub>8</sub> and reversed-phase C<sub>18</sub> types 869(2000)57

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by  $\gamma$ -radiation on titanium-grafted silica 869(2000)137

- Development of a new cyano-bonded column for high-performance liquid chromatography 869(2000)143
- Characterization of an avidin-bonded column for direct injection in reversed-phase high-performance liquid chromatography 869(2000)151
- Optimized stationary phases for the high-performance liquid chromatography–electrospray ionization mass spectrometric analysis of basic pharmaceuticals 869(2000)159
- Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. IV. Results obtained with Luna C<sub>18</sub> (2) columns 869(2000)181
- Control of column temperature in reversed-phase liquid chromatography 869(2000)211
- Ion-pair chromatography on a porous graphitic carbon stationary phase for the analysis of twenty underivatized protein amino acids 870(2000)245
- Separation of polyesters by gradient reversed-phase high-performance liquid chromatography on a 1.5 μm non-porous column 871(2000)243
- Chromatographic resolution of tryptophan enantiomers with L-Leu–L-Leu–L-Leu peptide. Effects of mobile phase composition and chromatographic support 874(2000)21
- Characterization of C<sub>18</sub>-bonded liquid chromatographic stationary phases by Raman spectroscopy: The effect of mobile phase composition 877(2000)25
- Characterization of C<sub>18</sub>-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41
- Separation of aromatic isomers on cyclophane-bonded stationary phases 877(2000)61
- Retention versatility of silica-supported comb-shaped crystalline and non-crystalline phases in high-performance liquid chromatography 877(2000)71
- C<sub>30</sub> Stationary phases for the analysis of food by liquid chromatography 880(2000)189
- Hydrodynamic aspects of slurry packing processes in microcolumn liquid chromatography 883(2000)11
- New stationary phase for liquid chromatography with chemically bonded pinane ligand: synthesis and characterization by nuclear magnetic resonance and high-performance liquid chromatographic investigations 883(2000)27
- Pore size distributions of cation-exchange adsorbents determined by inverse size-exclusion chromatography 883(2000)39
- Fullerenes separation with monomeric type C<sub>30</sub> stationary phase in high-performance liquid chromatography 883(2000)55
- Ion chromatographic selectivity of polyelectrolyte sorbents based on some aliphatic and aromatic ionenes 884(2000)31
- Application of neutral hydrophobic hypercrosslinked polystyrene to the separation of inorganic anions by ion chromatography 884(2000)41
- Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113
- Ion-chromatographic behavior of alkali metal cations and ammonium ion on zirconium-adsorbing silica gel 884(2000)123
- Suppressed electrostatic ion chromatography with tetraborate as eluent and its application to the determination of inorganic anions in snow and rainwater 884(2000)161
- Elution mechanism in electrostatic ion chromatography with histidine as an isoelectric ampholytic mobile phase 884(2000)287
- Low-capacity cation-exchange chromatography of ultraviolet-absorbing urinary basic metabolites using a reversed-phase column coated with hexadecylsulfonate 886(2000)297
- Capillary electrochromatography of proteins and peptides with a cationic acrylic monolith 887(2000)465
- Antibody immobilization to high-performance liquid chromatography supports. Characterization of maximum loading capacity for intact immunoglobulin G and Fab fragments 888(2000)13
- High-speed recovery of germanium in a convection-aided mode using functional porous hollow-fiber membranes 888(2000)43
- Behavior of lanthanides in countercurrent chromatography using dihexyl-*N,N*-diethylcarbamoyl methylene phosphonate as stationary phase 888(2000)137
- Investigations on the behaviour of acidic, basic and neutral compounds in capillary electrochromatography on a mixed-mode stationary phase 888(2000)267
- New configurations and applications of molecularly imprinted polymers 889(2000)15
- Superficially porous silica microspheres for fast high-performance liquid chromatography of macromolecules 890(2000)3
- Use of spray-dried zirconia microspheres in the separation of immunoglobulins from cell culture supernatant 890(2000)15

- Discoloration of ceramic hydroxyapatite used for protein chromatography 891(2000)93
- Synthesis of spherical porous silicas in the micron and submicron size range: challenges and opportunities for miniaturized high-resolution chromatographic and electrokinetic separations 892(2000)47
- Chromatographic silanol activity test procedures: the quest for a universal test 892(2000)57
- Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435
- Preparation and evaluation of packed capillary columns for the separation of nucleic acids by ion-pair reversed-phase high-performance liquid chromatography 893(2000)23
- Characterization of reversed-phase columns using the linear free energy relationship. III. Effect of the organic modifier and the mobile phase composition 893(2000)215
- Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy 893(2000)235
- Dependence of cyano bonded phase hydrolytic stability on ligand structure and solution pH 893(2000)245
- Comparison of phenyl-type columns in the development of a fast liquid chromatographic system for eighteen opiates commonly found in forensic toxicology 896(2000)311
- Optimisation and characterisation of silica-based reversed-phase liquid chromatographic systems for the analysis of basic pharmaceuticals 897(2000)1
- Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. I. Optimizing selectivity and resolution 897(2000)37
- Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. II. Minimizing column reproducibility problems 897(2000)51
- Protein adsorption on novel acrylamido-based polymeric ion-exchangers. I. morphology and equilibrium adsorption 897(2000)65
- Protein adsorption on novel acrylamido-based polymeric ion exchangers. II. Adsorption rates and column behavior 897(2000)81
- Comparison of chromatographic ion-exchange resins. I. Strong anion-exchange resins 897(2000)99
- Laterally attached liquid crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. II. Optimization of the molecular parameter of the polymer 897(2000)131
- Reversed-phase high-performance liquid chromatography on porous copolymers of different chemical structure 898(2000)13
- Phenylaminopropyl silica – a new specific stationary phase for high-performance liquid chromatography of phenols 898(2000)23
- New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent 898(2000)35
- Statistical analysis**  
Spectrophotometric and coulometric detection in the high-performance liquid chromatography of flavonoids and optimization of sample treatment for the determination of quercetin in orange juice 881(2000)449
- Ion-chromatographic study of the possible absorption of copper and zinc by the skin of *Rana pipiens* 884(2000)143
- Ion-chromatographic assay of chromium(VI)-containing semiconductor etchants. Statistical comparison of two columns 884(2000)151
- Statistical evaluation of linear solvation energy relationship models used to characterize chemical selectivity in micellar electrokinetic chromatography 886(2000)245
- Statistical optimization**  
Statistics aided optimization for high-performance liquid chromatographic analysis of organic acids in tobacco 859(1999)153
- Steady-state recycling**  
Modeling of separations by closed-loop steady-state recycling chromatography of a racemic pharmaceutical intermediate 867(2000)1
- Steam distillation–extraction**  
Determination of phenols using simultaneous steam distillation–extraction 867(2000)281
- Stepwise capillary electrophoresis**  
Stepwise capillary electrophoretic separation of DNA fragments using poly(ethylene oxide) solutions in the presence of electroosmotic flow 853(1999)337
- Stereoisomer separation**  
Reproducibility of the separation of astaxanthin stereoisomers on Pirkle covalent L-leucine and D-phenylglycine columns 855(1999)157
- Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107
- Stereoselective metabolism**  
Enantioselective determination of drugs in body fluids by capillary electrophoresis 875(2000)27

**Steric mass action**

Optimization of ion-exchange displacement separations. I. Validation of an iterative scheme and its use as a methods development tool 876(2000)51

Optimization of ion-exchange displacement separations. II. Comparison of displacement separations on various ion-exchange resins 876(2000)63

**Stoichiometric displacement model**

Reversed-phase high-performance liquid chromatography of ionogenic compounds: comparison of retention models 859(1999)1

**Stopped-flow multidimensional gas chromatography**

Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487

***Strategus aloeus***

Role of solid-phase microextraction in the identification of highly volatile pheromones of two Rhinoceros beetles *Scapanes australis* and *Strategus aloeus* (Coleoptera, Scarabaeidae, Dynastinae) 885(2000)433

**Streamline chelating**

Immobilised metal affinity chromatography of  $\beta$ -galactosidase from unclarified *Escherichia coli* homogenates using expanded bed adsorption 874(2000)27

**Structure–activity relationships**

Microscale characterization of the structure–activity relationship of a heparin-binding glycopeptide using affinity capillary electrophoresis and immobilized enzymes 853(1999)189

**Structure–chromatographic parameters relationships**

Modeling chromatographic parameters by a novel graph theoretical sub-structural approach 858(1999)187

**Structure–mobility relationships**

Studies on the relationship between structure and electrophoretic mobility of  $\alpha$ -helical and  $\beta$ -sheet peptides using capillary zone electrophoresis 857(1999)263

**Structure–retention relationships**

Revision of a theoretical expression for gas–liquid chromatographic retention 852(1999)583

Molecular mechanism of retention in reversed-phase high-performance liquid chromatography and classification of modern stationary phases by using quantitative structure–retention relationships 855(1999)455

Relationship between the retention characteristics on an alumina column and physico-chemical parameters of some environmental pollutants 869(2000)41

Stereoselective binding of 2,3-substituted 3-hydroxypropionic acids on an immobilised human serum albumin chiral stationary phase: stereochemical characterisation and quantitative structure–retention relationship study 876(2000)75

Exploratory chemometric analysis of the classification of pharmaceutical substances based on chromatographic data 897(2000)23

Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227

**Subcritical fluid chromatography**

Study of dead volume measurement in packed subcritical fluid chromatography with ODS columns and carbon dioxide–modifier mobile phases 866(2000)241

Retention behaviour of ceramides in sub-critical fluid chromatography in comparison with non-aqueous reversed-phase liquid chromatography 883(2000)211

**Subcritical water extraction**

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

On-line coupling of subcritical water extraction with high-performance liquid chromatography via solid-phase trapping 873(2000)175

Static subcritical water extraction with simultaneous solid-phase extraction for determining polycyclic aromatic hydrocarbons on environmental solids 886(2000)237

Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421

Extraction of cloransulam-methyl from soil with subcritical water and supercritical CO<sub>2</sub> 897(2000)405

**Successive multiple ionic–polymer layer coated capillaries**

Enantiomeric separation by capillary electrophoresis with an electroosmotic flow-controlled capillary 875(2000)315

**Sugar cane**

Solid-phase extraction of sugar cane soot extract for analysis by gas chromatography with flame ionisation and mass spectrometric detection 889(2000)281

**Sugarcane juices**

Changes in free and conjugated polyamines during starvation of sugarcane juices as analyzed by high-performance liquid chromatography 881(2000)531

**Sulfonic acid bonded phase capillaries**

Applications of a sulfonated-polymer wall-modified open-tubular fused-silica capillary in capillary zone electrophoretic separations 884(2000)297

**Supercritical fluid chromatography**

Electrospray mass spectrometry and supercritical fluid chromatography of methylated  $\beta$ -cyclodextrins 855(1999)645

Supercritical fluids in separation science – the dreams, the reality and the future 856(1999)83

Separation of stereoisomers in a simulated moving bed-supercritical fluid chromatography plant 865(1999)175

**Supercritical fluid chromatography–mass spectrometry**

see also Mass spectrometry and Special Issue Index

Capillary column supercritical fluid chromatography–atmospheric pressure ionisation mass spectrometry. Interface performance of atmospheric pressure chemical ionisation and electrospray ionisation 855(1999)317

**Supercritical fluid derivatization and extraction**

Simultaneous supercritical fluid derivatization and extraction of formaldehyde by the Hantzsch reaction 896(2000)51

**Supercritical fluid extraction**

see also Extraction methods

Quantitative determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls and organochlorine pesticides in sewage sludges using supercritical fluid extraction and mass spectrometric detection 852(1999)545

Supercritical fluid extraction of nylon 6,6 oligomers and their characterization via liquid chromatography coupled with mass spectrometry 855(1999)715

Supercritical fluids in separation science – the dreams, the reality and the future 856(1999)83

Esterification of decanoic acid during supercritical fluid extraction employing either methanol-modified carbon dioxide or a methanol trap 858(1999)201

Comparison of methods for quantitative analysis of additives in low-density polyethylene using supercritical fluid and enhanced solvent extraction 858(1999)209

Simultaneous extraction and derivatization of 2-chlorovinylarsonous acid from soils using supercritical and pressurized fluids 888(2000)327

Comparison of two extraction methods independently developed on two conceptually different automated supercritical fluid extraction systems for the determination of polychlorinated biphenyls in sediments 891(2000)195

Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: recovery, selectivity and effects on sample matrix 892(2000)421

Supercritical fluid extraction of hydrocarbons and 2-alkylcyclobutanones for the detection of irradiated foodstuffs 897(2000)259

Extraction of cloransulam-methyl from soil with subcritical water and supercritical CO<sub>2</sub> 897(2000)405

**Supercritical fluids**

Supercritical fluids in separation science – the dreams, the reality and the future 856(1999)83

**Superheated water chromatography**

Selective deuterium exchange during superheated heavy water chromatography–nuclear magnetic resonance spectroscopy–mass spectrometry of sulfonamides 886(2000)289

**Supported liquid membrane extraction**

Extraction of glyphosate by a supported liquid membrane technique 889(2000)93

Determination of thiophanate-methyl and its metabolites at trace level in spiked natural water using the supported liquid membrane extraction and the microporous membrane liquid–liquid extraction techniques combined on-line with high-performance liquid chromatography 893(2000)123

**Surface characteristics**

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

**Surface composition**

Unexpected surface chemistry in capillaries for electrophoresis 894(2000)73

Unexpected surface chemistry in capillaries for electrophoresis 894(2000)73

**Surface coverage**

Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375

Studies of retention and stability of a horizontally polymerized bonded phase for reversed-phase liquid chromatography 868(2000)153

**Surface heterogeneity**

Effect of surface heterogeneity on adsorption on solid surfaces. Application of inverse gas chromatography in the studies of energetic heterogeneity of adsorbents 886(2000)133

**Surface ionization detection**

Surface-ionization methods and devices of indication and identification of nitrogen-containing base molecules 896(2000)3

**Surface plasmon resonance detection**

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59



Erratum to: "Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes". [J. Chromatogr. A, 852 (1999) 59-71] 872(2000)323

#### Surface treatment

Moderation of the electroosmotic flow in capillary electrophoresis by chemical modification of the capillary surface with tentacle-like oligourethanes 894(2000)79

#### Surfactant-mediated extraction

Alkyltrimethylammonium surfactant-mediated extractions: Characterization of surfactant-rich and aqueous layers, and extraction performance 877(2000)87

#### Suspended bed chromatography

Suspended bed chromatography, a new approach in downstream processing 890(2000)45

#### Swing injection system

Introduction of large volumes of water-containing samples into a gas chromatograph. Improved retention of volatile solutes through the swing system 876(2000)135

#### Synthesis

Solid-phase extraction for combinatorial libraries 885(2000)305

## T

#### Tandem mass spectrometry

see also Mass spectrometry and Special Issue Index

Quantitative analysis of mutagenic heterocyclic aromatic amines in cooked meat using liquid chromatography–atmospheric pressure chemical ionisation tandem mass spectrometry 883(2000)89

On-line dual-precolumn-based trace enrichment for the determination of polar and acidic microcontaminants in river water by liquid chromatography with diode-array UV and tandem mass spectrometric detection 885(2000)377

#### Tapers

Improved column preparation and performance in capillary electrochromatography 887(2000)367

#### *Taxus wallichiana*

Analysis of taxol and major taxoids in Himalayan yew, *Taxus wallichiana* 858(1999)239

#### Tea

High-performance liquid chromatographic determination of  $\Delta^9$ -tetrahydrocannabinol and the corresponding acid in hemp containing foods with special regard to the fluorescence properties of  $\Delta^9$ -tetrahydrocannabinol 872(2000)101

Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography 876(2000)235

Electrophoretic methods in the analysis of beverages 880(2000)243

Determination of tea catechins 881(2000)411

Analysis of tea components by high-performance liquid chromatography and high-performance capillary electrophoresis 881(2000)425

Comparative analysis of tea catechins and theaflavins by high-performance liquid chromatography and capillary electrophoresis 881(2000)439

Effect of sample handling on alkaloid and mineral content of aqueous extracts of greater celandine (*Chelidonium majus* L.) 889(2000)69

Supercritical fluid extraction of catechins from *Cratogeomys prunifolium* Dyer and subsequent purification by high-speed counter-current chromatography 898(2000)75

#### Temperature effects

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. I. Peak capacity limitations 857(1999)1

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. II. Two-run assay procedures 857(1999)21

Reversed-phase liquid chromatographic separation of complex samples by optimizing temperature and gradient time. III. Improving the accuracy of computer simulation 857(1999)41

Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. I. Optimizing selectivity and resolution 897(2000)37

Selectivity differences for C<sub>18</sub> and C<sub>8</sub> reversed-phase columns as a function of temperature and gradient steepness. II. Minimizing column reproducibility problems 897(2000)51

Retention and separation studies of cholesterol and bile acids using thermostated thin-layer chromatography 857(1999)255

Temperature-induced inversion of elution order in the enantioseparation of sotalol on a cellobiohydrolase I-based stationary phase 858(1999)33

Analysis of the equations for the temperature dependence of the retention index. II. Physico-chemical meaning of the parameters 859(1999)49

Ion exchange on resins with temperature-responsive selectivity. III. Influence of complex formation stoichiometry on temperature dependence of resin selectivity 868(2000)143

Low-energy interactions in high-performance liquid chromatography 872(2000)49

Characterization of C<sub>18</sub>-bonded liquid chromatographic stationary phases by Raman spectroscopy: the effect of temperature 877(2000)41

Fullerenes separation with monomeric type C<sub>30</sub> stationary phase in high-performance liquid chromatography 883(2000)55

Investigation into the effects of temperature and stirring rate on the solid-phase extraction of diuron from water using a C<sub>18</sub> extraction disk 885(2000)41

High temperature and temperature programming in capillary electrochromatography 887(2000)245

Reversed-phase separation of achiral isomers by varying temperature and either gradient time or solvent strength 892(2000)107

Effects of DNA topology, temperature and solvent viscosity on DNA retardation in slalom chromatography 893(2000)115

Influence of temperature on the behaviour of small linear peptides in capillary electrochromatography 893(2000)401

### Temperature programming

Determination of low levels of an antioxidant in polyolefins by large-volume injection temperature-programmed packed capillary liquid chromatography 864(1999)103

Hindered amine stabilizers investigated by the use of packed capillary temperature-programmed liquid chromatography. I. Poly(6-(1,1,3,3-tetramethylbutyl)-amino)-1,3,5-triazine-2,4-diyl)(2,2,6,6-tetramethyl-4-piperidyl)imino)-1,6-hexanedyl(2,2,6,6-tetramethyl-4-piperidyl)imino) 874(2000)65

High temperature and temperature programming in capillary electrochromatography 887(2000)245

Determination of 1-(2-methoxyphenyl)piperazine derivatives of isocyanates at low concentrations by temperature-programmed miniaturized liquid chromatography 892(2000)67

### Tenax

Surface characterization of industrial fibers with inverse gas chromatography 888(2000)175

### Test atmosphere systems

Dynamic system for the calibration of semi-volatile carbonyl compounds in air 883(2000)171

### Textiles

High-performance liquid chromatographic determination of some anthraquinone and naphthoquinone dyes occurring in historical textiles 863(1999)235

### Thermal conductivity detection

Prediction of thermal conductivity detection response factors using an artificial neural network 897(2000)227

### Thermal degradation

Determination of basic nitrogen-containing polynuclear aromatic hydrocarbons formed during thermal degradation of polymers by high-performance liquid chromatography–fluorescence detection 878(2000)171

### Thermal evaporation

Gas chromatographic–mass spectrometric separation, identification and determination of C<sub>6</sub>–C<sub>12</sub> cyclic imides in thermally treated epoxy and alkyd resins to locate hot spots inside large electricity generators 862(1999)17

### Thermal lens detection

Integration of a microextraction system. Solvent extraction of a Co–2-nitroso-5-dimethylaminophenol complex on a microchip 894(2000)19

Non-contact photothermal control of enzyme reactions on a microchip by using a compact diode laser 894(2000)45

### Thermodynamic parameters

Comparison between the isocratic and gradient retention behaviour of polypeptides in reversed-phase liquid chromatographic environments 852(1999)43

Revision of a theoretical expression for gas–liquid chromatographic retention 852(1999)583

Temperature-induced inversion of elution order in the enantioseparation of sotalol on a cellobiohydrolase I-based stationary phase 858(1999)33

Analysis of the equations for the temperature dependence of the retention index. I. Relation between equations 858(1999)65

Analysis of the equations for the temperature dependence of the retention index. II. Physico-chemical meaning of the parameters 859(1999)49

Synthesis and properties of two new liquid crystals: an analytical and thermodynamic study 859(1999)59

Heats of adsorption of some organic compounds on β-cyclodextrin determined by gas–solid chromatography 864(1999)293

Determination of the thermodynamic contribution to peak asymmetry of basic solutes in reversed-phase liquid chromatography 865(1999)97

Study of hydrophobic interaction adsorption of bovine serum albumin under overloaded conditions using flow microcalorimetry 865(1999)111

Chiral separation of four 1,3-dioxolane derivatives by supercritical fluid chromatography on an amylose-based column 871(2000)127

Microcalorimetric studies of interactions between proteins and hydrophobic ligands in hydrophobic interaction chromatography: effects of ligand chain length, density and the amount of bound protein 872(2000)37

Interpreting the gas chromatographic retention of *n*-alkanes 873(2000)209

Characterization of partition and thermodynamic properties of cephalosporins using micellar electrokinetic chromatography in glycodeoxycholic acid solution 873(2000)237

Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography 875(2000)179

Distribution coefficients of *n*-alkanes measured on wall-coated capillary columns 879(2000)157

Enantioselective stopped-flow multidimensional gas chromatography. Determination of the inversion barrier of 1-chloro-2,2-dimethylaziridine 892(2000)487

Gas chromatographic study of the hydrogen bonding of aliphatic alcohols to tri-*n*-octylphosphine oxide 898(2000)103

#### **Thermo-optical absorbance detection**

Capillary electrochromatography with thermo-optical absorbance detection for the analysis of phenylthiohydantoin-amino acids 853(1999)131

Non-aqueous capillary electrophoresis of tamoxifen and its acid hydrolysis products 895(2000)81

#### **Thermospray injection**

Video-taped sample evaporation in hot chambers simulating gas chromatography split/splitless injectors. I. Thermospray injection 897(2000)237

#### **Thermostripping**

Ion exchange on resins with temperature-responsive selectivity. IV. Influence of solution and column parameters on efficiency of reagentless separation of copper and zinc using thermo-induced concentration waves technique 867(2000)57

#### **Thin-layer chromatography**

see also Planar chromatography

Planar chromatography at the turn of the century 856(1999)399

Thin-layer chromatography in food and agricultural analysis 880(2000)129

Practitioner's guide to method development in thin-layer chromatography 892(2000)123

#### ***Thlaspi caerulescens***

Determination of glucosinolates in rapeseed and *Thlaspi caerulescens* plants by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 889(2000)75

#### **Three-dimensional capillary electrophoresis**

Novel three-dimensional capillary electrophoresis system for complex and trace analysis 894(2000)117

#### ***Thymus mastichina***

Comparison of continuous subcritical water extraction and hydrodistillation of marjoram essential oil 855(1999)625

#### **Time-of-flight mass spectrometry**

Liquid chromatography–electrospray time-of-flight mass spectrometry for on-line accurate mass determination and identification of cyclodepsipeptides in a crude extract of the fungus *Metarrhizium anisopliae* 872(2000)85

#### **Time-weighted average sampling**

Time-weighted average sampling of volatile and semi-volatile airborne organic compounds by the solid-phase microextraction device 892(2000)455

#### **Titania**

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

#### **Titanium-grafted silica**

Effects in high-performance liquid chromatography of a high pH in the mobile phase on poly(methyloctylsiloxane) immobilized by  $\gamma$ -radiation on titanium-grafted silica 869(2000)137

#### **Tobacco**

see also Cigarette smoke

Optimization study for the reversed-phase ion-pair liquid chromatographic determination of nicotine in commercial tobacco products 852(1999)451

Application of non-aqueous capillary electrophoresis with electrochemical detection to the determination of nicotine in tobacco 853(1999)27

Determination of polyphenols by high-performance liquid chromatography with inhibited chemiluminescence detection 855(1999)171

Solid phase microextraction of alkenylbenzenes and other flavor-related compounds from tobacco for analysis by selected ion monitoring gas chromatography–mass spectrometry 858(1999)79

Statistics aided optimization for high-performance liquid chromatographic analysis of organic acids in tobacco 859(1999)153

Gas-liquid chromatographic determination of major tobacco alkaloids 898(2000)235

#### Topological indices

Modeling chromatographic parameters by a novel graph theoretical sub-structural approach 858(1999)187

#### Total organic carbon

Efficient solid-phase extraction procedures for trace enrichment of priority phenols from industrial effluents with high total organic carbon content 857(1999)97

#### Toxicological analysis

Fractionation and toxicity evaluation of waste waters 889(2000)149

#### Transdermal patch

Headspace gas chromatography method for the analysis of volatile impurities in a hormone replacement transdermal patch 859(1999)113

#### Transducers

Capillary electrophoresis coupled to biosensor detection 892(2000)143

#### Transient state

Transient state of chiral recognition in a binary mixture of cyclodextrins in capillary electrophoresis 888(2000)241

#### Trapping

Sorbent trapping of volatile organic compounds from air 885(2000)129

#### Triazine stationary phase

Synthesis of a silica-bonded bovine serum albumin *s*-triazine chiral stationary phase for high-performance liquid chromatographic resolution of enantiomers 866(2000)173

#### *Trichoderma harzianum*

Colour pigments of *Trichoderma harzianum*. Preliminary investigations with thin-layer chromatography–Fourier transform infrared spectroscopy and high-performance liquid chromatography with diode array and mass spectrometric detection 896(2000)61

#### *Trichoderma reesei*

Studies on the chromatographic fractionation of *Trichoderma reesei* cellulases by hydrophobic interaction 865(1999)123

#### *Trichoderma viride*

Detection of new sequences of peptaibol antibiotics trichotoxins A-40 by on-line liquid chromatography–electrospray ionization mass spectrometry 862(1999)179

#### Trimethylsilyl groups

Equation for calculating surface coverage from end-capping of chromatographic bonded phases 852(1999)375

#### *Triticum aestivum*

Simultaneous determination of phenolic acids and 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one in wheat (*Triticum aestivum* L.) by gas chromatography–tandem mass spectrometry 864(1999)315

#### Troubleshooting

Investigations of artifact peaks in sensitive high-performance liquid chromatography methods 869(2000)85

#### True moving beds

Optimization of simulated moving bed plants with low efficient stationary phases: separation of fructose and glucose 868(2000)169

#### Turbulent flow chromatography

Comparison of turbulent-flow chromatography with automated solid-phase extraction in 96-well plates and liquid–liquid extraction used as plasma sample preparation techniques for liquid chromatography–tandem mass spectrometry 854(1999)23

#### Two-dimensional liquid chromatography

Protein mapping by two-dimensional high-performance liquid chromatography 893(2000)293

## U

#### Ultrasonic irradiation

Evaluation of the secondary consolidation of columns for liquid chromatography by ultrasonic irradiation 878(2000)153

Effect of linear polymer additives on the electroosmotic characteristics of agarose gels in ultrathin-layer electrophoresis 853(1999)511

High-throughput genotyping of factor V Leiden mutation by ultrathin-layer agarose gel electrophoresis 853(1999)519

**Ultra-thin-layer sodium dodecylsulfate gel electrophoresis**

Rapid analysis of covalently and non-covalently fluorophore-labeled proteins using ultra-thin-layer sodium dodecylsulfate gel electrophoresis 894(2000)329

**UV detection**

Identification of compounds and specific functional groups in the wavelength region 168–330 nm using gas chromatography with UV detection (2000)187

**V****Vaccines**

Determination of free carrier protein in protein–polysaccharide conjugate vaccines by micellar electrokinetic chromatography 894(2000)311

**Validation**

Analysis of anions in aqueous samples by ion chromatography and capillary electrophoresis. A comparative study of peak modeling and validation criteria 852(1999)487

Development and validation of a capillary electrophoresis method for the characterization of protegrin IB-367 853(1999)197

Elimination of adsorption effects in the analysis of water-soluble vitamins in pharmaceutical formulations by capillary electrophoresis 853(1999)391

Comparison of turbulent-flow chromatography with automated solid-phase extraction in 96-well plates and liquid–liquid extraction used as plasma sample preparation techniques for liquid chromatography–tandem mass spectrometry 854(1999)23

Chromatographic separation of 3,4-difluorophenylacetic acid and its positional isomers using five different techniques 866(2000)281

Use of optimization software to determine rugged analysis conditions in high-performance liquid chromatography 868(2000)13

Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463

Determination of *N*-acetylcysteine in human plasma by liquid chromatography coupled to tandem mass spectrometry 870(2000)13

Full and fractionated experimental designs for robustness testing in the high-performance liquid chromatographic analysis of codeine phosphate, pseudoephedrine hydrochloride and chlorpheniramine maleate in a pharmaceutical preparation 870(2000)45

Validation of the removal of acetylsalicylic acid. Recovery and determination of residues on various surfaces by high performance liquid chromatographic 870(2000)69

Simultaneous determination of paracetamol and chlorpheniramine in human plasma by liquid chromatography–tandem mass spectrometry 870(2000)77

Optimization and validation of a method for the determination of caffeine, 8-chlorotheophylline and diphenhydramine by isocratic high-performance liquid chromatography. Stress test for stability evaluation 870(2000)97

Quantitation of nifedipine in human plasma by on-line solid-phase extraction and high-performance liquid chromatography 870(2000)115

Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated  $\beta$ -cyclodextrin 875(2000)389

Analysis of polymyxin B sulfate by capillary zone electrophoresis with cyclodextrin as additive. Method development and validation 879(2000)211

Development and validation of a capillary electrophoresis method for direct measurement of isocitric, citric, tartaric and malic acids as adulteration markers in orange juice 881(2000)395

Quantitation of nine quinolones in chicken tissues by high-performance liquid chromatography with fluorescence detection 882(2000)63

Validation of a capillary electrophoresis method for determination of 5-aminolevulinic acid and degradation products 883(2000)285

Performance evaluation of a method for the determination of bromate in drinking water by ion chromatography (EPA Method 317.0) and validation of EPA Method 324.0 884(2000)201

Validation of an analytical procedure for polychlorinated biphenyls, coplanar polychlorinated biphenyls and polycyclic aromatic hydrocarbons in environmental samples 886(2000)153

Determination of clonixin in plasma and urine by reversed-phase high-performance liquid chromatography 889(2000)135

Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the ng l<sup>-1</sup> concentration level 893(2000)367

Development and validation of a capillary zone electrophoretic method for the determination of bisphosphonate and phosphonate impurities in clodronate 893(2000)411

Development and validation of a capillary electrophoresis method for the measurement of short-chain organic acids in natural rubber latex 894(2000)135

Micellar electrokinetic and high-performance liquid chromatographic determination of potential manufacturing impurities in pholcodine 895(2000)9

Analysis of doxycycline by capillary electrophoresis. Method development and validation 895(2000)43

Analysis of 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-fluorophenyl)-*N*-(3-iodo-*E*-allyl)nortropine in rat plasma. I. Method development and validation by capillary electrophoresis 895(2000)87

Quantitative analysis of *N*-acetylcysteine and its pharmacopeial impurities in a pharmaceutical formulation by liquid chromatography–UV detection–mass spectrometry 896(2000)191

Determination of deramciclone and *N*-desmethylderamciclone in human plasma by liquid chromatography–tandem mass spectrometry using off-line robotic sample pretreatment 896(2000)279

Optimization strategy and validation of one chromatographic method as approach to determine the phenolic compounds from different sources 897(2000)161

#### Vaporising injection

Video-taped sample evaporation in hot chambers simulating gas chromatography split/splitless injectors. I. Thermospray injection 897(2000)237

Video-taped sample evaporation in hot chambers simulating gas chromatography split and splitless injectors. II. Injection with band formation 897(2000)247

#### Vegetable juices

Separation and identification of various carotenoids by C<sub>30</sub> reversed-phase high-performance liquid chromatography coupled to UV and atmospheric pressure chemical ionization mass spectrometric detection 854(1999)37

#### Vegetable oils

see also Oils

Non-volatile products of triolein produced at frying temperatures characterized using liquid chromatography with online mass spectrometric detection 852(1999)417

Determination of tocopherols and sterols in vegetable oils by solid-phase extraction and subsequent capillary gas chromatographic analysis 857(1999)231

Authentication of vegetable oils by chromatographic techniques 881(2000)93

Chromatographic analysis of minor constituents in vegetable oils 881(2000)131

Automated analysis of vitamin E isomers in vegetable oils by continuous membrane extraction and liquid chromatography–electrochemical detection 881(2000)229

Solid-phase microextraction with on-fibre derivatisation applied to the analysis of volatile carbonyl compounds 886(2000)175

Microwave-assisted derivatization of volatile carbonyl compounds with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine 891(2000)267

#### Vegetables

Determination of folate vitamers in food and in Italian reference diet by high-performance liquid chromatography 855(1999)237

Inclusion of the fungicide chlorzoxinate in a multiresidue method 858(1999)123

Determination of carbamate residues in fruits and vegetables by matrix solid-phase dispersion and liquid chromatography–mass spectrometry 871(2000)43

Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables 881(2000)309

Pesticide residue determination in fruit and vegetables by liquid chromatography–mass spectrometry 882(2000)153

Analysis of herbicide residues in cereals, fruits and vegetables 882(2000)175

Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in some fruits and vegetables using high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 888(2000)113

Gas chromatographic determination of the fatty-acid content of heat-treated green beans 891(2000)367

Determination of pesticides in vegetables using large-volume injection column liquid chromatography–electrospray tandem mass spectrometry 892(2000)379

#### Venoms

Partial primary structure of a fibrinogenase from the venom of the snake *Lachesis stenophrys* 852(1999)237

#### *Veronica longifolia*

Extraction of iridoid glycosides and their determination by micellar electrokinetic capillary chromatography 868(2000)73

**Vesicles**

Micellar electrokinetic chromatography with bis(2-ethylhexyl)sodium sulfosuccinate vesicles. Determination of synthetic food antioxidants 871(2000)403

**Viscosity**

Different elution modes and field programming in gravitational field-flow fractionation. 2. Experimental verification of the range of conditions for flow-rate and carrier liquid density programming 868(2000)217

Mobile-phase-viscosity dependence on DNA separation in slalom chromatography 886(2000)1

Effects of DNA topology, temperature and solvent viscosity on DNA retardation in slalom chromatography 893(2000)115

**Viscous fingering**

Visualization of viscous fingering in high-performance liquid chromatographic columns. Influence of the header design 865(1999)73

**Visualization of sample introduction**

Visualization of sample introduction in liquid chromatography columns. The effect of the frit diameter 855(1999)367

Visualization of sample introduction in liquid chromatographic columns. Contribution of a flow distributor on the sample band shape 865(1999)83

**Visualization of solute migration**

Visualization of solute migration in chromatographic columns. Quantitation of the concentration in a migrating zone 867(2000)71

**Visualisation of wall effects**

Physical evidence of two wall effects in liquid chromatography 888(2000)1

**Void volume**

Study of dead volume measurement in packed subcritical fluid chromatography with ODS columns and carbon dioxide–modifier mobile phases 866(2000)241

**Voltage gradient gel electrophoresis**

Modified apparatus for voltage gradient gel electrophoresis 855(1999)723

Use of voltage gradient gel electrophoresis in apoptotic DNA analysis 890(2000)371

**Von Willebrand factor**

Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes 852(1999)59

Use of high-resolution techniques for the characterization of clotting factor VIII 852(1999)175

Erratum to: “Use of surface plasmon resonance for studies of protein–protein and protein–phospholipid membrane interactions. Application to the binding of factor VIII to von Willebrand factor and to phosphatidylserine-containing membranes”. [J. Chromatogr. A, 852 (1999) 59-71] 872(2000)323

**W****Wall effects**

Physical evidence of two wall effects in liquid chromatography 888(2000)1

**Warfare agents**

Rapid screening procedures for the hydrolysis products of chemical warfare agents using positive and negative ion liquid chromatography–mass spectrometry with atmospheric pressure chemical ionisation 862(1999)169

Multiple-technique analytical characterization of a mixture containing chemical-weapons simulant from a munition 883(2000)185

Efficiency of pretreatment of aqueous samples using a macroporous strong anion-exchange resin on the determination of nerve gas hydrolysis products by gas chromatography–mass spectrometry after *tert.*-butyldimethylsilylation 891(2000)295

Methanol chemical ionization quadrupole ion trap mass spectrometry of *O*-ethyl *S*-[2-(diisopropylamino)ethyl] methylphosphonothiolate (VX) and its degradation products 893(2000)393

**Wastewater recycling**

Determination of anionic surfactants during wastewater recycling process by ion pair chromatography with suppressed conductivity detection 874(2000)207

**Water analysis**

Quantification of phenylurea herbicides and their free and humic acid-associated metabolites in natural waters 852(1999)465

Quantification of anions and cations in environmental water samples. Measurements with capillary electrophoresis and indirect-UV detection 853(1999)403

Determination of flotation reagents used in tin-mining by capillary electrophoresis 853(1999)497

- Detection and identification of sulphonamide drugs in municipal waste water by liquid chromatography coupled with electrospray ionisation tandem mass spectrometry 854(1999)163
- Use of volatile amines as ion-pairing agents for the high-performance liquid chromatographic–tandem mass spectrometric determination of aromatic sulfonates in industrial wastewater 854(1999)175
- On-line solid-phase extraction–ion-pair liquid chromatography–electrospray mass spectrometry for the trace determination of naphthalene monosulphonates in water 854(1999)187
- Simultaneous determination of antifouling herbicides in marina water samples by on-line solid-phase extraction followed by liquid chromatography–mass spectrometry 854(1999)197
- Investigations on the metabolism of alkyl polyglucosides and their determination in waste water by means of liquid chromatography–electrospray mass spectrometry 854(1999)221
- Analysis of low-molecular-mass inorganic and organic anions by ion chromatography–atmospheric pressure ionization mass spectrometry 854(1999)275
- On-line combination of aqueous-sample preparation and capillary gas chromatography 856(1999)279
- Efficient solid-phase extraction procedures for trace enrichment of priority phenols from industrial effluents with high total organic carbon content 857(1999)97
- Ultra-trace-level determination of polar pesticides and their transformation products in surface and estuarine water samples using column liquid chromatography–electrospray tandem mass spectrometry 857(1999)157
- Analysis of methyl *tert.*-butyl ether and its degradation products by direct aqueous injection onto gas chromatography with mass spectrometry or flame ionization detection systems 857(1999)205
- Simultaneous determination of halogenated neutral and acidic disinfection by-products in drinking water by closed-loop stripping extraction and capillary gas chromatography 857(1999)217
- Study of applicability of various solid-phase extraction materials for sample handling in screening analysis of organic micropollutants in water 857(1999)247
- Liquid chromatographic determination of imazosulfuron in drinking water and in soil using ultraviolet detection 857(1999)327
- Potential for chlorate interference in ion chromatographic determination of total nitrogen in natural waters following alkaline persulfate digestion 857(1999)337
- Analysis of linear alkylbenzenesulfonates in water samples by large-volume injection-port derivatization and gas chromatography–mass spectrometry 857(1999)359
- Semi-permeable surface analytical reversed-phase column for the improved trace analysis of acidic pesticides in water with coupled-column reversed-phase liquid chromatography with UV detection. Determination of bromoxynil and bentazone in surface water 858(1999)45
- Selective immunoclean-up followed by liquid or gas chromatography for the monitoring of polycyclic aromatic hydrocarbons in urban waste water and sewage sludges used for soil amendment 859(1999)29
- Solid-phase microextraction coupled with gas chromatography–ion trap mass spectrometry for the analysis of haloacetic acids in water 859(1999)159
- Analysis of some pesticides in water samples using solid-phase microextraction–gas chromatography with different mass spectrometric techniques 859(1999)193
- Sample cleanup and reversed-phase high-performance liquid chromatographic analysis of polar aromatic compounds in groundwater samples from a former gas plant 862(1999)137
- Analysis of anatoxin-a in freshwaters by automated on-line derivatization–liquid chromatography–electrospray mass spectrometry 862(1999)191
- Trapping efficiency of aqueous pollutants in multichannel thick-film silicone-rubber traps for capillary gas chromatography 863(1999)57
- Comparison of the performance of graphitized carbon black and poly(styrene–divinylbenzene) cartridges for the determination of pesticides and industrial phosphates in environmental waters 864(1999)121
- Ion chromatography determination of trace level bromate by large volume injection with conductivity and spectrophotometric detection after post column derivatisation 864(1999)263
- Determination of halogenated mono-alcohols and diols in water by gas chromatography with electron-capture detection 866(2000)65
- Determination of pesticides in waters by automatic on-line solid-phase extraction–capillary electrophoresis 866(2000)137
- Tandem solid-phase extraction of atrazine ozonation products in water 866(2000)195
- On-line coupling of equilibrium-sorptive enrichment to gas chromatography to determine low-molecular-mass pollutants in environmental water samples 867(2000)207



- Pesticide trace analysis using solid-phase extraction and gas chromatography with electron-capture and tandem mass spectrometric detection in water samples 867(2000)235
- Determination of ammonium in seawater by column-switching ion chromatography 868(2000)209
- On-line ion-pair solid-phase extraction–liquid chromatography–mass spectrometry for the analysis of quaternary ammonium herbicides 869(2000)441
- Determination of triazine herbicides in natural waters by solid-phase extraction and non-aqueous capillary zone electrophoresis 869(2000)451
- Use of solid-phase extraction and high-performance liquid chromatography for the determination of triazine residues in water: validation of the method 869(2000)463
- Evaluation of surface- and ground-water pollution due to herbicides in agricultural areas of Zamora and Salamanca (Spain) 869(2000)471
- Microporous membrane liquid–liquid extraction coupled on-line with normal-phase liquid chromatography for the determination of cationic surfactants in river and waste water 869(2000)523
- Trace-level determination of triazines and several degradation products in environmental waters by disk solid-phase extraction and micellar electrokinetic chromatography 872(2000)299
- Analysis of steroids in environmental water samples using solid-phase extraction and ion-trap gas chromatography–mass spectrometry and gas chromatography–tandem mass spectrometry 872(2000)309
- Determination of butyltin species in water and sediment by solid-phase microextraction–gas chromatography–flame ionization detection 873(2000)63
- Solid-phase microextraction–gas chromatography–mass spectrometry of biologically active substances in water samples 873(2000)95
- Development of a headspace solid-phase microextraction procedure for the determination of free volatile fatty acids in waste waters 873(2000)107
- Determination of silicate in water by ion exclusion chromatography with conductivity detection 874(2000)143
- Trace analysis of ten chlorinated benzenes in water by headspace solid-phase microextraction 874(2000)149
- Solid-phase extraction followed by high-performance liquid chromatography–ionspray interface–mass spectrometry for monitoring of herbicides in environmental water 874(2000)187
- Determination of chlorophenoxy acid herbicides in water by in situ esterification followed by in-vial liquid–liquid extraction combined with large-volume on-column injection and gas chromatography–mass spectrometry 877(2000)153
- Simultaneous determination of acidic and non-acidic pesticides in natural waters by liquid chromatography–mass spectrometry 878(2000)87
- Determination of rice herbicides, their transformation products and clofibric acid using on-line solid-phase extraction followed by liquid chromatography with diode array and atmospheric pressure chemical ionization mass spectrometric detection 879(2000)3
- Monitoring of priority pesticides and other organic pollutants in river water from Portugal by gas chromatography–mass spectrometry and liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)13
- Part-per-trillion level determination of antifouling pesticides and their byproducts in seawater samples by off-line solid-phase extraction followed by high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 879(2000)27
- Comparison of various liquid chromatographic methods involving UV and atmospheric pressure chemical ionization mass spectrometric detection for the efficient trace analysis of phenylurea herbicides in various types of water samples 879(2000)39
- Application of liquid chromatography with mass spectrometry combined with photodiode array detection and tandem mass spectrometry for monitoring pesticides in surface waters 879(2000)51
- Stability of organophosphorus insecticides on graphitized carbon black extraction cartridges used for large volumes of surface water 879(2000)73
- Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants 879(2000)83
- Determination of endocrine disruptors in water after derivatization with *N*-methyl-*N*-(*tert*-butyldimethyltrifluoroacetamide) using gas chromatography with mass spectrometric detection 879(2000)97
- Solid-phase microextraction for determining the distribution of sixteen US Environmental Protection Agency polycyclic aromatic hydrocarbons in water samples 879(2000)177
- Advances in the determination of inorganic anions by ion chromatography 881(2000)607
- Eliminating the chlorite interference in US Environmental Protection Agency Method 317.0 permits analysis of trace bromate levels in all drinking water matrices 882(2000)309

- On-line microdialysis–ion chromatographic determination of inorganic anions in olive-oil mill wastewater 882(2000)321
- Improved scheme of chelation ion chromatography with a mixed eluent for the simultaneous analysis of transition metals at  $\mu\text{g l}^{-1}$  levels 883(2000)127
- Determination of sulfonated azo dyes in river water samples by capillary zone electrophoresis 883(2000)277
- Sensitive determination of dissolved sulfide in estuarine water by solid-phase extraction and high-performance liquid chromatography of methylene blue 883(2000)305
- Simplified column-switching technology for the determination of traces of anions in the presence of high concentrations of other anions 884(2000)53
- Determination of thiosulfate, thiocyanate and polythionates in a mixture by ion-pair chromatography with ultraviolet absorbance detection 884(2000)81
- Indirect ultraviolet spectrophotometric detection in the ion chromatography of common mono- and divalent cations on an aluminium adsorbing silica gel column with tyramine-containing crown ethers as eluent 884(2000)113
- Suppressed electrostatic ion chromatography with tetraborate as eluent and its application to the determination of inorganic anions in snow and rainwater 884(2000)161
- Simultaneous ion-exclusion/cation-exchange chromatography of anions and cations in acid rain waters on a weakly acidic cation-exchange resin by elution with sulfosalicylic acid 884(2000)167
- Determination of inorganic anions in environmental waters with a hydroxide-selective column 884(2000)175
- Determination of chloride, sulfate and nitrate in groundwater samples by ion chromatography 884(2000)185
- Comparison of on-line coupling of ion-chromatography with atmospheric pressure ionization mass spectrometry and with inductively coupled plasma mass spectrometry as tools for the ultra-trace analysis of bromate in surface water samples 884(2000)191
- Performance evaluation of a method for the determination of bromate in drinking water by ion chromatography (EPA Method 317.0) and validation of EPA Method 324.0 884(2000)201
- Determination of inorganic and organic anionic arsenic species in water by ion chromatography coupled to hydride generation–inductively coupled plasma atomic emission spectrometry 884(2000)211
- Trace analysis of heavy metals in groundwater samples by ion chromatography with post-column reaction and ultraviolet–visible detection 884(2000)229
- Determination of epichlorohydrin by ion chromatography 884(2000)251
- Fifty years of solid-phase extraction in water analysis – historical development and overview 885(2000)3
- Investigation into the effects of temperature and stirring rate on the solid-phase extraction of diuron from water using a  $\text{C}_{18}$  extraction disk 885(2000)41
- Solid-phase extraction for multiresidue analysis of organic contaminants in water 885(2000)195
- Multiresidue methods using solid-phase extraction techniques for monitoring priority pesticides, including triazines and degradation products, in ground and surface waters 885(2000)217
- Solid-phase extraction of quaternary ammonium herbicides 885(2000)251
- Solid-phase extraction of polycyclic aromatic compounds 885(2000)273
- On-line dual-precursor-based trace enrichment for the determination of polar and acidic microcontaminants in river water by liquid chromatography with diode-array UV and tandem mass spectrometric detection 885(2000)377
- New methods for determination of glyphosate and (aminomethyl)phosphonic acid in water and soil 886(2000)207
- Identification of reactive dyes in spent dyebaths and wastewater by capillary electrophoresis–mass spectrometry 886(2000)271
- Improved method for the determination of trace perchlorate in ground and drinking waters by ion chromatography 888(2000)151
- Mixed-level orthogonal array design for the optimization of solid-phase extraction of some pesticides from surface water 888(2000)197
- Fractionation and toxicity evaluation of waste waters 889(2000)149
- Assessment of the pollutant elimination efficiency by gas chromatography/mass spectrometry, liquid chromatography–mass spectrometry and –tandem mass spectrometry. Comparison of conventional and membrane-assisted biological wastewater treatment processes 889(2000)155
- Inter-laboratory comparison of liquid chromatographic techniques and enzyme-linked immunosorbent assay for the determination of surfactants in wastewaters 889(2000)195
- Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids 889(2000)211

- Determination of linear alkylbenzenesulfonates in wastewater treatment plants and coastal waters by automated solid-phase extraction followed by capillary electrophoresis–UV detection and confirmation by capillary electrophoresis–mass spectrometry 889(2000)221
- Stability of sulfonated derivatives of benzene and naphthalene on disposable solid-phase extraction pre-columns and in an aqueous matrix 889(2000)231
- Unknown bisethylisooctanollactone isomers in industrial waste water. Isolation, identification and occurrence in surface water 889(2000)245
- Multiresidue method for the analysis of five antifouling agents in marine and coastal waters by gas chromatography–mass spectrometry with large-volume injection 889(2000)261
- Determination of naphthalenesulfonates in water by on-line ion-pair solid-phase extraction and ion-pair liquid chromatography with fast-scanning fluorescence detection 890(2000)289
- Simultaneous determination of inorganic disinfection by-products and the seven standard anions by ion chromatography 890(2000)295
- Indirect analysis of urea herbicides from environmental water using solid-phase microextraction 890(2000)303
- Analysis of agrochemicals by capillary electrophoresis (Review) 891(2000)45
- Application of gas chromatography coupled to chemical ionisation mass spectrometry following headspace solid-phase microextraction for the determination of free volatile fatty acids in aqueous samples 891(2000)287
- Determination of steroid sex hormones and related synthetic compounds considered as endocrine disrupters in water by liquid chromatography–diode array detection–mass spectrometry 892(2000)391
- Recent and future developments of liquid chromatography in pesticide trace analysis 892(2000)435
- Determination of fungicides in natural waters using solid-phase microextraction and gas chromatography coupled with electron-capture and mass spectrometric detection 893(2000)143
- On-line monitoring trihalomethanes in chlorinated water by membrane introduction–fast gas chromatography mass–spectrometry 893(2000)169
- High-performance liquid chromatography–diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation 893(2000)253
- Solid-phase microextraction for the determination of systemic and non-volatile pesticides in river water using gas chromatography with nitrogen–phosphorous and electron-capture detection 893(2000)347
- Evaluation of purge-and-trap–high-resolution gas chromatography–mass spectrometry for the determination of 27 volatile organic compounds in marine water at the  $\text{ng l}^{-1}$  concentration level 893(2000)367
- Determination of soluble anions and cations from waters of pulp and paper mills with on-line coupled capillary electrophoresis 895(2000)189
- Determination of aromatic amines in water samples by capillary electrophoresis with electrochemical and fluorescence detection 895(2000)197
- Simultaneous determination of alkali, alkaline earth and transition metal ions by capillary electrophoresis with indirect UV detection 895(2000)205
- Separation of two groups of oestrogen mimicking compounds using micellar electrokinetic chromatography 895(2000)237
- On-line coupling of solid-phase extraction to gas chromatography with fast solvent vaporization and concentration in an open injector liner. Analysis of pesticides in aqueous samples 896(2000)41
- Determination of aqueous chlorothalonil with solid-phase microextraction and gas chromatography 896(2000)105
- Analysis of chlorophenoxy acid herbicides in water by large-volume on-line derivatization and gas chromatography–mass spectrometry 896(2000)111
- Improved routine speciation of organotin compounds in environmental samples by pulsed flame photometric detection 896(2000)149
- Comparison of different sorbent materials for *on-line* solid-phase extraction with liquid chromatography–atmospheric pressure chemical ionization mass spectrometry of phenols 896(2000)159
- Optimisation of alachlor solid-phase microextraction from water samples using experimental design 896(2000)373
- Simultaneous determination of cyanogen chloride and cyanogen bromide in treated water at sub- $\mu\text{g/L}$  levels by a new solid-phase microextraction–gas chromatographic–electron-capture detection method 897(2000)307
- Wheat**  
Determination of nivalenol and deoxynivalenol in wheat using liquid chromatography–mass spectrometry with negative ion atmospheric pressure chemical ionisation 854(1999)45

Simultaneous determination of phenolic acids and 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one in wheat (*Triticum aestivum* L.) by gas chromatography–tandem mass spectrometry 864(1999)315

Continuous split-flow thin cell and gravitational field-flow fractionation of wheat starch particles 871(2000)449

Determination of aflatoxins in food by overpressured-layer chromatography 882(2000)11

Simultaneous determination of trichothecene mycotoxins and zearalenone in cereals by gas chromatography–mass spectrometry 882(2000)23

Interlaboratory study of identification and quantitation of multiresidue pyrethroids in agricultural products by gas chromatography–mass spectrometry 882(2000)231

### Whey

Analysis of bovine whey proteins in soybean dairy-like products by capillary electrophoresis 859(1999)77

Simultaneous separation and quantitation of the major bovine whey proteins including proteose peptone and caseinomacropptide by reversed-phase high-performance liquid chromatography on polystyrene–divinylbenzene 878(2000)183

### Wine

Determination of indole-3-acetic acid, tryptophan and other indoles in must and wine by high-performance liquid chromatography with fluorescence detection 855(1999)227

Liquid chromatographic–mass spectrometric determination of phenolic compounds using a capillary-scale particle beam interface 855(1999)515

Determination of oak lactones in barrel-aged wines and in oak extracts by stable isotope dilution analysis 857(1999)239

Semipreparative reversed-phase liquid chromatographic fractionation of aroma extracts from wine and other alcoholic beverages 864(1999)77

Determination of ochratoxin A in wine by means of immunoaffinity column clean-up and high-performance liquid chromatography 864(1999)89

Optimization of chromatographic parameters for the determination of biogenic amines in wines by reversed-phase high-performance liquid chromatography 871(2000)75

Determination of phenols in wines by liquid chromatography with photodiode array and fluorescence detection 871(2000)85

Method development for the determination of anthocyanins in red wines by high-performance liquid chromatography and classification of German red wines by means of multivariate statistical methods 871(2000)95

Optimization of the separation of phenolic compounds by micellar electrokinetic capillary chromatography 871(2000)427

Utilization of fluorescein sodium salt for the indirect fluorimetric detection in micellar electrokinetic chromatography 872(2000)247

Quantitative analysis of 4-ethylphenol and 4-ethylguaiacol in red wine 874(2000)101

Separation of polyphenols in Canary Islands wine by capillary zone electrophoresis without preconcentration 874(2000)111

Headspace solid-phase microextraction method for determining 3-alkyl-2-methoxypyrazines in musts by means of polydimethylsiloxane–divinylbenzene fibres 880(2000)93

Electrophoretic methods in the analysis of beverages 880(2000)243

High-performance liquid chromatographic determination of mono-, poly- and hydroxycarboxylic acids in foods and beverages as their 2-nitrophenylhydrazides 881(2000)365

Analysis of anthocyanins in foods by liquid chromatography, liquid chromatography–mass spectrometry and capillary electrophoresis 881(2000)403

Analysis of several phenolic compounds with potential antioxidant properties in grape extracts and wines by high-performance liquid chromatography–photodiode array detection without sample preparation 881(2000)461

Analysis of resveratrol in wine by capillary electrophoresis 881(2000)471

Monoterpenes in grape juice and wines 881(2000)557

Analysis of organic sulfur compounds in wine aroma 881(2000)569

Analysis of low-volatility organic sulphur compounds in wines by solid-phase microextraction and gas chromatography 881(2000)583

Determination of lithium in wines by ion chromatography 881(2000)645

Solid-phase extraction applied to the determination of ochratoxin A in wines by reversed-phase high-performance liquid chromatography 882(2000)29

Multiresidue analysis of seventeen pesticides in wine by gas chromatography with mass-selective detection 882(2000)205

Multiresidue method for the rapid determination of organophosphorus insecticides in grapes, must and wine 882(2000)213

Multiresidue method for the rapid determination – in grape, must and wine – of fungicides frequently used on vineyards 882(2000)221

Combined ion-pair extraction and gas chromatography–mass spectrometry for the simultaneous determination of diamines, polyamines and aromatic amines in Port wine and grape juice 886(2000)183

High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light 888(2000)121

Comparison of different immunoaffinity clean-up procedures for high-performance liquid chromatographic analysis of ochratoxin A in wines 888(2000)129

Multi-residue methodology for pesticide screening in wines 889(2000)59

### **Wood**

Determination of oak lactones in barrel-aged wines and in oak extracts by stable isotope dilution analysis 857(1999)239

Optimization of the extraction of polycyclic aromatic hydrocarbons from wood samples by the use of microwave energy 869(2000)505

Analysis of pitch deposits produced in Kraft pulp mills using a totally chlorine free bleaching sequence 874(2000)235

Analysis of carbohydrates in wood and pulps employing enzymatic hydrolysis and subsequent capillary zone electrophoresis 891(2000)157

## **X**

### **X-Ray photoelectron spectrometry**

Unexpected surface chemistry in capillaries for electrophoresis 894(2000)73

Unexpected surface chemistry in capillaries for electrophoresis 894(2000)73

## **Z**

### **Zero defects**

Towards a quantitative definition of perfection in chromatographic analyses 868(2000)305

### **Zirconia**

Titania as a sorbent in normal-phase liquid chromatography 888(2000)51

Use of spray-dried zirconia microspheres in the separation of immunoglobulins from cell culture supernatant 890(2000)15

### **Zirconium-adsorbing silica gel**

Ion-chromatographic behavior of alkali metal cations and ammonium ion on zirconium-adsorbing silica gel 884(2000)123

### **Zirconized silica**

Stability of high-performance liquid chromatography columns packed with poly(methyloctylsiloxane) sorbed and radiation-immobilized onto porous silica and zirconized silica 869(2000)129

### **Zwitterionic surfactant stationary phase**

Suppressed electrostatic ion chromatography with tetraborate as eluent and its application to the determination of inorganic anions in snow and rainwater 884(2000)161