



ELSEVIER

Journal of Chromatography A, 914 (2001) 337–352

JOURNAL OF
CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Subject Index Vols. 913–914

Acetonitrile

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Actinorhodin

Recovery of actinorhodin from fermentation broth 914(2001)67

Adsorbents

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Adsorption isotherms

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Affinity chromatography

Study on the degeneracy of antisense peptides using affinity chromatography 913(2001)423

Aflatoxins

Some applications of near-ultraviolet laser-induced fluorescence detection in nanomolar- and subnanomolar-range high-performance liquid chromatography or micro-high-performance liquid chromatography 913(2001)253

Agenerase

Liquid chromatographic–tandem mass spectrometric determination of amprenavir (agenerase) in serum/plasma of human immunodeficiency virus type-1 infected patients receiving combination antiretroviral therapy 914(2001)1

Alkali metals

Determination of inorganic cations and anions by ion-exchange chromatography with evaporative light-scattering detection 914(2001)167

Alkaline-earth metals

Determination of inorganic cations and anions by ion-exchange chromatography with evaporative light-scattering detection 914(2001)167

Alkylbenzamides

Comparison of selected retention models in reversed-phase liquid chromatography 913(2001)3

Alkylbenzene

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Alkylbenzenes

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Amides

Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities 913(2001)113

Amines, heterocyclic aromatic

Liquid chromatography–tandem mass spectrometry method of urine analysis for determining human variation in carcinogen metabolism 914(2001)95

Amino acids

Derivatization and chromatographic behavior of the *o*-phthaldialdehyde amino acid derivatives obtained with various SH-group-containing additives 913(2001)283

Analysis of amino acids in human serum by isocratic reversed-phase high-performance liquid chromatography with electrochemical detection 913(2001)303

Simultaneous determination of several amino acids, including homocysteine, cysteine and glutamic acid, in human plasma by isocratic reversed-phase high-performance liquid chromatography with fluorimetric detection 913(2001)309

Determination by liquid chromatography of free and total cysteine in human urine in the form of its *S*-quinolinium derivative 913(2001)319

Simultaneous analysis of underivatized chiral amino acids by liquid chromatography–ionspray tandem mass spectrometry using a teicoplanin chiral stationary phase 913(2001)331

Study on the degeneracy of antisense peptides using affinity chromatography 913(2001)423

Application of capillary electrophoresis with laser-induced fluorescence detection to the determination of biogenic amines and amino acids in brain microdialysate and homogenate samples 914(2001)293

Amino acids, OPA derivatives

Derivatization and chromatographic behavior of the *o*-phthalaldehyde amino acid derivatives obtained with various SH-group-containing additives 913(2001)283

Aminoglycosides

Sample preparation for residue determination of gentamicin and neomycin by liquid chromatography 914(2001)59

Aminomethylphenylimidazo[4,5-*b*]pyridine

Liquid chromatography–tandem mass spectrometry method of urine analysis for determining human variation in carcinogen metabolism 914(2001)95

Amino thiols

Determination of biologically active low-molecular-mass thiols in human blood. III. Highly sensitive narrow-bore isocratic reversed-phase high-performance liquid chromatography with fluorescence detection 913(2001)315

Amprenavir

Liquid chromatographic–tandem mass spectrometric determination of amprenavir (agenerase) in serum/plasma of human immunodeficiency virus type-1 infected patients receiving combination antiretroviral therapy 914(2001)1

Anilines

Effect of the eluent pH and acidic modifiers in high-performance liquid chromatography retention of basic analytes 913(2001)173

Effect of counter-anion concentration on retention in high-performance liquid chromatography of protonated basic analytes 913(2001)189

Antibiotics

Liquid chromatography of troleandomycin 914(2001)53

Sample preparation for residue determination of gentamicin and neomycin by liquid chromatography 914(2001)59

Recovery of actinorhodin from fermentation broth 914(2001)67

Determination and pharmacokinetic study of unbound cefepime in rat bile by liquid chromatography with on-line microdialysis 914(2001)77

On-line microdialysis coupled with microbore liquid chromatography with ultraviolet detection for continuous monitoring of free cefsulodin in rat blood 914(2001)83

Antibodies

Rapid method for monitoring galactosylation levels during recombinant antibody production by electrospray mass spectrometry with selective-ion monitoring 913(2001)437

Antidepressants

Bupropion hydrochloride: the development of a chiral separation using an ovomucoid column 913(2001)275

Antioxidants

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Antisense peptides

Study on the degeneracy of antisense peptides using affinity chromatography 913(2001)423

Aromatic compounds

Acrylamide-based monoliths as robust stationary phases for capillary electrochromatography 914(2001)211

Arylpropionic acids

Uniformly sized molecularly imprinted polymer for (*S*)-naproxen. Retention and molecular recognition properties in aqueous mobile phase 913(2001)141

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Association constants

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

Band profiles

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. V. Results obtained with Vydac 218TP C₁₈ columns 913(2001)89

Barbiturates

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

Basic drugs

Effect of liposome type and membrane fluidity on drug–membrane partitioning analyzed by immobilized liposome chromatography 913(2001)123

Beclomethasone

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

Benzene

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Benzoic acid

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Benzylamines

Effect of counter-anion concentration on retention in high-performance liquid chromatography of protonated basic analytes 913(2001)189

Beta-blockers

Effect of liposome type and membrane fluidity on drug–membrane partitioning analyzed by immobilized liposome chromatography 913(2001)123

Betacyanins

Application of high-performance liquid chromatography to the characterization of the betalain pigments in prickly pear fruits 913(2001)415

Betamethasones

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

Betaxanthins

Application of high-performance liquid chromatography to the characterization of the betalain pigments in prickly pear fruits 913(2001)415

Biogenic amines

Application of capillary electrophoresis with laser-induced fluorescence detection to the determination of biogenic amines and amino acids in brain microdialysate and homogenate samples 914(2001)293

Bipyridyls

Removal of metallic impurities from mobile phases in reversed-phase high-performance liquid chromatography by the use of an in-line chelation column 913(2001)159

Bisphosphonates

Ion-exchange liquid chromatographic analysis of bisphosphonates by on-line post-column photochemical reaction and spectrophotometric detection 914(2001)105

Bupropion

Bupropion hydrochloride: the development of a chiral separation using an ovomucoid column 913(2001)275

Butylparaben

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Capillary electrophoresis–mass spectrometry

Pressure-assisted capillary electrophoresis–electrospray ion trap mass spectrometry for the analysis of heparin depolymerised disaccharides 914(2001)277

Carbohydrates

Application of high-performance liquid chromatography with isotope-ratio mass spectrometry for measuring low levels of enrichment of underivatized materials 913(2001)269

Artifact-free matrix-assisted laser desorption ionization time-of-flight mass spectra of *tert*-butyldimethylsilyl ether derivatives of cyclodextrins used for the synthesis of single-isomer, chiral resolving agents for capillary electrophoresis 914(2001)325

Carotenoids

High-performance liquid chromatography for the characterization of carotenoids in the new sweet orange (Earlygold) grown in Florida, USA 913(2001)371

Catechin

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Catecholamines

Some applications of near-ultraviolet laser-induced fluorescence detection in nanomolar- and subnanomolar-range high-performance liquid chromatography or micro-high-performance liquid chromatography 913(2001)253

Cefepime

Determination and pharmacokinetic study of unbound cefepime in rat bile by liquid chromatography with on-line microdialysis 914(2001)77

Cefsulodin

On-line microdialysis coupled with microbore liquid chromatography with ultraviolet detection for continuous monitoring of free cefsulodin in rat blood 914(2001)83

Cells

Proteolysis of whole cell extracts with immobilized enzyme columns as part of multidimensional chromatography 913(2001)429

Cephalosporins

Determination and pharmacokinetic study of unbound cefepime in rat bile by liquid chromatography with on-line microdialysis 914(2001)77

On-line microdialysis coupled with microbore liquid chromatography with ultraviolet detection for continuous monitoring of free cefsulodin in rat blood 914(2001)83

2-Chloro-1-methyloquinolinium tetrafluoroborate

Facile and sensitive method for the determination of mesna in plasma by high-performance liquid chromatography with ultraviolet detection 914(2001)29

2-Chloro-1-methylquinolinium tetrafluoroborate

Determination by liquid chromatography of free and total cysteine in human urine in the form of its *S*-quinolinium derivative 913(2001)319

Chelation

Removal of metallic impurities from mobile phases in reversed-phase high-performance liquid chromatography by the use of an in-line chelation column 913(2001)159

Chiral stationary phases, LC

Simultaneous analysis of underivatized chiral amino acids by liquid chromatography–ionspray tandem mass spectrometry using a teicoplanin chiral stationary phase 913(2001)331

Column switching

Isolation of the components of a complex mixture by means of column switching for their enhanced detection by mass spectrometry 913(2001)209

Complexation

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

Cyclodextrin

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

Cyclodextrins

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Study on the chiral recognition of the enantiomers of ephedrine derivatives with neutral and sulfated heptakis(2,3-*O*-diacetyl)- β -cyclodextrins using capillary electrophoresis, UV, nuclear magnetic resonance spectroscopy and mass spectrometry 914(2001)315

Artifact-free matrix-assisted laser desorption ionization time-of-flight mass spectra of *tert*-butyldimethylsilyl ether derivatives of cyclodextrins used for the synthesis of single-isomer, chiral resolving agents for capillary electrophoresis 914(2001)325

Cysteine

Simultaneous determination of several amino acids, including homocysteine, cysteine and glutamic acid, in human plasma by isocratic reversed-phase high-performance liquid chromatography with fluorimetric detection 913(2001)309

Determination by liquid chromatography of free and total cysteine in human urine in the form of its *S*-quinolinium derivative 913(2001)319

Cystine

Determination by liquid chromatography of free and total cysteine in human urine in the form of its *S*-quinolinium derivative 913(2001)319

Daidzein

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Daidzin

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Data analysis software

Validation of a chromatography data analysis software 913(2001)221

Degeneracy

Study on the degeneracy of antisense peptides using affinity chromatography 913(2001)423

Derivatization, electrophoresis

Application of capillary electrophoresis with laser-induced fluorescence detection to the determination of biogenic amines and amino acids in brain microdialysate and homogenate samples 914(2001)293

Derivatization, LC

Derivatization and chromatographic behavior of the *o*-phthaldialdehyde amino acid derivatives obtained with various SH-group-containing additives 913(2001)283

Analysis of amino acids in human serum by isocratic reversed-phase high-performance liquid chromatography with electrochemical detection 913(2001)303

Determination by liquid chromatography of free and total cysteine in human urine in the form of its *S*-quinolinium derivative 913(2001)319

Facile and sensitive method for the determination of mesna in plasma by high-performance liquid chromatography with ultraviolet detection 914(2001)29

Improved measurement of formaldehyde in water-soluble polymers by high-performance liquid chromatography coupled with post-column reaction detection 914(2001)123

Descarboethoxyloratadine

Sensitive liquid chromatography–tandem mass spectrometry method for the determination of loratadine and its major active metabolite descarboethoxyloratadine in human plasma 914(2001)37

Detection, LC

Some applications of near-ultraviolet laser-induced fluorescence detection in nanomolar- and subnanomolar-range high-performance liquid chromatography or micro-high-performance liquid chromatography 913(2001)253

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Fractionation and characterization of phenolic resins by high-performance liquid chromatography and gel-permeation chromatography combined with ultraviolet, refractive index, mass spectrometry and light-scattering detection 914(2001)147

Determination of inorganic cations and anions by ion-exchange chromatography with evaporative light-scattering detection 914(2001)167

Dexamethasone

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

Dinitrophenylhydrazine

Improved measurement of formaldehyde in water-soluble polymers by high-performance liquid chromatography coupled with post-column reaction detection 914(2001)123

Diphenylethylenediamine

Some applications of near-ultraviolet laser-induced fluorescence detection in nanomolar- and subnanomolar-range high-performance liquid chromatography or micro-high-performance liquid chromatography 913(2001)253

Disaccharides

Pressure-assisted capillary electrophoresis–electrospray ion trap mass spectrometry for the analysis of heparin depolymerised disaccharides 914(2001)277

DNA

On-line nanoliter cycle sequencing reaction with capillary zone electrophoresis purification for DNA sequencing 914(2001)245

Doxepin

Sensitive liquid chromatographic–tandem mass spectrometric method for the determination of fluoxetine and its primary active metabolite norfluoxetine in human plasma 914(2001)45

Efficiency

Volume-load capacity in fast-gradient liquid chromatography. Effect of sample solvent composition and injection volume on chromatographic performance 913(2001)233

Electrochromatography

Rapid separation of peptides and proteins by isocratic capillary electrochromatography at elevated temperature 914(2001)189

Size-exclusion electrochromatography with controlled pore flow 914(2001)201

Acrylamide-based monoliths as robust stationary phases for capillary electrochromatography 914(2001)211

Electrokinetic chromatography

Hydrophobic interaction electrokinetic chromatography for the separation of polycyclic aromatic hydrocarbons using non-aqueous matrices 914(2001)223

Electroosmotic flow

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

Electrospray ionization

Preliminary investigation of the application of on-line membrane extraction of trifluoroacetic acid as an aid to improvement of negative ion electrospray mass spectrometry data 913(2001)205

Enantiomer separation

Uniformly sized molecularly imprinted polymer for (*S*)-naproxen. Retention and molecular recognition properties in aqueous mobile phase 913(2001)141

Bupropion hydrochloride: the development of a chiral separation using an ovomucoid column 913(2001)275

Simultaneous analysis of underivatized chiral amino acids by liquid chromatography–ionspray tandem mass spectrometry using a teicoplanin chiral stationary phase 913(2001)331

Study on the chiral recognition of the enantiomers of ephedrine derivatives with neutral and sulfated heptakis(2,3-*O*-diacetyl)- β -cyclodextrins using capillary electrophoresis, UV, nuclear magnetic resonance spectroscopy and mass spectrometry 914(2001)315

Enantiomer separations

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Enzyme inhibitors

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Determination of serum levels of thirteen human immunodeficiency virus-suppressing drugs by high-performance liquid chromatography 913(2001)449

Liquid chromatographic–tandem mass spectrometric determination of amprenavir (agenerase) in serum/plasma of human immunodeficiency virus type-1 infected patients receiving combination antiretroviral therapy 914(2001)1

Ephedrines

Study on the chiral recognition of the enantiomers of ephedrine derivatives with neutral and sulfated heptakis(2,3-*O*-diacetyl)- β -cyclodextrins using capillary electrophoresis, UV, nuclear magnetic resonance spectroscopy and mass spectrometry 914(2001)315

Escherichia coli

Proteolysis of whole cell extracts with immobilized enzyme columns as part of multidimensional chromatography 913(2001)429

High-efficiency capillary isoelectric focusing of protein complexes from *Escherichia coli* cytosolic extracts 914(2001)257

Ethoxylates

Separation of isomers of nonylphenol and select nonylphenol polyethoxylates by high-performance liquid chromatography on a graphitic carbon column 914(2001)161

Ethylene glycol dimethacrylate

Uniformly sized molecularly imprinted polymer for (*S*)-naproxen. Retention and molecular recognition properties in aqueous mobile phase 913(2001)141

Evaporative light-scattering detection

Determination of inorganic cations and anions by ion-exchange chromatography with evaporative light-scattering detection 914(2001)167

Expanded bed chromatography

Recovery of actinorhodin from fermentation broth 914(2001)67

Fetuin

Determination of sialic acids by liquid chromatography–mass spectrometry 913(2001)365

Field-flow fractionation

Optimization of experimental conditions for the separation of small and large starch granules by gravitational field-flow fractionation 914(2001)183

Flavonoids

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Flow-rate

New micro-flow pumping system for liquid chromatography 913(2001)165

Flow-rate gradients

Optimization of experimental conditions for the separation of small and large starch granules by gravitational field-flow fractionation 914(2001)183

Food analysis

High-performance liquid chromatography for the characterization of carotenoids in the new sweet orange (Earlygold) grown in Florida, USA 913(2001)371

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Sample preparation for residue determination of gentamicin and neomycin by liquid chromatography 914(2001)59

Effects of the matrix and sample preparation on the determination of fluoroquinolone residues in animal tissues 914(2001)89

Foreword

Foreword 913(2001)1

Fluoroquinolones

Effects of the matrix and sample preparation on the determination of fluoroquinolone residues in animal tissues 914(2001)89

Fluoxetine

Sensitive liquid chromatographic–tandem mass spectrometric method for the determination of fluoxetine and its primary active metabolite norfluoxetine in human plasma 914(2001)45

Flutamide

Appropriate column configurations for the rapid analysis and semipreparative purification of the radiolabeled drug flutamide by high-performance liquid chromatography 914(2001)23

Formaldehyde

Improved measurement of formaldehyde in water-soluble polymers by high-performance liquid chromatography coupled with post-column reaction detection 914(2001)123

Fractionation and characterization of phenolic resins by high-performance liquid chromatography and gel-permeation chromatography combined with ultraviolet, refractive index, mass spectrometry and light-scattering detection 914(2001)147

Fruit juices

High-performance liquid chromatography for the characterization of carotenoids in the new sweet orange (Earlygold) grown in Florida, USA 913(2001)371

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Fruits

Application of high-performance liquid chromatography to the characterization of the betalain pigments in prickly pear fruits 913(2001)415

Genistein

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Genistin

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Gentamicin

Sample preparation for residue determination of gentamicin and neomycin by liquid chromatography 914(2001)59

Glucose

Simultaneous monitoring of extracellular glucose, pyruvate, lactate and glutamate in gerbil cortex during focal cerebral ischemia by dual probe microdialysis 913(2001)349

Glucuronides

Liquid chromatography–tandem mass spectrometry method of urine analysis for determining human variation in carcinogen metabolism 914(2001)95

Glutamate

Development of an immobilized brain glutamine synthetase liquid chromatographic stationary phase for on-line biochemical studies 913(2001)133

Simultaneous monitoring of extracellular glucose, pyruvate, lactate and glutamate in gerbil cortex during focal cerebral ischemia by dual probe microdialysis 913(2001)349

Glutamic acid

Simultaneous determination of several amino acids, including homocysteine, cysteine and glutamic acid, in human plasma by isocratic reversed-phase high-performance liquid chromatography with fluorimetric detection 913(2001)309

Glutamine synthetase

Development of an immobilized brain glutamine synthetase liquid chromatographic stationary phase for on-line biochemical studies 913(2001)133

Glycitein

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Glycitin

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Glycoproteins

Determination of sialic acids by liquid chromatography–mass spectrometry 913(2001)365

Rapid method for monitoring galactosylation levels during recombinant antibody production by electrospray mass spectrometry with selective-ion monitoring 913(2001)437

Glycosides

Sample preparation for residue determination of gentamicin and neomycin by liquid chromatography 914(2001)59

Gradient elution

Volume-load capacity in fast-gradient liquid chromatography. Effect of sample solvent composition and injection volume on chromatographic performance 913(2001)233

Ion-exchange liquid chromatographic analysis of bisphosphonates by on-line post-column photochemical reaction and spectrophotometric detection 914(2001)105

Optimization of experimental conditions for the separation of small and large starch granules by gravitational field-flow fractionation 914(2001)183

Henry constants

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Heparin

Pressure-assisted capillary electrophoresis–electrospray ion trap mass spectrometry for the analysis of heparin depolymerised disaccharides 914(2001)277

Homocysteine

Simultaneous determination of several amino acids, including homocysteine, cysteine and glutamic acid, in human plasma by isocratic reversed-phase high-performance liquid chromatography with fluorimetric detection 913(2001)309

Determination of biologically active low-molecular-mass thiols in human blood. III. Highly sensitive narrow-bore isocratic reversed-phase high-performance liquid chromatography with fluorescence detection 913(2001)315

Human immunodeficiency virus

Determination of serum levels of thirteen human immunodeficiency virus-suppressing drugs by high-performance liquid chromatography 913(2001)449

Liquid chromatographic–tandem mass spectrometric determination of amprenavir (agenerase) in serum/plasma of human immunodeficiency virus type-1 infected patients receiving combination antiretroviral therapy 914(2001)1

Hummel–Dreyer method

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

Hydrophilic interaction chromatography

Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities 913(2001)113

Hydrophobic interaction electrokinetic chromatography

Hydrophobic interaction electrokinetic chromatography for the separation of polycyclic aromatic hydrocarbons using non-aqueous matrices 914(2001)223

Immobilized liposomes

Effect of liposome type and membrane fluidity on drug–membrane partitioning analyzed by immobilized liposome chromatography 913(2001)123

Immobilized peptides

Study on the degeneracy of antisense peptides using affinity chromatography 913(2001)423

Immobilized proteins

Development of an immobilized brain glutamine synthetase liquid chromatographic stationary phase for on-line biochemical studies 913(2001)133

Proteolysis of whole cell extracts with immobilized enzyme columns as part of multidimensional chromatography 913(2001)429

Immobilized trypsin

Proteolysis of whole cell extracts with immobilized enzyme columns as part of multidimensional chromatography 913(2001)429

Indirect detection

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Injection volume

Volume-load capacity in fast-gradient liquid chromatography. Effect of sample solvent composition and injection volume on chromatographic performance 913(2001)233

Indomethacin

Isolation of the components of a complex mixture by means of column switching for their enhanced detection by mass spectrometry 913(2001)209

Inorganic anions

Determination of inorganic cations and anions by ion-exchange chromatography with evaporative light-scattering detection 914(2001)167

Differential elution of sodium or potassium dihydrogen- and hydrogenphosphate ions from a Sephadex G-15 column with sodium or potassium chloride solution 914(2001)175

Inorganic cations

Determination of inorganic cations and anions by ion-exchange chromatography with evaporative light-scattering detection 914(2001)167

Differential elution of sodium or potassium dihydrogen- and hydrogenphosphate ions from a Sephadex G-15 column with sodium or potassium chloride solution 914(2001)175

Instrumentation

New micro-flow pumping system for liquid chromatography 913(2001)165

Interfaces, LC–MS

Preliminary investigation of the application of on-line membrane extraction of trifluoroacetic acid as an aid to improvement of negative ion electrospray mass spectrometry data 913(2001)205

Ion–ion interactions

Differential elution of sodium or potassium dihydrogen- and hydrogenphosphate ions from a Sephadex G-15 column with sodium or potassium chloride solution 914(2001)175

Ion-pairing reagents

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Isoelectric focusing

High-efficiency capillary isoelectric focusing of protein complexes from *Escherichia coli* cytosolic extracts 914(2001)257

Isoflavones

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Isotope-ratio mass spectrometry

Application of high-performance liquid chromatography with isotope-ratio mass spectrometry for measuring low levels of enrichment of underivatized materials 913(2001)269

Job plots

Study on the chiral recognition of the enantiomers of ephedrine derivatives with neutral and sulfated heptakis(2,3-*O*-diacetyl)- β -cyclodextrins using capillary electrophoresis, UV, nuclear magnetic resonance spectroscopy and mass spectrometry 914(2001)315

Kinetic studies

Sensitive liquid chromatographic–tandem mass spectrometric method for the determination of fluoxetine and its primary active metabolite norfluoxetine in human plasma 914(2001)45

Determination and pharmacokinetic study of unbound cefepime in rat bile by liquid chromatography with on-line microdialysis 914(2001)77

On-line microdialysis coupled with microbore liquid chromatography with ultraviolet detection for continuous monitoring of free cefsulodin in rat blood 914(2001)83

Lactate

Determination of pyruvate and lactate in primary liver cell culture medium during hypoxia by on-line microdialysis–liquid chromatography 913(2001)341

Simultaneous monitoring of extracellular glucose, pyruvate, lactate and glutamate in gerbil cortex during focal cerebral ischemia by dual probe microdialysis 913(2001)349

Laser-induced fluorescence detection

Some applications of near-ultraviolet laser-induced fluorescence detection in nanomolar- and subnanomolar-range high-performance liquid chromatography or micro-high-performance liquid chromatography 913(2001)253

Light-scattering detection

Fractionation and characterization of phenolic resins by high-performance liquid chromatography and gel-permeation chromatography combined with ultraviolet, refractive index, mass spectrometry and light-scattering detection 914(2001)147

Liposomes

Effect of liposome type and membrane fluidity on drug–membrane partitioning analyzed by immobilized liposome chromatography 913(2001)123

Liquid chromatography

Comparison of selected retention models in reversed-phase liquid chromatography 913(2001)3

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Liquid chromatography at critical conditions

Separation and quantification of the linear and cyclic structures of polyamide-6 at the critical point of adsorption 914(2001)131

Liquid chromatography, high-temperature

Superheated water as eluent in high-temperature high-performance liquid chromatographic separations of steroids on a polymer-coated zirconia column 913(2001)197

Liquid chromatography–mass spectrometry

Preliminary investigation of the application of on-line membrane extraction of trifluoroacetic acid as an aid to improvement of negative ion electrospray mass spectrometry data 913(2001)205

Liquid crystalline stationary phases

Laterally attached liquid-crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. III. Effect of the local anisotropic order on the separation of polycyclic aromatic hydrocarbons 913(2001)147

Liquid crystals

Laterally attached liquid-crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. III. Effect of the local anisotropic order on the separation of polycyclic aromatic hydrocarbons 913(2001)147

Loading capacity

Volume-load capacity in fast-gradient liquid chromatography. Effect of sample solvent composition and injection volume on chromatographic performance 913(2001)233

Loratadine

Sensitive liquid chromatography–tandem mass spectrometry method for the determination of loratadine and its major active metabolite descarboethoxyloratadine in human plasma 914(2001)37

Lutidines

Effect of counter-anion concentration on retention in high-performance liquid chromatography of protonated basic analytes 913(2001)189

2-Mercaptoethanesulfonic acid

Facile and sensitive method for the determination of mesna in plasma by high-performance liquid chromatography with ultraviolet detection 914(2001)29

Mass spectrometry

Isolation of the components of a complex mixture by means of column switching for their enhanced detection by mass spectrometry 913(2001)209

Application of high-performance liquid chromatography with isotope-ratio mass spectrometry for measuring low levels of enrichment of underivatized materials 913(2001)269

Rapid method for monitoring galactosylation levels during recombinant antibody production by electrospray mass spectrometry with selective-ion monitoring 913(2001)437

Mass transfer

Size-exclusion electrochromatography with controlled pore flow 914(2001)201

Mathematical modeling

Effect of counter-anion concentration on retention in high-performance liquid chromatography of protonated basic analytes 913(2001)189

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Mathematical modelling

Comparison of selected retention models in reversed-phase liquid chromatography 913(2001)3

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Matrix-assisted laser desorption ionization mass spectrometry

Artifact-free matrix-assisted laser desorption ionization time-of-flight mass spectra of *tert*.-butyldimethylsilyl ether derivatives of cyclodextrins used for the synthesis of single-isomer, chiral resolving agents for capillary electrophoresis 914(2001)325

Membranes

Effect of liposome type and membrane fluidity on drug–membrane partitioning analyzed by immobilized liposome chromatography 913(2001)123

Mesna

Facile and sensitive method for the determination of mesna in plasma by high-performance liquid chromatography with ultraviolet detection 914(2001)29

Metabolic disorders

Potential of capillary electrophoresis, tandem mass spectrometry and coupled capillary electrophoresis–tandem mass spectrometry as diagnostic tools 914(2001)265

Metals

Removal of metallic impurities from mobile phases in reversed-phase high-performance liquid chromatography by the use of an in-line chelation column 913(2001)159

Methanol

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Methionine sulfoximine

Development of an immobilized brain glutamine synthetase liquid chromatographic stationary phase for on-line biochemical studies 913(2001)133

Method development

Ranitidine hydrochloride: development of an isocratic stability-indicating high-performance liquid chromatographic separation 914(2001)13

Micellar electrokinetic chromatography

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

Micellar liquid chromatography

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

Micro liquid chromatography

New micro-flow pumping system for liquid chromatography 913(2001)165

Microdialysis

Determination of pyruvate and lactate in primary liver cell culture medium during hypoxia by on-line microdialysis–liquid chromatography 913(2001)341

Simultaneous monitoring of extracellular glucose, pyruvate, lactate and glutamate in gerbil cortex during focal cerebral ischemia by dual probe microdialysis 913(2001)349

Determination and pharmacokinetic study of unbound cefepime in rat bile by liquid chromatography with on-line microdialysis 914(2001)77

On-line microdialysis coupled with microbore liquid chromatography with ultraviolet detection for continuous monitoring of free cefsulodin in rat blood 914(2001)83

Mixed-mode chromatography

Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities 913(2001)113

Mobile phase composition

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Removal of metallic impurities from mobile phases in reversed-phase high-performance liquid chromatography by the use of an in-line chelation column 913(2001)159

Effect of the eluent pH and acidic modifiers in high-performance liquid chromatography retention of basic analytes 913(2001)173

Effect of counter-anion concentration on retention in high-performance liquid chromatography of protonated basic analytes 913(2001)189

Superheated water as eluent in high-temperature high-performance liquid chromatographic separations of steroids on a polymer-coated zirconia column 913(2001)197

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Capillary electrochromatographic evaluation of vitamin E-active oil constituents: tocopherols and tocotrienols 913(2001)379

Molecular descriptors

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Molecular imprinting

Uniformly sized molecularly imprinted polymer for (*S*)-naproxen. Retention and molecular recognition properties in aqueous mobile phase 913(2001)141

Molecular volumes

Geometry of chemically modified silica 913(2001)49

Monolithic column

Rapid separation of peptides and proteins by isocratic capillary electrochromatography at elevated temperature 914(2001)189

Monolithic columns

Acrylamide-based monoliths as robust stationary phases for capillary electrochromatography 914(2001)211

Multiple reaction monitoring

Potential of capillary electrophoresis, tandem mass spectrometry and coupled capillary electrophoresis–tandem mass spectrometry as diagnostic tools 914(2001)265

Myricetin

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Naproxen

Uniformly sized molecularly imprinted polymer for (*S*)-naproxen. Retention and molecular recognition properties in aqueous mobile phase 913(2001)141

Nash reagent

Improved measurement of formaldehyde in water-soluble polymers by high-performance liquid chromatography coupled with post-column reaction detection 914(2001)123

Neomycin

Sample preparation for residue determination of gentamicin and neomycin by liquid chromatography 914(2001)59

Neural networks

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Neuraminic acids

Determination of sialic acids by liquid chromatography–mass spectrometry 913(2001)365

Non-aqueous capillary electrophoresis

Hydrophobic interaction electrokinetic chromatography for the separation of polycyclic aromatic hydrocarbons using non-aqueous matrices 914(2001)223

Nonsteroidal anti-inflammatory drugs

Acrylamide-based monoliths as robust stationary phases for capillary electrochromatography 914(2001)211

Nonylphenol

Separation of isomers of nonylphenol and select nonylphenol polyethoxylates by high-performance liquid chromatography on a graphitic carbon column 914(2001)161

Nonylphenol polyethoxylates

Separation of isomers of nonylphenol and select nonylphenol polyethoxylates by high-performance liquid chromatography on a graphitic carbon column 914(2001)161

Norfluoxetine

Sensitive liquid chromatographic–tandem mass spectrometric method for the determination of fluoxetine and its primary active metabolite norfluoxetine in human plasma 914(2001)45

Nortriptyline

Isolation of the components of a complex mixture by means of column switching for their enhanced detection by mass spectrometry 913(2001)209

Nuclear magnetic resonance spectrometry

Study on the chiral recognition of the enantiomers of ephedrine derivatives with neutral and sulfated heptakis(2,3-*O*-diacetyl)- β -cyclodextrins using capillary electrophoresis, UV, nuclear magnetic resonance spectroscopy and mass spectrometry 914(2001)315

Nucleotides

Application of high-performance liquid chromatography with isotope-ratio mass spectrometry for measuring low levels of enrichment of underivatized materials 913(2001)269

Octadecylmethylglycerophosphodeoxyinositol

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Oligosaccharides

Rapid method for monitoring galactosylation levels during recombinant antibody production by electrospray mass spectrometry with selective-ion monitoring 913(2001)437

Optimization

Optimization of experimental conditions for the separation of small and large starch granules by gravitational field-flow fractionation 914(2001)183

Opuntia ficus-indica

Application of high-performance liquid chromatography to the characterization of the betalain pigments in prickly pear fruits 913(2001)415

Organic acids

Determination of pyruvate and lactate in primary liver cell culture medium during hypoxia by on-line microdialysis–liquid chromatography 913(2001)341

Simultaneous monitoring of extracellular glucose, pyruvate, lactate and glutamate in gerbil cortex during focal cerebral ischemia by dual probe microdialysis 913(2001)349

Organophosphorous compounds

Ion-exchange liquid chromatographic analysis of bisphosphonates by on-line post-column photochemical reaction and spectrophotometric detection 914(2001)105

Ovomucoid stationary phases

Bupropion hydrochloride: the development of a chiral separation using an ovomucoid column 913(2001)275

Peak trapping

Isolation of the components of a complex mixture by means of column switching for their enhanced detection by mass spectrometry 913(2001)209

Pentanone

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Peptides

Study on the degeneracy of antisense peptides using affinity chromatography 913(2001)423

Proteolysis of whole cell extracts with immobilized enzyme columns as part of multidimensional chromatography 913(2001)429

Rapid separation of peptides and proteins by isocratic capillary electrochromatography at elevated temperature 914(2001)189

Pesticides

Determination of quaternary ammonium pesticides by liquid chromatography–electrospray tandem mass spectrometry 914(2001)111

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

pH effects

Effect of the eluent pH and acidic modifiers in high-performance liquid chromatography retention of basic analytes 913(2001)173

Pharmaceutical analysis

Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities 913(2001)113

Isolation of the components of a complex mixture by means of column switching for their enhanced detection by mass spectrometry 913(2001)209

Appropriate column configurations for the rapid analysis and semipreparative purification of the radiolabeled drug flutamide by high-performance liquid chromatography 914(2001)23

Sensitive liquid chromatographic–tandem mass spectrometric method for the determination of fluoxetine and its primary active metabolite norfluoxetine in human plasma 914(2001)45

Determination and pharmacokinetic study of unbound cefepime in rat bile by liquid chromatography with on-line microdialysis 914(2001)77

On-line microdialysis coupled with microbore liquid chromatography with ultraviolet detection for continuous monitoring of free cefsulodin in rat blood 914(2001)83

Phenolic acids

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Phenolic resins

Fractionation and characterization of phenolic resins by high-performance liquid chromatography and gel-permeation chromatography combined with ultraviolet, refractive index, mass spectrometry and light-scattering detection 914(2001)147

Phenols

Fractionation and characterization of phenolic resins by high-performance liquid chromatography and gel-permeation chromatography combined with ultraviolet, refractive index, mass spectrometry and light-scattering detection 914(2001)147

Phenylureas

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

Phosphatidyl inositol

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Phospholipids

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Picoline

Effect of counter-anion concentration on retention in high-performance liquid chromatography of protonated basic analytes 913(2001)189

Pigments

Application of high-performance liquid chromatography to the characterization of the betalain pigments in prickly pear fruits 913(2001)415

Polyamide

Separation and quantification of the linear and cyclic structures of polyamide-6 at the critical point of adsorption 914(2001)131

Polymers

Improved measurement of formaldehyde in water-soluble polymers by high-performance liquid chromatography coupled with post-column reaction detection 914(2001)123

Fractionation and characterization of phenolic resins by high-performance liquid chromatography and gel-permeation chromatography combined with ultraviolet, refractive index, mass spectrometry and light-scattering detection 914(2001)147

Polynuclear aromatic hydrocarbons

Laterally attached liquid-crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. III. Effect of the local anisotropic order on the separation of polycyclic aromatic hydrocarbons 913(2001)147

Hydrophobic interaction electrokinetic chromatography for the separation of polycyclic aromatic hydrocarbons using non-aqueous matrices 914(2001)223

Polystyrene

Size-exclusion electrochromatography with controlled pore flow 914(2001)201

Pore flow

Size-exclusion electrochromatography with controlled pore flow 914(2001)201

Preparative chromatography

Appropriate column configurations for the rapid analysis and semipreparative purification of the radiolabeled drug flutamide by high-performance liquid chromatography 914(2001)23

Principal component analysis

Identification of the factors that influence the reproducibility of chromatographic retention data 913(2001)23

Profens

Uniformly sized molecularly imprinted polymer for (*S*)-naproxen. Retention and molecular recognition properties in aqueous mobile phase 913(2001)141

Propionic acids

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Proteins

Application of high-performance liquid chromatography with isotope-ratio mass spectrometry for measuring low levels of enrichment of underivatized materials 913(2001)269

Determination of sialic acids by liquid chromatography–mass spectrometry 913(2001)365

Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography–mass spectrometry 913(2001)397

Proteolysis of whole cell extracts with immobilized enzyme columns as part of multidimensional chromatography 913(2001)429

Rapid method for monitoring galactosylation levels during recombinant antibody production by electrospray mass spectrometry with selective-ion monitoring 913(2001)437

Rapid separation of peptides and proteins by isocratic capillary electrochromatography at elevated temperature 914(2001)189

High-efficiency capillary isoelectric focusing of protein complexes from *Escherichia coli* cytosolic extracts 914(2001)257

Protriptyline

High-performance liquid chromatographic analysis for a non-chromophore-containing phosphatidyl inositol analog, 1-[(1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero)-phospho]-1*D*-3-deoxy-*myo*-inositol, using indirect UV detection 913(2001)355

Pumps

New micro-flow pumping system for liquid chromatography 913(2001)165

Purines

Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities 913(2001)113

Pyrene

Some applications of near-ultraviolet laser-induced fluorescence detection in nanomolar- and subnanomolar-range high-performance liquid chromatography or micro-high-performance liquid chromatography 913(2001)253

Pyridines

Effect of the eluent pH and acidic modifiers in high-performance liquid chromatography retention of basic analytes 913(2001)173

Effect of counter-anion concentration on retention in high-performance liquid chromatography of protonated basic analytes 913(2001)189

Liquid chromatography–tandem mass spectrometry method of urine analysis for determining human variation in carcinogen metabolism 914(2001)95

Pyrimidines

Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities 913(2001)113

Pyruvate

Determination of pyruvate and lactate in primary liver cell culture medium during hypoxia by on-line microdialysis–liquid chromatography 913(2001)341

Simultaneous monitoring of extracellular glucose, pyruvate, lactate and glutamate in gerbil cortex during focal cerebral ischemia by dual probe microdialysis 913(2001)349

Quaternary ammonium compounds

Determination of quaternary ammonium pesticides by liquid chromatography–electrospray tandem mass spectrometry 914(2001)111

Quercetin

Separation and determination of flavonoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography 913(2001)387

Raman spectrometry

Raman spectroscopy of octadecylsilane stationary phase conformational order. Effect of solvent 913(2001)243

Ranitidine hydrochloride

Ranitidine hydrochloride: development of an isocratic stability-indicating high-performance liquid chromatographic separation 914(2001)13

Refractive index detection

Fractionation and characterization of phenolic resins by high-performance liquid chromatography and gel-permeation chromatography combined with ultraviolet, refractive index, mass spectrometry and light-scattering detection 914(2001)147

Regression analysis

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Reproducibility

Identification of the factors that influence the reproducibility of chromatographic retention data 913(2001)23

Reserpine

Liquid chromatographic–tandem mass spectrometric determination of amprevir (agenerase) in serum/plasma of human immunodeficiency virus type-1 infected patients receiving combination antiretroviral therapy 914(2001)1

Retention behavior

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Retention behaviour

Effect of the eluent pH and acidic modifiers in high-performance liquid chromatography retention of basic analytes 913(2001)173

Retention data

Identification of the factors that influence the reproducibility of chromatographic retention data 913(2001)23

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. V. Results obtained with Vydac 218TP C₁₈ columns 913(2001)89

Retention factor

Volume-load capacity in fast-gradient liquid chromatography. Effect of sample solvent composition and injection volume on chromatographic performance 913(2001)233

Retention indices

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

Retention models

Comparison of selected retention models in reversed-phase liquid chromatography 913(2001)3

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Reversed-phase liquid chromatography

Comparison of selected retention models in reversed-phase liquid chromatography 913(2001)3

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Review

Derivatization and chromatographic behavior of the *o*-phthalaldehyde amino acid derivatives obtained with various SH-group-containing additives (Review) 913(2001)283

Sample solvent strength

Volume-load capacity in fast-gradient liquid chromatography. Effect of sample solvent composition and injection volume on chromatographic performance 913(2001)233

Sequencing methods

On-line nanoliter cycle sequencing reaction with capillary zone electrophoresis purification for DNA sequencing 914(2001)245

Sialic acids

Determination of sialic acids by liquid chromatography–mass spectrometry 913(2001)365

Silica

Geometry of chemically modified silica 913(2001)49

Silica, urea-functionalized

Development of new urea-functionalized silica stationary phases. Characterization and chromatographic performance 913(2001)65

Size exclusion electrochromatography

Size-exclusion electrochromatography with controlled pore flow 914(2001)201

Solid-phase extraction

Effects of the matrix and sample preparation on the determination of fluoroquinolone residues in animal tissues 914(2001)89

Solvation

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Solvent effects

Raman spectroscopy of octadecylsilane stationary phase conformational order. Effect of solvent 913(2001)243

Stability studies

Ranitidine hydrochloride: development of an isocratic stability-indicating high-performance liquid chromatographic separation 914(2001)13

Starch

Optimization of experimental conditions for the separation of small and large starch granules by gravitational field-flow fractionation 914(2001)183

Stationary phases, electrochromatography

Acrylamide-based monoliths as robust stationary phases for capillary electrochromatography 914(2001)211

Stationary phases, LC

Identification of the factors that influence the reproducibility of chromatographic retention data 913(2001)23

Geometry of chemically modified silica 913(2001)49

Development of new urea-functionalized silica stationary phases. Characterization and chromatographic performance 913(2001)65

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. V. Results obtained with Vydac 218TP C₁₈ columns 913(2001)89

Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities 913(2001)113

Development of an immobilized brain glutamine synthetase liquid chromatographic stationary phase for on-line biochemical studies 913(2001)133

Laterally attached liquid-crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. III. Effect of the local anisotropic order on the separation of polycyclic aromatic hydrocarbons 913(2001)147

Superheated water as eluent in high-temperature high-performance liquid chromatographic separations of steroids on a polymer-coated zirconia column 913(2001)197

Raman spectroscopy of octadecylsilane stationary phase conformational order. Effect of solvent 913(2001)243

Bupropion hydrochloride: the development of a chiral separation using an ovomucoid column 913(2001)275

Steroids

Superheated water as eluent in high-temperature high-performance liquid chromatographic separations of steroids on a polymer-coated zirconia column 913(2001)197

Determination of apparent association constants of steroid–cyclodextrin inclusion complexes using a modification of the Hummel–Dreyer method 913(2001)261

***Streptomyces coelicolor* A3(2)**

Recovery of actinorhodin from fermentation broth 914(2001)67

Structure–enantioselectivity relationships

Approaches to quantitative structure–enantioselectivity relationship modeling of chiral separations using capillary electrophoresis 914(2001)299

Sulphur compounds

Determination of biologically active low-molecular-mass thiols in human blood. III. Highly sensitive narrow-bore isocratic reversed-phase high-performance liquid chromatography with fluorescence detection 913(2001)315

Superheated water chromatography

Superheated water as eluent in high-temperature high-performance liquid chromatographic separations of steroids on a polymer-coated zirconia column 913(2001)197

Surfactants

Separation of isomers of nonylphenol and select nonylphenol polyethoxylates by high-performance liquid chromatography on a graphitic carbon column 914(2001)161

Tamoxifen

Analysis of tamoxifen and its metabolites in synthetic gastric fluid digests and urine samples using high-performance liquid chromatography with electrospray mass spectrometry 914(2001)5

***tert*-butyldimethylsilylcyclodextrins**

Artifact-free matrix-assisted laser desorption ionization time-of-flight mass spectra of *tert*-butyldimethylsilyl ether derivatives of cyclodextrins used for the synthesis of single-isomer, chiral resolving agents for capillary electrophoresis 914(2001)325

Testosterone

Superheated water as eluent in high-temperature high-performance liquid chromatographic separations of steroids on a polymer-coated zirconia column 913(2001)197

Tetrahydrofuran

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Thermodynamic parameters

Modeling the effect of solvation on solute retention in reversed-phase liquid chromatography 913(2001)15

Thiols

Determination of biologically active low-molecular-mass thiols in human blood. III. Highly sensitive narrow-bore isocratic reversed-phase high-performance liquid chromatography with fluorescence detection 913(2001)315

Facile and sensitive method for the determination of mesna in plasma by high-performance liquid chromatography with ultraviolet detection 914(2001)29

Thiomalic acid

Facile and sensitive method for the determination of mesna in plasma by high-performance liquid chromatography with ultraviolet detection 914(2001)29

Tocopherols

Capillary electrochromatographic evaluation of vitamin E-active oil constituents: tocopherols and tocotrienols 913(2001)379

Tocotrienols

Capillary electrochromatographic evaluation of vitamin E-active oil constituents: tocopherols and tocotrienols 913(2001)379

Tolbutamide

Isolation of the components of a complex mixture by means of column switching for their enhanced detection by mass spectrometry 913(2001)209

Toluene

Interpretation of the excess adsorption isotherms of organic eluent components on the surface of reversed-phase adsorbents. Effect on the analyte retention 913(2001)75

Triazines

Characterisation of retention in micellar high-performance liquid chromatography, in micellar electrokinetic chromatography and in micellar electrokinetic chromatography with reduced flow 914(2001)233

Trifluoroacetic acid

Preliminary investigation of the application of on-line membrane extraction of trifluoroacetic acid as an aid to improvement of negative ion electrospray mass spectrometry data 913(2001)205

Troleandomycin

Liquid chromatography of troleandomycin 914(2001)53

Validation

Validation of a chromatography data analysis software 913(2001)221

Liquid chromatographic–tandem mass spectrometric determination of amprenavir (agenerase) in serum/plasma of human immunodeficiency virus type-1 infected patients receiving combination antiretroviral therapy 914(2001)1

Sensitive liquid chromatography–tandem mass spectrometry method for the determination of loratadine and its major active metabolite descarboethoxyloratadine in human plasma 914(2001)37

Sensitive liquid chromatographic–tandem mass spectrometric method for the determination of fluoxetine and its primary active metabolite norfluoxetine in human plasma 914(2001)45

Vinylpyridine

Uniformly sized molecularly imprinted polymer for (*S*)-naproxen. Retention and molecular recognition properties in aqueous mobile phase 913(2001)141

Vitamins

Capillary electrochromatographic evaluation of vitamin E-active oil constituents: tocopherols and tocotrienols 913(2001)379

Vydac 218TP C₁₈ columns

Repeatability and reproducibility of retention data and band profiles on reversed-phase liquid chromatography columns. V. Results obtained with Vydac 218TP C₁₈ columns 913(2001)89

Water analysis

Determination of quaternary ammonium pesticides by liquid chromatography–electrospray tandem mass spectrometry 914(2001)111

Zirconia–polybutadiene stationary phases

Superheated water as eluent in high-temperature high-performance liquid chromatographic separations of steroids on a polymer-coated zirconia column 913(2001)197