ELSEVIER

Journal of Chromatography A, 940 (2001) B119–B239

JOURNAL OF CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

985 Gilar, M., Bouvier, E.S.P. and Compton, B.J.: Advances in sample preparation in electromigration, chromatographic and mass spectrometric separation methods. *J. Chromatogr. A,* 909 (2001) 111-135 - a review with 226 refs.

986 Kirkland, J.J.: Ultrafast reversed-phase high-performance liquid chromatographic separations: An overview. *J. Chromatogr. Sci.,* 38 (2000) 535-544.

987 Leon-Gonzalez, M.E. and Perez-Arribas, L.V.: Chemically modified polymeric sorbents for sample preconcentration. *J. Chromatogr. A,* 902 (2000) 3-16 - a review with 42 refs.

988 Miyabe, K. and Guiochon, G.: Fundamental interpretation of the peak profiles in linear reversed-phase liquid chromatography. *Adv. Chromatogr. (N.Y.),* 40 (2000) 1-113; *C.A.,* 133 (2000) 243104f - a review with 126 refs.

989 Scott, R.P.W.: Moving bed chromatography. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3069-3082.

990 Siouffi, A.-M.: HPLC. *Food Sci. Technol. (N.Y.),* 100(Food Analysis by HPLC; 2nd Edition) (2000) 1-54; *C.A.,* 133 (2000) 368856j - a review with 29 refs.

991 Sparr Eskilsson, C. and Björklund, E.: Analytical-scale microwave-assisted extraction. *J. Chromatogr. A,* 902 (2000) 227-250 - a review with 139 refs.

See also 994, 1000, 1032, 1044, 1051, 1054, 1071, 1078, 1098, 1099, 1109, 1111, 1113, 1128, 1130, 1140, 1141, 1142, 1143, 1144, 1149, 1150, 1151, 1152, 1153, 1157, 1158, 1159, 1160, 1161, 1165, 1256, 1276, 1347, 1446, 1451, 1601, 1665, 1691, 1747, 1764, 1799, 1815, 1819, 1821, 1825, 1926, 1927, 1946, 2025, 2029, 2030, 2037, 2039, 2045, 2088, 2104.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

992 Canals, I., Oumada, F.Z., Roses, M. and Bosch, E.: Retention of ionizable compounds on HPLC. 6. pH measurements with the glass electrode in methanol-water mixtures. *J. Chromatogr. A,* 911 (2001) 191-202.

993 Cherrak, D.E. and Guiochon, G.: Phenomenological study of the bed-wall friction in axially compressed packed chromatographic columns. *J. Chromatogr. A,* 911 (2001) 147-166.

994 Cordero, B.M., Perez Pavon, J.L., Garcia Pinto, C., Fernandez Laespada, E., Carabias Martinez, R. and Rodriguez Gonzalo, E.: Analytical applications of membrane extraction in chromatography and electrophoresis. *J. Chromatogr. A,* 902 (2000) 195-204 - a review with 103 refs.

995 Dolan, J.W.: Reducing column diameter in gradient elution - a case study. *LC-GC Eur.,* 14 (2001) 14-17.

996 Dolan, J.W.: Starting out right, Part V - Changing column conditions. *LC-GC Eur.,* 13 (2000) 302-308.

997 Dolan, J.W.: Unexpected peaks. *LC-GC Eur.,* 14 (2001) 76-81.

998 Du, C.M., Valko, K., Bevan, C., Reynolds, D. and Abraham, M.H.: Characterizing the selectivity of stationary phases and organic modifiers in reversed-phase high-performance liquid chromatographic systems by a general solvation equation using gradient elution. *J. Chromatogr. Sci.,* 38 (2000) 503-511.

999 Halmagyi, B. and Holz, B.: (HPLC in a minimum of space). *LaborPraxis,* 24 (2000) 50-51; *C.A.,* 133 (2000) 246610d.

1000 Jönsson, J.Å. and Mathiasson, L.: Membrane-based techniques for sample enrichment. *J. Chromatogr. A,* 902 (2000) 205-225 - a review with 106 refs.

1001 Kalász, H.: Welcome on board. *LC-GC Eur.,* 14 (2001) 12.

1002 Kanazawa, H., Sunamoto, T., Matsushima, Y., Kikuchi, A. and Okano, T.: Temperature-responsive chromatographic separation of amino acid phenylthiohydantoins using aqueous media as the mobile phase. *Anal. Chem.,* 72 (2000) 5961-5966.

1003 Kehl, K.G. and Meyer, V.R.: Argentometic titration for the determination of liquid chromatographic injection reproducibility. *Anal. Chem.,* 73 (2001) 131-133.

1004 Lewis, R. and Dolan, J.W.: The problem of late elution - A case study. *LC-GC Eur.,* 13 (2000) 882-886.

1005 Merdink, J.L. and Dolan, J.W.: Scaling gradient LC methods to LC-MS. *LC-GC Eur.,* 13 (2000) 806-811.

1006 Nigam, S., Stephens, M., de Juan, A. and Rutan, S.C.: Characterization of the polarity of reversed-phase liquid chromatography stationary phases in the presence of 1-propanol using solvatochromism and multivariate curve resolution. *Anal. Chem.,* 73 (2001) 290-297.

1007 Pace, G., Berton, A. and Bergamaschi, L.: Bi-quadratic surface response for quantitative determination of analytes leading to partially overlapped chromatographic peaks. *J. Chromatogr. A,* 907 (2001) 81-88.

1008 Wu, N., Lippert, J.A. and Lee, M.L.: Practical aspects of ultrahigh pressure capillary liquid chromatography. *J. Chromatogr. A,* 911 (2001) 1-12.

0021-9673/01/$ - see front matter © 2001 Elsevier Science B.V. All rights reserved.
PII: S0021-9673(01)00891-3

See also 991, 1075, 1106, 1110, 1115, 1128.

2b. Thermodynamics and theoretical relationships

1009 Baczek, T., Markuszewski, M., Kaliszan, R., van Straten, M. and Claessens, H.A.: Linear and quadratic relationships between retention and organic modifier content in eluent in reversed phase high-performance liquid chromatography: A systematic comparative statistical study. *J. High Resolut. Chromatogr.,* 23 (2000) 667-676.

1010 Brinkman, U.A.Th., Hill, H.M., Borner, K. and Reid, E.: Unorthodox temperature conditions: opinions and supplementary observations from bioanalytical forum participants. *Chromatographia,* 52(Supplement) (2000) S57-S59.

1011 Guillaume, Y.C., Guinchard, C., Robert, J.F. and Berhelot, A.: Ionic binding of human serum albumin - dependence on pH and ionic strength: a chromatographic approach. *Chromatographia,* 52 (2000) 575-578.

1012 Jandera, P., Skavrada, M., Andel, L., Komers, D. and Guiochon, G.: Description of adsorption equilibria in liquid chromatography systems with binary mobile phases. *J. Chromatogr. A,* 908 (2001) 3-17.

1013 Kwietniewski, L., Javicka, H. and Rozylo, J.K.: Modelling of RP chromatographic systems. *Chromatographia,* 52 (2000) 559-563.

1014 Lisec, O., Hugo, P. and Seidel-Morgenstern, A.: Frontal analysis method to determine competitive adsorption isotherms. *J. Chromatogr. A,* 908 (2001) 19-34.

1015 Meyers, J.J., Crosser, O.K. and Liapis, A.I.: Pore network modelling: determinaion of the dynamic profiles of the pore diffusivity and its effect on column performance as the loading of the solute in the adsorbed phase varies with time. *J. Chromatogr. A,* 908 (2001) 35-47.

1016 Miyabe, K. and Guiochon, G.: Influence of the column radial heterogeneity on the determination of single-component isotherms by the elution by characteristic point method. *J. Chromatogr. A,* 891 (2000) 217-233.

1017 Nikitas, P. and Pappa-Louisi, A.: Non-linear least-squares fitting with microsoft excel solver and related routines in HPLC modelling of retention I. Considerations of the problems of the method. *Chromatographia,* 52 (2000) 477-486.

1018 Pappa-Louisi, A. and Nikitas, P.: Non-linear least-squares fitting with micrisoft excel solver and related routines in HPLC modelling of retention II. Considerations of selection of optimal fit. *Chromatographia,* 52 (2000) 487-493.

1019 Ranatunga, R.P.J. and Carr, P.W.: A study of the enthalpy and entropy contributions of the stationary phase in reversed-phase liquid chromatography. *Anal. Chem.,* 72 (2000) 5679-5692.

1020 Ravald, L. and Fornstedt, T.: Theoretical study of the accuracy of the elution by characteristic points method for bi-Langmuir isotherms. *J. Chromatogr. A,* 908 (2001) 111-130.

1021 Reid, E.: Use of non-ambient temperatures in separatory runs. *Chromatographia,* 52(Supplement) (2000) S55-S56.

1022 Rudakov, O.B. and Bochrova, O.N.: (Boiling point of binary mobile phases for high-performance liquid chromatography). *Zh. Fiz. Khim.,* 74 (2000) 1093-1098; *C.A.,* 133 (2000) 272217a.

1023 Sandi, A., Nagy, M. and Szepesy, L.: Characterization of reversed-phase columns using the linear free energy relationship. III. Effect of the organic modifier and the mobile phase composition. *J. Chromatogr. A,* 893 (2000) 215-234.

1024 Scott, R.P.W.: The thermodynamic properties of tetrahydrofuran-water association and its influence on retention in liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3083-3093.

1025 Van der Hoff, G.R., Hoogerbrugge, R., Baumann, R.A., Brinkman, U.A.Th. and van Zoonen, P.: An empirical retention model for the optimization of on-line normal-phase LC-GC for the multi-analyte determination of hydrophobic compounds in fatty samples. *Chromatographia,* 52 (2000) 433-438.

See also 988, 1027, 1028, 1090, 1091, 1093, 1094, 1106, 1147, 1156, 1163, 1181, 1405, 1435, 1697, 1884, 1902.

2c. Relationship between structure and chromatographic behaviour

1026 Dzido, T.H.: Modifier influence on selectivity of reversed-phase HPLC systems. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2773-2788.

1027 Torrens, F.: Calculation of partition coefficient and hydrophobic moment of he secondary structure of lysozyme. *J. Chromatogr. A,* 908 (2001) 215-221.

See also 992, 1009, 1386, 1671, 2092.

2d. Measurement of physico-chemical and related values

1028 Arvanitopoulou, E., Nikolaropoulos, S.S., Bazoti, F. and Katsanos, N.A.: Physicochemical measurements in liquids by simple HPLC instrumentation. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2911-2928.

1029 Romanelli, G.P., Cafferata, L.R.F. and Castro, E.A.: Application of improved quantitative structure retention relationships to study the molecular mechanism of stationary phases for HPLC. *Khim. Fiz.,* 19, No. 9 (2000) 105-109; *C.A.,* 134 (2001) 21883r.

1030 Storm, T., Hartig, C., Reemtsma, T. and Jekel, M.: Exact mass measurements on-line with high-performance liquid chromatography on a quadrupole mass spectrometer. *Anal. Chem.,* 73 (2001) 589-595.

1031 Trapp, O. and Schurig, V.: Approximation function for the direct calculation of rate constants and Gibbs activation energies of enantiomerization of racemic mixtures from chromatographic parameters in dynamic chromatography. *J. Chromatogr. A,* 911 (2001) 167-175.

See also 992, 1011, 1015, 1018, 1342, 1435.

3. GENERAL TECHNIQUES

1032 Sessler, J.L., Genge, J.W., Gale, P.A. and Kral, V.: Calix[4]pyrrole-functionalized silica gels: novel supports for the HPLC-based separation of anions. *ACS Symp. Ser.,* 757(Calixarenes for Separations) (2000) 238-254; *C.A.,* 133 (2000) 316962w - a review with 34 refs.

3a. Apparatus and accessories

1033 Baker, T.R. and Pinkston, J.D.: Preliminary investigation of supercritical-fluid-assisted nebulization for enhanced response in electrospray mass spectrometry. *Analusis,* 27 (1999) 701-705; *C.A.*, 134 (2001) 50804w.

1034 Jönsson, J.A. and Mathiasson, L.: Membrane extraction techniques in bioanalysis. *Chromatographia,* 52(Supplement) (2000) S8-S11.

1035 Küppers, S., Renger, B. nad Meyer, V.R.: In response. *LC-GC Eur.,* 13 (2000) 300.

1036 Paul, P.H.: Electrokinetic pumps and actuators. *Sandia Natl. Lab. [Tech. Rep.] SAND,* (SAND2000-8218) (2000) 5-30; *C.A.*, 133 (2000) 272209z.

1037 Rule, G., Chapple, M. and Henion, J.: A 384-well solid-phase extraction for LC/MS/MS determination of methotrexate and its 7-hydroxy metabolite in human urine and plasma. *Anal. Chem.,* 73 (2001) 439-443.

1038 Tommasi, R.A., Whaley, L.W. and Marepalli, H.R.: AutoChem: Automated solution-phase parallel synthesis and purification via HPLC. *J. Comb. Chem.,* 2 (2000) 447-449; *C.A.*, 133 (2000) 252086d.

1039 Wu, J. and Pawliszyn, J.: Polypyrrole-coated capillary coupled to HPLC for in-tube solid-phase microextraction and analysis of aromatic compounds in aqueous samples. *Anal. Chem.,* 73 (2001) 55-63.

1040 Zhang, X. and Huang, S.: Single step on-column frit making for capillary high-performance liquid chromatography using sol-gel technology. *J. Chromatogr. A,* 910 (2001) 13-18.

See also 1083, 1084, 1101, 1318, 1784, 1817.

3b. Detectors and detection reagents

1041 Jin, J., Miwa, T., Shi, G., Zhang, W. and Jin, L.: (Study of electrochemical detector for micro high performance liquid chromatography and capillary electrophoresis (CE). *Huaxue Chuanganqi,* 20, No. 2 (2000) 18-24; *C.A.*, 134 (2001) 10680w.

1042 Peters, H.L., Levine, K.E. and Jones, B.T.: An inductively coupled plasma carbon emission detector for aqueous carbohydrate separations by liquid chromatography. *Anal. Chem.,* 73 (2001) 453-457.

1043 Torabi, K., Karami, A., Balke, S.T. and Schunk, T.C.: Quantitative FTIR detection in size-exclusion chromatography. *J. Chromatogr. A,* 910 (2001) 19-30.

1044 Wei, Y. and Ding, M.-y.: (Evaporative light-scattering detection technology). *Sepu,* 18 (2000) 398-401; *C.A.*, 133 (2000) 358677n - a review with 45 refs.

See also 1033, 1186, 1187, 1196, 1201, 1235, 1243, 1246, 1250, 1289, 1310, 1351, 1367, 1372, 1468, 1476, 1629, 1652, 1664, 1685, 1901, 2007, 2043, 2089, 2094.

3c. Sorbents and columns, packing procedures

1045 Castells, C.B. and Carr, P.W.: A study of the thermodynamics and influence of temperature on chiral high-performance liquid chromatographic separations using cellulose tris(3,5-dimethylphenylcarbamate) coated zirconia stationary phases. *Chromatographia,* 52 (2000) 535-542.

1046 Chen, X., Yang, L., Zou, H., Zhang, Q. and Ni, J.: (Preparation of the chmically bonded cellulose phenylcarbamate chiral stationary phases for the separation of enantiomers). *Fenxi Huaxue,* 28 (2000) 1074-1078; *C.A.*, 134 (2001) 24996r.

1047 Dolan, J.W., Snyder, L.R. and Blanc, T.: Selectivity differences for C_{18} and C_8 reversed-phase columns as a function of temperature and gradient steepness. II. Minimizing column reproducibility problems. *J. Chromatogr. A,* 897 (2000) 51-63.

1048 Dolan, J.W., Snyder, L.R. and van Heukelem, L.: Selectivity differences for C_{18} and C_8 reversed-phase columns as a function of temperature and gradient steepness. I. Optimizing selectivity and resolution. *J. Chromatogr. A,* 897 (2000) 37-50.

1049 Du, J., Zhang, X.-S. and Zhang, H.: (Study on new packing material in low-pressure ion chromatography). *Sichuan Daxue Xuebao, Gongcheng Kexueban,* 32, No. 2 (2000) 48-50; *C.A.*, 133 (2000) 174107v.

1050 Gritti, F., Felix, G., Achard, M.F. and Hardouin, F.: Laterally attached liquid crystalline polymers as stationary phases in reversed-phase high-performance liquid chromatography. II. Optimization of the molecular parameter of the polymer. *J. Chromatogr. A,* 897 (2000) 131-143.

1051 Gruner, R. and Engelhardt, H.: (Use and characterization of polar reversed-phase (RP) HPLC phases). *LaborPraxis,* 24, No. 9 (2000) 40-45; *C.A.*, 133 (2000) 343970d - a review with 22 refs.

1052 Guo, Y., Yang, Y. and Su, T.: (Preparation and characterization of monosized crosslinked PMMA resins with hydrophilic surface). *Gaofenzi Xuebao,* No.5 (2000) 641-643; *C.A.*, 134 (2001) 57325z.

1053 Hahn, R. and Jungbauer, A.: Control method for integrity of continuous beds. *J. Chromatogr. A,* 908 (2001) 179-184.

1054 Huang, Y. and Mou, S.-f.: (Recent advances in the ion chromatographic stationary phases). *Sepu,* 18 (2000) 412-419; *C.A.*, 133 (2000) 358678p - a review with 96 refs.

1055 Hunter, A.K. and Carta, G.: Protein adsorption on novel acrylamido-based polymeric ion exchangers. II. Adsorption rates and column behavior. *J. Chromatogr. A,* 897 (2000) 81-97.

1056 Hunter, A.K. and Carta, G.: Protein adsorption on novel acrylamido-based polymeric ion-exchangers. I. Morphology and equilibrium adsorption. *J. Chromatogr. A,* 897 (2000) 65-80.

1057 Jiang, Z.-T. and Zuo, Y.-M.: Synthesis of porous titania microspheres for HPLC packings by polymerization-induced colloid aggregation (PICA). *Anal. Chem.,* 73 (2001) 686-688.

1058 Ljunglöf, A., Larsson, M., Knuuttila, K.-G. and Lindgren, J.: Measurement of ligand distribution in individual adsorbent particles using confocal scanning laser microscopy and confocal micro-Raman spectroscopy. *J. Chromatogr. A,* 893 (2000) 235-244.

1059 Ludemann-Hombourger, O., Bailly, M. and Nicoud, R.M.: Design of a simulated moving bed: optimal particle size of the stationary phase. *Sep. Sci. Technol.,* 35 (2000) 1285-1305; *C.A.*, 133 (2000) 271600h.

1060 Massolini, G., de Lorenzi, E., Calleri, E., Bertucci, C., Monaco, H.L., Pereuca, M., Caccialanza, G. and Wainer, I.W.: Properties of a stationary phase based on immobilised chicken liver basic fatty acid-binding protein. *J. Chromatogr. B,* 751 (2001) 117-130.

1061 Mayfield, S., Cabrera, K. and Lubda, D.: Chromoligh SpeedROD - a new monolithic HPLC column for rapid separations. *CAST, Chromatogr. Sep. Technol.,* 15 (2000) 5-6; *C.A.,* 134 (2001) 80318h.

1062 Mayr, B. and Buchmeiser, M.R.: Influences of surface chemistry on the separation behavior of stationary phases for reversed-phase and ion-exchange chromatography: a comparison of coated and grafted supports prepared by ring-opening metathesis polymerization. *J. Chromatogr. A,* 907 (2001) 73-80.

1063 Mayr, B., Sinner, F. and Buchmeiser, M.R.: Chiral β-cyclodextrin-based polymer supports prepared via ring-opening metathesis graft-polymerization. *J. Chromatogr. A,* 907 (2001) 47-56.

1064 Miyabe, K. and Guiochon, G.: Influence of the modification conditions of alkyl bonded ligands on the characteristics of reversed-phase liquid chromatography. *J. Chromatogr. A,* 903 (2000) 1-12.

1065 O'Gara, J.E., Alden, B.A., Gendreau, C.A., Iraneta, P.C. and Walter, T.H.: Dependence of cyano bonded phase hydrolytic stability on ligand structure and solution pH. *J. Chromatogr. A,* 893 (2000) 245-251.

1066 Ousalem, M., Zhu, X.X. and Hradil, J.: Evaluation of the porous structures of new polymer packing materials by inverse size-exclusion chromatography. *J. Chromatogr. A,* 903 (2000) 13-19.

1067 Pesek, J.J., Matyska, M.T. and Muley, S.: Synthesis and characterization of a new type of chemically bonded liquid crystal stationary phase for HPLC. *Chromatographia,* 52 (2000) 439-444.

1068 Podgornik, A., Barut, M., Strancar, A., Josic, D. and Koloini, T.: Construction of large-volume monolithic columns. *Anal. Chem.,* 72 (2000) 5693-5699.

1069 Pursch, M., Vanderhart, D.L., Sander, L.C., Gu, X., Nguyen, T., Wise, S.A. and Gajewski, D.A.: C_{30} Self-assembled monolayers on silica, titania, and zirconia: HPLC performance, atomic force microscopy, ellipsometry, and NMR studies of molecular dynamics and uniformity of coverage. *J. Am. Chem. Soc.,* 122 (2000) 6997-7011; *C.A.,* 133 (2000) 199220q.

1070 Ross, P.: The role of porous graphitic carbon in HPLC. *LC-GC Eur.,* 13 (2000) 310-319.

1071 Ross, P. and Ax, O.: (Porous graphitized carbon for HPLC and solid-phase extraction (SPE)). *LaborPraxis,* 24, No. 9 (2000) 28-35; *C.A.,* 133 (2000) 343969k - a review with 26 refs.

1072 Schmiding, T. and Malordy, B.: (On the way to development of a universal HPLC column). *LaborPraxis,* 24, No. 9 (2000) 36-39; *C.A.,* 133 (2000) 343982j.

1073 Staby, A. and Jensen, I.H.: Comparison of chromatographic ion-exchange resins II. More strong anion-exchange resins. *J. Chromatogr. A,* 908 (2001) 149-161.

1074 Staby, A., Jensen, I.H. and Mollerup, I.: Comparison of chromatographic ion-exchange resins. I. Strong anion-exchange resins. *J. Chromatogr. A,* 897 (2000) 99-111.

1075 Turowski, M., Morimoto, T., Kimata, K., Monde, H., Ikegami, T., Hosoya, K. and Tanaka, N.: Selectivity of stationary phases in reversed-phase liquid chromatography based on the dispersion interactions. *J. Chromatogr. A,* 911 (2001) 177-190.

1076 Viklund, C., Sjögren, A., Irgum, K. and Nes, I.: Chromatographic interactions between proteins and sulfoalkylbetaine-based zwitterionic copolymers in fully aqueous low-salt buffers. *Anal. Chem.,* 73 (2001) 444-452.

1077 Willetts, M., Brown, V.A. and Clench, M.R.: Influence of surface carbon coverage of C_1 (TMS) stationary phases on the separation of nonylphenol ethoxylate ethoxymers. *J. Chromatogr. A,* 903 (2000) 33-40.

1078 Yang, X.-l., Wang, J.-d. and Xiong, B.-h.: (Recent advances of silica packing in high performance liquid chromatography). *Sepu,* 18 (2000) 308-312; *C.A.,* 133 (2000) 187215d - a review with 65 refs.

1079 Yu, X., Zhao, R. and Liu, G.Q.: A novel method for the preparation of C_{18} ester-bonded RP-HPLC packings. *Chromatographia,* 52 (2000) 517-519.

1080 Zeng, R., Pang, Z. and Zhu, H.: Modification of a Nafion® ion exchange membrane by a plasma polymerization process. *J. Electroanal. Chem.,* 490 (2000) 102-106; *C.A.,* 134 (2001) 80233b.

1081 Zhang, C., Zhang, L., Yang, J. and Liu, Z.: (Application of high performance gel permeation chromatography in characterization of a number of heavy industry chemical products). *Fenxi Huaxue,* 28 (2000) 1248-1251; *C.A.,* 134 (2001) 80319j.

1082 Zhang, Q., Zou, H., Chen, X., Wang, H., Luo, Q. and Ni, J.: Synthesis and characteristics of the human serum albumin-triazine chiral stationary phase. *Chirality,* 12 (2000) 714-719; *C.A.,* 134 (2001) 33075s.

1083 Zhu, C., Guan, Y., Wang, H. and Zhu, D.: (Packed capillary liquid chromatographic columns with conical ends). *Fenxi Huaxue,* 28 (2000) 1204-1208; *C.A.,* 133 (2000) 358691n.

1084 Zhu, C.-r., Guan, Y.-f. and Wu, J.-q.: (Conical column end for wide bore packed capillary liquid chromatographic column). *Sepu,* 18 (2000) 383-386; *C.A.,* 133 (2000) 358689t.

1085 Ziao, X.-Z., Feng, Y.-Q., Da, S.-L. and Zhang, Y.: Preparation and evaluation of *p-tert.*-butyl-calix[8]arene-bonded silica stationary phase for high performance liquid chromatography. *Anal. Lett.,* 33 (2000) 3355-3372.

1086 Zobel, H.: (Improvement of the applicability of reversed phases). *GIT Spez. Sep.,* 20, No. 2 (2000) 72-74; *C.A.,* 133 (2000) 343981h.

See also 986, 995, 996, 1006, 1011, 1015, 1019, 1091, 1106, 1124, 1134, 1138, 1140, 1142, 1143, 1144, 1145, 1152, 1155, 1157, 1164, 1170, 1181, 1341, 1363, 1415, 1435, 1463, 1527, 1819, 1867, 1902, 1912, 2008, 2044, 2071, 2088.

3d. Quantitative analysis

See 1007.

3e. Preparative scale chromatography

1087 Huthmann, E. and Juza, M.: Modification of a commercial chiral stationary phase. Influences of enantiomer separations using simulated moving bed chromatography. *J. Chromatogr. A,* 908 (2001) 185-200.

1088 Knudsen, H.L., Fahrner, R.L., Xu, Y., Norling, L.A. and Blank, G.S.: Membrane ion-exchange chromatography for process-scale antibody purification. *J. Chromatogr. A,* 907 (2001) 145-154.

1089 Lorenz, H., Sheehan, P. and Seidel-Morgenstern, A.: Coupling of simulated moving bed chromatography and fractional crystallisation for efficient enantioseparation. *J. Chromatogr. A,* 908 (2001) 201-214.

1090 Mihlbachler, K., Fricke, J., Yun, T., Seidel-Morgenstern, A., Schmidt-Traub, H. and Guiochon, G.: Effect of the homogeneity of the column set on the performance of a simulated moving bed unit I. Theory. *J. Chromatogr. A,* 908 (2001) 49-70.

1091 Moscariello, J., Purdom, G., Coffman, J., Root, T.W. and Lightfoot, E.N.: Characterizing the performance of industrial-scale columns. *J. Chromatogr. A,* 908 (2001) 131-141.

1092 Natarajan, V. and Purdom, G.: Advances in process chromatography gradient elution using binary linear gradients. *J. Chromatogr. A,* 908 (2001) 163-167.

1093 Nicolaos, A., Muhr, L., Gotteland, P., Nicoud, R.-M. and Bailly, M.: Application of equilibrium theory to ternary moving bed configurations (four+four, five+four, eight and nine zones) I. Linear case. *J. Chromatogr. A,* 908 (2001) 71-86.

1094 Nicolaos, A., Muhr, L., Gotteland, P., Nicoud, R.-M. and Bailly, M.: Application of the equilibrium theory to ternary moving bed configurations (4+4, 5+4, 8 and 9 zones) II. Langmuir case. *J. Chromatogr. A,* 908 (2001) 87-109.

1095 Nicoud, R.-M. and Majors, R.E.: Simulated moving bed chromatography for preparative separations. *LC-Gc Eur.,* 13 (2000) 887-891.

1096 Renold, P., Madero, E. and Maetzke, T.: Automated normal-phase preparative high-performance liquid chromatography as a substitute for flash chromatography in the synthetic research laboratory. *J. Chromatogr. A,* 908 (2001) 143-148.

1097 Rittgerodt, B. and Clerc, M.: (Introduction to preparative HPLC). *GIT Labor-Fachz,* 44 (2000) 1139-1140; *C.A.,* 134 (2001) 2186z.

1098 Schulte, M. and Strube, J.: Preparative enantioseparation by simulated moving bed chromatography. *J. Chromatogr. A,* 906 (2001) 399-416 - a review with 97 refs.

See also 989, 999, 1020, 1139, 1141, 1148, 1167, 1193, 1314, 1319, 1334, 1449, 1463, 1464, 1477, 1478, 1495, 1526, 1546, 1733, 1751.

3f. Programmed temperature, pressure, vapors, gradients

1099 Mutton, I.M.: Fast generic HPLC methods. *Handb. Anal. Sep.,* 1(Separation Methods in Drug Synthesis and Purification) (2000) 73-85; *C.A.,* 134 (2001) 80305b - a review with 15 refs.

See also 1022.

4. SPECIAL TECHNIQUES

4a. Automation

1100 Gan, F., Xu, Q.-S. and Liang, Y.-Z.: Two novel procedures for automatic resolution of two-way data from coupled chromatography. *Analyst (Cambridge),* 126 (2001) 161-168.

1101 Peng, S.X., Branch, T.M. and King, S.L.: Fully automated 96-well liquid-liquid extraction for analysis of biological samples by liquid chromatography with tandem mass spectrometry. *Anal. Chem.,* 73 (2001) 708-714.

1102 Raezke, K.-P.: (Automated on-line solid-phase extraction). *LaborPraxis,* 24 (2000) 52-54; *C.A.,* 133 (2000) 246611e.

See also 1003, 1038, 1039, 1096, 1115, 1118, 1618, 1933.

4b. Computerization and modelling

1103 Gergely, A., Horvath, P. and Noszal, B.: Deconvolution of composite chromatographic peaks by simultaneous dual detections. *J. Chromatogr. Sci.,* 38 (2000) 425-429.

1104 Haber, P., Baczek, T., Kaliszan, R., Snyder, L.R., Dolan, J.W. and Wehr, C.T.: Computer simulation for the simultaneous optimization of any two variables and any chromatographic procedure. *J. Chromatogr. Sci.,* 38 (2000) 386-392.

1105 McDowall, R.D.: Have you checked your device? *LC-GC Eur.,* 14 (2001) 28-32.

1106 Meyers, J.J., Nahar, S., Ludlow, D.K. and Liapis, A.I.: Determination of the pore connectivity and pore size distribution and pore spatial distribution of porous chromatographic particles from nitrogen sorption measurements and pore network modelling theory. *J. Chromatogr. A,* 907 (2001) 57-71.

1107 Novic, M., Kunaver, M. and Novic, M.: (Computerized simulation of separation in ion chromatography: continuous study of the separation of anions in mobile and stationary phases of a chromatographic column). In: Glavic, P. and Brodnjak-Voncina, D. (Editors), *Zb. Ref. Posvetovanja Slov. Kem. Dnevi,* Univerza v Mariboru, Fakulteta za Kemijo in Kemijsko Tehnologijo, Mribor, Pt. 1, 2000, pp. 328-335; *C.A.,* 134 (2001) 24908p.

1108 Reijenga, J.C.: Training software for high-performance liquid chromatography. *J. Chromatogr. A,* 903 (2000) 41-48.

See also 1006, 1150, 1154, 1414, 1803, 2092.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

1109 Al-Dirbashi, O.Y. and Nakashima, K.: Hyphenated chromatographic methods for biomaterials. *Biomed. Chromatogr.,* 14 (2000) 406-421; *C.A.,* 134 (2001) 27025j - a review with 90 refs.

1110 Cech, N.B., Krone, J.R. and Enke, C.G.: Predicting electrospray response from chromatographic retention time. *Anal. Chem.,* 73 (2001) 208-213.

1111 Dreher, W.: (Fundamentals of HPLC-MS coupling. Ionization methods and coupling techniques). *Biol. Abwasserreinig.,* 11(Anwendung der LC-MS in der Wasseranalytik) (1999) 3-23; *C.A.,* 133 (2000) 316963x - a review with 22 refs.

1112 Karami, A., Balke, S.T. and Schunk, T.C.: Quantitative interpretation of Fourier-transform infrared spectroscopic data from a size-exclusion chromatography solvent-evaporation interface. *J. Chromatogr. A,* 911 (2000) 27-37.

1113 Kiyonami, R.: (Technology of ion-trap type HPLC-MS system). *J. Mass Spectrom. Soc. Jpn.,* 48 (2000) 202-206; *C.A.,* 133 (2000) 258515q - a review without refs.

1114 Moberg, M., Bylnd, D., Danielsson, R. and Markides, K.: Optimization strategy for liquid chromatography-electrospray ionisation mass spectrometry methods. *Analyst (Cambridge),* 125 (2000) 1970-1976.

1115 Moy, F.J., Haraki, K., Mobilio, D., Walker, G., Powers, R., Tabei, K., Tong, H. and Siegel, M.M.: MS/NMR: a structure-based approach for discovering protein ligands and for drug design by coupling size exclusion chromatography, mass spectrometry, and nuclear magnetic resonance spectroscopy. *Anal. Chem.,* 73 (2001) 571-581.

1116 Ooms, J.A.B., van Gils, G.J.M., Duninkeren, A.R. and Halmingh, O.: Development and validation of protocols for solid-phase extraction coupled to LC and LC-MS. *Am. Lab. (Shelton),* 32, No. 18 (2000) 52-57; *C.A.,* 134 (2001) 12910v.

1117 Seifar, R.M., Altelaar, M.A.F., Dijkstra, R.J., Ariese, F., Brinkman, U.A.Th. and Gooijer, C.: Surface-enhanced resonance Raman spectroscopy as an identification tool in column liquid chromatography. *Anal. Chem.,* 72 (2000) 5718-5724.

1118 Van Pelt, C.K., Corso, T.N., Schultz, G.A., Lowes, S. and Henion, J.: A four-column parallel chromatography system for isocratic or gradient LC/MS analyses. *Anal. Chem.,* 73 (2001) 582-588.

1119 Verhaert, P., Uttenweiler-Joseph, S., de Vries, M., Loboda, A., Ens, W. and Standing, K.G.: Matrix-assisted laser desorption/ionization quadrupole time-of-flight mass spectrometry: An elegant tool for peptidomics. *Proteomics (Weinheim),* 1 (2001) 118-131.

1120 Volmer, D.A.: Analysing thermally unstable compounds by LC-SSI-MS. *LC-GC Eur.,* 13 (2000) 838-847.

1121 Zhao, J.J., Xie, I.H., Yang, A.Y., Roadcap, B.A. and Rogers, J.D.: Quantitation of simvastatin and its β-hydroxy acid in human plasma by liquid-liquid cartridge extraction and liquid chromatography/tandem mass spectrometry. *J. Mass Spectrom.,* 35 (2000) 1133-1143; *C.A.,* 134 (2001) 36638p.

1122 Zimmer, D., Pickard, V., Czembor, W. and Müller, C.: Turbulent flow chromatography combined with tandem mass spectrometry for directly injected raw plasma samples derived from pharmacokinetic studies. *Chromatographia,* 52(Supplement) (2000) S26-S27.

See also 985, 1005, 1010, 1030, 1033, 1037, 1042, 1044, 1101, 1127, 1136, 1173, 1185, 1188, 1199, 1208, 1222, 1224, 1226, 1234, 1241, 1282, 1288, 1289, 1292, 1294, 1300, 1310, 1320, 1321, 1322, 1324, 1333, 1339, 1343, 1344, 1347, 1356, 1369, 1401, 1409, 1413, 1427, 1433, 1436, 1438, 1443, 1458, 1460, 1469, 1487, 1507, 1514, 1516, 1530, 1540, 1543, 1595, 1597, 1600, 1612, 1618, 1627, 1634, 1637, 1649, 1652, 1661, 1662, 1667, 1668, 1670, 1672, 1687, 1688, 1692, 1700, 1702, 1704, 1712, 1718, 1733, 1735, 1741, 1759, 1760, 1765, 1766, 1770, 1777, 1779, 1785, 1788, 1789, 1790, 1791, 1795, 1797, 1800, 1805, 1807, 1811, 1816, 1818, 1823, 1826, 1827, 1829, 1848, 1854, 1858, 1859, 1868, 1877, 1881, 1883, 1891, 1892, 1893, 1905, 1915, 1928, 1933, 1945, 1946, 1949, 1955, 1968, 1976, 1977, 1989, 1998, 1999, 2007, 2029, 2031, 2034, 2039, 2045, 2046, 2053, 2058, 2062, 2063, 2067, 2068, 2070, 2083, 2084, 2105, 2113.

4d. Affinity chromatography (advances)

1123 Jiang, H., Zou, H., Wang, H., Ni, J., Zhang, Q. and Zhang, Y.: On-line characterization of the activity and reaction kinetics of immobilized enzyme by high-performance frontal analysis. *J. Chromatogr. A,* 903 (2000) 77-84.

1124 Lopez-Mas, J.A., Streitenberger, S.A., Garcia-Carmona, F. and Sanchez-Ferrer, A.: Cyclodextrin biospecific-like displacement in dye-affinity chromatography. *J. Chromatogr. A,* 911 (2001) 47-53.

1125 Ye, M., Zou, H., Liu, Z., Lei, Z. and Ni, J.: Study on open tubular capillary affinity liquid chromatography. *J. Chromatogr. Sci.,* 38 (2000) 517-520.

See also 1258, 1415, 1448, 1477, 1501, 1506, 1509, 1522, 1575, 1579, 1618, 1804.

4f. Trace analysis and preseparation techniques

1126 Bogenschutz, G., Schafer, H. and Viehweger, K.-H.: Trace analysis in difficult matrices: new sample preparation methods for ion chromatography. *GIT Lab. J.,* 4, No. 3 (2000) 156-159; *C.A.,* 133 (2000) 286048h.

1127 Cappiello, A., Berloni, A., Famiglini, G., Mangani, F. and Palma, P.: Micro-SPE method for sample introduction in capillary HPLC/MS. *Anal. Chem.,* 73 (2001) 298-302.

1128 Carabias-Martinez, R., Rodriguez-Gonzalo, E., Moreno-Cordero, B., Perez-Pavon, J.L., Garcia-Pinto, C. and Fernandez Laespada, E.: Surfactant cloud point extraction and preconcentration of organic compounds prior to chromatography and capillary electrophoresis. *J. Chromatogr. A,* 902 (2000) 251-265 - a review with 81 refs.

1129 Dos Santos, H.C., Korn, M.G.A. and Ferreira, S.L.C.: Enrichment and determination of molybdenum in geological samples and seawater by ICP-AES using calmagite and activated carbon. *Anal. Chim. Acta,* 426 (2001) 79-84.

1130 Fritz, J.S. and Macka, M.: Solid-phase trapping of solutes for further chromatographic or electrophoretic analysis. *J. Chromatogr. A,* 902 (2000) 137-166 - a review with 352 refs.

1131 Groom, C.A., Beaudet, S., Halasz, A., Paquet, L. and Hawari, J.: Detection of the cyclic nitramine explosives hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) and octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazine (HMX) and their degradation products in soil environments. *J. Chromatogr. A,* 909 (2001) 53-60.

1132 Schäfer, C. and Lubda, D.: Alkyl diol silica: restricted access pre-column packings for fast liquid chromatography-integrated sample preparation of biological fluids. *J. Chromatogr. A,* 909 (2001) 73-78.

1133 Schutze, D., Neumayer, J. and Bartl, R.: (High throughput solid phase extraction in 96-well plates). *LaborPraxis,* 24, No. 6 (2000) 64-67; *C.A.,* 133 (2000) 249221p.

1134 Sergeyeva, T.A., Matuschewski, H., Piletsky, S.A., Bending, J., Schedler, U. and Ulbricht, M.: Molecularly imprinted polymer membranes for substance-selective solid-phase extraction from water by surface photo-grafting polymerization. *J. Chromatogr. A,* 907 (2001) 89-99.

1135 Wu, J. and Pawliszyn, J.: Preparation and applications of polypyrrole films in solid-phase microextraction. *J. Chromatogr. A,* 909 (2001) 37-52.

See also 987, 991, 994, 1000, 1071, 1102, 1116, 1121, 1184, 1189, 1199, 1200, 1760, 1762, 1775, 1825, 2005, 2025.

4g. Enantiomers, separation

1136 Bakhtiar, R. and Tse, F.L.S.: High-throughput chiral liquid chromatography/tandem mass spectrometry. *Rapid Commun. Mass Spectrom.,* 14 (2000) 1128-1135; *C.A.,* 133 (2000) 198792r.

1137 Belloli, E., Foulon, C., Yous, S., Vaccher, M.-P., Bonte, J.-P. and

1146 Kazusaki, M., Kawabata, H. and Matsukura, H.: Comparative study of amylose and cellulose derivatized chiral stationary phases in the reversed-phase mode. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2819-2828.

1147 Kazusaki, M., Kawabata, H. and Matsukura, H.: Influcence of temperature on enantioseparation employing an amylose-derivative stationary phase. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2937-2946.

1148 Khattabi, S., Cherrak, D.E., Mihlbachler, K. and Guiochon, G.: Enantioseparation of 1-phenyl-1-propanol by simulated moving bed under linear and nonlinear conditions. *J. Chromatogr. A,* 893 (2000) 307-319.

1149 Kurganov, A.: Chiral chromatographic separations based on ligand exchange. *J. Chromatogr. A,* 906 (2001) 51-71 - a review with 64 refs.

1150 Lipkowitz, K.B.: Atomistic modeling of enantioselection in chromatography. *J. Chromatogr. A,* 906 (2001) 417-442 - a review with 90 refs.

1151 Maier, N.M., Franco, P. and Lindner, W.: Separation of enantio-

tography and centrifugal partition chromatography. *J. Chromatogr. A,* 906 (2001) 365-378 - a review with 47 refs.

1142 Franco, P., Senso, A., Oliveros, L. and Minguillon, C.: Covalently bonded polysaccharide derivatives as chiral stationary phases in high-performance liquid chromatography. *J. Chromatogr. A,* 906 (2001) 155-170 - a review with 44 refs.

1143 Gasparrini, F., Misiti, D. and Villani, C.: High-performance liquid chromatography chiral stationary phases based on low-molecular-mass selectors. *J. Chromatogr. A,* 906 (2001) 35-50 - a review with 80 refs.

1144 Haginaka, J.: Protein-based chiral stationary phases for high-performance liquid chromatography enantioseparations. *J. Chromatogr. A,* 906 (2001) 253-273 - a review with 113 refs.

1145 Horvath, G., Huszthy, P., Szarvas, S., Szokan, G., Redd, J.T., Bradshaw, J.S. and Izatt, R.M.: Preparation of a new chiral pyridino-crown ether-based stationary phase for enantioseparation of racemic primary organic ammonium salts. *Ind. Eng. Chem. Res.,* 39 (2000) 3576-3581; *C.A.,* 133 (2000) 362762d.

1157 Sellergren, B.: Imprinted chiral stationary phases in high-performance liquid chromatography. *J. Chromatogr. A,* 906 (2001) 227-252 - a review with 103 refs.

1158 Tachibana, K. and Ohnishi, A.: Reversed-phase liquid chromatographic separation of enantiomers on polysaccharide type chiral stationary phases. *J. Chromatogr. A,* 906 (2001) 127-154 - a review with 107 refs.

1159 Ward, T.J. and Farris, A.B., III.: Chiral separations using the macrocyclic antibiotics: a review. *J. Chromatogr. A,* 906 (2001) 73-89 - a review with 61 refs.

1160 Yamamoto, C. and Okamoto, Y.: (Design of materials for chiral recognition). *Maku,* 25, No. 6 (2000) 277-279; *C.A.,* 134 (2001) 24974g - a review with 7 refs.

1161 Yashima, E.: Polysaccharide-based chiral stationary phases for high-performance liquid chromatographic enantioseparation. *J. Chromatogr. A,* 906 (2001) 105-125 - a review with 133 refs.

See also 1021, 1031, 1045, 1046, 1059, 1060, 1063, 1082, 1087, 1089, 1098, 1229, 1239, 1302, 1365, 1384, 1386, 1390, 1392, 1517, 1527, 1628, 1633, 1655, 1678, 1752, 1795, 1815, 1821, 1824, 1830, 1838, 1839, 1842, 1843, 1855, 1862, 1867, 1871, 1879, 1883, 1912, 1917, 1944, 1947, 1964.

4h. Other special techniques

1162 Battu, S., Roux, A., Delebasee, S., Bosgiraud, C. and Cardot, P.J.P.: Sedimentation field-flow fractionation device cleaning, decontamination and sterilization procedures for cellular analysis. *J. Chromatogr. B,* 751 (2001) 131-141.

1163 Du, Q.-Z., Wu, P.-D. and Ito, Y.: Van Deemter plot in high-speed countercurrent chromatography with a fixed volume of stationary phase. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2929-2935.

1164 Hoyos, M., Moore, L.R., McCloskey, K.E., Margel, S., Zuberi, M., Chalmers, J.J. and Zborowski, M.: Study of magnetic particles pulse-injected into an annular SPLITT-like channel inside a quadrupole magnetic field. *J. Chromatogr. A,* 903 (2000) 99-116.

1165 Karg, J.: (Micro-HPLC - a newly revived old trend). *GIT Spez. Sep.,* 20, No.2 (2000) 70-71; *C.A.,* 133 (2000) 316850h - a review with 3 refs.

1166 Li, X., Mohan, S., Gu, W., Miyakoshi, N. and Baylink, D.J.: Differential protein profile in the ear-punched tissue of regeneration and non-regeneration strains of mice: a novel approach to explore the candidate genes for soft-tissue regeneration. *Biochim. Biophys. Acta,* 1524 (2000) 102-109.

1167 Quinones-Garcia, I., Rayner, I., Levison, P.R., Dickson, N. and Purdom, G.: Performance comparison of suspended bed and batch contactor chromatography. *J. Chromatogr. A,* 908 (2001) 169-178.

See also 989, 1088, 1095, 1123, 1141, 1190, 1458, 1733, 1810.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

1168 Pasadakis, N. and Varotsis, N.: A novel approach for the quantitation of the hydrocarbon groups in heavy petroleum fractions by HPLC-RI analysis. *Energy Fuels,* 14 (2000) 1184-1187; *C.A.,* 133 (2000) 323775r.

5b. Cyclic hydrocarbons, fullerenes

1169 Dmitrikov, V.P. and Nabivach, V.M.: (Chromatographic analysis of polycyclic arenes with various degrees of hydrogenation). *Koks Khim.,* No. 6 (2000) 28-32; *C.A.,* 134 (2001) 50807z.

1170 Ferroukhi, O., Atik, N., Guermouche, S., Guermouche, M.H., Berdague, P., Judenstein, P. and Bayle, J.P.: High-performance liquid chromatography of aromatic and polyaromatic hydrocarbons on a new chemically bonded liquid crystal phase. *Chromatographia,* 52 (2000) 564-568.

1171 Kayali-Sayadi, M.N., Rubio-Barroso, S., Diaz-Diaz, C.A. and Polo-Diez, L.M.: Rapid determination of PAHs in soil samples by HPLC with fluorometric detection following sonication extraction. *Fresenius J. Anal. Chem.,* 368 (2000) 697-701.

1172 Law, R.J., Kelly, C.A. and Nicholson, M.D.: The QUASIMEME laboratory performance study of polycyclic aromatic hydrocarbons (PAHs): Assessment for the period 1996-1999. *J. Environ. Monit.,* 2 (2000) 517-523; *C.A.,* 133 (2000) 339088e.

1173 Li, Y., Zhang, X., Xu, R., Cai, W., Wang, F. and Song, Y.: (Quantitative analysis of polynuclear aromatic hydrocarbons in water by solid phase microextraction/high performance liquid chromatography). *Huanjing Huaxue,* 19 (2000) 382-384; *C.A.,* 133 (2000) 182542y.

1174 Miyake, Y., Minami, T., Kikuchhi, K., Kainosho, M. and Achiba, Y.: Trends in structure and growth of higher fullerenes isomer structure of C_{86} and C_{88}. *Mol. Cryst. Liq. Cryst. Sci. Technol., Sect. A,* 340 (2000) 553-558; *C.A.,* 134 (2001) 65391h.

1175 Popp, P., Bauer, C., Möder, M. and Paschke, A.: Determination of polycyclic aromatic hydrocarbons in waste water by off-line coupling of solid-phase microextraction with column liquid chromatography. *J. Chromatogr. A,* 897 (2000) 153-159.

1176 Sueki, K.: (Chemistry of endohedral actinoids metallofullerene). *JAERI-Conf.,* 2000-2002(Proceedings of the Workshop on the Nuclear Sciences of the Heaviest Elements, 1999) (2000) 86-89; *C.A.,* 134 (2001) 12628j.

1177 Tagmatarchis, N. and Shinohara, H.: Production, separation, isolation, and spectroscopic study of dysprosium endohedral metallofullerenes. *Chem. Mater.,* 12 (2000) 3222-3226; *C.A.,* 134 (2001) 12668x.

1178 Tagmatarchis, N., Aslanis, E., Shinohara, H. and Prassides, K.: Isolation and spectroscopic study of a series of mon- and dierbium endohedral C_{82} and C_{84} metallofullerenes. *J. Phys. Chem. B,* 104 (2000) 11010-11012; *C.A.,* 134 (2001) 80060t.

1179 Yan, X.-h., Zhang, B.-k., Li, H.-d. and Ma, H.-x.: (HPLC determination of polycyclic aromatic hydrocarbons in power plant emissions). *Sepu,* 18 (2000) 432-435; *C.A.,* 133 (2000) 368832y.

1180 Yang, Z.-F., Li, M. and Zhang, S.-S.: (Analysis of polycyclic aromatic hydrocarbons (PAHs) in water by high pressure liquid chromatography). *Qingdao Huagong Xueyuan Xuebao,* 21 (2000) 274-276; *C.A.,* 134 (2001) 8929r.

See also 1061, 1069, 1084, 1135, 1168, 1182, 1351.

5c. Halogen derivatives

1181 Pietrogrande, M.C., Benvenuti, A., Previato, S. and Dondi, F.: HPLC analysis of PCBs on porous graphitic carbon: retention behavior and gradient elution. *Chromatographia,* 52 (2000) 425-432.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

1182 Fu, X.-t., Wang, Z.-p. and Bi, L.-f.: (Prediction of output fraction from single layer to commingling oil well by high performance liquid chromatography (HPLC)). *Sepu,* 18 (2000) 318-321; *C.A.,* 133 (2000) 179974k.

1183 Sarowha, S.L.S., Sharma, B.K., Sharma, C.D., Sharma, R.L. and Madhwal, D.C.: Hydrocarbon class type distriburion of crude oils by high performance liquid chromatography. *Indian J. Chem. Technol.*, 7, No. 4 (2000) 172-177; *C.A.*, 134 (2001) 6700k.

1184 Wang, X., Tian, S.-b., Lu, W.-z. and Qiu, Y.-m.: (Preliminary study on detection of composition of lubricating oil by supercritical fluid extraction). *Shiyou Xuebao, Shiyou Jiagong*, 16, No. 5 (2000) 70-75; *C.A.*, 134 (2001) 6748g.

6. ALCOHOLS

1185 Terado, I., Inoue, H., Tao, Y. and Tsuruta, Y.: 3-[(1-{[4-(5,6-Dimethoxy-1-oxoisoindolin-2-yl)-2-methoxyphenyl]sulfonyl}-pyrrolidin-2-yl)carbonylamino]phenylboronic acid as a fluorescent labeling reagent for determination of diol compounds by HPLC. *Anal. Sci.*, 16 (2000) 881-883; *C.A.*, 133 (2000) 316981b.

1186 Toussaint, B., Duchateau, A.L.L., van der Wal, S., Albert, A., Hubert, P. and Crommen, J.: Comparative evaluation of four detectors in the high-performance liquid chromatographic analysis of chiral nonaromatic alcohols. *J. Chromatogr. Sci.*, 38 (2000) 450-457.

1187 You, J., Zhang, B., Zhang, W., Yang, P. and Zhang, Y.: Determination of alcohols by high-performance liquid chromatography with fluorimetric detection after pre-column derivatisation with carbazole-9-*N*-acetic acid and chromatographic behaviour of alcoholic derivatives. *J. Chromatogr. A*, 909 (2001) 171-182.

See also 1148, 2026.

7. PHENOLS

1188 Andres-Lacueva, C., Lamuela-Raventós, R.M., Jáuregui, O., Casals, I., Izquierdo-Pulido, M. and Permanyer, J.: An LC method for the analysis of cocoa phenolics. *LC-GC Eur.*, 13 (2000) 902-904.

1189 Baciocchi, R., Attina, M., Lombardi, G. and Boni, M.R.: Fast determination of phenols in contaminated soils. *J. Chromatogr. A*, 911 (2001) 135-141.

1190 Chen, L., Han, Y., Yang, F. and Zhang, T.: High-speed countercurrent chromatography separation and purification of resveratrol and piceid from *Polygonum cuspidatum*. *J. Chromatogr. A*, 907 (2001) 343-346.

1191 Chen, S., Yang, Y., Zhang, L., Pang, X., Dai, Z. and Zeng, F.: A sensitive HPLC for the quantitation of dauricine. *J. Tongji Med. Univ.*, 20 (2000) 39-42; *C.A.*, 133 (2000) 256918.

1192 Coman, V. and Modovan, Z.: RP-HPLC method for the separation of some phenol derivatives using gradient elution and UV detection. *J. High Resolut. Chromatogr.*, 23 (2000) 699-701.

1193 Degenhardt, A., Engelhardt, U.H., Lakenbrink, C. and Winterhalter, P.: Preparative separation of polyphenols from tea by high-speed countercurrent chromatography. *J. Agric. Food Chem.*, 48 (2000) 3425-3430.

1194 Escarpa, A. and Gonzalez, M.C.: Optimization strategy and validation of one chromatographic method as approach to determine the phenolic compounds from different sources. *J. Chromatogr. A*, 897 (2000) 161-170.

1195 Fang, C.-m. and Sun, Y.-p.: (Simultaneous determination of phenol and its photo-catalyzed degradation intermediates by HPLC). *Fenxi Ceshi Xuebao*, 19 (2000) 48-50; *C.A.*, 133 (2000) 198224g.

1196 Inoue, K., Kato, K., Yoshimura, Y., Makino, T. and Nakazawa, H.: Determination of bisphenol A in human serum by high-performance liquid chromatography with multi-electrode electrochemical detection. *J. Chromatogr. B*, 749 (2000) 17-23.

1197 Inoue, K., Yoshimura, Y., Makino, T. and Nakazawa, H.: Determination of 4-nonylphenol and 4-octylphenol in human blood samples by high-performance liquid chromatography with multi-electrode electrochemical coulometric-array. *Analyst (Cambridge)*, 125 (2000) 1959-1961.

1198 Li, H., Krieger, R.I. and Li, Q.X.: Improved HPLC method for analysis of 1-hydroxypyrene in human urine specimens of cigarette smokers. *Sci. Total Environ.*, 257, No. 2-3 (2000) 147-153; *C.A.*, 133 (2000) 241739t.

1199 Rosenberg, E., Wissiack, R. and Grasserbauer, M.: (Possibilities and problems in determining phenols by on-line SPE-HPLC-MS). *Biol. Abwasserreinig.*, 11(Anwendung der LC-MS in der Wasseranalytik) (1999) 73-91; *C.A.*, 133 (2000) 242053b.

1200 Seronero, L.C., Fernandez Laespada, M.E., Perez Pavon, J.L. and Cordero, B.M.: Cloud point preconcentration of rather polar compounds: application to the high-performance liquid chromatographic determination of priority pollutant chlorophenols. *J. Chromatogr. A*, 897 (2000) 171-176.

1201 Sun, Y., Wada, M., Al-Dirbashi, O., Kuroda, N., Nakazawa, H. and Nakashima, K.: High-performance liquid chromatography with peroxyoxalate chemiluminescence detection of bisphenol A migrated from polycarbonate baby bottles using 4-(4,5-diphenyl-1*H*-imidazol-2-yl)benzoyl chloride as a label. *J. Chromatogr. B*, 749 (2000) 49-56.

1202 Will, F., Bauckhage, K. and Dietrich, H.: Apple pomace liquefaction with pectinases and celulases: analytical data of the corresponding juices. *Eur. Food Res. Technol.*, 211, No. 4 (2000) 291-297; *C.A.*, 134 (2001) 55759v.

See also 1086, 1163, 1203, 1230, 1234, 1291, 1365, 1794.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

1203 Areias, F., Valentao, P., Andrade, P.B., Ferreres, F. and Seabra, R.M.: Flavonoids and phenolic acids of sage. Influence of some agricultural factors. *J. Agric. Food Chem.*, 48 (2000) 6081-6084.

1204 Baugh, S. and Ignelzi, S.: Hydrolysis and redox factors affecting analysis of common phenolic marker compounds in botanical extracts and finished products. *J. Assoc. Off. Anal. Chem.*, 83 (2000) 1135-1140.

1205 De Pascual-Teresa, S., Santos-Buelga, C. and Rivas-Gonzalo, J.C.: Quantitative analysis of flavan-3-ols in Spanish foodstuffs and beverages. *J. Agric. Food Chem.*, 48 (2000) 5331-5337.

1206 Deggenhardt, A., Engelhardt, U.H., Wendt, A.-S. and Winterhalter, P.: Isolation of black tea pigments using high speed countercurrent chromatography and studies of properties of black tea polymers. *J. Agric. Food Chem.*, 48 (2000) 5200-5205.

1207 Huang, W. and Liang, X.: (Determination of two flavone glycosides in *Trollius ledebourii* by HPLC). *Zhongguo Yaoxue Zazhi (Beijing),* 35 (2000) 658-659; *C.A.,* 134 (2001) 33066q.

1208 Li, S., Li, G., Li, Z., Han, F. and Chen, Y.: (Rapid separation and identification of isoflavonoids from Chinese herb Radix Puerariae and determination of puerarin from Chinese traditional drug Xiao'er Fu Xie Ning effervescent pill by RP-HPLC-MS-MS).

1221 Munimbazi, C. and Bullerman, L.B.: Chromatographic method for the determination of the mycotoxin moniliformin in corn. *Methods Mol. Biol. (Totowa),* 157(Mycotoxin Protocols) (2001) 131-145; *C.A.,* 134 (2001) 52346x.

1222 Richard, J.L., Plattner, R.D., May, J. and Liska, S.L.: The occurrence of ochratoxin A in dust collected from a problem household. *Mycopathologia,* 146, No. 2 (2000) 99-103; *C.A.,* 133

1215 Dorner, J.W., Sobolev, V.S., Yu, W. and Chu, F.S.: Immunochemical method for cyclopiazonic acid. *Methods Mol. Biol. (Totowa),* 157(Mycotoxin Protocols) (2001) 71-80; *C.A.,* 134 (2001) 38013e.

1216 Entwisle, A.C., Williams, A.C., Mann, P.J., Slack, P.T. and Gilbert, J.: Liquid chromatographic method with immunoaffinity column cleanup for determination of ochratoxin A in barley: collaborative study. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1377-1383.

1217 Eppley, R.M.: Preparatory isolation of mycotoxins from solid phase fungal cultures. *Methods Mol. Biol. (Totowa),* 157(Mycotoxin Protocols) (2001) 25-29; *C.A.,* 134 (2001) 38010b.

1218 Festas, I., Hervert, P., Santos, L., Cabral, M., Barros, P. and Alves, A.: Ochratoxin A in some Portuguese wines. Method validation and screening in Port wine and Vinho Verde. *Am. J. Enol. Vitic.,* 51, No. 2 (2000) 150-154; *C.A.,* 133 (2000) 334200a.

1219 Lai, C.-L., Fuh, Y.-M. and Shih, D.Y.-C.: Detection of mycotoxin patulin in apple juice. *Yaowu Shipin Fenxi,* 8, No.2 (2000) 85-96; *C.A.,* 133 (2000) 295545v.

1220 MacDonald, S., Long, M., Gilbert, J. and Felgueiras, I.: Liquid chromatographic method for determination of patulin in clear and cloudy apple juices and apple puree: collaborative study. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1387-1394.

See also 1347, 1637.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

1229 Butkus, E., Berg, U., Zilinskas, A., Kubilius, R. and Stoncius, S.: Enantiomer separation and absolute configuration of densely functionalized 2-oxatricyclo[4.3.1.$0^{3,8}$]decanes by CD spectroscopy and chemical correlation. *Tetrahedron: Asymmetry,* 11 (2000) 3053-3057; *C.A.,* 134 (2001) 29273n.

1230 Fukumoto, L.R. and Mazza, G.: Assessing antioxidant and prooxidant activities of phenolic compounds. *J. Agric. Food Chem.,* 48 (2000) 3597-3604.

1231 Gioacchini, A.M., Pezzetta, D., Calonghi, N., Boga, C., Cappadone, C., Masotti, L. and Traldi, P.: Tandem mass spectrometry in the determination of 4-hydroxy-2-nonenal at the cellular level. *Rapid Commun. Mass Spectrom.,* 14 (2000) 1954-1956; *C.A.,* 134 (2001) 53436p.

1232 Guo, Q. and Lu, J.: (HPLC determination of anthraquinones in Radix Polygoni multiflori and Radix Polygoni multiflori preparations). *Yaowu Fenxi Zazhi,* 20 (2000) 326-328; *C.A.,* 134 (2001) 33064n.

1233 Huang, X. and Zhang, D.: (Determination of formaldehyde in rainwater by HPLC). *Huanjing Huaxue,* 19 (2000) 566-571; *C.A.,* 134 (2001) 75994w.

1234 Kempen, E.C., Brodbelt, J.S., Bartsch, R.A., Blanda, M.T. and Farmer, D.B.: Screening metal binding selectivities of macrocycle mixtures by HPLC-ESI-MS and postcolumn reactions. *Anal. Chem.,* 73 (2001) 384-390.

1235 Kobayashi, K., Yamamoto, T., Taguchi, M. and Chiba, K.: High-performance liquid chromatography determination of N- and O-demethylase activities of chemicals in human liver microsomes: applications of postcolumn fluorescence derivatization using Nash reagent. *Anal. Biochem.,* 284 (2000) 342-347.

1236 Li, L., Ding, G., Wu, J., Qian, G. and Liu, S.: (Determination of osthole in Fengshi oral liquid by HPLC). *Huaxi Yaoxue Zazhi,* 15 (2000) 203-204; *C.A.,* 133 (2000) 198797w.

1237 Luo, J., Zhang, Z., Tian, C. and Li, X.: (Determination of cinnamaldehyde in *Cinnamonum cassia* Predl by HPLC). *Zhongguo Zhongyao Zazhi,* 25 (2000) 544-545; *C.A.,* 133 (2000) 340341p.

1238 Schramm, C. and Rinderer, B.: Determination of cotton-bound glyoxal *via* an internal Cannizzaro reaction by means of high-performance liquid chromatography. *Anal. Chem.,* 72 (2000) 5829-5833.

1239 Steinreiber, A., Strauss, U., Mayer, S.F. and Uray, G.: Simultaneous direct high-performance liquid chromatographic enantioseparation of 2-methylglycidol-1-benzyl ether and 2-methylglycerol-1-benzyl ether using a solvent-switching technique. *J. Chromator. A,* 911 (2001) 295-298.

1240 Suttnar, J., Masova, L. and Dyr, J.E.: Influence of citrate and EDTA anticoagulants on plasma malondialdehyde concentrations estimated by high-performance liquid chromatography. *J. Chromatogr. B,* 751 (2001) 193-197.

1241 Wang, X.-c. and Jin, Y.: (Analysis of carbon number and ethylene oxide number distribution in alcohol ethoxylates by HPLC electrospray ionization mass spectrometry). *Sepu,* 18 (2000) 332-334; *C.A.,* 133 (2000) 268569z.

See also 1051, 1137, 1195, 1293, 1656, 1658, 1998, 2027.

10. CARBOHYDRATES

See 2098.

10a. Mono and oligosaccharides. Structural studies

1242 An, Y., Young, S.P., Hillman, S.L., van Hove, J.L.K., Chen, Y.-T. and Millington, D.S.: Liquid chromatographic assay for a glucose tetrasaccharide, a putative biomarker for the diagnosis of Pompe disease. *Anal. Biochem.,* 287 (2000) 136-143.

1243 Andersen, R. and Sørensen, A.: Separation and determination of alditols and sugars by high-pH anion-exchange chromatography with pulsed amperometric detection. *J. Chromatogr. A,* 897 (2000) 195-204.

1244 Boon, M.A., Van't Riet, K. and Janssen, A.E.M.: Enzymatic synthesis of oligosaccharides: product removal during a kinetically controlled reation. *Biotechnol. Bioeng.,* 70 (2000) 411-420; *C.A.,* 134 (2001) 56883t.

1245 Cataladi, T.R.I., Campa, C. and Benedetto, G.: Carbohydrate analysis by high performance anion-exchange chromatography with pulsed amperometric detection: The potential is still growing. *Fresenius J. Anal. Chem.,* 368 (2000) 739-758.

1246 Cheng, X. and Kaplan, L.A.: Improved analysis of dissolved carbohydrates in stream water with HPLC-PAD. *Anal. Chem.,* 73 (2001) 458-461.

1247 Duedahl-Olesen, L., Larsen, K.L. and Zimmermann, W.: Rapid detection of malto-oligosaccharide-forming bacterial amylases by high performance anion-exchange chromatography. *Lett. Appl. Microbiol.,* 30 (2000) 312-316; *C.A.,* 133 (2000) 277874x.

1248 Fontaine, T., Simenel, C., Dubreuq, G., Adam, O., Delpierre, M., Lemoine, J., Vorgais, C.E., Diaquin, M. and Latge, J.P.: Molecular organization of the alkali-insoluble fraction of *Aspergillus fumigatus* cell wall. *J. Biol. Chem.,* 275 (2000) 27594-27607.

1249 Fukuta, K., Abe, R., Yokomatsu, T., Omal, F., Asanagi, M. and Makino, T.: Control of bisecting GlcNAc addition to N-linked sugar chains. *J. Biol. Chem.,* 275 (2000) 23456-23461.

1250 Grard, S., Elfakir, C. and Dreux, M.: Characterization of sulfobutyl ether-β-cyclodextrins mixtures by anion-exchange chromatography using evaporative light scattering detection. *J. Chromatogr. A,* 897 (2000) 185-193.

1251 Guenu, S., Seigneuret, J.M. and Dreux, M.: Analysis of maltodextrins in plant extracts by LC using evaporative light scattering detection. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2883-2895.

1252 Hogarth, A.J.C.L., Hunter, D.E., Jacobs, W.A., Garleb, K.A. and Wolf, B.W.: Ion chromatographic determination of three fructooligosaccharide oligomers in prepared and preserved foods. *J. Agric. Food Chem.,* 48 (2000) 5326-5330.

1253 Hudina, M. and Stampar, F.: Sugars and organic acids contents of European (*Pyrus communis*) and Asian (*Pyrus serotina*) pear cultivars. *Acta Aliment.,* 29 (2000) 217-230; *C.A.,* 134 (2001) 70597j.

1254 Lopez-Hernandez, J., Gonzalez-Castro, M.J. and Pereda-Gonzalez, M.: High-performance liquid chromatographic determination of glycogen in sea urchin gonads with refractive index detection. *J. Chromatogr. Sci.,* 38 (2000) 383-385.

1255 Matsui, T., Hamako, J., Ozeki, Y. and Titani, K.: Comparative study of blood group-recognizing lectins toward ABO blood group antigens on neoglycoproteins, glycoproteins and complex-type oligosaccharides. *Biochim. Biophys. Acta,* 1525 (2001) 50-57.

1256 Michalski, J.-C. and Capon, C.: Monosaccharide composition of mucins. *Methods Mol. Biol. (Totowa),* 125(Glycoprotein Methods and Protocols) (2000) 159-180; *C.A.,* 133 (2000) 278095f - a review with 48 refs.

1257 Nakamura, T., Amikawa, S., Harada, T., Saito, T., Arai, I. and Urashima, T.: Occurrence of an unusual phosphorylated *N*-acetyllactosamine in horse colostrum. *Biochim. Biophys. Acta,* 1525 (2001) 13-18.

1258 Okada, T., Sato, Y., Kobayashi, N., Sumida, K., Satomura, S., Matsuura, S., Takasaki, M. and Endo, T.: Structural characteristics of the *N*-glycans of two isoforms of prostate-specific antigens purified from human seminal fluid. *Biochim. Biophys. Acta,* 1525 (2001) 149-160.

1259 Pellegrino, L., de Noni, I. and Cattaneo, S.: Formation of protein bound lysine-derived galactosyl and glucosyl pyrroles in heated model systems. *Nahrung,* 44 (2000) 193-200; *C.A.,* 134 (2001) 41309m.

1260 Sakaguchi, H., Watanabe, M., Ueoka, C., Sugiyama, E., Taketomi, T., Yamada, S. and Sugahara, K.: Isolation of reducing oligosaccharide chains from the chondroitin/dermatan sulfate-protein linkage region and preparation of analytical probes by fluorescent labeling with 2-aminobenzamide. *J. Biochem. (Tokyo),* 129 (2001) 107-118.

1261 Toyoda, H., Kinoshita-Toyoda, A., Fox, B. and Selleck, S.B.: Structural analysis of glycosaminoglycans in animal bearing mutations in sugarless, sulfateless, and tout-velu. *J. Biol. Chem.,* 275 (2000) 21856-21861.

1262 Way, W.K., Gibson, K.G. and Breite, A.G.: Determination of glucosamine in nutritional supplements by reversed-phase ion-pairing HPLC. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2861-2871.

1263 Zhou, L. and Ji, H.: (Determination of lactulose in oral solution by HPLC). *Yaowu Fenxi Zazhi,* 20 (2000) 174-176; *C.A.,* 133 (2000) 313735p.

See also 1042, 1120, 1207, 1266, 1296, 1669, 1784, 2011.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

1264 Bayliss, M.A., Gage, R., Edgington, A.M. and Venn, R.F.: Fluorescence determination of sulphobutylether-β-cyclodextrin sodium in human plasma and urine by size-exclusion chromatography with inclusion formation. *Chromatographia,* 52(Supplement) (2000) S83-S86.

1265 Eklund, E., Roden, L., Malmström, M. and Malmström, A.: Dermatan is a better substrate for 4-O-sulfation than chondroitin: implications in the generation of 4-O-sulfated, L-iduronate-rich galactosaminoglycans. *Arch. Biochem. Biophys.,* 383 (2000) 171-177.

1266 Guerardel, Y., Kol, O., Maes, E., Lefebvre, T., Boilly, B., Davril, M. and Strecker, G.: O-glycan variability of egg-jelly mucins from *Xenopus laevis*: characterization of four phenotypes that differ by the terminal glycosylation of their mucins. *Biochem. J.,* 352 (2000) 449-463.

1267 Kariya, Y., Kyogashima, M., Suzuki, K., Isomura, T., Sakamoto, T., Horie, K., Ishihara, M., Takamo, R., Kamei, K. and Hara, S.: Preparation of completely 6-O-desulfated heparin and its ability to enhance activity of basic fibroblast growth factor. *J. Biol. Chem.,* 275 (2000) 25949-25958.

1268 Mukhiddinov, Z.K., Khalikov, D.K., Abdusamiev, F.T. and Avloev, C.C.: Isolation and structural characterization of a pectin homo- and ramnogalacturonan. *Talanta,* 53 (2000) 171-176; *C.A.,* 133 (2000) 336765n.

1269 Pastore, A., Bernardini, S., Dello Strologo, L., Rizzoni, G., Cortese, C. and Federici, G.: Simultaneous determination of inulin and *p*-aminohippuric acid in plasma and urine by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B,* 751 (2001) 187-191.

1270 Shang, P., Mei, Q., Cao, Z. and Zhao, D.: (Analysis of Angelica polysaccharides constituents by HPLC). *Zhongguo Yaoxue Zazhi (Beijing),* 35 (2000) 332-335; *C.A.,* 133 (2000) 183122y.

1271 Warnken, U.H., Asskali, F. and Forster, H.: Molar substitution of clinically used acetylated starch by size exclusion chromatography using multiple detection. *Starch/Staerke,* 52 (2000) 261-266; *C.A.,* 134 (2001) 65743z.

1272 Way, W.K., Gibson, K.G. and Breite, A.G.: Determination of chondroitin sulfate in nutritional supplements by liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2851-2860.

See also 1160.

10c. Glycoproteins and their constituents

1273 Brucato, S., Fagnen, G., Villers, C., Bonnamy, P.-J., Langris, M. and Bocquet, J.: Biochemical characterization of integral membrane heparan sulfate proteoglycans in Sertoli cells from immature rat testis. *Biochim. Biophys. Acta,* 1510 (2001) 474-487.

1274 Cavada, B.S., Madeira, S.V.F., Calvete, J.J., Souza, L.A.G., Bomfim, L.R., Dantas, A.R., Loex, M.C., Grangeiro, T.B., Freitas, B.T., Pinto, V.P.T., Leite, K.B. and Ramos, M.V.: Purification, chemical, and immunochemical properties of a new lectin from Mimosoidae (*Parkia discolor*). *Prepar. Biochem. Biotechnol.,* 30 (2000) 271-280.

1275 Cochet, S., Volet, G., Cartron, J.-P. and Bertrand, O.: New procedures for glycophorin A purification with high yield and high purity. *J. Chromatogr. B,* 750 (2001) 109-119.

1276 Huet, G., Delannoy, P., Lesuffleur, T., Hennebicq, S. and Degand, P.: Inhibition of mucin glycosylation. *Methods Mol. Biol. (Totowa),* 125(Glycoprotein Methods and Protocols) (2000) 261-271; *C.A.,* 133 (2000) 292973x - a review with 15 refs.

1277 Inoue, S., Lin, S.-L. and Inoue, Y.: Chemical analysis of the developmental pattern of polysialylation in chicken brain. *J. Biol. Chem.,* 275 (2000) 29968-29979.

1278 Itoh, Y., Inuzuka, K., Kohno, I., Wada, H., Shiku, H., Ohkura, N. and Kato, H.: Highly increased plasma concentrations of the nicked form of β_2 glycoprotein I in patients with leukemia and with lupus anticoagulant: measurement with a monoclonal antibody specific for a nicked form of domain V. *J. Biochem. (Tokyo),* 128 (2000) 1017-1024.

1279 Li, Y.C., Larsson, E.L., Jungvid, H., Galaev, I.Y. and Mattiasson, B.: Shielding of protein-boronate interactions during boronate chromatography of neoglycoproteins. *J. Chromatogr. A,* 909 (2001) 137-145.

1280 Mori, S., Nishibori, M., Yamaoka, K. and Okamoto, M.: One-step purification of rabbit histidine rich glycoprotein by dye-ligand affinity chromatography with metal requirement. *Arch. Biochem. Biophys.,* 383 (2000) 191-196.

1281 Nakano, T. and Ozimek, L.: Purification of glycomacropeptide from dialyzed and non-dialyzed sweet whey by anion-exchange chromatography at different pH values. *Biotechnol. Lett.,* 22 (2000) 1081-1086; *C.A.,* 133 (2000) 265928e.

1282 Ohta, M., Kawasaki, N., Hyuga, S., Hyuga, M. and Hayakawa, T.: Selective glycopeptide mapping of erythropoietin by on-line high-performance liquid chromatography-electrospray ionization mass spectrometry. *J. Chromatogr. A,* 910 (2001) 1-11.

1283 Ooi, L.S.M., Ng, T.B., Geng, Y. and Ooi, V.E.C.: Lectins from bulks of the Chinese daffodil *Narcissus tazetta* (family Amaryllidaceae). *Biochem. Cell Biol.,* 78 (2000) 463-468.

1284 Patra, A.K., Gahlay, G.K., Reddy, B.V.V., Gupta, S.K. and Panda, A.K.: Refolding, structural transition and spermatozoa-binding of recombinant bonnet monkey (*Macaca radiata*) zona pellucida glycoprotein-C expressed in *Escherichia coli*. *Eur. J. Biochem.*, 267 (2000) 7075-7081.

1285 Svitacheva, N. and Davies, J.R.: Mucin biosynthesis and secre-

1295 Harukaze, A., Sugiyama, S., Iwamoto, Y., Murata, M. and Homma, S.: Convenient analysis and quantification of diferulic acids in foods. *Food Sci. Technol. Res.*, 6, No. 2 (2000) 122-125; *C.A.*, 133 (2000) 321118e.

1296 Hirayama, F., Ogata, T., Yano, H., Arima, H., Udo, K., Takano, M. and Uekama, K.: Release characteristics of a short-chain

the simultaneous determination of methylxanthines, chlorogenic acid, some hydrobyvenoic and hydroxy-cinnamic acids. Application to cocoa, coffee, tea and cola-drinks. *Chromatographia*, 52 (2000) 579-583.

1287 Baram, G.I., Azarova, I.N., Gorshkov, A.G., Vereshchagin, A.L., Lang, B. and Kiryukhina, E.D.: Determination of bis(2-ethylhexyl) phthalate in water by high-performance liquid chromatography with direct on-column preconcentration. *J. Anal. Chem.*, 55 (2000) 750-754; *C.A.*, 133 (2000) 198220c.

1288 Bylund, J., Bylund, M. and Oliw, E.H.: cDNA cloning and expression of CYP4F12. A novel human cytochrome P450. *Biochem. Biophys. Res. Commun.*, 280 (2001) 892-897.

1289 Claassen, F.W., van de Haar, C., van Beek, T.A., Dorado, J., Martinez-Inigo, M.-J. and Sierra-Alvarez, R.: Rapid analysis of apolar low-molecular-weight contituents in wood using high-pressure liquid chromatography with evaporative light scattering detection. *Phytochem. Amal.*, 11 (2000) 251-256; *C.A.*, 133 (2000) 239549n.

1290 Czarnecka, M., Czarnecki, Z. and Roszyk, H.: (Evaluation of selected methods for lactic acid determination). *Zywnosc*, 7 (2000) 92-101; *C.A.*, 133 (2000) 236978r.

1291 Deng, K., Liang, Y. and Qin, J.: (Determination of gallic acid in Jiechangning suppositories by HPLC). *Yaowu Fenxi Zazhi*, 20, No.6 (2000) 406-408; *C.A.*, 134 (2001) 61668f.

1292 Endo, Y., Tagiri-Endo, M., Seo, H.-S. and Fujimoto, K.: Identification and quantification of molecular species of diacyl glyceryl ether by reversed-phase high-performance liquid chromatography with refractive index detection and mass spectrometry. *J. Chromatogr. A*, 911 (2001) 39-45.

1293 Feng, S., Chen, L., Zhao, F., Wang, X. and Zhao, W.: (Determination of rhein, emodin and tanshinone II4A in Dhuangjiangzhi powder by HPLC). *Yaowu Fenxi Zazhi*, 20 (2000) 368-370; *C.A.*, 134 (2001) 46885f.

1294 Frauendoft, H. and Herzschuh, R.: (Use of HPLC electrospray MS to identify carboxylic acids from brown coal in an aqueous matrix). *Biol. Abwasserreinig.*, 11(Anwendung der LC-MS in der Wasseranalytik) (1999) 107-123; *C.A.*, 133 (2000) 242226k.

of EETs using microbore liquid chromatography with fluorescence detection. *Am. J. Physiol.*, 279 (2000) H857-H862; *C.A.*, 133 (2000) 249143q.

1301 Pan, R., Chen, D., Shen, L., Si, J., Hu, Z. and Zhu, C.: (Determination of ferulic acid and isoferulic acid in rhizoma cimicifugae by HPLC). *Yaowu Fenxi Zazhi*, 20 (2000) 396-398; *C.A.*, 134 (2001) 46887h.

1302 Schneider, C., Boeglin, W.E. and Brash, A.R.: Enantiomeric separation of hydroxy eicosanoids by chiral column chromatography: effect of the alcohol modifier. *Anal. Biochem.*, 287 (2000) 186-189.

1303 Sun, T., Shi, C., Siong, Y., Qi, H. and Mu, X.: (Determination of trace chloroacetic acids in water by high performance ion chromatography). *Fenxi Huaxue*, 28 (2000) 1443; *C.A.*, 134 (2001) 75996y.

1304 Vodopivec, M., Podgornik, A., Berovic, M. and Strancar, A.: Application of convective interaction media (CIM) disk monolithic columns for fast separation and monitoring of organic acids. *J. Chromatogr. Sci.*, 38 (2000) 489-495.

1305 Wu, Y., Zhang, X. and Jiang, X.: (Rapid determination of acetic acid, chloroacetic acid, dichloroacetic acid, and chloride ion in chemicals by ion chromatography). *Huaxue Yanhiu Yu Yingyong*, 12 (2000) 458-460; *C.A.*, 134 (2001) 12918d.

1306 Zhang, J., Xie, F., Li, N., Chen, R., Zhou, Z. and Zhou, T.: (HPLC determination of 2 main active constituents in *Ligusticum species*). *Yaowu Fenxi Zazhi*, 20 (2000) 301-303; *C.A.*, 134 (2001) 33063m.

1307 Zhang, Z., Diao, X., Yuan, C., Pei, G. and Wang, Z.: (Determination of ferulic acid in Tongjing tablets by HPLC). *Zhongcaoyao*, 31 (2000) 743-744; *C.A.*, 134 (2001) 46926v.

See also 1203, 1238, 1253, 1330, 1333, 1533, 1544, 1655, 1658, 1994, 2008, 2026, 2047, 2090, 2092, 2099.

11b. Prostaglandins

1308 Bylund, J., Hidestrand, M., Ingelman-Sundberg, M. and Oliw, E.H.: Identification of CYP4F8 in human seminal vesicles as a prominent 19-hydroxylase of prostaglandin endoperoxides. *J. Biol. Chem.*, 275 (2000) 21844-21849.

11c. Lipids and their constituents

1309 Araki, S., Yamada, S., Abe, S., Waki, H., Kon, K., Itonori, S., Sugita, M. and Ando, S.: Characterization of a novel triphosphonooctaosylceramide from the eggs of the sea hare, *Aplysia kurodai*. *J. Biochem. (Tokyo)*, 129 (2001) 93-100.

1310 Bonis, F.T.: (Evaporative light scattering detectors: an ideal tool for HPLC analysis of triglycerides). *Tec. Lab.*, 22, No. 248 (2000) 27-29; *C.A.*, 134 (2001) 27092d.

1311 Gertz, C., Klostermann, S. and Kochhar, S.P.: Testing and comparing oxidative stability of vegetable oils and fats at frying temperature. *Eur. J. Lipid Sci. Technol.*, 102 (2000) 543-551; *C.A.*, 134 (2001) 28598k.

1312 Itabashi, Y., Myher, J.J. and Kuksis, A.: High-performance liquid chromatographic resolution of reverse isomers of 1,2-diacyl-rac-glycerols as 3,5-dinitrophenylurethanes. *J. Chromatogr. A*, 893 (2000) 261-279.

1313 Lin, J.-Y., Lew, K.M., Chen, J.M. and McKeon, T.A.: Separation of the molecular species of intact phosphatidylethanolamines and their N-monomethyl and N,N-dimethyl derivatives by high-performance liquid chromatography of a C_8 column. *J. Chromatogr. A*, 891 (2000) 349-353.

1314 Wuhrer, M., Dennis, R.D., Doenhoff, M.J. and Geyer, R.: Stage-associated expression of ceramide structures in glycosphingolipids from the human trematode parasite *Schistosoma mansoni*. *Biochim. Biophys. Acta*, 1524 (2000) 155-161.

See also 1120, 1289, 1292, 1665, 1666, 1689.

11d. Lipoproteins and their constituents

1315 Batthyany, C., Santos, C.X.C., Botti, H., Cervenansky, C., Radi, R., Augusto, O. and Rubbo, H.: Direct evidence for apo B-1000-mediated copper reduction: studies with purified apo B-100 and detection of tryptophanyl radicals. *Arch. Biochem. Biophys.*, 384 (2000) 335-340.

1316 Haginaka, J., Yamaguchi, Yu. and Kunitono, M.: Anion-exchange high-performance liquid chromatography assays of plasma lipoproteins and modified low-density lipoproteins using a ProtEx-DEAE column. *J. Chromatogr. B*, 751 (2001) 161-167.

1317 Nofer, J.-K., Fobker, M., Höbbel, G., Voss, R., Wolinska, I., Tepel, M., Zidek, W., Junker, R., Seedorf, N., von Eckardstein, A., Assmann, G. and Walter, M.: Activation of phosphatidylinositol-specific phospholipase C by HDL-associated lysosphingolipid. Involvement in mitogenesis but not in cholesterol efflux. *Biochemistry*, 39 (2000) 15199-15207.

12. ORGANIC PEROXIDES

See 1533.

13. STEROIDS

13a. General techniques

1318 Pfeffer, M.: (Automation in the development of HPLC methods). *GIT Labor-Fachz.*, 44 (2000) 900-903; *C.A.*, 133 (2000) 340329r.

1319 Szanya, T., Argyelan, J., Kovats, S. and Hanak, L.: Separation of steroid compounds by overloaded preparative chromatography with percipitation in the fluid phase. *J. Chromatogr. A*, 908 (2001) 265-272.

See also 1831, 1885.

13b. Pregnane and androstane derivatives

1320 Coldham, N.G. and Sauer, M.J.: Analysis of [^{3}H]19-nortestosterone metabolism by bovine hepatocytes using narrow bore radio-HPLC-MS. *CAST, Chromatogr. Sep. Technol.*, 13 (2000) 18-19; *C.A.*, 133 (2000) 232957k.

1321 De Menezes, F.G., Ribeiro, W., Ifa, D.R., Amaral de Moraes, M.E., Amaral de Moraes, M.O. and de Nucci, G.: Bioequivalence study of finasteride. Determination in human plasma by high-performance liquid chromatography coupled to tandem mass spectrometry. *Arzneim.-Forsch.*, 51 (2001) 145-150.

1322 Jia, Q., Hong, M.-F., Pan, Z.-X. and Orndorff, S.: Quantification of urine 17-ketosteroid sulfates and glucuronides by high-performance liquid chromatography-ion trap mass spectroscopy. *J. Chromatogr. B*, 750 (2001) 81-91.

1323 Kao, Y.-C., Korzekwa, K.R., Laughton, C.A. and Chen, S.: Evaluation of the mechanism of aromatase cytochrome P450. A site-directed mutagenesis study. *Eur. J. Biochem.*, 268 (2001) 243-251.

1324 Wu, Z., Zhang, Ch., Yang, Ch., Zhang, X. and Wu, E.: Simultaneous quantitative determination of norgestrel and progesterone in human serum by HPLC-tandem MS with atmospheric pressure chemical ionization. *Analyst (Cambridge)*, 125 (2000) 2201-2205.

See also 1328, 1531, 1750.

13c. Estrogens

1325 Blom, M.J., Wassink, M.G., Kloosterboer, H.J., Ederveen, A.G.H., Lambert, J.G.D. and Goos, H.J.Th.: Metabolism of estradiol, ethynylestradiol, and moxestrol in rat uterus, vagina, and aorta: influence os sex steroid treatment. *Drug Metab. Disp.*, 29 (2001) 76-81.

1326 Kuang, P., Li, P., Luo, X. and Li, Y.: (Determination of diethylstilbestrol in pastes by HPLC). *Yaowu Fenxi Zazhi,k* 20 (2000) 351-352; *C.A.*, 134 (2001) 21575s.

1327 Lopez de Alda, M.J. and Barcelo, D.: Determination of steroid sex hormones and related synthetic compounds considered as endocrine disrupters in water by fully automated on-line solid-phase extraction-liquid chromatography-diode array detection. *J. Chromatogr. A*, 911 (2001) 203-210.

1328 Naga, S., Shimizu, J., Okada, N., Yokota, C., Tamaru, M. and Maruta, K.: Determination of estrone, estradiol, and progesterone in human saliva by three-column-switching high-performance liquid chromatography. *Seibutsu Shiryo Bunseki,* 23 (2000) 252-257; *C.A.,* 133 (2000) 247376u.

1329 Satoh, T., Fujita, K.-i., Munakata, H., Itoh, S., Nakamura, K., Kamataki, T., Itoh, S. and Yoshizawa, I.: Studies on the interactions between drugs and estrogen: analytical method for prediction system of gynecomastia induced by drugs on the inhibitory metabolism of estradiol using *Escherichia coli* coexpressing human CYP3A4 with human NADPH-cytochrome P450 reductase. *Anal. Biochem.,* 286 (2000) 179-186.

1330 Xu, S., Zhu, B.T. and Conney, A.H.: Stimulatory effect of clofibrate and gemfibrozil administration on the formation of fatty acid esters of estradiol by rat liver microsomes. *J. Pharmacol. Exp. Ther.,* 296 (2001) 188-197.

1331 Zhu, J. and Luo, X.-P.: (HPLC determination of estradiol benzoate injection). *Zhongguo Yiyao Gongye Zazhi,* 31, No. 6 (2000) 272-273; *C.A.,* 134 (2001) 21558p.

See also 1323.

13d. Sterols

1332 Brown, A.J., Watts, G.F., Burnett, J.R., Dean, R.T. and Jessup, W.: Sterol 27-hydroxylase acts on 7-ketocholesterol in human atherosclerotic lessions and macrophages in culture. *J. Biol. Chem.,* 275 (2000) 27627-27633.

1333 Havrilla, C.M., Hachey, D.L. and Porter, N.A.: Coordination (Ag^+) ion spray-mass spectrometry of peroxidation products of cholesterol linoleate and cholesterol arachidonate: High-performance liquid chromatography-mass spectrometry analysis of peroxide products from polyunsaturated lipid autoxidation. *J. Am. Chem. Soc.,* 122 (2000) 8042-8055; *C.A.,* 133 (2000) 292403m.

1334 Kenig, F., Popp, B.N. and Summons, R.E.: Preparative HPLC with ultrastable-Y zeolite for compound-specific carbon isotopic analyses. *Org. Geochem.,* 31 (2000) 1087-1094; *C.A.,* 134 (2001) 58795q.

1335 Ma, Y., Bi, K., Wang, X. and Zhou, X.: (Determination of cyasterone in medicinal cyathula *(Cyathula officinalis)* root by RP-HPLC). *Zhongcaoyao,* 31 (2000) 427-428; *C.A.,* 133 (2000) 256887p.

See also 1289.

13e. Bile acids and alcohols

1336 Constantino, G., Wolf, C., Natalini, B. and Pellicciari, R.: Evaluation of hydrophotic-hydrophilic balance of bile acids by comparative molecular field analysis (CoMFA). *Steroids,* 65 (2000) 483-489; *C.A.,* 134 (2001) 2276d.

1337 Yeh, Y.H. and Hwang, D.-F.: High-performance liquid chromatographic determination for bile components in fish, chicken and duck. *J. Chromatogr. B,* 751 (2001) 1-8.

14. STEROID GLYCOSIDES AND SAPONINS

1338 Eder, M. and Mehnert, W.: Lösyngsverhgalten von Digoxin in Gegenwart pflanzlicher Begleitstoffe aus *Digitalis-lanata*-Trockenextrakten. *Pharmazie,* 55 (2000) 928-933.

1339 Li, J.: (Application of HPLC-ELSD on quality control of aescin for injection). *Zhongcaoyao,* 31 (2000) 582-583; *C.A.,* 133 (2000) 366507x.

1340 Vakkuri, O., Arnason, S.S., Joensuu, P., Jalonen, J., Vuolteenaho, O. and Lepaluoto, J.: Radioiodinated tyrosyl-ouabain and measurement of a circulating ouabain-like compound. *Clin. Chem. (Washington),* 47 (2001) 95-101.

See also 1983, 2011, 2021.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

1341 Clarot, I., Cledat, D., Battu, S. and Cardot, P.J.P.: Chromatographic study of terpene derivatives on porous graphitic carbon stationary phase with β-cyclodextrin as mobile phase modifier. *J. Chromatogr. A,* 903 (2000) 67-76.

1342 Jouenne, E. and Crouzet, J.: Determination of apparent binding constants for aroma compounds with β-lactoglobulin by dynamic coupled column liquid chromatograpy. *J. Agric. Food Chem.,* 48 (2000) 5396-5400.

1343 Ortelli, D., Rudaz, S., Cognard, E. and Veuthey, J.-L.: Analysis of dihydroartemisin in plasma by liquid chromatography-mass spectrometry. *Chromatographia,* 52 (2000) 445-450.

1344 Pinto, A.C., Braga, W.F., Rezende, C.M., Garrido, F.M.S., Veiga, V.F., Jr., Bergter, L., Patitucci, M.L. and Antunes, O.A.C.: Separation of acid diterpenes of *Copaifera cearensis* Huber ex Ducke by flash chromatography using potassium hydroxide impregnated silica gel. *J. Braz. Chem. Soc.,* 11 (2000) 355-360; *C.A.,* 134 (2001) 32839g.

1345 Sessa, R.A., Bennett, M.H., Lewis, M.J., Mansfield, J.W. and Beale, M.H.: Metabolite profiling of sesquiterpene lactones from Lactuca species. *J. Biol. Chem.,* 275 (2000) 26877-26884.

1346 Yuping, T. and Fengchang, L.: Separation and isolation of terpene lactones from *Ginkgo biloba* L. by direct high performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2897-2900.

See also 1334, 2046.

15b. Essential oils

1347 Dugo, G., Bartle, K.D., Bonaccorsi, I., Catalfamo, M., Cotroneo, A., Dugo, P., Lamonica, G., McNair, H., Mondello, L., Previti, P., D'Alcontres, I.S., Trozzi, A. and Verzera, A.: Advanced analytical techniques for the analysis of citrus essential oils. Part 3. Oxygen heterocyclic compounds: HPLC, HPLC/MS, OPLC, SFC, fast HPLC analysis. *Essenze, Deriv. Agrum.,* 69, No. 4 (1999) 251-283; *C.A.,* 133 (2000) 300904t - a review with refs.

15c. Bitter substances

1348 Harms, D. and Nitzsche, F.: (New ways to determine bitter substances in beer and wort). *Monatsschr. Brauwiss.*, 53, No. 7/8 (2000) 143-146; *C.A.*, 134 (2001) 41242j.

1349 Lin, Y., Shan, L., Pei, Z. and Qian, Z.: (Determination of α-acids and β-acids in hops, hop pellets, and hop extracts by high performance liquid chromatography). *Fenxi Huaxue*, 28 (2000) 1047; *C.A.*, 133 (2000) 334207h.

16. NITRO AND NITROSO COMPOUNDS

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

1358 Faaland, C.A., Thomas, T.J., Balabhadrapathruni, S., Langer, T., Miam, S., Shirahata, A., Gallo, M.A. and Thhomas, T.: Molecular correlates of the action of bis(ethyl)polyamines in breast cancer cell growth inhibition and apoptosis. *Biochem. Cell Biol.*, 78 (2000) 415-426.

1359 Gadzala-Kopciuch, R.M., Buszewska, T. and Buszewski, B.: Isolation and determination of denatonium benzoate in antifreezing agents by solid phase extraction and HPLC. *J. Liq. Chroma-*

fluorenylmethyloxycarbonyl chloride derivatization. *J. Chromatogr. B*, 749 (2000) 179-189.

1369 Hatsuda, K., Tsuji, T. and Katayama, Y.: (Evaluation of the analytical performance of TOSOH catecholamine analyzer HCL-725CA improved type.) *Seibutsu Shiryo Bunseki,* 22 (1999) 343-350; *C.A.,* 133 (2000) 249150q.

1370 Lu, J., Zhang, S., Wang, A., Zhang, W. and Jin, L.: *In vivo* monitoring of the monoamine neurotransmitters in the rat brain by microdialysis coupled with the liquid chromatography dual electrochemical detector. *Talanta,* 52 (2000) 807-815; *C.A.,* 133 (2000) 232963j.

1371 Suciu, A.O. and Vilceanu, N.: (Purity evaluation study of methyldopa lot I USP-standard substance. Part 1. HPLC purity testing of methyldopa). *Farmacia (Bucharest),* 48 (2000) 63-72; *C.A.,* 134 (2001) 21547j.

See also 1041.

17c. Urea and guanidine derivatives

1372 Nohara, Y., Hanai, T., Kinoshita, T. and Watanbe, M.: Chemiluminescence-HPLC for the assay of guanidino compounds. *Chem. Pharm. Bull.,* 48 (2000) 1841-1842.

17d. Other amine derivatives and amides (excl. peptides)

1373 Evgen'ev, M.I., Evgen'eva, I.I. and Ismailova, R.N.: Extraction-chromatographic determination of hydrazine in natural water as a 5,7-dinitrobenzofurazan derivative using diode-array detection. *J. Anal. Chem.,* 55 (2000) 933-937; *C.A.,* 133 (2000) 313266m.

1374 Keating, T.A., Marshall, C.G. and Walsh, Ch.T.: Reconstitution and characterization of the *Vibrio cholerae* vibriobactin synthetase from VibB, VibE, VibF and VibH. *Biochemistry,* 39 (2000) 15522-15530.

1375 Keating, T.A., Marshall, C.G. and Walsh, Ch.T.: Vibriobactin biosynthesis in *Vibrio cholerae*: VibH is an amide synthase homologous to nonribosomal peptide synthetase condensation domains. *Biochemistry,* 39 (2000) 15513-15521.

1376 Santolaya, C., Bernal, F., Guardino, X. and Arias, M.C.: Determination of isocyanates in spray paint booths of auto repair shops using HPLC, UV, and electrochemical detection. *Am. Lab. (Shelton),* 32, No. 17 (2000) 44-46; *C.A.,* 133 (2000) 339773z.

See also 1067, 1191, 1313, 1614.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

1377 Beauge, L., Siravegna, M. and Barberian, G.: Enzymatic synthesis of ^{32}P-labeled phosphoarginine. *Anal. Biochem.,* 286 (2000) 306-208.

1378 Guo, J., Xiao, F. and Tang, Z.: (Determination of homocysteine in plasma by high performance liquid chromatography). *Zhonghua Yixue Jianyan Zazhi,* 23 (2000) 217-219; *C.A.,* 133 (2000) 171693x.

1379 Hermann, K. and Abeck, D.: Analysis of histidine and urocainic acid isomers by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B,* 750 (2001) 71-80.

1380 Hojo, K., Maeda, M., Iguchi, S., Smith, T., Okamoto, H. and Kawasaki, K.: Amino acids and peptides. XXXV. Facile preparation of *p*-nitroanilide analogs by the solid-phase method. *Chem. Pharm. Bull.,* 48 (2000) 1740-1744.

1381 Huang, J. and Williams, M.: Total plasma homocysteine determination using ion-exchange chromatography. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3143-3153.

1382 Jandik, P., Cheng, J., Jensen, D., Manz, S. and Avdalovic, N.: New technique for increasing retention of arginine on an anion-exchange column. *Anal. Biochem.,* 287 (2000) 38-44.

1383 Kawatsu, K., Hamano, Y. and Noguchi, T.: Determination of domoic acid in Japanese mussels by enzyme immunoassay. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1384-1386.

1384 McMurtrey, K., Strawbridge, C. and McCoy, J.: HPLC resolution of the enantiomers of dihydroxyphenylalanine and selected salsolinol derivatives using sulfated β-cyclodextrin. *Enantiomer,* 5 (2000) 377-383; *C.A.,* 134 (2001) 76473n.

1385 Minniti, G., Cerone, R., Piana, A., Armani, U. and Lorini, R.: Plasma and serum total homocysteine concentrations in pediatric patients, evaluated by high-performance liquid chromatography with fluorescence. *Clin. Chem. Lab. Med.,* 38 (2000) 675-676; *C.A.,* 134 (2001) 68220n.

1386 Miyabe, K. and Guiochon, G.: Kinetic study of the concentration dependence of the mass transfer rate coefficient in enantiomeric separation on a polymeric imprinted stationary phase. *Anal. Sci.,* 16 (2000) 719-730; *C.A.,* 133 (2000) 257343v.

1387 Morikawa, A., Hamase, K., Konno, R. and Zaitsu, K.: (Determination of free D-Asp, D-Ser and D-Ala in the brain of mutant mouse lacking D-amino acid oxidase activity). *Chromatography,* 21, No. 2 (2000) 157-158; *C.A.,* 134 (2001) 68212m.

1388 Nagren, K., Sutinen, E. and Jyrkkio, S.: [N-Methyl-^{11}C]MeAIB, a tracer for system a amino acid transport: preparation from [^{11}C]methyl triflate and HPLC metabolite analysis of plasma samples after intravenous administration in man. *J. Labelled Compd. Radiopharm.,* 43 (2000) 1013-1021; *C.A.,* 134 (2001) 2139m.

1389 Okubo, S., Mashige, F., Hashimoto, Y., Nakahara, K., Kanazawa, H., Matsushima, Y., Fukushima, T., Huang, Y. and Imai, K.: D-serine concentration in blood serum of liver-transplanted patients. *Chromatography,* 21 (2000) 43-47; *C.A.,* 133 (2000) 249100y.

1390 Schlauch, M., Volk, F.-J., Fondekar, K.P., Wede, J. and Frahm, A.W.: Enantiomeric and diastereomeric high-performance liquid chromatographic separation of cyclic β-substituted α-amino acids on a copper(II)-D-penicillamine chiral stationary phase. *J. Chromatogr. A,* 897 (2000) 145-152.

1391 Stancher, B. and Calabrese, M.: (Presence of homocysteine in beer samples). *Ind. Bevande,* 29 (2000) 137-146; *C.A.,* 133 (2000) 280867c.

1392 Wang, J., Zhou, L., Liu, X., Wang, J. and Shang, Z.: (Simultaneous chiral separation of two pairs of enantiomers using series column of molecular imprinted phase). *Fenxi Huaxue,* 28 (2000) 1096-1098; *C.A.,* 134 (2001) 36510r.

1393 Werner, U., Brune, K., Layh, B. and Guehring, H.: Determination of L-iminoethyl-L-lysine in serum by liquid chromatography. *J. Chromatogr. B,* 749 (2000) 35-40.

1394 Whittle, M.C., Stobaugh, J.F. and Wong, O.S.: High performance liquid chromatographic method for the determination of homocysteine in human plasma using a water-soluble phosphine reducing agent and fluorogenic derivatization. *Chromatographia,* 52(Supplement) (2000) S78-S82.

1395 Woo, K.-L.: Determination of amino acids in foods by reversed-phase high-performance liquid chromatography with new precolumn derivatives, butylthiocarbamyl, and benzylthiocarbamyl derivatives compared to the phenylthiocarbamyl derivative and ion exchange chromatography. *Methods Mol. Biol. (Totowa),* 159(Amino Acid Analysis Protocols) (2001) 141-147; *C.A.,* 134 (2001) 70465q.

1396 Yan, J.X. and Packer, N.H.: Amino acid analysis of mucins. *Methods Mol. Biol. (Totowa),* 125(Glycoprotein Methods and Protocols) (2000) 113-119; *C.A.,* 133 (2000) 278164c.

1397 Zachara, N.E. and Gooley, A.A.: Identification of glycosylation sites in mucin peptides by Edman degradation. *Methods Mol. Biol. (Totowa),* 125(Glycoprotein Methods and Protocols) (2000) 121-128; *C.A.,* 133 (2000) 278165d.

See also 1002, 1041, 1155, 1156, 1269, 1365, 1399, 1435, 1452, 1841.

18b. Peptides, peptidic and proteinous hormones, growth factors

1398 Asami, Y., Nagano, H., Ikematsu, S. and Murasugi, A.: An approach to the removal of yeast specific *O*-linked oligo-mannoses from human midkine expressed *Pichia pastoris* using site-specific mutagenesis. *J. Biochem. (Tokyo),* 128 (2000) 823-826.

1399 Bar-Or, D., Curtis, G., Rao, N., Bampos, N. and Lau, E.: Characterization of the Co^{2+} and Ni^{2+} binding amino-acid residues of the N-terminus of human albumin. An insight into the mechanism of a new assay for myocardial ischemia. *Eur. J. Biochem.,* 268 (2001) 42-47.

1400 Belozersky, M.A., Dunaevsky, Y.E., Musolyamov, A.K. and Egorov, T.A.: (Complete amino acid sequence of the anionic protease inhibitor BWI-4a from buckwheat seeds). *Biokhimiya (Moscow),* 65 (2000) 1347-1352.

1401 Bertola, F., Banigand, C., Picard, P., Belghazi, M. and Precigoux, G.: Human T-lymphotrophic virus type I nucleocapsid protein NCp15: structural study and stability of the N-terminal zinc-finger. *Biochem. J.,* 352 (2000) 293-300.

1402 Bidnychenko, Y.: HPLC determination of mushroom peptide toxins in forensic-chemical examination. *Am. Lab. (Shelton),* 32, No. 9 (2000) 14; *C.A.,* 133 (2000) 306436p.

1403 Bonetto, V., Eriste, E., Jonsson, A.P., Efendic, S., Jörnvall, H. and Sillard, R.: Isolation of peptides from porcine intestinal tissue that induce extracellular acidification in CHO cells: identification of the active peptide as IGF-I and characterization of a fragment of calponin H1 processed at a dibasic site. *Arch. Biochem. Biophys.,* 385 (2001) 276-282.

1404 Brittain, S.M., Wang, J., Babock-Jackson, L., Carmichael, W.W., Rinehart, K.L. and Culver, D.A.: Isolation and characterization of microcystins, cyclic heptapeptide hepatotoxins from a Lake Erie strain of *Microcystis aeruginosa. J. Great Lakes Res.,* 26 (2000) 241-249; *C.A.,* 133 (2000) 354816j.

1405 Byun, C.K., Song, J.H., Lee, S.K., Kim, D.H. and Lee, D.W.: Salt effects on the retention of peptides in hydrophobic interaction chromatography. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2963-2978.

1406 Deacon, C.F., Nauck, M.A., Meier, J., Hucking, K. and Holst, J.J.: Degradation of endogenous and exogenous gastric inhibitory polypeptide in healthy and in type 2 diabetic subjects as revealed using a new assay for the intact peptide. *J. Clin. Endocrinol. Metab.,* 85 (2000) 3575-3581; *C.A.,* 133 (2000) 344749a.

1407 Desportes, C., Charpentier, M., Duteurtre, B., Maujean, A. and Duchiron, F.: Liquid chromatographic fractionation of small peptides from wine. *J. Chromatogr. A,* 893 (2000) 281-291.

1408 Filipe, S.R., Pinko, M.G. and Tomasz, A.: Characterization of the murMN operon involved in the synthesis of branched peptidoglycan peptides in *Streptococcus pneumoniae. J. Biol. Chem.,* 275 (2000) 27768-27774.

1409 Hale, J.E., Butler, J.P., Knierman, M.D. and Becker, G.W.: Increased sensitivity of tryptic peptide detection by MALDI-TOF mass spectrometry is achieved by conversion of lysine to homoarginine. *Anal. Biochem.,* 287 (2000) 110-117.

1410 Hattori, A., Kitatani, K., Matsumoto, H., Miyazawa, S., Rogi, T., Tsuruoka, N., Mizutani, S., Natori, Y. and Tsujimoto, M.: Characterization of recombinant human adipocyte-derived leucine aminopeptidase expressed in Chinese hamster ovary cells. *J. Biochem. (Tokyo),* 128 (2000) 755-762.

1411 Hidaka, Y., Shimono, Ch., Ohno, M., Okumura, N., Adermann, K., Forssman, W.-G. and Shimonishi, Y.: Dual function of the propeptide of prouroguanylin in the folding of the mature peptide. *J. Biol. Chem.,* 275 (2000) 25155-25162.

1412 Hidalgo, F.J., Alaiz, M. and Zamora, R.: Determination of peptides and proteins in fats and oils. *Anal. Chem.,* 73 (2001) 698-702.

1413 Ishihama, Y., Katayama, H. and Asakawa, N.: Surfactants usable for electrospray ionization mass spectrometry. *Anal. Biochem.,* 287 (2000) 45-54.

1414 Lee, K.R., Bongers, J., Gulati, D. and Burman, S.: Statistical validation of reproducibility of HPLC peptide mapping for the identity of an investigational drug compound based on principal component analysis. *Drug Dev. Ind. Pharm.,* 26 (2000) 1045-1057; *C.A.,* 134 (2001) 25083c.

1415 Liu, X.-Y., Yang, Q., Nakamura, C. and Miyake, J.: Avidin-biotin-immobilized liposome column for chromatographic fluorescence on-line analysis of solute-membrane interactions. *J. Chromatogr. B,* 750 (2001) 51-60.

1416 Menyes, U., Kessler, D. and Roth, U.: (Separation of peptides by HPLC Caltrex partition phases). *GIT Spez. Sep.,* 20, No. 2 (2000) 84-85; *C.A.,* 133 (2000) 346617e.

1417 Moffatt, F., Senkans, P. and Ricketts, D.: Approaches towards the quantitative analysis of peptides and proteins by reversed-phase high-performance liquid chromatography in the absence of a pure reference sample. *J. Chromatogr. A,* 891 (2000) 235-242.

1418 Nagy, I.B., Varga, I. and Hudecz, F.: Preparation of indium-115-labeled diethylenetriaminotetraacetic acid monoacetamide peptides purified by 8-hydroxyquinoline. *Anal. Biochem.,* 287 (2000) 17-24.

1419 Nardai, G., Sass, B., Eber, J., Orosz, G. and Csermely, P.: Reactive cysteines of the 90-kDa heat shock protein, Hsp90. *Arch. Biochem. Biophys.,* 384 (2000) 59-67.

1420 Neal, R., Zigler, J.S., Jr. and Bettelheim, F.A.: On the equilibrium between monomeric α-lactalbumin and the chaperoning complex of α-crystallin. *Biochem. Biophys. Res. Commun.,* 280 (2001) 14-18.

1421 Nemeth, J., Vecsernyes, M., Oroszi, G., Balla, Z., Helyes, Z., Farkas, B. and Szilvassy, Z.: Preparation of mono-^{125}I-labelled gastrin-17 for radioimmunoassay measurements. *J. Labelled Compd. Radiopharm.*, 43 (2000) 855-863; *C.A.*, 133 (2000) 278296x.

1422 Odani, S., Baba, K., Tsuchida, Y., Aoyagi, Y., Wakui, S. and Takahashi, Y.: Hepatic fatty acid-binding proteins of a teleost, *Lateolabrax japonicus*. The primary structures and location of a disulfide bond. *J. Biochem. (Tokyo)*, 129 (2001) 69-76.

1423 Oh, J.-E. and Lee, K.-H.: Characterization of the unique function of a reduced amide bond in a cytolytic peptide that acts on phospholipid membranes. *Biochem. J.*, 352 (2000) 659-666.

1424 Oliva, A., Farina, J. and Llabres, M.: Development of two high-performance liquid chromatographic methods for the analysis and characterization of insulin and its degradation products in pharmaceutical preparations. *J. Chromatogr. B*, 749 (2000) 25-34.

1425 Sack, R., Willi, A. and Hunziker, P.E.: Determination of total glutathione in cell lysates by high performance liquid chromatography with *o*-phthalaldehyde precolumn derivatization in the presence of tris(2-carboxyethyl)-phosphine. *J. Liq. Chromatogr. Relat. Technol.*, 23 (2000) 2947-2962.

1426 Sasaki, H., Yamamura, K., Mukai, T., Nishida, K., Nakamura, J., Nakashima, M. and Ichikawa, M.: Modification of ocular permeability of peptide drugs by absorption promoters. *Biol. Pharm. Bull.*, 23 (2000) 1524-1527.

1427 Sergeev, N.V., Gloukhova, N.S., Nazimov, I.V., Gulyaev, V.A., Shvets, S.V., Donetsky, I.A. and Miroshnikov, A.I.: Monitoring of recombinant human insulin production by narrow-bore reversed-phase high-performance liquid chromatography, high-performance capillary electrophoresis and matrix-assisted laser desorption ionisation time-of-fligh mass spectrometry. *J. Chromatogr. A*, 907 (2001) 131-144.

1428 Sneller, F.E.C., van Heerwaarden, L.M., Koevoets, P.L.M., Vooijs, R., Schat, H. and Verkleij, J.A.C.: Derivatization of phytochelatins from *Silene vulgaris*, induced upon exposure to arsenate and cadmium: comparison of derivatization with Ellman's reagent and monobromobimane. *J. Agric. Food Chem.*, 48 (2000) 4014-4019.

1429 Su, G.: (Determination of ornithinaspartate in hepamerz injection by precolumn derivatization of HPLC). *Guangdong Yaoxueyuan Xuebao*, 16, No. 2 (2000) 100-104; *C.A.*, 133 (2000) 271777w.

1430 Taguchi, S., Kuwasako, K., Suenaga, A., Okada, M. and Momose, H.: Functional mapping against *Escherichia coli* for the broad-spectrum antimicrobial peptide, thanatin, based on an *in vivo* monitoring assay system. *J. Biochem. (Tokyo)*, 128 (2000) 745-754.

1431 Torfs, P., Nieto, J., Cerstiaens, A., Boon, D., Baggerman, G., Poulos, C., Waelkens, E., Derua, R., Calderon, J., de Loof, A. and Schoofs, L.: Pyrokinin neuropeptides in a crustacean. Isolation and identification in the white shrimp *Penaeus vannamei*. *Eur. J. Biochem.*, 268 (2001) 149-154.

1432 Wagner, K., Racaityte, K., Unger, K.K., Miliotis, T., Edholm, L.E., Bischoff, R. and Marko-Varga, G.: Protein mapping by two-dimensional high performance liquid chromatography. *J. Chromatogr. A*, 893 (2000) 293-305.

1433 Wang, Z., Vandenberghe, I., Depreitere, J., Devreese, B., Clerens, S., Neuwen, E.J., van Beemen, J. and de Potter, W.: Identification and characterization of novel chromogranin B-derived peptides from porcine chromaffin granules by liquid chromatography/electrospray tandem MS. *Eur. J. Biochem.*, 268 (2001) 235-242.

1434 Xia, C.Q., Wang, J. and Shen, W.-C.: Hypoglycemic effect of insulin-transferrin conjugate in streptozotocin-induced diabetic rats. *J. Pharmacol. Exp. Ther.*, 295 (2000) 594-600.

1435 Yoshida, T., Okada, T., Hobo, T. and Chiba, R.: Calculation of amino acid hydrophilicity indices for retention of peptides on amide, diol and silica columns in normal-phase liquid chromatography. *Chromatographia*, 52 (2000) 418-424.

1436 Ysay, Y.-G., Wang, Y.-H., Chiu, C.-M., Shen, B.-J. and Lee, S.-C.: A strategy for identification and quantitation of phosphopeptides by liquid chromatography-tandem mass spectrometry. *Anal. Biochem.*, 287 (2000) 55-64.

See also 1110, 1354, 1380, 1514, 1516, 1603.

18c. Elucidation of structure of proteins and enzymes

1437 Bingham, R., Ekunwe, S.I.N., Falk, S., Snyder, L. and Kleanthous, C.: The mayor head protein of bacteriophage T4 binds specifically to elongation factor Tu. *J. Biol. Chem.*, 275 (2000) 23219-23226.

1438 Braun, K., Kuhl, P., Bernd, M. and Kutscher, B.: Stability of several LHRH antagonists against proteolytic enzymes and identification of degradation products by mass spectrometry. *Pharmazie*, 56 (2001) 45-49.

1439 Cairns, J.R.K., Champattanachai, V., Srisomsap, C., Wittman-Liebold, B., Thiede, B. and Svasti, J.: Sequence and expression of Thai rosewood β-glucosidase/β-fucosidase, a family 1 glycosyl hydrolase glycoprotein. *J. Biochem. (Tokyo)*, 128 (2000) 999-1008.

1440 Daoud, R., Kast, Ch., Gros, P. and Georges, E.: Rhodamine 123 binds to multiple sites in the multidrug resistance proteins (MRP1). *Biochemistry*, 39 (2000) 15344-15352.

1441 Takemoto, L. and Boyle, D.: Specific glutamine and asparagine residues of γ-S crystallin are resitant to *in vivo* deamination. *J. Biol. Chem.*, 275 (2000) 26109-26112.

1442 Tung, P.P. and Alter, G.M.: Substrate and cofactor binding to fluorescently labeled cytoplasmic malate dehydrogenase. *Biochim. Biophys. Acta*, 1545 (2001) 132-145.

See also 1397, 1414, 1422, 1432, 1483, 1494, 1514, 1516.

19. PROTEINS

19a. General techniques

1443 Bures, E.J., Courchesne, P.L., Douglass, J., Chen, K., Davis, M.T., Jones, M.D., McGinley, M.D., Robinson, J.H., Spahr, C.S., Sun, J., Wahl, R.C. and Patterson, S.D.: Identification of incompletely processed potential carboxypeptidase E substrates from CpE^{fat}/CpE^{fat} mice. *Proteomics (Weinheim)*, 1 (2001) 79-92.

1444 Chen, S.-H., Ho, C.-T., Hsiao, K.-Y. and Chen, J.-M.: Pressure-induced retention of the lysozyme on reversed-phase liquid chromatography. *J. Chromatogr. A*, 891 (2000) 207-215.

1445 Flurkey, W.H.: An inexpensive gradient maker for the biochemistry laboratory. *J. Chem. Educ.,* 77 (2000) 1041; *C.A.,* 133 (2000) 266224j.

1446 Hearn, M.T.W.: Physicochemical factors in polypeptide and protein purification and analysis by high-performance liquid chromatographic techniques. Current status and challenges for the future. *Sep. Sci. Technol. (San Diego),* 2(Handbook of Bioseparations) (2000) 71-235; *C.A.,* 134 (2001) 2148 - a review with 543 refs.

1447 Ishihara, T. and Yamamoto, S.: Separation behavior of proteins near the isoelectric points in electrostatic interaction (ion exchange) chromatography. *Prog. Biotechnol.,* 16(Bioseparation Engineering) (2000) 93-98; *C.A.,* 133 (2000) 319180a.

1448 Ivanov, A.E., Galaev, I.Y., Kazakov, S.V. and Mattiasson, B.: Thermosensitive copolymers of *N*-vinylimidazole as displacers of proteins in immobilised metal affinity chromatography. *J. Chromatogr. A,* 907 (2001) 115-130.

1449 Seely, J.E. and Richey, C.W.: Use of ion-exchange chromatography and hydrophobic interaction chromatography in the preparation and recovery of polyethylene glycol-linked proteins. *J. Chromatogr. A,* 908 (2001) 235-241.

1450 Shan, L. and Anderson, D.J.: Effect of buffer concentration of gradient chromatofocusing performance separating proteins on a high-performance DEAE column. *J. Chromatogr. A,* 909 (2001) 191-205.

1451 Tachibana, H. and Oka, T.: (Separation of disulfide-intermediates with reversed-phase HPLC). *Seibutsu Butsuri,* 40 (2000) 398-399; *C.A.,* 134 (2001) 2163 - a review with 6 refs.

1452 Walsh, M.K. and Brown, R.J.: Use of amino acid analysis for estimating the individual concentrations of proteins in mixtures. *J. Chromatogr. A,* 891 (2000) 355-360.

1453 Yamamoto, S. and Ishihara, T.: Resolution and retention of proteins near isoelectric points in ion-exchange chromatography. Molecular recognition in electrostatic interaction chromatography. *Sep. Sci. Technol.,* 35 (2000) 1707-1717; *C.A.,* 133 (2000) 331612a.

See also 1027, 1049, 1055, 1056, 1076, 1119, 1125, 1166, 1412, 1413, 1417, 1667.

19b. Proteins of cells, viruses and subcellular particles

1454 Bouvier, M. and Stafford, W.F.: Probing the three-dimensional structure of human calreticulin. *Biochemistry,* 39 (2000) 14950-14959.

See also 1437, 1479.

19c. Proteins synthesized by genetic manipulation, monoclonal antibodies

1455 Scott, M.P. and Miller, W.T.: A peptide model system for processive phosphorylation by Src family kinases. *Biochemistry,* 39 (2000) 14531-14537.

1456 Shen, Y.H., Zhang, Q.D., Geng, X.D., Zhang, Z.Q. and Hou, Y.D.: Purification of RhIFN- in bacteria bodies with acetic acid-water as mobile phase in RPLC. *Chin. Chem. Lett.,* 11 (2000) 799-802; *C.A.,* 133 (2000) 280236w.

See also 1466, 1484.

19d. Microbial and plant proteins

1457 Bolla, J.M., De, E., Dorez, A. and Pages, J.M.: Purification, characterization and sequence analysis of Omp50, a new purin isolated from *Campylobacter jejuni. Biochem. J.,* 352 (2000) 637-643.

1458 Ferrarello, C.N., Fernandez de la Campa, M.R., Muniz, C.S. and Sanz-Mendel, A.: Metal distribution patterns in the mussel *Mytilus edulis* cytosols using size-exclusion chromatography and double focusing ICP-MS detection. *Analyst (Cambridge),* 125 (2000) 2223-2229.

1459 Garcia, M.C., Marina, M. and Torre, M.: Determination of the soybean protein content in soybean liquid milks by reversed-phase HPLC. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3165-3174.

1460 Hanna, S.L., Sherman, N.E., Kinter, M.T. and Goldberg, J.B.: Comparison of proteins expressed by *Pseudomonas aeruginosa* strains representing initial and chronic isolates from a cystic fibrosis patient: an analysis by 2-D gel electophoresis and capillary column liquid chromatography tandem mass spetrometry. *Microbiology (Reading),* 146 (2000) 2495-2508; *C.A.,* 134 (2001) 68395y.

1461 Takagi, K., Yamamoto, K., Kano, K. and Ideda, T.: New pathway of amine oxidation respiratory chain of *Paracoccus denitrificans* IFO 12442. *Eur. J. Biochem.,* 268 (2001) 470-476.

1462 Wang, H.X. and Ng, T.B.: Flammulin: a novel ribosome-inactivating protein from fruiting bodies of the winter mushroom *Flammulina velutipes. Biochem. Cell Biol.,* 78 (2000) 699-702.

19e. Proteins of blood, serum and blood cells

1463 Branovic, K., Buchacher, A., Barut, M., Strancar, A. and Josic, D.: Application of monoliths for downstream processing of clotting factor IX. *J. Chromatogr. A,* 903 (2000) 21-32.

1464 Denton, G., Murray, A., Price, M.R. and Levison, P.R.: Direct isolation of monoclonal antibodies from tissue culture supernatant using the cation-exchange cellulose Express-Ion S. *J. Chromatogr. A,* 908 (2001) 223-234.

1465 Dubnovitsky, A.P., Kravchuk, Z.I., Chumanevich, A.A., Cozzi, A., Arosio, P. and Martsev, S.P.: (Expression, refolding and ferritin-binding activity of isolated VL-domain of monoclonal antibody F11). *Biokhimiya (Moscow),* 65 (2000) 1196-1204.

1466 Gibson, A.D. and Peterson, C.B.: Full-length and truncated forms of vitronectin provide insight into effects of proteolytic processing on function. *Biochim. Biophys. Acta,* 1545 (2001) 289-304.

1467 Grippa, E., Pavone, F., Gatto, M.T., Petrucci, R., Marrosu, G., Silvestrini, B. and Saso, L.: *In vitro* evaluation of antioxidant activity by electrophoresis and high performance liquid chromatography. *Biochim. Biophys. Acta,* 1524 (2000) 171-177.

1468 Hanai, T., Uchida, M., Amao, M., Ikeda, C., Koizumi, K. and Kinoshita, T.: Selective chemiluminescence analysis of Amadori form of glycated human serum albumin. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3119-3131.

1469 Inagaki, K., Umemura, T., Matsuura, H. and Haraguchi, H.: Speciation of trace elements, binding and non-binding with proteins in human blood serum, by surfactant-mediated HPLC with element-selective detection by ICP-MS. *Anal. Sci.,* 16 (2000) 787-788; *C.A.,* 133 (2000) 331687d.

1470 Katsumi, A., Tuley, E.A., Bodo, I. and Sadler, J.E.: Localization of disulfide bonds in the cystine knot domain of human von Willebrand factor. *J. Biol. Chem.*, 275 (2000) 25585-25594.

1471 Kouoh Elombo, F., Radosevich, M., Pulle, M., Descamps, J., Chtourou, S., Burnouf, T., Catteau, J.-P., Bernier, J.-L. and Cotelle, N.: Purification of human ceruloplasmin as a by-product of CI-inhibitor. *Biol. Pharm. Bull.*, 23 (2000) 1406-1409.

1472 Li, W., Zhang, D., Lin, B. and Su, Z.: Purification and identification of PEGlated hemoglobin, a potential blood substitute, by chromatography and capillary electrophoresis. *Chromatographia*, 52 (2000) 451-454.

1473 Matsumoto, A., Kojima, N., Takeuchi, F. and Mizuochi, T.: Kinetic analysis of interaction of different types of rheumatoid factors with immobilized IgG using surface plasmon resonance. *J. Biochem. (Tokyo)*, 128 (2000) 1009-1016.

1474 Mills, C.E., Sedelnikova, S., Soballe, B., Hughes, M.N. and Poole, R.K.: *Escherichia coli* flavohaemoglobin (Hmp) with equistoichiometric FAD and haem contents has a low affinity for dioxygen in the absence or presence of nitric acid. *Biochem. J.*, 353 (2001) 207-213.

1475 Minchiotti, L., Campagnoli, M., Rossi, A., Cosulich, M.E., Monti, M., Pucci, P., Kragh-Hansen, U., Granel, B., Disdier, P., Weiler, P.-J. and Galliano, M.: A nucleotide insertion and frameshift cuase albumin Kenitra, an extended and O-glycosylated mutant of human serum albumin with two additional disulfide bridges. *Eur. J. Biochem.*, 268 (2001) 344-352.

1476 Padrós, J. and Pelletier, É.: Subpicogram determination of (+)-anti-benzo[*a*]pyrene diol-epoxide adducts in fish albumin and globin by high-performance liquid chromatography with fluorescence detection. *Anal. Chim. Acta*, 426 (2001) 71-77.

1477 Schwartz, W., Judd, D., Wysocki, M., Guerrier, L., Birck-Wilson, E. and Boschetti, E.: Comparison of hydrophobic charge induction chromatography with affinity chromatography on protein A for harvest and purification of antibodies. *J. Chromatogr. A*, 908 (2001) 251-263.

1478 Thrash, M.E., Jr. and Pinto, N.G.: Flow microcalorimetric measurements for bovine serum albumin on reversed-phase and anion-exchange supports under overloaded conditions. *J. Chromatogr. A*, 908 (2001) 293-299.

1479 Yamamoto, S. and Miyagawa, E.: Dynamic binding performance of large biomolecules such as γ-globulin, viruses and virus-like particles on various chromatographic supports. *Prog. Biotechnol.*, 16(Bioseparation Engineering) (2000) 81-86; *C.A.*, 134 (2001) 14830t.

1480 Yan, Y., Hao, Q., Fan, G.-l., Zhao, Q.-w., Liu, Y.-w., Wang, J.-l. and Wan, D.-l.: (Preliminary study on the purification on IL-2.) *Sepu*, 18 (2000) 325-327; *C.A.*, 133 (2000) 265365u.

See also 1088, 1447, 1453.

19f. Structural and muscle proteins

1481 Au, V. and Madison, S.A.: Effects of singlet oxygen on the extracellular matrix protein collagen: oxidation of the collagen crosslink histidinohydroxylysinonorleucine and histidine. *Arch. Biochem. Biophys.*, 384 (2000) 133-142.

1482 Brandt, N.R., Franklin, G., Brunschwig, J.P. and Caswell, A.H.: The role of mitsugumin 29 in transverse tubules of rabbit skeletal muscle. *Arch. Biochem. Biophys.*, 385 (2001) 406-409.

1483 Luo, Y., Leszyk, J., Li, B., Gergely, J. and Tao, T.: Proximity relationships between residue 6 of troponin I and residues in troponin C: further evidence for extended conformation of troponin C in the troponin complex. *Biochemistry*, 39 (2000) 15306-15315.

1484 Piatibratov, M., Hou, S., Brooun, A., Yang, J., Chen, H. and Alam, M.: Expression and fast-flow purification of a polyhistidine-tagged myoglobin-like aerotaxis transducer. *Biochim. Biophys. Acta*, 1524 (2000) 149-154.

1485 Ward, D.G., Ashton, P.R., Trayer, H.R. and Trayer, I.P.: Additional PKA phosphorylation sites in human cardiac troponin I. *Eur. J. Biochem.*, 268 (2001) 179-185.

19g. Protamines, histones and other nuclear proteins

See 1441.

19h. Chromoproteins and metalloproteins

1486 Belzer, C.A., Tabatabai, L.B. and Frank, G.H.: Purification and characterization of the *Pasteurella haemolytica* 35 kiloDalton periplasmic iron-regulated protein. *Prepar. Biochem. Biotechnol.*, 30 (2000) 343-355.

1487 Ferrarello, C.N., Fernandez de la Campa, M.R., Carrasco, J.F. and Sanz-Medel, A.: Speciation of metallothionein-like proteins of the jussel *Mytilus edulis* at a basal level by chromatographic separations coupled to quadrupole and double-focusing magnetic sector ICPMS. *Anal. Chem.*, 72 (2000) 5874-5880.

1488 Grace, J.E., Jr., van Eden, M.E. and Aust, S.D.: Production of recombinant human apoferritin heteromers. *Arch. Biochem. Biophys.*, 384 (2000) 116-122.

1489 Hara, K. and Takahashi, H.: (Basic evaluation of hemoglobin A_{1c} measurement by ADAMS-A1c HA-8160). *Igaku to Yakugaku*, 43 (2000) 547-554; *C.A.*, 134 (2001) 68225t.

1490 Molon, A., Di Muro, P., Bubacco, L., Vasilyev, V., Salvato, B., Beltramini, M., Conze, W., Hellmann, N. and Decker, H.: Molecular heterogeneity of the hemocyanin isolated from the king crab *Paralighodes camtschaticae*. *Eur. J. Biochem.*, 267 (2000) 7046-7057.

1491 Seo, D., Tomioka, A., Kusumoto, N., Kamo, M., Enami, I. and Sakurai, H.: Purification of ferredoxins and their reaction with purified reaction center complex from the green sulfur bacterium *Chlorobium tepidum*. *Biochim. Biophys. Acta*, 1503 (2001) 377-384.

1492 Shokrani, M., Terrell, F., Turner, E.A. and Del Pilar Aguinaga, M.: Chromatographic measurements of hemoglobin A_2 in blood samples that contain sickle hemoglobin. *Ann. Clin. Lab. Sci.*, 30 (2000) 191-194; *C.A.*, 133 (2000) 249146t.

See also 1401, 1434, 1513, 1541, 1545, 2063.

19i. Proteins of glands, gland products, various zymogens (incl. milk proteins)

1493 Castagnola, M., Congiu, D., Denotti, G., di Nunzio, A., Fadda, M.B., Melis, S., Messana, I., Misiti, F., Murtas, R., Olianas, A., Piras, V., Pittau, A. and Puddu, G.: Determination of the human salivary peptides histatins 1, 3, 5 and statherin by high-performance liquid chromatography and by diode-array detection. *J. Chromatogr. B*, 751 (2001) 153-160.

1494 Cavagnari, B.M., Sterin-Speziale, N., Affanni, J.M., Knudsen, J. and Santome, J.A.: Acyl-CoA-binding protein in the armadillo Harderian gland: its primary structure and possible role in lipid secretion. *Biochim. Biophys. Acta,* 1545 (2001) 314-325.

1495 Couriol, C., Le Quellec, S., Guihard, L., Molle, D., Chaufer, B. and Prigent, Y.: Separation of acid whey proteins on the preparative scale by hyperdiffusive anion exchange chromatography. *Chromatographia,* 52 (2000) 465-472.

1496 Horn, F., Coutinho dos Santos, R. and Terminoni, C.: Recombi-

19k. Proteins of neoplastic tissue and transformed cells

1506 Lesniak, W., Jezierska, A. and Kuznicki, J.: Upstream stimulatory factor is involved in the regulation of the human calcyclin (S100A6) gene. *Biochim. Biophys. Acta,* 1517 (2000) 73-81.

1507 Schirle, M., Keilholz, W., Weber, B., Gouttefangeas, C., Dumrese, T., Becker, H.D., Stevanovic, S. and Rammensee, H.-G.: Identification of tumor-associated MHC class I ligands by a novel T cell-independent approach. *Eur. J. Immunol.,* 30

amino acid oxidase. *Biochem. Biophys. Res. Commun.,* 280 (2001) 620-624.

1501 Verdoliva, A., Basile, G. and Fassina, G.: Affinity purification of immunoglobulins from chicken egg yolk using a new synthetic ligand. *J. Chromatogr. B,* 749 (2000) 233-242.

See also 1342.

19j. Proteins of brain, cerebrospinal fluid and eye

1502 Boelens, W.C., Croes, Y. and de Jong, W.W.: Interaction between αB-crystallin and the human 20S proteasomal subunit C8/α7. *Biochim. Biophys. Acta,* 1544 (2001) 311-319.

1503 Fujii, N., Takemoto, L.J., Matsumoto, S., Hiroki, K., Boyle, D. and Akaboshi, M.: Comparison of D-aspartic acid contents in αA-crystallin from normal and age-matched cataractous human lenses. *Biochem. Biophys. Res. Commun.,* 278 (2000) 408-413.

1504 Takaki, Y., Iwata, N., Tsubuki, S., Taniguchi, S., Toyoshima, S., Lu, B., Gerard, N.P., Gerard, C., Lee, H.-J., Shirotani, K. and Saido, T.C.: Biochemical identification of the neutral endopeptidase family member responsible for the catabolism of amyloid β peptide in the brain. *J. Biochem. (Tokyo),* 128 (2000) 897-902.

1505 Theodorsson, E. and Rugarn, O.: Radioimmunoassay for rat galanin: immunochemical and chromatographic characterization of immunoreactivity in tissue extracts. *Scand. J. Clin. Lab. Invest.,* 60 (2000) 411-418; *C.A.,* 134 (2001) 13408z.

1512 Wilk, S., Chen, W.-L. and Magnusson, R.P.: Properties of the nuclear proteasome activator PA28γ (REGγ). *Arch. Biochem. Biophys.,* 383 (2000) 265-271.

1513 Xhang, L.N., Yang, X. and Hua, Z.C.: Expression and purification of recombinant human annexin. *Prepar. Biochem. Biotechnol.,* 30 (2000) 305-312.

See also 1422, 1469, 1494, 1497.

19m. Urinary proteins

1514 Davis, M.T., Spahr, C.S., McGinley, M.D., Robinson, J.H., Bures, E.J., Beierle, J., Mort, J., Yu, W., Luethy, R. and Patterson, S.D.: Towards defining the urinary proteome using liquid chromatography-tandem mass spectrometry II. Limitations of complex mixture analyses. *Proteomics (Weinheim),* 1 (2001) 108-117.

1515 Iwata, S.: Recommended method of the Japanese Association of Medical Technologists for the determination of protein in urine by HPLC. *Clin. Chim. Acta,* 303 (2001) 95-104.

1516 Spahr, C.S., Davis, M.T., McGinley, M.D., Robinson, J.H., Bures, E.J., Beierle, J., Mort, J., Courchesne, P.L., Chen, K., Wahl, R.C., Yu, W., Luethy, R. and Petterson, S.D.: Towards defining the urinary proteome using liquid chromatography-tandem mass spectrometry I. Profiling an unfractionated tryptic digest. *Proteomics (Weinheim),* 1 (2001) 93-107.

19n. Other proteins (incl. proteinous inhibitors of enzymic activity)

1517 Aboul-Enein, H.Y., Ali, I., Gubitz, G., Simons, C. and Nicholls, P.J.: HPLC enantiomeric resolution of novel aromatase inhibitors on cellulose- and amylose-based chiral stationary phases under reversed phase mode. *Chirality,* 12 (2000) 727-733; *C.A.,* 134 (2001) 21572p.

1518 Chen, T.-b., Xu, X.-h. and Rao, P.-f.: (Improved method for the separation and purification of IgY from egg by one step elution anion-exchange chromatography). *Fuzhou Daxue Xuebao, Ziran Kexueban,* 28, No. 2 (2000) 99-102; *C.A.,* 134 (2001) 69948t.

1519 Croguennec, T., Nau, F., Pezennec, S. and Brule, G.: Simple rapid procedure for preparation of large quantities of ovalbumin. *J. Agric. Food Chem.,* 48 (2000) 4883-4889.

1520 Gebruers, K., Debyser, W., Goesaert, H., Proost, P., van Damme, J. and Delcour, J.A.: *Triticum aestivum* L. endoxylanase inhibitor (TAXI) constists of two inhibitors, TAXI I and TAXI II, with different specificities. *Biochem. J.,* 353 (2001) 239-244.

1521 Härndahl, U., Kokke, B.P.A., Gustavsson, N., Linse, S., Berggren, K., Tjerneld, F., Boelens, W.C. and Sundby, C.: The chaperone-like activity of a small heat shock protein is lost after sulfoxidation of conserved methionines in a surface-exposed amphipathic α-helix. *Biochim. Biophys. Acta,* 1545 (2001) 227-237.

1522 Li, Y., Agrawal, A., Sakon, J. and Beitle, R.R.: Characterization of metal affinity of green fluorescent protein and its purification through salt promoted, immobilized metal affinity chromatography. *J. Chromatogr. A,* 909 (2001) 183-190.

1523 Nakazawa, M., Mizukoshi, H., Makino, T., Harada, K., Miyagi, A., Deguchi, Y., Osada, K., Watanuki, M., Nagaoka, M. and Aiyama, R.: (Assay method for α-amylase inhibitors in the *Psidium guajava* L tea). *Chromatography,* 21, No. 2 (2000) 155-156; *C.A.,* 133 (2000) 280711x.

1524 Salzet, M., Chopin, V., Baest, J.-L., Matias, I. and Malecha, J.: Theromin, a novel leech thrombin inhibitor. *J. Biol. Chem.,* 275 (2000) 30774-20780.

1525 Tran, S.T. and Shrake, A.: The folding of alpha-1-proteinase inhibitor: kinetic vs equilibrium control. *Arch. Biochem. Biophys.,* 385 (2001) 322-331.

1526 Uretschläger, A., Einhauer, A. and Jungbauer, A.: Continuous separation of green fluorescent protein by annular chromatography. *J. Chromatogr. A.,* 908 (2001) 243-250.

1527 Vaccher, C., Wallez, V., Carato, P., Vaccher, M.P. and Bonte, J.P.: Enantiomeric separation of precursors of ho-kinase inhibitors by HPLC on polysaccharide-based chiral stationary phase. *Chromatographia,* 52 (2000) 513-516.

See also 1167.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

See 1413.

20a. Oxidoreductases

1528 Chen, R. and Yang, H.: A highly specific monomeric isocitrate dehydrogenase from *Corynebacterium glutamicum. Arch. Biochem. Biophys.,* 383 (2000) 238-245.

1529 Cvrk, T. and Stroebel, H.W.: Role of LYS271 and LYS279 residues in the interaction of cytochrome P4501A1 with NADPH-cytochrome P450 reductase. *Arch. Biochem. Biophys.,* 385 (2001) 290-300.

1530 Dierks, A., Stams, K.R., Lim, H.-K., Cornelius, G., Zhang, H. and Ball, S.E.: A method for the simultaneous evaluation of the activities of seven major human drug-metabolizing cytochrome P450s using an *in vitro* cocktail of probe substrates and fast gradient liquid chromatography tandem mass spectrometry. *Drug Metab. Disp.,* 29 (2001) 23-29.

1531 Felicia, N.D., Rekha, G.K. and Murply, S.E.: Characterization of cytochrome p450 2A4 and 2A5-catalyzed 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) metabolism. *Arch. Biochem. Biophys.,* 384 (2000) 418-424.

1532 Galuszka, P., Frebort, I., Sebela, M., Sauer, P., Jacobsen, S. and Pec, P.: Cytokinin oxidase or dehydrogenase? Mechanism of cytokinin degradation in cereals. *Eur. J. Biochem.,* 268 (2001) 450-461.

1533 Grechkin, A.N., Mukhtarova, L.S. and Hamberg, M.: The lipoxygenase pathway in tulip (*Tulipa gesneriana*): detection of the ketol route. *Biochem. J.,* 352 (2000) 501-509.

1534 Hughes, R.K., West, S.I., Hornostaj, A.R., Lawson, D.M., Fairhurst, S.A., Sanchez, R.O., Hough, P., Robinson, B.H. and Casey, R.: Probing a novel potato lipoxygenase with dual positional specificity reveals primary determinants of substrate binding and requirements for surface hydrophobic loop and has implications for the role of lipoxygenases in tubers. *Biochem. J.,* 353 (2001) 345-355.

1535 Incharoensakdi, A., Matsuda, N., Hibino, T., Meng, Y.-L., Ishikawa, H., Hara, A., Funaguma, T., Takabe, T. and Takabe, T.: Overproduction of spinal betaine aldehyde dehydrogenase in *Escherichia coli*. Structural and functional properties of wild-type, mutants and *E. coli* enzymes. *Eur. J. Biochem.,* 267 (2000) 7015-7023.

1536 Luo, Y.-S., Wang, H.-J., Gopalan, K.V., Srivastava, D.K., Nicaud, J.-M. and Chardot, T.: Purification and characterization of the recombinant form of acyl CoA oxidase 3 from the yeast *Yarrowia lipolytica. Arch. Biochem. Biophys.,* 384 (2000) 1-8.

1537 McGinnity, D.F., Parker, A.J., Soars, M. and Riley, R.J.: Automated definition of the enzymology of drug oxidation by the major human lung metabolizing cytochrome P450s. *Drug Metab. Disp.,* 28 (2000) 1327-1334.

1538 Regal, K.A. and Nelson, S.D.: Orientation of caffeine within the active site of human cytochrome P450 1A2 based on NMR longitudinal (T_1) relaxation measurements. *Arch. Biochem. Biophys.,* 384 (2000) 47-58.

1539 Shin, K.-S. and Lee, Y.-J.: Purification and characterization of a new member of the laccase family from the white-rot basidiomycete *Coriolus hirsutus. Arch. Biochem. Biophys.,* 384 (2000) 109-115.

1540 Sorianello, E.M. and Mazzetti, M.B.: Simple and rapid method for the determination of coproporphyrinogen oxidase activity. *J. Biochem. Biophys. Methods,* 45 (2000) 75-86; *C.A.,* 133 (2000) 307012j.

1541 Stewart, J.M., Blakely, J.A. and Johnson, M.D.: The interaction of ferrocytochrome c with long-chain fatty acids and their CoA and carnitine esters. *Biochem. Cell Biol.,* 78 (2000) 675-681.

1542 Stock, J.E. and de Voss, J.J.: Expression, purification, and characterization of biol: a carbon-carbon bond cleaving cytochrome P450 involved in biotin biosynthesis in *Bacillus subtilis*. *Arch. Biochem. Biophys.*, 384 (2000) 3511-360.

1543 Stresser, D.M., Blanchard, A.P., Turner, S.D., Erve, J.C.L., Dandeneau, A.A., Miller, V.P. and Crespi, C.L.: Substrate-dependent modulation of CYP3A4 catalytic activity: analysis of 27 test compounds with four fluorometric substrates. *Drug Metab. Disp.*, 28 (2000) 1440-1448.

1544 Sutherland, M., Shankaranarayanan, P., Schewe, T. and Nigam, S.: Evidence for the presence of phospholipid hydroperoxide glutathione peroxidase in human platelets: implications for its involvement in the regulatory network of the 12-lipoxygenase pathway of arachidonic acid metabolism. *Biochem. J.*, 353 (2001) 911-100.

1545 Sutter, B., Bounds, P. and Koppenol, W.H.: The preparation of apo-Cu,Zn superoxide dismutase by ion-exchange chromatography on iminodiacetic acid-sepharose. *Protein Expression Purif.*, 19 (2000) 53-56; *C.A.*, 133 (2000) 173784b.

1546 Takatsuka, H., Sakurai, Y., Yoshioka, A., Kokubo, T., Usami, Y., Suzuki, M., Matsui, T., Titani, K., Yagi, H., Matsumoto, M. and Fujimura, Y.: Molecular characterization of L-amino acid oxidase from *Agkistrodon halys blomhoffii* with special reference to platelet aggregation. *Biochim. Biophys. Acta*, 1544 (2001) 267-277.

1547 Ten Brink, H.B., Schoemaker, H.E. and Wever, R.: Sulfoxidation mechanism of vanadium bromoperoxidase from *Ascophyllum nodosum*. Evidence for direct oxygen transfer catalysis. *Eur. J. Biochem.*, 268 (2001) 132-138.

1548 Venkatakrishnan, K., von Moltke, L.L., Court, M.H., Harmatz, J.S., Crespi, C.L. and Greenblatt, D.J.: Comparison between cytochrome P450 (CYP) content and relative activity approaches to scaling from cDNA-expressed CYPs to human liver microsomes ratios of accessory proteins as sources of discrepancies between the approaches. *Drug Metab. Disp.*, 28 (2000) 1493-1504.

1549 Yang, Y.-H., Miranda, C.L., Henderson, M.C., Wang-Buhler, J.-L. and Buhler, D.R.: Heterologous expression of CYP2K1 and identification of the expressed protein (BV-CYP2K1) as lauric acid (-1)-hydroxylase and aflatoxin B1 exo-epoxidase. *Drug Metab. Disp.*, 28 (2000) 1279-1283.

See also 1442, 1656.

20b. Transferases (excl. E.C. 2.7.-.-)

1550 Lim, I.K., Park, T.J. and Paik, W.K.: Phosphorylation of methylated-DNA-protein-cysteine S-methyltransferase at serine-204 significantly increases its resistance to proteolytic digestion. *Biochem. J.*, 352 (2000) 801-808.

1551 Shibata, K., Fukuwatari, T. and Sugimoto, E.: Reversed-phase high-performance liquid chromatography of nicotinic acid mononucleotide for measurement of quinolinate phosphoribosyltransferase. *J. Chromatogr. B*, 749 (2000) 281-285.

20c. Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)

1552 Dewaste, V., Pouillon, M.C., Shears, S., Takazawa, K. and Erneux, C.: Cloning and expression of a cDNA encoding human inositol 1,4,5-trisphosphate 3-kinase C. *Biochem. J.*, 352 (2000) 343-351.

1553 Finch, A., Davies, W., Carter, W.G. and Saklatvala, J.: Analysis of mitogen-activated protein kinase pathways used by interleukin 1 in tissues *in vivo*: activation of hepatic c-Jun N-terminal kinases 1 and 2, and mitogen-activated protein kinase kinases 4 and 7. *Biochem. J.*, 353 (2001) 275-281.

1554 Huang, K., Park, Y.-D., Cao, Z.-F. and Zhou, H.-M.: Reactivation and refolding of rabbit muscle creatine kinase denatured in 2,2,2-trifluoroethanol solutions. *Biochim. Biophys. Acta*, 1545 (2001) 305-313.

1555 Knoche, M., Mönnich, K., Schäfer, H.-J. and Kopperschläger, G.: Photoaffinity labeling and photoaffinity cross-linking of phosphofructokinase-1 from *Saccharomyces cerevisiae* by 8-azidoadeninenucleotides. *Arch. Biochem. Biophys.*, 385 (2001) 301-310.

1556 Millard, T.H., Cullen, P.J. and Banting, G.: Effects of elevated expression of inositol 1,4,5-trisphosphate 3-kinase B on Ca^{2+} homeostasis in HeLa cells. *Biochem. J.*, 352 (2000) 709-715.

1557 Turner, W.L. and Plaxton, W.C.: Purification and characterization of cytosolic pyruvate kinase from banana fruit. *Biochem. J.*, 352 (2000) 875-882.

1558 Vitale, G., Bernardi, L., Napolitani, G., Mock, M. and Montecucco, C.: Susceptibility of mitogen-activated protein kinase kinase family members to proteolysis by anthrax lethal factor. *Biochem. J.*, 352 (2000) 739-745.

1559 Wugsintaweekul, J., Herz, S., Hecht, S., Eisenreich, W., Feicht, R., Rohdich, F., Bacher, A. and Zenk, M.H.: Phosphorylation of 1-deoxy-D-xylulose by D-xylulokinase of *Escherichia coli*. *Eur. J. Biochem.*, 268 (2001) 310-316.

20d. Hydrolases, acting on ester bonds (E.C. 3.1.-.-)

1560 Billecke, S., Draganov, D., Counsell, R., Stetson, P., Watson, C., Hso, C. and la Du, B.N.: Human serum paraoxonase (PON1) isozymes Q and R hydrolyze lactones and cyclic carbonate esters. *Drug Metab. Disp.*, 28 (2000) 1335-1342.

1561 Gao, R., Kini, R.M., Li, G., Luo, R., Selvanayagam, Z.E. and Gopalakrishnakone, P.: Purification and properties of three new phospholipase A_2 isoenzymes from *Micropechis ikaheka* venom. *Biochim. Biophys. Acta*, 1545 (2001) 30-40.

1562 Horn, J.M., Lehman, J.A., Alter, G., Horwitz, J. and Gomez-Cambronero, J.: Presence of a phospholipase D (PLD) distinct from PLD1 or PLD2 in human neutrophils: immunobiochemical characterization and initial purification. *Biochim. Biophys. Acta*, 1530 (2001) 97-110.

1563 Janckila, A.J., Takahashi, K., Sun, S.Z. and Yam, L.T.: Tartrate-resistant acid phosphatase isoform 5b as serum marker for osteoclastic activity. *Clin. Chem. (Washington)*, 47 (2001) 74-80.

1564 Kemken, D., Mier, K., Katus, H.A., Richardt, G. and Kurz, T.: A HPLC-fluorescence detection method for determination of cardiac phospholipase D activity *in vitro*. *Anal. Biochem.*, 286 (2000) 277-281.

1565 Kerovuo, J., Rouvinen, J. and Hatzack, F.: Analysis of myo-inositol hexakisphosphate hydrolysis by Bacillus phytase: indication of a novel reaction mechanism. *Biochem. J.,* 352 (2000) 623-628.

1566 Le Petit-Thevenin, J., Verine, A., Nganga, A., Nobili, O., Lombardo, D. and Bruneau, N.: Impairment of bile salt-dependent lipase secretion in AR4-2J rat pancreatic cells induces its degradation by the proteasome. *Biochim. Biophys. Acta,* 1530 (2001) 184-198.

1567 Ogino, C., Negi, Y., Daido, H., Kanemasu, M., Kondo, A., Kuroda, S., Tanizawa, K., Shimizu, N. and Fukuda, H.: Identification of novel membrane-bound phospholipase D from *Streptoverticillium cinnamoneum*, possessing only hydrolytic activity. *Biochim. Biophys. Acta,* 1530 (2001) 23-31.

1568 Taes, Y.E., Louagie, H., Yvergneux, J.P., de Buyzere, M.L., de Puydt, H., Delanghe, J.R. and Lott, J.A.: Prolonged hyperlipasemia attributable to a novel type of macrolipase. *Clin. Chem. (Washington),* 46 (2000) 2008-2013.

1569 Yusupova, R.A., Gulevich, A.Yu. and Drygin, Yu.F.: (Isolation of an unlinking enzyme, which hydrolyzes the covalent bond between picornavirus RNA and VPg, from ascites carcinoma Krebs II cells). *Biokhimiya (Moscow),* 65 (2000) 1440-1449.

20e. Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)

1570 Barbieri, L., Valbonesi, P., Righi, F., Zuccheri, G., Monti, F., Gorini, P., Samori, B. and Stirpe, F.: Polynucleotide:adenosine glycosidase is the sole activity of ribosome-inactivating proteins on DNA. *J. Biochem. (Tokyo),* 128 (2000) 883-889.

1571 Burtseva, Yu.V., Sova, V.V., Pivkin, M.V. and Zvyagintseva, T.N.: (Carbohydrate metabolism enzymes of micelial fungi living in a marine environment. β-1,3-Glucanase from marine *Chaetomium indicum*). *Biokhimiya (Moscow),* 65 (2000) 1389-1399.

1572 Diaz Arribas, J.C., Herrero, A.G., Martin-Lomas, M., Canada, F.J., He, S. and Withers, S.G.: Differential mechanism-based labeling and unequivocal activity assignment of the two active sites of intestinal lactase/phlorizin hydrolase. *Eur. J. Biochem.,* 267 (2000) 6996-7005.

1573 Ketesztessy, Z., Brown, K., Dunn, M.A. and Hughes, M.A.: Identification of essential active-site residues in the cyanogenic β-glucosidase (linamarase) from cassava (*Manihot esculenta* Crantz) by site-directed mutagenesis. *Biochem. J.,* 353 (2001) 199-205.

1574 Parry, N.J., Beever, D.E., Owen, E., Bandenberghe, I., van Beeumen, J. and Bhat, M.K.: Biochemical characterization and mechanism of activation of a thermostable β-glucosidase purified from *Thermoascus aurantiacus. Biochem. J.,* 353 (2001) 117-127.

1575 Takesue, Y., Yokota, K., Oda, S.-i. and Takesue, S.: Comparison of sucrase-free isomaltase with sucrase-isomaltase purified from the house musk shrew *Suncus murinus. Biochim. Biophys. Acta,* 1544 (2001) 341-349.

1576 Totani, K., Yasutake, N., Ohi, H., Murata, T. and Usui, T.: Enzymatic synthesis of aliphatic β-lactosides as mimic units of glycosphingolipids by use of *Trichoderma reesei* cellulase. *Arch. Biochem. Biophys.,* 385 (2001) 70-77.

1577 Van Vaeck, C., Wera, S., van Dijck, P. and Thevelein, J.M.: Analysis and modification of trehalose 6-phosphate levels in the yeast *Saccharomyces cerevisiae* with the use of *Bacillus subtilis* phosphotrehalase. *Biochem. J.,* 353 (2001) 157-162.

See also 1027, 1439, 1444.

20f. Other hydrolases

1578 Al-Jassabi, S.: (Purification and characterization of cathepsin L from skeletal muscle of the lizard *Agama stellio stellio*). *Biokhimiya (Moscow),* 65 (2000) 1130-1134.

1579 Arnostova, H., Kucerova, Z., Tislerova, I., Trnka, T. and Ticha, M.: Affinity chromatography of porcine pepsin of different types of immobilized 3,5-diiodo-L-tyrosine. *J. Chromatogr. A,* 911 (2001) 211-216.

1580 Benrezzak, O., Grondin, G., Proulx, J., Rousseau, E., D'Orléans-Juste, P. and Beaudoin, A.R.: Characterization and immunohistochemical localization of nucleoside triphosphate diphosphohydrolase (NTPDase) in pig adrenal glands (presence of a non-sedimentable isoform). *Biochim. Biophys. Acta,* 1524 (2000) 94-101.

1581 Bose, S., Brooks, P., Mason, G.G.F. and Rivett, A.J.: γ-Interferon decreases the level of 26 S proteasomes and changes the pattern of phosphorylation. *Biochem. J.,* 353 (2001) 291-297.

1582 Grünberg, R., Dongall, I., Günther, R., Rall, K., Hofmann, H.-J. and Bordusa, F.: Peptide bond formation, mediated by substrate mimetics. Structure-guided optimization of trypsin for synthesis. *Eur. J. Biochem.,* 267 (2000) 7024-7030.

1583 Ikuzawa, M., Inokuchi, T., Kobayashi, K.-i. and Yasumasu, S.: Amphibian pepsinogens: purification and characterization of Xenopus pepsinogens, and molecular cloning of Xenopus and bullfrog. *J. Biochem. (Tokyo),* 129 (2001) 147-153.

1584 Jacobsen, W., Christians, U. and Benet, L.Z.: *In vitro* evaluation of the disposition of a novel cysteine protease inhibitor. *Drug Metab. Disp.,* 28 (2000) 1343-1351.

1585 Khan, M.T. and Joseph, S.K.: Characterization of membrane-associated proteasomes in WB rat liver epithelial cells. *Arch. Biochem. Biophys.,* 385 (2001) 99-107.

1586 Kiel, M.C. and Ganoza, M.C.: Functional interaction of an *Escherichia coli* ribosomal ATPase. *Eur. J. Biochem.,* 268 (2001) 278-286.

1587 Oneda, H. and Inouye, K.: Effects of dimethyl sulfoxide, temperature, and sodium chloride on the activity of human matrix metalloproteinase 7 (matrilysin). *J. Biochem. (Tokyo),* 128 (2000) 785-791.

1588 Seo, S.-b., Tan-Wilson, A. and Wilson, K.A.: Protease C2, a cysteine endopeptidase involved in the continuing mobilization of soybean β-conglycinin seed proteins. *Biochim. Biophys. Acta,* 1545 (2001) 192-206.

1589 Takao, K., Takai, S., Ishihara, T., Mita, S. and Miyazaki, M.: Isolation of chymase complexed with physiological inhibitor similar to secretory leukocyte protease inhibitor (SLPI) from hamster cheek pouch tissues. *Biochim. Biophys. Acta,* 1545 (2001) 146-152.

1590 Wilk, S., Chen, W.-E. and Magnusson, R.P.: Properties of the beta subunit of the proteasome activator PA28 (11S REG). *Arch. Biochem. Biophys.,* 384 (2000) 174-180.

See also 1123, 1499.

20g. Lyases

1591 Bertoldi, M. and Borri Voltattorni, C.: Reaction of dopa decarboxylase with L-aromatic amino acids under aerobic and anaerobic conditions. *Biochem. J.,* 352 (2000) 533-538.

1592 Hans, M., Buckel, W. and Bill, E.: The iron-sulfur clusters in 2-hydroxyglutaryl-CoA dehydratase from *Acidaminococcus fermentans*. Biochemical and spectroscopic investigations. *Eur. J. Biochem.,* 267 (2000) 7082-7093.

1593 Jensen, P.E., Reid, J.D. and Hunter, C.N.: Modification of cysteine residues in the ChlI and CHlH subunits of magnesium chelatase results in enzyme inactivation. *Biochem. J.,* 352 (2000) 435-441.

20h. Isomerases

1594 Takahashi, S., Kumagai, M., Shindo, S., Saito, K. and Kawamura, Y.: Renin inhibits N-acetyl-D-glucosamine 2-epimerase (renin-binding protein). *J. Biochem. (Tokyo),* 128 (2000) 951-956.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

1595 Aussenac, J., Chassagne, D., Claparols, C., Charpentier, M., Duteurtre, B., Feuillat, M. and Charpentier, C.: Purification method for the isolation of monophosphate nucleotides from Champagne wine and their identification by mass spectrometry. *J. Chromatogr. A,* 907 (2001) 155-164.

1596 Cherepanov, A., Yildirim, E. and de Vries, S.: Joining of short DNA oligonucleotides with base pair mismatches by T4 DNA ligase. *J. Biochem. (Tokyo),* 129 (2001) 61-68.

1597 Collins, A.: Comparison of different methods of measuring 8-oxoguanine as a marker of oxidative DNA damage. *Free Radical Res.,* 32 (2000) 333-341; *C.A.,* 134 (2001) 53451q.

1598 Di Pietro, M.C., Vannoni, D., Leoncini, R., Liso, G., Guerranti, R. and Marinello, E.: Determination of urinary methylated purine pattern by high-performance liquid chromatography. *J. Chromatogr. B,* 751 (2001) 87-92.

1599 Duez, P., Helson, M., Some, T.I., Dubois, J. and Hanocq, M.: Chromatographic determination of 8-oxo-7,8-dihydro-2'-deoxyguanosine in cellular DNA: a validation study. *Free Radical Res.,* 33 (2000) 243-260; *C.A.,* 134 (2001) 53284n.

1600 Ito, T., van Kuilenburg, A.B.P., Bootsma, A.H., Haasnoot, A.J., van Cruchten, A., Wada, Y. and van Gennip, A.H.: The application of HPLC/ESI tandem mass spectrometry on urine-soaked filter-paper strips for the screening of disorders of purine and pyrimidine metabolism. *J. Inherited Metab. Dis.,* 23 (2000) 434-437; *C.A.,* 133 (2000) 331609e.

1601 Jameson, P.E., Zhang, H. and Lewis, D.H.: Cytokinins: extraction, separation, and analysis. *Methods Mol. Biol. (Totowa),* 141(Plant Hormone Protocols) (2000) 101-121; *C.A.,* 133 (2000) 292966x - a review with 36 refs.

1602 Junghans, M., Kreuter, J. and Zimmer, A.: Phosphodiester and phosphorothioate oligonucleotide condensation and preparation of antisense nanoparticles. *Biochim. Biophys. Acta,* 1544 (2001) 177-188.

1603 Landry, J., Slama, J.T. and Sternglanz, R.: Role of NAD^+ in the deacetylase activity of the SIR2-like proteins. *Biochem. Biophys. Res. Commun.,* 278 (2000) 685-690.

1604 Lazarowski, E.R., Boucher, R.C. and Harden, T.K.: Constitutive release of ATP and evidence for major contribution of ecto-nucleotide pyrophosphatase and nucleoside diphosphokinase to extracellular nucleotide concentration. *J. Biol. Chem.,* 275 (2000) 31061-31068.

1605 Lengger, C., Schöch, G. and Topp, H.: A high-performance liquid chromatographic method for the determination of 8-oxo-7,8-dihydro-2'-deoxyguanosine in urine from man and rat. *Anal. Biochem.,* 287 (2000) 65-72.

1606 Liu, L.-p., Lu, Y.-q. and Liu, J.-f.: (Estimation of phosphoriboxyl pyrophosphate (PRPP) synthetase activity in BFU-Es and CFU-Es from mice bone marrows by ion-pair rHPLC). *Shengming Kexue Yanjiu,* 4 (2000) 124-129; *C.A.,* 133 (2000) 234245a.

1607 Moon, K.-Y. and Kim, Y.S.: Synthesis and characterization of oligonucleotides containing site-specific bulky N^2-aralkylated guanines and N^6-aralkylated adenines. *Arch. Pharmacal Res.,* 23, No. 2 (2000) 139-146; *C.A.,* 133 (2000) 177401x.

1608 Nakanishi, M., Iwata, A., Yatome, C. and Kitade, Y.: Purification and properties of recombinant *Plasmodium falciparum* S-adenosyl-L-homocysteine hydrolase. *J. Biochem. (Tokyo),* 129 (2001) 101-105.

1609 Petit, P.X., Gendron, M.-C., Schrantz, N., metivier, D., Kroemer, G., Maciorowska, Z., Sureau, F. and Koester, S.: Oxidation of pyridine nucleotides during Fas- and ceramide-induced apoptosis in Jurkat cells: correlation with changes in mitochondria, glutathione depletion, intracellular acidification and caspase 3 activation. *Biochem. J.,* 353 (2001) 357-367.

1610 Safranow, K., Machoy, Z. and Ciechanowski, K.: Analysis of purines in urinary calculi by high-performance liquid chromatography. *Anal. Biochem.,* 286 (2000) 224-230.

1611 Somsen, O.J.G., Heoben, M.A., Esgalhado, E., Snoep, J.L., Visser, D., van der Heijden, R.T.J.M., Heijnen, J.J. and Westerhoff, H.V.: Glucose and the ATP paradox in yeast. *Biochem. J.,* 352 (2000) 593-599.

1612 Van Lenthe, H., van Kuilenburg, A.B.P., Ito, T., Bootsma, A.H., van Cruchten, A., Wada, Y. and van Gennip, A.H.: Defects in purimidine degradation identified by HPLC-electrospray tandem mass spectrometry of urine specimens or urine-soaked filter paper strips. *Clin. Chem. (Washington),* 46 (2000) 1916-1922.

1613 Zhang, L., Tang, S. and Li, S.: (Determination of inosine in injections by high-performance liquid chromatography). *Zhongguo Shenghua Yaowu Zazhi,* 21, No. 4 (2000) 193-194; *C.A.,* 133 (2000) 340342q.

1614 Zhang, W., Zhang, P. and Yang, H.: (Determination of cytidine diphosphate choline by RP-HPLC). *Yaowu Fenxi Zazhi,* 20, No. 6 (2000) 394-396; *C.A.,* 134 (2001) 61637v.

1615 Zhang, Y.-w., Long, C. and Shi, S.-y.: (Determination of the adenosine triphosphate in myocardial tissue by ion-pair reversed-phase high performance liquid chromatography). *Sepu,* 18 (2000) 322-324; *C.A.,* 133 (2000) 174109x.

See also 985, 1032, 1286, 1931.

21c. Nucleic acids, DNA

1616 Stellwagen, N.C., Bossi, A., Gelfi, C. and Righetti, P.G.: DNA and buffers: are there any noninteracting, neutral pH buffers? *Anal. Biochem.*, 287 (2000) 167-175.

See also 1620.

21e. Structural studies on DNA and DNA mapping

1617 Kratzmeier, M., Albig, W., Hänecke, K. and Doenecke, D.: Rapid dephosphorylation of H1 histones after apoptosis induction. *J. Biol. Chem.*, 275 (2000) 30478-30486.

1618 Roberts, D.W., Churchwell, M.I., Beland, F.A., Fang, J.-L. and Doerge, D.R.: Quantitative analysis of etheno-2'-deoxycytidine DNA adducts using on-line immunoaffinity chromatography coupled with LC/ES-MS/MS detection. *Anal. Chem.*, 73 (2001) 303-309.

See also 1596.

21f. Complex mixtures of nucleic acids, their fragments and PCR products

1619 Devaney, J.M., Pettit, E.L., Kaler, S.G., Vallone, P.M., Butler, J.M. and Marino, M.A.: Genotyping of two mutations in the HFE gene using single-base extension and high-performance liquid chromatography. *Anal. Chem.*, 73 (2001) 620-624.

1620 Katz, E.D.: HPLC of DNA and PCR products. In: Rapley R. (Editor), *Nucleic Acid Protoc. Handb.*, Humana Press Inc., Totowa, 2000, pp. 105-113; *C.A.*, 134 (2001) 2163.

22. ALKALOIDS

1621 Ammon, S., von Richter, O., Hofmann, U., Thon, K.-P., Eichelbaum, M. and Mikus, G.: *In vitro* interaction of codeine and diclofenac. *Drug Metab. Disp.*, 28 (2000) 1149-1152.

1622 Antonilli, L., Suriano, C., Grassi, M.C. and Nencini, P.: Analysis of cocaethylene, bonzoylecgonine and cocaine in human urine by high-performance thin-layer chromatography with ultraviolet detection: a comparison with high-performance liquid chromatography. *J. Chromatogr. B*, 751 (2001) 19-27.

1623 Chen, Z.: (Determination of methylephedrine hydrochloride and chlorpheniramine maleate in Shangfengyou antiinfluenza virus solution by HPLC). *Yaowu Fenxi Zazhi*, 20 (2000) 349-350; *C.A.*, 134 (2001) 21574r.

1624 Cui, Y. and Lin, G.: Simultaneous analysis of clivorine and its four microsomal metabolites by high-performance liquid chro-

1627 Katagi, M., Nishikawa, M., Tatsuno, M., Miki, A. and Tsuchihashi, H.: Column-switching high-performance liquid chromatographyy-electropsray ionization mass spectrometry for identification of heroin metabolites in human urine. *J. Chromatogr. B*, 751 (2001) 177-185.

1628 Kudo, K., Iwaya, K., Yomota, C., Morris, S. and Saito, M.: Determination of enantiomeric purity of hyoscyamine from scopolia extract using HPLC-CD system without chiral separation. *Enantiomer*, 5 (2000) 369-375; *C.A.*, 134 (2001) 61662z.

1629 Li, S.-L., Lin, G., Chan, S.-W. and Li, P.: Determination of the major isosteroidal alkaloids in bulbs of Fritillaria by high-performance liquid chromatography coupled with evaporative light scattering detection. *J. Chromatogr. A*, 909 (2001) 207-214.

1630 Lin, G., Cui, Y.-Y. and Hawes, E.M.: Characterization of rat liver microsomal metabolites of clivorine, an hepatotoxic otonecine-type pyrrolizidine alkaloid. *Drug Metab. Disp.*, 28 (2000) 1475-1483.

1631 Lindahl, R., Wasterby, A. and Levin, J.-O.: Determination of morpholine in air by derivatisation with 1-naphtylisothiocyanate and HPLC analysis. *Analyst (Cambridge)*, 126 (2001) 152-154.

1632 Ma, C. and Cao, T.: (Determination of alkaloids in *Sophora tonkinensis* by HPLC). *Yaowu Fenxi Zazhi*, 20, No. 6 (2000) 408-410; *C.A.*, 134 (2001) 61638w.

1633 Meng, Z., Wang, J., Zhou, L., Wang, Q. and Zhu, D.: Beaded molecule imprinted polymer for stereo isomer separation. *Chromatography*, 21 (2000) 217-219; *C.A.*, 134 (2001) 65630k.

1634 Oertel, R., Richter, K., Ebert, U. and Kirch, W.: Determination of scopolamine in human serum and microdialysis samples by liquid chromatography-tandem mass spectrometry. *J. Chromatogr. B*, 750 (2001) 121-128.

1635 Wang, Y.-j., Yang, X.-j., Zhang, Y.-h. and Zhao, W.-g.: (Determination of berberine hydrochloride and puerarin in Tangwei capsules by RP-HPLC-gradient elution method). *Sepu*, 18 (2000) 465-476; *C.A.*, 133 (2000) 313728p.

1636 Ware, G.M., Price, G., Carter, L., Jr. and Eitenmiller, R.R.: Liquid chromatographic preparative method for isolating ergot alkaloids, using a particle-loaded membrane extracting disk. *J. Assoc. Off. Anal. Chem.*, 83 (2000) 1395-1399.

1637 Wu, J., Xie, W. and Pawliszyn, J.: Automated in-tube solid phase microextraction coupled with HPLC-ES-MS for the determination of catechins and caffeine in tea. *Analyst (Cambridge)*, 125 (2000) 2216-2222.

1638 Xie, X., Zheng, P., Zhu, W. and Zhao, G.: (Separation and determination of nicotine in total particulate matter of cigarette smoke by high performance liquid chromatography). *Fenxi Huaxue*, 28 (2000) 1085-1087; *C.A.*, 133 (2000) 241824s.

1662 Storm, T., Reemtsma, T. and Jekel, M.: Analysis of monomeric aromatic sulfonates in complex industrial wastewater matrices by high performance liquid chromatography - mass spectrometry. *Biol. Abwasserreinig.,* 11(Anwendung der LC-MS in der Wasseranalytik) (1999) 57-68; *C.A.,* 133 (2000) 271226r.

See also 1394, 1451, 1813, 1826, 1920.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

1663 Delpierre, G., Vanstapel, F., Stroobant, V. and van Schaftingen, E.: Conversion of a synthetic fructosamine into its 3-phospho derivative in human erythrocyte. *Biochem. J.,* 352 (2000) 835-839.

1664 Hooijschuur, E.W.J., Kientz, C.E. and Brinkman, U.A.T.: Determination of alkylphosphonic acid by microcolumn liquid chromatography with gradient elution coupled on-line with flame photometric detection. *J. Chromatogr. A,* 907 (2001) 165-172.

1665 Ostrowska, J., Skrzydlewska, E. and Figaszewski, Z.A.: Isolation and analysis of phospholipids. *Chem. Anal. (Warsaw),* 45 (2000) 613-629; *C.A.,* 134 (2001) 53197m - a review with 93 refs.

1666 Schwedt, G.: (HPLC of inositol phosphates). *CLB, Chem. Labor Biotech.,* 51, No. 7 (2000) 257-261; *C.A.,* 133 (2000) 307194v.

1667 Wind, M., Edler, M., Jakubowski, N., Linscheid, M., Wesch, H. and Lehmann, W.D.: Analysis of protein phosphorylation by capillary liquid chromatography coupled to element mass spectrometry with ^{31}P detection and to electrospray mass spectrometry. *Anal. Chem.,* 73 (2001) 29-35.

See also 1257, 1313, 1377, 1565, 1606, 1615, 1936.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

1668 Angeles Suner, M., Devesa, V., Rivas, I., Velez, D. and Montoro, R.: Speciation of cationic arsenic species in seafood by coupling liquid chromatography with hydride generation atomic fluorescence detection. *J. Anal. At. Spectrom.,* 15 (2000) 1501-1507; *C.A.,* 134 (2001) 55701v.

1669 Fan, H., Liu, J. and Xu, K.: (Quality control of iron dextran by high performance gel permeation chromatography). *Yaowu Fenxi Zazhi,* 20 (2000) 254-257; *C.A.,* 133 (2000) 325717x.

1670 Mandel, F., Helle, N., Kruse, R. and Wendt, J.: (Tributyl tin in mussels and fish). *GIT Labor-Fachz.,* 44 (2000) 732-735; *C.A.,* 133 (2000) 251395s.

See also 1059, 1399, 1947, 2068.

26c. Coordination compounds

1671 Lazareva, E.E., Alpeeva, I.S. and Brykina, G.S.: (HPLC study of adsorption of tetrakis(*p*-methoxyphenyl)porphyrin and its complex with Pd(II).) *Vestn. Mosk. Univ., Ser. 2: Khim.,* 41 (2000) 231-232; *C.A.,* 134 (2001) 23a.

1672 Lemr, K., Holcapek, M., Jandera, P. and Lycka, A.: Analysis of metal complex azo dyes by high-performance liquid chromatography/electrospray ionization mass spectrometry and multistage mass spectrometry. *Rapid Commun. Mass Spectrom.,* 14 (2000) 1881-1888; *C.A.,* 134 (2001) 18543z.

1673 Yang, Z.-J., Liang, F., Wang, C.-Y., Zhang, H.-S. and Cheng, J.-K.: (Reversed-phase HPLC determination of niobium, vanadium, rhodium and chromium as their (5-bromo-2-thiazolylazo)resorcinol chelates). *Fenxi Kexue Xuebao,* 16 (2000) 379-382; *C.A.,* 134 (2001) 38a.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

1674 Alyshakov, V.I., Alpeeva, I.S., Brykina, G.D. and Lazareva, E.E.: (Determination of vitamin E by HPLC with indirect spectrophotometric detection). *Vestn. Mosk. Univ., Ser. 2: Khim.,* 41 (2000) 233-235; *C.A.,* 134 (2001) 21544f.

1675 Arkhapchev, Y.P., Nozdrin, V.I. and Guzev, K.S.: (Pharmacokinetics study of retinoic acid isobutyl ester using HPLC). *Farmatsiya (Moscow),* 49 (2000) 39-40; *C.A.,* 133 (2000) 286325w.

1676 Chung, H.J., Kim, Y.-A., Kim, Y.J., Choi, Y.K., Hwang, Y.K. and Park, Y.S.: Purification and characterization of UDP-glucose:tetrahydrobiopterin glucosyltransferase from Synechococcus sp. PCC 7942. *Biochim. Biophys. Acta,* 1524 (2000) 183-188.

1677 Craft, N.E., Bulux, J., Valdez, C., Li, Y. and Solomons, N.W.: Retinol concentrations in capillary dried blood spots from healthy volunteers: method validation. *Am. J. Clin. Nutr.,* 72 (2000) 450-454; *C.A.,* 133 (2000) 234568h.

1678 Drotleff, A.M. and Ternes, W.: Determination of RS,E/Z-tocotrienols by HPLC. *J. Chromatogr. A,* 909 (2001) 215-223.

1679 Filimonov, V.N., Zamuraev, O.V., Balyatinskaya, L.N. and Kolosova, I.F.: Selection of a ternary eluant for the determination of fat-soluble vitamins in multivitamin pharmaceutical preparations by high-performance liquid chromatography. *J. Anal. Chem.,* 55 (2000) 657-662; *C.A.,* 133 (2000) 198781m.

1680 Gama, P., Casal, S., Oliveira, B. and Ferreira, M.A.: Development of an HPLC/diode-array/fluorimetric detector method for monitoring tocopherols and tocotrienols in edible oils. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3011-3022.

1681 Hattori, A., Fukushima, T., Yoshimura, H., Abe, K. and Imai, K.: Production of LLU-α following an oral administration of γ-tocotrienol or γ-tocopherol to rats. *Biol. Pharm. Bull.,* 23 (2000) 1395-1397.

1682 Klaczkow, G. and Anuszewska, E.L.: Simultaneous determination of vitamins A, D_3 and E in multiple pharmaceutical preparations by HPLC method. *Acta Pol. Pharm.,* 57, No. 3 (2000) 167-170; C.A., 133 (2000) 301281z.

1683 Koiwu-Tikkanen, T.J., Ollilainen, V. and Piironen, V.I.: Determination of phylloquinone and menaquinones in animal products with fluorescence detection after postcolumn reduction with metallic zinc. *J. Agric. Food Chem.,* 48 (2000) 6325-6331.

1684 Lampen, A., Meyer, S., Arnhold, T. and Nau, H.: Metabolism of vitamin A and its active metabolite all-trans-retinoic acid in small intestinal enterocytes. *J. Pharmacol. Exp. Ther.,* 295 (2000) 979-985.

1685 Li, H.-B., Chen, F. and Jiang, Y.: Determination of vitamin B_{12} in multivitamin tablets and fermentation medium by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A,* 891 (2000) 243-247.

1686 Luo, G.: (Determination of vitamin E in Weishenxin capsules by HPLC). *Yaowu Fenxi Zazhi,* 20, No. 6 (2000) 424-425; *C.A.*, 134 (2001) 61639x.

1687 Morita, H., Takahashi, Y., Noguchi, H. and Abe, I.: Enzymatic formation of unnatural aromatic polyketides by chalcone synthase. *Biochem. Biophys. Res. Commun.,* 279 (2000) 190-195.

1688 Ndaw, S., Bergaentzle, M., Aoude-Werner, D. and Hasselmann, C.: Extraction procedures for the liquid chromatographic determination of thiamin, riboflavin and vitamin B_6 in foodstuffs. *Food Chem.,* 71 (2000) 129-138; *C.A.*, 133 (2000) 295509m.

1689 Ötles, S.: Applications of HPLC and GC to vitamin and lipid analysis. *LC-GC Eur.,* 13 (2000) 905-908.

1690 Reddy, G.S., Rao, D.S., Siu-Caldera, M.-L., Astecker, N., Weiskopf, A., Vouros, P., Sasso, G.J., Manchand, P.S. and Uskokovic, M.R.: 1a,25-Dihydroxy-16-ene-23-yne-vitamin D_3 and 1α,25-dihydroxy-16-ene-23-yne-2-epi-vitamin D_3: analogs of 1α,25-dihydroxyvitamin D_3 that resist metabolism through the C-24 oxidation pathway are metabolized through the C-3 epimerization pathway. *Arch. Biochem. Biophys.,* 383 (2000) 197-205.

1691 Saari, J.C., Garwin, G.G., Haeseleer, F., Jang, G.-F. and Palczewski, K.: Phase partition and high-performance liquid chromatography assays of retinoid dehydrogenases. *Methods Enzymmol.,* 316(Vertebrate Phototransduction and the Visual Cycle, Pt.B) (2000) 359-371; *C.A.*, 133 (2000) 292694g - a review with 27 refs.

1692 Shahdeo, K., Anderson, F.P. and Karnes, H.T.: Post-column reaction detection of biotin in human plasma ultrafiltrate based on laser-induced fluorescence energy transfer in the far-red spectral region. *Biomed. Chromatogr.,* 15 (2000) 311-316; *C.A.*, 133 (2000) 293139s.

1693 Valls, F., Sancho, M.T., Fernandez-Muino, M.A. and Checa, M.A.: Determination of vitamin B_6 in cooked sausages. *J. Agric. Food Chem.,* 49 (2001) 38-41.

1694 Wang, Y., Liu, Y. and Wang, M.: (Study on quality of retinamidic acid). *Yaoxue Xeobao,* 35 (2000) 688-691; *C.A.*, 143 (2001) 61644v.

1695 Wei, L. and Wang, D.: (Determination of vitamin B_6 in lysozyme oral pellicles by HPLC). *Zhongguo Yiyuan Yaoxue Zazhi,* 20 (2000) 613-614; *C.A.*, 134 (2001) 46875c.

1696 Ye, L., Landen, W.O. and Eitenmiller, R.R.: Liquid chromatographic analysis of all-trans-retinyl palmitate β-carotene, and vitamin E in fortified foods and the extraction of encapsulated and nonencapsulated retinyl palmitate. *J. Agric. Food Chem.,* 48 (2000) 4003-4008.

See also 1551, 1657, 1797, 1799, 1801, 1802.

28. ANTIBIOTICS

1697 Bentley, L., Harden, J., Browne, D. and Hill, H.M.: One method for all species: two case studies. *Chromatographia,* 52(Supplement) (2000) S49-S52.

1698 Brozmanova, H., Grundmann, M., Safarcik, K. and Jegorov, A.: High-performance liquid chromatographic method for therapeutic drug monitoring of cyclosporine A and its two metabolites in renal transplant patients. *J. Chromatogr. B,* 749 (2000) 93-100.

1699 Chang, Y.L., Chou, M.H., Lin, M.F., Chen, Y.F., Chen, C.F., Cheng, F.C. and Tsai, T.H.: Simultaneous determination of unbound cefoperazone in rat blood and brain using microdialysis. *J. Pharm. Pharmacol.,* 52 (2000) 963-968; *C.A.*, 133 (2000) 329079p.

1700 Chen, G., Wang, G.-Y.-s., Li, X., Waters, B. and Davies, J.: Enhanced production of microbial metabolites in the presence of dimethyl sulfoxide. *J. Antibiot.,* 53 (2000) 1145-1153.

1701 Chen, Z., Zhen, J. and Zhang, T.: (Determination of cefpiramide in human serum and urine by HPLC). *Zhongguo Yoaxue Zazhi (Beijing),* 35 (2000) 765-767; *C.A.*, 134 (2001) 65769n.

1702 Daeseleire, E., de Ruyck, H. and van Renterghem, R.: Confirmatory assay for the simultaneous detection of penicillins and cephalosporins in milk using liquid chromatography/tandem mass spectrometry. *Rapid Commun. Mass Spectrom.,* 14 (2000) 1404-1409; *C.A.*, 133 (2000) 280726f.

1703 Ding, X. and Mou, S.: Ion chromatographic analysis of tetracyclines using polymeric column and acidic eluent. *J. Chromatogr. A,* 897 (2000) 205-214.

1704 Eilbert, F., Anke, H. and Sterner, O.: Neobulgarones A~F from cultures of *Neobulgaria pura,* new inhibitors of apperssorium formation of *Magnaporthe grisea.* *J. Antibiot.,* 53 (2000) 1123-1129.

1705 Eldem, R. and Arican-Cellat, N.: High-performance liquid chromatographic determination of amphotericin B in a liposomal pharmaceutical product and validation of the assay. *J. Chromatogr. Sci.,* 38 (2000) 338-344.

1706 Feng, J. and Gu, J.: (HPLC method for determination of norvancomycin in plasma). *Zhongguo Yiyuan Yaoxue Zazhi,* 20 (2000) 458-459; *C.A.*, 133 (2000) 290596c.

1707 Fu, C., Li, Z., Hong, Z. and Liu, S.: (Determination of levofloxacin hydrochloride in human plasma by RP-HPLC). *Yaowu Fenxi Zazhi,* 20 (2000) 373-375; *C.A.*, 134 (2001) 36647r.

1708 Fujie, A., Iwamoto, T., Muramatsu, H., Okudaira, T., Nitta, K., Nakanishi, T., Sakamoto, K., Hori, Y., Hino, M., Hashimoto, S. and Okuhara, M.: FR901469, a novel antifungal antibiotic from an unidentified fungus No. 11243. I. Taxonomy, fermentation, isolation, physico-chemical properties and biological properties. *J. Antibiot.,* 53 (2000) 912-919.

1709 Fujii, M., Bouno, M., Fujita, S., Yoshida, M., Watanabe, Y. and Matsumoto, M.: Preparation of griseofulvin for topical application using N-methyl-L-pyrrolidone. *Biol. Pharm. Bull.,* 23 (2000) 1341-1345.

1710 Furusawa, N.: Rapid liquid chromatographic determination of residual penicillin G in milk. *Fresenius J. Anal. Chem.,* 368 (2000) 624-626.

1711 Hosokawa, N., Naganawa, H., Hamada, M., Iinuma, H. and Takeuchi, T.: New triene-ansamycins, thiazinotrienomycins F and G and a diene-ansamycin, benzoxamycin. *J. Antibiot.,* 53 (2000) 886-894.

1712 Howells, L. and Sauer, M.J.: Multi-residue analysis of avermectins and moxidectin by ion-trap LC-MS. *Analyst (Cambridge),* 126 (2001) 155-160.

1713 Hu, J.-F., Wunderlich, D., Sattler, I., Härtl, A., Papastavrou, I., Grond, S., Grablej, S., Feng, X.-Z. and Thiericke, R.: New 1-O-acyl α-L-rhamnopyranosides and rhamnosylated lactones from Streptomyces sp., inhibitors of 3α-hydroxysteroid-dehydrogenase (3α-HSD). *J. Antibiot.*, 53 (2000) 944-953.

1714 Hu, Y.-I., Feng, Y.-q., Zhang, Q.-h. and Da, S.-l.: (Chromatographic characterization of fosfomycin-modified zirconia support). *Fenxi Kexue Xuebao,* 16 (2000) 274-277; *C.A.*, 133 (2000) 328969s.

1715 Hwang, E.-I., Yun, B.-S., Kim, Y.-K., Kwon, B.-M., Kim, H.-G., Lee, H.-B., Jeong, W.-J. and Kim, S.-U.: Phellinsin A, a novel chitin synthase inhibitor produced by Phellinus sp. PL3. *J. Antibiot.*, 53 (2000) 903-911.

1716 Igarashi, M., Hayashi, C., Homma, Y., Hattori, S., Kinoshita, N., Hamada, M. and Takeuchi, T.: Tubelactomicin A, a novel 16-membered lactone antibiotic, from Nocardia sp. I. Taxonomy, production, isolation and biological properties. *J. Antibiot.*, 53 (2000) 1096-1101.

1717 Iijima, M., Someno, T., Amemiya, M., Ishizuka, M., Naganawa, H. and Takeuchi, T.: Biosynthesis of IC202C, a new siderophore with immunosuppressive activity. *J. Antibiot.*, 53 (2000) 1411-1415.

1718 Ito, Y., Ikai, Y., Oka, H., Matsumoto, H., Miyazaki, Y., Takeba, K. and Nagase, H.: Application of ion-exchange cartridge clean-up in food analysis IV. Confirmatory assay of benzylpenicillin, phenoxymethylpenicillin, oxacillin, cloxacillin, nafcillin and dicloxacillin in bovine tissues by liquid chromatography-electrospray ionization tandem mass spectrometry. *J. Chromatogr. A,* 911 (2001) 217-223.

1719 Kakinuma, N., Iwai, H., Takahashi, S., Hamano, K., Yanaagisawa, T., Nagai, K., Tanaka, K., Suzuki, K., Kirakae, F., Kirikae, T. and Nakagawa, A.: Quinolactacins A, B and C: novel quinolone compounds from Penicillium sp. EPF-6 I. Taxonomy, production, isolation and biological properties. *J. Antibiot.*, 53 (2000) 1247-1251.

1720 Kanai, Y., Ishiyama, D., Senda, H., Iwatani, W., Takahashi, H., Konno, H., Tokumasu, S. and Kanazawa, S.: Novel human topoisomerase I inhibitors, topopyrones A, B, C and D I. Producing strain, fermentation, isolation, physico-chemical properties and biological activity. *J. Antibiot.*, 53 (2000) 863-872.

1721 Kanazaki, H., Yanagisawa, S. and Nitoda, T.: Biosynthetic intermediates of the tetradehydro cyclic dipeptide albonoursin produced by *Streptomyces albulus* KO-23. *J. Antibiot.*, 53 (2000) 1257-1264.

1722 Khedr, A. and Darwish, I.: A validated stability-indicating high performance liquid chromatographic assay of amoxicillin. *Bull. Pharm. Sci., Assiut Univ.*, 23 (2000) 11-21; *C.A.*, 133 (2000) 271767t.

1723 Kitahashi, T.: (Rapid determination of vancomycin by serum direct injection high-performance liquid chromatography). *Igaku to Seibutsugaku,* 141 (2000) 23-25; *C.A.*, 133 (2000) 171703a.

1724 Kitahashi, T.: (Significance on determination of vancomycin levels in human serum using HPLC as clinical examination). *Igaku to Seibutsugaku,* 141, No. 4 (2000) 133-135; *C.A.*, 133 (2000) 358832j.

1725 Kochansky, J.: Analysis of oxytetracycline in extender patties. *Apidologie,* 31 (2000) 517-524; *C.A.*, 133 (2000) 340339u.

1726 Lacramioara-Stefania, M. and Nacu, A.: Study of oral active penicillin esters by high-pressure liquid chromatography. *Bul. Inst. Politeh. Iasi, Sect. 2: Chim. Ing. Chim.,* 45 (1999) 5-10; *C.A.*, 133 (2000) 198779s.

1727 Lei, J., Mei, Z., Yu, J., Cai, H., Luo, S. and Zhang, Y.: (Determination of erythromycin ethylsuccinate and its reklated substances by ion suppression chromatography). *Yaoxue Xuebao,* 35 (2000) 514-517; *C.A.*, 133 (2000) 366501r.

1728 Mariappan, T.T., Singh, B. and Singh, S.: A validate reversed-phase (C_{18}) HPLC method for simultaneous determination of rifampicin, isoniazid and pyrazinamide in USP dissolution medium and simulated gastric fluid. *Pharm. Pharmacol. Commun.*, 6, No. 8 (2000) 345-349; *C.A.*, 133 (2000) 286587h.

1729 Miyazaki, T., Yomota, C. and Okada, S.: Hyaluronate depolymerization following thermal decomposition of oxytetracycline. *Chem. Pharm. Bull.*, 49 (2001) 118-122.

1730 Muller-Serieys, C., Bergogne-Berezin, E., Farinotti, R. and Massias, L.: Plasma pharmacokinetics of cefuroxime after IM injection of 750 mg with and without lidocaine. *Antibiotiques,* 1 (1999) 33-37; *C.A.*, 133 (2000) 261115q.

1731 Nishihara, Y., Tsujii, E., Takase, S., Tsurumi, Y., Kino, T., Hino, M., Yamashita, M. and Hashimoto, S.: FR191512, a novel anti-influenza agent isolated from a fungus strain No. 17415 I. Taxonomy, formation, isolation, physico-chemical properties and structure elucidation. *J. Antibiot.*, 53 (2000) 1333-1340.

1732 Nishino, I., Fujitomo, H. and Umeda, T.: Determination of a new oral cephalosporin, cefmatilen hydrochloride hydrate, and its seven metabolites in human and animal plasma and urine by coupled systems of ion-exchange and reversed-phase high-performance liquid chromatography. *J. Chromatogr. B,* 749 (2000) 101-110.

1733 Oka, H., Harada, K.-i., Suzuki, M. and Ito, Y.: Separation of spiramycin components using high-speed counter-current chromatography. *J. Chromatogr. A,* 903 (2000) 93-98.

1734 Otani, T., Yoshida, K.-i., Kubota, H., Kawai, S., Ito, S., Hori, H., Ishiyama, T. and Oki, T.: Novel triene-β-lactone antibiotics, oxazolomycin derivative and its isomer, produced by Streptomyces sp. KSM-2690. *J. Antibiot.*, 53 (2000) 1397-1400.

1735 Rees, L.W., Munn, A.D., Maddock, J. and Robinson, P.R.: The rapid determination of tobramycin in microlitre quantities of human tear fluid using LC-MS-MS. *Chromatographia,* 52(Supplement) (2000) S98-S1000.

1736 Renner, U.D., Thiede, C., Bornhäuser, M., Ehninger, G. and Thiede, H.-M.: Determination of mycophenolic acid and mycophenolate mofetil by high-performance liquid chromatography using postcolumn derivatization. *Anal. Chem.*, 73 (2001) 41-46.

1737 Sakuda, S., Ikeda, H., Nakamura, T., Kawachi, R., Kondo, T., Ono, M., Sakurada, M., Inagaki, H., Ito, R. and Nagawawa, H.: Blastaicidin A derivatives with highly specific inhibitory activity toward aflatoxin production in *Aspergillus parasiticus. J. Antibiot.*, 53 (2000) 1378-1384.

1738 Sasse, F., Steinmetz, H., Heil, J., Höfle, G. and Reichenbach, H.: Tubulysins, new cytostatic peptides from myxobacteria on microtubuli. Production, isolation, physico-chemical and biological properties. *J. Antibiot.*, 53 (2000) 879-885.

1739 Scanes, T., Hundt, A.F., Swart, K.J. and Hundt, H.K.L.: Simultaneous determination of cefotaxime and desacetylcefotaxime in human plasma and cerebrospinal fluid by high-performance liquid chromatography. *J. Chromatogr. B,* 750 (2001) 171-176.

1740 Stefanska, A.L., Fulston, M., Houge-Frydrych, C.S.V., Jones, J.J. and Warr, S.R.: A potent seryl tRNA synthetase inhibitor SB-217452 isolated from a Streptomyces sp. *J. Antibiot.*, 53 (2000) 1346-1353.

1741 Stout, S.J., Wickremesinhe, E., da Cunhga, A.R. and Khunachak, A.: Confirmation of moxidectin residues in cattle fat by liquid chromatography/mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 83 (2000) 1446-1450.

1742 Su, Y.: (Determination of clarithromycin in its suspension by RP-HPLC). *Guangdong Yaoxueyuan Xuebao*, 16 (2000) 201-202; *C.A.*, 133 (2000) 301280y.

1743 Tipre, D.N. and Kasture, A.V.: Simultaneous estimation of ciprofloxacin and tinidazole in tablet dosage form using spectrophotometric and HPLC methods. *Indian Drugs*, 37 (2000) 148-154; *C.A.*, 133 (2000) 256904s.

1744 Trucksess, M.W. and Tang, Y.: Solid phase extraction method for patulin in apple juice and unfiltered apple juice. *Methods Mol. Biol. (Totowa)*, 157(Mycotoxin Protocols) (2001) 205-213; *C.A.*, 134 (2001) 70468t.

1745 Tsai, T.-H., Kao, H.-Y. and Chen, C.-F.: Measurement and pharmacokinetic analysis of unbound ceftazidime in rat blood using microdialysis and microbore liquid chromatography. *J. Chromatogr. B*, 750 (2001) 93-98.

1746 Tsai, T.-H., Liu, S.-C., Ho, L.-K. and Chen, C.-F.: Determination of unbound ampicillin in rat blood by microdialysis and liquid chromatography. *Anal. Lett.*, 33 (2000) 3213-3224.

1747 Ueno, R.: Rapid determination of residual antibiotics and antibacterial agents in cultured fish by HPLC. *ITE Lett. Batteries, New Technol. Med.*, 1, No. 2 (2000) 37-42; *C.A.*, 133 (2000) 334176x - a review with 25 refs.

1748 Van Wensveen, P.: New paradigm for HPLC method development. *Chem. Aust.*, 67, No. 4 (2000) 17-19; *C.A.*, 134 (2001) 9423h.

1749 Wang, J.: (Determination of impurities in cefetamet pivoxil hydrochloride). *Zhongguo Yiyao Gongye Zazhi*, 31, No. 7 (2000) 308-310; *C.A.*, 133 (2000) 340363x.

1750 Wu, X., Li, Z. and Hu, J.: (Determination of chloramphenicol and hydrocortisone in ear drops by HPLC). *Zhongguo Yiyuan Yaoxue Zazhi*, 20 (2000) 476-477; *C.A.*, 133 (2000) 301279e.

1751 Xie, Y., Van de Sandt, E., de Weerd, T. and Wang, N.-H.L.: Purification of adipoyl-7-amino-3-deacetoxycephalosporanic acid from fermentation broth using stepwise elution with a synergistically adsorbed modulator. *J. Chromatogr. A*, 908 (2001) 273-291.

1752 Yan, X., Gu, D., Hu, X., Cao, G., He, X. and Song, Y.: (Study on enantiomeric separation of levofloxacin hydrochloride injection by HPLC). *Huaxi Yaoxue Zazhi*, 15, No. 4 (2000) 287-288; *C.A.*, 133 (2000) 355319m.

1753 Yanai, M., Suzuki, M., Kawamura, K., Oshida, N., Hiramoto, S., Yasuda, O., Shingai, A. and Suguro, T.: Synthesis of N-4909 analogs part I. A stimulant of apolipoprotein E secretion in human hepatoma G2 cells. *J. Antibiot.*, 53 (2000) 1385-1396.

1754 Yang, Z., Chen, J., Chen, W. and Zeng, H.: (Simultaneous determination of ampicillin and cloxacillin for injection by RP-HPLC). *Zhongguo Yiyuan Yaoxue Zazhi*, 20 (2000) 529-531; *C.A.*, 133 (2000) 301285d.

1755 Zajac, M., Musial, W. and Pawlowski, L.: Stability of cefotaxime sodium in solid state. *Pharmazie*, 55 (2000) 917-918.

1756 Zarapakar, S.S. and Rane, S.H.: Reverse phase high-performance liquid chromatographic determination of Amoxycillin and Ambroxol HCl in tablets. *Indian Drugs*, 37 (2000) 246-250; *C.A.*, 133 (2000) 256881g.

1757 Zhang, B., Yan, X., Liu, X., Zhang, S. and Wu, H.: (Determination of pingyangmycin in magnetic-microspheres and rat liver by HPLC). *Yaowu Fenxi Zazhi*, 20 (2000) 149-151; *C.A.*, 133 (2000) 313715g.

1758 Zhang, J.-j., Gao, Y. and Tu, X.-d.: (Simultaneous determination of ampicillin and cloxacillin in compound capsules by HPLC). *Zhongguo Yaoke Daxue Xuebao*, 31 (2000) 358-360; *C.A.*, 134 (2001) 46923s.

See also 1132, 1800, 1975.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

1759 Chatani, Y., Yashima, S., Kitano, R., Ohfuji, M. and Komatsu, M.: (Studies on simultaneous determination of pesticide residues in agricultural products by GC/MS (V) - citrus fruits, banana). *Kyoto-fu Hoken Kankyo Kenkyusho Nenpo*, 44 (1999) 14-19; *C.A.*, 133 (2000) 176352v.

1760 Eisert, R., Jackson, S. and Krotzky, A.: Application of on-site solid-phase microextraction in aquatic dissipation studies of profoxydim in rice. *J. Chromatogr. A*, 909 (2001) 29-36.

1761 Galeano-Diaz, T., Guiberteau-Cabanillas, A., Mora-Diez, N., Parrilla-Vazquez, P. and Salinas-Lopez, F.: Rapid and sensitive determination of 4-nitrophenol, 3-methyl-4-nitrophenol, 4,6-dinitro-o-cresol, parathion-methyl, fenitrothion, and parathion-ethyl by liquid chromatography with electrochemical detection. *J. Agric. Food Chem.*, 48 (2000) 4508-4513.

1762 Kaihara, A., Yoshii, K., Tsumura, Y., Nakamura, Y., Ishimitsu, S. and Tonogai, Y.: Multiresidue analysis of pesticides in fresh fruits and vegetables by supercritical fluid extraction and HPLC. *J. Health Sci.*, 46 (2000) 336-342; *C.A.*, 133 (2000) 251418b.

1763 Lopez-Mesas, M., Crespi, M., Brach, J. and Mullender, J.P.: Clean-up of a pesticide-lanolin mixture by gel permeatin chromatography. *J. Chromatogr. Sci.*, 38 (2000) 551-555.

1764 Stalikas, C.D. and Konidari, C.N.: Analytical methods to determine phosphonic and amino acid group-containing pesticides. *J. Chromatogr. A*, 907 (2001) 1-19 - a review with 104 refs.

See also 1127, 1134, 2038.

29b. Chlorinated insecticides

1765 Hogenboom, A.C., Niessen, W.M.A. and Brinkman, U.A.Th.: Characterization of photodegradation products of alachlor in water by on-line solid-phase extraction liquid chromatography combined with tandem mass spectrometry and orthogonal-acceleration time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.*, 14 (2000) 1914-1924; *C.A.*, 143 (2001) 8994h.

1766 Sancho, J.V., Pozo, O.J. and Hernandez, F.: Direct determination of chlorpyrifos and its main metabolite 3,5,6-trichloro-2-pyridinol in human serum and urine by coupled-column liquid chromatography/electrospray-tandem mass spectrometry. *Rapid Commun. Mass Spectrom.,* 14 (2000) 1485-1490; *C.A.,* 133 (2000) 277247v.

1767 Vera-Avila, L.E., Vazuez-Lira, J.C., Covarrubias-Herrera, R., Garcia-Camacho, M.P. and Nunez-Gaytan, A.M.: Stability of parathion-methyl, methiocarb, DDT and 2,4-D in solid-phase extraction precolumns packed with a polymeric reversed phase. *Int. J. Environ. Anal. Chem.,* 77 (2000) 255-268; *C.A.,* 134 (2001) 1535a.

See also 1025.

29c. Phosphorus insecticides

1768 Ismail, N., Vairamani, M. and Rao, R.N.: Determination of *cis* and *trans* isomers of monocrotophos in technical products by reversed-phase column liquid chromatography. *J. Chromatogr. A,* 903 (2000) 255-260.

1769 Mori, H., Nagase, H., Nakamura, M. and Yamazaki, F.: (Quantitative analysis of fenitrothion and its metabolites in serum and urine by column-switching HPLC-DAD method and application to poisoning cases). *Jpn. J. Toxicol.,* 13, No. 2 (2000) 165-174; *C.A.,* 134 (2001) 1414k.

1770 Yokley, R.A., Shen, N. and Cheung, M.W.: Determination of two oxy-pyrimidine metabolites of diazinon in urine by gas chromatography/mass selective detection and liquid chromatography/electrospray ionization/mass spectrometry/mass spectrometry. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1229-1238.

See also 1761.

29d. Carbamates

1771 Gebreegzi, Y.Y., Foster, G.D. and Khan, S.U.: Simultaneous determination of carbaryl, malathion, fenitrothion, and diazinon residues in sesame seeds (*Sesamum indicum* L.) *J. Agric. Food Chem.,* 48 (2000) 5165-5168.

1772 Hagiwara, T., Yasuno, T., Haneishi, N. and Saito, K.: (Analysis of N-methyl carbamate pesticide residues in natural food additives by high-performance liquid chromatography). *Tokyo-toritsu Eisei Kenkyusho Kenkyu Nenpo,* 50 (1999, Pub. 2000) 193-197; *C.A.,* 134 (2001) 28611j.

1773 She, Y., Gao, Y. and Wang, Y.: (Determination of fenoxycarb by HPLC). *Henan Huangong,* No. 9 (2000) 25-26; *C.A.,* 134 (2001) 67401k.

1774 Shimizu, T., Nakaya, M., Miura, H. and Takahashi, M.: (Studies on rapid simultaneous analytical methods for N-methyl-carbamate pesticides with GPC clean-up). *Aomori-ken Kankyo Hoken Senta Kenkyu Hokoku,* 10 (1999, Pub. 2000) 55-58; *C.A.,* 134 (2001) 55654g.

1775 Soriano, J.M., Jimenez, B., Molto, J.C. and Font, G.: Behaviour of graphitized carbon black in the extraction of polar non-ionic nitrogen-containing pesticides. A checking of hypotheses. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2829-2838.

1776 Toscano, I.A.S., Ribeiro, M.L., Rocha, J.C., Nunes, G.S. and Barcelo, D.: (Determination of carbaryl by ELISA (enzyme-linked immunosorbent assay) and HPLC with photodiode array detection). *Quim. Nova,* 23 (2000) 466-471; *C.A.,* 134 (2001) 9006z.

See also 1767.

29e. Herbicides

1777 Brausen, G. and Reupert, R.: (Analysis of acid herbicides in water). *LaborPraxis,* 24, No. 9 (2000) 70-78; *C.A.,* 133 (2000) 339784d.

1778 Curren, M.S.S. and King, J.W.: Solubility of triazine pesticides in pure and modified subcritical water. *Anal. Chem.,* 73 (2001) 740-745.

1779 Garrido Frenich, A., Gil Garcia, M.D., Arrebela, F.J., Martinez Vidal, J.L., Martinez Galerz, M. and Lopez, T.: Determination of parts per trillion levels of benzoylurea pesticides in groundwater by high-performance liquid chromatography-electrospray ionization mass spectrometry. *Chromatographia,* 52 (2000) 569-574.

1780 Gennari, M., Ferraris, L., Negre, M. and Cignetti, A.: Liquid chromatographic determination of triasulfuron in soil. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1076-1081.

1781 Hudecova, T., Tekel, J., Hatrik, S., Vaverkova, S. and Havranek, E.: (Determination of trifluralin residues in the plant drug of *Mentha piperita* L.) *Ceska Slov. Farm.,* 49, No. 5 (2000) 247-250; *C.A.,* 133 (2000) 301286e.

1782 Krieger, M.S., Wynn, J.L. and Yoder, R.N.: Extraction of cloransulam-methyl from soil with subcritical water and supercritical CO_2. *J. Chromatogr. A,* 897 (2000) 405-413.

1783 Lagana, A., Bacaloni, A., Pago, G. and Marino, A.: Development of solid extraction unit for determining multiple herbicides in maize. *Chromatographia,* 52 (2000) 552-558.

1784 Moody, R.P.: Automated In Vitro Dermal Absorption (AIVDA): predicting skin permeation of atrazine with finite and infinite (swimming/bathing) exposure models. *Toxicol. in Vitro,* 14 (2000) 467-474; *C.A.,* 133 (2000) 345635x.

1785 Okihashi, M., Akutsu, K., Obana, H. and Hori, S.: Determination of triazine herbicides in food with liquid chromatography mass spectrometry. *Analyst (Cambridge),* 125 (2000) 1966-1969.

29f. Fungicides

1786 Martel, A.-C. and Porthault, M.: Pesticide residues in raspberries and lettuce: extraction and comparison of three chromatographic methods: HPLC, HPTLC and GC. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3042-3058.

1787 Negeri, B.O., Foster, G.D. and Khan, S.U.: Determination of chlorothalonil residues in coffee. *Toxicol. Environ. Chem.,* 77 (2000) 41-47; *C.A.,* 133 (2000) 321130c.

29g. Other types of pesticides and various agrochemicals

1788 Choi, B.K., Hercules, D.M. and Gusev, A.I.: Effect of liquid chromatography separation of complex matrices on liquid chromatography-tandem mass spectrometry signal suppression. *J. Chromatogr. A,* 907 (2001) 337-342.

1789 Evans, C.S., Startin, J.R., Goodall, D.M. and Keely, B.J.: Im-

1800 Luetzhoft, H.-C.H., Vaes, W.-H.J., Freidig, A.P., Halling-Sorensen,

and evaluation of a new method for the determination of the carotenoid content in selected vegetables by HPLC and HPLC-MS-MS. *J. Chromatogr. Sci.,* 38 (2000) 441-449.

1798 Kohata, K., Yamashita, Y., Horie, H. and Yamauchi, Y.: (Investigation of the methods for converting percentage of chlorophyll to pheophytin in green tea leaves). *Chagyo Kenkyu Hokoku,* 88 (2000) 45-48; *C.A.,* 134 (2001) 39128h.

1799 Landrum, J.T. and Bone, R.A.: Lutein, zeaxanthin, and the macular pigment. *Arch. Biochem. Biophys.,* 385 (2001) 28-40 - a minireview.

terization of polystyrene and polyisoprene by normal-phase temperature gradient interaction chromatography. *J. Chromatogr. A,* 910 (2001) 51-60.

1810 Mes, E.P.C., Tijssen, R. and Kok, W.T.: Influence of the carrier composition on thermal field-flow fractionation for the characterisation of sub-micron polystyrene latex particles. *J. Chromatogr. A,* 907 (2001) 201-209.

1811 Nielen, M.W.F. and Buijtenhuijs, A.(F.A.): Coupling of liquid chromatographic polymer separations with time-of-flight mass spectrometry. *LC-GC Eur.,* 14 (2001) 82-88.

1812 Stella, B., Arpicco, S., Peracchia, M.T., Desmaele, D., Hoebeke, J., Renoir, M., D'Angelo, J., Cattel, L. and Couvreur, P.: Design of folic acid-conjugated nanoparticles for drug targeting. *J. Pharm. Sci.*, 89 (2000) 14522-1464.

1813 Vonk, E.C., Langeveld-Voss, B.M.W., van Dongen, J.L.J., Janssen, R.A.J., Claessens, H.A. and Cramers, C.A.: Separation and characterization of oligomers by reversed-phase high-performance liquid chromatography; a study on well-defined oligothiophenes. *J. Chromatogr. A*, 911 (2001) 13-26.

1814 Wang, J., Li, M. and He, G.: (Determination of cyclic oligomers in nylon 6 samples by HPLC). *Hecheng Xianwei Gonbye*, 23, No. 2 (2000) 50-55; *C.A.*, 134 (2001) 5318e.

See also 1112.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

1815 Aboul-Enein, H.Y.: High-performance liquid chromatographic enantioseparation of drugs containing multiple chiral centers on polysaccharide-type chiral stationary phases. *J. Chromatogr. A*, 906 (2001) 185-193 - a review with 30 refs.

1816 Ahrer, W., Scherwenk, E. and Buchberger, W.: Determination of drug residues in water by the combination of liquid chromatography or capillary electrophoresis with electrospray mass spectrometry. *J. Chromatogr. A*, 910 (2001) 69-78.

1817 Anton, K. and Siffrin, C.: Packed-column SFC in the pharmaceutical industry: cGMP aspects. *Analusis*, 27 (1999) 691-701; *C.A.*, 133 (2000) 329082j.

1818 Cai, Z., Sinhababu, A.K. and Harrelson, S.: Simultaneous quantitative cassette analysis of drugs and detection of their metabolites by high performance liquid chromatography/ion trap mass spectrometry. *Rapid Commun. Mass Spectrom.*, 14 (2000) 1637-1643; *C.A.*, 133 (2000) 301305k.

1819 Cervoort, R.J.M., Debets, A.J.J., Claessens, H.A., Cramers, C.A. and de Jong, G.J.: Optimisation and characterisation of silica-based reversed-phase liquid chromatographic systems for the analysis of basic pharmaceuticals. *J. Chromatogr. A*, 897 (2000) 1-22 - a review with 112 refs.

1820 Detroyer, A., Schoonjans, V., Questier, F., Vander Heyden, Y., Borosy, A.P., Guo, Q. and Massart, D.L.: Exploratory chemometric analysis of the classification of pharmaceutical substances based on chromatographic data. *J. Chromatogr. A*, 897 (2000) 23-36.

1821 Francotte, E.R.: Enantioselective chromatography as a powerful alternative for the preparation of drug enantiomers. *J. Chromatogr. A*, 906 (2001) 379-397 - a review with 132 refs.

1822 Kobayashi, A., Koyama, K., Maihara, A., Kadowaki, T. and Yoden, K.: (Economical and ecologically sound assay of drugs by HPLC-ECD method with recycling of mobile phase). *Iyakuhin Kenkyu*, 31 (2000) 668-673; *C.A.*, 134 (2001) 13049h.

1823 Lazou, K., de Geyter, T., de Reu, L., Zhao, Y. and Sandra, P.: Applicability of a benchtop LC-DAD-MSD for quality control in the pharmaceutical industry. *LC-GC Eur.*, 13 (2000) 340-356.

1824 Liu, Y.-q., Cai, Q.-s., Yu, Z.-w., Chang, J.-b., Jiang, S.-x. and Chen, L.-r.: (Direct optical resolution of acidic biphenyl drugs by high performance liquid chromatography on tris(3,5-dimethylphenylcarbamate) of cellulose). *Sepu*, 18 (2000) 287-290; *C.A.*, 133 (2000) 24763h.

1825 Lord, H. and Pawliszyn, J.: Microextraction of drugs. *J. Chromatogr. A*, 902 (2000) 17-63 - a review with 131 refs.

1826 Mascher, D.G., Mascher, H.J., Scherer, G. and Schmid, E.R.: High-performance liquid chromatographic-tandem mass spectrometric determination of 3-hydroxypropylmercapturic acid in human urine. *J. Chromatogr. B*, 750 (2001) 163-169.

1827 Needham, S.R.: Directions in discovery: Fast, efficient separations in drug discovery-LC-MS analysis using column switching and rapid gradients. *LC-GC*, 18 (2000) 1156-1161; *C.A.*, 134 (2001) 36643m.

1828 Owada, K., Sato, S., Horiike, A., Uchida, M., Sano, S. and Nagano, T.: (Problems shown by HPLC chromatograms at the quantitative test of drugs). *Shizuoka-ken Kankyo Eisei Kagaku Kenkyusho Hokoku*, 40 (1997) 39-43; *C.A.*, 133 (2000) 340331k.

See also 1060, 1070, 1105, 1122, 1136, 2038, 2048.

32b. Antirheumatics and antiinflammatory drugs

1829 Abdel-Hamid, M.E.: LC-MS analysis of selected sulfur-containing non-steroid anti-inflammatory agents: applications to pharmaceutical products. *J. Liq. Chromatogr. Relat. Technol.*, 23 (2000) 3095-3107.

1830 Ching, C.B., Artl, W., Lisso, M. and Wozny, G.: Kinetic and equilibrium study of the enantioseparation of fenoprofen in a batch setup. *Ind. Eng. Chem. Res.*, 39 (2000) 4365-4369; *C.A.*, 134 (2001) 21418t.

1831 Flood, K.G., Reynolds, E.R. and Snow, N.H.: Characterization of inclusion complexes of betamethasone-related steroids with cyclodextrins using high-performance liquid chromatography. *J. Chromatogr. A*, 903 (2000) 49-65.

1832 Halpin, R.A., Geer, L.A., Zhang, K.E., Marks, T.M., Dean, D.C., Jones, A.N., Melillo, D., Doss, G. and Vyas, K.P.: The absorption, distribution, metabolism and excretion of rofecoxib, a potent and selective cyclooxygenase-2 inhibitor, in rats and dogs. *Drug Metab. Disp.*, 28 (2000) 1244-1254.

1833 Hutzler, J.M., Frye, R.F. and Tracy, T.S.: Sensitive and specific high-performance liquid chromatographic assay for 4'-hydroxyflurbiprofen and flurbiprofen in human urine and plasma. *J. Chromatogr. B*, 749 (2000) 119-125.

1834 Li, J., Gao, Y., Gao, Y. and Li, X.: (Simultaneous determination of ibuprofen and pseudoephedrine in granules by HPLC). *Zhongguo Yaoxue Zazhi (Beijing)*, 35 (2000) 623-624; *C.A.*, 133 (2000) 301311j.

1835 Liu, J., Li, J., Li, D. and You, G.: (HPLC determination of tinidazole and salicylic acid in compound tinidazole liniments). *Yaowu Fenxi Zazhi*, 20 (2000) 329-331; *C.A.*, 134 (2001) 9453t.

1836 Liu, Q., Zhang, Z., Bo, H. and Guo, X.: (Determination of guaizaulene sodium sulfonate in marzulene-S by high performance liquid chromatography). *Fenxi Huaxue*, 28 (2000) 1319; *C.A.*, 134 (2001) 9444r.

1837 Sochor, J., Klimes, J., Sedlacek, J., Somolikova, B. and Slovencik, D.: HPLC analysis of tiaprofenic acid in the samples of whole blood using L-L and S-L extractions. *J. Liq. Chromatogr. Relat. Technol.*, 23 (2000) 3191-3201.

1838 Uray, G. and Kosjek, B.: Simultaneous enantioseparation of flobufen and its diastereomeric metabolites on ULMO, a pirkle type chiral stationary phase. *Enantiomer*, 5 (2000) 329-332; *C.A.*, 134 (2001) 76472m.

1839 Vermeulen, B. and Remon, J.P.: Validation of a high-performance liquid chromatographic method for the determination of ibuprofen enantiomers in plasma of broiler chickens. *J. Chromatogr. B*, 749 (2000) 243-251.

1840 Wang, Q.-F. and Mei, H-H.: (Determination of indomethacin in glucosamine enteric-coated tablets by HPLC). *Zhongguo Yiyao Gongye Zazhi*, 31 (2000) 362-363; *C.A.*, 133 (2000) 301313m.

1841 Xu, H. and Hu, Z.: (An ion-pair RP-HPLC determination of bendazac lysine in its eye drops). *Yaowu Fenxi Zazhi*, 20 (2000) 243-245; *C.A.*, 133 (2000) 183116z.

1842 Zhao, H., Mu, Z., Li, H. and Qiu, Z.: (Determination of the optical purity of ketoprofen by reversed-phase HPLC after derivatization with R-(+)-α-methylbenzylamine). *Xinan Shifan Daxue Xuebao, Ziran Kexueban*, 25 (2000) 160-163; *C.A.*, 134 (2001) 61650u.

1843 Zhou, Q., Yao, T. and Zeng, S.: (Chiral RP-HPLC assay for propafenone enantiomers in rat liver microsomal incubates and its application *in vitro* metabolism). *Yaoxue Xuebao*, 35 (2000) 370-373; *C.A.*, 133 (2000) 187536j.

See also 1621, 1931.

32c. Autonomic and cardiovascular drugs

1844 Aramaki, S., Mori, M., Nakata, M., Shinohara, T. and Koizumi, T.: Pharmacokinetics of propranolol and its metabolites in horses after intravenous or oral administration. *Biol. Pharm. Bull.*, 23 (2000) 1333-1340.

1845 Belal, F., Al-Malaq, H.A., Al-Majed, A.A. and Gadkariem, E.A.: A stability-indicating HPLC method for the determination of isoxsuprine in dosage forms. Application to kinetic studies of isoxsuprine. *J. Liq. Chromatogr. Relat. Technol.*, 23 (2000) 3175-3189.

1846 Catalano, T., Madsen, G. and Demarest, C.: Systematic approach to the development of HPLC test methods. *Am. Pharm. Rev.*, 3, No. 3 (2000) 62-71; *C.A.*, 134 (2001) 61608m.

1847 Chen, G.-L., Li, Y.-F. and Lee, F.-Y.: A stability indicating method for verapamil by high-performance liquid chromatography/diode array detector. *Chin. Pharm. J. (Taipei)*, 52 (2000) 113-122; *C.A.*, 134 (2001) 61645w.

1848 Gergov, M., Robson, J.N., Duchoslav, E. and Ojanpera, I.: Automated liquid chromatographic/tandem mass spectrometric method for screening β-blocking drugs in urine. *J. Mass Spectrom.*, 35 (2000) 912-918; *C.A.*, 133 (2000) 263481e.

1849 Hassib, S.T., El-Sherif, Z.A., El-Bagary, R.I. and Youssef, N.F.: Reversed-phase high performance liquid chromatographic and thin layer chromatographic methods for the simultaneous determination of benazepril hydrochloride and hydrochlorothiazide in Cibadrex tablets. *Anal. Lett.*, 33 (2000) 3225-3237.

1850 Huo, X.: (Determination of timolol maleate and related substances in eye drops by reversed phase ion-pair HPLC). *Yaowu Fenxi Zazhi*, 20 (2000) 181-184; *C.A.*, 133 (2000) 313739t.

1851 Iyer, R.A., Mitroka, J., Malhotra, B., Monacorsi, S., Jr., Waller, S.C., Rinehart, J.K., Roongta, V.A. and Kripalani, K.: Metabolism of [^{14}C]omapatrilat, a sulfhydryl-containing vasopeptidase inhibitor in humans. *Drug Metab. Disp.*, 29 (2001) 60-69.

1852 Jacobsen, W., Kuhn, B., Soldner, A., Kirchner, G., Sewing, K.-F., Kollman, P.A., Benet, L.Z. and Crhistians, U.: Lactonization is the critical first step in the disposition of the 3-hydroxy-3-methylglutaryl-CoA reductase inhibitor atorvastatin. *Drug Metab. Disp.*, 28 (2000) 1369-1378.

1853 Kramer, S. and Blaschke, G.: High-performance liquid chromatographic determination of the β_2-selective adrenergic agonist fenoterol in human plasma after fluorescence derivatization. *J. Chromatogr. B*, 751 (2001) 169-175.

1854 Lausecker, B., Hess, B., Fischer, G., Mueller, M. and Hopfgartner, G.: Simultaneous determination of bosentan and its three major metabolites in various biological matrices and species using narrow bore liquid chromatography with ion spray tandem mass spectrometric detection. *J. Chromatogr. B*, 749 (2000) 67-83.

1855 Li, X., Liu, W. and Zeng, S.: (Resolution of enantiomers of atenolol and its determination in rats hepatic microsome by RP-HPLC). *Yaowu Fenxi Zazhi*, 20 (2000) 219-223; *C.A.*, 133 (2000) 171696a.

1856 Li, Y., Zheng, Q. and Huang, Z.: (Determination of nitrendipine in tablets by HPLC). *Zhongguo Yiyuan Yaoxue Zazhi*, 20 (2000) 602-603; *C.A.*, 134 (2001) 9454u.

1857 Liu, X., Yu, X., Wang, E., Wang, G. and Qian, Z.: (Determination of nimodipine and its dehydrogenation metabolite in human liver microsome by HPLC and its metabolic kinetics). *Yaoxue Xuebao*, 35, No. 4 (2000) 257-260; *C.A.*, 133 (2000) 290600z.

1858 Påhlman, I., Kankaanranta, S. and Palmer, L.: Pharmacokinetics of tolterodine, a muscarinic receptor antagonist, in mouse, rat and dog. *Arzneim.-Forsch.*, 51 (2001) 134-144.

1859 Pereira, V.A., Auler, J.O., Jr., Carmona, M.J., Mateus, F.H., Lanchote, V.L., Breimer, D.D. and Santos, S.R.C.J.: A micromethod for quantitation of debrisoquine and 4-hydroxydebrisoquine in urine by liquid chromatography. *Braz. J. Med. Biol. Res.*, 33 (2000) 509-514; *C.A.*, 133 (2000) 190017w.

1860 Qin, Y., Zou, Y., Liang, M., Huang, Y., Yu, Q. and Wang, J.: (Determination of isosorbide-5-mononitrate in human plasma by HPLC). *Yaowu Fenxi Zazhi*, 20, No. 3 (2000) 165-167; *C.A.*, 133 (2000) 305225g.

1861 Qin, Y., Zou, Y., Liang, M., Wang, J., Huang, Y., Yu, Q. and Li, T.: (Determination of salbutamol in human plasma by reversed-phase HPLC with UV detection). *Zhongguo Yaoxue Zazhi (Beijing)*, 35 (2000) 475-477; *C.A.*, 133 (2000) 171692w.

1862 Santoro, M.I.R.M., Cho, H.S. and Kedor-Hackmann, E.R.M.: Enantiomeric separation and quantitative determination of atenolol in tablets by chiral high-performance liquid chromatography. *Drug Dev. Ind. Pharm.*, 26 (2000) 1107-1110; *C.A.*, 134 (2001) 21559q.

1863 Sasaki, H., Yamamura, K., Mukai, T., Nishida, K., Nakamura, J., Nakashima, M. and Ichikawa, M.: Pharmacokinetic prediction of the ocular absorption of an instilled drug with opthalmic viscous vehicle. *Biol. Pharm. Bull.*, 23 (2000) 1352-1356.

1864 Sheu, M.-T., Yeh, G.-C., Ke, W.-T. and Ho, H.-O.: Development of a high-performance liquid chromatographic method for bioequivalence study of flavoxate tablets. *J. Chromatogr. B,* 751 (2001) 79-86.

1865 Wang, S. and He, L.: (Determination of nicardipine in plasma of rabbit by HPLC and study on its pharmacokinetids). *Yaowu Fenxi Zazhi,* 20, No.3 (2000) 160-162; *C.A.,* 133 (2000) 305224f.

See also 1061, 1338, 1634, 1931, 1949.

32d. Central nervous system drugs

1866 Ahrens, B., Schwandt, H.-J. and Schütz, H.: Screening, identification and quantitation of benzodiazepines in serum by solid phase extraction on a cyanopropyl phase using high performance liquid chromatography and photodiode array detection. *Arzneim.-Forsch.,* 50 (2000) 1057-1062.

1867 Alvarez, C., Sanchez-Brunete, J.A., Torrado-Santiago, S., Cadorniga, R. and Torrado, J.J.: Enantisellective separation of thalidomide on an immobilized α_1-acid glycoprotein chiral stationary phase. *Chromatographia,* 52 (2000) 455-458.

1868 Barker, S.A., Littlefield-Chabaud, M.A. and David, C.: Distribution of the hallucinogens N,N-dimethyltryptamine and 5-methoxy-N,N-dimethyltryptamine in rat brain following intraperitoneal injection: application of a new solid-phase extraction LC-APcI-MS-MS-isotope dilution method. *J. Chromatogr. B,* 751 (2001) 37-47.

1869 Bereczki, A., Horvath, V. and Horvai, G.: Immunoassay-based determination of phenobarbital using size-exclusion chromatography. *J. Chromatogr. B,* 749 (2000) 215-223.

1870 Bolner, A., Tagliaro, F. and Lomeo, A.: Optimised determination of clobazam in human plasma with extraction and high-performance liquid chromatography analysis. *J. Chromatogr. B,* 750 (2001) 177-180.

1871 Brunnenberg, M. and Kovar, K.-A.: Stereospecific analysis of ecstasy-like N-ethyl-3,4-methylenedioxyamphetamine and its metabolites in humans. *J. Chromatogr. B,* 751 (2001) 9-18.

1872 Cox, D.S., Du, J., Scott, K.R., Gao, H. and Eddington, N.D.: Determination of the enaminone DM5, an anti-epileptic agent, in mouse plasma and brain tissue by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B,* 749 (2000) 191-196.

1873 Davies, P.C.: Thirteen years of quetiapine ("seroquel") bioanalysis. *Chromatographia,* 52(Supplement) (2000) S103.

1874 De Jong, G.J. and Koster, E.H.M.: Solid-phase micro-extraction in bioanalysis, exemplified by lidocaine determination. *Chromatographia,* 52(Supplement) (2000) S12-S13.

1875 Faucette, S.R., Hawke, R.L., Lecluyse, E.L., Shord, S.S., Yan, B., Laethem, R.M. and Lindley, C.M.: Validation of bupropion hydroxylation as a selective marker of human cytochrome P450 2B6 catalytic activity. *Drug Metab. Disp.,* 28 (2000) 1222-1230.

1876 Gil-Augustini, M., Carda-Broch, S., Garcia-Alvarez-Coque, M.C. and Esteve-Romero, J.: Use of micellar mobile phases for the chromatographic determination of cloracepate, diazepam, and diltiazem in pharmaceuticals. *J. Chromatogr. Sci.,* 38 (2000) 521-527.

1877 Heizmann, P., Gössl, R. and Herisse, R.: Application of LC-SIM-MS in drug assay development. *Chromatographia,* 52(Supplement) (2000) S46-S48.

1878 Hesse, L.M., Venkatakrishnan, K., Court, M.H., von Moltke, L.L., Duan, S.X., Shader, R.I. and Greenblatt, D.J.: CYP2B6 Mediates the *in vitro* hydroxylation of bupropion: potential drug interactions with other antidepressants. *Drug Metab. Disp.,* 28 (2000) 1176-1183.

1879 Kim, K.H., Kim, H.J., Kim, J.-H. and Shin, S.D.: Determination of terbutaline enantiomers in human urine by coupled achiral-chiral high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. B,* 751 (2001) 69-77.

1880 Kunicki, P.K.: Simple and sensitive high-performance liquid chromatographic method for the determination of 1,5-benzodiazepine clobazam and its active metabolite *N*-desmethylclobazam in human serum and urine with application to 1,4-benzodiazepines analysis. *J. Chromatogr. B,* 750 (2001) 41-49.

1881 Locker, K.L., Morrison, D. and Watt, A.P.: Quantitative determination of L-775,606, a selective 5-hydroxytryptamine 1D agonist, in rat plasma using automated sample preparation and detection by liquid chromatography-tandem mass spectrometry. *J. Chromatogr. B,* 750 (2001) 13-23.

1882 Malaviya, R., Chen, C.-L., Navara, C., Malaviya, R., Liu, X.-P., Keenan, M., Waurzyniak, B. and Uckun, F.M.: Treatment of allergic asthma by targeting Janus kinase 3-dependent leukotriene synthesis in mast cells with 4-(3',5'-dibromo-4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline (WHI-P97). *J. Pharmacol. Exp. Ther.,* 295 (2000) 912-926.

1883 Margolis, J.M., O'Donell, J.P., Mankowski, D.C., Ekins, S. and Obach, R.S.: (R)-, (S)-, and racemic fluoxetine N-demethylation by human cytochrome P450 enzymes. *Drug Metab. Disp.,* 28 (2000) 1187-1191.

1884 Marshall, D., Robinson, M., Hincks, P., Dumasia, M., Teale, P. and Houghton, E.: The application of *in vitro* models for production of metabolites: isolation and characterisation of hydroxysecobarbital. *Chromatographia,* 52(Supplement) (2000) S35-S38.

1885 Martin, V.: Determination of tree anabolic compounds in calf urine by liquid chromatography with photodiode-arrray detection. *Analyst (Cambridge),* 125 (2000) 2230-2235.

1886 Mittal, S., Alexander, K.S. and Dollimore, D.: A high-performance liquid chromatography assay for yohimbine-HCl analysis. *Drug Dev. Ind. Pharm.,* 26 (2000) 1059-1065; *C.A.,* 134 (2001) 9422g.

1887 Mushiroda, T., Douya, R., Takahara, E. and Nagata, O.: The involvement of flavin-containing monooxygenase but not CYP3A4 in metabolism of itopride hydrochloride, a gastroprokinetic agent: comparison with cisapride and mosapride citrate. *Drug Metab. Disp.,* 28 (2000) 1231-1237.

1888 Pallant, A., Mathian, B., Prost, L., Theodore, C. and Patricot, M.-C.: Determination of plasma methoxyamines. *Clin. Chem. Lab. Med.,* 38 (2000) 513-517; *C.A.,* 134 (2001) 53271f.

1889 Paw, B. and Misztal, G.: Rapid and simple method for the determination of zopiclone in plasma by HPLC. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2873-2882.

1890 Raggi, M.A., Casamenti, G., Mandrioli, R. and Volterra, V.: A sensitive high-performance liquid chromatographic method using electrochemical detection for the analysis of olanzapine and desmethylolanzapine in plasma of schizophrenic patients using a new solid-phase extraction procedure. *J. Chromatogr. B,* 750 (2001) 137-146.

1891 Ramu, K.: A high-performance liquid chromatography-tandem mass spectrometric method for the determination of pharmacokinetics of ganaxolone in rat, monkey, dog and human plasma. *J. Chromatogr. B,* 751 (2001) 49-59.

1892 Ramu, K., Lam, G.N. and Chien, B.: Development of a high-performance liquid chromatographic-tandem mass spectrometric method for the determination of pharmacokinetics of Co 102862 in mouse, rat, monkey and dog plasma. *J. Chromatogr. B,* 749 (2000) 1-15.

1893 Ramu, K., Lam, G.N. and Hughes, H.: *In vivo* metabolism and mass balance of 4-[4-fluorophenoxy]benzaldehyde semicarbazone in rats. *Drug Metab. Disp.,* 28 (2000) 1153-1161.

1894 Reist, M., Roy-De-Vos, M., Montseny, J.-P., Mayer, J.M., Carrupt, P.-A., Berger, Y. and Testa, B.: Very slow chiral inversion of clopidogrel in rats: a pharmacokinetic and mechanistic investigation. *Drug Metab. Disp.,* 28 (2000) 1405-1410.

1895 Rowlands, J.C., Wang, H., Hakkak, R., Ronis, M.J.J., Strobel, H.W. and Badger, T.M.: Chromic intragastric infusion of ethanol-containing diets induces CYP3A9 while decreasing CYP3A2 in male rats. *J. Pharmacol. Exp. Ther.,* 295 (2000) 747-752.

1896 Störmer, E., von Moltke, L.L., Shader, R.I. and Greenblatt, D.J.: Metabolism of the antidepressant mirtazapine *in vitro*: contribution of cytochrome P-450 1A2, 2D6, and 3A4. *Drug Metab. Disp.,* 28 (2000) 1164-1175.

1897 Suzuki, T., Shimizu, R., Suganuma, T., Nishino, J.-i., Tomono, K., Hanano, M. and Watanabe, J.: Pharmacokinetic/pharmacodynamic relationship of eptazocine, a narcotic-antagonist analgesic, in rats. *Biol. Pharm. Bull.,* 23 (2000) 1504-1510.

1898 Tan, G., Dai, Z., Xu, P., Lei, P. and Yuan, S.: (Determination of tetrandrine in human plasma by high-performance liquid chromatography). *Hunan Yike Daxue Xuebao,* 25 (2000) 409-410; *C.A.,* 133 (2000) 290598e.

1899 Unterhalt, B. and Wenning, C.: Zur Trennung von Oxetacain und seinen Metaboliten. *Pharmazie,* 56 (2001) 58-60.

1900 Wang, Z., Hu, C., Chen, Z., Yu, R. and Xie, X.: (Determination of carbamazepine in finger-prick blood by reversed-phase high-performance liquid chromatography). *Zhongguo Yiyuan Yaoxue Zazhi,* 20 (2000) 534-536; *C.A.,* 134 (2001) 36641j.

1901 Wilhelm, M., Battista, H.-J. and Obendorf, D.: Selective and sensitive assay for the determination of benzodiazepines by high-performance liquid chromatography with simultaneous ultraviolet and reductive electrochemical detection at the hanging mercury drop electrode. *J. Chromatogr. A,* 897 (2000) 215-225.

1902 Wilson, I.D.: Investigation of a range of stationary phases for the separation of model drugs by HPLC using superheated water as the mobile phase. *Chromatographia,* 52(Supplement) (2000) S28-S34.

1903 Yu, J., Wen, M. and Wu, G.: (Study on content uniformity of scopolamine hydrobromide in compound carbamazepine tablets). *Zhongguo Yiyuan Yaoxue Zazhi,* 20 (2000) 611-613; *C.A.,* 134 (2001) 9430h.

1904 Zhang, X., Liu, G., Zhao, X. and Zhang, R.: (Determination of naphazoline hydrochloride in Dibejing by HPLC). *Yaowu Fenxi Zazhi,* 20 (2000) 277-278; *C.A.,* 133 (2000) 183120w.

See also 1621, 1627, 1628, 1830, 1840, 1842, 1861, 1974, 1976, 1992, 1996.

32e. Chemotherapeutics (exc. cytostatics and antibiotics)

1905 Allen, G.D. and Brookes, S.T.: Determination of zanamavir, a viral neuramidase inhibitor, in human serum by LC-MS-MS. *Chromatographia,* 52(Supplement) (2000) S-108.

1906 Aoyagi, Y., Taguchi, N., Oishi, M. and Suzuki, S.: (Determination of thiabendazole in antimicrobial sponge brushes for kitchen use). *Tokyo-toritsu Eisei Kenkyusho Kenkyu Nenpo,* 50 (1999, Pub. 2000) 215-219; *C.A.,* 134 (2001) 52519f.

1907 Borner, K., Hartwig, H. and Lode, H.: Determination of garifloxacin in human serum and urine by HPLC. *Chromatographia,* 52 (2000) S105-S107.

1908 Chen, X.-F. and Tong, Z.-M.: (Determination of rifampin and isoniazid in compound capsules by RP-HPLC). *Zhongguo Kangshengsu Zazhi,* 25 (2000) 274-277; *C.A.,* 134 (2001) 46881b.

1909 Cheng, H.: (Determination of aminosalicylic acid and isoniazid in pasiniazide tablets by HPLC). *Yaowu Fenxi Zazhi,* 20 (2000) 186-188; *C.A.,* 133 (2000) 313740m.

1910 Dong, W., Sun, X., Ma, P., Shi, L. and Tang, W.: (Determination of chlorhexidine acetate in solutions and nasal cream drop by HPLC). *Zhongguo Yaoxue Zazhi (Beijing),* 35 (2000) 550-552; *C.A.,* 133 (2000) 301303h.

1911 Furusawa, N.: Simultaneous high-performance liquid chromatographic determination of sulfamonomethoxine and its hydroxy N^4-acetyl metabolites following centrifugal ultra-filtration in animal blood plasma. *Chromatographia,* 52 (2000) 653-656.

1912 Hyun, M.H., Han, S.C., Jin, J.S. and Lee, W.: Separation of the steroisomers of racemic fluoroquinolone antibacterial agents on a crown-ether-based chiral HPLC stationary phase. *Chromatographia,* 52 (2000) 473-476.

1913 Krabel, B.J., Dickson, D.A., Zimmermann, A.G. and Coleman, M.R.: Liquid chromatographic determination of nicarbazin in feeds. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1027-1038.

1914 Li, J., Kang, J.-x. and Wu, D.-n.: (Determination of 4 antiseptics and 6 sunscreen agents in cosmetic products by reversed-phase high performance liquid chromatography). *Sepu,* 18 (2000) 462-464; *C.A.,* 133 (2000) 339936e.

1915 Matabudul, D., Conway, B. and Lumbley, I.: A rapid method for the determination of lasalocid in animal tissues and eggs by high-performance liquid chromatography with fluorescence detection and confirmation by LC-MS-MS. *Analyst (Cambridge),* 125 (2000) 2196-2200.

1916 Milovanovic, L., Kapetanovic, V., Dugumovic, Z., Miljevic, D. and Aleksic, M.: Spectrophotometric and HPLC determination of fleroxacin in tablets. *Pharmazie,* 56 (2001) 150-151.

1917 Moeder, C.W., Whitener, M.A. and Sowa, J.R., Jr.: Quantitative stereochemical analysis of a reagent that exhibits asymmetric amplification, B-chlorodiisopinocampheylborane (Dip-Cl). *J. Am. Chem. Sci.,* 122 (2000) 7218-7225; *C.A.,* 133 (2000) 238036f.

1918 Moreno, M.A., Castro, D., Frutos, P., Ballesteros, M.P. and Lastres, J.L.: Liquid chromatographic determination of methylparaben and propylparaben in nortriptyline hydrochloride oil-water microemulsions. *Chromatographia,* 52 (2000) 589-592.

1919 Papanagiotou, E.P., Iossifidou, E.G., Psomas, I.E. and Photis, G.: Simultaneous HPLC determination of sulfadiazine and trimethoprim in cultured gilthead sea bream (*Sparus aurata,* L.) tissues. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 2839-2849.

1920 Rao, T.N., Sarada, B.V., Tryk, D.A. and Fujishima, A.: Electroanalytical study of sulfa drugs at diamond electrodes and their determination by HPLC with amperometric detection. *J. Electroanal. Chem.,* 491 (2000) 175-181; *C.A.,* 134 (2001) 46930s.

1921 Saunders, K.C., Venn, R.F. and Fane, J.A.: Analysis of CP-122,288 at femto-molar levels from human plasma using solid-supported extraction and HPLC column switching. *Chromatographia,* 52(Supplement) (2000) S87-S91.

1922 Schneider, M.J. and Donoghue, D.J.: Multiresidue determination of fluoroquinolones in eggs. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1306-1312.

1923 Shiraki, N., Hamada, A., Yasuda, K., Fujii, J., Arimoori, K. and Nakano, M.: Inhibitory effect of human immunodeficiency virus protease inhibitors on multidrug resistence transporter P-glycoprotein. *Biol. Pharm. Bull.,* 23 (2000) 1528-1531.

1924 Umehara, K., Kudo, S., Hirao, Y., Morita, S., Ohtani, T., Uchida, M. and Miyamoto, G.: *In vitro* characterization of the oxidative cleavages of the octyl side chain of olanexidine, a novel antimicrobial agent, in dog liver microsomes. *Drug Metab. Disp.,* 28 (2000) 1417-1424.

See also 1383, 1633, 1741.

32f. Cytostatics

1925 Allain, P., Huedi, O., Cailleux, A., le Bouil, A., Larra, F., Boisdron-Celle, M. and Gamelin, E.: Early biotransformation of oxaliplatin after its intravenous administration to cancer patients.

1930 Komatsu, T., Yamazaki, H., Shimada, N., Nakajima, M. and Yokoi, T.: Roles of cytochromes P450 1A2, 2A6, and 2C8 in 5-fluorouracil formation from tegafur, an anticancer prodrug, in human liver microsomes. *Drug Metab. Disp.,* 28 (2000) 1457-1463.

1931 Micoli, G., Turci, R., Arpellini, M. and Minoia, C.: Determination of 5-fluorouracil in environmental samples by solid-phase extraction and high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B,* 750 (2001) 25-32.

1932 Nguyen, T., Eshraghi, J., Gonyea, G., Ream, R. and Smith, R.: Studies on factors influencing stability and recovery of paclitaxel from suspension media and cultures of *Taxus cuspidata* cv *Densiformis* by high-performance liquid chromatography. *J. Chromatogr. A,* 911 (2001) 55-61.

1933 Onorato, J.M., Henion, J.D., Legebvre, P.M. and Kiplinger, J.P.: Selected reaction monitoring LC-MS determination of idoxifene and its pyrrolidinone metabolite in human plasma using robotic high-throughput, sequential sample injection. *Anal. Chem.,* 73 (2001) 119-125.

1934 Parise, R.A., Miles, D.R. and Egorin, M.J.: Sensitive high-performance liquid chromatographic assay for motexafin gadolinium and motexafin lutetium in human plasma. *J. Chromatogr. B,* 749 (2000) 145-152.

1935 Singh, J., Rastogi, S.K., Singh, S.N. and Bhatia, J.S.: Quantitation of leuprolide acetate by high performance liquid chromatography. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3023-3031.

1936 Thaler, A., Hottkowitz, T. and Eibl, H.: Separation and quantification of alkylphosphocholines by reversed phase high performance liquid chromatography. *Chem. Phys. Lipids,* 107 (2000) 131-139; *C.A.,* 133 (2000) 286582c.

1937 Zhang, H., Wen, L., Zhang, S. and Du, P.: (HPLC assay for methotrexate utilizing solid phase extraction and purification and separation of biological sample). *Yaowu Fenxi Zazhi,* 20, No.6 (2000) 401-404; *C.A.,* 134 (2001) 61667e.

See also 1037, 1132, 2024.

1971 Yokoyama, T., Mori, K., Nakamura, Y., Ohnuki, N., Matsushima, Y. and Fujii, T.: (Simultaneous determination of indicative ingredients and restrictive ingredients in cosmetics by HPLC). *Tokyo-toritsu Eisei Kenkyusho Kenkyu Nenpo,* 50 (1999, Pub. 2000) 65-69; *C.A.*, 134 (2001) 46614s.

1972 Yuan, B., Li, W. and Li, L.: (Determination of nicomol in tablets by HPLC). *Yaowu Fenxi Zazhi,* 20 (2000) 418-419; *C.A.*, 134 (2001) 46928x.

1973 Yuan, H., Xiao, X., Liu, F., He, C., Cai, G., Zhang, G. and Liu, Z.: (Determination of emodin in Lede capsules by HPLC). *Zhongguo Zhongyao Zazhi,* 25 (2000) 533-535; *C.A.*, 133 (2000) 340340n.

See also 1146, 1291, 1414, 1633, 1717, 1849, 1858, 2027.

32h. Toxicological and forensic applications

1974 Aalberg, L., DeRuiter, J., Noggle, F.T., Sippola, E. and Clark, C.R.: Chromatographic and mass spectral methods of identification for the side-chain and ring regioisomers of methylenedioxymethamphetamine. *J. Chromatogr. Sci.,* 38 (2000) 329-337.

1975 Backmund, M., Meyer, K., von Zielonka, M. and Eichenlaub, D.: Ofloxacin causes false-positive immunoassay results for urine opiates. *Addict. Biol.,* 5 (2000) 319-320; *C.A.*, 134 (2001) 36630e.

1976 Clauwaert, K.M., van Bocxlaer, J.F., de Letter, E.A., van Calenbergh, S., Lambert, W.E. and de Leenheer, A.P.: Determination of the designer drugs 3,4-methylenedioxy-methamphetamine, 3,4-methylenedioxyethylamphetamine, and 3,4-methylenedioxyamphetamine with HPLC and fluorescence detection in whole blood, serum, vitreous humor, and urine. *Clin. Chem. (Washington),* 46 (2000) 1968-1977.

1977 Goto, H., Igarashi, T., Yamamoto, M., Yasuda, M., Sekiguchi, R., Watai, M., Tanno, K. and Yasumoto, T.: Quantitative determination of marine toxins associated with diarrhetic shellfish poisoning by liquid chromatography coupled with mass spectrometry. *J. Chromatogr. A,* 907 (2001) 181-189.

1978 Haffen, E., Paintaud, G., Berard, M., Masuyer, C., Bechtel, Y. and Bechtel, P.R.: On the assessment of drug metabolism by assays of codeine and its main metabolites. *Ther. Drug Monit.,* 22 (2000) 258-265; *C.A.*, 133 (2000) 275784u.

1979 Spoof, L., Karlsson, K. and Meriluoto, J.: High-performance liquid chromatographic separation of microcystins and nodularin, cyanobacterial peptide toxins, on C_{18} and amide C_{16} sorbents. *J. Chromatogr. A,* 909 (2001) 225-236.

1980 Vale, P. and Sampayo, M.A.de M.: Determination of paralytic shellfish toxins in Portuguese shellfish by automated pre-column oxidation. *Toxicon,* 39 (2001) 561-571; *C.A.*, 134 (2001) 67255r.

See also 1120, 1402, 1404, 1660, 2050.

32i. Plant extracts and traditional Chinese medicines

1981 Arai, T., Zhao, C., Mizukami, H., Kojima, K., Yasui, T., Furuhashi, S., Sekita, S., Satake, M. and Ogihara, Y.: (Quantitative determination of shikonin in Shiunko). *Nat. Med. (Tokyo),* 54, No. 2 (2000) 81-85; *C.A.*, 133 (2000) 183140c.

1982 Cai, X., Cao, Z. and Liu, Z.: (Determination of emodin in Gongkangshu by RP-HPLC). *Huaxi Yaoxue Zazhi,* 15 (2000) 210-211; *C.A.*, 133 (2000) 198785r.

1983 Chen, W., Hu, G.-l., Wang, Y.-r. and Wang, X.-r.: (Determination of six ginsenosides in Panax species by high performance liquid chromatography). *Sepu,* 18 (2000) 439-441; *C.A.*, 134 (2001) 61619r.

1984 Chen, X., Li, W., Zhu, J. and Cui, H.: (RP-HPLC determination of cyasterone in radix *Cyathula officinalis*). *Yaowu Fenxi Zazhi,* 20 (2000) 234-237; *C.A.*, 133 (2000) 183115y.

1985 Chen, Y., Mo, Z. and Lu, M.: (Determination of the pueraria content in Xinkeshu tablets by RP-HPLC). *Guangxi Yike Daxue Xuebao,* 17 (2000) 414-415; *C.A.*, 134 (2001) 46884e.

1986 Cheng, B., Liu, Q. and Yu, X.: (HPLC determination of dissolution rates for terfenadine tablets). *Zhongguo Yiyuan Yaoxue Zazhi,* 20 (2000) 471-473; *C.A.*, 133 (2000) 301021q.

1987 Deliorman, D. and Ergun, F.: High performance liquid chromatographic determination of suringin in *Viscum album* L. ssp. *album* samples collected from different host plants. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3033-3042.

1988 Dou, H., Zeng, S., Wang, Y. and Shang, M.: (HPLC determination of kaerophyllin in *Bupleurum smithii* var. *parvifolium*. *Zhongguo Zhongyao Zazhi,* 25, No. 8 (2000) 488-489; *C.A.*, 133 (2000) 325721u.

1989 Draves, A.H. and Walker, S.E.: Determination of hypericin and pseudohypericin in pharmaceutical preparations by liquid chromatography with fluorescence detection. *J. Chromatogr. B,* 749 (2000) 57-66.

1990 Fang, Y. and Chen, Z.: (Determination of homoplantaginin in compound *Salvia plebeia* granules by HPLC). *Yaowu Fenxi Zazhi,* 20, No. 6 (2000) 388-390; *C.A.*, 134 (2001) 61637t.

1991 Fu, G. and Zhao, S.: (HPLC determination of paeoniflorin in Jiechangan enema). *Yaowu Fenxi Zazhi,* 20 (2000) 338-340; *C.A.*, 134 (2001) 21552g.

1992 Gao, L. and Wang, W.: (HPLC determination of content uniformity in naloxone hydrochloride tablets). *Yaowu Fenxi Zazhi,* 20 (2000) 318-320; *C.A.*, 134 (2001) 9452s.

1993 Gonzalez, A.M., Salager, J.L., Brunetto, M.R., Callignani, M., Burguera, J.L. and Petit de Pena, Y.: Determination of molecular weight distribution of lignin derivatives by aqueous phase high performance size exclusion chromatography (HPSEC). *J. High Resolut. Chromatogr.,* 23 (2000) 693-696.

1994 Hoberg, E., Meier, B. and Sticher, O.: An analytical high performance liquid chromatographic method for the determination of agnuside and *p*-hydroxybenzoic acid contents in agni-casti fructus. *Phytochem. Anal.,* 11 (2000) 327-329; *C.A.*, 133 (2000) 313755v.

1995 Hou, X., Yu, Z., Xu, Z., Wang, Y., Dai, F. and Bi, K.: (Determination of contents of evodiamine and rutaecarpine in 34 species of *Evodia rutaecarpa* by HPLC). *Shenyang Yaoke Daxue Xuebao,* 17 (2000) 334-337; *C.A.*, 134 (2001) 61616n.

1996 Huang, M.-z.: (Determination of chlorophenamine maleate and paracetamol of Xiaoer Suang Pu Pian by HPLC). *Guizhou Shifan Daxue Xuebao, Ziran Kexueban,* 18, No. 2 (2000) 30-31; *C.A.*, 133 (2000) 286586g.

1997 Li, F.-M., Qiu, F., Wang, W.-h., Guo, X.-t. and Di, X.: (Determination of three principal components in Chuanjing tablets by using high performance liquid chromatography). *Sepu,* 18 (2000) 442-444; *C.A.*, 134 (2001) 9440m.

1998 Li, W., Chan, C.-L. and Lueng, H.-W.: Liquid chromatography-atmospheric pressure chemical ionization mass spectrometry as a tool for the characterization of anthraquinone derivatives from Chinese herbal medicine. *J. Pharm. Pharmacol.,* 52 (2000) 723-729; *C.A.,* 133 (2000) 183103t.

1999 Linh, P.T., Kim, Y.H., Hong, S.P., Jian, J.K. and Kang, J.S.: Quantitative determination of salidroside and tyrosol from the underground part of *Rhodiola rosea* by high performance liquid chromatography. *Arch. Pharmacal Res.,* 23 (2000) 349-352; *C.A.,* 133 (2000) 313748v.

2000 Lu, Y., Liu, G., Li, M., Li, D. and Yu, X.: (Determination of oridonin in leaves and stems of blushred Ragdosia by HPLC). *Yaowu Fenxi Zazhi,* 20, No. 6 (2000) 390-392; *C.A.,* 143 (2001) 61636u.

2001 Okamura, M., Maki, T., Ishida, S., Uraguchi, C., Onishi, Y., Sadasue, E., Tsuruta, Y. and Yagi, A.: Dissolution profiles of principal ingredients in Kampo medicinal powders by high-performance liquid chromatography. *Chem. Pharm. Bull.,* 48 (2000) 1782-1785.

2002 Qiao, F., Zhang, Q., Lu, S. and Feng, Z.: (Determination of isofraxidin in Ciwujia preparation by HPLC). *Yaowu Fenxi Zazhi,* 20, No.3 (2000) 168-169; *C.A.,* 133 (2000) 313716h.

2003 Qin, L. and Xu, X.: (Determination of rheum emodin and chrysophanol in sodium hydrogen carbonate tablets of *Rheum officinalis* by HPLC). *Yaowu Fenxi Zazhi,* 20 (2000) 274-275; *C.A.,* 133 (2000) 183119c.

2004 Schmoltzi, P. and Scherges, M.: (Confusion of Chinese herbal drugs). *Dtsch. Apoth. Ztg.,* 140 (2000) 4094-4103; *C.A.,* 133 (2000) 325545q.

2005 Shen, G. and Yao, W.: (Determination of ginkgolic acid using supercritical fluid extraction and high performance liquid chromatography). *Fenxi Huaxue,* 28 (2000) 985-988; *C.A.,* 133 (2000) 256928c.

2006 Tang, C. and Zhou, X.: (HPLC determination of paeonol in Erlong Zuoci pills). *Yaowu Fenxi Zazhi,* 20 (2000) 320-321; *C.A.,* 134 (2001) 21551f.

2007 Tian, N., Yang, G., Fang, Y., Li, J. and Zhang, K.: (Determination of astragaloside A in *Astragalus membranaceous* and its preparations by HPLC with ELSD). *Yaowu Fenxi Zazhi,* 20 (2000) 199-200; *C.A.,* 133 (2000) 325715v.

2008 Wang, H., Zou, H., Ni, J. and Guo, B.: Comparative separation of biologically active compounds in *Rhizoma chuanxiong* by affinity chromatography with α_1-acid glycoprotein and human serum albumin as stationary phases. *Chromatographia,* 52 (2000) 459-464.

2009 Wang, J., Xie, L., Liu, H., Wang, X. and Cai, S.: (HPLC determination of harpagoside and cinnamic acid in Xuanshen after processing with different methods). *Zhongguo Yaoxue Zazhi (Beijing),* 35 (2000) 375-378; *C.A.,* 133 (2000) 183113w.

2010 Wang, J., Zhang, Z.-h., Dai, Y.-j. and Wu, R.-j.: (Determination of picroside II in *Picrorhiza scrophulariiflora* by HPLC). *Zhongguo Yaoke Daxue Xuebao,* 31 (2000) 368-370; *C.A.,* 134 (2001) 46879g.

2011 Wang, Q., Jiang, Y., Ma, S. and Dang, X.: (Determination of notoginsenoside R_1 in radix *Panax notoginseng* by HPLC-ELSD). *Zhongguo Zhongyao Zazhi,* 25 (2000) 617-618; *C.A.,* 134 (2001) 33068s.

2012 Wang, X.: (HPLC determination of salidroside in 6 *Rhodiola species* growing in Qinghai). *Zhongguo Yaoxue Zazhi (Beijing),* 35 (2000) 513-514; *C.A.,* 133 (2000) 301276b.

2013 Weng, X. and Jin, M.: (Determination of Ciwujia injection by HPLC). *Zhongguo Yiyuan Yaoxue Zazhi,* 20 (2000) 474-476; *C.A.,* 134 (2001) 9432k.

2014 Wu, X., Hu, Z. and Xiong, Y.: (Determination of paeoniflorin and naringin in complex Ganxiankang oral solution by RP-HPLC). *Huaxi Yaoxue Zazhi,* 15 (2000) 373-374; *C.A.,* 134 (2001) 46876d.

2015 Xiao, J., Wang, X., Ma, P. and Li, C.: (Determination of astragaloside IV in Yupingfeng granules by RP-HPLC). *Huaxi Yaoxue Zazhi,* 15 (2000) 216-217; *C.A.,* 133 (2000) 198775n.

2016 Xiu, J., Che, Z. and Bi, K.: (Pre-column derivatization HPLC method for determination of asiaticoside in *Centella asiatica* and Sanjinpian). *Yaoxue Xuebao,* 35 (2000) 605-608; *C.A.,* 133 (2000) 366512v.

2017 Xu, F., Zhao, X., Qin, G. and Cui, D.: (Determination of dehydrocorydaline in Yuanhuzhitong tablets by HPLC). *Zhongguo Zhongyao Zazhi,* 25, No. 8 (2000) 478-479; *C.A.,* 133 (2000) 325736c.

2018 Yan, C., Su, B., Wei, L., Guo, L., Wang, R. and Liu, W.: (Determination of danshensu in Danshen-Tongmai oral liquid by HPLC). *Zhongguo Yiyuan Yaoxue Zazhi,* 20 (2000) 600-602; *C.A.,* 134 (2001) 9429q.

2019 Yang, Y., Wang, S., Yan, B. and Zhang, L.: (High performance liquid chromatographic determination of gardenoside in Tuihuang oral solution). *Zhongguo Yaoxue Zazhi (Beijing),* 35 (2000) 470-471; *C.A.,* 133 (2000) 183114x.

2020 Yao, M., Qi, Y., Bi, K., Wang, X. and Luo, X.: A precolumn derivatization high-performance liquid chromatographic method with improved sensitivity and specificity for the determination of astragaloside IV in Radix Astragali. *J. Chromatogr. Sci.,* 38 (2000) 325-328.

2021 Yu, C., Diao, C., Xia, W. and Zhou, B.: (Determination of ginsenoside Rg1 in Shenfu injection by RP-HPLC). *Yaowu Fenxi Zazhi,* 20 (2000) 263-265; *C.A.,* 133 (2000) 183117a.

2022 Zhang, L., Song, Y., Sun, C., Ren, B. and Fu, D.: (Determination of tanshinone IIa in Huozhen capsules by RP-HPLC). *Zhongguo Yoaxue Zazhi (Beijing),* 35 (2000) 552-554; *C.A.,* 133 (2000) 301304j.

2023 Zhang, M., Liu, X., Ju, D. and Zhu, Z.: (Determination of tanshinone II_A in Quxianruangan capsules by HPLC). *Shenyang Yaoke Daxue Xuebao,* 17 (2000) 353-354; *C.A.,* 133 (2000) 313726m.

2024 Zhao, D., Yang, X., Chen, L., Wang, B., Xu, J. and Zhu, H.: (Extraction and separation of antitumor components from *Ganoderma lucidum* (Leuss. ex Fr.) karst). *Zhongguo Zhongyao Zazhi,* 25 (2000) 288-290; *C.A.,* 134 (2001) 15038c.

See also 1188, 1228, 1236, 1293, 1335, 1338, 1346, 1639, 1846, 1932, 2046.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

2025 Ulrich, S.: Solid-phase microextraction in biomedical analysis. *J. Chromatogr. A,* 902 (2000) 167-194 - a review with 100 refs.

33b. Complex mixtures and profiling (single compounds by cross-reference only)

See 1240, 1340, 1381, 1515, 1563, 1568, 1612.

34. FOOD ANALYSIS

34a. General papers and reviews

See 990, 1718, 1797.

34b. Complex mixtures (single compounds by cross-reference only)

2026 Methacanon, P., Kennedy, J.F., Lloyd, L. and Paterson, M.: Application of gas and liquid chromatography in analysis of wine components. *Khimiya (Sofiya),* 9, No. 2 (2000) 97-100; *C.A.*, 133 (2000) 349728c.

See also 1213, 1286, 1295, 1391, 1412, 1459, 1680, 1786, 1913, 1919, 2049, 2086.

34c. Organoleptically important compounds (flavors, odors, volatiles)

2027 Conkins, D., Economou, J.E., Boersma, J.A., Dedhiya, M.G. and Hansen, G.: Reversed phase-high performance liquid chromatographic method to measure migration of semivolatile compound, vanillin, in ipratropium bromide inhalation solution. *PharmSci,* 1, No. 3 (1999) no pp. given; *C.A.*, 134 (2001) 61344x.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

2028 Fischer, K., Kotalik, J. and Kettrup, A.: Chromatographic properties of the ion-exclusion column IonPac ICE-AS6 and its application in environmental analysis, Part II: Application in environmental analysis. *J. Chromatogr. Sci.,* 38 (2000) 409-415.

2029 Yonekubo, J.: (Pretreatment of environmental endocrine disruptors and examples of analysis by LC/MS). *Eco Ind.*, 5, No. 11 (2000) 34-42; *C.A.*, 133 (2000) 316969d - a review with 21 refs.

See also 1360.

35b. Air pollution (complex mixtures; single compounds by cross-reference only)

See 1179, 1360, 2069.

35c. Water pollution (complex mixtures; single compounds by cross-reference only)

2030 Bruzzoniti, M.C., Sarzanini, C. and Mentasti, E.: Preconcentration of contaminants in water analysis. *J. Chromatogr. A,* 902 (2000) 289-309 - a review with 150 refs.

2031 Castillo, M. and Barceló, D.: Characterisation of organic pollutants in textile wastewaters and landfill leachate by using toxicity-based fractionation methods followed by liquid and gas chromatography coupled to mass spectrometric detection. *Anal. Chim. Acta,* 426 (2001) 253-264.

2032 Castillo, M., Alonso, M.C., Riu, J., Reinke, M., Klöter, G., Dizer, H., Fischer, B., Hansen, P.D. and Barceló, D.: Identification of cytotoxic compounds in European wastewaters during a field experiment. *Anal. Chim. Acta,* 426 (2001) 265-277.

2033 Hodge, E.M.: On-line monitoring of wastewater using ion chromatography. *J. Chromatogr. Sci.,* 38 (2000) 353-356.

2034 Hormazabal, V., Ostensvik, O., Underdal, B. and Skulberg, O.M.: Simultaneous determination of the cyanotoxins anatoxin A, microcystin desmethyl-3-RR, LR, RR and YR in water using liquid chromatography-mass spectrometry. *J. Liq. Chromatogr. Relat. Technol.,* 23 (2000) 3155-3164.

2035 Jackson, P.E., Weigert, C., Pohl, C.A. and Saini, C.: A new approach to the determination of inorganic anions in environmental waters by ion chromatography. *Proc.-Water Qual. Technol. Conf.* (computer optical disk) (1999) 1073-1083; *C.A.*, 133 (2000) 256438t.

2036 Kim, M.-H., Kim, T.-S., Kim, T.-K. and Park, S.K.: (Application of reversed-phase extraction for the HPLC analysis of microcystins in water). *Anal. Sci. Technol.,* 13 (2000) 399-402; *C.A.*, 133 (2000) 182546c.

2037 Newton, B. and Somerville, K.H.: Trends in the use of in chromatography for monitoring high-purity water at semiconductor plants. *Ultrapure Water,* 17, No. 6 (2000) 52-54; *C.A.*, 133 (2000) 198025t - a review with 7 refs.

2038 Öllers, S., Singer, H.P., Fässler, P. and Müller, S.R.: Simultaneous quantification of neutral and acidic pharmaceuticals and pesticides at the low-ng/l level in surface and waste water. *J. Chromatogr. A,* 911 (2001) 225-234.

2039 Preiss, A. and Levsen, K.: (Use of high-performance liquid chromatography complex with nuclear resonance spectroscopy (HPLC/NMR) and with mass spectrometry (HPLC/MS) for analysis of complex water samples). *Biol. Abwasserreinig.,* 11(Anwendung der LC-MS in der Wasseranalytik) (1999) 169-200; *C.A.*, 133 (2000) 285854f - a review with 27 refs.

2040 Reemtsma, T.: Prospects of toxicity-directed wastewater analysis. *Anal. Chim. Acta,* 426 (2001) 279-287.

2041 Zima, S.V., Savranskii, L.I. and Sukhan, V.V.: (Determination of toxic metals in natural waters by high-performance liquid chromatography). *Khim. Tekhnol. Vody,* 21 (1999) 485-493; *C.A.*, 133 (2000) 300780z.

See also 1039, 1172, 1173, 1175, 1180, 1200, 1233, 1246, 1287, 1303, 1327, 1350, 1404, 1651, 1662, 1761, 1765, 1776, 1779, 1804, 2058, 2061, 2064, 2068, 2079, 2080, 2083, 2087, 2093, 2097, 2102, 2103, 2104, 2105, 2106, 2108.

35d. Soil pollution (complex mixtures; single compounds by cross-reference only)

See 1171, 1189, 1780, 2058.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

2042 Cserhati, T. and Forgacs, E.: Retention behavior of oligomer surfactants on polyethylene-coated zirconia HPLC column. *Toxicol. Environ. Chem.,* 76 (2000) 29-34; *C.A.,* 133 (2000) 336884a.

2043 Field, J.A. and Sawyer, T.E.: High-performance liquid chromatography-diode array detection of trichloroethene and aromatic and aliphatic anionic surfactants used for surfactant-enhanced aquifer remediation. *J. Chromatogr. A,* 893 (2000) 253-260.

2044 Hu, W. and Haddad, P.R.: Absorption behaviour of zwitterionic surfactant onto a reversed-phase stationary phase. *Chromatographia,* 52 (2000) 547-551.

2045 Schroder, H.F. and Ventura, F.: Applications of liquid chromatography-mass spectrometry in environmental chemistry: characterization and determination of surfactants and their metabolites in water samples by modern mass spectrometric techniques. *Tech. Instrum. Anal. Chem.,* 21(Sample Handling and Trace Analysis of Pollutants) (2000) 827-933; *C.A.,* 133 (2000) 194970z - a review with 159 refs.

See also 1070, 1077, 1413, 2032.

36b. Antioxidants and preservatives

2046 Bicchi, C., Binello, A. and Rubiolo, P.: Determination of phenolic diterpene antioxidants in rosemary (*Rosmarinus officinalis* L.) with different methods of extraction and analysis. *Phytochem. Anal.,* 11 (2000) 236-242; *C.A.,* 133 (2000) 249n.

2047 Boyce, M. and Spickett, E.: Separation and quantification of preservatives using ion pair HPLC and CZE: an extended investigation of separation mechanisms. *J. Chem. Educ.,* 77 (2000) 740-742; *C.A.,* 133 (2000) 192703r.

2048 Costa, L., Vega, M., Diaz, Y., Marcelo, J.L., Hernandez, J.M. and Martino, T.: Reversed-phase high-performance liquid chromatography versus spectrophotometric assay for thimerosal in Cuban recombinant hepatitis B vaccine. *J. Chromatogr. A,* 907 (2001) 173-179.

2049 Oshima, H., Ueno, E., Saito, I. and Matsumoto, H.: Determination of sodium azide in beverages by ion chromatography. *J. Assoc. Off. Anal. Chem.,* 83 (2000) 1410-1414.

See also 1230, 1291, 1361.

36c. Complex mixtures, technical products and unidentified compounds

2050 Furton, K.G., Wu, L. and Almirall, J.R.: Optimization of solid-phase microextraction (SPME) for the recovery of explosives from aqueous and post-explosion debris followed by gas and liquid chromatographic analysis. *J. Forensic Sci.,* 45 (2000) 857-864; *C.A.,* 133 (2000) 262412w.

See also 1081, 1376, 2049, 2075.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

2051 Chiu, G.N.C., Bally, M.B. and Mayer, L.D.: Selective protein interactions with phosphatidylserine containing liposomes alter the steric stabilization properties of poly(ethylene glycol). *Biochim. Biophys. Acta,* 1510 (2001) 56-69.

2052 Ujam, L.B., Clemmitt, R.H. and Chase, H.A.: Cell separation by expanded bed adsorption: use of ion exchange chromatography for the separation of *E. coli* and *S. cerevisiae*. *Bioprocess Eng.,* 23 (2000) 245-250; *C.A.,* 134 (2001) 16563g.

See also 1162, 1479, 1810, 1812.

38. INORGANIC COMPOUNDS

38a. Cations

2053 Awadallah, R.M., Bobleter, O., Sherif, M.K., Belal, A.A. and Taha, G.M.H.: High performance liquid chromatographic (HPLC)-atomic absorption spectroscopic (AAS) determination of trace elements in some plant samples. *Aswan Sci. Technol. Bull.,* 18 (1997) 26-40; *C.A.,* 134 (2001) 53918x.

2054 Bashir, W. and Paull, B.: Determination of trace alkaline earth metals in brines using chelation ion chromatography with an iminodiacetic acid bonded silica column. *J. Chromatogr. A,* 907 (2001) 191-200.

2055 Bohrer, D., do Nascimento, P.C. and Mendonca, J.K.A.: On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography. *J. Chromatogr. B,* 750 (2001) 61-69.

2056 Chongyang, L., Jinzhang, G., Wu, Y., Boyang, L. and Huitao, L.: Separation and detection of lanthanide ions with nitrilotri(methylenephosphonic) acid as complexing agent and eluent by IPC. *Rare Met. (Beijing),* 19 (2000) 136-140; *C.A.,* 133 (2000) 187260q.

2057 Cowan, J., Shaw, M.J., Achterberg, E.P., Jones, P. and Nesterenko, P.N.: The ion chromatographic separation of high balence metal cations using a neutral polystyren resin dynamically modified with dipicolinic acid. *Analyst (Cambridge),* 125 (2000) 2157-2159.

2058 Guerin, T., Molenat, N., Astruc, A. and Pinel, R.: Arsenic speciation in some environmental samples: a comparative study of HG-GC-QFAAS and HPLC-ICP-MS methods. *Appl. Organomet. Chem.,* 14, No. 8 (2000) 401-410; *C.A.,* 133 (2000) 290482n.

2059 He, B., Jiang, G. and Xu, X.: Arsenic speciation based on ion exchange high-performance liquid chromatography hyphenated with hydride generation atomic fluorescence and on-line UV photo oxidation. *Fresenius J. Anal. Chem.,* 368 (2000) 803-808.

2060 Hu, N., Hasebe, K., Tanaka, K., Inoue, S. and Nagai, M.: Ion chromatographic analysis of hydrogenium ion using ODS modified with dodecylsulfate as stationary phase. *Analyst (Cambridge),* 125 (2000) 2160-2162.

2061 Itoyama, T., Kumanomidou, H., Higuti, T. and Ishii, T.: (On the basic study of the deep sea water). *Mizu Shori Gijutsu,* 41 (2000) 367-370; *C.A.,* 133 (2000) 198228m.

2062 Milacic, R., Scancar, J.: Determination of hexavalent chromium in lime-treated sewage sludge by anion-exchange fast protein liquid chromatography with electrothermal atomic absorption spectrometry detection. *Analyst (Cambridge),* 125 (2000) 1938-1942.

2063 Nagaoka, M.H. and Maitani, T.: Binding patterns of co-existing aluminium and iron to human serum transferin studied by HPLC-high resolution ICP-MS. *Analyst (Cambridge),* 125 (2000) 1962-1965.

2064 Raessler, M., Michalke, B., Schulte-Hostede, S. and Kettrup, A.: Long-term monitoring of arsenic and selenium species in contaminated groundwaters by HPLC and HG-AAS. *Sci. Total Environ.,* 258, No. 3 (2000) 171-181; *C.A.,* 133 (2000) 256417k.

2065 Sanchez, J.M., Hidalgo, M. and Salvado, V.: The separation of Au(III) and Pd(II) in hydrochloric acid solutions by strong anion type II exchange resins: the effect of counter ion concentration and temperature. *Solvent Extr. Ion Exch.,* 18 (2000) 1199-1217; *C.A.,* 133 (2000) 328886n.

2066 Sekiguchi, Y.: (Study for improvement of precision for ammonium ion by a gradient elution and a recycle technique in ion chromatography). *Kogyo Yosui,* 505 (2000) 28-32; *C.A.,* 133 (2000) 366072b.

2067 Svedsen, R. and Lund, W.: Speciation of Cu, Fe and Mn in beer using ion exchange separation and size-exclusion chromatography in combination with electrothermal atomic absorption spectrometry. *Analyst (Cambridge),* 125 (2000) 1833-1837.

2068 Szpunar, J., McSheehy, S., Polec, K., Vacchia, V., Mounicou, S., Rodriquez, I. and Lobinski, R.: Gas and liquid chromatography with inductively coupled plasma mass spectrometry detection for environmental speciation analysis - advances and limitations. *Spectrochim. Acta, Part B,* 55B (2000) 779-793; *C.A.,* 133 (2000) 277229r.

2069 Tanaka, K., Ohta, K., Hu, W.Z., Hasebe, K., Fritz, J.S., Haddad, P.R., Lee, K.-P. and Sarzanini, C.: (Monitoring of acid rain in East Asia by ion-exclusion/cation-exchange ion chromatography). *Nagoya Kogyo Gijutsu Kenkyusho Hokoku,* 49 (2000) 71-76; *C.A.,* 133 (2000) 242266y.

2070 Tirez, K., Brusten, W., van Roy, S., de Brucker, N. and Diels, L.: Characterization of inorganic selenium species by ion chromatography with ICP-MS detection in microbial-treated industrial waste water. *J. Anal. At. Spectrom.,* 15 (2000) 1087-1092.

2075 Wilcoxon, J.P., Martin, J.E. and Provencio, P.: Size distributions of gold nanoclusters studied by liquid chromatography. *Langmuir,* 16 (2000) 9912-9920; *C.A.,* 133 (2000) 355747z.

2076 Wolf, Ch., Rösick, U. and Brätter, P.: Quantification of the metal distribution in metallothioneins of the human liner by HPLC coupled with ICP-AES. *Fresenius J. Anal. Chem.,* 368 (2000) 839-843.

2077 Xu, Q., Xu, C., Wang, Y., Zhang, W., Jin, L., Tanaka, K., Haraguchi, H. and Itoh, A.: Polydiphenylamine-dodecyl sulfate films for the simultaneous amperometric determination of electroinactive anions and cations in ion-exclusion cation-exchange chromatography. *Fresenius J. Anal. Chem.,* 368 (2000) 791-796.

2078 Zhang, Z., Gao, J. and Wang, J.: (Determination of microaluminum by using on-line ion exchange-flow injection spectrophotometry). *Yejin Fenxi,* 20, No.5 (2000) 48-49; *C.A.,* 134 (2001) 65551k.

See also 1126, 2037, 2041, 2087, 2093, 2097, 2102.

38b. Anions

2079 Achilli, M. and Romele, L.: Ion chromatographic determination of bromate in drinking water by postcolumn reaction with fuchsin. *Am. Lab. (Shelton),* 32, No. 17 (2000) 60-61; *C.A.,* 133 (2000) 354856x.

2080 Clewell, R.A., Chaudhuri, S., Dickson, S., Cassady, R.S., Wallner, W.M., Eldridge, J.E. and Tsui, D.T.: Analysis of trace level perchlorate in drinking water and ground water by electrospray mass spectrometry. *Environ. Sci. Res.,* 57(Perchlorate in the Environment) (2000) 45-58; *C.A.,* 133 (2000) 286055h.

2081 Echigo, S., Minear, R.A. and Yamada, H.: Ultra-low bromate detection in drinking water: a post-column derivatization method without anion suppressors for bromate analysis. *ACS Symp. Ser.,* 761(Natural Organic Matter and Disinfection By-Products) (2000) 316-329; *C.A.,* 133 (2000) 242205c.

2082 Echigo, S., Zhang, X., Minear, R.A. and Plewa, M.J.: Differentiation of total organic brominated and chlorinated compounds in total organic halide measurement: a new approach with an ion-chromatographic technique. *ACS Symp. Ser.,* 761(Natural Organic Matter and Disinfection By-Products)