



ELSEVIER

Journal of Chromatography A, 949 (2002) 379–392

JOURNAL OF
CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Subject Index Vols. 948 and 949

Acetylcysteine

Advances in the evaluation of the stability and characteristics of the amino acid and amine derivatives obtained with the *o*-phthalaldehyde/3-mercaptopropionic acid and *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents. High-performance liquid chromatography–mass spectrometry study 949(2002)99

H-acid

Monitoring of azo dye degradation processes in a bioreactor by on-line high-performance liquid chromatography 949(2002)263

Affinity sorbents

Multifunctional fractionation of polyclonal antibodies by immunoaffinity high-performance monolithic disk chromatography 949(2002)163

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Albumin

Multifunctional fractionation of polyclonal antibodies by immunoaffinity high-performance monolithic disk chromatography 949(2002)163

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Alkaloids

Enantiomeric separation of *N*-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Simple high-performance liquid chromatography method for the simultaneous determination of ketoconazole and piperine in rat plasma and hepatocyte culture 949(2002)43

Simultaneous determination of *N*-oxides and free bases of pyrrolizidine alkaloids by cation-exchange solid-phase extraction and ion-pair high-performance liquid chromatography 949(2002)249

Alumina, polyethylene-coated

Quantitative relationship between the retention of peptides on a reversed-phase alumina support and their physicochemical parameters 948(2002)69

Amines

Advances in the evaluation of the stability and characteristics of the amino acid and amine derivatives obtained with the *o*-phthalaldehyde/3-mercaptopropionic acid and *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents. High-performance liquid chromatography–mass spectrometry study 949(2002)99

Behavior and characteristics of amine derivatives obtained with *o*-phthalaldehyde/3-mercaptopropionic acid and with *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents (Review) 949(2002)235

Amines, heterocyclic aromatic

Ion-trap tandem mass spectrometry for the determination of heterocyclic amines in food 948(2002)267

Amino acids

Application of a new chiral derivatizing agent to the enantioseparation of secondary amino acids 948(2002)283

Enantiomeric separation of *N*-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Advances in the evaluation of the stability and characteristics of the amino acid and amine derivatives obtained with the *o*-phthalaldehyde/3-mercaptopropionic acid and *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents. High-performance liquid chromatography–mass spectrometry study 949(2002)99

Indirect high-performance liquid chromatographic separation of stereoisomers of β -alkyl-substituted amino acids by the application of (*S*)-*N*-(4-nitrophenoxycarbonyl)phenylalanine methoxyethyl ester as chiral derivatizing agent 949(2002)125

Behavior and characteristics of amine derivatives obtained with *o*-phthalaldehyde/3-mercaptopropionic acid and with *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents (Review) 949(2002)235

Aminohydroxynaphthalenedisulfonic acid

Monitoring of azo dye degradation processes in a bioreactor by on-line high-performance liquid chromatography 949(2002)263

Aminoimidazozaarenes

Ion-trap tandem mass spectrometry for the determination of heterocyclic amines in food 948(2002)267

Angiotensin II receptor antagonists

Fast screening method for the determination of angiotensin II receptor antagonists in human plasma by high-performance liquid chromatography with fluorimetric detection 949(2002)49

Antibiotics

Preparation of monodisperse porous polymethacrylate microspheres and their application in the capillary electrochromatography of macrolide antibiotics 948(2002)193

Antibodies

Multifunctional fractionation of polyclonal antibodies by immunoaffinity high-performance monolithic disk chromatography 949(2002)163

Automation

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Azo dyes

Monitoring of azo dye degradation processes in a bioreactor by on-line high-performance liquid chromatography 949(2002)263

Background electrolyte composition

Enantiomeric separation of *N*-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Basic drugs

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Chiral separation of basic drugs by capillary electrophoresis with carboxymethylcyclodextrins 948(2002)331

Benzimidazoles

Determination of omeprazole and its metabolites in human plasma by liquid chromatography–mass spectrometry 949(2002)1

Bioequivalence

High-performance liquid chromatographic determination of tramadol and its *O*-desmethylated metabolite in blood plasma. Application to a bioequivalence study in humans 949(2002)11

Biogenic amines

Behavior and characteristics of amine derivatives obtained with *o*-phthalaldehyde/3-mercaptopropionic acid and with *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents (Review) 949(2002)235

β-Blockers

Quantitative determination of oxprenolol and timolol in urine by capillary zone electrophoresis 949(2002)91

Buffer composition

Effects of substituted cyclodextrins on the separation of aromatic sulphonic acids by capillary zone electrophoresis 948(2002)235

Separation of polycyclic aromatic hydrocarbons with sodium dodecylbenzenesulfonate in electrokinetic chromatography 949(2002)301

Caffeine

Selective solid-phase extraction sorbent for caffeine made by molecular imprinting 948(2002)163

Calixarenes

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Carboxylic acids

Relationship between chromatographic properties of aromatic carboxylic acids and their structure 948(2002)65

Catechins

Effect of particle size in preparative reversed-phase high-performance liquid chromatography on the isolation of epigallocatechin gallate from Korean green tea 949(2002)275

Ceramides

Separation, identification and quantitation of ceramides in human cancer cells by liquid chromatography–electrospray ionization tandem mass spectrometry 949(2002)225

2-Chloro-1-methylquinolinium tetrafluoroborate

Determination of different species of homocysteine in human plasma by high-performance liquid chromatography with ultraviolet detection 949(2002)141

Chip technology

Capillary electrochromatography of peptides on microfabricated poly(dimethylsiloxane) chips modified by cerium(IV)-catalyzed polymerization 948(2002)225

Chiral selectors

Enantiomeric separation of N-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Chlorpheniramine

Uniformly sized molecularly imprinted polymer for *d*-chlorpheniramine. Evaluation of retention and molecular recognition properties in an aqueous mobile phase 948(2002)77

Cinchona alkaloids

Enantiomeric separation of N-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Circular dichroism detection

Stereospecific analysis of loxoprofen in plasma by chiral column liquid chromatography with a circular dichroism-based detector 948(2002)303

Clotrimazole

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Cloud-point extraction

Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration 949(2002)291

Colchicine

Highly specific and sensitive liquid chromatography–tandem mass spectrometry method for the determination of 3-desmethylthiocolchicine in human plasma as analyte for the assessment of bioequivalence after oral administration of thiocolchicoside 949(2002)71

Column switching

Increasing sample throughput in pharmacological studies by using dual-column liquid chromatography with tandem mass spectrometry 948(2002)187

Computational fluid dynamics

Shear-flow-based chromatographic separations as an alternative to pressure-driven liquid chromatography 948(2002)19

Computer simulation

Two-dimensional optimization using different pairs of variables for the reversed-phase high-performance liquid chromatographic separation of a mixture of acidic compounds 948(2002)35

Advanced high-performance liquid chromatography method development. Discovering unexpected choices in chromatography 948(2002)43

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Critical chromatography

Endgroup-based separation and quantitation of polyamide-6,6 by means of critical chromatography 949(2002)337

Cyclodextrins

Effects of substituted cyclodextrins on the separation of aromatic sulphonic acids by capillary zone electrophoresis 948(2002)235

Prediction of selectivity for enantiomeric separations of uncharged compounds by capillary electrophoresis involving dual cyclodextrin systems 948(2002)321

Cysteine

Quantitative proteomics strategy involving the selection of peptides containing both cysteine and histidine from tryptic digests of cell lysates 949(2002)153

Cytochromes

Determination of omeprazole and its metabolites in human plasma by liquid chromatography–mass spectrometry 949(2002)1

Deacetylbaccatin III

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

Dendrimers

Application of capillary zone electrophoresis to the separation and characterization of poly(amidoamine) dendrimers with an ethylenediamine core 949(2002)351

3-Desmethylthiocolchicine

Highly specific and sensitive liquid chromatography–tandem mass spectrometry method for the determination of 3-desmethylthiocolchicine in human plasma as analyte for the assessment of bioequivalence after oral administration of thiocolchicoside 949(2002)71

O-desmethyltramadol

High-performance liquid chromatographic determination of tramadol and its *O*-desmethylated metabolite in blood plasma. Application to a bioequivalence study in humans 949(2002)11

Derivatization, LC

Application of a new chiral derivatizing agent to the enantioseparation of secondary amino acids 948(2002)283

Advances in the evaluation of the stability and characteristics of the amino acid and amine derivatives obtained with the *o*-phthalaldehyde/3-mercaptopropionic acid and *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents. High-performance liquid chromatography–mass spectrometry study 949(2002)99

Indirect high-performance liquid chromatographic separation of stereoisomers of β -alkyl-substituted amino acids by the application of (*S*)-*N*-(4-nitrophenoxycarbonyl)phenylalanine methoxyethyl ester as chiral derivatizing agent 949(2002)125

Determination of different species of homocysteine in human plasma by high-performance liquid chromatography with ultraviolet detection 949(2002)141

Behavior and characteristics of amine derivatives obtained with *o*-phthaldialdehyde/3-mercaptopropionic acid and with *o*-phthaldialdehyde/*N*-acetyl-L-cysteine reagents 949(2002)235

Detection, LC

Shear-flow-based chromatographic separations as an alternative to pressure-driven liquid chromatography 948(2002)19

Fourier transform infrared spectroscopy with a sample deposition interface as a quantitative detector in size-exclusion chromatography 948(2002)257

Stereospecific analysis of loxoprofen in plasma by chiral column liquid chromatography with a circular dichroism-based detector 948(2002)303

Dexamethasone sodium phosphate

High-performance liquid chromatographic method for the assay of dexamethasone and xylometazoline in nasal drops containing methyl *p*-hydroxybenzoate 949(2002)79

Dialysis

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Dibenzoxepins

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Dimethylaniline

Poly(alkylmethylsiloxanes) thermally immobilized on silica as stationary phases for high-performance liquid chromatography 948(2002)121

Dispersion

Mass transfer in rectangular chromatographic channels 948(2002)3

Domperidone

Rapid and sensitive liquid chromatography–tandem mass spectrometry method for the quantitation of domperidone in human plasma 949(2002)65

Dopamine

Biosynthesis in an on-line immobilized-enzyme reactor containing phenylethanolamine *N*-methyltransferase in single-enzyme and coupled-enzyme formats 948(2002)249

Dopamine β -hydroxylase

Biosynthesis in an on-line immobilized-enzyme reactor containing phenylethanolamine *N*-methyltransferase in single-enzyme and coupled-enzyme formats 948(2002)249

Doxepin

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Econazole

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Electrochromatography

Preparation of monodisperse porous polymethacrylate microspheres and their application in the capillary electrochromatography of macrolide antibiotics 948(2002)193

Capillary electrochromatography of peptides on microfabricated poly(dimethylsiloxane) chips modified by cerium(IV)-catalyzed polymerization 948(2002)225

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Electrospray ionization

On-line microdialysis of proteins with high-salt buffers for direct coupling of electrospray ionization mass spectrometry and liquid chromatography 948(2002)139

Enantiomer separation

Uniformly sized molecularly imprinted polymer for *d*-chlorpheniramine. Evaluation of retention and molecular recognition properties in an aqueous mobile phase 948(2002)77

Application of a new chiral derivatizing agent to the enantioseparation of secondary amino acids 948(2002)283

Enantiomeric separation of *N*-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Stereospecific analysis of loxoprofen in plasma by chiral column liquid chromatography with a circular dichroism-based detector 948(2002)303

Prediction of selectivity for enantiomeric separations of uncharged compounds by capillary electrophoresis involving dual cyclodextrin systems 948(2002)321

Chiral separation of basic drugs by capillary electrophoresis with carboxymethylcyclodextrins 948(2002)331

Indirect high-performance liquid chromatographic separation of stereoisomers of β -alkyl-substituted amino acids by the application of (*S*)-*N*-(4-nitrophenoxycarbonyl)phenylalanine methoxyethyl ester as chiral derivatizing agent 949(2002)125

Enzyme inhibitors

Determination and pharmacokinetic profile of omeprazole in rat blood, brain and bile by microdialysis and high-performance liquid chromatography 949(2002)35

Simple high-performance liquid chromatography method for the simultaneous determination of ketoconazole and piperine in rat plasma and hepatocyte culture 949(2002)43

Improved procedure for the the determination of rofecoxib in human plasma involving 96-well solid-phase extraction and fluorescence detection 949(2002)83

Enzymes

Biosynthesis in an on-line immobilized-enzyme reactor containing phenylethanolamine *N*-methyltransferase in single-enzyme and coupled-enzyme formats 948(2002)249

Global internal standard technology for comparative proteomics 949(2002)173

Epigallocatechin gallate

Effect of particle size in preparative reversed-phase high-performance liquid chromatography on the isolation of epigallocatechin gallate from Korean green tea 949(2002)275

Epinephrine

Biosynthesis in an on-line immobilized-enzyme reactor containing phenylethanolamine *N*-methyltransferase in single-enzyme and coupled-enzyme formats 948(2002)249

Erythromycins

Preparation of monodisperse porous polymethacrylate microspheres and their application in the capillary electrochromatography of macrolide antibiotics 948(2002)193

Indirect high-performance liquid chromatographic separation of stereoisomers of β -alkyl-substituted amino acids by the application of (*S*)-*N*-(4-nitrophenoxycarbonyl)phenylalanine methoxyethyl ester as chiral derivatizing agent 949(2002)125

Escherichia coli

Quantitative proteomics strategy involving the selection of peptides containing both cysteine and histidine from tryptic digests of cell lysates 949(2002)153

Ethylenediamine

Application of capillary zone electrophoresis to the separation and characterization of poly(amidoamine) dendrimers with an ethylenediamine core 949(2002)351

Extraction methods

Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration 949(2002)291

Flow profiles

Mass transfer in rectangular chromatographic channels 948(2002)3

Fluorimetric detection

Determination of reboxetine, a recent antidepressant drug, in human plasma by means of two high-performance liquid chromatography methods 949(2002)23

Food analysis

Ion-trap tandem mass spectrometry for the determination of heterocyclic amines in food 948(2002)267

Effect of reduced glutathione on the stability of pigments in paprika powders studied by multiwavelength spectrometry and high-performance liquid chromatography 949(2002)269

Analysis of thiabendazole and procymidone in fruits and vegetables by capillary electrophoresis–electrospray mass spectrometry 949(2002)359

Fruits

Analysis of thiabendazole and procymidone in fruits and vegetables by capillary electrophoresis–electrospray mass spectrometry 949(2002)359

Gamma irradiation

Poly(methyloctadecylsiloxane) immobilized on silica by γ radiation for use in solid-phase extraction 948(2002)171

Geometrical isomers

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Geothermal fluids

Suppressed ion chromatography for monitoring chemical impurities in steam for geothermal power plants 949(2002)281

Global internal standard technology

Global internal standard technology for comparative proteomics 949(2002)173

Glutathione

Determination of different species of homocysteine in human plasma by high-performance liquid chromatography with ultraviolet detection 949(2002)141

Effect of reduced glutathione on the stability of pigments in paprika powders studied by multiwavelength spectrometry and high-performance liquid chromatography 949(2002)269

Gradient elution

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. III. Separation mechanism and gradient optimization 949(2002)307

Endgroup-based separation and quantitation of polyamide-6,6 by means of critical chromatography 949(2002)337

Green tea

Effect of particle size in preparative reversed-phase high-performance liquid chromatography on the isolation of epigallocatechin gallate from Korean green tea 949(2002)275

Hexanesulfonic acid

Simultaneous determination of *N*-oxides and free bases of pyrrolizidine alkaloids by cation-exchange solid-phase extraction and ion-pair high-performance liquid chromatography 949(2002)249

Histidine

Quantitative proteomics strategy involving the selection of peptides containing both cysteine and histidine from tryptic digests of cell lysates 949(2002)153

Homocysteine

Determination of different species of homocysteine in human plasma by high-performance liquid chromatography with ultraviolet detection 949(2002)141

HPLC fluorescence

Fast screening method for the determination of angiotensin II receptor antagonists in human plasma by high-performance liquid chromatography with fluorimetric detection 949(2002)49

Human plasma

Determination of reboxetine, a recent antidepressant drug, in human plasma by means of two high-performance liquid chromatography methods 949(2002)23

Imino acids

Application of a new chiral derivatizing agent to the enantioseparation of secondary amino acids 948(2002)283

Immobilization

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. I. Physical and chemical characterizations 948(2002)97

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. II. Chromatographic evaluation 948(2002)109

Immobilized enzymes

Biosynthesis in an on-line immobilized-enzyme reactor containing phenylethanolamine *N*-methyltransferase in single-enzyme and coupled-enzyme formats 948(2002)249

Immobilized metal affinity chromatography

Quantitative proteomics strategy involving the selection of peptides containing both cysteine and histidine from tryptic digests of cell lysates 949(2002)153

Immunoaffinity chromatography

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

Multifunctional fractionation of polyclonal antibodies by immunoaffinity high-performance monolithic disk chromatography 949(2002)163

Immunoglobulins

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Infrared detection

Fourier transform infrared spectroscopy with a sample deposition interface as a quantitative detector in size-exclusion chromatography 948(2002)257

Injection methods

Shear-flow-based chromatographic separations as an alternative to pressure-driven liquid chromatography 948(2002)19

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. III. Separation mechanism and gradient optimization 949(2002)307

Inorganic ions

Suppressed ion chromatography for monitoring chemical impurities in steam for geothermal power plants 949(2002)281

Instrumentation

Shear-flow-based chromatographic separations as an alternative to pressure-driven liquid chromatography 948(2002)19

Interfaces, LC-IR

Fourier transform infrared spectroscopy with a sample deposition interface as a quantitative detector in size-exclusion chromatography 948(2002)257

Interfaces, LC-MS

On-line microdialysis of proteins with high-salt buffers for direct coupling of electrospray ionization mass spectrometry and liquid chromatography 948(2002)139

Determination of omeprazole and its metabolites in human plasma by liquid chromatography-mass spectrometry 949(2002)1

Internal standards

Global internal standard technology for comparative proteomics 949(2002)173

Ion-pair chromatography

Simultaneous determination of *N*-oxides and free bases of pyrrolizidine alkaloids by cation-exchange solid-phase extraction and ion-pair high-performance liquid chromatography 949(2002)249

Isotope analysis

Quantitative proteomics strategy involving the selection of peptides containing both cysteine and histidine from tryptic digests of cell lysates 949(2002)153

Kava pyrones

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Ketoconazole

Simple high-performance liquid chromatography method for the simultaneous determination of ketoconazole and piperine in rat plasma and hepatocyte culture 949(2002)43

Latex

Capillary electrophoresis for the determination of new markers of natural latex quality 949(2002)367

Leucotrienes

Increasing sample throughput in pharmacological studies by using dual-column liquid chromatography with tandem mass spectrometry 948(2002)187

Lipids

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Separation, identification and quantitation of ceramides in human cancer cells by liquid chromatography–electrospray ionization tandem mass spectrometry 949(2002)225

Liquid chromatography

Advanced high-performance liquid chromatography method development. Discovering unexpected choices in chromatography 948(2002)43

Liquid chromatography under critical conditions

Quantitation of functionality of poly(methyl methacrylate) by liquid chromatography under critical conditions followed by evaporative light-scattering detection. Comparison with NMR and titration 949(2002)327

Loxoprofen

Stereospecific analysis of loxoprofen in plasma by chiral column liquid chromatography with a circular dichroism-based detector 948(2002)303

Macrolide antibiotics

Preparation of monodisperse porous polymethacrylate microspheres and their application in the capillary electrochromatography of macrolide antibiotics 948(2002)193

Mass transfer

Mass transfer in rectangular chromatographic channels 948(2002)3

Mathematical modelling

Prediction of selectivity for enantiomeric separations of uncharged compounds by capillary electrophoresis involving dual cyclodextrin systems 948(2002)321

Melperone

Rapid and sensitive liquid chromatography–tandem mass spectrometry method for the quantitation of domperidone in human plasma 949(2002)65

Membranes

Preparation and characterization of polyalkene membranes modified with four different ion-exchange groups by radiation-induced graft polymerization 948(2002)129

2-Mercaptopropionic acid

Determination of different species of homocysteine in human plasma by high-performance liquid chromatography with ultraviolet detection 949(2002)141

Mercaptopropionic acid

Advances in the evaluation of the stability and characteristics of the amino acid and amine derivatives obtained with the *o*-phthalaldehyde/3-mercaptopropionic acid and *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents. High-performance liquid chromatography–mass spectrometry study 949(2002)99

Method development

Two-dimensional optimization using different pairs of variables for the reversed-phase high-performance liquid chromatographic separation of a mixture of acidic compounds 948(2002)35

Advanced high-performance liquid chromatography method development. Discovering unexpected choices in chromatography 948(2002)43

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Methyl-*p*-hydroxybenzoate

High-performance liquid chromatographic method for the assay of dexamethasone and xylometazoline in nasal drops containing methyl *p*-hydroxybenzoate 949(2002)79

Miconazole

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Microdialysis

On-line microdialysis of proteins with high-salt buffers for direct coupling of electrospray ionization mass spectrometry and liquid chromatography 948(2002)139

Determination and pharmacokinetic profile of omeprazole in rat blood, brain and bile by microdialysis and high-performance liquid chromatography 949(2002)35

Mobile phase composition

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Application of a new chiral derivatizing agent to the enantioseparation of secondary amino acids 948(2002)283

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Effect of particle size in preparative reversed-phase high-performance liquid chromatography on the isolation of epigallocatechin gallate from Korean green tea 949(2002)275

Molecular imprinting

Uniformly sized molecularly imprinted polymer for *d*-chlorpheniramine. Evaluation of retention and molecular recognition properties in an aqueous mobile phase 948(2002)77

Selective solid-phase extraction sorbent for caffeine made by molecular imprinting 948(2002)163

Monolith columns

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Monolithic discs

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Monolithic disks

Multifunctional fractionation of polyclonal antibodies by immunoaffinity high-performance monolithic disk chromatography 949(2002)163

Naphthalene

Poly(alkylmethylsiloxanes) thermally immobilized on silica as stationary phases for high-performance liquid chromatography 948(2002)121

Naphthalenesulfonic acids

Effects of substituted cyclodextrins on the separation of aromatic sulphonic acids by capillary zone electrophoresis 948(2002)235

Nitrophenoxycarbonylphenylalanine methoxyethyl ester

Indirect high-performance liquid chromatographic separation of stereoisomers of β -alkyl-substituted amino acids by the application of (*S*)-*N*-(4-nitrophenoxycarbonyl)phenylalanine methoxyethyl ester as chiral derivatizing agent 949(2002)125

Application of a new chiral derivatizing agent to the enantioseparation of secondary amino acids 948(2002)283

Non-aqueous capillary electrophoresis

Enantiomeric separation of *N*-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Nonsteroidal anti-inflammatory drugs

Stereospecific analysis of loxoprofen in plasma by chiral column liquid chromatography with a circular dichroism-based detector 948(2002)303

Prediction of selectivity for enantiomeric separations of uncharged compounds by capillary electrophoresis involving dual cyclodextrin systems 948(2002)321

Oligomers

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. III. Separation mechanism and gradient optimization 949(2002)307

Omeprazole

Determination of omeprazole and its metabolites in human plasma by liquid chromatography–mass spectrometry 949(2002)1

Determination and pharmacokinetic profile of omeprazole in rat blood, brain and bile by microdialysis and high-performance liquid chromatography 949(2002)35

Optimization

Two-dimensional optimization using different pairs of variables for the reversed-phase high-performance liquid chromatographic separation of a mixture of acidic compounds 948(2002)35

Advanced high-performance liquid chromatography method development. Discovering unexpected choices in chromatography 948(2002)43

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. III. Separation mechanism and gradient optimization 949(2002)307

Organic acids

Two-dimensional optimization using different pairs of variables for the reversed-phase high-performance liquid chromatographic separation of a mixture of acidic compounds 948(2002)35

Monitoring of azo dye degradation processes in a bioreactor by on-line high-performance liquid chromatography 949(2002)263

Capillary electrophoresis for the determination of new markers of natural latex quality 949(2002)367

Organosulfur compounds

Effects of substituted cyclodextrins on the separation of aromatic sulphonic acids by capillary zone electrophoresis 948(2002)235

Highly specific and sensitive liquid chromatography–tandem mass spectrometry method for the determination of 3-desmethylthiocolchicine in human plasma as analyte for the assessment of bioequivalence after oral administration of thiocolchicoside 949(2002)71

Determination of different species of homocysteine in human plasma by high-performance liquid chromatography with ultraviolet detection 949(2002)141

Monitoring of azo dye degradation processes in a bioreactor by on-line high-performance liquid chromatography 949(2002)263

Oxprenolol

Quantitative determination of oxprenolol and timolol in urine by capillary zone electrophoresis 949(2002)91

Packed columns

Streaming potential in open and packed fused-silica capillaries 948(2002)203

Paclitaxel

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

Pancuronium

Validation of a high-performance liquid chromatography method for the determination of pancuronium in Pavulon injections 949(2002)61

Particle size

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Effect of particle size in preparative reversed-phase high-performance liquid chromatography on the isolation of epigallocatechin gallate from Korean green tea 949(2002)275

Partition coefficients

Relationship between chromatographic properties of aromatic carboxylic acids and their structure 948(2002)65

Peptides

Quantitative relationship between the retention of peptides on a reversed-phase alumina support and their physicochemical parameters 948(2002)69

Capillary electrochromatography of peptides on microfabricated poly(dimethylsiloxane) chips modified by cerium(IV)-catalyzed polymerization 948(2002)225

Quantitative proteomics strategy involving the selection of peptides containing both cysteine and histidine from tryptic digests of cell lysates 949(2002)153

Multifunctional fractionation of polyclonal antibodies by immunoaffinity high-performance monolithic disk chromatography 949(2002)163

Global internal standard technology for comparative proteomics 949(2002)173

Pesticides

Poly(methyloctadecylsiloxane) immobilized on silica by γ radiation for use in solid-phase extraction 948(2002)171

Analysis of thiabendazole and procymidone in fruits and vegetables by capillary electrophoresis–electrospray mass spectrometry 949(2002)359

pH effects

Quantitative relationship between the retention of peptides on a reversed-phase alumina support and their physicochemical parameters 948(2002)69

Pharmaceutical analysis

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Increasing sample throughput in pharmacological studies by using dual-column liquid chromatography with tandem mass spectrometry 948(2002)187

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Prediction of selectivity for enantiomeric separations of uncharged compounds by capillary electrophoresis involving dual cyclodextrin systems 948(2002)321

Chiral separation of basic drugs by capillary electrophoresis with carboxymethylcyclodextrins 948(2002)331

Determination of reboxetine, a recent antidepressant drug, in human plasma by means of two high-performance liquid chromatography methods 949(2002)23

Validation of a high-performance liquid chromatography method for the determination of pancuronium in Pavulon injections 949(2002)61

High-performance liquid chromatographic method for the assay of dexamethasone and xylometazoline in nasal drops containing methyl *p*-hydroxybenzoate 949(2002)79

Pharmacokinetics in humans

High-performance liquid chromatographic determination of tramadol and its *O*-desmethylated metabolite in blood plasma. Application to a bioequivalence study in humans 949(2002)11

Phenol

Poly(alkylmethylsiloxanes) thermally immobilized on silica as stationary phases for high-performance liquid chromatography 948(2002)121

Phenylethanolamine *N*-methyltransferase

Biosynthesis in an on-line immobilized-enzyme reactor containing phenylethanolamine *N*-methyltransferase in single-enzyme and coupled-enzyme formats 948(2002)249

Phosphatidylcholine

Phosphatidylcholine isolation from egg yolk phospholipids by high-performance liquid chromatography 949(2002)209

Phosphatidylethanolamine

Phosphatidylcholine isolation from egg yolk phospholipids by high-performance liquid chromatography 949(2002)209

Phospholipids

Phosphatidylcholine isolation from egg yolk phospholipids by high-performance liquid chromatography 949(2002)209

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Phthaldialdehyde

Advances in the evaluation of the stability and characteristics of the amino acid and amine derivatives obtained with the *o*-phthaldialdehyde/3-mercaptopropionic acid and *o*-phthaldialdehyde/*N*-acetyl-L-cysteine reagents. High-performance liquid chromatography-mass spectrometry study 949(2002)99

Pigments

Effect of reduced glutathione on the stability of pigments in paprika powders studied by multiwavelength spectrometry and high-performance liquid chromatography 949(2002)269

Piper methysticum

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Piperine

Simple high-performance liquid chromatography method for the simultaneous determination of ketoconazole and piperine in rat plasma and hepatocyte culture 949(2002)43

Plant materials

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

Simultaneous determination of *N*-oxides and free bases of pyrrolizidine alkaloids by cation-exchange solid-phase extraction and ion-pair high-performance liquid chromatography 949(2002)249

Plasma

Fast screening method for the determination of angiotensin II receptor antagonists in human plasma by high-performance liquid chromatography with fluorimetric detection 949(2002)49

Plate height

Mass transfer in rectangular chromatographic channels 948(2002)3

Poly(alkylmethylsiloxanes)

Poly(alkylmethylsiloxanes) thermally immobilized on silica as stationary phases for high-performance liquid chromatography 948(2002)121

Polyamide

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. III. Separation mechanism and gradient optimization 949(2002)307

Endgroup-based separation and quantitation of polyamide-6,6 by means of critical chromatography 949(2002)337

Poly(amidoamine) dendrimers

Application of capillary zone electrophoresis to the separation and characterization of poly(amidoamine) dendrimers with an ethylenediamine core 949(2002)351

Poly(dimethylsiloxane)

Capillary electrochromatography of peptides on microfabricated poly(dimethylsiloxane) chips modified by cerium(IV)-catalyzed polymerization 948(2002)225

Poly(glycidyl methacrylate-divinylbenzene) stationary phases

Preparation of monodisperse porous polymethacrylate microspheres and their application in the capillary electrochromatography of macrolide antibiotics 948(2002)193

Polymers

Fourier transform infrared spectroscopy with a sample deposition interface as a quantitative detector in size-exclusion chromatography 948(2002)257

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. III. Separation mechanism and gradient optimization 949(2002)307

Poly(methyl methacrylate)

Quantitation of functionality of poly(methyl methacrylate) by liquid chromatography under critical conditions followed by evaporative light-scattering detection. Comparison with NMR and titration 949(2002)327

Poly(methyloctylsiloxane)-silica stationary phases

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. II. Chromatographic evaluation 948(2002)109

Poly(methyloctadecylsiloxane)

Poly(methyloctadecylsiloxane) immobilized on silica by γ radiation for use in solid-phase extraction 948(2002)171

Poly(methyloctylsiloxane)-silica stationary phases

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. I. Physical and chemical characterizations 948(2002)97

Poly(methyloctylsiloxane)

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. I. Physical and chemical characterizations 948(2002)97

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. II. Chromatographic evaluation 948(2002)109

Polymer stationary phases

Preparation of monodisperse porous polymethacrylate microspheres and their application in the capillary electrochromatography of macrolide antibiotics 948(2002)193

Polymerization

Preparation and characterization of polyalkene membranes modified with four different ion-exchange groups by radiation-induced graft polymerization 948(2002)129

Polynuclear aromatic hydrocarbons

Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration 949(2002)291

Separation of polycyclic aromatic hydrocarbons with sodium dodecylbenzenesulfonate in electrokinetic chromatography 949(2002)301

Polyoxyethylene-10-lauryl ether

Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration 949(2002)291

Positional isomers

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Preparative chromatography

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Effect of particle size in preparative reversed-phase high-performance liquid chromatography on the isolation of epigallocatechin gallate from Korean green tea 949(2002)275

Principal component analysis

Quantitative relationship between the retention of peptides on a reversed-phase alumina support and their physicochemical parameters 948(2002)69

Effect of reduced glutathione on the stability of pigments in paprika powders studied by multiwavelength spectrometry and high-performance liquid chromatography 949(2002)269

Process control

Suppressed ion chromatography for monitoring chemical impurities in steam for geothermal power plants 949(2002)281

Process monitoring

Monitoring of azo dye degradation processes in a bioreactor by on-line high-performance liquid chromatography 949(2002)263

Procymidone

Analysis of thiabendazole and procymidone in fruits and vegetables by capillary electrophoresis-electrospray mass spectrometry 949(2002)359

Profens

Stereospecific analysis of loxoprofen in plasma by chiral column liquid chromatography with a circular dichroism-based detector 948(2002)303

Prediction of selectivity for enantiomeric separations of uncharged compounds by capillary electrophoresis involving dual cyclodextrin systems 948(2002)321

Propiverine

Increasing sample throughput in pharmacological studies by using dual-column liquid chromatography with tandem mass spectrometry 948(2002)187

Protein G

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Proteins

On-line microdialysis of proteins with high-salt buffers for direct coupling of electrospray ionization mass spectrometry and liquid chromatography 948(2002)139

Multifunctional fractionation of polyclonal antibodies by immunoaffinity high-performance monolithic disk chromatography 949(2002)163

Global internal standard technology for comparative proteomics 949(2002)173

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Pyrones

Computer-assisted optimization in the development of a high-performance liquid chromatographic method for the analysis of kava pyrones in *Piper methysticum* preparations 948(2002)51

Pyrrolizidines

Simultaneous determination of *N*-oxides and free bases of pyrrolizidine alkaloids by cation-exchange solid-phase extraction and ion-pair high-performance liquid chromatography 949(2002)249

Quinidines

Enantiomeric separation of *N*-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Quinines

Enantiomeric separation of *N*-protected amino acids by non-aqueous capillary electrophoresis with dimeric forms of quinine and quinidine derivatives serving as chiral selectors 948(2002)295

Reboxetine

Determination of reboxetine, a recent antidepressant drug, in human plasma by means of two high-performance liquid chromatography methods 949(2002)23

Rectangular channels

Mass transfer in rectangular chromatographic channels 948(2002)3

Resolution

Two-dimensional optimization using different pairs of variables for the reversed-phase high-performance liquid chromatographic separation of a mixture of acidic compounds 948(2002)35

Advanced high-performance liquid chromatography method development. Discovering unexpected choices in chromatography 948(2002)43

Resorcinarenes

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Retention factor

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Reviews

Behavior and characteristics of amine derivatives obtained with *o*-phthalaldehyde/3-mercaptopropionic acid and with *o*-phthalaldehyde/*N*-acetyl-L-cysteine reagents (Review) 949(2002)235

Robustness testing

Advanced high-performance liquid chromatography method development. Discovering unexpected choices in chromatography 948(2002)43

Rofecoxib

Improved procedure for the determination of rofecoxib in human plasma involving 96-well solid-phase extraction and fluorescence detection 949(2002)83

Sample handling

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

Sample preparation

Strategy for the development of automated methods involving dialysis and trace enrichment as on-line sample preparation for the determination of basic drugs in plasma by liquid chromatography 948(2002)151

Sandwich injection

Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. III. Separation mechanism and gradient optimization 949(2002)307

Scopolamine

Increasing sample throughput in pharmacological studies by using dual-column liquid chromatography with tandem mass spectrometry 948(2002)187

Selectivity

Prediction of selectivity for enantiomeric separations of uncharged compounds by capillary electrophoresis involving dual cyclodextrin systems 948(2002)321

Shear-driven chromatography

Shear-flow-based chromatographic separations as an alternative to pressure-driven liquid chromatography 948(2002)19

Silica, urea-modified

Preparation of a new C₁₈ stationary phase containing embedded urea groups for use in high-performance liquid chromatography 948(2002)85

Size-exclusion chromatography

Fourier transform infrared spectroscopy with a sample deposition interface as a quantitative detector in size-exclusion chromatography 948(2002)257

Sodium dodecylbenzenesulfonate

Separation of polycyclic aromatic hydrocarbons with sodium dodecylbenzenesulfonate in electrokinetic chromatography 949(2002)301

Solid-phase extraction

Selective solid-phase extraction sorbent for caffeine made by molecular imprinting 948(2002)163

Poly(methyloctadecylsiloxane) immobilized on silica by γ radiation for use in solid-phase extraction 948(2002)171

Improved procedure for the the determination of rofecoxib in human plasma involving 96-well solid-phase extraction and fluorescence detection 949(2002)83

Solvent elimination interfaces

Fourier transform infrared spectroscopy with a sample deposition interface as a quantitative detector in size-exclusion chromatography 948(2002)257

Sonic spray ionization

Determination of omeprazole and its metabolites in human plasma by liquid chromatography–mass spectrometry 949(2002)1

Sorbents

Poly(methyloctadecylsiloxane) immobilized on silica by γ radiation for use in solid-phase extraction 948(2002)171

Soya bean

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Sphingolipids

Separation, identification and quantitation of ceramides in human cancer cells by liquid chromatography–electrospray ionization tandem mass spectrometry 949(2002)225

Stationary phases, electrochromatography

Preparation of monodisperse porous polymethacrylate microspheres and their application in the capillary electrochromatography of macrolide antibiotics 948(2002)193

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Stationary phases, LC

Uniformly sized molecularly imprinted polymer for *d*-chlorpheniramine. Evaluation of retention and molecular recognition properties in an aqueous mobile phase 948(2002)77

Preparation of a new C₁₈ stationary phase containing embedded urea groups for use in high-performance liquid chromatography 948(2002)85

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. I. Physical and chemical characterizations 948(2002)97

High-performance liquid chromatographic stationary phases based on poly(methyloctylsiloxane) immobilized on silica. II. Chromatographic evaluation 948(2002)109

Poly(alkylmethylsiloxanes) thermally immobilized on silica as stationary phases for high-performance liquid chromatography 948(2002)121

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Quantitative investigation of the affinity properties of different recombinant forms of protein G by means of high-performance monolithic chromatography 949(2002)185

Fractionation of soybean phospholipids by preparative high-performance liquid chromatography with sorbents of various particle size 949(2002)217

Sterols

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Streaming potential

Streaming potential in open and packed fused-silica capillaries 948(2002)203

Structure–retention relationships

Relationship between chromatographic properties of aromatic carboxylic acids and their structure 948(2002)65

Sulfonic acids

Effects of substituted cyclodextrins on the separation of aromatic sulphonic acids by capillary zone electrophoresis 948(2002)235

Surface characterization

Streaming potential in open and packed fused-silica capillaries 948(2002)203

Surfactants

Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration 949(2002)291

Separation of polycyclic aromatic hydrocarbons with sodium dodecylbenzenesulfonate in electrokinetic chromatography 949(2002)301

Talinolol

Increasing sample throughput in pharmacological studies by using dual-column liquid chromatography with tandem mass spectrometry 948(2002)187

Taxanes

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

Taxol

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

***Taxus* spp**

Immunoaffinity chromatography for the sample pretreatment of *Taxus* plant and cell extracts prior to analysis of taxanes by high-performance liquid chromatography 948(2002)177

Terephthalic acid

Preparation and characterization of polyalkene membranes modified with four different ion-exchange groups by radiation-induced graft polymerization 948(2002)129

Theobromine

Selective solid-phase extraction sorbent for caffeine made by molecular imprinting 948(2002)163

Theophylline

Selective solid-phase extraction sorbent for caffeine made by molecular imprinting 948(2002)163

Thiabendazole

Analysis of thiabendazole and procymidone in fruits and vegetables by capillary electrophoresis–electrospray mass spectrometry 949(2002)359

Thiocolchicoside

Highly specific and sensitive liquid chromatography–tandem mass spectrometry method for the determination of 3-desmethylthiocolchicine in human plasma as analyte for the assessment of bioequivalence after oral administration of thiocolchicoside 949(2002)71

Thioxanthenes

Separation of *cis*- and *trans*-isomers of thioxanthene and dibenz[*b,e*]oxepin derivatives on calixarene- and resorcinarene-bonded high-performance liquid chromatography stationary phases 948(2002)309

Timolol

Quantitative determination of oxprenolol and timolol in urine by capillary zone electrophoresis 949(2002)91

Tocopherols

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Tocotrienols

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Tramadol

High-performance liquid chromatographic determination of tramadol and its *O*-desmethylated metabolite in blood plasma. Application to a bioequivalence study in humans 949(2002)11

UV detection

Determination of reboxetine, a recent antidepressant drug, in human plasma by means of two high-performance liquid chromatography methods 949(2002)23

Validation

Validation of a high-performance liquid chromatography method for the determination of pancuronium in Pavulon injections 949(2002)61

Vegetable oils

Elution behavior of unsaponifiable lipids with various capillary electrochromatographic stationary phases 949(2002)195

Vegetables

Analysis of thiabendazole and procymidone in fruits and vegetables by capillary electrophoresis–electrospray mass spectrometry 949(2002)359

Water analysis

Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration 949(2002)291

Xanthines

Selective solid-phase extraction sorbent for caffeine made by molecular imprinting 948(2002)163

Xylometazoline hydrochloride

High-performance liquid chromatographic method for the assay of dexamethasone and xylometazoline in nasal drops containing methyl *p*-hydroxybenzoate 949(2002)79

Zeta potential

Streaming potential in open and packed fused-silica capillaries 948(2002)203