

Journal of Chromatography A, 955 (2002) 283-288

JOURNAL OF CHROMATOGRAPHY A

www.elsevier.com/locate/chroma

Subject Index Vol. 955

Activity coefficients

Gas chromatographic determination of vapour pressure and related thermodynamic properties of monoterpenes and biogenically related compounds 955(2002)105

Adsorption isotherm

Measurement and modeling of the equilibrium behavior of the Tröger's base enantiomers on an amylose-based chiral stationary phase 955(2002)35

Air analysis

Gas chromatographic determination of glutaraldehyde in the workplace atmosphere after derivatization with O-(2,3,4,5,6pentafluorobenzyl)hydroxylamine on a solid-phase microextraction fibre 955(2002)117

Albumin

Use of a linear gradient flow program for liquid chromatography-mass spectrometry protein-binding studies 955(2002)237

Alcohols

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Aldehydes

Gas chromatographic determination of glutaraldehyde in the workplace atmosphere after derivatization with O-(2,3,4,5,6pentafluorobenzyl)hydroxylamine on a solid-phase microextraction fibre 955(2002)117

Alkaline earth ions

Ethylenediamine as eluent component in cation chromatography. Predictive and comparative study for analysis of alkaline earth ions 955(2002)1

Alkaloids

Screening of Catharanthus roseus secondary metabolites by high-performance liquid chromatography 955(2002)87

Amines, aromatic

Harmful azo colorants in leather. Determination based on their cleavage and extraction of corresponding carcinogenic aromatic amines using modern extraction techniques 955(2002)215

Amino acids

New triazine spectroscopic reagent for the separation of DLamino acids by micellar electrokinetic chromatography 955(2002)125

Chiral separation of amino acids derivatized with fluoresceine-5isothiocyanate by capillary electrophoresis and laser-induced fluorescence detection using mixed selectors of β-cyclodextrin and sodium taurocholate 955(2002)133

Direct high-performance liquid chromatographic enantioseparation of apolar β-amino acids on a quinine-derived chiral anion-exchanger stationary phase 955(2002)141

Anion-exchangers

Ion-exchange separation of proteins by polyallylamine-grafted cellulose gel 955(2002)191

Studies on separation of rare earth elements on various types of anion-exchangers in the C₃H₇OH-7 M HNO₃ systems 955(2002)257

Azo dves

Harmful azo colorants in leather. Determination based on their cleavage and extraction of corresponding carcinogenic aromatic amines using modern extraction techniques 955(2002)215

Carbon loads

Chromatographic behaviour of selected steroids and their inclusion complexes with β-cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Carbonyl compounds

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Catharanthus roseus

Screening of Catharanthus roseus secondary metabolites by high-performance liquid chromatography 955(2002)87

Ion-exchange separation of proteins by polyallylamine-grafted cellulose gel 955(2002)191

0021-9673/02/\$ - see front matter © 2002 Elsevier Science B.V. All right reserved PII: S0021-9673(02)00421-1

Chiral stationary phases, LC

Measurement and modeling of the equilibrium behavior of the Tröger's base enantiomers on an amylose-based chiral stationary phase 955(2002)35

Separation of chiral sulfoxides by liquid chromatography using macrocyclic glycopeptide chiral stationary phases 955(2002)53

Amino-bonded silica as stationary phase for liquid chromatographic determination of cyclopiazonic acid in fungal extracts 955(2002)79

Direct high-performance liquid chromatographic enantioseparation of apolar β -amino acids on a quinine-derived chiral anion-exchanger stationary phase 955(2002)141

Continuous-flow liquid membrane extraction

Determination of sulfonylurea herbicides by continuous-flow liquid membrane extraction on-line coupled with high-performance liquid chromatography 955(2002)183

Critical interaction mode

Theory of liquid chromatography of mono- and difunctional macromolecules. I. Studies in the critical interaction mode 955(2002)9

β-Cyclodextrin

Chiral separation of amino acids derivatized with fluoresceine-5-isothiocyanate by capillary electrophoresis and laser-induced fluorescence detection using mixed selectors of β -cyclodextrin and sodium taurocholate 955(2002)133

Cyclodextrins

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Cyclopiazonic acid

Amino-bonded silica as stationary phase for liquid chromatographic determination of cyclopiazonic acid in fungal extracts 955(2002)79

Derivatisation, LC

Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains 955(2002)245

Derivatization, electrophoresis

New triazine spectroscopic reagent for the separation of DLamino acids by micellar electrokinetic chromatography 955(2002)125

Derivatization, GC

Gas chromatographic determination of glutaraldehyde in the workplace atmosphere after derivatization with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine on a solid-phase microextraction fibre 955(2002)117

Derivitisation, electrophoresis

Chiral separation of amino acids derivatized with fluoresceine-5-isothiocyanate by capillary electrophoresis and laser-induced fluorescence detection using mixed selectors of β -cyclodextrin and sodium taurocholate 955(2002)133

Dichlorotriazinylamino-dimethylaminomethylphenazine

New triazine spectroscopic reagent for the separation of DLamino acids by micellar electrokinetic chromatography 955(2002)125

Drugs

Use of a linear gradient flow program for liquid chromatography-mass spectrometry protein-binding studies 955(2002)237

Dual-field and flow-programming

Field and flow programming in frit-inlet asymmetrical flow field-flow fractionation 955(2002)263

Enantiomer separation

Measurement and modeling of the equilibrium behavior of the Tröger's base enantiomers on an amylose-based chiral stationary phase 955(2002)35

Separation of chiral sulfoxides by liquid chromatography using macrocyclic glycopeptide chiral stationary phases 955(2002)53

New triazine spectroscopic reagent for the separation of DLamino acids by micellar electrokinetic chromatography 955(2002)125

Chiral separation of amino acids derivatized with fluoresceine-5-isothiocyanate by capillary electrophoresis and laser-induced fluorescence detection using mixed selectors of β -cyclodextrin and sodium taurocholate 955(2002)133

Direct high-performance liquid chromatographic enantioseparation of apolar β -amino acids on a quinine-derived chiral anion-exchanger stationary phase 955(2002)141

Equilibrium-dispersive model

Measurement and modeling of the equilibrium behavior of the Tröger's base enantiomers on an amylose-based chiral stationary phase 955(2002)35

Esters

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Estrogens

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Ethylenediamine

Ethylenediamine as eluent component in cation chromatography. Predictive and comparative study for analysis of alkaline earth ions 955(2002)1

Extraction methods

Harmful azo colorants in leather. Determination based on their cleavage and extraction of corresponding carcinogenic aromatic amines using modern extraction techniques 955(2002)215

Field programming

Different elution modes and field programming in gravitational field-flow fractionation. IV. Field programming achieved with channels of non-constant cross-sections 955(2002)95

Field-flow fractionation

Different elution modes and field programming in gravitational field-flow fractionation. IV. Field programming achieved with channels of non-constant cross-sections 955(2002)95

Field and flow programming in frit-inlet asymmetrical flow field-flow fractionation 955(2002)263

Flow-velocity gradients

Different elution modes and field programming in gravitational field-flow fractionation. IV. Field programming achieved with channels of non-constant cross-sections 955(2002)95

Frit inlet asymmetrical flow field-flow fractionation

Field and flow programming in frit-inlet asymmetrical flow field-flow fractionation 955(2002)263

Fumonisins

Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains 955(2002)245

Fusarium spp.

Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains 955(2002)245

Glutaraldehyde

Gas chromatographic determination of glutaraldehyde in the workplace atmosphere after derivatization with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine on a solid-phase microextraction fibre 955(2002)117

Glycopeptides

Separation of chiral sulfoxides by liquid chromatography using macrocyclic glycopeptide chiral stationary phases 955(2002)53

Gradient elution

Use of a linear gradient flow program for liquid chromatography-mass spectrometry protein-binding studies 955(2002)237

Gravitational field-flow fractionation

Different elution modes and field programming in gravitational field-flow fractionation. IV. Field programming achieved with channels of non-constant cross-sections 955(2002)95

Hold-up time

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Honey

Rapid determination of minority organic acids in honey by high-performance liquid chromatography 955(2002)207

Hormones

Separation and detection of neuroactive steroids from biological matrices (Review) 955(2002)151

Reversed-phase high-performance liquid chromatography method for the determination of prolactin in bacterial extracts and in its purified form 955(2002)229

Hydrodynamic lift forces

Different elution modes and field programming in gravitational field-flow fractionation. IV. Field programming achieved with channels of non-constant cross-sections 955(2002)95

Hydrodynamic relaxation

Field and flow programming in frit-inlet asymmetrical flow field-flow fractionation 955(2002)263

Immobilized proteins

Use of a linear gradient flow program for liquid chromatography-mass spectrometry protein-binding studies 955(2002)237

Indoles

Screening of *Catharanthus roseus* secondary metabolites by high-performance liquid chromatography 955(2002)87

Iridoids

Screening of *Catharanthus roseus* secondary metabolites by high-performance liquid chromatography 955(2002)87

Lipids

Separation and detection of neuroactive steroids from biological matrices (Review) 955(2002)151

Mathematical modelling

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Microwave-assisted extraction

Harmful azo colorants in leather. Determination based on their cleavage and extraction of corresponding carcinogenic aromatic amines using modern extraction techniques 955(2002)215

Mobile phase composition

Ethylenediamine as eluent component in cation chromatography. Predictive and comparative study for analysis of alkaline earth ions 955(2002)1

Studies on separation of rare earth elements on various types of anion-exchangers in the C₃H₇OH-7 *M* HNO₃ systems 955(2002)257

Molecular descriptors

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Monoterpenes

Gas chromatographic determination of vapour pressure and related thermodynamic properties of monoterpenes and biogenically related compounds 955(2002)105

Multilayer adsorption

Measurement and modeling of the equilibrium behavior of the Tröger's base enantiomers on an amylose-based chiral stationary phase 955(2002)35

Mycotoxins

Amino-bonded silica as stationary phase for liquid chromatographic determination of cyclopiazonic acid in fungal extracts 955(2002)79

Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains 955(2002)245

Neural networks, artificial

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Organic acids

Rapid determination of minority organic acids in honey by high-performance liquid chromatography 955(2002)207

Pentafluorobenzylhydroxylamine

Gas chromatographic determination of glutaraldehyde in the workplace atmosphere after derivatization with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine on a solid-phase microextraction fibre 955(2002)117

Peptides

Reversed-phase high-performance liquid chromatography method for the determination of prolactin in bacterial extracts and in its purified form 955(2002)229

Pesticides

Determination of sulfonylurea herbicides by continuous-flow liquid membrane extraction on-line coupled with high-performance liquid chromatography 955(2002)183

Plant materials

Screening of *Catharanthus roseus* secondary metabolites by high-performance liquid chromatography 955(2002)87

Polarity measurements

Prediction of the retention in reversed-phase liquid chromatography using solute-mobile phase-stationary phase polarity parameters 955(2002)19

Polyallylamine

Ion-exchange separation of proteins by polyallylamine-grafted cellulose gel 955(2002)191

Polyethylene glycols

Theory of liquid chromatography of mono- and difunctional macromolecules. I. Studies in the critical interaction mode 955(2002)9

Polymers

Theory of liquid chromatography of mono- and difunctional macromolecules. I. Studies in the critical interaction mode 955(2002)9

Polystyrene sulfonate

Field and flow programming in frit-inlet asymmetrical flow field-flow fractionation 955(2002)263

Porous graphitic carbon

Role of the Na⁺ ion on phenol derivatives/hydroxypropyl-β-cyclodextrin complex formation on porous graphitic carbon phase 955(2002)197

Progestogens

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Prolectin

Reversed-phase high-performance liquid chromatography method for the determination of prolactin in bacterial extracts and in its purified form 955(2002)229

1-Propanol

Studies on separation of rare earth elements on various types of anion-exchangers in the $C_3H_7OH-7\ M$ HNO $_3$ systems 955(2002)257

2-Propanol

Studies on separation of rare earth elements on various types of anion-exchangers in the $C_3H_7OH-7\ M\ HNO_3$ systems 955(2002)257

Proteins

Ion-exchange separation of proteins by polyallylamine-grafted cellulose gel 955(2002)191

Reversed-phase high-performance liquid chromatography method for the determination of prolactin in bacterial extracts and in its purified form 955(2002)229

Use of a linear gradient flow program for liquid chromatography-mass spectrometry protein-binding studies 955(2002)237

Quantitative structure-property relationships

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Rare earth elements

Studies on separation of rare earth elements on various types of anion-exchangers in the $C_3H_7OH-7\ M\ HNO_3$ systems 955(2002)257

Regression analysis

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Retention indices

Gas chromatographic determination of vapour pressure and related thermodynamic properties of monoterpenes and biogenically related compounds 955(2002)105

Simultaneous modeling of the Kovats retention indices on OV-1 and SE-54 stationary phases using artificial neural networks 955(2002)273

Retention mechanisms

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Retention models

Ethylenediamine as eluent component in cation chromatography. Predictive and comparative study for analysis of alkaline earth ions 955(2002)1

Theory of liquid chromatography of mono- and difunctional macromolecules. I. Studies in the critical interaction mode 955(2002)9

Retention prediction

Prediction of the retention in reversed-phase liquid chromatography using solute-mobile phase-stationary phase polarity parameters 955(2002)19

Reviews

Separation and detection of neuroactive steroids from biological matrices (Review) 955(2002)151

Sodium ion

Role of the Na⁺ ion on phenol derivatives/hydroxypropyl-β-cyclodextrin complex formation on porous graphitic carbon phase 955(2002)197

Sodium taurocholate

Chiral separation of amino acids derivatized with fluoresceine-5-isothiocyanate by capillary electrophoresis and laser-induced fluorescence detection using mixed selectors of β -cyclodextrin and sodium taurocholate 955(2002)133

Solid-phase microextraction

Gas chromatographic determination of glutaraldehyde in the workplace atmosphere after derivatization with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine on a solid-phase microextraction fibre 955(2002)117

Stationary phases, LC

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Ion-exchange separation of proteins by polyallylamine-grafted cellulose gel 955(2002)191

Steroid hormones

Separation and detection of neuroactive steroids from biological matrices (Review) 955(2002)151

Steroids

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Separation and detection of neuroactive steroids from biological matrices (Review) 955(2002)151

Sulfonvlureas

Determination of sulfonylurea herbicides by continuous-flow liquid membrane extraction on-line coupled with high-performance liquid chromatography 955(2002)183

Sulfoxides

Separation of chiral sulfoxides by liquid chromatography using macrocyclic glycopeptide chiral stationary phases 955(2002)53

Supercritical fluid extraction

Harmful azo colorants in leather. Determination based on their cleavage and extraction of corresponding carcinogenic aromatic amines using modern extraction techniques 955(2002)215

Temperature effects

Chromatographic behaviour of selected steroids and their inclusion complexes with β -cyclodextrin on octadecylsilica stationary phases with different carbon loads 955(2002)71

Terpenes

Gas chromatographic determination of vapour pressure and related thermodynamic properties of monoterpenes and biogenically related compounds 955(2002)105

Terpenoids

Screening of *Catharanthus roseus* secondary metabolites by high-performance liquid chromatography 955(2002)87

Thermodynamic parameters

Gas chromatographic determination of vapour pressure and related thermodynamic properties of monoterpenes and biogenically related compounds 955(2002)105

Trace enrichment

Determination of sulfonylurea herbicides by continuous-flow liquid membrane extraction on-line coupled with high-performance liquid chromatography 955(2002)183

Triazines

New triazine spectroscopic reagent for the separation of DLamino acids by micellar electrokinetic chromatography 955(2002)125

Trichothecenes

Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains 955(2002)245

Tröger's base

Measurement and modeling of the equilibrium behavior of the Tröger's base enantiomers on an amylose-based chiral stationary phase 955(2002)35

Vapour pressures

Gas chromatographic determination of vapour pressure and related thermodynamic properties of monoterpenes and biogenically related compounds 955(2002)105

Zearalenone

Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains 955(2002)245