

Subject Index Vol. 1044

Affinity capillary electrochromatography

Approaches to imprinted stationary phases for affinity capillary electrochromatography (Review) 1044(2004)53

Affinity chromatography with monolithic capillary columns. I. Poly-methacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Affinity chromatography

Affinity chromatography with monolithic capillary columns. I. Poly-methacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Agarose thermogels

Homogeneous reversed-phase agarose thermogels for electrochromatography 1044(2004)223

Alkylbenzenes

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Alkylthiols

On-column derivatization–capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Amines

On-column derivatization–capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Amino acids

Polystyrene monolithic column functionalized with copper–iminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

Silica sol–gel/organic hybrid material for protein encapsulated column of capillary electrochromatography 1044(2004)267

Angiotensins

Comparison of a three-peptide separation by capillary electrochromatography, voltage-assisted liquid chromatography and nano-high-performance liquid chromatography 1044(2004)201

Anilines

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Polymer-brush stationary phases for open-tubular capillary electrochromatography 1044(2004)323

Aspect ratio

Study of the influence of the aspect ratio on efficiency, flow resistance and retention factors of packed capillary columns in pressure- and electrically-driven liquid chromatography 1044(2004)311

Automation

Sample injection in capillary electrochromatography by heart-cut technique 1044(2004)259

Barbitals

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Benzenes

Influence of moderate Joule heating on electroosmotic flow velocity, retention, and efficiency in capillary electrochromatography 1044(2004)287

Benzoates

Influence of moderate Joule heating on electroosmotic flow velocity, retention, and efficiency in capillary electrochromatography 1044(2004)287

Benzoic acids

Homogeneous reversed-phase agarose thermogels for electrochromatography 1044(2004)223

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Biogenic amines

On-column derivatization–capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Caffeine

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Carbohydrates

Affinity chromatography with monolithic capillary columns. I. Poly-methacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

CEC–MS

Review coupling of capillary electrochromatography to mass spectrometry (Review) 1044(2004)131

Chip technology

Recent highlights in stationary phase design for open-tubular capillary electrochromatography (Review) 1044(2004)67

Chip electrochromatography (Review) 1044(2004)97

On-chip electrochromatography using sol-gel immobilized stationary phase with UV absorbance detection 1044(2004)277

Complexation

Capillary electrochromatography with polymeric continuous beds synthesized via free radical polymerization in aqueous media using derivatized cyclodextrins as solubilizing agents 1044(2004)211

Cyclodextrins

Capillary electrochromatography with polymeric continuous beds synthesized via free radical polymerization in aqueous media using derivatized cyclodextrins as solubilizing agents 1044(2004)211

Derivatization, electrochromatography

On-column derivatization-capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Detection, electrochromatography

On-chip electrochromatography using sol-gel immobilized stationary phase with UV absorbance detection 1044(2004)277

Dynamic field gradient focusing

Towards a miniaturised system for dynamic field gradient focused separation of proteins 1044(2004)253

Efficiency

Influence of moderate Joule heating on electroosmotic flow velocity, retention, and efficiency in capillary electrochromatography 1044(2004)287

Study of the influence of the aspect ratio on efficiency, flow resistance and retention factors of packed capillary columns in pressure- and electrically-driven liquid chromatography 1044(2004)311

Electrochromatography

Development and application of polymeric monolithic stationary phases for capillary electrochromatography (Review) 1044(2004)3

Sol-gel stationary phases for capillary electrochromatography (Review) 1044(2004)23

Approaches to imprinted stationary phases for affinity capillary electrochromatography (Review) 1044(2004)53

Recent highlights in stationary phase design for open-tubular capillary electrochromatography (Review) 1044(2004)67

Planar electrochromatography (Review) 1044(2004)83

Chip electrochromatography (Review) 1044(2004)97

Capillary electrochromatography of peptides and proteins (Review) 1044(2004)113

Review coupling of capillary electrochromatography to mass spectrometry (Review) 1044(2004)131

Capillary electrochromatography in the size-exclusion mode (Review) 1044(2004)145

Peak compression effects in capillary electrochromatography (Review) 1044(2004)153

Developments in the use of soluble ionic polymers as pseudo-stationary phases for electrokinetic chromatography and stationary phases for electrochromatography (Review) 1044(2004)159

Affinity chromatography with monolithic capillary columns. I. Poly-methacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Development of reactive thiol-modified monolithic capillaries and in-column surface functionalization by radical addition of a chromatographic ligand for capillary electrochromatography 1044(2004)187

Comparison of a three-peptide separation by capillary electrochromatography, voltage-assisted liquid chromatography and nano-high-performance liquid chromatography 1044(2004)201

Capillary electrochromatography with polymeric continuous beds synthesized via free radical polymerization in aqueous media using derivatized cyclodextrins as solubilizing agents 1044(2004)211

Homogeneous reversed-phase agarose thermogels for electrochromatography 1044(2004)223

Polystyrene monolithic column functionalized with copper-iminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Polymerised bicontinuous microemulsions as stationary phases for capillary electrochromatography. Effect of pore size on chromatographic performance 1044(2004)245

Sample injection in capillary electrochromatography by heart-cut technique 1044(2004)259

Silica sol-gel/organic hybrid material for protein encapsulated column of capillary electrochromatography 1044(2004)267

On-column derivatization-capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Analysis of ketorolac and its related impurities by capillary electrochromatography 1044(2004)295

Study of flow rate in pressurized gradient capillary electrochromatography using splitter and separation of peptides using an Amide stationary phase 1044(2004)305

Study of the influence of the aspect ratio on efficiency, flow resistance and retention factors of packed capillary columns in pressure- and electrically-driven liquid chromatography 1044(2004)311

Separation of polyethylene glycol-modified proteins by open tubular capillary electrochromatography 1044(2004)317

Electrokinetic chromatography

Developments in the use of soluble ionic polymers as pseudo-stationary phases for electrokinetic chromatography and stationary phases for electrochromatography (Review) 1044(2004)159

Electroosmosis

Homogeneous reversed-phase agarose thermogels for electrochromatography 1044(2004)223

Electroosmotic flow

Influence of moderate Joule heating on electroosmotic flow velocity, retention, and efficiency in capillary electrochromatography 1044(2004)287

Enantiomer separation

Development of reactive thiol-modified monolithic capillaries and in-column surface functionalization by radical addition of a chromatographic ligand for capillary electrochromatography 1044(2004)187

Silica sol-gel/organic hybrid material for protein encapsulated column of capillary electrochromatography 1044(2004)267

Enzymes

Separation of polyethylene glycol-modified proteins by open tubular capillary electrochromatography 1044(2004)317

Flow rate

Study of flow rate in pressurized gradient capillary electrochromatography using splitter and separation of peptides using an Amide stationary phase 1044(2004)305

Flow resistance

Study of the influence of the aspect ratio on efficiency, flow resistance and retention factors of packed capillary columns in pressure- and electrically-driven liquid chromatography 1044(2004)311

Gradient elution

Sample injection in capillary electrochromatography by heart-cut technique 1044(2004)259

Histamine

On-column derivatization-capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Hydrophilic interaction chromatography

Study of flow rate in pressurized gradient capillary electrochromatography using splitter and separation of peptides using an Amide stationary phase 1044(2004)305

Hydroxybenzoic acids

Homogeneous reversed-phase agarose thermogels for electrochromatography 1044(2004)223

Immobilized carbohydrates

Affinity chromatography with monolithic capillary columns. I. Polymethacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Injection methods

Sample injection in capillary electrochromatography by heart-cut technique 1044(2004)259

Instrumentation

Towards a miniaturised system for dynamic field gradient focused separation of proteins 1044(2004)253

On-chip electrochromatography using sol-gel immobilized stationary phase with UV absorbance detection 1044(2004)277

Interfaces

Review coupling of capillary electrochromatography to mass spectrometry (Review) 1044(2004)131

Joule heating

Influence of moderate Joule heating on electroosmotic flow velocity, retention, and efficiency in capillary electrochromatography 1044(2004)287

Ketorolac

Analysis of ketorolac and its related impurities by capillary electrochromatography 1044(2004)295

Ligand-exchange electrochromatography

Polystyrene monolithic column functionalized with copper-aminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

Mannan

Affinity chromatography with monolithic capillary columns. I. Polymethacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Mass spectrometry

Review coupling of capillary electrochromatography to mass spectrometry (Review) 1044(2004)131

Methionine enkephalin

Comparison of a three-peptide separation by capillary electrochromatography, voltage-assisted liquid chromatography and nano-high-performance liquid chromatography 1044(2004)201

Methylhistamine

On-column derivatization-capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Microfluidics

On-chip electrochromatography using sol-gel immobilized stationary phase with UV absorbance detection 1044(2004)277

Miniaturisation

Towards a miniaturised system for dynamic field gradient focused separation of proteins 1044(2004)253

Mixed-mode separations

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Molecular imprinting

Approaches to imprinted stationary phases for affinity capillary electrochromatography (Review) 1044(2004)53

Recent highlights in stationary phase design for open-tubular capillary electrochromatography (Review) 1044(2004)67

Monolithic channels

Towards a miniaturised system for dynamic field gradient focused separation of proteins 1044(2004)253

Monolithic columns

Development and application of polymeric monolithic stationary phases for capillary electrochromatography (Review) 1044(2004)3

Capillary electrochromatography of peptides and proteins (Review) 1044(2004)113

Affinity chromatography with monolithic capillary columns. I. Polymethacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Development of reactive thiol-modified monolithic capillaries and in-column surface functionalization by radical addition of a chromatographic ligand for capillary electrochromatography 1044(2004)187

Capillary electrochromatography with polymeric continuous beds synthesized via free radical polymerization in aqueous media using derivatized cyclodextrins as solubilizing agents 1044(2004)211

Polystyrene monolithic column functionalized with copper–iminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Polymerised bicontinuous microemulsions as stationary phases for capillary electrochromatography. Effect of pore size on chromatographic performance 1044(2004)245

Silica sol–gel/organic hybrid material for protein encapsulated column of capillary electrochromatography 1044(2004)267

Nano-scale liquid chromatography

Affinity chromatography with monolithic capillary columns. I. Polymethacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Oligopeptides

Polystyrene monolithic column functionalized with copper–iminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

Peak compression

Peak compression effects in capillary electrochromatography (Review) 1044(2004)153

Peptides

Development and application of polymeric monolithic stationary phases for capillary electrochromatography (Review) 1044(2004)3

Capillary electrochromatography of peptides and proteins (Review) 1044(2004)113

Comparison of a three-peptide separation by capillary electrochromatography, voltage-assisted liquid chromatography and nano-high-performance liquid chromatography 1044(2004)201

Polystyrene monolithic column functionalized with copper–iminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

On-chip electrochromatography using sol–gel immobilized stationary phase with UV absorbance detection 1044(2004)277

Study of flow rate in pressurized gradient capillary electrochromatography using splitter and separation of peptides using an Amide stationary phase 1044(2004)305

Pharmaceutical analysis

Analysis of ketorolac and its related impurities by capillary electrochromatography 1044(2004)295

Phenols

Polymer-brush stationary phases for open-tubular capillary electrochromatography 1044(2004)323

Phthalaldehyde

On-column derivatization–capillary electrochromatography with *o*-phthalaldehyde/alkylthiol for assay of biogenic amines 1044(2004)271

Planar electrochromatography

Planar electrochromatography (Review) 1044(2004)83

Poly(2-hydroxyethyl methacrylate)

Polymer-brush stationary phases for open-tubular capillary electrochromatography 1044(2004)323

Polymers

Capillary electrochromatography in the size-exclusion mode (Review) 1044(2004)145

Developments in the use of soluble ionic polymers as pseudo-stationary phases for electrokinetic chromatography and stationary phases for electrochromatography (Review) 1044(2004)159

Polynuclear aromatic hydrocarbons

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Polysaccharides

Capillary electrochromatography in the size-exclusion mode (Review) 1044(2004)145

Polystyrene

Polystyrene monolithic column functionalized with copper–iminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

Pore size

Polymerised bicontinuous microemulsions as stationary phases for capillary electrochromatography. Effect of pore size on chromatographic performance 1044(2004)245

Proteins

Development and application of polymeric monolithic stationary phases for capillary electrochromatography (Review) 1044(2004)3

Capillary electrochromatography of peptides and proteins (Review) 1044(2004)113

Capillary electrochromatography in the size-exclusion mode (Review) 1044(2004)145

Affinity chromatography with monolithic capillary columns. I. Polymethacrylate monoliths with immobilized mannan for the separation of mannose-binding proteins by capillary electrochromatography and nano-scale liquid chromatography 1044(2004)177

Towards a miniaturised system for dynamic field gradient focused separation of proteins 1044(2004)253

Separation of polyethylene glycol-modified proteins by open tubular capillary electrochromatography 1044(2004)317

Pseudo-stationary phases

Developments in the use of soluble ionic polymers as pseudo-stationary phases for electrokinetic chromatography and stationary phases for electrochromatography (Review) 1044(2004)159

Retention factor

Influence of moderate Joule heating on electroosmotic flow velocity, retention, and efficiency in capillary electrochromatography 1044(2004)287

Study of the influence of the aspect ratio on efficiency, flow resistance and retention factors of packed capillary columns in pressure- and electrically-driven liquid chromatography 1044(2004)311

Reviews

Development and application of polymeric monolithic stationary phases for capillary electrochromatography (Review) 1044(2004)3

Sol-gel stationary phases for capillary electrochromatography (Review) 1044(2004)23

Approaches to imprinted stationary phases for affinity capillary electrochromatography (Review) 1044(2004)53

Recent highlights in stationary phase design for open-tubular capillary electrochromatography (Review) 1044(2004)67

Planar electrochromatography (Review) 1044(2004)83

Chip electrochromatography (Review) 1044(2004)97

Capillary electrochromatography of peptides and proteins (Review) 1044(2004)113

Review coupling of capillary electrochromatography to mass spectrometry (Review) 1044(2004)131

Peak compression effects in capillary electrochromatography (Review) 1044(2004)153

Developments in the use of soluble ionic polymers as pseudo-stationary phases for electrokinetic chromatography and stationary phases for electrochromatography (Review) 1044(2004)159

Saccharides

Development and application of polymeric monolithic stationary phases for capillary electrochromatography (Review) 1044(2004)3

Sample handling

Sample injection in capillary electrochromatography by heart-cut technique 1044(2004)259

Size-exclusion chromatography

Capillary electrochromatography in the size-exclusion mode (Review) 1044(2004)145

Size-exclusion electrochromatography

Capillary electrochromatography in the size-exclusion mode (Review) 1044(2004)145

Sol-gel

Sol-gel stationary phases for capillary electrochromatography (Review) 1044(2004)23

Silica sol-gel/organic hybrid material for protein encapsulated column of capillary electrochromatography 1044(2004)267

On-chip electrochromatography using sol-gel immobilized stationary phase with UV absorbance detection 1044(2004)277

Stationary phases, electrochromatography

Development and application of polymeric monolithic stationary phases for capillary electrochromatography (Review) 1044(2004)3

Sol-gel stationary phases for capillary electrochromatography (Review) 1044(2004)23

Approaches to imprinted stationary phases for affinity capillary electrochromatography (Review) 1044(2004)53

Recent highlights in stationary phase design for open-tubular capillary electrochromatography (Review) 1044(2004)67

Chip electrochromatography (Review) 1044(2004)97

Developments in the use of soluble ionic polymers as pseudo-stationary phases for electrokinetic chromatography and stationary phases for electrochromatography (Review) 1044(2004)159

Development of reactive thiol-modified monolithic capillaries and in-column surface functionalization by radical addition of a chromatographic ligand for capillary electrochromatography 1044(2004)187

Homogeneous reversed-phase agarose thermogels for electrochromatography 1044(2004)223

Polystyrene monolithic column functionalized with copper-iminodiacetate complex as a stationary phase for open tubular capillary electrochromatography 1044(2004)229

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Polymerised bicontinuous microemulsions as stationary phases for capillary electrochromatography. Effect of pore size on chromatographic performance 1044(2004)245

Silica sol-gel/organic hybrid material for protein encapsulated column of capillary electrochromatography 1044(2004)267

On-chip electrochromatography using sol-gel immobilized stationary phase with UV absorbance detection 1044(2004)277

Polymer-brush stationary phases for open-tubular capillary electrochromatography 1044(2004)323

Terephthalic acid

Monolithic columns with mixed modes of reversed-phase and anion-exchange stationary phase for capillary electrochromatography 1044(2004)237

Thermodynamic parameters

Polymerised bicontinuous microemulsions as stationary phases for capillary electrochromatography. Effect of pore size on chromatographic performance 1044(2004)245

Thermogels

Homogeneous reversed-phase agarose thermogels for electrochromatography 1044(2004)223

Tryptophan

Silica sol-gel/organic hybrid material for protein encapsulated column of capillary electrochromatography 1044(2004)267

Voltage-assisted liquid chromatography

Comparison of a three-peptide separation by capillary electrochromatography, voltage-assisted liquid chromatography and nano-high-performance liquid chromatography 1044(2004)201

Wall effects

Study of the influence of the aspect ratio on efficiency, flow resistance and retention factors of packed capillary columns in pressure- and electrically-driven liquid chromatography 1044(2004)311