

# Available online at www.sciencedirect.com



JOURNAL OF CHROMATOGRAPHY A

Journal of Chromatography A 1086 (2005) 195-199

www.elsevier.com/locate/chroma

# Subject Index Vol. 1086

#### APCI-MS

Silver-ion reversed-phase comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection in lipidic food analysis 1086(2005)91

# Aromatic compounds

Analysis of aromatic compounds in gasoline with flow-switching comprehensive two-dimensional gas chromatography 1086(2005)115

#### **ASTM** methods

Characterisation of middle-distillates by comprehensive two-dimensional gas chromatography (GC  $\times$  GC): A powerful alternative for performing various standard analysis of middle-distillates 1086(2005)21

#### **Butter extracts**

Improved determination of flavour compounds in butter by solid-phase (micro)extraction and comprehensive two-dimensional gas chromatography 1086(2005)99

# Chemometrics

Evaluation of the DotMap algorithm for locating analytes of interest based on mass spectral similarity in data collected using comprehensive two-dimensional gas chromatography coupled with time-of-flight mass spectrometry 1086(2005)185

#### Chlorinated paraffins

Characterization of polychlorinated *n*-alkanes using comprehensive two-dimensional gas chromatography–electron-capture negative ionisation time-of-flight mass spectrometry 1086(2005)71

# Column combinations

Twin comprehensive two-dimensional gas chromatographic system: concept and applications 1086(2005)128

#### Column dimensions

Flow regime at ambient outlet pressure and its influence in comprehensive two-dimensional gas chromatography 1086(2005)141

#### Column flow

Flow model for coupled-column gas chromatography systems 1086(2005)135

# Compound identification

Peak pattern variations related to comprehensive two-dimensional gas chromatography acquisition 1086(2005)165

# Comprehensive two-dimensional chromatography

Analysis of aromatic compounds in gasoline with flow-switching comprehensive two-dimensional gas chromatography 1086(2005)115

Method for reducing the ambiguity of comprehensive two-dimensional chromatography retention times 1086(2005)171

# Comprehensive two-dimensional gas chromatography (GC $\times$ GC)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

#### Comprehensive two-dimensional gas chromatography ( $GC \times GC$ )

Peak pattern variations related to comprehensive two-dimensional gas chromatography acquisition 1086(2005)165

# Comprehensive two-dimensional gas chromatography

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

Trace analysis of polychlorinated dibenzo-*p*-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

Characterization of polychlorinated *n*-alkanes using comprehensive two-dimensional gas chromatography–electron-capture negative ionisation time-of-flight mass spectrometry 1086(2005)71

Improved determination of flavour compounds in butter by solid-phase (micro)extraction and comprehensive two-dimensional gas chromatography 1086(2005)99

Quantitative determination of compounds in tobacco essential oils by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry 1086(2005)107

Twin comprehensive two-dimensional gas chromatographic system: concept and applications 1086(2005)128

Flow model for coupled-column gas chromatography systems 1086(2005)135

Deconvolution of overlapped peaks based on the exponentially modified Gaussian model in comprehensive two-dimensional gas chromatography 1086(2005)160

Resolution prediction and optimization of temperature programme in comprehensive two-dimensional gas chromatography 1086(2005)175

# Comprehensive two-dimensional liquid chromatography

Silver-ion reversed-phase comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection in lipidic food analysis 1086(2005)91

# Comprehensive two-dimensional

Characterization of lipids in complex samples using comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry 1086(2005)2

doi:10.1016/S0021-9673(05)01449-4

#### **Curve fitting**

Deconvolution of overlapped peaks based on the exponentially modified Gaussian model in comprehensive two-dimensional gas chromatography 1086(2005)160

#### Data analysis

Evaluation of the DotMap algorithm for locating analytes of interest based on mass spectral similarity in data collected using comprehensive two-dimensional gas chromatography coupled with time-of-flight mass spectrometry 1086(2005)185

# Data mining

Evaluation of the DotMap algorithm for locating analytes of interest based on mass spectral similarity in data collected using comprehensive two-dimensional gas chromatography coupled with time-of-flight mass spectrometry 1086(2005)185

#### Data processing

Statistical methods for comparing comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry results: Metabolomic analysis of mouse tissue extracts 1086(2005)83

#### Deconvolution

Deconvolution of overlapped peaks based on the exponentially modified Gaussian model in comprehensive two-dimensional gas chromatography 1086(2005)160

# Dioxin-responsive chemically activated luciferase gene expression (DR-CALUX)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

# **Dioxins**

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

# μECD

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

# ECNI-TOF-MS

Characterization of polychlorinated *n*-alkanes using comprehensive two-dimensional gas chromatography–electron-capture negative ionisation time-of-flight mass spectrometry 1086(2005)71

# EMG

Deconvolution of overlapped peaks based on the exponentially modified Gaussian model in comprehensive two-dimensional gas chromatography 1086(2005)160

#### environmental pollutants

Dynamic ultrasound-assisted extraction of environmental pollutants from marine sediments for comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometric detection 1086(2005)122

#### **Essential oils**

Quantitative determination of compounds in tobacco essential oils by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry 1086(2005)107

#### Flavour compounds

Improved determination of flavour compounds in butter by solid-phase (micro)extraction and comprehensive two-dimensional gas chromatography 1086(2005)99

#### Flow-switching modulation

Analysis of aromatic compounds in gasoline with flow-switching comprehensive two-dimensional gas chromatography 1086(2005)115

#### Food extracts

Twin comprehensive two-dimensional gas chromatographic system: concept and applications 1086(2005)128

#### Food

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

#### Gas chromatography, comprehensive two-dimensional

Statistical methods for comparing comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry results: Metabolomic analysis of mouse tissue extracts 1086(2005)83

#### Gas chromatography

Characterization of lipids in complex samples using comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry 1086(2005)2

Band acceleration device for enhanced selectivity with tandem-column gas chromatography 1086(2005)151

Deconvolution of overlapped peaks based on the exponentially modified Gaussian model in comprehensive two-dimensional gas chromatography 1086(2005)160

# Gas chromatography-mass spectrometry

Quantitative determination of compounds in tobacco essential oils by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry 1086(2005)107

# Gasoline analysis

Analysis of aromatic compounds in gasoline with flow-switching comprehensive two-dimensional gas chromatography 1086(2005)115

#### $GC \times GC$

Characterization of polychlorinated *n*-alkanes using comprehensive two-dimensional gas chromatography–electron-capture negative ionisation time-of-flight mass spectrometry 1086(2005)71

Improved determination of flavour compounds in butter by solid-phase (micro)extraction and comprehensive two-dimensional gas chromatography 1086(2005)99

Flow model for coupled-column gas chromatography systems 1086(2005)135

Flow regime at ambient outlet pressure and its influence in comprehensive two-dimensional gas chromatography 1086(2005)141

# $GC \times GC$ -TOF-MS

Dynamic ultrasound-assisted extraction of environmental pollutants from marine sediments for comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometric detection 1086(2005)122

Evaluation of the DotMap algorithm for locating analytes of interest based on mass spectral similarity in data collected using comprehensive two-dimensional gas chromatography coupled with time-of-flight mass spectrometry 1086(2005)185

# GC-HRMS

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

#### $GC \times GC$

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

#### Group type separation

Characterisation of middle-distillates by comprehensive two-dimensional gas chromatography (GC  $\times$  GC): A powerful alternative for performing various standard analysis of middle-distillates 1086(2005)21

# High-resolution mass spectrometry (HRMS)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

#### Hydrocarbons

Characterisation of middle-distillates by comprehensive two-dimensional gas chromatography (GC  $\times$  GC): A powerful alternative for performing various standard analysis of middle-distillates 1086(2005)21

#### Lipids

Characterization of lipids in complex samples using comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry 1086(2005)2

# Loop-type modulator

Flow model for coupled-column gas chromatography systems 1086(2005)135

# Marine sediment

Dynamic ultrasound-assisted extraction of environmental pollutants from marine sediments for comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometric detection 1086(2005)122

# Mass spectrometry, time-of-flight

Statistical methods for comparing comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry results: Metabolomic analysis of mouse tissue extracts 1086(2005)83

#### Metabolomics

Statistical methods for comparing comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry results: Metabolomic analysis of mouse tissue extracts 1086(2005)83

# Modeling

Band acceleration device for enhanced selectivity with tandem-column gas chromatography 1086(2005)151

#### **Modulation criterion**

Flow regime at ambient outlet pressure and its influence in comprehensive two-dimensional gas chromatography 1086(2005)141

# Multi-dimensional chromatographic studies

Comprehensive multi-dimensional chromatographic studies on the separation of saturated hydrocarbon ring structures in petrochemical samples 1086(2005)12

#### Multiple internal standards calibration

Quantitative determination of compounds in tobacco essential oils by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry 1086(2005)107

#### **Optimization**

Resolution prediction and optimization of temperature programme in comprehensive two-dimensional gas chromatography 1086(2005)175

#### **Optimum flow**

Flow regime at ambient outlet pressure and its influence in comprehensive two-dimensional gas chromatography 1086(2005)141

# Organobromine compounds

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

#### Organochlorine compounds

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

#### Organohalogenated pesticides

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

#### Orthogonality

Twin comprehensive two-dimensional gas chromatographic system: concept and applications 1086(2005)128

# Overlapped peak

Deconvolution of overlapped peaks based on the exponentially modified Gaussian model in comprehensive two-dimensional gas chromatography 1086(2005)160

#### **PCB**

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

# PCDD/F

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

#### Peak pattern matching

Peak pattern variations related to comprehensive two-dimensional gas chromatography acquisition 1086(2005)165

# Peak pattern variation

Peak pattern variations related to comprehensive two-dimensional gas chromatography acquisition 1086(2005)165

# Petrochemical samples

Comprehensive multi-dimensional chromatographic studies on the separation of saturated hydrocarbon ring structures in petrochemical samples 1086(2005)12

# Polybrominated biphenyls

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Polybrominated diphenyl ethers

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Polychlorinated alkanes

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

Characterization of polychlorinated *n*-alkanes using comprehensive two-dimensional gas chromatography–electron-capture negative ionisation time-of-flight mass spectrometry 1086(2005)71

# Polychlorinated biphenyls (PCBs)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

# Polychlorinated biphenyls

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Polychlorinated dibenzofurans (PCDFs)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

#### Polychlorinated dibenzofurans

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

#### Polychlorinated dibenzo-p-dioxins (PCDDs)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

#### Polychlorinated dibenzo-p-dioxins

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Polychlorinated dibenzothiophenes

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Polychlorinated diphenyl ethers

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Polychlorinated naphthalenes

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Polychlorinated terphenyls

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Quadrupole ion storage mass spectrometry (QIST-MS)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

### Quantification

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

# **Resolution prediction**

Resolution prediction and optimization of temperature programme in comprehensive two-dimensional gas chromatography 1086(2005)175

#### Resolution

Flow regime at ambient outlet pressure and its influence in comprehensive two-dimensional gas chromatography 1086(2005)141

#### **Retention time analysis**

Method for reducing the ambiguity of comprehensive two-dimensional chromatography retention times 1086(2005)171

# Sample capacity

Flow regime at ambient outlet pressure and its influence in comprehensive two-dimensional gas chromatography 1086(2005)141

# Sample dimensionality

Comprehensive multi-dimensional chromatographic studies on the separation of saturated hydrocarbon ring structures in petrochemical samples 1086(2005)12

# Saturated hydrocarbon ring structures

Comprehensive multi-dimensional chromatographic studies on the separation of saturated hydrocarbon ring structures in petrochemical samples 1086(2005)12

#### Selectivity

Evaluation of the DotMap algorithm for locating analytes of interest based on mass spectral similarity in data collected using comprehensive two-dimensional gas chromatography coupled with time-of-flight mass spectrometry 1086(2005)185

#### Simulated distillation

Characterisation of middle-distillates by comprehensive two-dimensional gas chromatography (GC × GC): A powerful alternative for performing various standard analysis of middle-distillates 1086(2005)21

#### SPE

Improved determination of flavour compounds in butter by solid-phase (micro)extraction and comprehensive two-dimensional gas chromatography 1086(2005)99

# **SPME**

Improved determination of flavour compounds in butter by solid-phase (micro)extraction and comprehensive two-dimensional gas chromatography 1086(2005)99

# TAG

Silver-ion reversed-phase comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection in lipidic food analysis 1086(2005)91

# Tandem-in-time mass spectrometry (MS/MS)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

#### TEF

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

# Temperature program

Resolution prediction and optimization of temperature programme in comprehensive two-dimensional gas chromatography 1086(2005)175

# **Temperature-programming**

Band acceleration device for enhanced selectivity with tandem-column gas chromatography 1086(2005)151

### TEO

Trace analysis of polychlorinated dibenzo-p-dioxins, dibenzofurans and WHO polychlorinated biphenyls in food using comprehensive two-dimensional gas chromatography with electron-capture detection 1086(2005)61

# Time-of-flight mass spectrometry (TOF-MS)

Comprehensive two-dimensional gas chromatography with isotope dilution time-of-flight mass spectrometry for the measurement of dioxins and polychlorinated biphenyls in foodstuffs. Comparison with other methods 1086(2005)45

#### Time-of-flight mass spectrometry

Characterization of lipids in complex samples using comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry 1086(2005)2

Quantitative determination of compounds in tobacco essential oils by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry 1086(2005)107

#### Tobacco

Quantitative determination of compounds in tobacco essential oils by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry 1086(2005)107

#### **Toxaphene**

Group separation of organohalogenated compounds by means of comprehensive two-dimensional gas chromatography 1086(2005)29

# Transformation model

Peak pattern variations related to comprehensive two-dimensional gas chromatography acquisition 1086(2005)165

#### Triacylglycerols

Silver-ion reversed-phase comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection in lipidic food analysis 1086(2005)91

#### Twin system

Twin comprehensive two-dimensional gas chromatographic system: concept and applications 1086(2005)128

# Ultrasound-assisted extraction

Dynamic ultrasound-assisted extraction of environmental pollutants from marine sediments for comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometric detection 1086(2005)122

#### Wrap-around

Method for reducing the ambiguity of comprehensive two-dimensional chromatography retention times 1086(2005)171