

Tetrakis(triphenylarsine oxide)copper(II) Bis[dichlorocuprate(I)]

BY R. H. P. FRANCISCO, R. H. DE ALMEIDA SANTOS AND J. R. LECHAT

Instituto de Física e Química de São Carlos, Universidade de São Paulo, CP 369, 13560 São Carlos, São Paulo, Brazil

AND A. C. MASSABNI

Instituto de Química de Araraquara, UNESP, CP 174, 14800 Araraquara, São Paulo, Brazil

(Received 23 May 1980; accepted 19 August 1980)

Abstract. $[\text{Cu}(\text{C}_{18}\text{H}_{15}\text{AsO})_4]^{2+} \cdot 2[\text{CuCl}_2]^-$, monoclinic, $P2_1/c$, $a = 13.315$ (2), $b = 20.162$ (4), $c = 17.761$ (2) Å, $\beta = 134.25$ (3)°, $V = 3415$ (1) Å³, $Z = 2$, $D_m = 1.59$ (floatation), $D_x = 1.577$ Mg m⁻³. The final R was 0.045 for 3357 observed reflections. The Cu^{II} is coordinated to four O in a square-planar arrangement and the Cu^I is coordinated to two Cl in a linear structure.

Introduction. The preparation and the spectral properties of a yellow-brown complex with empirical formula $[\text{Cu}(\text{tpa}_5\text{O})_2\text{Cl}_2]$ ($\text{tpa}_5\text{O} = \text{triphenylarsine oxide}$) have been reported (Goodgame & Cotton, 1961; Rodley, Goodgame & Cotton, 1965). In an attempt to prepare this compound according to the procedure described by Goodgame & Cotton a yellow-brown complex was in fact obtained, but it analysed as $2\text{CuCl}_2 \cdot 3(\text{tpa}_5\text{O})$. On the other hand, recrystallization of this complex from anhydrous acetone yields blue crystals of a new complex with empirical formula $\text{Cu}_3\text{Cl}_4 \cdot 4(\text{tpa}_5\text{O})$. A complex with the same colour and empirical formula was studied by Nyholm (1951) starting from CuCl_2 and methyltriphenylarsine. The complex was formulated as $[\text{Cu}(\text{mdpa}_5\text{O})_4][\text{CuCl}_2]_2$, where mdpa_5O is methyltriphenylarsine oxide, suggesting that copper would be present as Cu^I and Cu^{II}. Although this formulation seems plausible and could also apply to the $\text{Cu}_3\text{Cl}_4 \cdot 4(\text{tpa}_5\text{O})$ complex it was felt that more conclusive evidence should be obtained concerning the copper coordination. Hence, an X-ray investigation of the latter complex was undertaken to elucidate its structural situation and to confirm the presence of copper in two oxidation states.

A crystal $0.2 \times 0.3 \times 0.15$ mm was mounted at random on an automatic CAD-4 diffractometer. Cell dimensions were determined by a least-squares fit of settings for 25 reflections ($\pm hkl$). Intensity measurements were carried out up to $\theta = 75^\circ$ using the $\omega/2\theta$ scan mode and graphite-monochromated Cu $K\alpha$ radiation. 4345 unique reflections were measured, of

which 3357 were considered observed [$I > 3\sigma(I)$] and retained for use in the structure analysis.

Lorentz and polarization corrections were applied but no absorption correction [$\mu(\text{Cu } K\alpha) = 4.02 \text{ mm}^{-1}$] was made.

The structure was solved by application of the multi-solution weighted tangent formula (Germain, Main & Woolfson, 1971) to the reflections with $E > 1.70$. An E map based on the solution with the highest absolute figure of merit (1.008) and the lowest residual index (25.88) revealed the positions of one Cu and two As atoms.

The remaining non-hydrogen atoms of the organic-ligand molecules were located by successive difference-Fourier calculations. At this point, a difference-Fourier synthesis unambiguously showed the dichlorocuprate ion.

The refinement was by minimization of $\sum w(k|F_o| - |F_c|)^2$ through iterative full-matrix least-squares calculations including all data. Weights $w_i = [\sigma_{oi}^2 + (0.05F_{oi}^2)^2]^{-1}$ (where σ_{oi}^2 was the variance based on counting statistics) were assigned to observed reflections ($w_i = 0$ for unobserved).

The refinement was conducted until all atomic-parameter shifts were smaller than their standard deviations. The final unweighted R factor omitting unobserved reflections was 0.045, and including them, 0.048.*

Anisotropic temperature factors were assigned to all non-hydrogen atoms. Neutral-atom complex scattering factors were used (*International Tables for X-ray Crystallography*, 1974). Table 1 gives the heavy-atom coordinates.

* Lists of structure factors, anisotropic thermal parameters and bond distances and angles in the organic-ligand molecule have been deposited with the British Library Lending Division as Supplementary Publication No. SUP 35598 (22 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

Table 1. Fractional atomic coordinates ($\times 10^5$ for As and Cu, $\times 10^4$ for Cl, O and C) with their e.s.d.'s in parentheses

The B_{eq} values are the equivalent isotropic temperature factors (Hamilton, 1959).

	x	y	z	B_{eq} (Å ²)
As(1)	11510 (7)	5075 (5)	21946 (5)	2.4
As(2)	27997 (7)	5764 (5)	6625 (5)	2.3
Cu(1)	0	0	0	2.2
Cu(2)	80410 (10)	16588 (8)	33116 (9)	4.5
Cl(1)	9579 (2)	933 (1)	4212 (2)	5.2
Cl(2)	6499 (3)	2380 (2)	2382 (2)	7.8
O(1)	1426 (4)	277 (3)	1445 (3)	2.7
O(2)	1271 (4)	151 (3)	-142 (3)	2.6
C(1)	2854 (7)	924 (5)	3450 (5)	3.2
C(2)	2762 (8)	1494 (5)	3841 (6)	4.0
C(3)	4010 (9)	1771 (5)	4759 (6)	4.9
C(4)	5290 (9)	1493 (5)	5246 (6)	4.8
C(5)	5367 (8)	922 (6)	4830 (6)	4.8
C(6)	4128 (7)	625 (5)	3918 (6)	3.7
C(7)	-301 (7)	1172 (4)	1469 (5)	2.6
C(8)	-1000 (7)	1331 (5)	1773 (6)	3.7
C(9)	-2035 (8)	1831 (5)	1203 (7)	4.5
C(10)	-2368 (9)	2134 (5)	353 (7)	5.0
C(11)	-1643 (9)	1965 (5)	61 (7)	4.8
C(12)	-598 (8)	1477 (5)	626 (6)	3.7
C(13)	844 (7)	-254 (4)	2664 (5)	2.6
C(14)	927 (8)	-878 (5)	2380 (6)	3.8
C(15)	734 (10)	-1428 (5)	2747 (7)	5.5
C(16)	459 (9)	-1348 (6)	3365 (7)	5.8
C(17)	402 (8)	-722 (6)	3645 (6)	5.2
C(18)	618 (7)	-165 (5)	3307 (5)	3.9
C(19)	4235 (7)	75 (4)	1916 (5)	2.7
C(20)	3908 (7)	-555 (4)	2005 (5)	3.1
C(21)	4968 (8)	-935 (5)	2902 (6)	4.4
C(22)	6337 (8)	-674 (5)	3678 (6)	4.6
C(23)	6658 (8)	-34 (6)	3559 (6)	4.7
C(24)	5588 (7)	344 (5)	2654 (6)	3.4
C(25)	3294 (6)	681 (4)	-116 (5)	2.4
C(26)	3993 (7)	1252 (5)	9 (6)	3.8
C(27)	4302 (8)	1320 (6)	-589 (6)	5.1
C(28)	3921 (7)	831 (6)	-1304 (6)	5.0
C(29)	3245 (8)	251 (6)	-1404 (6)	5.1
C(30)	2939 (7)	180 (5)	-795 (5)	3.9
C(31)	2619 (6)	1456 (4)	982 (5)	2.9
C(32)	1711 (8)	1873 (5)	125 (7)	5.0
C(33)	1527 (9)	2515 (6)	300 (8)	6.3
C(34)	2248 (9)	2725 (5)	1303 (7)	6.2
C(35)	3195 (10)	2311 (6)	2155 (7)	6.4
C(36)	3392 (8)	1651 (5)	1993 (5)	4.3

Discussion. The structure consists of discrete [Cu^{II}{OAs(C₆H₅)₃]₄]²⁺ cations and [Cu^ICl₂]⁻ anions, as can be seen in Fig. 1.

The Cu^{II} ions are located at centres of inversion. Four organic ligands are coordinated to the metal, *via* oxygen, in a square-planar geometry, according to symmetry conditions. The rings of the organic part are planar within standard deviations. All distances and angles are normal. The most important bond distances and angles are indicated in Fig. 2.




Fig. 1. Representation of the molecule giving atomic numbering.




Fig. 2. Representation of the environment of the Cu^{II} ion with bond lengths (Å) and angles (°).

The dichlorocuprate ion is practically linear. The Cu—Cl distance is 2.079 (2) Å for the two bonds and the central angle is 178.65 (8)°.

The occurrence of this kind of ion in the solid state has been definitely established in chlorododekakis(dimethylamino)cyclohexa(phosphazene)-*N*¹,*N*²,-*N*⁴,*N*⁵]copper(II) dichlorocuprate(I) where the Cu—Cl bond length is 2.06 (1) Å and the Cl—Cu—Cl angle is 179.5 (9)° (Marsh & Trotter, 1971).

All computer calculations were performed using the Enraf–Nonius structure determination package. The figures were drawn with the *ORTEP* program (Johnson, 1965).

This work was sponsored by grants from BID-FINEP, CNPq and FAPESP.

References

- GERMAIN, G., MAIN, P. & WOOLFSON, M. M. (1971). *Acta Cryst.* **A27**, 368–376.
- GOODGAME, D. M. L. & COTTON, F. A. (1961). *J. Chem. Soc.* pp. 2298–2304.
- HAMILTON, W. C. (1959). *Acta Cryst.* **12**, 609–610.
- International Tables for X-ray Crystallography* (1974). Vol. IV. Birmingham: Kynoch Press.
- JOHNSON, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee.
- MARSH, W. C. & TROTTER, J. (1971). *J. Chem. Soc. A*, pp. 1482–1486.
- NYHOLM, R. S. (1951). *J. Chem. Soc.* pp. 1767–1774.
- RODLEY, G. A., GOODGAME, D. M. L. & COTTON, F. A. (1965). *J. Chem. Soc.* pp. 1499–1505.

Acta Cryst. (1981). **B37**, 234–237

Cyano[3,3'-(1,3-propanediamino)bis(3-methyl-2-butanone oximato)]copper(II)– ½ Acetonitrile

BY E. O. SCHLEMPER, M. SAKHAWAT HUSSAIN AND R. KENT MURMANN

Department of Chemistry, University of Missouri, Columbia, Missouri 65211, USA

(Received 1 April 1980; accepted 19 August 1980)

Abstract. [Cu(C₁₃H₂₇N₄O₂)CN] · ½ C₂H₃N, [Cu(CN)(pnao-H)] · ½ CH₃CN, orthorhombic, *Fdd2*, *a* = 18.993 (2), *b* = 32.352 (2), *c* = 11.799 (1) Å, *V* = 7250.0 Å³, *M_r* = 381.48, *Z* = 16, *ρ_{calc}* = 1.40 (1), *ρ_{obs}* = 1.41 (1) Mg m⁻³. Three-dimensional counter X-ray data were used to solve and refine the structure to a final *R(F²)* of 0.054 for 2600 independent reflections. The structure consists of discrete monomeric square-pyramidal Cu complex molecules with a strong intramolecular hydrogen bond. The α-amine-oxime ligand is quadridentate with an average Cu–N(amine) distance = 2.043 (3) Å and Cu–N(oxime) = 1.978 (3) Å. The cyanide ion is carbon bonded at the apical position with a Cu–C bond distance of 2.154 (5) Å. The strong Cu–CN bond pulls the Cu²⁺ ion out of the plane of the four N atoms by 0.499 (6) Å. The short intramolecular hydrogen bond of 2.475 (4) Å is not appreciably modified by the presence of CN⁻. The acetonitrile molecule is not coordinated to the Cu^{II} atom, but packs between the Cu complexes as solvent of crystallization.

Introduction. The deep-purple crystals (Murmman & Monzyk, 1978) were consistently twinned when grown from aqueous media but gave satisfactory crystals from anhydrous acetonitrile. They crystallize with primarily {010}, {011}, and {110} facial development. The crystal selected was semi-spherical and had a volume of about 0.023 mm³. The space group and cell dimensions were first determined by precession film data, and the final cell dimensions were obtained with a Picker diffractometer from a least-squares refinement of the

setting angles of 18 manually centered reflections. Intensity data were collected for 4094 reflections by the *θ*–2*θ* scan method using Mo *K*α radiation (*λ* = 0.7107 Å) filtered with Nb. The data were corrected for background, *L_p* and absorption (*μ* = 1.27 mm⁻¹)*. The transmission factor ranged from 0.89 to 0.73. Standards monitored about every 75 reflections showed no significant systematic variation with time.

The structure was solved by Patterson and Fourier methods. The acetonitrile was revealed by a difference Fourier map and its occupancy factor refined to 0.5. After a few cycles of least-squares refinement with all non-hydrogen atoms present, a difference Fourier map revealed most of the H atoms. When ambiguities occurred, idealized positions for these H atoms were calculated and were found to be consistent with difference Fourier peaks. Refinement was made using anisotropic temperature factors for all non-hydrogen atoms, with fixed H atoms. H atoms on the CH₃CN molecule were treated as disordered around the twofold axis. Full-matrix refinement minimizing $\sum w(F_o^2 - kF_c^2)^2$, using 2600 independent reflections, 213 variables, scattering factors of Cromer & Waber (1962), and anomalous scattering for Cu (Cromer, 1962), gave

* All calculations were performed on the University of Missouri IBM 370/168 computer system with the following programs: *ANGSET* (angle settings program); *SORTH* (sorting program; W. C. Hamilton); *HORSE* (general absorption-correction programs; W. C. Hamilton); *FORDAP* (Fourier synthesis program; A. Zalkin); *NUCLS* (least-squares program; R. Doedens and J. A. Ibers); a modification of *ORFLS* (Busing, Martin & Levy, 1962); *ORFFE* (Busing, Martin & Levy, 1964); *ORTEP* (Johnson, 1965).