

References

- AHMED, F. R., HALL, S. R., PIPPY, M. E. & HUBER, C. P. (1966). NRC Crystallographic Programs, Ottawa, variously modified for use on the Univ. of Aberdeen Honeywell 66/80 computer by R. A. HOWIE, J. S. KNOWLES & H. F. W. TAYLOR.
- BLAKE, A. J. (1980). Ph.D. Thesis, Aberdeen Univ.
- BLAKE, A. J., HOWIE, R. A. & McQUILLAN, G. P. (1979). *J. Appl. Cryst.* **12**, 414–415.
- BLAKE, A. J., HOWIE, R. A. & McQUILLAN, G. P. (1981). *Acta Cryst.* **B37**, 997–999.
- CAMERON, T. S. & DAHLÉN, B. (1975). *J. Chem. Soc. Perkin Trans. 2*, pp. 1737–1751.
- DREISSIG, W., PLIETH, K. & ZÄSKE, P. (1972). *Acta Cryst.* **B28**, 3473–3483.
- International Tables for X-ray Crystallography* (1974). Vol. IV. Birmingham: Kynoch Press.
- LEE, J. D. & GOODACRE, G. W. (1969). *Acta Cryst.* **B25**, 2127–2131.
- LEE, J. D. & GOODACRE, G. W. (1970). *Acta Cryst.* **B26**, 507–515.
- LEE, J. D. & GOODACRE, G. W. (1971). *Acta Cryst.* **B27**, 302–307.
- MACHIN, P. A. & ELDER, M. (1977). *Notes for Weissenberg and Precession Film Processing*. SRC Daresbury Laboratory.
- WILKINS, C. J., HAGEN, K., HEDBERG, L., QUANG SHEN & HEDBERG, K. (1975). *J. Am. Chem. Soc.* **97**, 6352–6358.

SHORT COMMUNICATION

Contributions intended for publication under this heading should be expressly so marked; they should not exceed about 1000 words; they should be forwarded in the usual way to the appropriate Co-editor; they will be published as speedily as possible.

Acta Cryst. (1981), **B37**, 1962

X-ray structure analysis and molecular conformation of *N*-propionylproline: erratum. By M. E. KAMWAYA,* O. OSTER and H. BRADACZEK. *Institut für Kristallographie, Freie Universität Berlin, Takustrasse 6, D-1000 Berlin 33, Federal Republic of Germany*

(Received 23 July 1981)

Abstract

Corrections are given to the paper by Kamwaya, Oster & Bradaczek [*Acta Cryst.* (1981), **B37**, 364–367]. In Table 2

the atomic nomenclature for three entries requires correction: the correct entries are N'–C₄^α–C₅^β 101.8 (3), O₃–C₈^γ–O₂ 124.0 (3)°, C₄^α–C₅^β 1.527 (5) Å. In Table 3 the torsion angle C₃'–N'–C₄^α–C₅^β should be designated ω . See also *Acta Cryst.* (1981), **B37**, 1161.

* Present address: School of Physics, Universiti Sains Malaysia, Minden, Penang, Malaysia.

All the relevant information is given in the *Abstract*.