Subject Index

Acylaminoethylarylsulphonylurea TLC determination of acylaminoethylarylsulphonylureas in serum after hydrolysis and dansylation 533

Adrenaline Quantitation of adrenaline and noradrenaline from human plasma by combined GC—high-resolution mass fragmentography 221

Adrenal steroids

Covenient system for the simultaneous separation of 11-deoxycortisol and aldosterone by Sephadex LH-20 multiple column chromatography 333

Albumin

Electrophoretic bimorphism of serum albumin in the presence of indocyanine green 77

Aldosterone

Convenient system for the simultaneous separation of 11-deoxycortisol and aldosterone by Sephadex LH-20 multiple column chromatography 333

Aldosterone

Determination of aldosterone in urine and plasma 273

Amino acids

MS study of the transesterification of O,N^{α} -bis(trifluoroacetyl)*n*-butyl esters of some hydroxylated amino acids into their O-carbethoxy, N^{α} -trifluoroacetyl *n*-butyl ester derivatives during the N^{*}-carbethoxy derivatization of histidine 413

Aminophenazone

GC determination of phenazone derivatives in human plasma. I. Aminophenazone 503

Amitriptyline

Simplified method for monitoring tricyclic antidepressant therapy using GLC with nitrogen detection 449

Anaesthetics

Rapid GC analysis of plasma levels of ketamine and major metabolites employing either nitrogen selective or MS detection 344 Anaesthetics

Routine direct injection GLC procedure for the analysis of volatile halogenated anaesthetics in whole blood using a new external injection port 297

Analgesic drugs

Fluorimetric determination of flufenamic acid from plasma by direct measurement of TLC plates 169

Analgesic drugs

GC determination of phenazone derivatives in human plasma. I. Aminophenazone 503

Analgesic drugs

GC determination of phenazone derivatives in human plasma. II. Propyphenazone 508

Anti-anginal drugs

Determination of molsidomine in plasma by column HPLC 465

Anti-anginal drugs

Improved method for the rapid determination of isosorbide dinitrate in human plasma and its application in pharmacokinetic studies 457

Anti-anxiety drugs

Analysis of lorazepam and its glucuronide metabolite by electron-capture GLC. Use in pharmacokinetic studies of lorazepam 311

Anti-asthmatic drugs

Micro-method for the determination of caffeine and theophylline allowing direct application of biological fluids to TLC plates 175

Anticoagulants

Comparison of HPLC and a spectrophotometric technique for determining plasma warfarin 522

Anticonvulsant drugs

New direct micro-method for determination of valproic acid in serum by GC 494

Anticonvulsant drugs

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin,

phenobarbital and primidone in serum by TLC 167

Anticonvulsant drugs

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC. Improvement of the buffer system 361

Antidepressant drugs

Quantitation of zimelidine and norzimelidine in plasma using HPLC 113 Antidepressants, tricyclic

Simplified method for monitoring tricyclic antidepressant therapy using GLC with nitrogen detection 449

Antidiabetic drugs TLC determination of acylaminoethylarylsulphonylureas in serum after hydrolysis and dansylation 533

Antihistamine drugs

HPLC determination of cimetidine sulphoxide in human blood and urine 354

Antihypertensive drugs

Determination of the thiazide diuretic bemetizide in the plasma and urine of humans by HPLC 152

Anti-inflammatory drugs GC determination of phenazone derivatives in human plasma. I. Aminophenazone 503

Anti-inflammatory drugs

GC determination of phenazone derivatives in human plasma. II. Propyphenazone 508

Antileukemic drugs

Determination of cytosine arabinoside in human plasma by GC with a nitrogen-sensitive detector and by GC-MS 283

Antimetabolites

Determination of cytosine arabinoside in human plasma by HPLC 350

Antiprotozoal drugs

Determination of metronidazole and misonidazole and their metabolites in plasma and urine by HPLC 163 Antipyretic drugs

GC determination of phenazone derivatives in human plasma. I. Aminophenazone 503 Antipyretic drugs

GC determination of phenazone derivatives in human plasma. II. Propyphenazone 508

Antirheumatic drugs

Fluorimetric determination of flufenamic acid from plasma by direct measurement of TLC plates 169

Antitumour agents

Determination of cytosine arabinoside in human plasma by HPLC 350

Antitumour drugs

Quantitative analysis of 5-fluorouracil in human serum by selected ion monitoring GC-MS 512

Antitumour drugs

Rapid determination of 1-(2-tetrahydrofuryl)-5-fluorouracil in human blood by HPLC 518

Arylsulphonylurea, acylaminoethyl derivatives

TLC determination of acylaminoethylarylsulphonylureas in serum after hydrolysis and dansylation 533

Assays, mass fragmentographic

Automation of HPLC sample clean-up for mass fragmentographic assays 425 Auto-sampling

Automation of HPLC sample clean-up for mass fragmentographic assays 425

Bemetizide

Determination of the thiazide diuretic bemetizide in the plasma and urine of humans by HPLC 152

Benzodiazepines

Analysis of lorazepam and its glucuronide metabolite by electron-capture GLC. Use in pharmacokinetic studies of lorazepam 311

Benzodiazepines

Quantitation of chlordiazepoxide and its metabolites in biological fluids by TLC 473

Benzodiazepines

Quantitative analysis of prazepam and its metabolites by electron capture GC and selected ion monitoring. Application to diaplacental passage and fetal hepatic metabolism in early human pregnancy 227

Bile salts

Quantitative separation of bile salts from mixtures by gradient elution HPLC 137 Cadaverine

Monoacylcadaverines in the blood of schizophrenic patients 67

Caffeine

Microdetermination of caffeine in blood by GC-MS 85

Caffeine

Micro-method for the determination of caffeine and theophylline allowing direct application of biological fluids to TLC plates 175

Carbamazepine

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC 167

Carbamazepine

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC. Improvement of the buffer system 361

Carboxy tolbutamide

Simple, rapid and micro HPLC method for the simultaneous determination of tolbutamide and carboxy tolbutamide in plasma 121

Catecholamines

Fluorimetric detection of octopamine in HPLC and its application to the assay of dopamine β -monooxygenase in human serum 433

Catecholamines

Quantitation of adrenaline and noradrenaline from human plasma by combined GC-high-resolution mass fragmentography 221

Chemotherapeutic drugs Determination of sulfapyridine and its major metabolites by HPLC 157

Chemotherapeutic drugs Determination of trimethoprim and sulfamethoxazole (Co-Trimoxazole) in body fluids of man by means of HPLC 103

Chemotherapeutic drugs

GLC analysis of trimethoprim in plasma and urine 143

Chlordiazepoxide Quantitation of chlordiazepoxide and its metabolites in biological fluids by TLC 473 Cholesterol

Class - gar

Automated quantitative GLC of intact lipids. I. Preparation and calibration of the column 241

Cholesterol

Comparative determination of plasma cholesterol and triacylglycerol levels by automated GLC and AutoAnalyzer methods 393

Cholesterol

Quantitative analysis of human and rabbit tear cholesterol by GLC 321

Chondroitin sulphates

Rapid and sensitive determination of enzymatic degradation products of isomeric chondroitin sulfates by HPLC 439

Cimetidine

HPLC determination of cimetidine sulphoxide in human blood and urine 354

Clinical chemistry

Place of GC-MS in clinical chemistry 185

Co-enzymes

New method for quantitative analysis of pyridoxal-5'-phosphate in biological material 485

Co-Trimoxazole

Determination of trimethoprim and sulfamethoxazole (Co-Trimoxazole) in body fluids of man by means of HPLC 103

Cytosine arabinoside

Determination of cytosine arabinoside in human plasma by GC with a nitrogen-sensitive detector and by GC-MS 283

Cytosine arabinoside

Determination of cytosine arabinoside in human plasma by HPLC 350

Deoxycortisol

Convenient system for the simultaneous separation of 11-deoxycortisol and aldosterone by Sephadex LH-20 multiple column chromatography 333

Diphenylhydantoin

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC 167 Diphenylhydantoin

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC. Improvement of the buffer system 361

Dipropylacetic acid

New direct micro-method for determination of valproic acid in serum by GC 494

Diuretic drugs

Determination of the thiazide diuretic bemetizide in the plasma and urine of humans by HPLC 152

Dopamine β -monooxygenase

Fluorimetric detection of octopamine in HPLC and its application to the assay of dopamine β -monooxygenase in human serum 433

Drug monitoring

Comparison of HPLC and a spectrophotometric technique for determining plasma warfarin 522

Drug monitoring

Determination of cytosine arabinoside in human plasma by GC with a nitrogen-sensitive detector and by GC-MS 283

Drug monitoring

Determination of metformin in plasma at therapeutic levels by GLC using a nitrogen detector 148

Drug monitoring

Determination of metronidazole and misonidazole and their metabolites in plasma and urine by HPLC 163

Drug monitoring

Determination of molsidomine in plasma by column HPLC 465

Drug monitoring

Determination of sulfapyridine and its major metabolites by HLPC 157

Drug monitoring

Determination of the thiazide diuretic bemetizide in the plasma and urine of humans by HPLC 152

Drug monitoring

Determination of trimethoprim and sulfamethoxazole (Co-Trimoxazole) in body fluids of man by means of HPLC 103

Drug monitoring

Fluorimetric determination of flufenamic acid from plasma by direct measurement of TLC plates 169 Drug monitoring

GC determination of phenazone derivatives in human plasma. I. Aminophenazone 503

Drug monitoring

GC determination of phenazone derivatives in human plasma. II. Propyphenazone 508

Drug monitoring

GC-MS analysis of optically active metabolites and drugs on a novel chiral stationary phase 197

Drug monitoring

GLC analysis of trimethoprim in plasma and urine 143

Drug monitoring

HPLC determination of cimetidine sulphoxide in human blood and urine 354

Drug monitoring

Improved method for the rapid determination of isosorbide dinitrate in human plasma and its application in pharmacokinetic studies 457

Drug monitoring

Micro-method for the determination of caffeine and theophylline allowing direct application of biological fluids to TLC plates 175

Drug monitoring

New direct micro-method for determination of valproic acid in serum by GC 494

Drug monitoring

Quantitation of chlordiazepoxide and its metabolites in biological fluids by TLC 473

Drug monitoring

Quantitation of zimelidine and norzimelidine in plasma using HPLC 113

Drug monitoring

Quantitative analysis of 5-fluorouracil in human serum by selected ion monitoring GC—MS 512

Drug monitoring

Quantitative analysis of prazepam and its metabolites by electron capture GC and selected ion monitoring. Application to diaplacental passage and fetal hepatic metabolism in early human pregnancy 227

Drug monitoring

Rapid determination of 1-(2-tetrahydrofuryl)-5-fluorouracil in human blood by HPLC 518

Drug monitoring

Rapid GC analysis of plasma levels of ketamine and major metabolites employing either nitrogen selective or mass spectroscopic detection 344

Drug monitoring

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC 167

Drug monitoring

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC. Improvement of the buffer system 361

Drug monitoring

Rapid, sensitive analysis of 8-methoxypsoralen in human plasma 498

Drug monitoring

Routine direct injection GLC procedure for the analysis of volatile halogenated anaesthetics in whole blood using a new external injection port 297

Drug monitoring

Simple, rapid and micro HPLC method for the simultaneous determination of tolbutamide and carboxy tolbutamide in plasma 121

Drug monitoring

Simplified method for monitoring tricyclic antidepressant therapy using GLC with nitrogen detection 449

Drug monitoring

TLC determination of acylaminoethylarylsulphonylureas in serum after hydrolysis and dansylation 533

Enzymes

Fluorimetric detection of octopamine in HPLC and its application to the assay of dopamine β -monooxygenase in human serum 433

Ergenyl

New direct micro-method for determination of valproic acid in serum by GC 494

Flufenamic acid

Fluorimetric determination of flufenamic acid from plasma by direct measurement of TLC plates 169

5-Fluorouracil

Quantitative analysis of 5-fluorouracil

in human serum by selected ion monitoring GC-MS 512

Furocoumarins

Rapid, sensitive GC analysis of 8-methoxy-psoralen in human plasma 498 Glycopeptides

Two-dimensional separation of glycopeptides and charged oligosaccharides on silica thin layers 337

Glycosaminoglycans

Rapid and sensitive determination of enzymatic degradation products of isomeric chondroitin sulfates by HPLC 439

Halogenated anaesthetics

Routine direct injection GLC procedure for the analysis of volatile halogenated anaesthetics in whole blood using a new external injection port 297

Hawkinsin

New tyrosine metabolites in humans: hawkinsin and *cis*- and *trans*-4-hydroxycyclohexylacetic acids. Unusual adsorption of deuterated and non-deuterated hawkinsin during GC 207 Histamine

Determination of histamine in plasma by high-speed LC 490

Histidine

MS study of the transesterification of O,N^{α} -bis(trifluoroacetyl)*n*-butyl esters of some hydroxylated amino acids into their O-carbethoxy, N^{α} -trifluoroacetyl, *n*-butyl ester derivatives during the N^{*}-carbethoxy derivatization of histidine 413

Hormones, steroid

Determination of aldosterone in urine and plasma 273

Hydrocarbons, saturated

Paraffin oil pneumonia. Analysis of saturated hydrocarbons in different human tissues 91

Hydroxycyclohexylacetic acids

New tyrosine metabolites in humans: hawkinsin and *cis*- and *trans*-4-hydroxycyclohexylacetic acids. Unusual adsorption of deuterated and non-deuterated hawkinsin during GC 207 Hydroxylated amino acids

MS study of the transesterification of O, N^{α} -bis(trifluoroacetyl)*n*-butyl esters of some hydroxylated amino acids into their O-carbethoxy, N^{α} -trifluoroacetyl *n*-butyl ester derivatives during the N⁻-carbethoxy derivatization of histidine 413

Hydroxyproline

MS study of the transesterification of O_N^{α} -bis(trifluoroacetyl)n-butyl esters of some hydroxylated amino acids into their O-carbethoxy.Na-trifluoroacetyl n-butyl ester derivatives during the N^{*}-carbethoxy derivatization of histidine 413

Hypoglycaemic drugs

Determination of metformin in plasma at therapeutic levels by GLC using a nitrogen detector 148

Hypoglycaemic drugs

_ Simple, rapid and micro HPLC method for the simultaneous determination of tolbutamide and carboxy tolbutamide in plasma 121

Hypoglycaemic drugs

TLC determination of acylaminoethylarylsulphonylureas in serum after hydrolysis and dansylation 533

Immunoreactive substances

Isolation of immunoreactive components from experimental and human tumour tissues and serums by HPLC 253

Indocyanine green

Electrophoretic bimorphism of serum albumin in the presence of indocyanine green 77

Isosorbide dinitrate

Improved method for the rapid determination of isosorbide dinitrate in human plasma and its application in pharmacokinetic studies 457

Ketamine

Rapid GC analysis of plasma levels of ketamine and major metabolites employing either nitrogen selective or mass spectroscopic detection 344

Ketone bodies

GC head-space analysis of clinically interesting ketone bodies 327

Kvnurenine

Determination of kynurenine by a simple gas-liquid chromatographic method applicable to urine, plasma, brain and cerebrospinal fluid 33

Lipids

Automated quantitative GLC of intact lipids. I. Preparation and calibration of the column 241

Lipids

Comparative determination of plasma cholesterol and triacylglycerol levels by automated GLC and AutoAnalyzer methods 393

Lipids

Quantitative HPTLC of lipids in plasma and liver homogenates after direct application of 0,5-µl samples to the silica-gel layer 261

and a second second

Lorazepam

Analysis of lorazepam and its glucuronide metabolite by electron-capture GLC. Use in pharmacokinetic studies of lorazepam 311

Mephenytoin

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10.11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC 167

Mephenytoin

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC. Improvement of the buffer system 361

Metabolic disorders

New tyrosine metabolites in humans: hawkinsin and cis- and trans-4-hydroxycyclohexylacetic acids. Unusual adsorption of deuterated and non-deuterated hawkinsin during GC 207 Metabolic disorders

Quantitative analysis of β -phenylpyruvic acid by single ion monitoring. Evaluation of isomeric internal standards 213

Metabolic disorders

Rapid and simplified extraction procedure for GC-MS profiling of urinary organic acids 481

Metformin

Determination of metformin in plasma at therapeutic levels by GLC using a nitrogen detector 148

Methoxypsoralen

Rapid, sensitive GC analysis of 8methoxy-psoralen in human plasma 498

Methylene diphosphonate

In vitro and in vivo studies with sodium pertechnetate and technetiummethylene labelled diphosphonate 527

Metronidazole

Determination of metronidazole and misonidazole and their metabolites in plasma and urine by HPLC 163

"Middle molecules"

Determination of "middle molecules" presenting vitamin B_{12} molecular size in normal and uremic body fluids 55 Misonidazole

. . . .

Determination of metronidazole and misonidazole and their metabolites in plasma and urine by HPLC 163

Molsidomine

Determination of molsidomine in plasma by column HPLC 465

Monoacylcadaverines

Monoacylcadaverines in the blood of schizophrenic patients 67

Mucopolysaccharides

Rapid and sensitive determination of enzymatic degradation products of isomeric chondroitin sulfates by HPLC 439

Nitroimidazole derivatives

Determination of metronidazole and misonidazole and their metabolites in plasma and urine by HPLC 163

Noradrenaline

Quantitation of adrenaline and noradrenaline from human plasma by combined GC—high-resolution mass fragmentography 221

Nortriptyline

Simplified method for monitoring tricyclic antidepressant therapy using GLC with nitrogen detection 449 Norzimelidine

Quantitation of zimelidine and norzimelidine in plasma using HPLC 113

Octopamine

Fluorimetric detection of octopamine in HPLC and its application to the assay of dopamine β -monooxygenase in human serum 433

Oligosaccharides

Two-dimensional separation of glycopeptides and charged oligosaccharides on silica thin layers 337

Organic acids

Rapid and simplified extraction procedure for GC-MS profiling of urinary organic acids 481

Paraffin oil pneumonia

Paraffin oil pneumonia. Analysis of saturated hydrocarbons in different human tissues 91

Phenazone derivatives

GC determination of phenazone derivatives in human plasma. I. Aminophenazone 503 Phenazone derivatives

۰.

GC determination of phenazone derivatives in human plasma. II. Propyphenazone 508

Phenobarbital

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC 167

Phenobarbital

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC. Improvement of the buffer system 361

Phenylpyruvic acid

Quantitative analysis of β -phenylpyruvic acid by single ion monitoring. Evaluation of isomeric internal standards 213

Plasma lipids

Automated quantitative GLC of intact lipids. I. Preparation and calibration of the column 241

Plasma lipids

Quantitative HPTLC of lipids in plasma and liver homogenates after direct application of $0.5 - \mu l$ samples to the silica-gel layer 261

Plasma proteins

Applications of polyacrylamide gel electrophoresis and polyacrylamide gel isoelectric focusing in clinical chemistry 1

Polyacrylamide gel electrophoresis

Applications of polyacrylamide gel electrophoresis and polyacrylamide gel isoelectric focusing in clinical chemistry 1

Quantitative analysis of prazepam and its metabolites by electron capture GC and selected ion monitoring. Application to diaplacental passage and fetal hepatic metabolism in early human pregnancy 227

Primidone

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC 167

Prazepam

Primidone

Rapid quantitative method for the simultaneous determination of carbamazepine, carbamazepine-10,11-epoxide, diphenylhydantoin, mephenytoin, phenobarbital and primidone in serum by TLC. Improvement of the buffer system 361

Profiles, chromatographic

Analysis of sulfonic acids by GC-MS of trimethylsilyl derivatives 132

Profiles, chromatographic

Determination of aldosterone in urine and plasma 273

Profiles, chromatographic

Determination of kynurenine by a simple gas—liquid method applicable to urine, plasma, brain and cerebrospinal fluid 33

Profiles, chromatographic Determination of "middle molecules" presenting vitamin B_{12} molecular size in normal and uremic body fluids 55 Profiles, chromatographic

- Determination of urinary thiamine by HPLC utilizing the thiochrome fluo-
- rescent method 43
- Profiles, chromatographic GC head-space analysis of clinically interesting ketone bodies 327

Profiles, chromatographic GC-MS analysis of optically active

metabolites and drugs on a novel chiral stationary phase 197

Profiles, chromatographic

Isolation of immunoreactive components from experimental and human tumour tissues and serums by highperformance gel chromatography 253

Profiles, chromatographic Monoacylcadaverines in the blood of schizophrenic patients 67

Profiles, chromatographic

Place of GC-MS in clinical chemistry 185

Profiles, chromatographic

Profiles in chronic diseases. I. Investigations of steroid profiles in uremia 381

Profiles, chromatographic

Quantitative analysis of β -phenylpyruvic acid by single ion monitoring. Evaluation of isomeric internal standards 213 Profiles, chromatographic

Quantitative analysis of steroid profiles from urine by capillary GC. I. Accuracy and reproducibility of the sample preparation 365

Profiles, chromatographic Quantitative HPTLC of lipids in plasma and liver homogenates after direct application of 0.5-µl samples to the silica-gel layer 261

Profiles, chromatographic

Rapid and simplified extraction procedure for GC-MS profiling of urinary organic acids 481

Propyphenazone

GC determination of phenazone derivatives in human plasma. II. Propyphenazone 508

Proteins

Applications of polyacrylamide gel electrophoresis and polyacrylamide gel isoelectric focusing in clinical chemistry 1

Pyridoxal-5'-phosphate

New method for quantitative analysis of pyridoxal-5'-phosphate in biological material 485

Pyrimidines

Determination of cytosine arabinoside in human plasma by HPLC 350

Pyrimidines

Quantitative analysis of 5-fluorouracil in human serum by selected ion monitoring GC-MS 512

Sample clean-up

Automation of HPLC sample clean-up for mass fragmentographic assays 425

Sedatives

Analysis of lorazepam and its glucuronide metabolite by electron-capture GLC. Use in pharmacokinetic studies of lorazepam 311

Serine

MS study of the transesterification of O,N^{α} -bis(trifluoroacetyl)*n*-butyl esters of some hydroxylated amino acids into their O-carbethoxy, N^{α} -trifluoroacetyl *n*-butyl ester derivatives during the N^{*}-carbethoxy derivatization of histidine 413

Serum albumin Electrophoretic bimorphism of serum albumin in the presence of indocyanine green 77 Sodium pertechnetate

In vitro and in vivo studies with sodium pertechnetate and technetiumlabelled methylene diphosphonate 527

Stationary phases, chiral

GC-MS analysis of optically active metabolites and drugs on a novel chiral stationary phase 197

Steroids

Convenient system for the simultaneous separation of 11-deoxycortisol and aldosterone by Sephadex LH-20 multiple column chromatography 333

Steroids

Determination of adosterone in urine and plasma 273

Steroids

Profiles in chronic diseases. I. Investigations of steroid profiles in uremia 381

Steroids

Quantitative analysis of steroid profiles from urine by capillary GC. I. Accuracy and reproducibility of the sample preparation 365

Sterols

Quantitative analysis of human and rabbit tear cholesterol by GLC 321 Stimulants

Microdetermination of caffeine in blood in GC-MS 85

Stimulants

Micro-method for the determination of caffeine and theophylline allowing direct application of biological fluids to TLC plates 175

Sulphamethoxazole

Determination of trimethoprim and sulfamethoxazole (Co-Trimoxazole) in body fluids of man by means of HPLC 103

Sulphonamides

Determination of trimethoprim and sulfamethoxazole (Co-Trimoxazole) in body fluids of man by means of HPLC 103

Sulphonic acids

Analysis of sulfonic acids by GC–MS of trimethylsilyl derivatives 132 Tagamet

HPLC determination of cimetidine sulphoxide in human blood and urine 354

Tear cholesterol

Quantitative analysis of human and rabbit tear cholesterol by GLC 321 Technetium-99

In vitro and in vivo studies with sodium pertechnetate and technetiumlabelled methylene diphosphonate 527

1-(2-Tetrahydrofuryl)-5-fluoroacyl Rapid determination of 1-(2-tetrahydrofuryl)-5-fluorouracil in human blood by HPLC 518

Theophylline

Micro-method for the determination of caffeine and theophylline allowing direct application of biological fluids to TLC plates 175

Thiamine -

Determination of urinary thiamine by HPLC utilizing the thiochrome fluorescent method 43

Threonine

MS study of the transesterification of O, N^{α} -bis(trifluororacetyl)*n*-butyl esters of some hydroxylated amino acids into their O-carbethoxy, N^{α} -trifluoro-acetyl *n*-butyl ester derivatives during the N^r-carbethoxy derivatization of histidine 413

Tissue proteins

Applications of polyacrylamide gel electrophoresis and polyacrylamide gel isoelectric focusing in clinical chemistry 1

Tolbutamide

Simple, rapid and micro HPLC method for the simultaneous determination of tolbutamide and carboxy tolbutamide in plasma 121

Tranquilizers

Quantitation of chlordiazepoxide and its metabolites in biological fluids by TLC 473

Tranquilizers

Quantitative analysis of prazepam and its metabolites by electron capture GC and selected ion monitoring. Application to diaplacental passage and fetal hepatic metabolism in early human pregnancy 227

Triacylglycerol

Comparative determination of plasma cholesterol and triacylglycerol levels by automated GLC and AutoAnalyzer methods 393 Triglycerides

Automated quantitative GLC of intact lipids. I. Preparation and calibration of the column 241

Trimethoprim

Determination of trimethoprim and sulfamethoxazole (Co-Trimoxazole) in body fluids of man by means of HPLC 103

Trimethoprim

GLC analysis of trimethoprim in plasma and urine 143

Tyrosine

MS study of the transesterification of O_*N^{α} -bis(trifluoroacetyl)*n*-butyl esters of some hydroxylated amino acids into their O-carbethoxy, N^{α} trifluoroacetyl *n*-butyl ester derivatives during the N^{*}-carbethoxy derivatization of histidine 413

Tyrosine metabolites

New tyrosine metabolites in humans: hawkinsin and *cis*- and *trans*-4-hydroxycyclohexylacetic acids. Unusual adsorption of deuterated and non-deuterated hawkinsin during GC 207

Uremia

Determination of "middle molecules" presenting vitamin B_{12} molecular size in normal and uremic body fluids 55 Urinary metabolites

Rapid and simplified extraction proce-

dure for GC-MS profiling of urinary organic acids 481

Valproie acid

New direct micro-method for determination of valproic acid in serum by GC 494

Vasodilators

Improved method for the rapid determination of isosorbide dinitrate in human plasma and its application in pharmacokinetic studies 457

Vitamins

Determination of urinary thiamine by HPLC utilizing the thiochrome

fluorescent method 43

Vitamins

New method for quantitative analysis of pyridoxal-5'-phosphate in biological material 485

Warfarin

Comparison of HPLC and spectrophotometric technique for determining plasma warfarin 522

Xanthine derivatives

Micro-method for the determination of caffeine and theophylline allowing direct application of biological fluids to TLC plates 175

Zimelidine

Quantitation of zimelidine and norzimelidine in plasma using HPLC 113