

Related articles published in *Journal of Chromatography*, Vols. 248 and 249

Volume 248

Correlation between the retention of cardiac glycosides in reversed-phase high-performance liquid chromatography with a diphenylsilyl stationary phase, the structure of their molecules and their biological activity by V.Ya. Davydov, M.E. Gonzalez and A.V. Kiselev (Moscow, U.S.S.R.)	49
Gas chromatographic identification of common drugs by their multiple peaks and those of their trimethylsilyl derivatives by M.C. Dutt (Singapore)	115
Determination of impurities in levodopa and carbidopa by high-performance liquid chromatography with electrochemical detection by L.A. Rihbany (Winchester, MA, U.S.A.) and M.F. Delaney (Boston, MA, U.S.A.)	125
Rapid high-performance liquid chromatographic determination of acetaminophen in dosage form using a totally aqueous mobile phase by R.K. Gilpin and M.H. Gaudet (Danbury, CT, U.S.A.)	160
Liquid chromatographic separation and detection of epinephrine and its oxidation products, adrenochrome and adrenolutin by E.C. Juenge, P.E. Flinn and W.B. Furman (St. Louis, MO, U.S.A.)	297
Analysis of the macrolide antibiotics propionylmaridomycins and related compounds by reversed-phase high-performance liquid chromatography by G.A. Bens, W. Van den Bossche, G. Van der Weken and P. De Moerloose (Ghent, Belgium)	312
Study of the stability of cocaine and benzoylecgonine, its major metabolite, in blood samples by Y. Liu, R.D. Budd and E.C. Griesemer (Los Angeles, CA, U.S.A.)	318
Rapid separation of biologically active Syrian hamster lymphotoxin in high yield by size-exclusion high-performance liquid chromatography by J.P. Fuhrer and C.H. Evans (Bethesda, MD, U.S.A.)	427
Purification of polyadenylated mRNA on three oligo(dT)-substituted gels: a comparative study by C. Sene, P. Girot and E. Boschetti (Villeneuve la Garenne, France) and J.L. Plassat, J. Bloch and J.M. Egly (Strasbourg, France)	441
Thin-layer chromatographic identification of aminoglycoside antibiotics by P.J. Claes and H. Vanderhaeghe (Leuven, Belgium)	483

Volume 249

Kováts' indices of trimethylsilylated amino acids on fused-silica capillary columns by E. Gajewski, M. Dizzaroglu and M.G. Simic (Washington, DC, U.S.A.)	41
Hydrogenation of cholesteryl esters during gas chromatography on a polar fused-silica capillary column by N.B. Smith (London, Canada)	57
Determination of specificity of endopeptidases by combined high-performance liquid chromatography and amino acid analysis by S. Wilk and M. Orlowski (New York, NY, U.S.A.)	121
High-performance liquid chromatographic determination of hydroxyproline after derivatization with 4-dimethylaminoazobenzene-4-sulphonyl chloride by A. Casini, F. Martini, S. Nieri, D. Ramari, F. Franconi and C. Surrenti (Florence, Italy) . .	187
Facile, semi-preparative, high-performance liquid chromatographic separation of synthetic peptides using ammonium bicarbonate buffers by D.R. Knighton, D.R.K. Harding, J.R. Napier and W.S. Hancock (Palmerston North, New Zealand)	193
Nachweis von LSD in Körperflüssigkeiten mit Hochleistungsflüssigkeitschromatographie von K. Harzer (Stuttgart, B.R.D.)	205

Determination of bromhexine hydrochloride in pharmaceutical preparations by reversed-phase ion-pair high-performance liquid chromatography by J.L. Kumar, W.C. Mann and A. Rozanski (Queenborough, Great Britain)	373
Fast protein liquid chromatography of pyridoxalated and glutaraldehyde-treated human haemoglobin by M. Kramlová, T.I. Přistoupil, V. Fríčová and J. Kraml (Prague, Czechoslovakia) and R. Berglund and N. Kantardjiev (Uppsala, Sweden).	403
Isotachophoretic analysis of two synthetic peptides by P. Stehle, B. Kühne, P. Pfaender and P. Fürst (Stuttgart, G.F.R.).	408