

Author Index

- Ageta, T.
 —, Mikasa, H., Kojima, K. and Kodama, H.
 Determination of γ -aminobutyric acid in rat brain using an isotachophoretic analyzer 361
- Ahmed, A.E.
 —, Koenig, M. and Farrish, Jr., H.H.
 Studies on the quantitation of chlorambucil in plasma by reversed-phase high-performance liquid chromatography 392
- Albani, F., see Riva, R. 371
- Alberts, D.S., see Peng, Y.-M. 235
- Aravind, M.K.
 —, Miceli, J.N. and Kauffman, R.E.
 Analysis of piperacillin using high-performance liquid chromatography 423
- Arima, M., see Nakazawa, K. 313
- Arita, T., see Takikawa, K. 343
- Assenza, S.P., see Nakano, K. 51
- Astier, A.
 — and Renat, B.
 Sensitive high-performance liquid chromatographic determination of indomethacin in human plasma. Pharmacokinetic studies after single oral dose 279
- Baltassat, P., see Boulieu, R. 131
- Baruzzi, A., see Riva, R. 371
- Basseches, P.J.
 —, Durski, A. and Powis, G.
 High-performance liquid chromatographic assay of the antineoplastic agent tricyclic nucleoside 5'-phosphate and its disposition in rabbit 227
- Bergamo, N., see Pastorello, L. 398
- Berge, R.J.M. ten, see Jonkers, R.E. 249
- Berthon, G., see Brisson, A.M. 386
- Bisker, A., see Monboisse, J.C. 355
- Bogema, S.C.
 —, Narasimhachari, N., Mumtaz, M., Goldin, S. and Friedel, R.O.
 Separation and quantitation of *cis*- and *trans*-triothixene in human plasma by high-performance liquid chromatography 257
- Boré, P., see Goetz, N. 19
- Borel, J.P., see Monboisse, J.C. 355
- Bory, C., see Boulieu, R. 131
- Boulieu, R.
 —, Bory, C., Baltassat, P. and Gonnet, C. 131
- Boxtel, C.J. van, see Jonkers, R.E. 249
- Brechbühler, S., see Degen, P.H. 375
- Brisson, A.M.
 —, Fourtillan, J.B. and Berthon, G.
 Sensitive high-performance liquid chromatographic analysis of moxalactam in biological fluids 386
- Brown, P.R., see Nakano, K. 51
- Burbach, J.P.H., see Verhoef, J. 317
- Capet-Antonini, F., see Ong, H. 97
- Carlson, J.H., see Elchisak, M.A. 79
- Chiu, T.S.K., see Umans, J.G. 213
- Codd, E.E., see Verhoef, J. 317
- Cooper, J.K.
 —, McKay, G., Hawes, E.M. and Midha, K.K.
 High-performance liquid chromatographic assay of indomethacin and its application in pharmacokinetics in healthy volunteers 289
- , see McKay, G. 417
- Corkill, J.A.
 — and Crout, D.H.G.
 Simultaneous analysis of methyl methacrylate and methacrylic acid in blood by double isotope derivative dilution analysis 404
- Corona, G.L., see Santagostino, G. 89
- Crout, D.H.G., see Corkill, J.A. 404
- Cucchi, M.L., see Santagostino, G. 89
- Dalgaard, L., see Nordholm, L. 427
- D'Alterio, M., see Pastorello, L. 398
- Danks, M.K.
 Two simple high-performance liquid chromatographic methods for simultaneous determination of 2'-deoxycytidine 5'-triphosphate and cytosine arabinoside 5'-triphosphate concentrations in biological samples 141
- Davis, T.P., see Peng, Y.-M. 235
- Degen, P.H.
 —, Brechbühler, S., Schneider, W. and Zbinden, P.
 Determination of apparent dihydralazine in plasma by gas-liquid chromatography and electron-capture detection 375

- De Grazia, F., see Rubio, F. 149
- De Jong, A.P.J.M.
 Analysis of oxime-trimethylsilyl derivatives of organic acids on OV-1701 fused silica capillary column 297
- DiFazio, C.A., see Lake, C.L. 410
- Duckworth, E.N., see Lake, C.L. 410
- Durbin, C.G., see Lake, C.L. 410
- Durski, A., see Basseches, P.J. 227
- Eklund, P., see Torelli, J.A. 334
- Elchisak, M.A.
 — and Carlson, J.H.
 Assay of free and conjugated catecholamines by high-performance liquid chromatography with electrochemical detection 79
- Engle, J.S., see Lake, C.L. 410
- Farrish, Jr., H.H., see Ahmed, A.E. 392
- Felver, M.E., see Needham, L.L. 9
- Fourtillan, J.B., see Brisson, A.M. 386
- Frattini, P., see Santagostino, G. 89
- Friedel, R.O., see Bogema, S.C. 257
- Frigerio, A., see Pastorello, L. 398
- Fujimura, K.-I., see Horiuchi, M. 123
- Garland, W.A., see Rubio, F. 149, 157, 167
- Gelpi, E., see Mahy, N. 115
- Gill, R.
 — and Moffat, A.C.
 Analysis of barbiturates in blood by high-performance liquid chromatography 438
- Gilmour, W.J.
 — and Leary, J.R.
 Gas-liquid chromatographic determination of etoperidone in plasma, serum, and urine 381
- Goetz, N.
 —, Kaba, G. and Boré, P.
 Capillary gas chromatographic analysis of human skin surface lipids after microsampling on ground-glass platelets 19
- Goldin, S., see Bogema, S.C. 257
- Gonnet, C., see Bouliou, R. 131
- Grazia, F. De, see Rubio, F. 149
- Guerret, M.
 —, Julien-Larose, C., Kiechel, J.R. and Lavène, D.
 Determination of 3-hydroxy-guanfacine in biological fluids by electron-capture gas-liquid chromatography 181
- Gupta, R.N.
 Analysis of barbiturates in blood by high-performance liquid chromatography 437
- Haefelfinger, P.
 Sensitive assay for the tricyclic antidepressant Ro 11-2465 in biological fluids by high-performance liquid chromatography and fluorescence detection 269
- Hall, K., see McKay, G. 417
- Hall, M.E., see Metz, S.A. 193
- Han Jui
 — and Roboz, J.
 Quantitation of harringtonine and homoharringtonine in serum by high-performance liquid chromatography 203
- Harper, T.W., see Metz, S.A. 193
- Harzer, G., see Haug, M. 349
- Haug, M.
 —, Reinhardt, D. and Harzer, G.
 Rapid method for the determination of the phospholipid subclass distribution in human breast milk samples 349
- Hawes, E.M., see Cooper, J.K. 289
 —, see McKay, G. 417
- Hayashi, Y., see Udagawa, M. 338
- Hill, Jr., R.H., see Needham, L.L. 9
- Hirayama, C., see Udagawa, M. 338
- Horiuchi, M.
 —, Fujimura, K.-I., Terashima, T. and Iso, T.
 Method for determination of angiotensin-converting enzyme activity in blood and tissue by high-performance liquid chromatography 123
- Inturrisi, C.E., see Umans, J.G. 213
- Iso, T., see Horiuchi, M. 123
- Johnson, E.A.
 Simple electrophoresis of glycosaminoglycuronans and the distinction of lung heparins from mucosal heparins 365
- Jong, A.P.J.M. de, see De Jong, A.P.J.M. 297
- Jonkers, R.E.
 —, Oosterhuis, B., Ten Berge, R.J.M. and Van Bortel, C.J.
 Analysis of 6-mercaptopurine in human plasma with a high-performance liquid chromatographic method including post-column derivatization and fluorimetric detection 249
- Julien-Larose, C., see Guerret, M. 181
- Kaba, G., see Goetz, N. 19
- Kataoka, H., see Yamamoto, S. 29
- Kato, N., see Trocewicz, J. 328
- Kauffman, R.E., see Aravind, M.K. 423
- Kawasaki, T.

- , Maeda, M. and Tsuji, A.
Determination of 17-oxosteroid glucuronides and sulfates in urine and serum by fluorescence high-performance liquid chromatography using dansyl hydrazine as a pre-labeling reagent 61
- Kiechel, J.R., see Guerret, M. 181
- Kiyama, S., see Yamamoto, S. 39
- Knauf, H., see Reuter, K. 432
- Kodama, H., see Ageta, T. 361
- Koenig, M., see Ahmed, A.E. 392
- Kojima, K., see Ageta, T. 361
- Lake, C.L.
- , DiFazio, C.A., Duckworth, E.N., Moscicki, J.C., Engle, J.S. and Durbin, C.G.
High-performance liquid chromatographic analysis of plasma levels of nalbuphine in cardiac surgical patients 410
- Lamontagne, D., see Ong, H. 97
- Lavène, D., see Guerret, M. 181
- Leary, J.R., see Gilmour, W.J. 381
- Levy, E., see Torelli, J.A. 334
- Liddle, J.A., see Needham, L.L. 9
- Lindstrom, T.D.
- and Wolen, R.L.
Determination of clofilium, a new antifibrillatory agent, in plasma by gas chromatography—mass spectrometry 175
- Lipman, R.A., see Umans, J.G. 213
- Mabuchi, H.
- and Nakahashi, H.
Profiling of urinary medium-sized peptides in normal and uremic urine by high-performance liquid chromatography 107
- McKay, G.
- , Cooper, J.K., Midha, K.K., Hall, K. and Hawes, E.M.
Simple and sensitive high-performance liquid chromatographic procedure with electrochemical detection for the determination of plasma concentrations of trimetoprim following single oral doses 417
- , see Cooper, J.K. 289
- Maeda, K., see Ohki, T. 1
- Maeda, M., see Kawasaki, T. 61
- Mahy, N.
- , Tusell, J.M. and Gelpi, E.
Study of leucine-enkephalin in rat brain by a new high-performance liquid chromatographic method 115
- Makita, M., see Yamamoto, S. 29, 39
- Mantegazza, P., see Zecca, L. 307
- Metz, S.A.
- , Hall, M.E., Harper, T.W. and Murphy, R.C.
Rapid extraction of leukotrienes from biologic fluids and quantitation by high-performance liquid chromatography 193
- Miceli, J.N., see Aravind, M.K. 423
- Midha, K.K., see Cooper, J.K. 289
- , see McKay, G. 417
- Mikasa, H., see Ageta, T. 361
- Miwa, B.J., see Rubio, F. 149, 157, 167
- Miyazaki, K., see Takikawa, K. 432
- Moffat, A.C., see Gill, R. 438
- Monboisse, J.C.
- , Pierrelee, P., Bisker, A., Pailler, V., Randoux, A. and Borel, J.P.
Quantitation of the urinary methyl-histidine isomers by a combination of thin-layer and fluorometric techniques 355
- Moscicki, J.C., see Lake, C.L. 410
- Mumtaz, M., see Bogema, S.C. 257
- Murphy, R.C., see Metz, S.A. 193
- Mutschler, E., see Reuter, K. 432
- Nagatsu, T., see Trocewicz, J. 328
- Nakahashi, H., see Mabuchi, H. 107
- Nakano, K.
- , Assenza, S.P. and Brown, P.R.
Reversed-phase liquid chromatographic investigation of UV-absorbing low-molecular-weight compounds in saliva 51
- Nakazawa, K.
- , Tanaka, H. and Arima, M.
Rapid, simultaneous and sensitive determination of free hydroxyproline and proline in human serum by high-performance liquid chromatography 313
- Narasimhachari, N., see Bogema, S.C. 257
- Needham, L.L.
- , Hill, Jr., R.H., Orti, D.L., Felver, M.E. and Liddle, J.A.
Electron-capture, capillary column gas chromatographic determination of low-molecular-weight diols in serum 9
- Niwa, T., see Ohki, T. 1
- Nohta, H.
- , Yamaguchi, M., Zaitzu, K. and Ohkura, Y.
Assay of dopamine β -hydroxylase in human serum as a modification of the assay for the enzyme in rat serum by high-performance liquid chromatography 324

- Nordholm, L.
 — and Dalgaard, L.
 Assay of trimethoprim in plasma and urine by high-performance liquid chromatography using electrochemical detection 427
- Ohki, T.
 —, Saito, A., Ohta, K., Niwa, T., Maeda, K. and Sakakibara, J.
 Amine metabolite profile of normal and uremic urine using gas chromatography—mass spectrometry 1
- Ohkura, Y., see Nohta, H. 324
 Ohta, K., see Ohki, T. 1
 Oka, K., see Trocewicz, J. 328
 Olivi, F., see Riva, R. 371
 Ong, H.
 —, Capet-Antonini, F., Yamaguchi, N. and Lamontagne, D.
 Simultaneous determination of free 3-methoxy-4-hydroxymandelic acid and free 3-methoxy-4-hydroxyphenylethyleneglycol in plasma by liquid chromatography with electrochemical detection 97
- Oosterhuis, B., see Jonkers, R.E. 249
 Ormberg, D., see Peng, Y.-M. 235
 Orti, D.L., see Needham, L.L. 9
 Paillet, V., see Monboisse, J.C. 355
 Paladino, R., see Pastorello, L. 398
 Pantaloni, M., see Riva, R. 371
 Pastorello, L.
 —, Tolentino, D., D'Alterio, M., Paladino, R., Frigerio, A., Bergamo, N. and Valli, A.
 Determination of α -amanitin by high-performance liquid chromatography 398
- Peng, Y.-M.
 —, Ormberg, D., Alberts, D.S. and Davis, T.P.
 Improved high-performance liquid chromatography of the new antineoplastic agents bisantrene and mitoxantrone 235
- Pierrelee, P., see Monboisse, J.C. 355
 Powis, G., see Basseches, P.J. 227
 Randoux, A., see Monboisse, J.C. 355
 Reinhardt, D., see Haug, M. 349
 Renat, B., see Astier, A. 279
 Reuter, K.
 —, Knauf, H. and Mutschler, E.
 Fluorimetrische Bestimmung von Amilorid in Humanplasma mittels Dünnschichtchromatographie 432
- Riva, R.
 —, Albani, F., Olivi, F., Pantaloni, M. and Baruzzi, A.
 Quantitative determination of *n*-di-propylacetamide in the plasma of epileptic patients by gas-liquid chromatography with nitrogen-selective detection 371
- Roboz, J., see Han Jui 203
 Rubio, F.
 —, De Grazia, F., Miwa, B.J. and Garland, W.A.
 Determination of (–)-*threo*-chlorocitric acid in human plasma by gas chromatography—positive chemical-ionization mass spectrometry 149
- , Miwa, B.J. and Garland, W.A.
 Determination of imidazobenzodiazepine-3-carboxamide, a new anxiolytic agent, in human plasma by gas chromatography—negative chemical-ionization mass spectrometry 167
- , Miwa, B.J. and Garland, W.A.
 Determination of midazolam and two metabolites of midazolam in human plasma by gas chromatography—negative chemical-ionization mass spectrometry 157
- Saito, A., see Ohki, T. 1
 Sakakibara, J., see Ohki, T. 1
 Santagostino, G.
 —, Frattini, P., Schinelli, S., Cucchi, M.L. and Corona, G.L.
 Urinary 3-methoxy-4-hydroxyphenylethyleneglycol determination using reversed-phase chromatography with amperometric detection 89
- Scheline, R., see Svardal, A. 301
 Schinelli, S., see Santagostino, G. 89
 Schneider, W., see Degen, P.H. 375
 Schultz, M.F., see Umans, J.G. 213
 Shin, S.-U., see Umans, J.G. 213
 Svardal, A.
 — and Scheline, R.
 Gas chromatographic—mass spectrometric determination of aromatization of cyclohexanecarboxylic acid in guinea pig liver 301
- Takikawa, K.
 —, Miyazaki, K. and Arita, T.
 High-performance liquid chromatographic determination of nicotinic acid and its metabolites, nicotinuric acid and nicotinamide, in plasma 343
- Tanaka, H., see Nakazawa, K. 313
 Ten Berge, R.J.M., see Jonkers, R.E. 249
 Terashima, T., see Horiuchi, M. 123
 Tolentino, D., see Pastorello, L. 398
 Torelli, J.A.
 —, Eklund, P. and Levy, E.

- Enhanced high-performance liquid chromatographic resolution of hemoglobin A_{1c} at low temperatures 334
Trocewicz, J.
- , Kato, N., Oka, K. and Nagatsu, T.
Determination of phenylethanolamine N-methyltransferase activity in rat brain by high-performance liquid chromatography with fluorometric detection 328
- Tsuji, A., see Kawasaki, T. 61
- Tusell, J.M., see Mahy, N. 115
- Udagawa, M.
- , Hayashi, Y. and Hirayama, C.
Determination of coproporphyrin I and III isomers by high-performance liquid chromatography 338
- Umans, J.G.
- , Chiu, T.S.K., Lipman, R.A., Schultz, M.F., Shin, S.-U. and Inturrisi, C.E.
Determination of heroin and its metabolites by high-performance liquid chromatography 213
- Valli, A., see Pastorello, L. 398
- Van Bostel, C.J., see Jonkers, R.E. 249
- Verhoef, J.
- , Codd, E.E., Burbach, J.P.H. and Witter, A.
Reversed-phase high-performance liquid chromatography of neuropeptides related to adrenocorticotropin, including a potent adrenocorticotropin 4-9 analogue (ORG 2766) 317
- Wakamatsu, K., see Yamamoto, S. 29
- Watanabe, Y., see Yamamoto, S. 39
- Westerink, B.H.C.
Correlation between high-performance liquid chromatography and automated fluorimetric methods for the determination of dopamine, 3,4-dihydroxyphenylacetic acid, homovanillic acid and 5-hydroxyindoleacetic acid in nervous tissue and cerebrospinal fluid 69
- Witter, A., see Verhoef, J. 317
- Wolen, R.L., see Lindstrom, T.D. 175
- Yamaguchi, M., see Nohta, H. 324
- Yamaguchi, N., see Ong, H. 97
- Yamamoto, S.
- , Kiyama, S., Watanabe, Y. and Makita, M.
Practical gas-liquid chromatographic method for the determination of amino acids in human serum 39
- , Yokogawa, M., Wakamatsu, K., Kataoka, H. and Makita, M.
Gas chromatographic method for the determination of urinary acetylpolymines 29
- Yokogawa, M., see Yamamoto, S. 29
- Zaitzu, K., see Nohta, H. 324
- Zambotti, F., see Zecca, L. 307
- Zbinden, P., see Degen, P.H. 375
- Zecca, L.
- , Zambotti, F., Zonta, N. and Mantegazza, P.
Determination of γ -aminobutyric acid in brain areas by high-performance liquid chromatography of dansyl derivatives with ultraviolet detection 307
- Zonta, N., see Zecca, L. 307