

Author Index

- Alasandro, M.
 —, Wiesler, D. and Novotny, M.
 Effects of starvation and refeeding on the excretion of urinary steroid metabolites in mice with different genetic background 1
- Anjaneyulu, B., see Nene, S. 334
- Antonsson, A.-M.
 —, Gyllenhaal, O., Kylberg-Hanssen, K., Johansson, L. and Vessman, J.
 Monitoring of *S*- and *R*-tocainide in human plasma after heptafluorobutyrylation, separation on Chirasil-Val® and electron-capture detection 181
- Aravind, M.K.
 —, Miceli, J.N. and Kauffman, R.E.
 Determination of ibuprofen by high-performance liquid chromatography 350
- Beland, F.A.
 —, Fullerton, N.F. and Heflich, R.H.
 Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA 121
- Beretta, E.
 — and Vanazzi, G.
 Determination of nanogram amounts of dicyclomine with gas chromatography and nitrogen-selective detection 341
- Bernard, D.
 —, Maurizis, J.C., Chollet, Ph., Chassagne, J. and Plagne, R.
 Isolation and purification of HLA-DR antigens from tumour cells by affinity chromatography and chromatofocusing 322
- Bianchetti, G., see Padovani, P. 229
- Bobik, A., see Jackman, G.P. 301
 —, see Oddie, C.J. 370
- Bock, J.L.
 —, Lam, S. and Karmen, A.
 Therapeutic drug monitoring using high-speed liquid chromatography and rapid sample preparation: an assay for serum theophylline 354
- Bourne, D.W.A., see Schneider, J.J. 359
- Breckenridge, A.M., see Coleman, M.D. 363
- Breimer, D.D., see Kleinbloesem, C.H. 209
- Brondz, I.
 — and Olsen, I.
 Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 19
- and Olsen, I.
 Determination of bound cellular fatty acids in *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* by gas chromatography and gas chromatography-mass spectrometry 282
- Brummelen, P. van, see Kleinbloesem, C.H. 209
- Burch, J.E.
 —, Roberts, S.G. and Raddats, M.A.
 Amitriptyline and its basic metabolites determined in plasma by gas chromatography 165
- Calvey, T.N., see Murray, G.R. 143
- Carr Williams, M.
 —, Kelley, J.L. and Kushwaha, R.S.
 Detection of an abnormal lipoprotein in a large colony of pedigreed baboons using high-performance gel exclusion chromatography 101
- Cautreels, W., see Fels, J.-P. 273
- Chan, K., see Murray, G.R. 143
- Chassagne, J., see Bernard, D. 322
- Choi, R.L., see Rosenberg, M. 382
- Chollet, Ph., see Bernard, D. 322
- Chubb, C.
 — and Nolan, C.
 Quantitative metabolic profiling of testicular steroid secretion with bonded-phase capillary gas chromatography 11
- Claus-Walker, J., see Rodriguez, G.P. 65
- Cockrell, C.S.
 — and Ellis, E.F.
 Simple single-step high-performance liquid chromatographic method for the separation of cyclooxygenase and lipoxygenase enzyme metabolites of arachidonic acid 316
- Coleman, M.D.
 —, Edwards, G., Mihaly, G.W., Howells, R.E. and Breckenridge, A.M.
 High-performance liquid chromatographic method for the determination of pyrimethamine and its 3-*N*-oxide metabolite in biological fluids 363
- Coletti, O., see Cremonesi, P. 310
- Cremonesi, P.
 —, Galimberti, G., Coletti, O. and Sproletti, G.
 High-performance liquid chromatographic determination of imidazole in biological fluids 310

- Demaster, E.G.
 —, Shirota, F.N., Redfern, B., Goon, D.J.W. and Nagasawa, H.T.
 Analysis of hepatic reduced glutathione, cysteine and homocysteine by cation-exchange high-performance liquid chromatography with electrochemical detection 83
- Deves, C., see Padovani, P. 229
- Dobrinska, M.R., see Musson, D.G. 251
- Edwards, G., see Coleman, M.D. 363
- Ellis, E.F., see Cockrell, C.S. 316
- Fels, J.-P.
 —, Lechat, P., Rispe, R. and Cautreels, W.
 Determination of glaucine in plasma and urine by high-performance liquid chromatography 273
- Fullerton, N.F., see Beland, F.A. 121
- Fung, H.-L., see Morrison, R.A. 153
- Galimberti, G., see Cremonesi, P. 310
- George, S., see McBurney, A. 387
- Goon, D.J.W., see Demaster, E.G. 83
- Grego, B., see Skinner, S.J.M. 111
- Gyllenhaal, O., see Antonsson, A.-M. 181
- Hagino, S., see Tanaka, E. 393
- Harten, J. van, see Kleinbloesem, C.H. 209
- Haviland, A.M., see Schneider, J.J. 359
- Hearn, M.T.W., see Skinner, S.J.M. 111
- Heflich, R.H., see Beland, F.A. 121
- Hosojima, H., see Imaizumi, N. 295
- Howells, R.E., see Coleman, M.D. 363
- Imaizumi, N.
 —, Morimoto, S., Kigoshi, T., Uchida, K., Hosojima, H. and Yamamoto, I.
 Application of reversed-phase high-performance liquid chromatography for radioimmunoassay of plasma 18-hydroxycorticosterone 295
- Imanari, T., see Toida, T. 133
- Ishizaki, T., see Minegishi, A. 55
- Isozaki, S., see Uchino, K. 241
- Jackman, G.P., see Oddie, C.J. 370
 —, Oddie, C.J., Skews, H. and Bobik, A.
 High-performance liquid chromatographic determination of plasma catecholamines during α -methyldopa therapy 301
- Johansson, L., see Antonsson, A.-M. 181
- Kadowaki, H.
 —, Shiino, M., Uemura, I. and Kobayashi, K.
 Sensitive method for the determination of diclofenac in human plasma by gas chromatography—mass spectrometry 329
- Kågedal, B.
 — and Källberg, M.
 Cysteinylglycine in urine determined by high-performance liquid chromatography 75
- Källberg, M., see Kågedal, B. 75
- Karmen, A., see Bock, J.L. 354
- Katayama, K., see Kuwada, M. 398
- Kauffman, R.E., see Aravind, M.K. 350
- Kelley, J.L., see Carr Williams, M. 101
- Kigoshi, T., see Imaizumi, N. 295
- Kinkel, A.W., see Randinitis, E.J. 345
- Kiss, B., see Oka, K. 43
- Kleinbloesem, C.H.
 —, Van Harten, J., Van Brummelen, P. and Breimer, D.D.
 Liquid chromatographic determination of nifedipine in plasma and of its main metabolite in urine 209
- Kobayashi, K., see Kadowaki, H. 329
- Kojima, K., see Oka, K. 43
- Kuroiwa, Y., see Tanaka, E. 393
- Kushwaha, R.S., see Carr Williams, M. 101
- Kuwada, M.
 — and Katayama, K.
 An improved method for the determination of γ -carboxyglutamic acid in human urine by high-performance liquid chromatography 398
- Kyberg-Hanssen, K., see Antonsson, A.-M. 181
- Lakings, D.B.
 — and Wozniak, J.M.
 High-performance liquid chromatographic methods for the determination of cefpimizole in plasma and urine 261
- Lam, S., see Bock, J.L. 354
- Lechat, P., see Fels, J.-P. 273
- Leidy, H.L., see Musson, D.G. 251
- Liggins, G.C., see Skinner, S.J.M. 111
- Lundmo, P.I.
 — and Sunde, A.
 Rapid analysis of C_{19} -steroid metabolism by high-performance liquid chromatography and in-line monitoring of radioactivity 289
- McBurney, A.
 — and George, S.
 High-performance liquid chromatography of haloperidol in serum at the concentrations achieved during chronic therapy 387
- Maurizis, J.C., see Bernard, D. 322
- Miceli, J.N., see Aravind, M.K. 350
- Midskov, C.
 High-performance liquid chromatographic assay of pyrimethamine, sulfadoxine and its N^4 -acetyl metabolite in serum and urine after ingestion of Suldox 217

- Mihaly, G.W., see Coleman, M.D. 363
- Minegishi, A.
- and Ishizaki, T.
Rapid and simple method for the simultaneous determination of 3,4-dihydroxyphenylacetic acid, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in human plasma by high-performance liquid chromatography with electrochemical detection 55
- Morimoto, S., see Imaizumi, N. 295
- Morrison, R.A.
- and Fung, H.-L.
Determination of the partitioning, stability, and metabolite formation of isosorbide dinitrate in human and rat blood using an improved gas-liquid chromatographic assay 153
- Morselli, P.L., see Padovani, P. 229
- Murray, G.R.
- , Calvey, T.N., Williams, N.E. and Chan, K.
Quantitative capillary column gas chromatographic method for the determination of glycopyrronium in human plasma 143
- Musson, D.G.
- , Vincek, W.C., Dobrinska, M.R., Vickers, S. and Leidy, H.L.
Sensitive high-performance liquid chromatographic assay using electrochemical detection for a novel prodrug ester of methyl dopa, pivaloyloxyethyl 3-(3,4-dihydroxyphenyl)-2-methylalaninate, in plasma and urine 251
- Nagasawa, H.T., see Demaster, E.G. 83
- Nagatsu, T., see Oka, K. 43
- Nakagawa, F., see Uchino, K. 241
- Nelson, C., see Randinitis, E.J. 345
- Nene, S.
- , Anjaneyulu, B. and Rajagopalan, T.G.
Determination of diethylcarbamazine in blood using gas chromatography with alkali flame ionization detection 334
- Nohta, H.
- , Noma, S., Ohkura, Y. and Yoo, B.-T.
Assay for catechol-O-methyltransferase in erythrocytes using a new fluorogenic substrate, 2-(3,4-dihydroxyphenyl)-naphtho[1,2-d]thiazole 93
- Nolan, C., see Chubb, C. 11
- Noma, S., see Nohta, H. 93
- Nousiainen, U.
- and Raatikainen, O.
High-performance liquid chromatography of propanidid in rat plasma 376
- Novotny, M., see Alasandro, M. 1
- Oddie, C.J., see Jackman, G.P. 301
- , Jackman, G.P. and Bobik, A.
Measurement of xamoterol in plasma and urine by high-performance liquid chromatography 370
- Ohkura, Y., see Nohta, H. 93
- Oka, K.
- , Kojima, K., Togari, A., Nagatsu, T. and Kiss, B.
An integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by liquid chromatography with electrochemical detection 43
- Olsen, I., see Brondz, I. 19, 282
- Østensen, E.T., see Øyehaug, E. 199
- Øyehaug, E.
- , Østensen, E.T. and Salvesen, B.
High-performance liquid chromatographic determination of citalopram and four of its metabolites in plasma and urine samples from psychiatric patients 199
- Padovani, P.
- , Deves, C., Bianchetti, G., Thenot, J.P. and Morselli, P.L.
Determination of progabide and its main acid metabolite in biological fluids using high-performance liquid chromatography and electrochemical detection. Application to the measurement of blood/plasma partition ratio 299
- Plagne, R., see Bernard, D. 322
- Raatikainen, O., see Nousiainen, U. 376
- Raddats, M.A., see Burch, J.E. 165
- Rajagopalan, T.G., see Nene, S. 334
- Randinitis, E.J.
- , Nelson, C. and Kinkel, A.W.
Gas chromatographic determination of bevantolol in plasma 345
- Redfern, B., see Demaster, E.G. 83
- Rispe, R., see Fels, J.-P. 273
- Roberts, S.G., see Burch, J.E. 165
- Rodriguez, G.P.
- and Claus-Walker, J.
Measurement of hydroxylysine glycosides in urine and its application to spinal cord injury 65
- Rose, S.M., see Tsao, C.S. 306
- Rosenberg, M.
- and Choi, R.L.
High-performance liquid chromatographic analysis of flordipine in human plasma 382
- Rydzik, R.
- , Terragno, A. and Tackett, R.
Picogram detection of eicosanoids by

- ultraviolet absorbance after narrow-bore high-performance liquid chromatography. Comparison with conventional-bore column 31
- Saitoh, Y., see Uchino, K. 241
- Salvesen, B., see Øyehaug, E. 199
- Schneider, J.J.
 —, Triggs, E.J., Bourne, D.W.A., Stephens, I.D. and Haviland, A.M.
 Determination of oxycodone in human plasma by high-performance liquid chromatography with electrochemical detection 359
- Shiino, M., see Kadowaki, H. 329
- Shirota, F.N., see Demaster, E.G. 83
- Shukla, U.A.
 —, Stetson, P.L. and Wagner, J.G.
 Sensitive gas-liquid chromatographic-electron-capture detection method for determination of sobrerol in biological fluids 189
- Skews, H., see Jackman, G.P. 301
- Skinner, S.J.M.
 —, Grego, B., Hearn, M.T.W. and Liggins, G.C.
 The separation of collagen α -chains by reversed-phase high-performance liquid chromatography. Comparison of column alkyl stationary phases and temperature effects 111
- Sportoletti, G., see Cremonesi, P. 310
- Stephens, I.D., see Schneider, J.J. 359
- Stetson, P.L., see Shukla, U.A. 189
- Sunde, A., see Lundmo, P.I. 289
- Tackett, R., see Rydzik, R. 31
- Tamura, Z., see Uchino, K. 241
- Tanabe, S., see Toida, T. 133
- Tanaka, E.
 —, Hagino, S., Yoshida, T. and Kuroiwa, Y.
 Simultaneous determination of trimethadione and its metabolite in rat and human serum by high-performance liquid chromatography 393
- Tanaka, N., see Uchino, K. 241
- Terragno, A., see Rydzik, R. 31
- Thenot, J.P., see Padovani, P. 229
- Togari, A., see Oka, K. 43
- Togawa, T., see Toida, T. 133
- Toida, T.
 —, Togawa, T., Tanabe, S. and Imanari, T.
 Determination of cyanide and thiocyanate in blood plasma and red cells by high-performance liquid chromatography with fluorometric detection 133
- Triggs, E.J., see Schneider, J.J. 359
- Tsao, C.S.
 —, Young, M. and Rose, S.M.
 Quantitative analysis of ascorbic acid-2-sulfate by high-performance liquid chromatography with electrochemical detection 306
- Uchida, K., see Imaizumi, N. 295
- Uchino, K.
 —, Isozaki, S., Saitoh, Y., Nakagawa, F., Tamura, Z. and Tanaka, N.
 Quantitative determination of furosemide in plasma, plasma water, urine and ascites fluid by high-performance liquid chromatography 241
- Uemura, I., see Kadowaki, H. 329
- Vanazzi, G., see Beretta, E. 341
- Van Brummelen, P., see Kleinbloesem, C.H. 209
- Van Harten, J., see Kleinbloesem, C.H. 209
- Vessman, J., see Antonsson, A.-M. 181
- Vickers, S., see Musson, D.G. 251
- Vincek, W.C., see Musson, D.G. 251
- Wagner, J.G., see Shukla, U.A. 189
- Wiesler, D., see Alasandro, M. 1
- Williams, M. Carr, see Carr Williams, M. 101
- Williams, N.E., see Murray, G.R. 143
- Wozniak, J.M., see Lakings, D.B. 261
- Yamamoto, I., see Imaizumi, N. 295
- Yoo, B.-T., see Nohta, H. 93
- Yoshida, T., see Tanaka, E. 393
- Young, M., see Tsao, C.S. 306