

Author Index Vol. 434

- Adam, M., see Macek, J. 423
- Allain, P., see Premel-Cabic, A. 187
- Anjaneyulu, B., see Bhatia, S.C. 288
- Aronsson, E., see Odham, G. 31
- Assmann, B.
- and Haas, H.J.
Determination of the age dependency of the creatinine-related pseudouridine excretion in children's urine by high-performance liquid chromatography 202
- Babušíková, O., see Boháček, J. 439
- Balíková, M.
- and Kohlíček, J.
Rapid determination of ethylene glycol at toxic levels in serum and urine 469
- Bamonte, F., see Dionisotti, S. 169
- Banavalikar, M.M., see Bhatia, S.C. 288
- Barnoux, H., see Bougerolle, A.M. 232
- Baveja, S.K., see Padmalatha Devi, K. 265
- Belenkii, B.G.
Review. Use of laser detectors in capillary liquid chromatography 337
- Berger, J.A., see Bougerolle, A.M. 232
- Bergqvist, Y.
- and Churchill, F.C.
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1
- Bharucha, E.D., see Bhatia, S.C. 296, 303
- Bhatia, K.L., see Kamar, S. 228
- Bhatia, S.C.
- , Revankar, S.N., Anjaneyulu, B., Desai, N.D., Banavalikar, M.M., Sahajwalla, C.F. and Subrahmanyam, D.
High performance liquid chromatography of antifilarials. I. Simultaneous determination of 1-4-(4-nitrophenylaminophenylthiocarbamido-4-methyl)piperazine and three of its metabolites in biological fluids 288
- , Revankar, S.N., Bharucha, E.D., Desai, N.D. and Sahajwalla, C.G.
High-performance liquid chromatography of antifilarials. III. Determination of 2-*tert*-butyl-6-isothiocyanato-5-(piperidin-1-yl)-benzothiazole in biological fluids 303
- , Revankar, S.N., Doshi, K.J., Desai, N.D., Bharucha, E.D. and Sahajwalla, C.G.
High-performance liquid chromatography of antifilarials. II. Determination of S-(2-carboxyethyl)-N-(2-*tert*-butyl-5-methoxybenzothiazol-6-yl) dithiocarbamate and its active isothiocyanate metabolite in biological fluids 296
- Bidault, R., see Le Coguic, A. 320
- Bisse, E.
- and Wieland, H.
High-performance liquid chromatographic separation of human haemoglobins. Simultaneous quantitation of foetal and glycosylated haemoglobins 95
- Bökens, H.
- , Bourscheidt, C. and Müller, R.F.
Determination of bumetanide in plasma by high-performance liquid chromatography 327
- Boháček, J.
- , Hošek, B., Babušíková, O. and Hrivňáková, A.
Determination of adenosine deaminase and purine nucleoside phosphorylase activities using high-performance liquid chromatographic and radiochromatographic methods 439
- Bougerolle, A.M.
- , Chabard, J.L., Jbilou, M., Barnoux, H., Petit, J., Berger, J.A. and Dordain, G.
Simultaneous determination of trimipramine and its demethylated metabolites in plasma by gas chromatography-mass spectrometry 232
- Bourscheidt, C., see Bökens, H. 327
- Caccamo, F.
- , Carfagnini, G., Di Corcia, A. and Samperi, R.
Measurement of urinary estriol glucuronides during the menstrual cycle by high-performance liquid chromatography 61
- Carfagnini, G., see Caccamo, F. 61
- Černý, M., see Jegorov, A. 417
- Chabard, J.L., see Bougerolle, A.M. 232
- Churchill, F.C., see Bergqvist, Y. 1
- Clark, B.J., see Fell, A.F. 377
- Clarke, G.S., see Sheridan, M.E. 253
- Clifford, A.J., see Smidt, C.R. 21
- Coguic, A. Le, see Le Coguic, A. 320
- Cook, C.E., see Nakahara, Y. 247
- Corcia, A. Di, see Caccamo, F. 61
- Cserháti, T.
- and Szögyi, M.
Amino acid-amino acid interactions studied by charge-transfer chromatography 455

- Curtis, M.A., see Pullen, R.H. 271
- Dauphin, A., see Le Coguic, A. 320
- Dayer, P., see Padmalatha Devi, K. 265
- Desai, N.D., see Bhatia, S.C. 288, 296, 303
- Devi, K. Padmalatha, see Padmalatha Devi, K. 265
- Di Corcia, A., see Caccamo, F. 61
- Dionisotti, S.
- , Bamonte, F., Gamba, M. and Ongini, E.
High-performance liquid chromatographic determination of netilmicin in guinea-pig and human serum by fluorodinitrobenzene derivatization with spectrophotometric detection 169
- Doormaal, J.J. van, see Hoving, E.B. 395
- Dordain, G., see Bougerolle, A.M. 232
- Doshi, K.J., see Bhatia, S.C. 296
- Dubiel, W., see Toguzov, R.T. 447
- Dubois, J.P., see Sioufi, A. 239
- Ehrsson, H., see Hassan, M. 330
- Farinotti, R., see Le Coguic, A. 320
- Fell, A.F.
- , Noctor, T.A.G., Mama, J.E. and Clark, B.J.
Computer-aided optimisation of drug enantiomer separation in chiral high-performance liquid chromatography 377
- Frič, P., see Kocna, P. 429
- Gamba, M., see Dionisotti, S. 169
- Gerber, G., see Toguzov, R.T. 447
- Gibassier, D., see Sado, P.A. 157
- Haas, H.J., see Assmann, B. 202
- Hack, M.H.
- and Helmy, F.M.
Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a thin-layer chromatographic evaluation 215
- Hais, I.M., see Nobilis, M. 363
- Hase, S.
- , Ikenaka, K., Mikoshiba, K. and Ikenaka, T.
Analysis of tissue glycoprotein sugar chains by two-dimensional high-performance liquid chromatographic mapping 51
- Hassan, M.
- and Ehrsson, H.
Determination of busulfan in human plasma by gas chromatography with electron-capture detection 330
- Hayashi, S., see Matsukura, H. 196
- Hekkens, W.Th.J.M., see Kocna, P. 429
- Helmy, F.M., see Hack, M.H. 215
- Hill, H.M., see Selinger, K. 259
- Hirata, N.
- , Kasai, M., Yanagihara, Y. and Noguchi, K.
Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 71
- Holthuis, J.J.M., see Vendrig, D.E.M.M. 145
- Hoogmartens, J., see Kibwage, I.O. 177
- Hošek, B., see Boháček, J. 439
- Hoving, E.B.
- , Jansen, G., Volmer, M., Van Doormaal, J.J. and Muskiet, F.A.J.
Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary gas chromatography, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 395
- , Prins, J., Rutgers, H.M. and Muskiet, F.A.J.
Behaviour of plasmalogens during high-performance liquid chromatography on a silica column with a mobile phase containing phosphoric acid 411
- Hrivňáková, A., see Boháček, J. 439
- Ichinose, H., see Naoi, M. 209
- Ikenaka, K., see Hase, S. 51
- Ikenaka, T., see Hase, S. 51
- Jansen, G., see Hoving, E.B. 395
- Jbilou, M., see Bougerolle, A.M. 232
- Jegorov, A.
- , Tříška, J., Trnka, T. and Černý, M.
Separation of α -amino acid enantiomers by reversed-phase high-performance liquid chromatography after derivatization with o-phthaldialdehyde and a sodium salt of 1-thio- β -D-glucose 417
- Jones, A.D., see Smidt, C.R. 21
- Kaiser, D.G.
- , Vangiessen, G.J., Shah, J.A. and Weber, D.J.
Gas chromatographic determination of guanadrel in plasma and urine 135
- Kaiser, G., see Sioufi, A. 239
- Kamar, S.
- and Bhatia, K.L.
Characterization of buffalo lactotransferrin by polyacrylamide gel electrophoresis in 6 M urea 228
- Karnes, H.T., see Sarkar, M.A. 324
- Kasaffrek, E., see Kocna, P. 429
- Kasai, M., see Hirata, N. 71
- Kennedy, C.M., see Pullen, R.H. 271
- Kibwage, I.O.
- , Roets, E., Verbruggen, A., Hoogmartens, J. and Vanderhaeghe, H.
Thin-layer chromatographic study of the metabolites of erythromycins in the Wistar rat 177

- Kocna, P.
- , Frič, P., Kočová-Holáková, M., Slabý, J., Kasafírek, E. and Hekkens, W.Th. J.M. Isolation and analysis of peptidic fragments of α -gliadin using reversed-phase high-performance liquid chromatography 429
- Kočová-Holáková, M., see Kocna, P. 429
- Koel, M., see Nebinger, P. 43
- Kohlčěk, J., see Balíková, M. 469
- Kurachi, M., see Kurata, K. 278
- Kurata, K.
- , Kurachi, M. and Tanii, Y. High-performance liquid chromatographic determination of mianserin in plasma and brain and its application to pharmacokinetic studies in the rat 278
- LaCreta, F.P.
- , Olszewski, J.J. and Tew, K.D. Purification of glutathione S-transferases from rat liver and Walker 256 mammary carcinoma cells by high-performance liquid chromatography and a glutathione affinity column 83
- Lamiable, D.
- , Vistelle, R., Nguyen-Khac, M. and Millart, H. High-performance liquid chromatographic determination of dexamethasone in cerebrospinal fluid and plasma in the rabbit 315
- Le Coguic, A.
- , Bidault, R., Farinotti, R. and Dauphin, A. Determination of ofloxacin in plasma and urine by liquid chromatography 320
- Leemann, T., see Padmalatha Devi, K. 265
- Leverge, R., see Sado, P.A. 157
- Lichý, A., see Macek, J. 423
- McDowall, M., see Odham, G. 31
- Macek, J.
- , Lichý, A., Tesařová, E. and Adam, M. Separation of type IX collagen from other cartilage collagens by hydrophobic interaction chromatography 423
- Mama, J.E., see Fell, A.F. 377
- Maniwa, M., see Umeno, Y. 123
- Marunaka, T., see Umeno, Y. 123
- Matheou, D., see Selinger, K. 259
- Matsukura, H.
- , Suzuki, Y., Okada, T., Naiki, S., Hayashi, S. and Yamaji, Y. High-performance liquid chromatographic determination of acid and alkaline phosphatases in urine 196
- Matsushima, E., see Umeno, Y. 123
- Meiborg, G., see Volmer, M. 385
- Mezzetti, T.
- , Proietti, G. and Sbordoni, A. Simple method for the determination of hormonal iodine and exogenous iodine from X-ray contrast media drugs 308
- Michelsen, P., see Odham, G. 31
- Mikoshiba, K., see Hase, S. 51
- Millart, H., see Lamiable, D. 315
- Muda, M.
- , Pelizzoni, F., Sello, G. and Mussini, E. Determination of ergothioneine in red blood cells by high-performance liquid chromatography 191
- Müller, R.F., see Bökens, H. 327
- Muskiet, F.A.J., see Hoving, E.B. 395, 411
- , see Volmer, M. 385
- Mussini, E., see Muda, M. 191
- Nagatsu, T., see Naoi, M. 209
- Naiki, S., see Matsukura, H. 196
- Nakahara, Y.
- and Cook, C.E. Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of Δ^9 -tetrahydrocannabinol in plasma by high-performance liquid chromatography with electrochemical detection 247
- Naoi, M.
- , Ichinose, H., Takahashi, T. and Nagatsu, T. Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using high-performance liquid chromatography after derivatization with cyanide 209
- Nebinger, P.
- and Koel, M. Automated analysis of urinary 3-methoxy-4-hydroxymandelic acid using ion-pair chromatography and fluorimetric detection 43
- Nguyen-Khac, M., see Lamiable, D. 315
- Nobilis, M.
- and Hais, I.M. Direct absorptiometric quantitation in high-performance liquid chromatography in the absence of concomitant reference standards 363
- Noctor, T.A.G., see Fell, A.F. 377
- Noguchi, K., see Hirata, N. 71
- Odham, G.
- , Valeur, A., Michelsen, P., Aronsson, E. and McDowall, M. Highly sensitive determination and characterization of intact cellular ester-linked phospholipids using liquid chromatography-plasma spray mass spectrometry 31

- Okada, T., see Matsukura, H. 196
- Olivares, J.
- and Verdys, M.
Isocratic high-performance liquid chromatographic method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
- Olszewski, J.J., see LaCreta, F.P. 83
- Ongini, E., see Dionisotti, S. 169
- Padmalatha Devi, K.
- , Ranga Rao, K.V., Baveja, S.K., Leemann, T. and Dayer, P.
Determination of alprenolol and metoprolol in plasma by column liquid chromatography 265
- Pelizzoni, F., see Muda, M. 191
- Peteghem, C. Van, see Rosier, J. 222
- Petit, J., see Bougerolle, A.M. 232
- Pimenov, A.M., see Toguzov, R.T. 447
- Pommier, F., see Sioufi, A. 239
- Premel-Cabic, A.
- and Allain, P.
Simultaneous liquid chromatographic determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 187
- Prins, J., see Hoving, E.B. 411
- Proietti, G., see Mezzetti, T. 308
- Prokudin, V.Yu., see Toguzov, R.T. 447
- Pullen, R.H.
- , Kennedy, C.M. and Curtis, M.A.
Direct plasma injection using internal surface reversed-phase high-performance liquid chromatography: feasibility study using propofol as a model compound 271
- Ranga Rao, K.V., see Padmalatha Devi, K. 265
- Rao, K.V. Ranga, see Padmalatha Devi, K. 265
- Revankar, S.N., see Bhatia, S.C. 288, 296, 303
- Richter, K.
Determination of clozapine in human serum by capillary gas chromatography 465
- Roets, E., see Kibwage, I.O. 177
- Rosier, J.
- and Van Peteghem, C.
Detection of oxidatively modified 2'-deoxyguanosine-3'-monophosphate, using ³²P-postlabelling and anion-exchange thin-layer chromatography 222
- Rutgers, H.M., see Hoving, E.B. 411
- Sado, P.A.
- , Gibassier, D. and Leverage, R.
Dosage simultané de la vinburnine et de la 6-hydroxyvinburnine par chromatographie liquide à haute performance 157
- Sahajwalla, C.G., see Bhatia, S.C. 288, 296, 303
- Samperi, R., see Caccamo, F. 61
- Sarkar, M.A.
- and Karnes, H.T.
Comparison of solid-phase extraction and liquid-liquid extraction for high-performance liquid chromatographic analysis of antipyrine in plasma 324
- Sbordoni, A., see Mezzetti, T. 308
- Selinger, K.
- , Matheou, D. and Hill, H.M.
High-performance liquid chromatographic method for the determination of miconazole in vaginal fluid 259
- Sello, G., see Muda, M. 191
- Shah, J.A., see Kaiser, D.G. 135
- Sheridan, M.E.
- and Clarke, G.S.
Improved high-performance liquid chromatographic determination of doxycycline in serum and urine using solid-phase extraction columns 253
- Shimizu, T., see Shioya, Y. 283
- Shioya, Y.
- and Shimizu, T.
High-performance liquid chromatographic procedure for the determination of a new anti-gastric ulcer agent, 2-(4-chlorobenzoylamino)-3-[2(1H)-quinolinon-4-yl]propionic acid, in human plasma and urine 283
- Sioufi, A.
- , Pommier, F., Kaiser, G. and Dubois, J.P.
Determination of benazepril, a new angiotensin-converting enzyme inhibitor, and its active metabolite, benazeprilat, in plasma and urine by capillary gas chromatography-mass-selective detection 239
- Slabý, J., see Kocna, P. 429
- Smidt, C.R.
- , Jones, A.D. and Clifford, A.J.
Gas chromatography of retinol and α -tocopherol without derivatization 21
- Subrahmanyam, D., see Bhatia, S.C. 288
- Suzuki, Y., see Matsukura, H. 196
- Szögyi, M., see Cserháti, T. 455
- Takahashi, T., see Naoi, M. 209
- Tanii, Y., see Kurata, K. 278
- Teeuwssen, J., see Vendrig, D.E.M.M. 145
- Tesařová, E., see Macek, J. 423
- Tew, K.D., see LaCreta, F.P. 83

- Tikhonov, Y.V., see Toguzov, R.T. 447
- Toguzov, R.T.
- , Tikhonov, Y.V., Pimenov, A.M., Prokudin, V.Yu., Dubiel, W., Ziegler, M. and Gerber, G.
Optimization of the ion-pair high-performance liquid chromatographic separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 447
- Tříska, J., see Jedorov, A. 417
- Trnka, T., see Jedorov, A. 417
- Umeno, Y.
- , Matsushima, E., Maniwa, M. and Marunaka, T.
Gas chromatographic-negative-ion chemical ionization mass spectrometric determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolites in human plasma and urine 123
- Valeur, A., see Odham, G. 31
- Vanderhaege, H., see Kibwage, I.O. 177
- Van Doormaal, J.J., see Hoving, E.B. 395
- Vangiessen, G.J., see Kaiser, D.G. 135
- Van Peteghem, C., see Rosier, J. 222
- Vendrig, D.E.M.M.
- , Teeuwesen, J. and Holthuis, J.J.M.
Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 145
- Verbruggen, A., see Kibwage, I.O. 177
- Verdys, M., see Olivares, J. 111
- Vistelle, R., see Lamiable, D. 315
- Volmer, M., see Hoving, E.B. 395
- , Meiborg, G. and Muskiet, F.A.J.
Simultaneous capillary gas chromatographic profiling of medium- and long-chain fatty acid methyl esters with split injection. Correction for injection-related discrimination by the 'bracketing' method 385
- Weber, D.J., see Kaiser, D.G. 135
- Wieland, H., see Bisse, E. 95
- Yamaji, Y., see Matsukura, H. 196
- Yanagihara, Y., see Hirata, N. 71
- Ziegler, M., see Toguzov, R.T. 447