

# Related articles published in *Journal of Chromatography A*, Vol. 705

High-speed high-performance liquid chromatography of peptides and proteins by H. Chen and Cs. Horváth (New Haven, CT, USA) . . . . .	3
Protein mass spectrometry: applications to analytical biotechnology (Review) by D.N. Nguyen, G.W. Becker and R.M. Riggin (Indianapolis, IN, USA) . . . . .	21
Chemical methods of protein sequence analysis (Review) by J.M. Bailey (Duarte, CA, USA) . . . . .	47
Isoelectric focusing as a tool for the investigation of post-translational processing and chemical modifications of proteins (Review) by E. Gianazza (Milan, Italy) . . . . .	67
Fluorophore-assisted carbohydrate electrophoresis. Technology and applications (Review) by G.-F. Hu (Boston, MA, USA) . . . . .	89
Host cell contaminant protein assay development for recombinant biopharmaceuticals (Review) by L.C. Eaton (Kalamazoo, MI, USA) . . . . .	105
Moisture content in proteins: its effects and measurement (Review) by J.K. Towns (Indianapolis, IN, USA) . . . . .	115
Processing of C-terminal lysine and arginine residues of proteins isolated from mammalian cell culture (Review) by R.J. Harris (San Francisco, CA, USA) . . . . .	129
Capillary electrophoresis of <i>S. nuclease</i> mutants by F. Kálmán, S. Ma, R.O. Fox and Cs. Horváth (New Haven, CT, USA) . . . . .	135
Analysis of recombinant human growth hormone in <i>Escherichia coli</i> fermentation broth by micellar high-performance liquid chromatography by M.A. Strega and A.L. Lagu (Indianapolis, IN, USA) . . . . .	155
Detection of neu differentiation factor with a biospecific affinity sensor during chromatography by H.S. Lu, D. Chang, D. Brankow and D. Wen (Thousand Oaks, CA, USA) . . . . .	163
Monitoring of optical isomers of some conformationally constrained amino acids with tetrahydroisoquinoline or tetraline ring structures. Part II by A. Péter, G. Tóth, E. Olajos and F. Fülöp (Szeged, Hungary) and D. Tourwé (Brussels, Belgium) . . . . .	257
Chromatographic behaviour of opioid peptides containing $\beta$ -methylphenylalanine isomers by A. Péter, G. Tóth and E. Olajos (Szeged, Hungary) and D. Tourwé (Brussels, Belgium) . . . . .	267
Enantiomeric separation of basic drugs using N-benzyloxycarbonylglycyl-L-proline as counter ion in methanol by N.-H. Huynh (Uppsala, Sweden), A. Karlsson (Mölnådal, Sweden) and C. Pettersson (Uppsala, Sweden) . . . . .	275
High-performance capillary electrophoretic separation of human serum albumin using a neutral coated capillary by K.A. Denton and R. Harris (Nottingham, UK) . . . . .	335
Separation of stereoisomers of aminoglutethimide using three capillary electrophoretic techniques by V.C. Anigbogu (Decatur, GA, USA) and C.L. Copper and M.J. Sepaniak (Knoxville, TN, USA) . . . . .	343
Application of modified cyclodextrins in capillary electrophoresis for enantiomeric resolution of propranolol and analogues by M.W. Matchett, S.K. Branch and T.M. Jefferies (Bath, UK) . . . . .	351
Purification of preparative quantities of group B <i>Streptococcus</i> type III oligosaccharides by L.C. Paoletti and K.D. Johnson (Boston, MA, USA) . . . . .	363