

## Related articles published in *Journal of Chromatography A*, Vols. 749–750

### Volume 749

- Immobilized liposome chromatography of drugs on capillary continuous beds for model analysis of drug–membrane interactions  
by Y. Zhang, C.-M. Zeng, Y.-M. Li, S. Hjertén and P. Lundahl (Uppsala, Sweden) . . . . . 13
- Isolation of dinophysistoxin-2 and the high-performance liquid chromatographic analysis of diarrhetic shellfish toxins using  
derivatisation with 1-bromoacetylpyrene  
by S.S. Kelly, A.G. Bishop, E.P. Carmody and K.J. James (Cork, Ireland) . . . . . 33
- Resolution of enantiomeric steroids by high-performance liquid chromatography on chiral stationary phases  
by M. Kummer and H.-J. Palme (Jena, Germany) and G. Werner (Leipzig, Germany) . . . . . 61
- Integrated derivatization–chemiluminescence detection system for the determination of  $\beta$ -carboline alkaloids by high-performance  
liquid chromatography  
by J. Cepas, M. Silva and D. Pérez-Bendito (Córdoba, Spain) . . . . . 73
- Liquid chromatographic method for the determination of nicotine in pharmaceutical formulations  
by A.K. Dash and S.-T. Wong (Omaha, NE, USA) . . . . . 81
- Packed-column supercritical fluid chromatography of a new dihydropyridine drug based on direct injection of emulsion samples  
by L. Karlsson, O. Gyllenhaal, A. Karlsson and J. Gottfries (Mölnådal, Sweden) . . . . . 193
- Capillary electrophoretic examination of underivatized oligosaccharide mixtures released from immunoglobulin G antibodies and  
CTLA4Ig fusion protein  
by K.F. Greve and D.E. Hughes (Syracuse, NY, USA) and B.L. Karger (Boston, MA, USA) . . . . . 237
- Separation of pharmaceutically important estrogens by micellar electrokinetic chromatography  
by S.K. Poole (London, UK) and C.F. Poole (London, UK and Detroit, MI, USA) . . . . . 247
- A more sensitive way to determine iron using an iron(II)–1,10-phenanthroline complex and capillary electrophoresis  
by J. Xu, P. Che and Y. Ma (Kirksville, MO, USA) . . . . . 287

### Volume 750

- Application of multidimensional affinity high-performance liquid chromatography and electrospray ionization liquid chromatography–  
mass spectrometry to the characterization of glycosylation in single-chain plasminogen activator. Initial results  
by A. Apffel, J.A. Chakel and W.S. Hancock (Palo Alto, CA, USA) and C. Souders, T. M'Timkulu and E. Pungor, Jr.  
(Brisbane, CA, USA) . . . . . 35
- Gas and liquid chromatography coupled to tandem mass spectrometry for the multiresidue analysis of  $\beta$ -agonists in biological matrices  
by G. Van Vyncht (Liège, Belgium), S. Preece (Altrincham, UK) and P. Gaspar, G. Maghuin-Rogister and E. DePauw (Liège,  
Belgium) . . . . . 43
- Hyphenation of capillary electrophoresis to inductively coupled plasma mass spectrometry as an element-specific detection method for  
metal specification  
by B. Michalke and P. Schramel (Oberschleißheim, Germany) . . . . . 51
- Quantitative determination of norethisterone acetate in human plasma by capillary gas chromatography with mass-selective detection  
by F. Pommier, A. Sioufi and J. Godbillon (Rueil Malmaison, France) . . . . . 75
- Comparison of the possibilities of gas chromatography–mass spectrometry and tandem mass spectrometry systems for the analysis of  
anabolics in biological material  
by H.F. De Brabander and P. Batjoens (Merelbeke, Belgium), D. Courtheyn and J. Vercammen (Gentbrugge, Belgium) and K.  
De Wasch (Merelbeke, Belgium) . . . . . 105
- Rapid and high-performance analysis of thyreostatic drug residues in urine using gas chromatography–mass spectrometry  
by P. Batjoens, H.F. De Brabander and K. De Wasch (Merelbeke, Belgium) . . . . . 127
- Gas chromatographic–tandem mass spectrometric analysis of clenbuterol residues in faeces  
by P. Batjoens (Merelbeke, Belgium), D. Courtheyn (Gentbrugge, Belgium), H.F. De Brabander (Merelbeke, Belgium), J.  
Vercammen (Gentbrugge, Belgium), K. De Wasch (Merelbeke, Belgium) and M. Logghe (Gentbrugge, Belgium) . . . . . 133
- Application of on-line capillary high-performance liquid chromatography–nuclear magnetic resonance spectrometry coupling for the  
analysis of vitamin A derivatives  
by K. Albert, G. Schlotterbeck and L.-H. Tseng (Tübingen, Germany) and U. Braumann (Rheinstetten, Germany) . . . . . 303

Determination of butyl- and phenyltin compounds in biological material by gas chromatography–flame photometric detection after ethylation with sodium tetraethylborate by C. Carlier-Pinasseau, A. Astruc, G. Lespes and M. Astruc (Pau, France) . . . . .	317
Automated determination of verapamil and norverapamil in human plasma with on-line coupling of dialysis to high-performance liquid chromatography and fluorometric detection by A. Ceccato, P. Chiap, Ph. Hubert, B. Toussaint and J. Crommen (Liège, Belgium) . . . . .	351
Quantification of cyclamate and cyclohexylamine in urine samples using high-performance liquid chromatography with trinitrobenzenesulfonic acid pre-column derivatization (Short communication) by I. Casals, M. Reixach, J. Amat, M. Fuentes and L. Serra-Majem (Barcelona, Spain) . . . . .	397
Toxicological evaluation of harmful substances by in situ enzymatic and biological detection in high-performance thin-layer chromatography (Short communication) by C. Weins and H. Jork (Saarbrücken, Germany) . . . . .	403