



ELSEVIER

Journal of Chromatography B, 730 (1999) 283–285

JOURNAL OF
CHROMATOGRAPHY B

Compound Index Vol. 730

Adenosine triphosphate

Improved high-performance liquid chromatographic assay for the determination of “high-energy” phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

Amino acids

High-performance liquid chromatographic analysis of physiological amino acids in human brain tumors by pre-column derivatization with phenylisothiocyanate 730(1999)81

Artemisinin

Rapid determination of artemisinin and related analogues using high-performance liquid chromatography and an evaporative light scattering detector 730(1999)71

Ascorbic acid

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Benzoin

Bioanalysis of *m*-iodobenzylguanidine in plasma by high-performance liquid chromatography after derivatization with benzoin 730(1999)193

Busulfan

Determination of tetrahydrothiophene formation as a probe of in vitro busulfan metabolism by human glutathione *S*-transferase A1-1: use of a highly sensitive gas chromatographic–mass spectrometric method 730(1999)25

Determination of busulfan in human plasma using high-performance liquid chromatography with pre-column derivatization and fluorescence detection 730(1999)33

1,3-Butadiene

Gas chromatographic determination of 3-butene-1,2-diol in urine samples after 1,3-butadiene exposure 730(1999)257

3-Butene-1,2-diol

Gas chromatographic determination of 3-butene-1,2-diol in urine samples after 1,3-butadiene exposure 730(1999)257

Catecholamines

Determination of catecholamines in human plasma by high-performance liquid chromatography with electrochemical detection 730(1999)201

Sensitive and specific method for the simultaneous determination of natural and synthetic catecholamines and 3,4-dihydroxyphenylglycol in microdialysis samples 730(1999)213

Chondroitin sulfate

Ultrasensitive capillary electrophoresis of sulfated disaccharides in chondroitin/dermatan sulfates by laser-induced fluorescence after derivatization with 2-aminoacridone 730(1999)129

Cortisol

Development of a urinary free cortisol assay using solid-phase extraction–capillary electrophoresis 730(1999)123

Creatine

Improved high-performance liquid chromatographic assay for the determination of “high-energy” phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

Creatinine

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Dehydroascorbic acid

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Dehydroisoascorbic acid

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Dermatan sulfate

Ultrasensitive capillary electrophoresis of sulfated disaccharides in chondroitin/dermatan sulfates by laser-induced fluorescence after derivatization with 2-aminoacridone 730(1999)129

3,4-Dihydroxyphenylglycol

Sensitive and specific method for the simultaneous determination of natural and synthetic catecholamines and 3,4-dihydroxyphenylglycol in microdialysis samples 730(1999)213

Disaccharides, sulfated

Ultrasensitive capillary electrophoresis of sulfated disaccharides in chondroitin/dermatan sulfates by laser-induced fluorescence after derivatization with 2-aminoacridone 730(1999)129

Domperidone

Determination of domperidone in human serum and human breast milk by high-performance liquid chromatography–electrospray mass spectrometry 730(1999)9

Efetirizine

Quantitation of efetirizine in human plasma and urine using automated solid-phase extraction and column-switching high-performance liquid chromatography 730(1999)239

Everninomycin

Simple and sensitive high-performance liquid chromatographic method for the determination of an everninomycin, SCH 27899, in rat plasma 730(1999)55

Furosemide

Simplified method to quantify furosemide in urine by high-performance liquid chromatography and ultraviolet detection 730(1999)49

Galanthamine

Enantiomeric resolution of galanthamine and related drugs used in anti-Alzheimer therapy by means of capillary zone electrophoresis employing derivatized cyclodextrin selectors 730(1999)167

Glutathione S-transferase A1-1

Determination of tetrahydrothiophene formation as a probe of in vitro busulfan metabolism by human glutathione S-transferase A1-1: use of a highly sensitive gas chromatographic–mass spectrometric method 730(1999)25

Hippuric acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

3-Hydroxypyridin-4-one

Gradient ion-pair high-performance liquid chromatographic method for analysis of 3-hydroxypyridin-4-one iron chelators 730(1999)135

Hydroxypyrimidine

Determination of metabolites of pirimicarb in human urine by gas chromatography–mass spectrometry 730(1999)229

Inosine monophosphate

Improved high-performance liquid chromatographic assay for the determination of “high-energy” phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

***m*-Iodobenzylguanidine**

Bioanalysis of *m*-iodobenzylguanidine in plasma by high-performance liquid chromatography after derivatization with benzoin 730(1999)193

Iron chelators

Gradient ion-pair high-performance liquid chromatographic method for analysis of 3-hydroxypyridin-4-one iron chelators 730(1999)135

Isoascorbic acid

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Mandelic acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Methylhippuric acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Methyl 6-mercaptopurine nucleotides

High-performance liquid chromatographic determination of methyl 6-mercaptopurine nucleotides (Me6-MPN) in red blood cells: analysis of Me6-MPN per se or Me6-MPN derivative? 730(1999)273

Comments on “High-performance liquid chromatographic determination of methyl 6-mercaptopurine nucleotides in red blood cells” Reply to R. Boulieu and T. Dervieux 730(1999)275

Modafinil

Determination of the D- and L-enantiomers of modafinil in human plasma utilizing liquid–liquid extraction and high-performance liquid chromatography 730(1999)1

1-Naphthylisothiocyanate

Semi-automatic liquid chromatographic analysis of pamidronate in urine after derivatization with 1-naphthylisothiocyanate 730(1999)95

Nefazodone

Determination of nefazodone and its pharmacologically active metabolites in human blood plasma and breast milk by high-performance liquid chromatography 730(1999)249

NE-100

Ultrasensitive determination of NE-100, a novel sigma ligand, in human plasma by liquid chromatography and electrospray ionization tandem mass spectrometry combined with a column-switching technique 730(1999)61

2-(2-Nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione

Determination of 2-(2-nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione in plasma by direct injection into a coupled column liquid chromatographic system 730(1999)177

Pamidronate

Semi-automatic liquid chromatographic analysis of pamidronate in urine after derivatization with 1-naphthylisothiocyanate 730(1999)95

Phenylisothiocyanate

High-performance liquid chromatographic analysis of physiological amino acids in human brain tumors by pre-column derivatization with phenylisothiocyanate 730(1999)81

Phosphocreatine

Improved high-performance liquid chromatographic assay for the determination of "high-energy" phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

Pirimicarb

Determination of metabolites of pirimicarb in human urine by gas chromatography-mass spectrometry 730(1999)229

1-Pyrenyldiazomethane

Headspace solid-phase microextraction with 1-pyrenyldiazomethane in-fibre derivatisation for analysis of faecal short-chain fatty acids 730(1999)113

Retinol

Simultaneous determination of serum retinol and α - and γ -tocopherol levels in type II diabetic patients using high-performance liquid chromatography with fluorescence detection 730(1999)265

Rofecoxib

Determination of rofecoxib, a cyclooxygenase-2 specific inhibitor, in human plasma using high-performance liquid chromatography with post-column photochemical derivatization and fluorescence detection 730(1999)221

Short-chain fatty acids

Headspace solid-phase microextraction with 1-pyrenyldiazomethane in-fibre derivatisation for analysis of faecal short-chain fatty acids 730(1999)113

Steroids

Development of a urinary free cortisol assay using solid-phase extraction-capillary electrophoresis 730(1999)123

Styrene

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Taurine

Ion chromatographic determination of taurine in medicine, nutrient capsule and human urine with electrochemical detection 730(1999)161

Tetrahydrothiophene

Determination of tetrahydrothiophene formation as a probe of in vitro busulfan metabolism by human glutathione *S*-transferase A1-1: use of a highly sensitive gas chromatographic-mass spectrometric method 730(1999)25

Tocopherol

Simultaneous determination of serum retinol and α - and γ -tocopherol levels in type II diabetic patients using high-performance liquid chromatography with fluorescence detection 730(1999)265

Tolcapone

Determination of the catechol-*O*-methyltransferase inhibitor tolcapone and three of its metabolites in animal and human plasma and urine by reversed-phase high-performance liquid chromatography with ultraviolet detection 730(1999)153

Toluene

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Uric acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Vitamins

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Application of two-dimensional high-performance liquid chromatography-mass spectrometry with particle beam interface 730(1999)141

Simultaneous determination of serum retinol and α - and γ -tocopherol levels in type II diabetic patients using high-performance liquid chromatography with fluorescence detection 730(1999)265

Xylene

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41