

CUMULATIVE SUBJECT INDEX

J. CHROMATOGR. B VOLS. 726–750

A

***Achillea* species**

Qualitative and quantitative determination of sesquiterpenoids in *Achillea* species by reversed-phase high-performance liquid chromatography, mass-spectrometry and thin-layer chromatography 729(1999)361

Affinity chromatography

Aptamer affinity chromatography: combinatorial chemistry applied to protein purification 731(1999)275

Affinity membranes

Human serum albumin chromatography by Cibacron Blue F3GA-derived microporous polyamide hollow-fiber affinity membranes 746(2000)123

Affinity partitioning

Purification of rabbit lacrimal gland plasma membranes by aqueous two-phase affinity partitioning 743(2000)397

Aggregation

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Allelic discrimination

Allelic discrimination by denaturing high-performance liquid chromatography 739(2000)345

Anaerobic bacteria

Routine analysis of short-chain fatty acids for anaerobic bacteria identification using capillary electrophoresis and indirect ultraviolet detection 741(2000)89

Aptamer affinity chromatography

Aptamer affinity chromatography: combinatorial chemistry applied to protein purification 731(1999)275

Aqueous two-phase systems

Partial purification of penicillin acylase from *Escherichia coli* in poly(ethylene glycol)–sodium citrate aqueous two-phase systems 734(1999)15

Effect of biological suspensions on the position of the binodal curve in aqueous two-phase systems 743(2000)5

Model process for separation based on unfolding and refolding of chymotrypsin inhibitor 2 in thermoseparating polymer two-phase systems 743(2000)13

Investigation of mathematical methods for efficient optimisation of aqueous two-phase extraction 743(2000)21

New aqueous two-phase system based on cashew-nut tree gum and poly(ethylene glycol) 743(2000)79

Drowning-out crystallisation of sodium sulphate using aqueous two-phase systems 743(2000)101

Naphthol- and resorcinol-based azo dyes as metal ion complexants in aqueous biphasic systems 743(2000)107

Polymer recycling in aqueous two-phase extractions using thermoseparating ethylene oxide–propylene oxide copolymers 743(2000)115

Investigation of aqueous biphasic systems for the separation of lignins from cellulose in the paper pulping process 743(2000)127

Solvatochromic studies in polyethylene glycol–salt aqueous biphasic systems 743(2000)137

- Viscosity prediction of polyethylene glycol–dextran–water solutions used in aqueous two-phase systems 743(2000)181
- Relative hydrophobicity of organic compounds measured by partitioning in aqueous two-phase systems 743(2000)187
- Kinetics of phase separation under different process and design parameters in aqueous two-phase systems 743(2000)195
- Protein refolding using stimuli-responsive polymer-modified aqueous two-phase systems 743(2000)215
- Determination of the dissociation constant of valine from acetohydroxy acid synthase by equilibrium partition in an aqueous two-phase system 743(2000)225
- Poly(ethylene glycol)-based aqueous biphasic systems: effect of temperature on phase equilibria and on partitioning of 1,10-phenanthroline–copper(II) sulphate complex 743(2000)231
- Partitioning of whey proteins, bovine serum albumin and porcine insulin in aqueous two-phase systems 743(2000)235
- Bioaffinity extraction of glucoamylase in aqueous two-phase systems using starch as free bioligand 743(2000)241
- Study on heterogeneity of β -glucosidase from *Aspergillus* species by using counter-current distribution 743(2000)247
- Bovine serum albumin partitioning in an aqueous two-phase system: Effect of pH and sodium chloride concentration 743(2000)255
- Partitioning of bovine serum albumin in an aqueous two-phase system: optimization of partition coefficient 743(2000)259
- Application of aqueous two-phase partition to the production of homogeneous preparations of fluorescently labelled human serum albumin 743(2000)263
- Peptides partitioning in an aqueous dextran–polyethylene glycol two-phase system 743(2000)271
- Purification of acetohydroxy acid synthase by separation in an aqueous two-phase system 743(2000)281
- Purification of soybean peroxidase (*Glycine max*) by metal affinity partitioning in aqueous two-phase systems 743(2000)287
- Partitioning of peptides and recombinant protein–peptide fusions in thermoseparating aqueous two-phase systems: effect of peptide primary structure 743(2000)295
- Partitioning behavior of amino acids in aqueous two-phase systems with recyclable volatile salts 743(2000)317
- Phase behavior and protein partitioning in aqueous two-phase systems of cationic–anionic surfactant mixtures 743(2000)327
- Partitioning of xylanolytic complex from *Penicillium janthinellum* by an aqueous two-phase system 743(2000)339
- Extraction in aqueous two-phase systems of alkaline xylanase produced by *Bacillus pumilus* and its application in kraft pulp bleaching 743(2000)349
- Applications of aqueous two-phase partition to isolation of membranes from plants: A periodic NADH oxidase activity as a marker for right side-out plasma membrane vesicles 743(2000)369
- Aqueous two-phase partition applied to the isolation of plasma membranes and Golgi apparatus from cultured mammalian cells 743(2000)377
- Affinity partitioning of biotinylated mixed liposomes: effect of charge on biotin–NeutrAvidin interaction 743(2000)389
- Purification of rabbit lacrimal gland plasma membranes by aqueous two-phase affinity partitioning 743(2000)397
- Aroma compounds recovery from mycelial cultures in aqueous two-phase processes 743(2000)403
- Recovery in aqueous two-phase systems of nanoparticulates applied as surrogate mimics for viral gene therapy vectors 743(2000)409
- Simultaneous accumulation of low-molecular-mass RNA at the interface along with accumulation of high-molecular-mass RNA on aqueous two-phase system partitioning 743(2000)421
- Cold cataracts: a naturally occurring aqueous two-phase systems 743(2000)431
- Artificial lectin**
Synthesis and evaluation of affinity adsorbents for glycoproteins: an artificial lectin 746(2000)265
- Aspergillus***
Study on heterogeneity of β -glucosidase from *Aspergillus* species by using counter-current distribution 743(2000)247
- Atherosclerosis**
High-performance liquid chromatographic assay of hydroperoxide levels in oxidatively modified lipoproteins 731(1999)223
- Automated in-tube solid-phase microextraction**
Automated in-tube solid-phase microextraction–liquid chromatography–electrospray ionization mass spectrometry for the determination of ranitidine 731(1999)353
- Automated protein precipitation**
Automated protein precipitation by filtration in the 96-well format 734(1999)257

B

Bacillus pumilus

Extraction in aqueous two-phase systems of alkaline xylanase produced by *Bacillus pumilus* and its application in kraft pulp bleaching 743(2000)349

Bacillus subtilis

New approach for separating *Bacillus subtilis* metalloprotease and α -amylase by affinity chromatography and for purifying neutral protease by hydrophobic chromatography 737(2000)277

***Bacillus subtilis* b213**

Purification of the fengycin synthetase multienzyme system from *Bacillus subtilis* b213 737(2000)267

Baculovirus

Multi-step purification strategy for RANTES wild-type and mutated analogues expressed in a baculovirus system 737(2000)47

Binding constants

Simultaneous determination of enantioselective plasma protein binding of aminohydantoins by ultrafiltration and chiral high-performance liquid chromatography 732(1999)31

Binodal curve

Effect of biological suspensions on the position of the binodal curve in aqueous two-phase systems 743(2000)5

Bioanalysis

Solid-phase microextraction for the analysis of biological samples 745(2000)49

Biochemical analysis

Solid-phase microextraction for the analysis of biological samples 745(2000)49

Biological monitoring

Biomonitoring of environmental polycyclic aromatic hydrocarbon exposure by simultaneous measurement of urinary phenanthrene, pyrene and benzo[*a*]pyrene hydroxides 739(2000)225

Biomimetic ligands

Affinity chromatography on immobilized “biomimetic” ligands. Synthesis, immobilization and chromatographic assessment of an immunoglobulin G binding ligand 740(2000)1

Bi-modal integrated amperometric detection

Analyzing mixtures of amino acids and carbohydrates using bi-modal integrated amperometric detection 732(1999)193

Blood

Simultaneous blood and brain sampling of cephalexin in the rat by microdialysis and microbore liquid chromatography: application to pharmacokinetics studies 740(2000)203

Brain

Simultaneous blood and brain sampling of cephalexin in the rat by microdialysis and microbore liquid chromatography: application to pharmacokinetics studies 740(2000)203

Breath sampling

Determination of isoprene in human expired breath using solid-phase microextraction and gas chromatography–mass spectrometry 739(2000)183

Breath tests

Variation in volatile organic compounds in the breath of normal humans 729(1999)75

Brush–particle interactions

Analysis of brush-particle interactions using self-consistent-field theory 743(2000)41

C

Candida intermedia

D-Xylose metabolism by *Candida intermedia*: isolation and characterisation of two forms of aldose reductase with different coenzyme specificities 737(2000)195

Cashew-nut tree gum

New aqueous two-phase system based on cashew-nut tree gum and poly(ethylene glycol) 743(2000)79

Cation-exchange

Determination of clenbuterol in bovine urine using gas chromatography–mass spectrometry following clean-up on an ion-exchange resin 728(1999)67

Cattle and diphasic dialysis

Determination of clenbuterol residues in bovine hair by using diphasic dialysis and gas chromatography–mass spectrometry 726(1999)133

Cell wall composition

Efficient method for preparation of highly purified lipopolysaccharides by hydrophobic interaction chromatography 732(1999)39

Chemical combinatorial library

New strategy for the design of ligands for the purification of pharmaceutical proteins by affinity chromatography 740(2000)17

Chemical development

Capillary electrophoresis in the pharmaceutical industry: applications in discovery and chemical development 745(2000)177

Chemical pulping

Investigation of aqueous biphasic systems for the separation of lignins from cellulose in the paper pulping process 743(2000)127

Chips

Nanotechnology in bio/clinical analysis 739(2000)139

Chip-based nanovials

Liquid lid for biochemical reactions in chip-based nanovials 745(2000)137

Chiral stationary phases, LC

Rapid chiral high-performance liquid chromatographic assay for salmeterol and α -hydroxysalmeterol. Application to in vitro metabolism studies 729(1999)225

Chromatographic investigation on the binding site characteristics of quail egg-white riboflavin binding protein as a chiral stationary phase 738(2000)343

Enantioselective analysis of disopyramide and mono-*N*-dealkyldisopyramide in plasma and urine by high-performance liquid chromatography on an amylose-derived chiral stationary phase 744(2000)299

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VIII. Chiral recognition ability of partially and completely deglycosylated ovoglycoprotein 745(2000)149

Chiral separation by simultaneous use of vancomycin as stationary phase chiral selector and chiral mobile phase additive 745(2000)159

Chloroplast stroma

Purification and characterization of an aminopeptidase from the chloroplast stroma of barley leaves by chromatographic and electrophoretic methods 737(2000)285

Cholesterol synthesis

Determination of isoprene in human expired breath using solid-phase microextraction and gas chromatography–mass spectrometry 739(2000)183

Chromatofocusing

Hydrophobic character of surface regions and total hydrophobicity of four variants of chromosomal class C β -lactamase from *Pseudomonas aeruginosa* are identical. Chromatographic comparison of the hydrophobic character of the variants and the effect of focusing buffer composition on the separation of the variants by chromatofocusing with internal and external pH gradients 746(2000)161

Circular dichroism

Conformational transition and mass transfer in extraction of proteins by AOT–alcohol–isooctane reverse micellar systems 743(2000)203

Cold cataracts

Cold cataracts: a naturally occurring aqueous two-phase systems 743(2000)431

Column overloading

Chromatographic assay of pharmaceutical compounds under column overloading 741(2000)55

Column switching

Achiral and chiral high-performance liquid chromatography of verapamil and its metabolites in serum samples 732(1999)395

Direct high-performance liquid chromatographic determination of (*R*)- and (*S*)-propranolol in rat microdialysate using on-line column switching procedures 739(2000)151

High-performance liquid chromatographic method for the determination of benzodiazepines in plasma or serum using the column-switching technique 742(2000)381

Trace determination of urinary 3-hydroxybenzo[*a*]pyrene by automated column-switching high-performance liquid chromatography 748(2000)337

Column-stitching

Automated column-switching high-performance liquid chromatography method for the determination of 1-hydroxypyrene in human urine 732(1999)91

Combinatorial chemistry

Affinity chromatography on immobilized “biomimetic” ligands. Synthesis, immobilization and chromatographic assessment of an immunoglobulin G binding ligand 740(2000)1

Directly coupled HPLC–NMR and HPLC–NMR–MS in pharmaceutical research and development 748(2000)233

Compression

Analysis of compression of polymer mushrooms using self-consistent field theory 743(2000)31

Corynebacterium

Detection of mycoloylglycerol by thin-layer chromatography as a tool for the rapid inclusion of corynebacteria of clinical origin in the genus *Corynebacterium* 738(2000)181

Counter-current distribution

Study on heterogeneity of β -glucosidase from *Aspergillus* species by using counter-current distribution 743(2000)247

Coupled columns

Determination of 2-(2-nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione in plasma by direct injection into a coupled column liquid chromatographic system 730(1999)177

Cryogenic oven trapping

Ultra-sensitive method for determination of ethanol in whole blood by headspace capillary gas chromatography with cryogenic oven trapping 727(1999)89

Cyanogen bromide peptide maps

Size-exclusion chromatography performed in capillaries. Studies by liquid chromatography–mass spectrometry 726(1999)37

D

Database retrieval

How to approach substance identification in qualitative bioanalysis 739(2000)125

Denaturing high-performance liquid chromatography

Allelic discrimination by denaturing high-performance liquid chromatography 739(2000)345

Derivatization

Determination of Guan-Fu Base A, a new anti-arrhythmic, in human plasma by gas chromatography and electron–capture detection 740(2000)273

Analysis of 4-hydroxyproline using 4-chloro-7-nitrobenzo-2-oxa-1,3-diazol derivatization and micellar electrokinetic chromatography combined with laser-induced fluorescence detection 744(2000)195

Hydrophylic surfactant–imidazole derivative association studied by RPLC using a hydroorganic solution 749(2000)225

Derivatization, GC

Headspace solid-phase microextraction with 1-pyrenyldiazomethane in-fibre derivatisation for analysis of faecal short-chain fatty acids 730(1999)113

Determination of metabolites of pirimicarb in human urine by gas chromatography–mass spectrometry 730(1999)229

Gas chromatographic–tandem mass spectrometric quantification of human plasma and urinary nitrate after its reduction to nitrite and derivatization to the pentafluorobenzyl derivative 731(1999)285

Gas chromatographic–mass spectrometric analysis of stable isotopes of cysteine and glutathione in biological samples 732(1999)127

Determination of lorazepam in plasma and urine as trimethylsilyl derivative using gas chromatography–tandem mass spectrometry 732(1999)509

Determination of Guan-Fu Base A, a new anti-arrhythmic, in human plasma by gas chromatography and electron–capture detection 740(2000)273

Gas chromatography–mass spectrometry with *tert*-butyldimethylsilyl derivatization: use of the simplified sample preparations and the automated data system to screen for organic acidemias 746(2000)63

Derivatization, HPCE

Direct detection of endogenous histamine in rat peritoneal mast cells by in-capillary derivatization high-performance capillary electrophoresis 736(1999)263

Derivatization, LC

Bioanalysis of apolidine, a new marine antitumoral depsipeptide, in plasma by high-performance liquid chromatography after derivatization with *trans*-4'-hydrazino-2-stilbazole 729(1999)43

Trace analysis of ethosuximide in human plasma with a chemically removable derivatizing reagent and high-performance liquid chromatography 729(1999)111

High-performance liquid chromatographic analysis of physiological amino acids in human brain tumors by pre-column derivatization with phenylisothiocyanate 730(1999)81

Semi-automatic liquid chromatographic analysis of pamidronate in urine after derivatization with 1-naphthylisothiocyanate 730(1999)95

Bioanalysis of *m*-iodobenzylguanidine in plasma by high-performance liquid chromatography after derivatization with benzoin 730(1999)193

Derivatization of thymine and thymine photodimers with 4-bromomethyl-7-methoxycoumarin for fluorescence detection in high-performance liquid chromatography 731(1999)179

On-line derivatization for continuous and automatic monitoring of brain extracellular glutamate levels in anesthetized rats: a microdialysis study 734(1999)1

Improved and simplified liquid chromatographic assay for adefovir, a novel antiviral drug, in human plasma using derivatization with chloroacetaldehyde 736(1999)115

4-Dimethylaminobenzylamine as a sensitive chemiluminescence derivatization reagent for 5-hydroxyindoles and its application to their quantification in human platelet-poor plasma 738(2000)199

Semi-automatic liquid chromatographic analysis of olpadronate in urine and serum using derivatization with (9-fluorenylmethyl)chloroformate 738(2000)331

Novel and sensitive method for the detection of anandamide by the use of its dansyl derivative 740(2000)93

Convenient method of threonine, methionine and their related amino compounds by high-performance liquid chromatography and its application to rumen fluid 741(2000)279

Measurement of free and bound malondialdehyde in plasma by high-performance liquid chromatography as the 2,4-dinitrophenylhydrazine derivative 742(2000)315

Reversed-phase high-performance liquid chromatographic analysis of atenolol enantiomers in rat hepatic microsome after chiral derivatization with 2,3,4,6-tetra-*O*-acetyl- β -D-glycopyranosyl isothiocyanate 742(2000)433

Sensitive determination of ephedrine and norephedrine in human plasma samples using derivatization with 9-fluorenylmethyl chloroformate and liquid chromatography 744(2000)25

High-performance liquid chromatographic assay for catecholamines and metanephrines using fluorimetric detection with pre-column 9-fluorenylmethyloxycarbonyl chloride derivatization 749(2000)179

Detection, electrophoresis

Determination of urine catecholamines by capillary electrophoresis with dual-electrode amperometric detection 750(2001)33

Detection, spectroscopic

Systematic toxicological analysis of drugs and poisons in biosamples by hyphenated chromatographic and spectroscopic techniques 733(1999)47

Dialysis

Automated liquid chromatographic determination of atenolol in plasma using dialysis and trace enrichment on a cation-exchange precolumn for sample handling 739(2000)205

Diffusion

Partitioning and diffusion of large molecules in fibrous structures 743(2000)169

Diffusion coefficients

Diffusion of lysozyme in gels and liquids. A general approach for the determination of diffusion coefficients using holographic laser interferometry 743(2000)151

Direct injection

Determination of 2-(2-nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione in plasma by direct injection into a coupled column liquid chromatographic system 730(1999)177

Direct injection of physiological fluids in micellar liquid chromatography 736(1999)1

Directed ligand evolution

Aptamer affinity chromatography: combinatorial chemistry applied to protein purification 731(1999)275

Displacement chromatography

Polymer displacement/shielding in protein chromatography 741(2000)103

Displacers

Phytic acid as an efficient low-molecular-mass displacer for anion-exchange displacement chromatography of proteins 741(2000)23

Dissociation constant

Determination of the dissociation constant of valine from acetohydroxy acid synthase by equilibrium partition in an aqueous two-phase system 743(2000)225

DNA sequencing

Improvements in DNA sequencing by capillary electrophoresis at elevated temperature using poly(ethylene oxide) as a sieving matrix 745(2000)221

Doping

Systematic toxicological analysis procedures for acidic drugs and/or metabolites relevant to clinical and forensic toxicology and/or doping control 733(1999)3

Gas chromatographic–tandem mass spectrometric determination of anabolic steroids and their esters in hair. Application in doping control and meat quality control 735(1999)189

Drowning-out crystallisation

Drowning-out crystallisation of sodium sulphate using aqueous two-phase systems 743(2000)101

Drug-binding affinities

Development of an immobilized P-glycoprotein stationary phase for on-line liquid chromatographic determination of drug-binding affinities 739(2000)33

Drug discovery

Capillary electrophoresis in the pharmaceutical industry: applications in discovery and chemical development 745(2000)177

Drug screening

Systematic toxicological analysis of drugs and poisons in biosamples by hyphenated chromatographic and spectroscopic techniques 733(1999)47

Testing hair for pharmaceuticals 733(1999)231

Drug screening assays

Chromatographic screening techniques in systematic toxicological analysis 733(1999)27

Dynamic coating

Dynamically-coated capillaries allow for capillary electrophoretic resolution of transferrin sialoforms via direct analysis of human serum 742(2000)79

E**Electrochromatography**

Capillary electrochromatography–electrospray ionization mass spectrometry for the qualitative investigation of the drug etodolac and its metabolites in biological samples 748(2000)213

Electroosmotic flow

External electric field control of electroosmotic flow in non-coated and coated fused-silica capillaries and its application for capillary electrophoretic separations of peptides 741(2000)43

Electrospray ionization

Ultrasensitive determination of NE-100, a novel sigma ligand, in human plasma by liquid chromatography and electrospray ionization tandem mass spectrometry combined with a column-switching technique 730(1999)61

ELISA

Enzyme-linked immunosorbent assay-based selection and optimization of elution buffer for TAG72-affinity chromatography 731(1999)299

Elution buffers

Enzyme-linked immunosorbent assay-based selection and optimization of elution buffer for TAG72-affinity chromatography 731(1999)299

Enantiomer separation

High-performance liquid chromatographic assay of (\pm)-lactic acid and its enantiomers in calf serum 727(1999)23

Stereoselective determination of fenfluramine enantiomers in rat liver microsomal incubates 727(1999)107

High-performance liquid chromatography separation and quantitation of ofloxacin enantiomers in rat microsomes 728(1999)151

Achiral and chiral determination of ciprofibrate and its glucuronide in human urine by capillary electrophoresis 729(1999)33

Direct determination of methamphetamine enantiomers in urine by liquid chromatography with a strong cation-exchange precolumn and phenyl- β -cyclodextrin-bonded semi-microcolumn 729(1999)97

Determination of the D- and L-enantiomers of modafinil in human plasma utilizing liquid–liquid extraction and high-performance liquid chromatography 730(1999)1

Enantiomeric resolution of galanthamine and related drugs used in anti-Alzheimer therapy by means of capillary zone electrophoresis employing derivatized cyclodextrin selectors 730(1999)167

Non-aqueous capillary electrophoresis chiral separations with sulfated β -cyclodextrin 731(1999)187

Modified gas chromatographic–mass spectrometric assay for the determination of gacyclidine enantiomers in human plasma 731(1999)317

Determination of the enantiomers of omeprazole in blood plasma by normal-phase liquid chromatography and detection by atmospheric pressure ionization tandem mass spectrometry 734(1999)191

Quantitative determination of isoflurane enantiomers in blood samples during and after surgery via headspace gas chromatography–mass spectrometry 735(1999)93

Enantioselective analysis of metoprolol in plasma using high-performance liquid chromatographic direct and indirect separations: applications in pharmacokinetics 738(2000)27

Determination of mandelic acid enantiomers in urine by gas chromatography and electron-capture or flame ionisation detection 738(2000)39

Enantioselective analysis of ketone bodies in patients with β -ketothiolase deficiency, medium-chain acyl coenzyme A dehydrogenase deficiency and ketonemic vomiting 739(2000)313

Chromatographic separation of chlorthalidone enantiomers using β -cyclodextrins as chiral additives 740(2000)169

Enantioselective analysis of (*R*)- and (*S*)-atenolol in urine samples by a high-performance liquid chromatography column-switching setup 740(2000)219

Determination of enantiomeric amphetamines as metabolites of illicit amphetamines and selegiline in urine by capillary

electrophoresis using modified β -cyclodextrin 741(2000)221

Sensitive stereospecific determination of acenocoumarol and phenprocoumon in plasma by high-performance liquid chromatography 742(2000)131

Reversed-phase high-performance liquid chromatographic analysis of atenolol enantiomers in rat hepatic microsome after chiral derivatization with 2,3,4,6-tetra-*O*-acetyl- β -D-glycopyranosyl isothiocyanate 742(2000)433

Enantioselective analysis of disopyramide and mono-*N*-dealkyldisopyramide in plasma and urine by high-performance liquid chromatography on an amylose-derived chiral stationary phase 744(2000)299

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VIII. Chiral recognition ability of partially and completely deglycosylated ovoglycoprotein 745(2000)149

Chiral separation by simultaneous use of vancomycin as stationary phase chiral selector and chiral mobile phase additive 745(2000)159

Enantioselective analysis of levetiracetam and its enantiomer *R*- α -ethyl-2-oxo-pyrrolidine acetamide using gas chromatography and ion trap mass spectrometric detection 745(2000)325

Chiral separation of amino acids in biological fluids by micellar electrokinetic chromatography with laser-induced fluorescence detection 745(2000)389

Enantiomeric determination of tramadol and its main metabolite *O*-desmethyltramadol in human plasma by liquid chromatography–tandem mass spectrometry 748(2000)65

Chiral determination of mirtazapine in human blood plasma by high-performance liquid chromatography 748(2000)439

Simultaneous chiral analysis of methamphetamine and related compounds by capillary electrophoresis 749(2000)111

Gas chromatography-electron capture determination of styrene-7,8-oxide enantiomers 749(2000)265

Enantiomeric separation

Validation of a high-performance liquid chromatographic method for the determination of ibuprofen enantiomers in plasma of broiler chickens 749(2000)243

Enzyme purification

Extraction in aqueous two-phase systems of alkaline xylanase produced by *Bacillus pumilus* and its application in kraft pulp bleaching 743(2000)349

Erythrocyte

Capillary gel electrophoresis: separation of major erythrocyte membrane proteins 742(2000)411

Escherichia coli

Preparative two-step purification of recombinant human basic fibroblast growth factor from high-cell-density cultivation of *Escherichia coli* 737(2000)25

Purification of the Ca^{2+} -binding protein S100A1 from myocardium and recombinant *Escherichia coli* 737(2000)39

Expression, purification and biological properties of the carboxyl half part of the HTLV-I surface envelope glycoprotein 737(2000)85

Development of a downstream process for the isolation of *Staphylococcus aureus* arsenate reductase overproduced in *Escherichia coli* 737(2000)167

Purification of recombinant hydantoinease and L-*N*-carbamoylase from *Arthrobacter aureus* expressed in *Escherichia coli*: comparison of wild-type and genetically modified proteins 737(2000)179

Expression of chymotrypsin(ogen) in the thioredoxin reductase deficient mutant strain of *Escherichia coli* AD494(DE3) and purification via a fusion product with a hexahistidine-tail 737(2000)213

Refolding and purification of a urokinase plasminogen activator fragment by chromatography 737(2000)225

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Expanded bed adsorption

Three-step chromatographic purification procedure for the production of a His-tag recombinant kinesin overexpressed in *E. coli* 737(2000)143

Experimental design

Partitioning of xylanolytic complex from *Penicillium janthinellum* by an aqueous two-phase system 743(2000)339

Extensive metabolizers

Enantioselectivity of debrisoquine 4-hydroxylation in Brazilian Caucasian hypertensive patients phenotyped as extensive metabolizers 749(2000)153

External electric field

External electric field control of electroosmotic flow in non-coated and coated fused-silica capillaries and its application for capillary electrophoretic separations of peptides 741(2000)43

F

Filtration

Automated protein precipitation by filtration in the 96-well format 734(1999)257

Fluorescence energy transfer

High-performance liquid chromatography with on-line post-column immunoreaction detection of digoxin and its metabolites based on fluorescence energy transfer in the far-red spectral region 745(2000)305

Fluorocarbon-bonded silica gel column

Determination of concentrations of flecainide in human serum by high-performance liquid chromatography on a fluorocarbon-bonded silica gel column 726(1999)219

Food analysis

Chromatographic and related techniques for the determination of aromatic heterocyclic amines in foods 747(2000)139

Forensic toxicology

Hyphenated liquid chromatographic techniques in forensic toxicology 733(1999)65

Functional genomics

Approaches to functional genomics: potential of matrix-assisted laser desorption ionization–time of flight mass spectrometry combined with separation methods for the analysis of DNA in biological samples 745(2000)231

G

Gamma-radiation

Analysis of gamma radiation-induced damage to plasmid DNA using dynamic size-sieving capillary electrophoresis 741(2000)243

Gene quantification

Direct quantification of AD-36 adenovirus DNA by capillary electrophoresis with laser-induced fluorescence 744(2000)1

Gene therapy vectors

Recovery in aqueous two-phase systems of nanoparticulates applied as surrogate mimics for viral gene therapy vectors 743(2000)409

Genes

Rapid method for relative gene expression determination in human tissues using automated capillary gel electrophoresis and multicolor detection 734(1999)47

Genetic algorithm

Investigation of mathematical methods for efficient optimisation of aqueous two-phase extraction 743(2000)21

Genetic typing

Genetic typing and HIV-1 diagnosis by using 96 capillary array electrophoresis and ultraviolet absorption detection 741(2000)15

Geometrical isomers

Determination of the geometrical diarylpropenamine isomers in feces by high-performance liquid chromatography 728(1999)117

Glass wool

Immobilization of antibodies onto glass wool 738(2000)217

Glomerular filtration rate

Determination of iothalamate in rat urine, plasma, and tubular fluid by capillary electrophoresis 728(1999)143

Glucuronidation

High-performance liquid chromatographic assay to detect hydroxylate and conjugate metabolites of propofol in human urine 742(2000)25

Glycine max

Purification of soybean peroxidase (*Glycine max*) by metal affinity partitioning in aqueous two-phase systems 743(2000)287

O-Glycosylation

Capillary zone electrophoresis and MALDI–mass spectrometry for the monitoring of in vitro *O*-glycosylation of a threonine/serine-rich MUC5AC hexadecapeptide 729(1999)65

Gobio gobio

Two-step purification method of vitellogenin from three teleost fish species: rainbow trout (*Oncorhynchus mykiss*), gudgeon (*Gobio gobio*) and chub (*Leuciscus cephalus*) 737(2000)3

Golgi apparatus

Aqueous two-phase partition applied to the isolation of plasma membranes and Golgi apparatus from cultured mammalian cells 743(2000)377

Grapefruit juice

Amounts and variation in grapefruit juice of the main components causing grapefruit–drug interaction 741(2000)195

H

Hair

Determination of clenbuterol residues in bovine hair by using diphase dialysis and gas chromatography–mass spectrometry 726(1999)133

Chromatographic procedures for determination of cannabinoids in biological samples, with special attention to blood and alternative matrices like hair, saliva, sweat and meconium 733(1999)119

Testing hair for pharmaceuticals 733(1999)231

Gas chromatographic–tandem mass spectrometric determination of anabolic steroids and their esters in hair. Application in doping control and meat quality control 735(1999)189

Hair analysis

Hair analysis for abused and therapeutic drugs 733(1999)161

Halomonas elongata

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Hapten conjugation

High-performance liquid chromatography with on-line post-column immunoreaction detection of digoxin and its metabolites based on fluorescence energy transfer in the far-red spectral region 745(2000)305

Helicobacter pylori

Purification of surface-associated urease from *Helicobacter pylori* 737(2000)202

Holographic laser interferometry

Diffusion of lysozyme in gels and liquids. A general approach for the determination of diffusion coefficients using holographic laser interferometry 743(2000)151

Homocystinuria

Differential diagnosis of homocystinuria by urease treatment, isotope dilution and gas chromatography–mass spectrometry 742(2000)59

HPLC

Simple and rapid determination of irinotecan and its metabolite SN-38 in plasma by high-performance liquid-chromatography: application to clinical pharmacokinetic studies 740(2000)159

On-line reversed-phase liquid chromatography hydride generation emission spectrometry: speciation of arsenic in urine of patients intravenously treated with As_2O_3 740(2000)179

Human hemofiltrate

Composition of the peptide fraction in human blood plasma: database of circulating human peptides 726(1999)25

Human plasma

Simultaneous quantitative determination of cilostazol and its metabolites in human plasma by high-performance liquid chromatography 728(1999)251

Hybridization

Monitoring hybridization during polymerase chain reaction 741(2000)3

Hydride generation

On-line reversed-phase liquid chromatography hydride generation emission spectrometry: speciation of arsenic in urine of patients intravenously treated with As_2O_3 740(2000)179

Hydrophobicity

Affinity chromatography for purification of two urokinases from human urine 742(2000)71

Relative hydrophobicity of organic compounds measured by partitioning in aqueous two-phase systems 743(2000)187

Hydroxyl free radical formation

Determination of hydroxyl free radical formation in human platelets using high-performance liquid chromatography with electrochemical detection 732(1999)213

I

ICP-AES

On-line reversed-phase liquid chromatography hydride generation emission spectrometry: speciation of arsenic in urine of patients intravenously treated with As_2O_3 740(2000)179

Identification

How to approach substance identification in qualitative bioanalysis 739(2000)125

Imaging detection

Effect of salt concentration on separation patterns in static capillary isoelectric focusing with imaging detection 729(1999)355

Immobilized artificial membrane surfaces

Development of predictive retention–activity relationship models of non-steroidal anti-inflammatory drugs by micellar liquid chromatography: comparison with immobilized artificial membrane columns 740(2000)59

Immobilized liposome chromatography

Stimuli-responsive separation of proteins using immobilized liposome chromatography 743(2000)85

Immobilized liposome chromatography for refolding and purification of protein 743(2000)93

Immunoaffinity

Comparison of different elution conditions for the immunopurification of recombinant hepatitis B surface antigen 735(1999)271

Immunoaffinity extraction

Immunoaffinity solid-phase extraction for the trace-analysis of low-molecular-mass analytes in complex sample matrices 745(2000)15

Immunoassays

Capillary electrophoretic immunoassays for digoxin and gentamicin with laser-induced fluorescence polarization detection 734(1999)31

Immuno-extraction

Immuno-affinity solid-phase extraction 745(2000)39

Inclusion bodies

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Injection methods

Determination of amoxicillin in plasma samples by capillary electrophoresis 731(1999)309

Ink

Chromatographic and electrophoretic approaches in ink analysis 733(1999)265

Intercalation dyes

Optimization of intercalation dye concentration for short tandem repeat allele genotyping using capillary electrophoresis with laser-induced fluorescence detection 732(1999)365

Isoelectric precipitation

Isoelectric precipitation of soybean protein using carbon dioxide as a volatile acid 743(2000)357

Isotope ratio

Automated isolation of high-purity plasma albumin for isotope ratio measurements 731(1999)199

Isotypes

Purification of horse immunoglobulin isotypes based on differential elution properties of isotypes from protein A and protein G columns 742(2000)327

J**JP-8 jet fuel**

Method for liquid–liquid extraction of blood surrogates for assessing human exposure to jet fuel 728(1999)193

L**Lacrimal gland**

Purification of rabbit lacrimal gland plasma membranes by aqueous two-phase affinity partitioning 743(2000)397

Laser induced fluorescence detection

On-column derivatization for the analysis of homocysteine and other thiols by capillary electrophoresis with laser-induced fluorescence detection 744(2000)149

LC

Hydrophilic surfactant–imidazole derivative association studied by RPLC using a hydroorganic solution 749(2000)225

LC–MS–MS

Liquid chromatographic–tandem mass spectrometric method for the determination of the neuraminidase inhibitor zanamivir (GG167) in human serum 732(1999)383

Leuciscus cephalus

Two-step purification method of vitellogenin from three teleost fish species: rainbow trout (*Oncorhynchus mykiss*), gudgeon (*Gobio gobio*) and chub (*Leuciscus cephalus*) 737(2000)3

Licorice

Determination of urinary 18 β -glycyrrhetic acid by gas chromatography and its clinical application in man 731(1999)323

Linear free energy relationships

Solvatochromic studies in polyethylene glycol–salt aqueous biphasic systems 743(2000)137

Lipid bilayer membranes

Immobilized liposome chromatography for refolding and purification of protein 743(2000)93

Lipid peroxidation

High-performance liquid chromatographic assay of hydroperoxide levels in oxidatively modified lipoproteins 731(1999)223

Liposomes

Affinity partitioning of biotinylated mixed liposomes: effect of charge on biotin–NeutrAvidin interaction 743(2000)389

Liquid crystals

Application of Pluronic copolymer liquid crystals for the capillary electrophoretic separation of collagen type I cyanogen bromide fragments 739(2000)109

Liquid lids

Liquid lid for biochemical reactions in chip-based nanovials 745(2000)137

Liquid-phase microextraction

Liquid-phase microextraction as a sample preparation technique prior to capillary gas chromatographic-determination of benzodiazepines in biological matrices 749(2000)85

Lithospermum erythrorhizon

Simultaneous analysis of shikimate-derived secondary metabolites in *Lithospermum erythrorhizon* cell suspension cultures by high-performance liquid chromatography 738(2000)3

Lymph

Application of atmospheric pressure chemical ionization liquid chromatography–mass spectrometry in identification of lymph triacylglycerols 748(2000)425

M**Maillard reaction**

Mass spectrometric study on the protein chemical modification of uremic patients in advanced Maillard reaction 731(1999)131

MALDI–TOF mass spectrometry

Approaches to functional genomics: potential of matrix-assisted laser desorption ionization–time of flight mass spectrometry combined with separation methods for the analysis of DNA in biological samples 745(2000)231

Mass spectrometry

Automated in-tube solid-phase microextraction–liquid chromatography–electrospray ionization mass spectrometry for the determination of ranitidine 731(1999)353

Determination of carnitine and acylcarnitines in biological samples by capillary electrophoresis–mass spectrometry 735(1999)171

Determination of raclopride in human plasma by on-column focusing packed capillary liquid chromatography–electrospray ionisation mass spectrometry 748(2000)189

Capillary electrochromatography–electrospray ionization mass spectrometry for the qualitative investigation of the drug etodolac and its metabolites in biological samples 748(2000)213

Capillary electrophoresis–mass spectrometry, liquid chromatography–mass spectrometry and nanoelectrospray-mass spectrometry of praziquantel metabolites 748(2000)221

Match factor

How to approach substance identification in qualitative bioanalysis 739(2000)125

Meconium

Chromatographic procedures for determination of cannabinoids in biological samples, with special attention to blood and alternative matrices like hair, saliva, sweat and meconium 733(1999)119

MEKC

Analysis of 4-hydroxyproline using 4-chloro-7-nitrobenzo-2-oxa-1,3-diazol derivatization and micellar electrokinetic chromatography combined with laser-induced fluorescence detection 744(2000)195

Membrane binding domain

Purification of the membrane binding domain of cytochrome *b*₅ by immobilised nickel chelate chromatography 737(2000)119

Membrane radial flow chromatography

Purification of human prothrombin from Nitschmann fraction III using DEAE membrane radial flow chromatography 742(2000)109

Metabolite

High-performance liquid chromatographic assay to detect hydroxylate and conjugate metabolites of propofol in human urine 742(2000)25

Metal affinity partitioning

Purification of soybean peroxidase (*Glycine max*) by metal affinity partitioning in aqueous two-phase systems 743(2000)287

Metastatic melanoma

Determination of 5-S-cysteinyldopa in plasma and urine using a fully automated solid-phase extraction–high-performance liquid chromatographic method for an improvement of specificity and sensitivity of this prognostic marker of malignant melanoma 727(1999)31

Micellar electrokinetic chromatography

Charged cyclodextrin-mediated sample stacking in micellar capillary electrophoresis. A simple method for enhancing the detection sensitivity of hydrophobic compounds 731(1999)369

Micellar liquid chromatography

Direct injection of physiological fluids in micellar liquid chromatography 736(1999)1

Microbore liquid chromatography

Simultaneous blood and brain sampling of cephalexin in the rat by microdialysis and microbore liquid chromatography: application to pharmacokinetics studies 740(2000)203

Microchip device

Integrated microchip-device for the digestion, separation and postcolumn labeling of proteins and peptides 745(2000)243

Microdialysis

Simultaneous blood and brain sampling of cephalexin in the rat by microdialysis and microbore liquid chromatography: application to pharmacokinetics studies 740(2000)203

High-performance liquid chromatographic analysis of muscular interstitial arginine and norepinephrine kinetics. A microdialysis study in rats 745(2000)279

Microemulsion electrokinetic chromatography

Rapid method for estimating the octanol–water partition coefficient ($\log P_{ow}$) by microemulsion electrokinetic chromatography 745(2000)117

Mimetic ligands

Affinity chromatography of plasma proteins (guanidinobenzoate): use of mimetic matrices and mimetic soluble ligands to prevent the binding of albumin on target affinity matrices 732(1999)165

Mimetic matrices

Affinity chromatography of plasma proteins (guanidinobenzoate): use of mimetic matrices and mimetic soluble ligands to prevent the binding of albumin on target affinity matrices 732(1999)165

Miniaturization

Nanotechnology in bio/clinical analysis 739(2000)139

Molecular imprinting

Separation and sensing based on molecular recognition using molecularly imprinted polymers 728(1999)1

Molecular imprinting: developments and applications in the analytical chemistry field 745(2000)3

Molecular recognition

Separation and sensing based on molecular recognition using molecularly imprinted polymers 728(1999)1

Molecular imprinting for drug bioanalysis. A review on the application of imprinted polymers to solid-phase extraction and binding assay 739(2000)163

Molecular sieve

Technique for a rapid and efficient purification of the SHV-1 and PSE-2 β -lactamases 737(2000)261

Molecular species

Effect of soybean oil and fish oil on individual molecular species of Atlantic salmon head kidney phospholipids determined by normal-phase liquid chromatography coupled to negative ion electrospray tandem mass spectrometry 748(2000)137

Multicolor detection

Rapid method for relative gene expression determination in human tissues using automated capillary gel electrophoresis and multicolor detection 734(1999)47

Multicomponent mixture

Fast liquid chromatographic–mass spectrometric determination of pharmaceutical compounds 732(1999)445

Mutation detection

High-throughput single-strand conformation polymorphism analysis by capillary electrophoresis 741(2000)115

Mutations

Establishment of a combined strategy of genetic and mass spectrometric analyses for characterizing hemoglobin mutations. An example of Hb Hoshida ($\beta 43\text{Glu} \rightarrow \text{Gln}$) 731(1999)125

N

Nanoelectrospray ion source

Capillary electrophoresis–mass spectrometry, liquid chromatography–mass spectrometry and nanoelectrospray-mass spectrometry of praziquantel metabolites 748(2000)221

Nanoparticles

Recovery in aqueous two-phase systems of nanoparticulates applied as surrogate mimics for viral gene therapy vectors 743(2000)409

Nanotechnology

Nanotechnology in bio/clinical analysis 739(2000)139

Natural products

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

Neonatal screening

Urine screening of five-day-old newborns: metabolic profiling of neonatal galactosuria 732(1999)469

Nitschmann fraction III

Purification of human prothrombin from Nitschmann fraction III using DEAE membrane radial flow chromatography 742(2000)109

O

Octanol–water partition coefficient

Rapid method for estimating the octanol–water partition coefficient ($\log P_{ow}$) by microemulsion electrokinetic chromatography 745(2000)117

Oil-in-water

Isolation and purification of serum and interfacial peptides of a trypsinolyzed β -lactoglobulin oil-in-water emulsion 744(2000)389

Oncorhynchus mykiss

Two-step purification method of vitellogenin from three teleost fish species: rainbow trout (*Oncorhynchus mykiss*), gudgeon (*Gobio gobio*) and chub (*Leuciscus cephalus*) 737(2000)3

On-line dialysis

On-line dialysis and quantitative high-performance liquid chromatography analysis of iodixanol in human, rat and monkey plasma 749(2000)135

On-line microdialysis

Measurement of unbound caffeic acid in rat blood by on-line microdialysis coupled with liquid chromatography and its application to pharmacokinetic study 729(1999)119

Determination of unbound cephalothin in rat blood by on-line microdialysis and microbore liquid chromatography 742(2000)125

Optimisation

Investigation of mathematical methods for efficient optimisation of aqueous two-phase extraction 743(2000)21

Organic acidemias

Gas chromatography–mass spectrometry with *tert*.-butyldimethylsilyl derivatization: use of the simplified sample preparations and the automated data system to screen for organic acidemias 746(2000)63

Organic modifiers

Capillary electrophoretic separation of vitamins in sodium dodecyl sulfate containing buffers with lower aliphatic alcohols and *n*-hexane as organic modifiers 741(2000)67

Oxidative stress

Fast quantification of the urinary marker of oxidative stress 8-hydroxy-2'-deoxyguanosine using solid-phase extraction and high-performance liquid chromatography with triple-stage quadrupole mass detection 738(2000)311

P

Packed capillary column

Determination of raclopride in human plasma by on-column focusing packed capillary liquid chromatography–electrospray ionisation mass spectrometry 748(2000)189

Palmaria palmata

One-step purification of R-phycoerythrin from the red macroalga *Palmaria palmata* using preparative polyacrylamide gel electrophoresis 739(2000)117

Particle beam interface

Application of two-dimensional high-performance liquid chromatography–mass spectrometry with particle beam interface 730(1999)141

Partition coefficient

Modelling alterations in the partition coefficient in in vitro biological systems using headspace gas chromatography 738(2000)207

Determination of the dissociation constant of valine from acetohydroxy acid synthase by equilibrium partition in an aqueous two-phase system 743(2000)225

Partitioning of bovine serum albumin in an aqueous two-phase system: optimization of partition coefficient 743(2000)259

Partitioning

Partial purification of penicillin acylase from *Escherichia coli* in poly(ethylene glycol)–sodium citrate aqueous two-phase systems 734(1999)15

Polymer recycling in aqueous two-phase extractions using thermoseparating ethylene oxide–propylene oxide copolymers 743(2000)115

Partitioning and diffusion of large molecules in fibrous structures 743(2000)169

Relative hydrophobicity of organic compounds measured by partitioning in aqueous two-phase systems 743(2000)187

Partitioning of whey proteins, bovine serum albumin and porcine insulin in aqueous two-phase systems 743(2000)235

Bioaffinity extraction of glucoamylase in aqueous two-phase systems using starch as free bioligand 743(2000)241

Study on heterogeneity of β -glucosidase from *Aspergillus* species by using counter-current distribution 743(2000)247

Bovine serum albumin partitioning in an aqueous two-phase system: Effect of pH and sodium chloride concentration 743(2000)255

Application of aqueous two-phase partition to the production of homogeneous preparations of fluorescently labelled human serum albumin 743(2000)263

Peptides partitioning in an aqueous dextran–polyethylene glycol two-phase system 743(2000)271

Purification of acetohydroxy acid synthase by separation in an aqueous two-phase system 743(2000)281

Partitioning of peptides and recombinant protein–peptide fusions in thermoseparating aqueous two-phase systems: effect of peptide primary structure 743(2000)295

Partitioning behavior of amino acids in aqueous two-phase systems with recyclable volatile salts 743(2000)317

Phase behavior and protein partitioning in aqueous two-phase systems of cationic–anionic surfactant mixtures 743(2000)327

Partitioning of xylanolytic complex from *Penicillium janthinellum* by an aqueous two-phase system 743(2000)339

Applications of aqueous two-phase partition to isolation of membranes from plants: A periodic NADH oxidase activity as a marker for right side-out plasma membrane vesicles 743(2000)369

Affinity partitioning of biotinylated mixed liposomes: effect of charge on biotin–NeutrAvidin interaction 743(2000)389

Simultaneous accumulation of low-molecular-mass RNA at the interface along with accumulation of high-molecular-mass RNA on aqueous two-phase system partitioning 743(2000)421

Pattern recognition

Reversed-phase high-performance liquid chromatographic investigation of urinary normal and modified nucleosides of cancer patients 732(1999)307

Peak selection

Mass directed peak selection, an efficient method of drug metabolite identification using directly coupled liquid chromatography–mass spectrometry–nuclear magnetic resonance spectroscopy 748(2000)281

Penicillium janthinellum

Partitioning of xylanolytic complex from *Penicillium janthinellum* by an aqueous two-phase system 743(2000)339

Peptide library

Use of liquid chromatography–mass spectrometry coupling for monitoring the serralsin-catalyzed hydrolysis of a peptide library 732(1999)271

Peptide mapping

Characterization of a solid-phase extraction device for discontinuous on-line preconcentration in capillary electrophoresis-based peptide mapping 736(1999)273

Two-step fast protein liquid chromatographic purification of the *Serratia marcescens* hemolysin and peptide mapping with mass spectrometry 737(2000)13

Peripheral benzodiazepine receptor

High-performance liquid chromatographic determination of [14 C]1-(2-chlorophenyl)-*N*-methyl-*N*-(1-methylpropyl)-3-isoquinoline carboxamide in mouse plasma and tissue and in human plasma 736(1999)61

Peroxyoxalate chemiluminescence detection

High-performance liquid chromatography with peroxyoxalate chemiluminescence detection of bisphenol A migrated from polycarbonate baby bottles using 4-(4,5-diphenyl-1*H*-imidazol-2-yl)benzoyl chloride as a label 749(2000)49

pH

Bovine serum albumin partitioning in an aqueous two-phase system: Effect of pH and sodium chloride concentration 743(2000)255

pH gradient

Hydrophobic character of surface regions and total hydrophobicity of four variants of chromosomal class C β -lactamase from *Pseudomonas aeruginosa* are identical. Chromatographic comparison of the hydrophobic character of the variants and the effect of focusing buffer composition on the separation of the variants by chromatofocusing with internal and external pH gradients 746(2000)161

Pharmaceutical quality control

System performance and variability of chromatographic techniques used in pharmaceutical quality control 745(2000)167

Pharmacokinetics

Determination of zolpidem in serum microsamples by high-performance liquid chromatography and its application to pharmacokinetics in rats 734(1999)299

Determination of tramadol by capillary gas chromatography with flame ionization detection. Application to human and rabbit pharmacokinetic studies 736(1999)89

Determination of unbound cefmetazole in rat blood by on-line microdialysis and microbore liquid chromatography: a pharmacokinetic study 736(1999)129

Enantioselective high-performance liquid chromatographic analysis of the 5-HT_{1A} receptor agonist 8-hydroxy-2-(di-*n*-propylamino)tetralin. Application to a pharmacokinetic–pharmacodynamic study in rats 738(2000)67

Simple and rapid determination of irinotecan and its metabolite SN-38 in plasma by high-performance liquid-chromatography: application to clinical pharmacokinetic studies 740(2000)159

Simultaneous blood and brain sampling of cephalexin in the rat by microdialysis and microbore liquid chromatography: application to pharmacokinetics studies 740(2000)203

Phase separation kinetics

Kinetics of phase separation under different process and design parameters in aqueous two-phase systems 743(2000)195

Physiological samples

Direct injection of physiological fluids in micellar liquid chromatography 736(1999)1

Plant materials

Analysis of amines in plant materials 747(2000)49

Plasma

Determination of 5-*S*-cysteinyldopa in plasma and urine using a fully automated solid-phase extraction–high-performance liquid chromatographic method for an improvement of specificity and sensitivity of this prognostic marker of malignant melanoma 727(1999)31

Quantification of topotecan and its metabolite *N*-desmethyltopotecan in human plasma, urine and faeces by high-performance liquid chromatographic methods 727(1999)191

Plasma membranes

Aqueous two-phase partition applied to the isolation of plasma membranes and Golgi apparatus from cultured mammalian cells 743(2000)377

Plasma protein binding

Simultaneous determination of enantioselective plasma protein binding of aminohydantoins by ultrafiltration and chiral high-performance liquid chromatography 732(1999)31

Plasma sample purification

Comparison of plasma sample purification by manual liquid–liquid extraction, automated 96-well liquid–liquid extraction and automated 96-well solid-phase extraction for analysis by high-performance liquid chromatography with tandem mass spectrometry 732(1999)501

Plasmids

Analysis of gamma radiation-induced damage to plasmid DNA using dynamic size-sieving capillary electrophoresis 741(2000)243

Pluronic modifier

Capillary electrophoretic separation of proteins and peptides using Pluronic liquid crystals and surface-modified capillaries 741(2000)37

Polyamide hollow-fibre

Human serum albumin chromatography by Cibacron Blue F3GA-derived microporous polyamide hollow-fiber affinity membranes 746(2000)123

Polycarbonate accumulation wall

Field- and flow-dependent trapping of red blood cells on polycarbonate accumulation wall in sedimentation field-flow fractionation 734(1999)91

Polycarbonate bottle

High-performance liquid chromatography with peroxyoxalate chemiluminescence detection of bisphenol A migrated from polycarbonate baby bottles using 4-(4,5-diphenyl-1*H*-imidazol-2-yl)benzoyl chloride as a label 749(2000)49

Polyionic sequence

Purification of a viral coat protein by an engineered polyionic sequence 737(2000)77

Polymer recycling

Polymer recycling in aqueous two-phase extractions using thermoseparating ethylene oxide–propylene oxide copolymers 743(2000)115

Polymerase chain reaction

Rapid method for relative gene expression determination in human tissues using automated capillary gel electrophoresis and multicolor detection 734(1999)47

Allelic discrimination by denaturing high-performance liquid chromatography 739(2000)345

Monitoring hybridization during polymerase chain reaction 741(2000)3

High-throughput single-strand conformation polymorphism analysis by capillary electrophoresis 741(2000)115

Polymorphonuclear leucocytes

Quantitative determination of free intracellular amino acids in single human polymorphonuclear leucocytes. Recent developments in sample preparation and high-performance liquid chromatography 728(1999)157

Preconcentration

Characterization of a solid-phase extraction device for discontinuous on-line preconcentration in capillary electrophoresis-based peptide mapping 736(1999)273

Pre-concentration

Molecular imprinting for drug bioanalysis. A review on the application of imprinted polymers to solid-phase extraction and binding assay 739(2000)163

Programmed temperature vaporiser

Evaluation of the programmed temperature vaporiser for large-volume injection of biological samples in gas chromatography 729(1999)199

Propofol

High-performance liquid chromatographic assay to detect hydroxylate and conjugate metabolites of propofol in human urine 742(2000)25

Protease inhibitors

Simultaneous determination of the HIV protease inhibitors indinavir, amprenavir, saquinavir, ritonavir, nelfinavir and the non-nucleoside reverse transcriptase inhibitor efavirenz by high-performance liquid chromatography after solid-phase extraction 740(2000)43

Protein binding

Determination of lidocaine in plasma by direct solid-phase microextraction combined with gas chromatography 739(2000)175

Protein-binding patterns

Protein-binding patterns of the antitumor antibiotic cryptophycin 52 as measured with a two-phase partitioning system 735(1999)121

Protein fingerprinting

Reversed-phase high-performance liquid chromatographic assay for the adenovirus type 5 proteome 732(1999)411

Protein folding

Model process for separation based on unfolding and refolding of chymotrypsin inhibitor 2 in thermoseparating polymer two-phase systems 743(2000)13

Protein refolding using stimuli-responsive polymer-modified aqueous two-phase systems 743(2000)215

Protein partitioning

Extraction in aqueous two-phase systems of alkaline xylanase produced by *Bacillus pumilus* and its application in kraft pulp bleaching 743(2000)349

Protein–peptide fusions

Partitioning of peptides and recombinant protein–peptide fusions in thermoseparating aqueous two-phase systems: effect of peptide primary structure 743(2000)295

Protein refolding

Immobilized liposome chromatography for refolding and purification of protein 743(2000)93

Protein removal

Free malondialdehyde determination in rat brain tissue by capillary zone electrophoresis: evaluation of two protein removal procedures 740(2000)87

Proteomics

Strategy for qualitative and quantitative analysis in proteomics based on signature peptides 745(2000)197

Pseudo-affinity

Preparation, evaluation and application of new pseudo-affinity chromatographic supports for penicillin acylase purification 739(2000)63

Purification

Two-step purification method of vitellogenin from three teleost fish species: rainbow trout (*Oncorhynchus mykiss*), gudgeon (*Gobio gobio*) and chub (*Leuciscus cephalus*) 737(2000)3

Two-step fast protein liquid chromatographic purification of the *Serratia marcescens* hemolysin and peptide mapping with mass spectrometry 737(2000)13

Preparative two-step purification of recombinant human basic fibroblast growth factor from high-cell-density cultivation of *Escherichia coli* 737(2000)25

Purification of the Ca²⁺-binding protein S100A1 from myocardium and recombinant *Escherichia coli* 737(2000)39

Multi-step purification strategy for RANTES wild-type and mutated analogues expressed in a baculovirus system 737(2000)47

Purification of the c-erbB2/*neu* membrane-spanning segment: a hydrophobic challenge 737(2000)55

Chromatographic purification of the C_H2 domain of the monoclonal antibody MAK33 737(2000)63

Purification of a viral coat protein by an engineered polyionic sequence 737(2000)77

Purification of respiratory syncytial virus F and G proteins 737(2000)97

Three-step purification of bacterially expressed human single-chain Fv antibodies for clinical applications 737(2000)107

Purification of the membrane binding domain of cytochrome *b*₅ by immobilised nickel chelate chromatography 737(2000)119

Novel, rapid purification of the membrane protein photosystem I by high-performance liquid chromatography on porous materials 737(2000)131

Three-step chromatographic purification procedure for the production of a His-tag recombinant kinesin overexpressed in *E. coli* 737(2000)143

Improved purification of the membrane-bound hydrogenase–sulfur-reductase complex from thermophilic archaea using ϵ -aminocaproic acid-containing chromatography buffers 737(2000)151

Highly efficient purification of porcine diamine oxidase 737(2000)161

Development of a downstream process for the isolation of *Staphylococcus aureus* arsenate reductase overproduced in *Escherichia coli* 737(2000)167

Purification of recombinant hydantoinase and L-N-carbamoylase from *Arthrobacter aurescens* expressed in *Escherichia coli*: comparison of wild-type and genetically modified proteins 737(2000)179

Purification of a D-hydantoinase using a laboratory-scale Streamline phenyl column as the initial step 737(2000)187

Purification of surface-associated urease from *Helicobacter pylori* 737(2000)202

Expression of chymotrypsin(ogen) in the thioredoxin reductase deficient mutant strain of *Escherichia coli* AD494(DE3) and purification via a fusion product with a hexahistidine-tail 737(2000)213

Refolding and purification of a urokinase plasminogen activator fragment by chromatography 737(2000)225

Purification of human adenosine deaminase for the preparation of a reference material 737(2000)237

Pullulanase from the hyperthermophilic bacterium *Thermotoga maritima*: purification by β -cyclodextrin affinity chromatography 737(2000)245

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Technique for a rapid and efficient purification of the SHV-1 and PSE-2 β -lactamases 737(2000)261

Purification of the fengycin synthetase multienzyme system from *Bacillus subtilis* b213 737(2000)267

New approach for separating *Bacillus subtilis* metalloprotease and α -amylase by affinity chromatography and for purifying neutral protease by hydrophobic chromatography 737(2000)277

Purification and characterization of an aminopeptidase from the chloroplast stroma of barley leaves by chromatographic and electrophoretic methods 737(2000)285

Three-step chromatographic purification of Cpr6, a cyclophilin from *Saccharomyces cerevisiae* 737(2000)295

Pyrococcus woesei

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Q

Qualitative analysis

How to approach substance identification in qualitative bioanalysis 739(2000)125

Quantitative retention–activity relationship

Development of predictive retention–activity relationship models of non-steroidal anti-inflammatory drugs by micellar liquid chromatography: comparison with immobilized artificial membrane columns 740(2000)59

Quantitative retention–structure relationship

Development of predictive retention–activity relationship models of non-steroidal anti-inflammatory drugs by micellar liquid chromatography: comparison with immobilized artificial membrane columns 740(2000)59

R

Radiochemistry

Radiochemical high-performance liquid chromatographic assay for the determination of catechol *O*-methyltransferase activity towards various substrates 736(1999)143

RANTES mutagenesis

Multi-step purification strategy for RANTES wild-type and mutated analogues expressed in a baculovirus system 737(2000)47

Rat

Simultaneous blood and brain sampling of cephalexin in the rat by microdialysis and microbore liquid chromatography: application to pharmacokinetics studies 740(2000)203

Refolding

Refolding and purification of a urokinase plasminogen activator fragment by chromatography 737(2000)225

Retro-Diels–Alder fragmentation

Identification of the principal circulating metabolite of a synthetic 5,4'-diaminoflavone (NSC 686288), an antitumor agent, in the rat 741(2000)205

Reviews

Separation and sensing based on molecular recognition using molecularly imprinted polymers (Review) 728(1999)1

Mass spectrometric analysis of neuropeptidergic systems in the human pituitary and cerebrospinal fluid (Review) 731(1999)3

3-Deoxyglucosone: metabolism, analysis, biological activity, and clinical implication (Review) 731(1999)23

Systematic toxicological analysis procedures for acidic drugs and/or metabolites relevant to clinical and forensic toxicology and/or doping control (Review) 733(1999)3

Chromatographic screening techniques in systematic toxicological analysis (Review) 733(1999)27

Systematic toxicological analysis of drugs and poisons in biosamples by hyphenated chromatographic and spectroscopic techniques (Review) 733(1999)47

Hyphenated liquid chromatographic techniques in forensic toxicology (Review) 733(1999)65

Liquid chromatography–mass spectrometry: potential in forensic and clinical toxicology (Review) 733(1999)93

Chromatographic procedures for determination of cannabinoids in biological samples, with special attention to blood and alternative matrices like hair, saliva, sweat and meconium (Review) 733(1999)119

Comparative analysis of heroin and cocaine seizures (Review) 733(1999)127

Determination of “Ecstasy” components in alternative biological specimens (Review) 733(1999)137

Recent advances in chromatographic and mass spectrometric methods for determination of LSD and its metabolites in physiological specimens (Review) 733(1999)145

Hair analysis for abused and therapeutic drugs (Review) 733(1999)161

Poisoning by plant material: review of human cases and analytical determination of main toxins by high-performance liquid chromatography–(tandem) mass spectrometry (Review) 733(1999)181

Testing hair for pharmaceuticals (Review) 733(1999)231

Sweat testing for cocaine, codeine and metabolites by gas chromatography–mass spectrometry (Review) 733(1999)247

Chromatographic and electrophoretic approaches in ink analysis (Review) 733(1999)265

Direct injection of physiological fluids in micellar liquid chromatography (Review) 736(1999)1

Advanced separation methods for collagen parent α -chains, their polymers and fragments (Review) 739(2000)3

High-performance affinity chromatography and immobilized serum albumin as probes for drug- and hormone-protein binding (Review) 739(2000)39

Molecular imprinting for drug bioanalysis. A review on the application of imprinted polymers to solid-phase extraction and binding assay (Review) 739(2000)163

Monitoring hybridization during polymerase chain reaction (Review) 741(2000)3

Polymer displacement/shielding in protein chromatography (Review) 741(2000)103

High-throughput single-strand conformation polymorphism analysis by capillary electrophoresis (Review) 741(2000)115

Molecular imprinting: developments and applications in the analytical chemistry field (Review) 745(2000)3

Immunoaffinity solid-phase extraction for the trace-analysis of low-molecular-mass analytes in complex sample matrices (Review) 745(2000)15

Solid-phase microextraction for the analysis of biological samples (Review) 745(2000)49

Cyclodextrins: a versatile tool in separation science (Review) 745(2000)83

Analysis of biogenic amines in microdialysates of the brain (Review) 747(2000)21

Clinical chemistry of serotonin and metabolites (Review) 747(2000)33

Analysis of amines in plant materials (Review) 747(2000)49

Current methodologies for the analysis of aminoglycosides (Review) 747(2000)69

Separation and assay methods for melatonin and its precursors (Review) 747(2000)95

Separation methods used in the determination of choline and acetylcholine (Review) 747(2000)111

Chromatographic and related techniques for the determination of aromatic heterocyclic amines in foods (Review) 747(2000)139

Separation of heteroaromatic amines in food products (Review) 747(2000)171

Chromatographic separation of 2,3-benzodiazepines (Review) 747(2000)203

Determination of uremic toxins in biofluids: creatinine, creatine, uric acid and xanthines (Review) 747(2000)217

Chromatographic and capillary electrophoretic methods for the analysis of nicotinic acid and its metabolites (Review) 747(2000)229

Separation methods for amino group-possessing pesticides in biological samples (Review) 747(2000)241

Liquid-phase microextraction as a sample preparation technique prior to capillary gas chromatographic-determination of benzodiazepines in biological matrices 749(2000)85

Sample pretreatment

Solid-phase microextraction for the analysis of biological samples 745(2000)49

Scophthalmus maximus

Determination of oxolinic acid in faeces and urine of turbot (*Scophthalmus maximus*) by high-performance liquid chromatography using fluorescence detection 749(2000)127

Sedimentation field-flow fractionation

Field- and flow-dependent trapping of red blood cells on polycarbonate accumulation wall in sedimentation field-flow fractionation 734(1999)91

Self-consistent field theory

Analysis of compression of polymer mushrooms using self-consistent field theory 743(2000)31

Analysis of brush-particle interactions using self-consistent-field theory 743(2000)41

Serratia marcescens

Two-step fast protein liquid chromatographic purification of the *Serratia marcescens* hemolysin and peptide mapping with mass spectrometry 737(2000)13

Short tandem repeat alleles

Optimization of intercalation dye concentration for short tandem repeat allele genotyping using capillary electrophoresis with laser-induced fluorescence detection 732(1999)365

Sigma ligand

Ultrasensitive determination of NE-100, a novel sigma ligand, in human plasma by liquid chromatography and electrospray ionization tandem mass spectrometry combined with a column-switching technique 730(1999)61

Similarity index

How to approach substance identification in qualitative bioanalysis 739(2000)125

Simplex

Investigation of mathematical methods for efficient optimisation of aqueous two-phase extraction 743(2000)21

Single-cell analysis

Single-cell analysis avoids sample processing bias 741(2000)31

Single-strand conformation polymorphism

High-throughput single-strand conformation polymorphism analysis by capillary electrophoresis 741(2000)115

Solid-phase extraction

Determination of amoxicillin in plasma samples by capillary electrophoresis 731(1999)309

S

Saccharomyces cerevisiae

Three-step chromatographic purification of Cpr6, a cyclophilin from *Saccharomyces cerevisiae* 737(2000)295

Saliva

Chromatographic procedures for determination of cannabinoids in biological samples, with special attention to blood and alternative matrices like hair, saliva, sweat and meconium 733(1999)119

Salt concentration

Effect of salt concentration on separation patterns in static capillary isoelectric focusing with imaging detection 729(1999)355

Sample preparation

Novel sample preparation method facilitating identification of urinary drug metabolites by liquid chromatography–tandem mass spectrometry 738(2000)259

Automated, fast and sensitive quantification of drugs in blood by liquid chromatography–mass spectrometry with on-line extraction: immunosuppressants 748(2000)41

Liquid chromatographic–tandem mass spectrometric method for the determination of the neuraminidase inhibitor zanamivir (GG167) in human serum 732(1999)383

Solid-phase microextraction

Fishing for a drug: solid-phase microextraction for the assay of clozapine in human plasma 731(1999)231

Automated in-tube solid-phase microextraction–liquid chromatography–electrospray ionization mass spectrometry for the determination of ranitidine 731(1999)353

Determination of lidocaine in plasma by direct solid-phase microextraction combined with gas chromatography 739(2000)175

Solid-phase microextraction for the analysis of biological samples 745(2000)49

High-temperature solid-phase microextraction procedure for the detection of drugs by gas chromatography–mass spectrometry 745(2000)399

Solloids

Surfactant aggregates (solloids) adsorbed on silica as stationary chromatographic phases: structures and properties 743(2000)65

Solubilization

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Solvation descriptors

Connection between chromatographic data and biological data 745(2000)103

Solvatochromic studies

Solvatochromic studies in polyethylene glycol–salt aqueous biphasic systems 743(2000)137

Stable isotope dilution

Combined method for the determination of γ -aminobutyric and β -alanine in cerebrospinal fluid by stable isotope dilution mass spectrometry 732(1999)245

Stable isotopic enrichment

Determination of stable isotopic enrichment and concentration of glycerol in plasma via gas chromatography–mass spectrometry for the estimation of lipolysis in vivo 744(2000)47

Stacking

Charged cyclodextrin-mediated sample stacking in micellar capillary electrophoresis. A simple method for enhancing the detection sensitivity of hydrophobic compounds 731(1999)369

Staphylococcus aureus

Development of a downstream process for the isolation of *Staphylococcus aureus* arsenate reductase overproduced in *Escherichia coli* 737(2000)167

Stationary phases, LC

Development of an immobilized P-glycoprotein stationary phase for on-line liquid chromatographic determination of drug-binding affinities 739(2000)33

Performance of a pentafluorophenylpropyl stationary phase for the electrospray ionization high-performance liquid chromatography–mass spectrometry–mass spectrometry assay of cocaine and its metabolite ecgonine methyl ester in human urine 748(2000)77

Stereoselectivity

Stereoselective quantification of methadone and its major oxidative metabolite, 2-ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine, in human urine using high-performance liquid chromatography 744(2000)165

Steroids

Screening urine for exogenous testosterone by isotope ratio mass spectrometric analysis of one pregnanediol and two androstane diols 727(1999)95

Stimuli-responsive separation

Stimuli-responsive separation of proteins using immobilized liposome chromatography 743(2000)85

Protein refolding using stimuli-responsive polymer-modified aqueous two-phase systems 743(2000)215

Structural elucidation

Application of directly coupled LC–NMR–MS to the structural elucidation of metabolites of the HIV-1 reverse-transcriptase inhibitor BW935U83 748(2000)269

Sulfation

High-performance liquid chromatographic assay to detect hydroxylate and conjugate metabolites of propofol in human urine 742(2000)25

Surfactant aggregates

Surfactant aggregates (solloids) adsorbed on silica as stationary chromatographic phases: structures and properties 743(2000)65

Sweat

Chromatographic procedures for determination of cannabinoids in biological samples, with special attention to blood and alternative matrices like hair, saliva, sweat and meconium 733(1999)119

Sweat testing for cocaine, codeine and metabolites by gas chromatography–mass spectrometry 733(1999)247

Synechocystis

Novel, rapid purification of the membrane protein photosystem I by high-performance liquid chromatography on porous materials 737(2000)131

System performance

System performance and variability of chromatographic techniques used in pharmaceutical quality control 745(2000)167

Systematic toxicological analysis

Chromatographic screening techniques in systematic toxicological analysis 733(1999)47

Systematic toxicological analysis of drugs and poisons in biosamples by hyphenated chromatographic and spectroscopic techniques 733(1999)47

T

Tandem mass spectrometry

Gas chromatographic–tandem mass spectrometric quantification of human plasma and urinary nitrate after its reduction to nitrite and derivatization to the pentafluorobenzyl derivative 731(1999)285

Novel sample preparation method facilitating identification of urinary drug metabolites by liquid chromatography–tandem mass spectrometry 738(2000)259

Simultaneous determination of ivabradine and its metabolites in human plasma by liquid chromatography–tandem mass spectrometry 745(2000)261

Temperature effects

Effect of temperature on the separation of DNA fragments by high-performance liquid chromatography and capillary electrophoresis: a comparative study 738(2000)243

Template polymerization

Separation and sensing based on molecular recognition using molecularly imprinted polymers 728(1999)1

Testosterone

Screening urine for exogenous testosterone by isotope ratio mass spectrometric analysis of one pregnanediol and two androstane diols 727(1999)95

Thermotoga maritima

Pullulanase from the hyperthermophilic bacterium *Thermotoga maritima*: purification by β -cyclodextrin affinity chromatography 737(2000)245

Toxic plants

Poisoning by plant material: review of human cases and analytical determination of main toxins by high-performance liquid chromatography–(tandem) mass spectrometry 733(1999)181

Toxicological analysis

Systematic toxicological analysis procedures for acidic drugs and/or metabolites relevant to clinical and forensic toxicology and/or doping control 733(1999)3

Trace enrichment

Automated liquid chromatographic determination of atenolol in plasma using dialysis and trace enrichment on a cation-exchange precolumn for sample handling 739(2000)205

Trace-analysis

Immunoaffinity solid-phase extraction for the trace-analysis of low-molecular-mass analytes in complex sample matrices 745(2000)15

Trapping of red blood cells

Field- and flow-dependent trapping of red blood cells on polycarbonate accumulation wall in sedimentation field-flow fractionation 734(1999)91

Turbot

Determination of oxolinic acid in faeces and urine of turbot (*Scophthalmus maximus*) by high-performance liquid chromatography using fluorescence detection 749(2000)127

Two-dimensional separation

Complete two-dimensional separation for analysis of acidic compounds in plasma using column-switching reversed-phase liquid chromatography 726(1999)225

Two-phase partitioning

Protein-binding patterns of the antitumor antibiotic cryptophycin 52 as measured with a two-phase partitioning system 735(1999)121

U

Urine

Determination of 5-S-cysteinyldopa in plasma and urine using a fully automated solid-phase extraction–high-performance liquid chromatographic method for an improvement of specificity and sensitivity of this prognostic marker of malignant melanoma 727(1999)31

Quantification of topotecan and its metabolite *N*-desmethyltopotecan in human plasma, urine and faeces by high-performance liquid chromatographic methods 727(1999)191

Determination of clenbuterol in bovine urine using gas chromatography–mass spectrometry following clean-up on an ion-exchange resin 728(1999)67

On-line reversed-phase liquid chromatography hydride generation emission spectrometry: speciation of arsenic in urine of patients intravenously treated with As₂O₃ 740(2000)179

V

Validation

Validated electrospray liquid chromatographic–mass spectrometric assay for the determination of the mushroom toxins α - and β -amanitin in urine after immunoaffinity extraction 748(2000)125

Viscosity prediction

Viscosity prediction of polyethylene glycol–dextran–water solutions used in aqueous two-phase systems 743(2000)181

Vitreous humour

Capillary zone electrophoresis of potassium in human vitreous humour: validation of a new method 733(1999)273

W

96-Well liquid–liquid extraction

Quantitative liquid chromatographic–tandem mass spectrometric determination of reserpine in FVB/N mouse plasma using a “chelating” agent (disodium EDTA) for releasing protein-bound analytes during 96-well liquid–liquid extraction 742(2000)369

Y

Yeast

Identification of yeast species by fatty acid profiling as measured by gas-liquid chromatography 742(2000)13