

CUMULATIVE COMPOUND INDEX

J. CHROMATOGR. B VOLS. 726–750

For all compound groups: see also sub-groups and individual compounds
For all compound sub-groups: see also main groups and individual compounds
For all individual compounds: see also compound groups and sub-groups

A

Aaptamine

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

Abacavir

Quantitative determination of abacavir (1592U89), a novel nucleoside reverse transcriptase inhibitor, in human plasma using isocratic reversed-phase high-performance liquid chromatography with ultraviolet detection 736(1999)123

Liquid chromatographic assay for simultaneous determination of abacavir and mycophenolic acid in human plasma using dual spectrophotometric detection 750(2001)155

Acenocoumarol

Sensitive stereospecific determination of acenocoumarol and phenprocoumon in plasma by high-performance liquid chromatography 742(2000)131

Acetaminophen

Simple and rapid assay for acetaminophen and conjugated metabolites in low-volume serum samples 732(1999)323

Acetohydroxy acid synthase

Determination of the dissociation constant of valine from acetohydroxy acid synthase by equilibrium partition in an aqueous two-phase system 743(2000)225

Purification of acetohydroxy acid synthase by separation in an aqueous two-phase system 743(2000)281

Acetylcholine

Effect of 2-deoxy-D-glucose on acetylcholine and histamine levels in gastric juice of pylorus-ligated rats anesthetized with urethane 742(2000)295

Separation methods used in the determination of choline and acetylcholine 747(2000)111

N-Acetyl-S-(N-methylcarbamoyl)cysteine

Simultaneous determination of two human urinary metabolites of *N,N*-dimethylformamide using gas chromatography–thermionic sensitive detection with mass spectrometric confirmation 734(1999)285

N-Acetylneuraminic acid

Importance of high-performance liquid chromatographic analysis of serum *N*-acetylneuraminic acids in evaluating surgical treatment in patients with early endometrial cancer 732(1999)375

Acidic compounds

Complete two-dimensional separation for analysis of acidic compounds in plasma using column-switching reversed-phase liquid chromatography 726(1999)225

Acidic drugs

Systematic toxicological analysis procedures for acidic drugs and/or metabolites relevant to clinical and forensic toxicology and/or doping control 733(1999)3

Acrolein

Quantitative high-performance liquid chromatographic determination of acrolein in plasma after derivatization with Luminarin® 3 739(2000)239

Acrosine-like protease

Inhibition of acrosine-like protease activity by a lectin affinity chromatographic bovine seminal plasma fraction containing the PDC-109 and aSFP proteins 746(2000)141

Acyclovir

Rapid determination of valaciclovir and acyclovir in human biological fluids by high-performance liquid chromatography using isocratic elution 732(1999)47

Rapid, simple and sensitive high-performance liquid chromatographic method for detection and determination of acyclovir in human plasma and its use in bioavailability studies 739(2000)231

Acylcarnitine

Effect of sports activity on carnitine metabolism. Measurement of free carnitine, γ -butyrobetaine and acylcarnitines by tandem mass spectrometry 731(1999)83

Modifications in electrospray tandem mass spectrometry for a neonatal-screening pilot study in Japan 731(1999)97

Acylcarnitines

Determination of carnitine and acylcarnitines in biological samples by capillary electrophoresis–mass spectrometry 735(1999)171

Acyl coenzyme A:cholesterol acyltransferase

Highly sensitive determination of TS-962 (HL-004), a novel acyl-CoA:cholesterol acyltransferase inhibitor, in rat and rabbit plasma by liquid chromatography and atmospheric pressure chemical ionization-tandem mass spectrometry combined with a column-switching technique 750(2001)99

Acyl glucuronides

Gemfibrozil and its oxidative metabolites: quantification of aglycones, acyl glucuronides, and covalent adducts in samples from preclinical and clinical kinetic studies 741(2000)129

Acylglycine

Screening for fatty acid beta oxidation disorders. Acylglycine analysis by electron impact ionization gas chromatography–mass spectrometry 731(1999)105

AD-36 Adenovirus

Direct quantification of AD-36 adenovirus DNA by capillary electrophoresis with laser-induced fluorescence 744(2000)1

Adefovir

Improved and simplified liquid chromatographic assay for adefovir, a novel antiviral drug, in human plasma using derivatization with chloroacetaldehyde 736(1999)115

Adenosine

Erratum to “Determination of adenosine and deoxyadenosine in urine by high-performance liquid chromatography with column switching”. [J. Chromatogr. B, 719 (1998) 55–61] 726(1999)317

Erratum to “Simultaneous determination of adenosine, S-adenosylhomocysteine and S-adenosylmethionine in biological samples using solid-phase extraction and high-performance liquid chromatography [J. Chromatogr. B, 724 (1999) 231–238] 729(1999)379

Adenosine deaminase

Erratum to “Determination of adenosine and deoxyadenosine in urine by high-performance liquid chromatography with column switching”. [J. Chromatogr. B, 719 (1998) 55–61] 726(1999)317

Purification of human adenosine deaminase for the preparation of a reference material 737(2000)237

Adenosine triphosphate

Improved high-performance liquid chromatographic assay for the determination of “high-energy” phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

S-Adenosylhomocysteine

Erratum to “Simultaneous determination of adenosine, S-adenosylhomocysteine and S-adenosylmethionine in biological samples using solid-phase extraction and high-performance liquid chromatography [J. Chromatogr. B, 724 (1999) 231–238] 729(1999)379

S-Adenosylmethionine

Erratum to “Simultaneous determination of adenosine, S-adenosylhomocysteine and S-adenosylmethionine in biological samples using solid-phase extraction and high-performance liquid chromatography [J. Chromatogr. B, 724 (1999) 231–238] 729(1999)379

Adenoviruses

Recovery in aqueous two-phase systems of nanoparticulates applied as surrogate mimics for viral gene therapy vectors 743(2000)409

Adenovirus type 5

Reversed-phase high-performance liquid chromatographic assay for the adenovirus type 5 proteome 732(1999)411

Adenylosuccinate

Analysis of diagnostic metabolites by capillary electrophoresis–mass spectrometry 727(1999)43

ADP

Assay of ATPase and Na,K-ATPase activity using high-performance liquid chromatographic determination of ADP derived from ATP 744(2000)19

Affinity partitioning

Affinity partitioning of a poly(histidine)-tagged integral membrane protein, cytochrome *bo*₃ ubiquinol oxidase, in a detergent–polymer aqueous two-phase system containing metal-chelating polymer 743(2000)307

Agarose

Essential role of the concentration of immobilized ligands in affinity chromatography: Purification of guanidinobenzoate on an ionized ligand 740(2000)211

Aglycones

Gemfibrozil and its oxidative metabolites: quantification of aglycones, acyl glucuronides, and covalent adducts in samples from preclinical and clinical kinetic studies 741(2000)129

Agmatine

Essential role of the concentration of immobilized ligands in affinity chromatography: Purification of guanidinobenzoate on an ionized ligand 740(2000)211

Airway surface fluid

Analysis of small-scale biological compartments by capillary electrophoresis 745(2000)127

β-Alanine

Combined method for the determination of γ-aminobutyric and β-alanine in cerebrospinal fluid by stable isotope dilution mass spectrometry 732(1999)245

Albumin

Automated isolation of high-purity plasma albumin for isotope ratio measurements 731(1999)199

Affinity chromatography of plasma proteins (guanidinobenzoate): use of mimetic matrices and mimetic soluble ligands to prevent the binding of albumin on target affinity matrices 732(1999)165

Alcohol

Conformational transition and mass transfer in extraction of proteins by AOT–alcohol–isooctane reverse micellar systems 743(2000)203

X-ALD

Gas chromatographic–mass spectrometric determination of plasma saturated fatty acids using pentafluorophenyldimethylsilyl derivatization 742(2000)37

Aldose reductase

D-Xylose metabolism by *Candida intermedia*: isolation and characterisation of two forms of aldose reductase with different coenzyme specificities 737(2000)195

Alinastine

Semi-automated solid-phase extraction procedure for the high-performance liquid chromatographic determination of alinastine in biological fluids 727(1999)131

Alkaline xylanase

Extraction in aqueous two-phase systems of alkaline xylanase produced by *Bacillus pumilus* and its application in kraft pulp bleaching 743(2000)349

Alkaloids

Poisoning by plant material: review of human cases and analytical determination of main toxins by high-performance liquid chromatography–(tandem) mass spectrometry 733(1999)181

Alkylaminoanthraquinone N-oxide

High-performance liquid chromatographic analysis of AQ4N, an alkylaminoanthraquinone *N*-oxide 742(2000)239

Alkylating agents

High-performance liquid chromatographic method for sensitive determination of the alkylating agent CB1954 in human plasma 731(1999)293

Alkylene bisdithiocarbamates

Determination of urinary 2-thiazolidinethione-4-carboxylic acid after exposure to alkylene bisdithiocarbamates using gas chromatography–mass spectrometry 726(1999)85

Alkylphenols

Determination of 4-nonylphenol, nonylphenol monoethoxylate, nonylphenol diethoxylate and other alkylphenols in fish and shellfish by high-performance liquid chromatography with fluorescence detection 746(2000)305

Allantoin

Determination of urea, allantoin and lysine pyroglutamate in cosmetic samples by hydrophilic interaction chromatography 742(2000)447

Quantification of allantoin, uric acid, xanthine and hypoxanthine in ovine urine by high-performance liquid chromatography and photodiode array detection 744(2000)129

Determination of minute amounts of D-leucine in various brain regions of rat and mouse using column-switching high-performance liquid chromatography 744(2000)213

Allysine

Bisphenol derivative of allysine for high-performance liquid chromatographic analysis of allysine residue of proteins 739(2000)273

Alphaxalone

High-performance liquid chromatography of the neuroactive steroids alphaxalone and pregnanolone in plasma using dansyl hydrazine as fluorescent label: application to a pharmacokinetic-pharmacodynamic study in rats 745(2000)357

Amanitin

Validated electrospray liquid chromatographic–mass spectrometric assay for the determination of the mushroom toxins α - and β -amanitin in urine after immunoaffinity extraction 748(2000)125

Ambroxol

Sensitive method for the determination of ambroxol in body fluids by capillary electrophoresis and fluorescence detection 742(2000)205

Amines

Analysis of amines in plant materials 747(2000)49

Amino acid digests

Rapid, single-step procedure for the identification of transglutaminase-mediated isopeptide crosslinks in amino acid digests 732(1999)65

Amino acids

Quantitative determination of free intracellular amino acids in single human polymorphonuclear leucocytes. Recent developments in sample preparation and high-performance liquid chromatography 728(1999)157

High-performance liquid chromatographic analysis of physiological amino acids in human brain tumors by pre-column derivatization with phenylisothiocyanate 730(1999)81

Modifications in electrospray tandem mass spectrometry for a neonatal-screening pilot study in Japan 731(1999)97

Analyzing mixtures of amino acids and carbohydrates using bimodal integrated amperometric detection 732(1999)193

Development of a protocol for the automated analysis of amino acids in brain tissue samples and microdialysates 735(1999)133

Simultaneous determination of the enantiomers of leucine and [$^2\text{H}_7$]leucine in plasma by capillary gas chromatography–mass spectrometry 735(1999)141

Convenient method of threonine, methionine and their related amino compounds by high-performance liquid chromatography and its application to rumen fluid 741(2000)279

Optimization of a capillary electrophoresis–electrospray mass spectrometry method for the quantitation of the 20 natural amino acids in children's blood 742(2000)163

Partitioning behavior of amino acids in aqueous two-phase systems with recyclable volatile salts 743(2000)317

Precise analysis of primary amino acids in urine by an automated high-performance liquid chromatography method: comparison with ion-exchange chromatography 744(2000)183

Determination of minute amounts of D-leucine in various brain regions of rat and mouse using column-switching high-performance liquid chromatography 744(2000)213

Chiral separation of amino acids in biological fluids by micellar electrokinetic chromatography with laser-induced fluorescence detection 745(2000)389

Determination of L-iminoethyl-L-lysine in serum by liquid chromatography 749(2000)35

p-Aminobenzamidine

Affinity chromatography for purification of two urokinases from human urine 742(2000)71

 γ -Aminobutyric acid

Simultaneous determination of γ -aminobutyric acid and glutamic acid in the brain of 3-mercaptopropionic acid-treated rats using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 726(1999)285

Combined method for the determination of γ -aminobutyric and β -alanine in cerebrospinal fluid by stable isotope dilution mass spectrometry 732(1999)245

On-line derivatization for continuous and automatic monitoring of brain extracellular glutamate levels in anesthetized rats: a microdialysis study 734(1999)1

Monitoring γ -aminobutyric acid in human brain and plasma microdialysates using micellar electrokinetic chromatography and laser-induced fluorescence detection 735(1999)1

Neutral amino acids monitoring in phenylketonuric plasma microdialysates using micellar electrokinetic chromatography and laser-induced fluorescence detection 739(2000)247

 ϵ -Aminocaproic acid

Improved purification of the membrane-bound hydrogenase–sulfur-reductase complex from thermophilic archaea using ϵ -aminocaproic acid-containing chromatography buffers 737(2000)151

6-Aminocaproic acid

Affinity chromatography for purification of two urokinases from human urine 742(2000)71

Aminoethylcysteine ketimine decarboxylated dimer

Identification of aminoethylcysteine ketimine decarboxylated dimer in human plasma 732(1999)137

Aminoglycosides

Current methodologies for the analysis of aminoglycosides 747(2000)69

Amino groups

Separation methods for amino group-possessing pesticides in biological samples 747(2000)241

***p*-Aminohippuric acid**

High-performance liquid chromatographic determination of *p*-aminohippuric acid and iothalamate in human serum and urine: comparison of two sample preparation methods 740(2000)81

Aminohydantoins

Simultaneous determination of enantioselective plasma protein binding of aminohydantoins by ultrafiltration and chiral high-performance liquid chromatography 732(1999)31

3-Amino-2-hydroxypropylmethylphosphinic acid

Determination of the GABA_B receptor agonist CGP 44532 (3-amino-2-hydroxypropylmethylphosphinic acid) in rat plasma after pre-column derivatization by micro-high-performance liquid chromatography combined with negative electrospray tandem mass spectrometry 748(2000)349

5-Aminolevulinic acid

Measurement of 4,5-dioxovaleric acid by high-performance liquid chromatography and fluorescence detection 729(1999)237

2-Amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine

Mass spectrometric detection and measurement of *N*²-(2'-deoxyguanosin-8-yl)PhIP adducts in DNA 744(2000)55

Aminopeptidase

Purification and characterization of an aminopeptidase from the chloroplast stroma of barley leaves by chromatographic and electrophoretic methods 737(2000)285

***S*-2-(3-Aminopropylamino)ethanethiol**

Measurement of both protein-bound and total *S*-2-(3-aminopropylamino)ethanethiol (WR-1065) in blood by high-performance liquid chromatography 739(2000)357

Amoxicillin

Application of a semipermeable surface column for the determination of amoxicillin in human blood serum 726(1999)169

Simultaneous quantification of amoxycillin and metronidazole in plasma using high-performance liquid chromatography with photodiode array detection 731(1999)261

Determination of amoxicillin in plasma samples by capillary electrophoresis 731(1999)309

Amperometric detection

A sensitive high-performance liquid chromatographic method using electrochemical detection for the analysis of olanzapine and desmethylolanzapine in plasma of schizophrenic patients using a new solid-phase extraction procedure 750(2001)137

Amphetamine

Simultaneous determination of dimethylamphetamine and its metabolites in rat hair by gas chromatography–mass spectrometry 741(2000)163

Determination of enantiomeric amphetamines as metabolites of illicit amphetamines and selegiline in urine by capillary electrophoresis using modified β -cyclodextrin 741(2000)221

Amphetamines

Routine analysis of amphetamine class drugs as their naphthaquinone derivatives in human urine by high-performance liquid chromatography 735(1999)229

Simple and reliable high-performance liquid chromatography fluorimetric procedure for the determination of amphetamine-derived designer drugs 735(1999)243

Ampicillin

Simple high-performance liquid chromatography determination of ampicillin in human serum using solid-phase extraction disk cartridges 727(1999)245

Amprenavir

Simultaneous determination of the HIV protease inhibitors indinavir, amprenavir, saquinavir, ritonavir, nelfinavir and the non-nucleoside reverse transcriptase inhibitor efavirenz by high-performance liquid chromatography after solid-phase extraction 740(2000)43

Sensitive liquid chromatographic assay for amprenavir, a human immunodeficiency virus protease inhibitor, in human plasma, cerebrospinal fluid and semen 742(2000)185

Simultaneous high-performance liquid chromatographic determination of the antiretroviral agents amprenavir, nelfinavir, ritonavir, saquinavir, delavirdine and efavirenz in human plasma 742(2000)453

 α -Amylase

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

New approach for separating *Bacillus subtilis* metalloprotease and α -amylase by affinity chromatography and for purifying neutral protease by hydrophobic chromatography 737(2000)277

Anabolic steroids

Comparison of sensitivity between gas chromatography–low-resolution mass spectrometry and gas chromatography–high-resolution mass spectrometry for determining metandienone metabolites in urine 734(1999)179

Metabolism of anabolic steroids by recombinant human cytochrome P450 enzymes. Gas chromatographic–mass spectrometric determination of metabolites 735(1999)73

Gas chromatographic–tandem mass spectrometric determination of anabolic steroids and their esters in hair. Application in doping control and meat quality control 735(1999)189

Testing of the anabolic stanozolol in human hair by gas chromatography–negative ion chemical ionization mass spectrometry 740(2000)265

High-performance liquid chromatographic optimization study for the separation of natural and synthetic anabolic steroids. Application to urine and pharmaceutical samples 742(2000)1

Anandamide

Novel and sensitive method for the detection of anandamide by the use of its dansyl derivative 740(2000)93

Anaplastic lymphoma kinase

Fractionation of polyclonal antibodies to fragments of a neuroreceptor using three increasingly chaotropic solvents 728(1999)49

Angiotensin

Determination of angiotensin metabolites in human plasma by fluorimetric high-performance liquid chromatography using a heart-cut column-switching technique 729(1999)89

Angiotensin II receptor antagonist

Determination of candesartan cilexetil, candesartan and a metabolite in human plasma and urine by liquid chromatography and fluorometric detection 731(1999)411

Anhydroecgonine methyl ester

Gas chromatographic–mass spectrometric detection of anhydroecgonine methyl ester (methylecgonidine) in human serum as evidence of recent smoking of crack 735(1999)127

Antagonist G

Development of a gradient elution high-performance liquid chromatography assay with ultraviolet detection for the determination in plasma of the anticancer peptide [Arg⁶, D-Trp^{7,9}, MePhe⁸]-substance P (6–11) (antagonist G), its major metabolites and a C-terminal pyrene-labelled conjugate 732(1999)277

Anthracyclines

Simultaneous determination of four anthracyclines and three metabolites in human serum by liquid chromatography–electrospray mass spectrometry 738(2000)281

Anthranilic acid derivatives

Simultaneous analysis of anthranilic acid derivatives in pharmaceuticals and human urine by high-performance liquid chromatography with isocratic elution 744(2000)81

Antibiotics

Simultaneous quantification of amoxycillin and metronidazole in plasma using high-performance liquid chromatography with photodiode array detection 731(1999)261

Determination of amoxicillin in plasma samples by capillary electrophoresis 731(1999)309

Antibodies

Immobilization of antibodies onto glass wool 738(2000)217

Antidepressants

Separation of eleven central nervous system drugs by capillary zone electrophoresis 728(1999)263

Antifungal agent

High-performance liquid chromatographic analysis of the antifungal agent SCH 56592 in dog serum 738(2000)93

Antifungal agents

Novel sample preparation method facilitating identification of urinary drug metabolites by liquid chromatography–tandem mass spectrometry 738(2000)259

Anti-methotrexate antibody fragments

Highly sensitive high-performance liquid chromatographic assay for methotrexate in the presence and absence of anti-methotrexate antibody fragments in rat and mouse plasma 736(1999)191

Anti-psychotic drugs

Electrospray ionisation–mass spectrometric characterisation of selected anti-psychotic drugs and their detection and determination in human hair samples by liquid chromatography–tandem mass spectrometry 740(2000)141

Antipsychotics

Separation of eleven central nervous system drugs by capillary zone electrophoresis 728(1999)263

Antipyrine

Determination of phenolic derivatives of antipyrine in plasma with solid-phase extraction and high-performance liquid chromatography–atmospheric-pressure chemical ionization mass spectrometry 732(1999)103

Aplidine

Bioanalysis of aplidine, a new marine antitumoral depsipeptide, in plasma by high-performance liquid chromatography after derivatization with *trans*-4'-hydrazino-2-stilbazole 729(1999)43

Analysis of aplidine (dehydrodidemnin B), a new marine-derived depsipeptide, in rat biological fluids by liquid chromatography–tandem mass spectrometry 731(1999)335

Apolipoprotein B-48

Sodium dodecyl sulphate gel electrophoretic preparation of protein standard human apolipoprotein B-48 742(2000)421

Aptamer

Detection and plasma pharmacokinetics of an anti-vascular endothelial growth factor oligonucleotide-aptamer (NX1838) in rhesus monkeys 732(1999)203

AQ4N

High-performance liquid chromatographic analysis of AQ4N, an alkylaminoanthraquinone *N*-oxide 742(2000)239

Arabinosylguanine

High-performance liquid chromatography method for the determination and quantitation of arabinosylguanine triphosphate and fludarabine triphosphate in human cells 745(2000)421

 α_1 -Acid glycoprotein

Hydrophobic interaction chromatography of human serum α_1 -antitrypsin and α_1 -acid glycoprotein 744(2000)73

 α_1 -Antitrypsin

Hydrophobic interaction chromatography of human serum α_1 -antitrypsin and α_1 -acid glycoprotein 744(2000)73

Arginine

Improved method for simultaneous determination of L-arginine and its mono- and dimethylated metabolites in biological samples by high-performance liquid chromatography 742(2000)199

High-performance liquid chromatographic analysis of muscular interstitial arginine and norepinephrine kinetics. A microdialysis study in rats 745(2000)279

Determination of arginine and methylated arginines in human plasma by liquid chromatography–tandem mass spectrometry 748(2000)157

[Arg⁶, D-Trp^{7,9}, MePhe⁸]-substance P (6–11)

Development of a gradient elution high-performance liquid chromatography assay with ultraviolet detection for the determination in plasma of the anticancer peptide [Arg⁶, D-Trp^{7,9}, MePhe⁸]-substance P (6–11) (antagonist G), its major metabolites and a C-terminal pyrene-labelled conjugate 732(1999)277

Aricept

Simultaneous determination of donepezil (aricept[®]) enantiomers in human plasma by liquid chromatography–electrospray tandem mass spectrometry 729(1999)147

Aroma compounds

Aroma compounds recovery from mycelial cultures in aqueous two-phase processes 743(2000)403

Aromatic amines

Haemoglobin adducts of aromatic amines: diamines and polyaromatic amines 744(2000)337

Arsanilic acid

Arsanilic acid–Sephacrose chromatography of pyruvate kinase from KB cells 740(2000)109

Arsenate reductase

Development of a downstream process for the isolation of *Staphylococcus aureus* arsenate reductase overproduced in *Escherichia coli* 737(2000)167

Arsenic

On-line reversed-phase liquid chromatography hydride generation emission spectrometry: speciation of arsenic in urine of patients intravenously treated with As₂O₃ 740(2000)179

Artemether

Simultaneous determination of artemether and its major metabolite dihydroartemisinin in plasma by gas chromatography–mass spectrometry–selected ion monitoring 731(1999)251

Artemisinin

Rapid determination of artemisinin and related analogues using high-performance liquid chromatography and an evaporative light scattering detector 730(1999)71

Direct analysis of artemisinin in plasma and saliva using coupled-column high-performance liquid chromatography with a restricted-access material pre-column 742(2000)155

Ascorbic acid

Rapid mass spectrometric analysis for ascorbate and related organic acids in small volumes of plasma for use in pediatric subjects 726(1999)79

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

aSFP protein

Inhibition of acrosine-like protease activity by a lectin affinity chromatographic bovine seminal plasma fraction containing the PDC-109 and aSFP proteins 746(2000)141

Asiaticoside

Column-switching high-performance liquid chromatographic assay for determination of asiaticoside in rat plasma and bile with ultraviolet absorbance detection 732(1999)357

Asparagine-linked sugar chain

Serial lectin affinity chromatography demonstrates altered asparagine-linked sugar-chain structures of prostate-specific antigen in human prostate carcinoma 727(1999)9

Aspartate

On-line derivatization for continuous and automatic monitoring of brain extracellular glutamate levels in anesthetized rats: a microdialysis study 734(1999)1

Asymmetric dimethylarginine

Assessment of nitric oxide synthase activity in vitro and in vivo by gas chromatography–mass spectrometry 742(2000)143

Atenolol

Two reproducible and sensitive liquid chromatographic methods to quantify atenolol and propranolol in human plasma and determination of their associated analytical error functions 738(2000)225

Automated liquid chromatographic determination of atenolol in plasma using dialysis and trace enrichment on a cation-exchange precolumn for sample handling 739(2000)205

Enantioselective analysis of (*R*)- and (*S*)-atenolol in urine samples by a high-performance liquid chromatography column-switching setup 740(2000)219

Atovaquone

Simultaneous separation of atovaquone, proguanil and its metabolites on a mixed mode high-performance liquid chromatographic column 741(2000)189

Capillary zone electrophoresis for the determination of atovaquone in serum 742(2000)441

Automated solid-phase extraction method for the determination of atovaquone in plasma and whole blood by rapid high-performance liquid chromatography 744(2000)9

ATPase

Assay of ATPase and Na,K-ATPase activity using high-performance liquid chromatographic determination of ADP derived from ATP 744(2000)19

Atractyloside

Poisoning by plant material: review of human cases and analytical determination of main toxins by high-performance liquid chromatography–(tandem) mass spectrometry 733(1999)181

Attachment protein G

Purification of respiratory syncytial virus F and G proteins 737(2000)97

Avidin–biotin-immobilized liposome

Avidin–biotin-immobilized liposome column for chromatographic fluorescence on-line analysis of solute–membrane interactions 750(2001)51

Azaperol

Gas chromatographic–mass spectrometric analysis of veterinary tranquillizers in urine: evaluation of method performance 734(1999)113

Determination of a method for detecting and quantifying azaperone, azaperol and carazolol in pig tissues by liquid chromatography–tandem mass spectrometry 744(2000)139

Azaperone

Gas chromatographic–mass spectrometric analysis of veterinary tranquillizers in urine: evaluation of method performance 734(1999)113

Determination of a method for detecting and quantifying azaperone, azaperol and carazolol in pig tissues by liquid chromatography–tandem mass spectrometry 744(2000)139

Azithromycin

Determination of erythromycin, clarithromycin, roxithromycin, and azithromycin in plasma by high-performance liquid chromatography with amperometric detection 738(2000)405

Azo dyes

Naphthol- and resorcinol-based azo dyes as metal ion complexants in aqueous biphasic systems 743(2000)107

Azacyclonal

Analysis of hydroxylated and *N*-dealkylated metabolites of terfenadine in microsomal incubates by liquid chromatography–mass spectrometry 741(2000)145

B

Benzimidazole

Determination of benzimidazole residues using liquid chromatography and tandem mass spectrometry 727(1999)167

Benzodiazepines

Determination of new 2,3-benzodiazepines in rat plasma using high-performance liquid chromatography with ultraviolet detection 731(1999)207

Fast liquid chromatographic–mass spectrometric determination of pharmaceutical compounds 732(1999)445

High-performance liquid chromatographic method for the determination of benzodiazepines in plasma or serum using the column-switching technique 742(2000)381

Liquid-phase microextraction as a sample preparation technique prior to capillary gas chromatographic-determination of benzodiazepines in biological matrices 749(2000)85

Simple and sensitive high-performance liquid chromatographic method for the determination of 1,5-benzodiazepine clobazam and its active metabolite *N*-desmethyloclobazam in human serum and urine with application to 1,4-benzodiazepines analysis 750(2001)41

2,3-Benzodiazepines

Chromatographic separation of 2,3-benzodiazepines
747(2000)203

Benzoic acid

Metabolism of an ionic contrast medium and the related agents
746(2000)25

Benzoin

Bioanalysis of *m*-iodobenzylguanidine in plasma by high-performance liquid chromatography after derivatization with benzoin 730(1999)193

Benzo[a]pyrene

Extraction and high-performance liquid chromatographic separation of selected pyrene and benzo[a]pyrene sulfates and glucuronides: preliminary application to the analysis of smokers' urine 728(1999)209

Biomonitoring of environmental polycyclic aromatic hydrocarbon exposure by simultaneous measurement of urinary phenanthrene, pyrene and benzo[a]pyrene hydroxides
739(2000)225

Benzothiazole

Analysis of 2-(3-methyl-4-aminophenyl)-benzothiazole (NSC 674495) in plasma by gas chromatography with mass-selective detection 732(1999)315

Benzoyllecgonine

High-performance liquid chromatographic determination of cocaine and its metabolites in serum microsamples with fluorimetric detection and its application to pharmacokinetics in rats 745(2000)315

Benzoylnorecgonine

High-performance liquid chromatographic determination of cocaine and its metabolites in serum microsamples with fluorimetric detection and its application to pharmacokinetics in rats 745(2000)315

Berenil

Determination by capillary zone electrophoresis of berenil, phenamidine, diampron and dibromopropamidine in serum and urine 738(2000)293

Betulinic acid

Determination of betulinic acid in mouse blood, tumor and tissue homogenates by liquid chromatography–electrospray mass spectrometry 732(1999)331

Bifonazole

Chromatographic approach to study β -cyclodextrin as a promoter of the penetration of bifonazole into keratinic tissue
735(1999)289

Bile acids

Determination of bile acids in human faecal samples using supercritical fluid extraction and high-performance liquid chromatography 726(1999)71

Bioanalysis

Simultaneous determination of bosentan and its three major metabolites in various biological matrices and species using narrow bore liquid chromatography with ion spray tandem mass spectrometric detection 749(2000)67

Biofluids

Analysis of small-scale biological compartments by capillary electrophoresis 745(2000)127

Biogenic amines

Reversed-phase high-performance liquid chromatography of catecholamines and indoleamines using a simple gradient solvent system and native fluorescence detection 744(2000)221

Analysis of biogenic amines in microdialysates of the brain
747(2000)21

Determination of biogenic amines in fish implicated in food poisoning by micellar electrokinetic capillary chromatography
749(2000)163

Bioligands, free

Bioaffinity extraction of glucoamylase in aqueous two-phase systems using starch as free bioligand 743(2000)241

Biotin

Affinity partitioning of biotinylated mixed liposomes: effect of charge on biotin–NeutrAvidin interaction 743(2000)389

Biphenyldimethyl dicarboxylate

Rapid microbore liquid chromatographic analysis of biphenyldimethyl dicarboxylate in human plasma with on-line column switching 738(2000)175

Bisphenol

Bisphenol derivative of allysine for high-performance liquid chromatographic analysis of allysine residue of proteins
739(2000)273

Bisphenol A

Determination of bisphenol A in human serum by high-performance liquid chromatography with multi-electrode electrochemical detection 749(2000)17

High-performance liquid chromatography with peroxyoxalate chemiluminescence detection of bisphenol A migrated from polycarbonate baby bottles using 4-(4,5-diphenyl-1*H*-imidazol-2-yl)benzoyl chloride as a label 749(2000)49

 β -Blockers

Two reproducible and sensitive liquid chromatographic methods to quantify atenolol and propranolol in human plasma and determination of their associated analytical error functions
738(2000)225

Blood group proteins

Immunopurification of the blood group RhD protein from human erythrocyte membranes 735(1999)207

Body fluids

Sensitive method for the determination of ambroxol in body fluids by capillary electrophoresis and fluorescence detection 742(2000)205

Bolasterone

Optimization of the high-performance liquid chromatographic separation of a complex mixture containing urinary steroids, boldenone and bolasterone: application to urine samples 742(2000)47

Boldenone

Optimization of the high-performance liquid chromatographic separation of a complex mixture containing urinary steroids, boldenone and bolasterone: application to urine samples 742(2000)47

Bosentan

Simultaneous determination of bosentan and its three major metabolites in various biological matrices and species using narrow bore liquid chromatography with ion spray tandem mass spectrometric detection 749(2000)67

Bovine serum albumin

Use of capillary electrophoresis-based competitive immunoassay for a large molecule 727(1999)63

Separation of bovine serum albumin and its monoclonal antibody from their immunocomplexes by sodium dodecyl sulfate–capillary gel electrophoresis and its application in capillary electrophoresis-based immunoassay 731(1999)389

Partitioning of whey proteins, bovine serum albumin and porcine insulin in aqueous two-phase systems 743(2000)235

Bovine serum albumin partitioning in an aqueous two-phase system: Effect of pH and sodium chloride concentration 743(2000)255

Partitioning of bovine serum albumin in an aqueous two-phase system: optimization of partition coefficient 743(2000)259

 β_2 -Agonists

Multiresidue analysis of β_2 -agonists in human and calf urine using multimodal solid-phase extraction and high-performance liquid chromatography with electrochemical detection 726(1999)149

Brequinar

Simultaneous separation by high-performance liquid chromatography of carbamoyl aspartate, carbamoyl phosphate and dihydroorotic acid 732(1999)487

Brombuterol

Multiresidue analysis of β_2 -agonists in human and calf urine using multimodal solid-phase extraction and high-performance liquid chromatography with electrochemical detection 726(1999)149

4-Bromomethyl-7-methoxycoumarin

Derivatization of thymine and thymine photodimers with 4-bromomethyl-7-methoxycoumarin for fluorescence detection of high-performance liquid chromatography 731(1999)179

***p*-Bromophenylacetylurea**

Application of the high-performance liquid chromatographic method for separation, purification and characterisation of *p*-bromophenylacetylurea and its metabolites (1999)349

4-(4-Bromophenyl)-3-oxapyrrolidine-2,5-dione

Application of the high-performance liquid chromatographic method for separation, purification and characterisation of *p*-bromophenylacetylurea and its metabolites (1999)349

(*E*)-5-(2-Bromovinyl)-2'-deoxyuridine

Determination of (*E*)-5-(2-bromovinyl)-2'-deoxyuridine in plasma and urine by capillary electrophoresis 726(1999)261

Development and validation of a bioanalytical assay for (*E*)-5-(2-bromovinyl)-2'-deoxyuridine in plasma by capillary zone electrophoresis 726(1999)269

Bupivacaine

Simultaneous analysis of ketamine and bupivacaine in plasma by high-performance liquid chromatography 728(1999)107

Determination of free concentrations of ropivacaine and bupivacaine in plasma from neonates using small-scale equilibrium–dialysis followed by liquid chromatography–mass spectrometry 748(2000)321

Busulfan

Determination of tetrahydrothiophene formation as a probe of in vitro busulfan metabolism by human glutathione *S*-transferase A1-1: use of a highly sensitive gas chromatographic–mass spectrometric method 730(1999)25

Determination of busulfan in human plasma using high-performance liquid chromatography with pre-column derivatization and fluorescence detection 730(1999)33

Rapid and sensitive high-performance liquid chromatographic method for busulfan assay in plasma 742(2000)427

1,3-Butadiene

Gas chromatographic determination of 3-butene-1,2-diol in urine samples after 1,3-butadiene exposure 730(1999)257

3-Butene-1,2-diol

Gas chromatographic determination of 3-butene-1,2-diol in urine samples after 1,3-butadiene exposure 730(1999)257

γ -Butyrobetaine

Effect of sports activity on carnitine metabolism. Measurement of free carnitine, γ -butyrobetaine and acylcarnitines by tandem mass spectrometry 731(1999)83

Measurement of carnitine precursors, ε -trimethyllysine and γ -butyrobetaine in human serum by tandem mass spectrometry 731(1999)89

Butyrophenones

Analyses of butyrophenones and their analogues in whole blood by high-performance liquid chromatography–electrospray tandem mass spectrometry 746(2000)3

BW935U83

Application of directly coupled LC–NMR–MS to the structural elucidation of metabolites of the HIV-1 reverse-transcriptase inhibitor BW935U83 748(2000)269

C**Cadmium**

Interference by *S*-nitroso compounds with the measurement of nitrate in methods requiring reduction of nitrate to nitrite by cadmium 729(1999)375

Caffeic acid

Measurement of unbound caffeic acid in rat blood by on-line microdialysis coupled with liquid chromatography and its application to pharmacokinetic study 729(1999)119

Simultaneous analysis of shikimate-derived secondary metabolites in *Lithospermum erythrorhizon* cell suspension cultures by high-performance liquid chromatography 738(2000)3

Caffeine

High-performance liquid chromatographic method for simultaneous determination of [1-methyl- 14 C]caffeine and its eight major metabolites in rat urine 726(1999)195

Liquid chromatographic method for the simultaneous determination of caffeine and fourteen caffeine metabolites in urine 746(2000)331

Calanolide A

Quantification of (+)-calanolide A, a novel and naturally occurring anti-HIV agent, by high-performance liquid chromatography in plasma from rat, dog and human 742(2000)267

Calcium

Chromatographic study of magnesium and calcium binding to immobilized human serum albumin 728(1999)167

On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography 750(2001)61

Calcium-binding proteins

Purification of the Ca^{2+} -binding protein S100A1 from myocardium and recombinant *Escherichia coli* 737(2000)39

Calmodulin antagonist

Determination of a novel calmodulin antagonist, 3-[2-[4-(3-chloro-2-methylphenyl)-1-piperazinyl]ethyl]-5,6-dimethoxy-1-(4-imidazolylmethyl)-1*H*-indazole dihydrochloride 3.5 hydrate, DY-9760e, in human plasma using solid-phase extraction and high-performance liquid chromatography with fluorescence detection 734(1999)39

Campestanol

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Campesterol

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Camphor

Sensitive and selective gas chromatographic methods for the quantitation of camphor, menthol and methyl salicylate from human plasma 729(1999)163

Camptothecin

Determination of unbound 20(*S*)-camptothecin in rat bile by on-line microdialysis coupled to microbore liquid chromatography with fluorescence detection 732(1999)221

Sensitive high-performance liquid chromatographic method for the determination of the lactone form and the lactone plus hydroxy-acid forms of the new camptothecin derivative DX-8951 in human plasma using fluorescence detection 740(2000)237

Candesartan

Determination of candesartan cilexetil, candesartan and a metabolite in human plasma and urine by liquid chromatography and fluorometric detection 731(1999)411

Cannabinoids

Chromatographic procedures for determination of cannabinoids in biological samples, with special attention to blood and alternative matrices like hair, saliva, sweat and meconium 733(1999)119

Capillary electrophoresis

Determination of asymmetrical dimethylarginine by capillary electrophoresis–laser-induced fluorescence 741(2000)77

Measurement of chloramphenicol by capillary zone electrophoresis following end-column amperometric detection at a carbon fiber micro-disk array electrode 741(2000)155

Sensitive method for the determination of ambroxol in body fluids by capillary electrophoresis and fluorescence detection 742(2000)205

Analysis of small-scale biological compartments by capillary electrophoresis 745(2000)127

Capillary zone electrophoresis

Kinetics of binding properties of 5'-GMP with cisplatin under simulated physiological conditions by capillary electrophoresis 745(2000)211

Carazolol

Determination of a method for detecting and quantifying azaperone, azaperol and carazolol in pig tissues by liquid chromatography–tandem mass spectrometry 744(2000)139

Carbamazepine

Fast liquid chromatographic–mass spectrometric determination of pharmaceutical compounds 732(1999)445

Carbamazepine-10,11-*trans*-dihydrodiol

Simultaneous stereoselective high-performance liquid chromatographic determination of 10-hydroxycarbamazepine and its metabolite carbamazepine-10,11-*trans*-dihydrodiol in human urine 738(2000)419

L-*N*-Carbamoylase

Purification of recombinant hydantoinase and L-*N*-carbamoylase from *Arthrobacter aurescens* expressed in *Escherichia coli*: comparison of wild-type and genetically modified proteins 737(2000)179

Carbamoyl aspartate

Simultaneous separation by high-performance liquid chromatography of carbamoyl aspartate, carbamoyl phosphate and dihydroorotic acid 732(1999)487

Carbamoyl phosphate

Simultaneous separation by high-performance liquid chromatography of carbamoyl aspartate, carbamoyl phosphate and dihydroorotic acid 732(1999)487

Carbapenems

Simple and rapid analytical method for carbapenems using capillary zone electrophoresis 727(1999)219

Carbohydrate-deficient transferrin

Improved method for carbohydrate-deficient transferrin determination in human serum by capillary zone electrophoresis 739(2000)81

Carbohydrates

Analyzing mixtures of amino acids and carbohydrates using bimodal integrated amperometric detection 732(1999)193

Carbohydrate release from picomole quantities of glycoprotein and characterisation of glycans by high-performance liquid chromatography and mass spectrometry 734(1999)169

Carbon dioxide

Isoelectric precipitation of soybean protein using carbon dioxide as a volatile acid 743(2000)357

Carboplatin

Validation of high-performance liquid chromatographic assay methods for the analysis of carboplatin in plasma ultrafiltrate 744(2000)367

Carboxyethylphosphoramidate mustard

Analysis of cyclophosphamide and five metabolites from human plasma using liquid chromatography–mass spectrometry and gas chromatography–nitrogen–phosphorus detection 732(1999)287

Carboxymefloquine

Improved validated assay for the determination of mefloquine and its carboxy metabolite in plasma, serum and whole blood using solid-phase extraction and high-performance liquid chromatography 727(1999)159

Carnitine

Effect of sports activity on carnitine metabolism. Measurement of free carnitine, γ -butyrobetaine and acylcarnitines by tandem mass spectrometry 731(1999)83

Determination of carnitine and acylcarnitines in biological samples by capillary electrophoresis–mass spectrometry 735(1999)171

Carnitine precursors

Measurement of carnitine precursors, ϵ -trimethyllysine and γ -butyrobetaine in human serum by tandem mass spectrometry 731(1999)89

Carotenoids

Stability of individual carotenoids, retinol and tocopherols in human plasma during exposure to light and after extraction 729(1999)191

Carvedilol

Rapid and simple micro-determination of carvedilol in rat plasma by high-performance liquid chromatography 732(1999)233

Catechin

Analysis of (+)-catechin, (–)-epicatechin and their 3'- and 4'-*O*-methylated analogs. A comparison of sensitive methods 726(1999)277

Catecholamines

Determination of differential activities of soluble and membrane-bound catechol-*O*-methyltransferase in tissues and erythrocytes 729(1999)347

Determination of catecholamines in human plasma by high-performance liquid chromatography with electrochemical detection 730(1999)201

Sensitive and specific method for the simultaneous determination of natural and synthetic catecholamines and 3,4-dihydroxyphenylglycol in microdialysis samples 730(1999)213

Simultaneous determination of catecholamines in rat brain tissue by high-performance liquid chromatography 731(1999)361

Reversed-phase high-performance liquid chromatography of catecholamines and indoleamines using a simple gradient solvent system and native fluorescence detection 744(2000)221

Liquid chromatography–electrospray tandem mass spectrometry of acidic monoamine metabolites 744(2000)423

High-performance liquid chromatographic separation and measurement of various biogenic compounds possibly involved in the pathomechanism of Parkinson's disease 746(2000)297

High-performance liquid chromatographic assay for catecholamines and metanephrines using fluorimetric detection with pre-column 9-fluorenylmethyloxycarbonyl chloride derivatization 749(2000)179

Determination of urine catecholamines by capillary electrophoresis with dual-electrode amperometric detection 750(2001)33

Catechol estrogens

Studies on neurosteroids XII. Determination of enzymatically formed catechol estrogens and guaiacol estrogens by rat brains using liquid chromatography–mass spectrometry–mass spectrometry 748(2000)89

Catechol *O*-methyltransferase

Radiochemical high-performance liquid chromatographic assay for the determination of catechol *O*-methyltransferase activity towards various substrates 736(1999)143

Catechol-*O*-methyltransferase

Determination of differential activities of soluble and membrane-bound catechol-*O*-methyltransferase in tissues and erythrocytes 729(1999)347

Gas chromatographic–mass spectrometric procedures for determination of the catechol-*O*-methyltransferase (COMT) activity and for detection of unstable catecholic metabolites in human and rat liver preparations after COMT catalyzed *in statu nascendi* derivatization using *S*-adenosylmethionine 739(2000)325

Cathepsin A

High-performance liquid chromatographic–fluorimetric assay for cathepsin A (lysosomal protective protein) activity 728(1999)59

Cationic–anionic surfactants

Phase behavior and protein partitioning in aqueous two-phase systems of cationic–anionic surfactant mixtures 743(2000)327

CB1954

High-performance liquid chromatographic method for sensitive determination of the alkylating agent CB1954 in human plasma 731(1999)293

Cefepime

Determination of cefepime and ceftiofime in human serum by high-performance liquid chromatography using an ultrafiltration for antibiotics serum extraction 734(1999)121

Cefmatilen

Determination of a new oral cephalosporin, cefmatilen hydrochloride hydrate, and its seven metabolites in human and animal plasma and urine by coupled systems of ion-exchange and reversed-phase high-performance liquid chromatography 749(2000)101

Cefmetazole

Determination of unbound cefmetazole in rat blood by on-line microdialysis and microbore liquid chromatography: a pharmacokinetic study 736(1999)129

Cefotaxime

Concurrent quantification and pharmacokinetic analysis of cefotaxime in rat blood and brain by microdialysis and microbore liquid chromatography 738(2000)75

Simultaneous determination of cefotaxime and desacetylcefotaxime in human plasma and cerebrospinal fluid by high-performance liquid chromatography 750(2001)171

Ceftazidimine

Measurement and pharmacokinetic analysis of unbound ceftazidime in rat blood using microdialysis and microbore liquid chromatography 750(2001)93

Ceftriaxone

Simple liquid chromatographic method for the analysis of the blood brain barrier permeability characteristics of ceftriaxone in an experimental rabbit meningitis model 728(1999)97

Cefuroxime

Simultaneous measurement of cefuroxime in rat blood and brain by microdialysis and microbore liquid chromatography. Application to pharmacokinetics 735(1999)25

Celecoxib

Determination of celecoxib in human plasma by normal-phase high-performance liquid chromatography with column switching and ultraviolet absorbance detection 738(2000)377

Cellulose

Investigation of aqueous biphasic systems for the separation of lignins from cellulose in the paper pulping process 743(2000)127

Centropazine

Determination of centropazine and its metabolite in rat serum by high-performance liquid chromatography and fluorescence detection 748(2000)331

Cephalexin

Simultaneous blood and brain sampling of cephalexin in the rat by microdialysis and microbore liquid chromatography: application to pharmacokinetics studies 740(2000)203

Cephalosporins

Determination of a new oral cephalosporin, S-1090, in human plasma and urine by direct injection high-performance liquid chromatography with ultraviolet detection and column switching 728(1999)125

Determination of the cephalosporin antibiotic cephadrine in human plasma by high-performance liquid chromatography with ultraviolet detection 740(2000)71

Determination of a new oral cephalosporin, cefmatilen hydrochloride hydrate, and its seven metabolites in human and animal plasma and urine by coupled systems of ion-exchange and reversed-phase high-performance liquid chromatography 749(2000)101

Cephalothin

Determination of unbound cephalothin in rat blood by on-line microdialysis and microbore liquid chromatography 742(2000)125

Cephadrine

Determination of the cephalosporin antibiotic cephadrine in human plasma by high-performance liquid chromatography with ultraviolet detection 740(2000)71

Cepirome

Determination of cefepime and cepirome in human serum by high-performance liquid chromatography using an ultrafiltration for antibiotics serum extraction 734(1999)121

Cerivastatin

Quantitation of cerivastatin and its seven acid and lactone biotransformation products in human serum by liquid chromatography–electrospray tandem mass spectrometry 736(1999)19

Cetirizine

Determination of cetirizine in human plasma by high-performance liquid chromatography 736(1999)231

Stereoselective determination of cetirizine and studies on pharmacokinetics in rat plasma 744(2000)201

Cetorelix

Measurement of the novel decapeptide cetorelix in human plasma and urine by liquid chromatography–electrospray ionization mass spectrometry 729(1999)245

Cetylpyridinium chloride

High-performance liquid chromatography determination of residue levels on chicken carcasses treated with cetylpyridinium chloride 728(1999)273

CGP 60536

Automated quantitative determination of the new renin inhibitor CGP 60536 by high-performance liquid chromatography 738(2000)129

Chemokines

High-performance liquid chromatographic purification and capillary electrophoresis quantification of the chemokine stromal cell-derived factor-1 729(1999)369

Multi-step purification strategy for RANTES wild-type and mutated analogues expressed in a baculovirus system 737(2000)47

Chlorambucil

Rapid determination of the anti-cancer drug chlorambucil (LeukeranTM) and its phenyl acetic acid mustard metabolite in human serum and plasma by automated solid-phase extraction and liquid chromatography–tandem mass spectrometry 732(1999)173

Chloramphenicol

Measurement of chloramphenicol by capillary zone electrophoresis following end-column amperometric detection at a carbon fiber micro-disk array electrode 741(2000)155

Chloroacetaldehyde

Improved and simplified liquid chromatographic assay for adefovir, a novel antiviral drug, in human plasma using derivatization with chloroacetaldehyde 736(1999)115

8-Chloroadenine

Determination of pharmacokinetics of 8-chloroadenosine and its two major metabolites in dogs by high-performance liquid chromatography 746(2000)319

8-Chloroadenosine

Determination of pharmacokinetics of 8-chloroadenosine and its two major metabolites in dogs by high-performance liquid chromatography 746(2000)319

8-Chloroinosine

Determination of pharmacokinetics of 8-chloroadenosine and its two major metabolites in dogs by high-performance liquid chromatography 746(2000)319

4-Chloro-7-nitrobenzo-2-oxa-1,3-diazol

Analysis of 4-hydroxyproline using 4-chloro-7-nitrobenzo-2-oxa-1,3-diazol derivatization and micellar electrokinetic chromatography combined with laser-induced fluorescence detection 744(2000)195

1-(2-Chlorophenyl)-*N*-methyl-*N*-(1-methylpropyl)-3-isoquinoline carboxamide

High-performance liquid chromatographic determination of [¹¹C]1-(2-chlorophenyl)-*N*-methyl-*N*-(1-methylpropyl)-3-isoquinoline carboxamide in mouse plasma and tissue and in human plasma 736(1999)61

Chlortetracycline

Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments. Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry 748(2000)369

Chlorthalidone

Chromatographic separation of chlorthalidone enantiomers using β -cyclodextrins as chiral additives 740(2000)169

Chlortoluron

Immuno-affinity solid-phase extraction 745(2000)39

Cholestanol

A new method for determination of serum cholestanol by high-performance liquid chromatography with ultraviolet detection 742(2000)345

Cholesterol

Purification of fatty acid methyl esters by high-performance liquid chromatography 732(1999)495

Separation and quantification of cholesterol and major phospholipid classes in human semen by high-performance liquid chromatography and light-scattering detection 740(2000)101

Choline

Separation methods used in the determination of choline and acetylcholine 747(2000)111

Chondroitin sulfate

Ultrasensitive capillary electrophoresis of sulfated disaccharides in chondroitin/dermatan sulfates by laser-induced fluorescence after derivatization with 2-aminoacridone 730(1999)129

Chymotrypsin

Expression of chymotrypsin(ogen) in the thioredoxin reductase deficient mutant strain of *Escherichia coli* AD494(DE3) and purification via a fusion product with a hexahistidine-tail 737(2000)213

Chymotrypsin inhibitor 2

Model process for separation based on unfolding and refolding of chymotrypsin inhibitor 2 in thermoseparating polymer two-phase systems 743(2000)13

Chymotrypsinogen

Expression of chymotrypsin(ogen) in the thioredoxin reductase deficient mutant strain of *Escherichia coli* AD494(DE3) and purification via a fusion product with a hexahistidine-tail 737(2000)213

Cibacron blue F3GA

Human serum albumin chromatography by Cibacron Blue F3GA-derived microporous polyamide hollow-fiber affinity membranes 746(2000)123

Cilostazol

Simultaneous quantitative determination of cilostazol and its metabolites in human plasma by high-performance liquid chromatography 728(1999)251

Cinnabarinic acid

3-Hydroxyanthranilic acid-derived compounds formed through electrochemical oxidation 736(1999)237

Ciprofibrate

Achiral and chiral determination of ciprofibrate and its glucuronide in human urine by capillary electrophoresis 729(1999)33

Ciprofloxacin

Simultaneous determination of enrofloxacin and ciprofloxacin in animal biological fluids by high-performance liquid chromatography. Application in pharmacokinetic studies in pig and rabbit 726(1999)175

Circulating human peptides

Composition of the peptide fraction in human blood plasma: database of circulating human peptides 726(1999)25

Multiresidue analysis of β_2 -agonists in human and calf urine using multimodal solid-phase extraction and high-performance liquid chromatography with electrochemical detection 726(1999)149

Cisapride

Microbore high-performance liquid chromatographic determination of cisapride in rat serum samples using column switching 727(1999)213

Stereoselective determination of cisapride, a prokinetic agent, in human plasma by chiral high-performance liquid chromatography with ultraviolet detection: application to pharmacokinetic study 744(2000)263

Cisplatin

Kinetics of binding properties of 5'-GMP with cisplatin under simulated physiological conditions by capillary electrophoresis 745(2000)211

Citalopram

Simultaneous determination of citalopram, fluoxetine, paroxetine and their metabolites in plasma and whole blood by high-performance liquid chromatography with ultraviolet and fluorescence detection 734(1999)229

Citrulline

Analysis of citrulline in rat brain tissue after perfusion with haloperidol by liquid chromatography–mass spectrometry 746(2000)33

Clarithromycin

Determination of erythromycin, clarithromycin, roxithromycin, and azithromycin in plasma by high-performance liquid chromatography with amperometric detection 738(2000)405

Clemastine

Determination of clemastine in human plasma by gas chromatography with nitrogen–phosphorus detection 744(2000)177

Clenbuterol

Determination of clenbuterol in bovine urine using gas chromatography–mass spectrometry following clean-up on an ion-exchange resin 728(1999)67

Quantitative analysis of clenbuterol in meat products using liquid chromatography–electrospray ionisation tandem mass spectrometry 736(1999)209

Immuno-affinity solid-phase extraction 745(2000)39

Clobazam

Simple and sensitive high-performance liquid chromatographic method for the determination of 1,5-benzodiazepine clobazam and its active metabolite *N*-desmethyloclobazam in human serum and urine with application to 1,4-benzodiazepines analysis 750(2001)41

Optimised determination of clobazam in human plasma with extraction and high-performance liquid chromatography analysis 750(2001)177

Clodronate

Analysis of an adenine nucleotide-containing metabolite of clodronate using ion pair high-performance liquid chromatography–electrospray ionisation mass spectrometry 738(2000)395

Clozapine

Fishing for a drug: solid-phase microextraction for the assay of clozapine in human plasma 731(1999)231

Coagulation proteins

Separation of human vitamin K-dependent coagulation proteins using hydrophobic interaction chromatography 736(1999)77

Cocaethylene

High-performance liquid chromatographic determination of cocaine and its metabolites in serum microsamples with fluorimetric detection and its application to pharmacokinetics in rats 745(2000)315

Cocaine

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

Comparative analysis of heroin and cocaine seizures 733(1999)127

Sweat testing for cocaine, codeine and metabolites by gas chromatography–mass spectrometry 733(1999)247

Gas chromatographic–mass spectrometric detection of anhydroecgonine methyl ester (methylecgonidine) in human serum as evidence of recent smoking of crack 735(1999)127

Liquid chromatography–tandem mass spectrometry analysis of cocaine and its metabolites from blood, amniotic fluid, placental and fetal tissues: study of the metabolism and distribution of cocaine in pregnant rats 745(2000)287

High-performance liquid chromatographic determination of cocaine and its metabolites in serum microsamples with fluorimetric detection and its application to pharmacokinetics in rats 745(2000)315

Performance of a pentafluorophenylpropyl stationary phase for the electrospray ionization high-performance liquid chromatography–mass spectrometry–mass spectrometry assay of cocaine and its metabolite ecgonine methyl ester in human urine 748(2000)77

Cocaine, metabolites

Liquid chromatography–tandem mass spectrometry analysis of cocaine and its metabolites from blood, amniotic fluid, placental and fetal tissues: study of the metabolism and distribution of cocaine in pregnant rats 745(2000)287

Codeine

Highly sensitive gas chromatographic–tandem mass spectrometric method for the determination of morphine and codeine in serum and urine in the femtomolar range 727(1999)81

Sweat testing for cocaine, codeine and metabolites by gas chromatography–mass spectrometry 733(1999)247

Collagen

Comparison of different electrokinetic separation modes applicable to a model peptide mixture (collagen type I and III CNBr fragments) 745(2000)251

Collagen parent α -chains

Advanced separation methods for collagen parent α -chains, their polymers and fragments 739(2000)3

Collagens

Application of Pluronic copolymer liquid crystals for the capillary electrophoretic separation of collagen type I cyanogen bromide fragments 739(2000)109

Column switching

Enantioselective analysis of (*R*)- and (*S*)-atenolol in urine samples by a high-performance liquid chromatography column-switching setup 740(2000)219

Corticosteroids

Liquid chromatography–mass spectrometry: potential in forensic and clinical toxicology 733(1999)93

Analysis of corticosteroids in hair by liquid chromatography–electrospray ionization mass spectrometry 740(2000)227

Corticosterone

Separation and identification of corticosterone metabolites by liquid chromatography–electrospray ionization mass spectrometry 726(1999)59

Quantitative analysis of human serum corticosterone by high-performance liquid chromatography coupled to electrospray ionization mass spectrometry 727(1999)227

Cortisol

Development of a urinary free cortisol assay using solid-phase extraction–capillary electrophoresis 730(1999)123

Simultaneous determination of endogenous and ^{13}C -labelled cortisols and cortisones in human plasma by stable isotope dilution mass spectrometry 738(2000)119

Simultaneous determination of urinary free cortisol and 6 β -hydroxycortisol by liquid chromatography–atmospheric pressure chemical ionization tandem mass spectrometry and its application for estimating hepatic CYP3A induction 742(2000)303

Specific determination of urinary 6 β -hydroxycortisol and cortisol by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 746(2000)95

Cortisone

Simultaneous determination of endogenous and ^{13}C -labelled cortisols and cortisones in human plasma by stable isotope dilution mass spectrometry 738(2000)119

Cotinine

Improved highly sensitive method for determination of nicotine and cotinine in human plasma by high-performance liquid chromatography 742(2000)211

High-performance liquid chromatographic determination of cotinine in urine in isocratic mode 746(2000)115

Coumarin 7-hydroxylase

Novel sensitive high-performance liquid chromatographic method for assay of coumarin 7-hydroxylation 734(1999)23

Co 102862

Development of a high-performance liquid chromatographic–tandem mass spectrometric method for the determination of pharmacokinetics of Co 102862 in mouse, rat, monkey and dog plasma 749(2000)1

Cpr6

Three-step chromatographic purification of Cpr6, a cyclophilin from *Saccharomyces cerevisiae* 737(2000)295

Creatine

Improved high-performance liquid chromatographic assay for the determination of “high-energy” phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

Analysis of creatine and creatinine in urine by capillary electrophoresis 732(1999)479

Determination of uremic toxins in biofluids: creatinine, creatine, uric acid and xanthines 747(2000)217

Creatinine

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Analysis of creatine and creatinine in urine by capillary electrophoresis 732(1999)479

Determination of uremic toxins in biofluids: creatinine, creatine, uric acid and xanthines 747(2000)217

Cryptophycin 52

Protein-binding patterns of the antitumor antibiotic cryptophycin 52 as measured with a two-phase partitioning system 735(1999)121

Cy5

High-performance liquid chromatography with on-line post-column immunoreaction detection of digoxin and its metabolites based on fluorescence energy transfer in the far-red spectral region 745(2000)305

Cyanogen bromide

Comparison of different electrokinetic separation modes applicable to a model peptide mixture (collagen type I and III CNBr fragments) 745(2000)251

Cyanogenic compounds

Poisoning by plant material: review of human cases and analytical determination of main toxins by high-performance liquid chromatography–(tandem) mass spectrometry 733(1999)181

Cyanogenic glycosides

Analysis of cyanogenic glycosides by micellar capillary electrophoresis 739(2000)95

Cyclodextrin

Cyclodextrins: a versatile tool in separation science
745(2000)83

Cyclodextrins

Separation of eleven central nervous system drugs by capillary zone electrophoresis 728(1999)263

Charged cyclodextrin-mediated sample stacking in micellar capillary electrophoresis. A simple method for enhancing the detection sensitivity of hydrophobic compounds 731(1999)369

Chromatographic approach to study β -cyclodextrin as a promoter of the penetration of bifonazole into keratinic tissue 735(1999)289

Cyclooxygenase-2 inhibitor

Determination of Rofecoxib (MK-0966), a cyclooxygenase-2 inhibitor, in human plasma by high-performance liquid chromatography with tandem mass spectrometric detection 748(2000)31

Cyclophilins

Three-step chromatographic purification of Cpr6, a cyclophilin from *Saccharomyces cerevisiae* 737(2000)295

Cyclophosphamide

Determination of cyclophosphamide and its metabolites in human plasma by high-performance liquid chromatography–mass spectrometry 729(1999)297

Analysis of cyclophosphamide and five metabolites from human plasma using liquid chromatography–mass spectrometry and gas chromatography–nitrogen–phosphorus detection 732(1999)287

Quantitative high-performance liquid chromatographic determination of acrolein in plasma after derivatization with Luminarin® 3 739(2000)239

Simple and selective determination of the cyclophosphamide metabolite phosphoramidate mustard in human plasma using high-performance liquid chromatography 745(2000)345

Cyclosporin A

Development of an immobilized P-glycoprotein stationary phase for on-line liquid chromatographic determination of drug-binding affinities 739(2000)33

Cyclosporine

Automated, fast and sensitive quantification of drugs in blood by liquid chromatography–mass spectrometry with on-line extraction: immunosuppressants 748(2000)41

Cyclosporine A

High-performance liquid chromatographic method for therapeutic drug monitoring of cyclosporine A and its two metabolites in renal transplant patients 749(2000)93

Cysteine

Determination of homocysteine and its related compounds by solid-phase microextraction–gas chromatography–mass spectrometry 727(1999)1

Gas chromatographic–mass spectrometric analysis of stable isotopes of cysteine and glutathione in biological samples 732(1999)127

Kinetics of binding properties of 5'-GMP with cisplatin under simulated physiological conditions by capillary electrophoresis 745(2000)211

5-S-cysteinyl-dopa

Determination of 5-S-cysteinyl-dopa in plasma and urine using a fully automated solid-phase extraction–high-performance liquid chromatographic method for an improvement of specificity and sensitivity of this prognostic marker of malignant melanoma 727(1999)31

Cytochalasin B

Chromatography on cells: analyses of solute interactions with the glucose transporter Glut1 in human red cells adsorbed on lectin-gel beads 739(2000)55

Cytochrome b_5

Purification of the membrane binding domain of cytochrome b_5 by immobilised nickel chelate chromatography 737(2000)119

Cytochrome P450 inhibitors

Amounts and variation in grapefruit juice of the main components causing grapefruit–drug interaction 741(2000)195

Cytochrome P450

Metabolism of anabolic steroids by recombinant human cytochrome P450 enzymes. Gas chromatographic–mass spectrometric determination of metabolites 735(1999)73

Metabolic profiling of valproic acid by cDNA-expressed human cytochrome P450 enzymes using negative-ion chemical ionization gas chromatography–mass spectrometry 742(2000)217

Determination of cytochrome P450 1A activities in mammalian liver microsomes by high-performance liquid chromatography with fluorescence detection 744(2000)399

Cytochromes

High-performance liquid chromatographic method for simultaneous determination of [1-methyl- ^{14}C]caffeine and its eight major metabolites in rat urine 726(1999)195

Purification of cytochrome P-450 from adult pig testis by hydroxylapatite and deoxycorticosterone affinity column chromatography 726(1999)291

Determination of testosterone and 6 β -hydroxytestosterone by gas chromatography–selected ion monitoring–mass spectrometry for the characterization of cytochrome p450 3a activity 734(1999)73

D

Dansyl hydrazine

High-performance liquid chromatography of the neuroactive steroids alphaxalone and pregnanolone in plasma using dansyl hydrazine as fluorescent label: application to a pharmacokinetic-pharmacodynamic study in rats 745(2000)357

Daunorubicin

Simultaneous determination of four anthracyclines and three metabolites in human serum by liquid chromatography–electrospray mass spectrometry 738(2000)281

DDT

High-performance liquid chromatographic method for determination of DDT and its degradation products in rat plasma, liver and brain: validation and application to a pharmacokinetic study 748(2000)361

17-Deacetylnorgestimate

Determination of norgestimate and its metabolites in human serum using high-performance liquid chromatography with tandem mass spectrometric detection 734(1999)247

Debrisoquine

Enantioselectivity of debrisoquine 4-hydroxylation in Brazilian Caucasian hypertensive patients phenotyped as extensive metabolizers 749(2000)153

Dehydroascorbic acid

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Dehydroisoascorbic acid

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Delavirdine

Rapid quantification of delavirdine, a novel non-nucleoside reverse transcriptase inhibitor, in human plasma using isocratic reversed-phase high-performance liquid chromatography with fluorescence detection 727(1999)151

Simultaneous high-performance liquid chromatographic determination of the antiretroviral agents amprenavir, nelfinavir, ritonavir, saquinavir, delavirdine and efavirenz in human plasma 742(2000)453

N-Demethyl-loperamide

Quantitation of loperamide and *N*-demethyl-loperamide in human plasma using electrospray ionization with selected reaction ion monitoring liquid chromatography–mass spectrometry 744(2000)323

Deoxyadenosine

Erratum to “Determination of adenosine and deoxyadenosine in urine by high-performance liquid chromatography with column switching”. [J. Chromatogr. B, 719 (1998) 55–61] 726(1999)317

Deoxyadenosine triphosphate

Separation of near-infrared fluorescent conjugates of dATP and related compounds by capillary electrophoresis 744(2000)315

3-Deoxyglucosone

3-Deoxyglucosone: metabolism, analysis, biological activity, and clinical implication (Review) 731(1999)23

Gas chromatographic–mass spectrometric analysis of erythrocyte 3-deoxyglucosone in hemodialysis patients (Review) 731(1999)37

2-Deoxy-D-glucose

Effect of 2-deoxy-D-glucose on acetylcholine and histamine levels in gastric juice of pylorus-ligated rats anesthetized with urethane 742(2000)295

2'-Deoxyuridine

Modified high-performance liquid chromatography assay for the measurement of 2'-deoxyuridine in human plasma and its application to pharmacodynamic studies of antimetabolite drugs 744(2000)351

Depsipeptide

Bioanalysis of aplidine, a new marine antitumoral depsipeptide, in plasma by high-performance liquid chromatography after derivatization with *trans*-4'-hydrazino-2-stilbazole 729(1999)43

Analysis of aplidine (dehydrodidemnin B), a new marine-derived depsipeptide, in rat biological fluids by liquid chromatography–tandem mass spectrometry 731(1999)335

Depsipeptides

Bioanalysis of thiocoraline, a new marine antitumoral depsipeptide, in plasma by high-performance liquid chromatography and fluorescence detection 726(1999)255

Dereplication

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

Dermatan sulfate

Ultrasensitive capillary electrophoresis of sulfated disaccharides in chondroitin/dermatan sulfates by laser-induced fluorescence after derivatization with 2-aminoacridone 730(1999)129

Desacetylcefotaxime

Simultaneous determination of cefotaxime and desacetylcefotaxime in human plasma and cerebrospinal fluid by high-performance liquid chromatography 750(2001)171

Deschloroethylcyclophosphamide

Analysis of cyclophosphamide and five metabolites from human plasma using liquid chromatography–mass spectrometry and gas chromatography–nitrogen–phosphorus detection 732(1999)287

Desglycylremacemide

Simultaneous determination of remacemide hydrochloride and desglycylremacemide (AR-R12495XX) in brain tissue by high-performance liquid chromatography 746(2000)325

Desmethyldoxepin

Stereoselective measurement of *E*- and *Z*-doxepin and its *N*-desmethyl and hydroxylated metabolites by gas chromatography–mass spectrometry 736(1999)201

Determination of doxepin and desmethyldoxepin in human plasma using liquid chromatography–tandem mass spectrometry 742(2000)91

Desmethyloanzapine

A sensitive high-performance liquid chromatographic method using electrochemical detection for the analysis of olanzapine and desmethyloanzapine in plasma of schizophrenic patients using a new solid-phase extraction procedure 750(2001)137

***N*-Desmethyltopotecan**

Quantification of topotecan and its metabolite *N*-desmethyltopotecan in human plasma, urine and faeces by high-performance liquid chromatographic methods 727(1999)191

***O*-Desmethyltramadol**

Enantiomeric determination of tramadol and its main metabolite *O*-desmethyltramadol in human plasma by liquid chromatography–tandem mass spectrometry 748(2000)65

Desmosterol

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Detergent–polymer aqueous two-phase system

Affinity partitioning of a poly(histidine)-tagged integral membrane protein, cytochrome *bo*₃ ubiquinol oxidase, in a detergent–polymer aqueous two-phase system containing metal-chelating polymer 743(2000)307

Detergents

Immunopurification of the blood group RhD protein from human erythrocyte membranes 735(1999)207

Dexamethasone

Simultaneous determination of hydrocortisone, dexamethasone, indomethacin, phenylbutazone and oxyphenbutazone in equine serum by high-performance liquid chromatography 738(2000)17

Dextran

Viscosity prediction of polyethylene glycol–dextran–water solutions used in aqueous two-phase systems 743(2000)181

High-performance size-exclusion chromatographic analysis of dextran–methylprednisolone hemisuccinate in rat plasma 744(2000)293

Diamine oxidase

Highly efficient purification of porcine diamine oxidase 737(2000)161

Diaminobutane

Dynamically-coated capillaries allow for capillary electrophoretic resolution of transferrin sialoforms via direct analysis of human serum 742(2000)79

5,4'-Diaminoflavone

Identification of the principal circulating metabolite of a synthetic 5,4'-diaminoflavone (NSC 686288), an antitumor agent, in the rat 741(2000)205

Diampron

Determination by capillary zone electrophoresis of berenil, phenamidine, diampron and dibromopropamidine in serum and urine 738(2000)293

Diarylpropenamine

Determination of the geometrical diarylpropenamine isomers in feces by high-performance liquid chromatography 728(1999)117

Diazepam

Evaluation of the programmed temperature vaporiser for large-volume injection of biological samples in gas chromatography 729(1999)199

Dibromopropamidine

Determination by capillary zone electrophoresis of berenil, phenamidine, diampron and dibromopropamidine in serum and urine 738(2000)293

Dibucaine

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

Dichloroacetic acid

Sensitive high-performance liquid chromatography method for the simultaneous determination of low levels of dichloroacetic acid and its metabolites in blood and urine 729(1999)271

Dichlorobenzene

Gas chromatographic–mass spectrometric analysis of dichlorobenzene isomers in human blood with headspace solid-phase microextraction 731(1999)217

2,2-Dichlorovinyl dimethylphosphate

Development, validation and application of assays to quantify metrifonate and 2,2-dichlorovinyl dimethylphosphate in human body fluids 741(2000)257

Diclofenac sodium

Study of the solid-phase extraction of diclofenac sodium, indomethacin and phenylbutazone for their analysis in human urine by liquid chromatography 729(1999)139

Didemnin

Analysis of apidine (dehydroididemnin B), a new marine-derived depsipeptide, in rat biological fluids by liquid chromatography–tandem mass spectrometry 731(1999)335

***N,N'*-Diethylene-*N''*-2-chloroethylphosphoramidate**

Simultaneous determination of thioTEPA, TEPA and a novel, recently identified thioTEPA metabolite, monochloroTEPA, in urine using capillary gas chromatography 742(2000)335

Diethylenetriaminepentaacetic acid

Simple and efficient method for the detection of diethylenetriaminepentaacetic acid 736(1999)299

1,3-Diethyl-2-thiobarbituric acid

High-performance liquid chromatographic determination of 1,3-diethyl-2-thiobarbituric acid–malonaldehyde adduct in fish meat 726(1999)313

***N,N*-Diethyl-*m*-toluamide**

Simultaneous determination of pyridostigmine bromide, *N,N*-diethyl-*m*-toluamide, permethrin, and their metabolites in rat plasma and urine by high-performance liquid chromatography 749(2000)171

Digoxin

Capillary electrophoretic immunoassays for digoxin and gentamicin with laser-induced fluorescence polarization detection 734(1999)31

High-performance liquid chromatography with on-line post-column immunoreaction detection of digoxin and its metabolites based on fluorescence energy transfer in the far-red spectral region 745(2000)305

Dihydroartemisinin

Simultaneous determination of artemether and its major metabolite dihydroartemisinin in plasma by gas chromatography–mass spectrometry–selected ion monitoring 731(1999)251

Dihydroergotamine

Determination of dihydroergotamine in human plasma by high-performance liquid chromatography with fluorescence detection 735(1999)151

Dihydroetorphine

Quantitative determination of dihydroetorphine in rat plasma and brain by liquid chromatography–tandem mass spectrometry 740(2000)253

Dihydroorotic acid

Simultaneous separation by high-performance liquid chromatography of carbamoyl aspartate, carbamoyl phosphate and dihydroorotic acid 732(1999)487

Dihydropyrimidinase

Radiochemical assay for determination of dihydropyrimidinase activity using reversed-phase high-performance liquid chromatography 729(1999)307

Dihydropyrimidine dehydrogenase

Measurement of plasma uracil using gas chromatography–mass spectrometry in normal individuals and in patients receiving inhibitors of dihydropyrimidine dehydrogenase 738(2000)249

Dihydroxybenzoate

Determination of hydroxyl free radical formation in human platelets using high-performance liquid chromatography with electrochemical detection 732(1999)213

3,4-Dihydroxyphenethylamine

Gas chromatographic–mass spectrometric procedures for determination of the catechol-*O*-methyltransferase (COMT) activity and for detection of unstable catecholic metabolites in human and rat liver preparations after COMT catalyzed *in situ* *nascendi* derivatization using *S*-adenosylmethionine 739(2000)325

3,4-Dihydroxyphenylacetic acid

Liquid chromatography–electrospray tandem mass spectrometry of acidic monoamine metabolites 744(2000)423

3,4-Dihydroxyphenylglycol

Sensitive and specific method for the simultaneous determination of natural and synthetic catecholamines and 3,4-dihydroxyphenylglycol in microdialysis samples 730(1999)213

4-Dimethylaminobenzylamine

4-Dimethylaminobenzylamine as a sensitive chemiluminescence derivatization reagent for 5-hydroxyindoles and its application to their quantification in human platelet-poor plasma 738(2000)199

Dimethylamphetamine

Simultaneous determination of dimethylamphetamine and its metabolites in rat hair by gas chromatography–mass spectrometry 741(2000)163

Dimethylamphetamine *N*-oxide

Simultaneous determination of dimethylamphetamine and its metabolites in rat hair by gas chromatography–mass spectrometry 741(2000)163

Dimethylarginine

Determination of asymmetrical dimethylarginine by capillary electrophoresis–laser-induced fluorescence 741(2000)77

***N,N*-Dimethylformamide**

Simultaneous determination of two human urinary metabolites of *N,N*-dimethylformamide using gas chromatography–thermionic sensitive detection with mass spectrometric confirmation 734(1999)285

Dimethyl sulfoxide

Quantification of residual dimethyl sulfoxide in supernatants of haematopoietic stem cells by capillary zone electrophoresis 728(1999)75

5,6-Dimethylxanthenone-4-acetic acid

Determination of two major metabolites of the novel anti-tumour agent 5,6-dimethylxanthenone-4-acetic acid in hepatic microsomal incubations by high-performance liquid chromatography with fluorescence detection 734(1999)129

2,4-Dinitrophenylhydrazine

Measurement of free and bound malondialdehyde in plasma by high-performance liquid chromatography as the 2,4-dinitrophenylhydrazine derivative 742(2000)315

2,3-Dinor-5,6-dihydro-8-isoprostaglandin F_{2α}

Tandem mass spectrometric quantification of 8-iso-prostaglandin F_{2α} and its metabolite 2,3-dinor-5,6-dihydro-8-iso-prostaglandin F_{2α} in human urine 744(2000)99

Diospyrin

Sensitive high-performance thin-layer chromatographic method for the estimation of diospyrin, a tumour inhibitory agent from the stem bark of *Diospyros montana* Roxb 744(2000)257

4,5-Dioxovaleric acid

Measurement of 4,5-dioxovaleric acid by high-performance liquid chromatography and fluorescence detection 729(1999)237

Dipalmitoylphosphatidylcholine

Quantitative determination of dipalmitoylphosphatidylcholine and palmitic acid in porcine lung surfactants used in the treatment of respiratory distress syndrome 744(2000)407

4-(4,5-Diphenyl-1*H*-imidazol-2-yl) benzoyl chloride

High-performance liquid chromatography with peroxyoxalate chemiluminescence detection of bisphenol A migrated from polycarbonate baby bottles using 4-(4,5-diphenyl-1*H*-imidazol-2-yl)benzoyl chloride as a label 749(2000)49

Disaccharides, sulfated

Ultrasensitive capillary electrophoresis of sulfated disaccharides in chondroitin/dermatan sulfates by laser-induced fluorescence after derivatization with 2-aminoacridone 730(1999)129

Disodium EDTA

Quantitative liquid chromatographic–tandem mass spectrometric determination of reserpine in FVB/N mouse plasma using a “chelating” agent (disodium EDTA) for releasing protein-bound analytes during 96-well liquid–liquid extraction 742(2000)369

Disopyramide

Enantioselective analysis of disopyramide and mono-*N*-dealkyldisopyramide in plasma and urine by high-performance liquid chromatography on an amylose-derived chiral stationary phase 744(2000)299

Diuretics

High-performance liquid chromatographic separation of a complex mixture of diuretics using a micellar mobile phase of sodium dodecyl sulphate. Application to human urine samples 748(2000)415

DMP 266

Quantitative determination of efavirenz (DMP 266), a novel non-nucleoside reverse transcriptase inhibitor, in human plasma using isocratic reversed-phase high-performance liquid chromatography with ultraviolet detection 734(1999)55

DNA

Optimization of intercalation dye concentration for short tandem repeat allele genotyping using capillary electrophoresis with laser-induced fluorescence detection 732(1999)365

Rapid method for relative gene expression determination in human tissues using automated capillary gel electrophoresis and multicolor detection 734(1999)47

Analysis of gamma radiation-induced damage to plasmid DNA using dynamic size-sieving capillary electrophoresis 741(2000)243

Direct quantification of AD-36 adenovirus DNA by capillary electrophoresis with laser-induced fluorescence 744(2000)1

Mass spectrometric detection and measurement of *N*²-(2'-deoxyguanosin-8-yl)PhIP adducts in DNA 744(2000)55

Approaches to functional genomics: potential of matrix-assisted laser desorption ionization–time of flight mass spectrometry combined with separation methods for the analysis of DNA in biological samples 745(2000)231

Comparison between capillary and nano liquid chromatography–electrospray mass spectrometry for the analysis of minor DNA–melfhalan adducts 748(2000)197

DNA adducts

Analysis of melfhalan adducts of 2'-deoxynucleotides in calf thymus DNA hydrolysates by capillary high-performance liquid chromatography–electrospray tandem mass spectrometry 736(1999)43

DNA fragments

Effect of temperature on the separation of DNA fragments by high-performance liquid chromatography and capillary electrophoresis: a comparative study 738(2000)243

Docetaxel

Liquid chromatographic analysis and preliminary pharmacokinetics of methotrexate in cancer patients co-treated with docetaxel 735(1999)111

Simple and sensitive high-performance liquid chromatography method for the determination of docetaxel in human plasma or urine 748(2000)383

Dolastatin-10

Improvement in assay sensitivity for plasma dolastatin-10 using capillary electrophoresis at elevated temperatures 729(1999)1

Domperidone

Determination of domperidone in human serum and human breast milk by high-performance liquid chromatography–electrospray mass spectrometry 730(1999)9

High-performance liquid chromatographic analysis for the determination of domperidone in human plasma 744(2000)207

Donepezil

Simultaneous determination of donepezil (aricept®) enantiomers in human plasma by liquid chromatography–electrospray tandem mass spectrometry 729(1999)147

L-DOPA

Determination of hydroxyl free radical formation in human platelets using high-performance liquid chromatography with electrochemical detection 732(1999)213

Dopamine

Simultaneous determination of catecholamines in rat brain tissue by high-performance liquid chromatography 731(1999)361

Determination of urine catecholamines by capillary electrophoresis with dual-electrode amperometric detection 750(2001)33

Dopamine D₄ receptor antagonist

High-performance liquid chromatographic analysis of the D₄ receptor antagonist SCH 66712 in rat plasma 735(1999)11

Doping control

Analysis of corticosteroids in hair by liquid chromatography–electrospray ionization mass spectrometry 740(2000)227

Doxepin

Stereoselective measurement of *E*- and *Z*-doxepin and its *N*-desmethyl and hydroxylated metabolites by gas chromatography–mass spectrometry 736(1999)201

Determination of doxepin and desmethyldoxepin in human plasma using liquid chromatography–tandem mass spectrometry 742(2000)91

Doxorubicin

Simultaneous determination of four anthracyclines and three metabolites in human serum by liquid chromatography–electrospray mass spectrometry 738(2000)281

Doxycycline

Determination of flumequine and doxycycline in milk by a simple thin-layer chromatographic method 734(1999)7

Simple and reliable method of doxycycline determination in human plasma and biological tissues 744(2000)359

Drug metabolites

Mass directed peak selection, an efficient method of drug metabolite identification using directly coupled liquid chromatography–mass spectrometry–nuclear magnetic resonance spectroscopy 748(2000)281

Drugs

High-performance affinity chromatography and immobilized serum albumin as probes for drug- and hormone-protein binding 739(2000)39

High-temperature solid-phase microextraction procedure for the detection of drugs by gas chromatography–mass spectrometry 745(2000)399

Factors to consider in the development of generic bioanalytical high-performance liquid chromatographic–mass spectrometric methods to support drug discovery 748(2000)21

Drugs of abuse

Hair analysis for abused and therapeutic drugs 733(1999)161

Liquid chromatography–mass spectrometry as a routine method in forensic sciences: a proof of maturity 748(2000)3

DY-9760e

Determination of a novel calmodulin antagonist, 3-[2-[4-(3-chloro-2-methylphenyl)-1-piperazinyl]ethyl]-5,6-dimethoxy-1-(4-imidazolylmethyl)-1*H*-indazole dihydrochloride 3.5 hydrate, DY-9760e, in human plasma using solid-phase extraction and high-performance liquid chromatography with fluorescence detection 734(1999)39

DY-9836

Determination of a novel calmodulin antagonist, 3-[2-[4-(3-chloro-2-methylphenyl)-1-piperazinyl]ethyl]-5,6-dimethoxy-1-(4-imidazolylmethyl)-1*H*-indazole dihydrochloride 3.5 hydrate, DY-9760e, in human plasma using solid-phase extraction and high-performance liquid chromatography with fluorescence detection 734(1999)39

E

E5564

Quantitative determination of a potent lipopolysaccharide antagonist, E5564, in rat and dog plasma by high-performance liquid chromatography with fluorescence detection 736(1999)67

EB 1089

Determination of the vitamin D analog EB 1089 (seocalcitol) in human and pig serum using liquid chromatography–tandem mass spectrometry 740(2000)117

Ecgonine methyl ester

Performance of a pentafluorophenylpropyl stationary phase for the electrospray ionization high-performance liquid chromatography–mass spectrometry–mass spectrometry assay of cocaine and its metabolite ecgonine methyl ester in human urine 748(2000)77

Echinofuran

Simultaneous analysis of shikimate-derived secondary metabolites in *Lithospermum erythrorhizon* cell suspension cultures by high-performance liquid chromatography 738(2000)3

Ecstasy

Determination of “Ecstasy” components in alternative biological specimens 733(1999)137

Ecstasy metabolite

On-line coupling of automated solid-phase extraction with high-performance liquid chromatography and electrochemical detection. Quantitation of oxidizable drugs of abuse and their metabolites in plasma and urine 731(1999)167

EDDP

Determination of methadone and its metabolites EDDP and EMDP in human hair by headspace solid-phase microextraction and gas chromatography–mass spectrometry 746(2000)255

EDTA

Rapid identification of metallothionein isoforms in liver cytosol fraction by capillary zone electrophoresis using EDTA 742(2000)193

Efavirenz

Quantitative determination of efavirenz (DMP 266), a novel non-nucleoside reverse transcriptase inhibitor, in human plasma using isocratic reversed-phase high-performance liquid chromatography with ultraviolet detection 734(1999)55

Simultaneous determination of the HIV protease inhibitors indinavir, amprenavir, saquinavir, ritonavir, nelfinavir and the non-nucleoside reverse transcriptase inhibitor efavirenz by high-performance liquid chromatography after solid-phase extraction 740(2000)43

Simultaneous high-performance liquid chromatographic determination of the antiretroviral agents amprenavir, nelfinavir, ritonavir, saquinavir, delavirdine and efavirenz in human plasma 742(2000)453

Efletirizine

Quantitation of efletirizine in human plasma and urine using automated solid-phase extraction and column-switching high-performance liquid chromatography 730(1999)239

Electrochemical detection

Measurement of chloramphenicol by capillary zone electrophoresis following end-column amperometric detection at a carbon fiber micro-disk array electrode 741(2000)155

EM574

Application of liquid chromatography–turbo ion spray tandem mass spectrometry for quantitative analysis of a potent motilin receptor agonist, EM574, and its metabolites in human plasma 734(1999)101

EMDP

Determination of methadone and its metabolites EDDP and EMDP in human hair by headspace solid-phase microextraction and gas chromatography–mass spectrometry 746(2000)255

Enaminone

Determination of the enaminone DM5, an anti-epileptic agent, in mouse plasma and brain tissue by high-performance liquid chromatography with ultraviolet detection 749(2000)191

β-Endorphin

Mass spectrometric analysis of neuropeptidergic systems in the human pituitary and cerebrospinal fluid (Review) 731(1999)3

Eniluracil

Measurement of plasma uracil using gas chromatography–mass spectrometry in normal individuals and in patients receiving inhibitors of dihydropyrimidine dehydrogenase 738(2000)249

Enrofloxacin

Simultaneous determination of enrofloxacin and ciprofloxacin in animal biological fluids by high-performance liquid chromatography. Application in pharmacokinetic studies in pig and rabbit 726(1999)175

Entacapone

Direct analysis of nitrocatechol-type glucuronides in urine by capillary electrophoresis–electrospray ionisation mass spectrometry and tandem mass spectrometry 749(2000)253

Enzymes

Purification of cytochrome P-450 from adult pig testis by hydroxylapatite and deoxycorticosterone affinity column chromatography 726(1999)291

Thiopurine methyl transferase activity: new extraction conditions for high-performance liquid chromatographic assay 727(1999)235

High-performance liquid chromatographic–fluorimetric assay for cathepsin A (lysosomal protective protein) activity 728(1999)59

Radiochemical assay for determination of dihydropyrimidinase activity using reversed-phase high-performance liquid chromatography 729(1999)307

Determination of differential activities of soluble and membrane-bound catechol-*O*-methyltransferase in tissues and erythrocytes 729(1999)347

Rapid, single-step procedure for the identification of transglutaminase-mediated isopeptide crosslinks in amino acid digests 732(1999)65

Affinity chromatography of plasma proteins (guanidinobenzoate): use of mimetic matrices and mimetic soluble ligands to prevent the binding of albumin on target affinity matrices 732(1999)165

Novel sensitive high-performance liquid chromatographic method for assay of coumarin 7-hydroxylation 734(1999)23

Metabolism of anabolic steroids by recombinant human cytochrome P450 enzymes. Gas chromatographic–mass spectrometric determination of metabolites 735(1999)73

Radiochemical high-performance liquid chromatographic assay for the determination of catechol *O*-methyltransferase activity towards various substrates 736(1999)143

Improved purification of the membrane-bound hydrogenase–sulfur-reductase complex from thermophilic archaea using ϵ -aminocaproic acid-containing chromatography buffers 737(2000)151

Highly efficient purification of porcine diamine oxidase 737(2000)161

Development of a downstream process for the isolation of *Staphylococcus aureus* arsenate reductase overproduced in *Escherichia coli* 737(2000)167

Purification of recombinant hydantoinase and L-*N*-carbamoylase from *Arthrobacter aurescens* expressed in *Escherichia coli*: comparison of wild-type and genetically modified proteins 737(2000)179

Purification of a D-hydantoinase using a laboratory-scale Streamline phenyl column as the initial step 737(2000)187

D-Xylose metabolism by *Candida intermedia*: isolation and characterisation of two forms of aldose reductase with different coenzyme specificities 737(2000)195

Purification of surface-associated urease from *Helicobacter pylori* 737(2000)202

Refolding and purification of a urokinase plasminogen activator fragment by chromatography 737(2000)225

Purification of human adenosine deaminase for the preparation of a reference material 737(2000)237

Pullulanase from the hyperthermophilic bacterium *Thermotoga maritima*: purification by β -cyclodextrin affinity chromatography 737(2000)245

Single-step purification of a recombinant thermostable α -amylase after solubilization of the enzyme from insoluble aggregates 737(2000)253

Purification of the fengycin synthetase multienzyme system from *Bacillus subtilis* b213 737(2000)267

New approach for separating *Bacillus subtilis* metalloprotease and α -amylase by affinity chromatography and for purifying neutral protease by hydrophobic chromatography 737(2000)277

Purification and characterization of an aminopeptidase from the chloroplast stroma of barley leaves by chromatographic and electrophoretic methods 737(2000)285

Three-step chromatographic purification of Cpr6, a cyclophilin from *Saccharomyces cerevisiae* 737(2000)295

Preparation, evaluation and application of new pseudo-affinity chromatographic supports for penicillin acylase purification 739(2000)63

Gas chromatographic–mass spectrometric procedures for determination of the catechol-*O*-methyltransferase (COMT) activity and for detection of unstable catecholic metabolites in human and rat liver preparations after COMT catalyzed *in statu nascendi* derivatization using *S*-adenosylmethionine 739(2000)325

Normal values and age-dependent changes in GTP cyclohydrolase I activity in stimulated mononuclear blood cells measured by high-performance liquid chromatography 740(2000)35

Arsanilic acid–Sepharose chromatography of pyruvate kinase from KB cells 740(2000)109

Essential role of the concentration of immobilized ligands in affinity chromatography: Purification of guanidinobenzoate on an ionized ligand 740(2000)211

Differential diagnosis of homocystinuria by urease treatment, isotope dilution and gas chromatography–mass spectrometry 742(2000)59

Affinity chromatography for purification of two urokinases from human urine 742(2000)71

Assessment of nitric oxide synthase activity in vitro and in vivo by gas chromatography–mass spectrometry 742(2000)143

Metabolic profiling of valproic acid by cDNA-expressed human cytochrome P450 enzymes using negative-ion chemical ionization gas chromatography–mass spectrometry 742(2000)217

Determination of the dissociation constant of valine from acetohydroxy acid synthase by equilibrium partition in an aqueous two-phase system 743(2000)225

Bioaffinity extraction of glucoamylase in aqueous two-phase systems using starch as free bioligand 743(2000)241

Study on heterogeneity of β -glucosidase from *Aspergillus* species by using counter-current distribution 743(2000)247

Purification of acetohydroxy acid synthase by separation in an aqueous two-phase system 743(2000)281

Purification of soybean peroxidase (*Glycine max*) by metal affinity partitioning in aqueous two-phase systems 743(2000)287

Applications of aqueous two-phase partition to isolation of membranes from plants: A periodic NADH oxidase activity as a marker for right side-out plasma membrane vesicles 743(2000)369

Assay of ATPase and Na,K-ATPase activity using high-performance liquid chromatographic determination of ADP derived from ATP 744(2000)19

Rapid, simplified and sensitive method for screening fructose-1,6-diphosphatase deficiency by analyzing urinary metabolites in urease/direct preparations and gas chromatography–mass spectrometry in the selected-ion monitoring mode 746(2000)75

Differential diagnosis of homocystinuria by urease treatment, isotope dilution and gas chromatography–mass spectrometry 746(2000)103

Inhibition of acrosine-like protease activity by a lectin affinity chromatographic bovine seminal plasma fraction containing the PDC-109 and aSFP proteins 746(2000)141

Hydrophobic character of surface regions and total hydrophobicity of four variants of chromosomal class C β -lactamase from *Pseudomonas aeruginosa* are identical. Chromatographic comparison of the hydrophobic character of the variants and the effect of focusing buffer composition on the separation of the variants by chromatofocusing with internal and external pH gradients 746(2000)161

Reversed-phase high-performance liquid chromatography of nicotinic acid mononucleotide for measurement of quinolinate phosphoribosyltransferase 749(2000)281

Ephedrine

Sensitive determination of ephedrine and norephedrine in human plasma samples using derivatization with 9-fluorenylmethyl chloroformate and liquid chromatography 744(2000)25

Simultaneous chiral analysis of methamphetamine and related compounds by capillary electrophoresis 749(2000)111

Epicatechin

Analysis of (+)-catechin, (–)-epicatechin and their 3'- and 4'-O-methylated analogs. A comparison of sensitive methods 726(1999)277

Epinephrine

Determination of urine catecholamines by capillary electrophoresis with dual-electrode amperometric detection 750(2001)33

Epirubicin

Simultaneous determination of four anthracyclines and three metabolites in human serum by liquid chromatography–electrospray mass spectrometry 738(2000)281

Ergovaline

Analysis of ergovaline in milk using high-performance liquid chromatography with fluorimetric detection 729(1999)255

Erythromycin

Determination of erythromycin, clarithromycin, roxithromycin, and azithromycin in plasma by high-performance liquid chromatography with amperometric detection 738(2000)405

Erythromycin derivative

Application of liquid chromatography–turbo ion spray tandem mass spectrometry for quantitative analysis of a potent motilin receptor agonist, EM574, and its metabolites in human plasma 734(1999)101

ESI

Factors to consider in the development of generic bioanalytical high-performance liquid chromatographic–mass spectrometric methods to support drug discovery 748(2000)21

Estradiol

Determination of estramustine phosphate and its metabolites estromustine, estramustine, estrone and estradiol in human plasma by liquid chromatography with fluorescence detection and gas chromatography with nitrogen-phosphorus and mass spectrometric detection 738(2000)267

17 β -Estradiol

High-performance liquid chromatographic determination of 17 β -estradiol and 17 β -estradiol-3-acetate solubilities and diffusion coefficients in silicone elastomeric intravaginal rings 744(2000)157

17 β -Estradiol-3-acetate

High-performance liquid chromatographic determination of 17 β -estradiol and 17 β -estradiol-3-acetate solubilities and diffusion coefficients in silicone elastomeric intravaginal rings 744(2000)157

Estramustine

Determination of estramustine phosphate and its metabolites estromustine, estramustine, estrone and estradiol in human plasma by liquid chromatography with fluorescence detection and gas chromatography with nitrogen-phosphorus and mass spectrometric detection 738(2000)267

Estramustine phosphate

Determination of estramustine phosphate and its metabolites estromustine, estramustine, estrone and estradiol in human plasma by liquid chromatography with fluorescence detection and gas chromatography with nitrogen-phosphorus and mass spectrometric detection 738(2000)267

Estrogens

Charged cyclodextrin-mediated sample stacking in micellar capillary electrophoresis. A simple method for enhancing the detection sensitivity of hydrophobic compounds 731(1999)369

Estromustine

Determination of estramustine phosphate and its metabolites estromustine, estramustine, estrone and estradiol in human plasma by liquid chromatography with fluorescence detection and gas chromatography with nitrogen-phosphorus and mass spectrometric detection 738(2000)267

Estrone

Determination of estramustine phosphate and its metabolites estromustine, estramustine, estrone and estradiol in human plasma by liquid chromatography with fluorescence detection and gas chromatography with nitrogen-phosphorus and mass spectrometric detection 738(2000)267

Ethanol

Determination of ethyl glucuronide, a minor metabolite of ethanol, in human serum by liquid chromatography–electrospray ionization mass spectrometry 726(1999)105

Ultra-sensitive method for determination of ethanol in whole blood by headspace capillary gas chromatography with cryogenic oven trapping 727(1999)89

Ethosuximide

Trace analysis of ethosuximide in human plasma with a chemically removable derivatizing reagent and high-performance liquid chromatography 729(1999)111

Ethoxyresorufin

Determination of cytochrome P450 1A activities in mammalian liver microsomes by high-performance liquid chromatography with fluorescence detection 744(2000)399

Ethyl alcohol

Liquid chromatographic determination of ethyl alcohol in body fluids 729(1999)103

Ethylene oxide–propylene oxide

Polymer recycling in aqueous two-phase extractions using thermoseparating ethylene oxide–propylene oxide copolymers 743(2000)115

17 α -Ethyl-5 β -estrane-3 α ,17 β -diol

17 α -Ethyl-5 β -estrane-3 α ,17 β -diol, a biological marker for the abuse of norethandrolone and ethylestrenol in slaughter cattle 728(1999)217

Ethylestrenol

17 α -Ethyl-5 β -estrane-3 α ,17 β -diol, a biological marker for the abuse of norethandrolone and ethylestrenol in slaughter cattle 728(1999)217

Ethyl glucuronide

Determination of ethyl glucuronide, a minor metabolite of ethanol, in human serum by liquid chromatography–electrospray ionization mass spectrometry 726(1999)105

2-Ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine

Stereoselective quantification of methadone and its major oxidative metabolite, 2-ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine, in human urine using high-performance liquid chromatography 744(2000)165

***R*-(α)-ethyl-2-oxo-pyrrolidine acetamide**

Enantioselective analysis of levetiracetam and its enantiomer *R*- α -ethyl-2-oxo-pyrrolidine acetamide using gas chromatography and ion trap mass spectrometric detection 745(2000)325

Etodolac

Capillary electrochromatography–electrospray ionization mass spectrometry for the qualitative investigation of the drug etodolac and its metabolites in biological samples 748(2000)213

Etoposide

Simultaneous quantitation of etoposide and its catechol metabolite in human plasma using high-performance liquid chromatography with electrochemical detection 728(1999)241

Highly sensitive liquid chromatography–electrospray mass spectrometry (LC–MS) method for the determination of etoposide levels in human serum and plasma 744(2000)91

Etoposide catechol

Simultaneous quantitation of etoposide and its catechol metabolite in human plasma using high-performance liquid chromatography with electrochemical detection 728(1999)241

Everninomycin

Simple and sensitive high-performance liquid chromatographic method for the determination of an everninomycin, SCH 27899, in rat plasma 730(1999)55

EXP-3174

Method for the simultaneous determination of losartan and its major metabolite, EXP-3174, in human plasma by liquid chromatography–electrospray ionization tandem mass spectrometry 734(1999)325

Ezlopitant

Assay of ezlopitant, a substance P receptor antagonist, and metabolites in biological matrices by gas chromatography with mass spectrometric detection: simultaneous analysis of a benzyl alcohol and alkene 744(2000)333

Narrow-bore high-performance liquid chromatography in combination with ionspray tandem mass spectrometry for the determination of the substance P receptor antagonist ezlopitant and its two active metabolites in plasma 748(2000)179

F**Famotidine**

Determination of famotidine in human plasma and urine by high-performance liquid chromatography 732(1999)239

Fatty acid methyl esters

Purification of fatty acid methyl esters by high-performance liquid chromatography 732(1999)495

Fatty acids

Application of high-performance liquid chromatography of plasma fatty acids as their phenacyl esters to evaluate splanchnic and renal fatty acid balance in vivo 727(1999)15

Screening for fatty acid beta oxidation disorders. Acylglycine analysis by electron impact ionization gas chromatography–mass spectrometry 731(1999)105

Routine analysis of short-chain fatty acids for anaerobic bacteria identification using capillary electrophoresis and indirect ultraviolet detection 741(2000)89

Identification of yeast species by fatty acid profiling as measured by gas-liquid chromatography 742(2000)13

Gas chromatographic–mass spectrometric determination of plasma saturated fatty acids using pentafluorophenyldimethylsilyl derivatization 742(2000)37

Fenfluramine

Stereoselective determination of fenfluramine enantiomers in rat liver microsomal incubates 727(1999)107

Fengycin synthetase

Purification of the fengycin synthetase multienzyme system from *Bacillus subtilis* b213 737(2000)267

Fenofibric acid

Determination of fenofibric acid in human plasma using automated solid-phase extraction coupled to liquid chromatography 742(2000)391

Fenoprofen

Determination of fenoprofen in serum by capillary isotachopheresis 729(1999)11

Fenproporex

Studies on the metabolism and toxicological detection of the amphetamine-like anorectic fenproporex in human urine by gas chromatography–mass spectrometry and fluorescence polarization immunoassay 738(2000)107

Finasteride

Determination of finasteride in human plasma by liquid–liquid extraction and high-performance liquid chromatography 738(2000)305

FK506 binding proteins

Three-step purification of a fragment of the large immunophilin FKBP52 737(2000)71

Flavin nucleotides

Analysis of riboflavin and riboflavin cofactor levels in plasma by high-performance liquid chromatography 739(2000)219

Flavones

Fast liquid chromatographic–mass spectrometric determination of pharmaceutical compounds 732(1999)445

Flecainide

Determination of concentrations of flecainide in human serum by high-performance liquid chromatography on a fluorocarbon-bonded silica gel column 726(1999)219

Therapeutic drug monitoring of flecainide in serum using high-performance liquid chromatography and electrospray mass spectrometry 746(2000)249

Flophemesyl

Gas chromatographic–mass spectrometric determination of plasma saturated fatty acids using pentafluorophenyldimethylsilyl derivatization 742(2000)37

Fludarabine

High-performance liquid chromatography method for the determination and quantitation of arabinosylguanine triphosphate and fludarabine triphosphate in human cells 745(2000)421

Flufenamic acid

Simultaneous analysis of anthranilic acid derivatives in pharmaceuticals and human urine by high-performance liquid chromatography with isocratic elution 744(2000)81

Flumequine

Determination of flumequine and doxycycline in milk by a simple thin-layer chromatographic method 734(1999)7

Flunarizine

Determination of flunarizine in rat brain by liquid chromatography–electrospray mass spectrometry 736(1999)167

Flunitrazepam

Validated method for the therapeutic drug monitoring of flunitrazepam in human serum using liquid chromatography–atmospheric pressure chemical ionization tandem mass spectrometry with an ion trap detector 738(2000)437

9-Fluorenylmethyl chloroformate

Convenient method of threonine, methionine and their related amino compounds by high-performance liquid chromatography and its application to rumen fluid 741(2000)279

Sensitive determination of ephedrine and norephedrine in human plasma samples using derivatization with 9-fluorenylmethyl chloroformate and liquid chromatography 744(2000)25

9-fluorenylmethyloxycarbonyl chloride

High-performance liquid chromatographic assay for catecholamines and metanephrines using fluorimetric detection with pre-column 9-fluorenylmethyloxycarbonyl chloride derivatization 749(2000)179

Fluoresceine isothiocyanate

Determination of asymmetrical dimethylarginine by capillary electrophoresis–laser-induced fluorescence 741(2000)77

5-Fluorodihydrouracil

Automated determination of 5-fluorouracil and its metabolite in urine by high-performance liquid chromatography with column switching 731(1999)267

5-Fluorouracil

Automated determination of 5-fluorouracil and its metabolite in urine by high-performance liquid chromatography with column switching 731(1999)267

Determination of 5-fluorouracil in microvolumes of human plasma by solvent extraction and high-performance liquid chromatography 735(1999)293

Simple and sensitive determination of 5-fluorouracil in plasma by high-performance liquid chromatography. Application to clinical pharmacokinetic studies 736(1999)97

Determination of 5-fluorouracil in environmental samples by solid-phase extraction and high-performance liquid chromatography with ultraviolet detection 750(2001)25

Fluorescent immunoassay

Immunoassay-based determination of phenobarbital using size-exclusion chromatography 749(2000)215

Fluoxetine

Simultaneous determination of citalopram, fluoxetine, paroxetine and their metabolites in plasma and whole blood by high-performance liquid chromatography with ultraviolet and fluorescence detection 734(1999)229

Fluprostenol

Use of short high-performance liquid chromatography columns and tandem-mass spectrometry for the rapid analysis of a prostaglandin analog, fluprostenol, in rat plasma 741(2000)213

Semi-automated 96-well solid-phase extraction and gas chromatography–negative chemical ionization tandem mass spectrometry for the trace analysis of fluprostenol in rat plasma 744(2000)283

Flurbiprofen

Sensitive and specific high-performance liquid chromatographic assay for 4'-hydroxyflurbiprofen and flurbiprofen in human urine and plasma 749(2000)119

Formic acid

Analysis of methanol or formic acid in body fluids by headspace solid-phase microextraction and capillary gas chromatography 734(1999)155

Formycin A

High-performance liquid chromatographic assay of formycin A in plasma after solid-phase extraction 731(1999)383

Fosinopril

Liquid chromatographic–electrospray tandem mass spectrometric method for the simultaneous quantitation of the prodrug fosinopril and the active drug fosinoprilat in human serum 739(2000)255

Fosinoprilat

Liquid chromatographic–electrospray tandem mass spectrometric method for the simultaneous quantitation of the prodrug fosinopril and the active drug fosinoprilat in human serum 739(2000)255

Frit FAB

Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments. Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry 748(2000)369

Fructose

Erratum to “Determination of adenosine and deoxyadenosine in urine by high-performance liquid chromatography with column switching”. [J. Chromatogr. B, 719 (1998) 55–61] 726(1999)317

Fructose-1,6-diphosphatase

Rapid, simplified and sensitive method for screening fructose-1,6-diphosphatase deficiency by analyzing urinary metabolites in urease/direct preparations and gas chromatography–mass spectrometry in the selected-ion monitoring mode 746(2000)75

Furosemide

Simplified method to quantify furosemide in urine by high-performance liquid chromatography and ultraviolet detection 730(1999)49

5-Furoylquinoline-3-carboxaldehyde

Reaction rate, activation energy, and detection limit for the reaction of 5-furoylquinoline-3-carboxaldehyde with neurotransmitters in artificial cerebrospinal fluid 741(2000)85

Fusion protein F

Purification of respiratory syncytial virus F and G proteins 737(2000)97

G

Gabapentin

Rapid high-performance liquid chromatographic determination of serum gabapentin 727(1999)119

Automated microanalysis of gabapentin in human serum by high-performance liquid chromatography with fluorometric detection 727(1999)125

Fast isocratic high-performance liquid chromatographic assay method for the simultaneous determination of gabapentin and vigabatrin in human serum 746(2000)311

Gacyclidine

Modified gas chromatographic–mass spectrometric assay for the determination of gacyclidine enantiomers in human plasma 731(1999)317

Galactose

Urine screening of five-day-old newborns: metabolic profiling of neonatal galactosuria 732(1999)469

Galanthamine

Enantiomeric resolution of galanthamine and related drugs used in anti-Alzheimer therapy by means of capillary zone electrophoresis employing derivatized cyclodextrin selectors 730(1999)167

GC–MS

Simultaneous determination of dimethylamphetamine and its metabolites in rat hair by gas chromatography–mass spectrometry 741(2000)163

Gas chromatographic–mass spectrometric determination of plasma saturated fatty acids using pentafluorophenyldimethylsilyl derivatization 742(2000)37

Inborn errors of metabolism discovered in Asian department of pediatrics and mental retardation research center 746(2000)41

Gemfibrozil

Gemfibrozil and its oxidative metabolites: quantification of aglycones, acyl glucuronides, and covalent adducts in samples from preclinical and clinical kinetic studies 741(2000)129

Gemifloxacin

Rapid determination of gemifloxacin in human plasma by high-performance liquid chromatography–tandem mass spectrometry 746(2000)191

GEM 231

Detection of a mixed-backbone oligonucleotide (GEM 231) in liver and tumor tissues by capillary electrophoresis 750(2001)129

Gentamicin

Capillary electrophoretic immunoassays for digoxin and gentamicin with laser-induced fluorescence polarization detection 734(1999)31

Gestrinone

Determination of gestrinone in human serum by liquid chromatography–electrospray tandem mass spectrometry 746(2000)151

GFAT

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

Ginkgolic acids

Determination of trace amounts of ginkgolic acids in *Ginkgo biloba* L. leaf extracts and phytopharmaceuticals by liquid chromatography–electrospray mass spectrometry 744(2000)249

Ginsenosides

Determination of ginsenoside Rg3 in plasma by solid-phase extraction and high-performance liquid chromatography for pharmacokinetic study 731(1999)403

Glucagon

Rapid simultaneous determination of glucagon and insulin by capillary electrophoresis immunoassays 742(2000)353

Glucoamylase

Bioaffinity extraction of glucoamylase in aqueous two-phase systems using starch as free bioligand 743(2000)241

Glucose

Chromatography on cells: analyses of solute interactions with the glucose transporter Glut1 in human red cells adsorbed on lectin-gel beads 739(2000)55

 β -Glucosidase

Study on heterogeneity of β -glucosidase from *Aspergillus* species by using counter-current distribution 743(2000)247

 β -Glucuronidase

High-performance liquid chromatographic determination of quercetin and isorhamnetin in rat tissues using β -glucuronidase and acid hydrolysis 738(2000)413

Glucuronides

Direct analysis of nitrocatechol-type glucuronides in urine by capillary electrophoresis–electrospray ionisation mass spectrometry and tandem mass spectrometry 749(2000)253

Glut1

Chromatography on cells: analyses of solute interactions with the glucose transporter Glut1 in human red cells adsorbed on lectin-gel beads 739(2000)55

Glutamate

On-line derivatization for continuous and automatic monitoring of brain extracellular glutamate levels in anesthetized rats: a microdialysis study 734(1999)1

Glutamic acid

Simultaneous determination of γ -aminobutyric acid and glutamic acid in the brain of 3-mercaptopropionic acid-treated rats using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 726(1999)285

Glutamine

On-line derivatization for continuous and automatic monitoring of brain extracellular glutamate levels in anesthetized rats: a microdialysis study 734(1999)1

Glutathione

Analysis of diagnostic metabolites by capillary electrophoresis–mass spectrometry 727(1999)43

Gas chromatographic–mass spectrometric analysis of stable isotopes of cysteine and glutathione in biological samples 732(1999)127

Glutathione S-transferase A1-1

Determination of tetrahydrothiophene formation as a probe of in vitro busulfan metabolism by human glutathione S-transferase A1-1: use of a highly sensitive gas chromatographic–mass spectrometric method 730(1999)25

Glutathionyl hemoglobin

Determination of glutathionyl hemoglobin in hemodialysis patients using electrospray ionization liquid chromatography–mass spectrometry 731(1999)121

Glutathionyl hemoglobin

Analysis of glutathionyl hemoglobin levels in diabetic patients by electrospray ionization liquid chromatography–mass spectrometry: effect of vitamin E administration 746(2000)91

Glycans

Carbohydrate release from picomole quantities of glycoprotein and characterisation of glycans by high-performance liquid chromatography and mass spectrometry 734(1999)169

Glycerol

Determination of stable isotopic enrichment and concentration of glycerol in plasma via gas chromatography–mass spectrometry for the estimation of lipolysis in vivo 744(2000)47

Glycerophospholipids

Oxidation of glycerophospholipids from biological membranes by reactive oxygen species: liquid chromatographic–mass spectrometric analysis of eicosanoid products 731(1999)59

Glycine

On-line derivatization for continuous and automatic monitoring of brain extracellular glutamate levels in anesthetized rats: a microdialysis study 734(1999)1

Glycine conjugates

Metabolism of an ionic contrast medium and the related agents 746(2000)25

Glycinexylidide

High-performance liquid chromatography–tandem electrospray mass spectrometry for the determination of lidocaine and its metabolites in human plasma and urine 741(2000)175

Glycolic acid

Sensitive high-performance liquid chromatography method for the simultaneous determination of low levels of dichloroacetic acid and its metabolites in blood and urine 729(1999)271

N-Glycolylneuraminic acid

Importance of high-performance liquid chromatographic analysis of serum N-acylneuraminic acids in evaluating surgical treatment in patients with early endometrial cancer 732(1999)375

Glycophorins

New procedures for glycophorin A purification with high yield and high purity 750(2001)109

P-Glycoprotein

Development of an immobilized P-glycoprotein stationary phase for on-line liquid chromatographic determination of drug-binding affinities 739(2000)33

Glycoproteins

Purification and quantitation of tumor necrosis factor receptor immunoadhesin using a combination of immunoaffinity and reversed-phase chromatography 728(1999)21

Discrimination of granulocyte colony-stimulating factor isoforms by high-performance capillary electrophoresis 732(1999)81

Carbohydrate release from picomole quantities of glycoprotein and characterisation of glycans by high-performance liquid chromatography and mass spectrometry 734(1999)169

Expression, purification and biological properties of the carboxyl half part of the HTLV-I surface envelope glycoprotein 737(2000)85

Purification of respiratory syncytial virus F and G proteins 737(2000)97

Synthesis and evaluation of affinity adsorbents for glycoproteins: an artificial lectin 746(2000)265

18 β -Glycyrrhetic acid

Determination of urinary 18 β -glycyrrhetic acid by gas chromatography and its clinical application in man 731(1999)323

Glycyrrhizin

Determination of 18 α -glycyrrhizin and 18 β -glycyrrhizin in dog plasma by high-performance liquid chromatography 738(2000)165

Glyoxylic acid

Sensitive high-performance liquid chromatography method for the simultaneous determination of low levels of dichloroacetic acid and its metabolites in blood and urine 729(1999)271

Gonadotropin-releasing hormone

Determination of immunoreactive gonadotropin-releasing hormone in serum and urine by on-line immunoaffinity capillary electrophoresis coupled to mass spectrometry 749(2000)197

Granulocyte colony-stimulating factor

Discrimination of granulocyte colony-stimulating factor isoforms by high-performance capillary electrophoresis 732(1999)81

Grepafloxacin

Determination of grepafloxacin in plasma and urine by a simple and rapid high-performance liquid chromatographic method 741(2000)295

Growth hormones

Quantitative analysis of clenbuterol in meat products using liquid chromatography–electrospray ionisation tandem mass spectrometry 736(1999)209

GTP cyclohydrolase I

Normal values and age-dependent changes in GTP cyclohydrolase I activity in stimulated mononuclear blood cells measured by high-performance liquid chromatography 740(2000)35

Guaiaicol estrogens

Studies on neurosteroids XII. Determination of enzymatically formed catechol estrogens and guaiaicol estrogens by rat brains using liquid chromatography–mass spectrometry–mass spectrometry 748(2000)89

Guan-Fu Base A

Determination of Guan-Fu Base A, a new anti-arrhythmic, in human plasma by gas chromatography and electron–capture detection 740(2000)273

Guanidinobenzoate

Affinity chromatography of plasma proteins (guanidinobenzoate): use of mimetic matrices and mimetic soluble ligands to prevent the binding of albumin on target affinity matrices 732(1999)165

Essential role of the concentration of immobilized ligands in affinity chromatography: Purification of guanidinobenzoate on an ionized ligand 740(2000)211

H

Haemoglobin adducts

Haemoglobin adducts of aromatic amines: diamines and polyaromatic amines 744(2000)337

Hair analysis

Analysis of corticosteroids in hair by liquid chromatography–electrospray ionization mass spectrometry 740(2000)227

Simultaneous determination of dimethylamphetamine and its metabolites in rat hair by gas chromatography–mass spectrometry 741(2000)163

Haloperidol

Determination of 1-methyl-1,2,3,4-tetrahydroisoquinoline in mouse brain after treatment with haloperidol by gas chromatography–selected ion monitoring 731(1999)53

Gas chromatographic–mass spectrometric analysis of veterinary tranquilizers in urine: evaluation of method performance 734(1999)113

Analysis of citrulline in rat brain tissue after perfusion with haloperidol by liquid chromatography–mass spectrometry 746(2000)33

Halothane

Solid-phase microextraction gas chromatographic–mass spectrometric method for the determination of inhalation anesthetics in urine 732(1999)115

Haptoglobin

Simple high-performance liquid chromatographic purification procedure for porcine plasma haptoglobin 731(1999)395

Hemoglobin β -chain

Determination of ionization efficiency of glycosylated and non-glycosylated peptides from the N-terminal of hemoglobin β -chain by electrospray ionization mass spectrometry 746(2000)83

Hemoglobins

Establishment of a combined strategy of genetic and mass spectrometric analyses for characterizing hemoglobin mutations. An example of Hb Hoshida (β 43Glu \rightarrow Gln) 731(1999)125

Hemolysin

Two-step fast protein liquid chromatographic purification of the *Serratia marcescens* hemolysin and peptide mapping with mass spectrometry 737(2000)13

Hepatitis B surface antigen

Aggregation of recombinant hepatitis B surface antigen induced in vitro by oxidative stress 736(1999)153

Heptafluorobutyric anhydride

Gas chromatographic method for the determination of toluidines in spiked urine samples 738(2000)427

Determination of paroxetine levels in human plasma using gas chromatography with electron-capture detection 749(2000)275

Heroin

Comparative analysis of heroin and cocaine seizures 733(1999)127

Heteroaromatic amines

Separation of heteroaromatic amines in food products 747(2000)171

Heterocyclic amines

Chromatographic and related techniques for the determination of aromatic heterocyclic amines in foods 747(2000)139

High throughput screening

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

Hinge glycopeptides

Mutual separation of hinge-glycopeptide isomers bearing five *N*-acetylgalactosamine residues from normal human serum immunoglobulin A1 by capillary electrophoresis 728(1999)175

Hippuric acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Histamine

Direct detection of endogenous histamine in rat peritoneal mast cells by in-capillary derivatization high-performance capillary electrophoresis 736(1999)263

Effect of 2-deoxy-D-glucose on acetylcholine and histamine levels in gastric juice of pylorus-ligated rats anesthetized with urethane 742(2000)295

Determination of biogenic amines in fish implicated in food poisoning by micellar electrokinetic capillary chromatography 749(2000)163

Histidine

Analysis of diagnostic metabolites by capillary electrophoresis–mass spectrometry 727(1999)43

Determination of histidine and urocanic acid isomers in the human skin by high-performance capillary electrophoresis 749(2000)41

Analysis of histidine and urocanic acid isomers by reversed-phase high-performance liquid chromatography 750(2001)71

Histone H1

Resolution of allelic and non-allelic variants of histone H1 by cation-exchange-hydrophilic-interaction chromatography 744(2000)33

HIV protease inhibitors

Determination of L-756 423, a novel HIV protease inhibitor, in human plasma and urine using high-performance liquid chromatography with fluorescence detection 732(1999)425

HIV-1 reverse-transcriptase inhibitor

Application of directly coupled LC–NMR–MS to the structural elucidation of metabolites of the HIV-1 reverse-transcriptase inhibitor BW935U83 748(2000)269

Homocysteine

Determination of homocysteine and its related compounds by solid-phase microextraction–gas chromatography–mass spectrometry 727(1999)1

Gas chromatographic–mass spectrometric determination of total homocysteine in human plasma by stable isotope dilution: method and clinical applications 729(1999)333

Determination of reduced and oxidized homocysteine and related thiols in plasma by thiol-specific pre-column derivatization and capillary electrophoresis with laser-induced fluorescence detection 735(1999)219

Determination of isotopic enrichments of [$1\text{-}^{13}\text{C}$]homocysteine, [$1\text{-}^{13}\text{C}$]methionine and [$2\text{H}_3\text{-methyl-1-}^{13}\text{C}$]methionine in human plasma by gas chromatography–negative chemical ionization mass spectrometry 738(2000)99

On-column derivatization for the analysis of homocysteine and other thiols by capillary electrophoresis with laser-induced fluorescence detection 744(2000)149

Homocystinuria

Differential diagnosis of homocystinuria by urease treatment, isotope dilution and gas chromatography–mass spectrometry 746(2000)103

Homogentisic acid

Analysis of diagnostic metabolites by capillary electrophoresis–mass spectrometry 727(1999)43

Homovanillic acid

Liquid chromatography–electrospray tandem mass spectrometry of acidic monoamine metabolites 744(2000)423

Hormones

High-performance affinity chromatography and immobilized serum albumin as probes for drug- and hormone-protein binding 739(2000)39

HPLC

Factors to consider in the development of generic bioanalytical high-performance liquid chromatographic–mass spectrometric methods to support drug discovery 748(2000)21

HPLC-NMR

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

5-HT_{1A} receptors

Enantioselective high-performance liquid chromatographic analysis of the 5-HT_{1A} receptor agonist 8-hydroxy-2-(di-*n*-propylamino)tetralin. Application to a pharmacokinetic–pharmacodynamic study in rats 738(2000)67

HTS

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

Human T-cell leukemia virus

Expression, purification and biological properties of the carboxyl half part of the HTLV-I surface envelope glycoprotein 737(2000)85

Human plasma

A sensitive high-performance liquid chromatographic method using electrochemical detection for the analysis of olanzapine and desmethylolanzapine in plasma of schizophrenic patients using a new solid-phase extraction procedure 750(2001)137

Human prothrombin

Purification of human prothrombin from Nitschmann fraction III using DEAE membrane radial flow chromatography 742(2000)109

Human serum albumin

Chromatographic study of magnesium and calcium binding to immobilized human serum albumin 728(1999)167

Non-covalent labeling of human serum albumin with indocyanine green: a study by capillary electrophoresis with diode laser-induced fluorescence detection 729(1999)55

Application of aqueous two-phase partition to the production of homogeneous preparations of fluorescently labelled human serum albumin 743(2000)263

Human serum albumin chromatography by Cibacron Blue F3GA-derived microporous polyamide hollow-fiber affinity membranes 746(2000)123

Hydantoin

Simultaneous determination of enantioselective plasma protein binding of aminohydantoins by ultrafiltration and chiral high-performance liquid chromatography 732(1999)31

Hydantoinase

Purification of recombinant hydantoinase and L-N-carbamoylase from *Arthrobacter aureus* expressed in *Escherichia coli*: comparison of wild-type and genetically modified proteins 737(2000)179

D-Hydantoinase

Purification of a D-hydantoinase using a laboratory-scale Streamline phenyl column as the initial step 737(2000)187

trans-4'-Hydrazino-2-stilbazole

Bioanalysis of aplidine, a new marine antitumoral depsipeptide, in plasma by high-performance liquid chromatography after derivatization with trans-4'-hydrazino-2-stilbazole 729(1999)43

Hydrocortisone

Simultaneous determination of hydrocortisone, dexamethasone, indomethacin, phenylbutazone and oxyphenbutazone in equine serum by high-performance liquid chromatography 738(2000)17

Hydrogenase–sulfur reductase complex

Improved purification of the membrane-bound hydrogenase–sulfur-reductase complex from thermophilic archaea using ϵ -aminocaproic acid-containing chromatography buffers 737(2000)151

Hydroperoxide

High-performance liquid chromatographic assay of hydroperoxide levels in oxidatively modified lipoproteins 731(1999)223

5-Hydroperoxyeicosatetraenoic acid

Determination of 5-hydroperoxyeicosatetraenoic acid produced in rat basophilic leukemia cell line RBL-2H3 by high-performance liquid chromatography with chemiluminescence detection 728(1999)35

3-Hydroxyanthranilic acid

3-Hydroxyanthranilic acid-derived compounds formed through electrochemical oxidation 736(1999)237

Hydroxyantipyrine

Determination of phenolic derivatives of antipyrine in plasma with solid-phase extraction and high-performance liquid chromatography–atmospheric-pressure chemical ionization mass spectrometry 732(1999)103

3-Hydroxybenz[a]anthracene

High-performance liquid chromatographic method with fluorescence detection for the determination of 3-hydroxybenzo[a]pyrene and 3-hydroxybenz[a]anthracene in the urine of polycyclic aromatic hydrocarbon-exposed workers 738(2000)47

***p*-Hydroxybenzoic acid-*O*-glucoside**

Simultaneous analysis of shikimate-derived secondary metabolites in *Lithospermum erythrorhizon* cell suspension cultures by high-performance liquid chromatography 738(2000)3

3-Hydroxybenzolketoprofen

Identification and confirmation of 3-hydroxy metabolite of ketoprofen in camels by gas chromatography–mass spectrometry and nuclear magnetic resonance spectroscopy 732(1999)299

3-Hydroxybenzo[a]pyrene

High-performance liquid chromatographic method with fluorescence detection for the determination of 3-hydroxybenzo[a]pyrene and 3-hydroxybenz[a]anthracene in the urine of polycyclic aromatic hydrocarbon-exposed workers 738(2000)47

Trace determination of urinary 3-hydroxybenzo[a]pyrene by automated column-switching high-performance liquid chromatography 748(2000)337

***N'*-Hydroxy-*p*-bromophenylacetylurea**

Application of the high-performance liquid chromatographic method for separation, purification and characterisation of *p*-bromophenylacetylurea and its metabolites (1999)349

3-Hydroxybutanoic acid

Enantioselective analysis of ketone bodies in patients with β -ketothiolase deficiency, medium-chain acyl coenzyme A dehydrogenase deficiency and ketonemic vomiting 739(2000)313

10-Hydroxycarbazepine

Simultaneous stereoselective high-performance liquid chromatographic determination of 10-hydroxycarbazepine and its metabolite carbamazepine-10,11-*trans*-dihydrodiol in human urine 738(2000)419

6 β -Hydroxycortisol

Simultaneous determination of urinary free cortisol and 6 β -hydroxycortisol by liquid chromatography–atmospheric pressure chemical ionization tandem mass spectrometry and its application for estimating hepatic CYP3A induction 742(2000)303

Specific determination of urinary 6 β -hydroxycortisol and cortisol by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 746(2000)95

Hydroxycortisol

Simultaneous determination of 6 β - and 6 α -hydroxycortisols and 6 β -hydroxycortisone in human urine by stable isotope dilution mass spectrometry 738(2000)367

Hydroxycortisone

Simultaneous determination of 6 β - and 6 α -hydroxycortisols and 6 β -hydroxycortisone in human urine by stable isotope dilution mass spectrometry 738(2000)367

4-Hydroxydebrisoquine

Enantioselectivity of debrisoquine 4-hydroxylation in Brazilian Caucasian hypertensive patients phenotyped as extensive metabolizers 749(2000)153

8-Hydroxy-2'-deoxyguanosine

Fast quantification of the urinary marker of oxidative stress 8-hydroxy-2'-deoxyguanosine using solid-phase extraction and high-performance liquid chromatography with triple-stage quadrupole mass detection 738(2000)311

8-Hydroxy-2-(di-*n*-propylamino)tetralin

Enantioselective high-performance liquid chromatographic analysis of the 5-HT_{1A} receptor agonist 8-hydroxy-2-(di-*n*-propylamino)tetralin. Application to a pharmacokinetic–pharmacodynamic study in rats 738(2000)67

Hydroxyethyl starch

Detection of the plasma volume expander hydroxyethyl starch in human urine 744(2000)345

4'-hydroxyflurbiprofen

Sensitive and specific high-performance liquid chromatographic assay for 4'-hydroxyflurbiprofen and flurbiprofen in human urine and plasma 749(2000)119

4-Hydroxyifosfamide

Determination of 4-hydroxyifosfamide concomitantly with ifosfamide and its dechloroethylated metabolites using gas chromatography and a nitrogen phosphorus-selective detector 732(1999)3

5-Hydroxyindoleacetic acid

Liquid chromatography–electrospray tandem mass spectrometry of acidic monoamine metabolites 744(2000)423

Clinical chemistry of serotonin and metabolites 747(2000)33

5-Hydroxyindoles

4-Dimethylaminobenzylamine as a sensitive chemiluminescence derivatization reagent for 5-hydroxyindoles and its application to their quantification in human platelet-poor plasma 738(2000)199

Hydroxyitraconazole

Liquid chromatographic–mass spectrometric determination of itraconazole and its major metabolite, hydroxyitraconazole, in dog plasma 745(2000)413

3-Hydroxyketoprofen

Identification and confirmation of 3-hydroxy metabolite of ketoprofen in camels by gas chromatography–mass spectrometry and nuclear magnetic resonance spectroscopy 732(1999)299

Hydroxyl radical

Measurement of hydroxyl radical in rat blood vessel by microbore liquid chromatography and electrochemical detection: an on-line microdialysis study 734(1999)277

3-Hydroxy-2-methylbutanoic acid

Enantioselective analysis of ketone bodies in patients with β -ketothiolase deficiency, medium-chain acyl coenzyme A dehydrogenase deficiency and ketonemic vomiting 739(2000)313

N-Hydroxymethyl-N-methylformamide

Simultaneous determination of two human urinary metabolites of N,N-dimethylformamide using gas chromatography–thermionic sensitive detection with mass spectrometric confirmation 734(1999)285

1-Hydroxymidazolam

Sensitive and specific determination of midazolam and 1-hydroxymidazolam in human serum by liquid chromatography–electrospray mass spectrometry 734(1999)137

4-(4'-Hydroxyphenyl)amino-6,7-dimethoxyquinazoline

Quantitative high-performance liquid chromatographic method for pharmacokinetic studies of the potent mast cell inhibitor 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline (WHI-P131) 727(1999)205

4-Hydroxyphenyllactic acid

Urine screening of five-day-old newborns: metabolic profiling of neonatal galactosuria 732(1999)469

4-Hydroxyproline

Analysis of 4-hydroxyproline using 4-chloro-7-nitrobenzo-2-oxa-1,3-diazol derivatization and micellar electrokinetic chromatography combined with laser-induced fluorescence detection 744(2000)195

5-Hydroxypropafenone

Determination of propafenone and its main metabolite 5-hydroxypropafenone in human serum with direct injection into a column-switching chromatographic system 726(1999)211

3-Hydroxypropylmercapturic acid

High-performance liquid chromatographic–tandem mass spectrometric determination of 3-hydroxypropylmercapturic acid in human urine 750(2001)163

Hydroxypropylphosphoramidate mustard

Analysis of cyclophosphamide and five metabolites from human plasma using liquid chromatography–mass spectrometry and gas chromatography–nitrogen–phosphorus detection 732(1999)287

1-Hydroxypyrene

Automated column-switching high-performance liquid chromatography method for the determination of 1-hydroxypyrene in human urine 732(1999)91

3-Hydroxypyridin-4-one

Gradient ion-pair high-performance liquid chromatographic method for analysis of 3-hydroxypyridin-4-one iron chelators 730(1999)135

Hydroxypyrimidine

Determination of metabolites of pirimicarb in human urine by gas chromatography–mass spectrometry 730(1999)229

9-Hydroxyrisperidone

Determination of risperidone and its major metabolite 9-hydroxyrisperidone in human plasma by reversed-phase liquid chromatography with ultraviolet detection 746(2000)173

 α -Hydroxysalmeterol

Rapid chiral high-performance liquid chromatographic assay for salmeterol and α -hydroxysalmeterol. Application to in vitro metabolism studies 729(1999)225

6 β -Hydroxytestosterone

Determination of testosterone and 6 β -hydroxytestosterone by gas chromatography–selected ion monitoring–mass spectrometry for the characterization of cytochrome p450 3a activity 734(1999)73

5-Hydroxytryptamine

Quantitative determination of L-775,606, a selective 5-hydroxytryptamine 1D agonist, in rat plasma using automated sample preparation and detection by liquid chromatography–tandem mass spectrometry 750(2001)13

5-Hydroxytryptophan

Clinical chemistry of serotonin and metabolites 747(2000)33

Hyperforin

Measurement of hyperforin a constituent of St. John's wort in plasma by high-performance liquid chromatography 735(1999)285

Hypericin

Determination of hypericin and pseudohypericin in pharmaceutical preparations by liquid chromatography with fluorescence detection 749(2000)57

Hypoxanthine

Quantification of allantoin, uric acid, xanthine and hypoxanthine in ovine urine by high-performance liquid chromatography and photodiode array detection 744(2000)129

I**Ibuprofen**

Validation of a high-performance liquid chromatographic method for the determination of ibuprofen enantiomers in plasma of broiler chickens 749(2000)243

Idarubicin

Determination of idarubicin and idarubicinol in rat plasma using reversed-phase high-performance liquid chromatography and fluorescence detection 728(1999)279

Simultaneous determination of four anthracyclines and three metabolites in human serum by liquid chromatography–electrospray mass spectrometry 738(2000)281

Idarubicinol

Determination of idarubicin and idarubicinol in rat plasma using reversed-phase high-performance liquid chromatography and fluorescence detection 728(1999)279

IDN5109

High-performance liquid chromatographic assay for the determination of the novel taxane derivative IDN5109 in mouse plasma 736(1999)135

Ifosfamide

Determination of 4-hydroxyifosfamide concomitantly with ifosfamide and its dechloroethylated metabolites using gas chromatography and a nitrogen phosphorus-selective detector 732(1999)3

Quantitative high-performance liquid chromatographic determination of acrolein in plasma after derivatization with Luminarin® 3 739(2000)239

Imidapril

Determination of imidapril and imidaprilat in human plasma by high-performance liquid chromatography–electrospray ionization tandem mass spectrometry 734(1999)145

Imidaprilat

Determination of imidapril and imidaprilat in human plasma by high-performance liquid chromatography–electrospray ionization tandem mass spectrometry 734(1999)145

Imidazole

Hydrophobic surfactant–imidazole derivative association studied by RPLC using a hydroorganic solution 749(2000)225

L-Iminoethyl-L-lysine

Determination of L-iminoethyl-L-lysine in serum by liquid chromatography 749(2000)35

Immunoaffinity capillary electrophoresis

Determination of immunoreactive gonadotropin-releasing hormone in serum and urine by on-line immunoaffinity capillary electrophoresis coupled to mass spectrometry 749(2000)197

Immunoglobulin

Affinity chromatography on immobilized “biomimetic” ligands. Synthesis, immobilization and chromatographic assessment of an immunoglobulin G binding ligand 740(2000)1

Affinity purification of immunoglobulins from chicken egg yolk using a new synthetic ligand 749(2000)233

Immunoglobulin A1

Mutual separation of hinge-glycopeptide isomers bearing five *N*-acetylgalactosamine residues from normal human serum immunoglobulin A1 by capillary electrophoresis 728(1999)175

Immunoglobulin G

Chromatographic behavior of mouse serum immunoglobulin G in protein G perfusion affinity chromatography 738(2000)149

Immunoglobulins

Carbohydrate release from picomole quantities of glycoprotein and characterisation of glycans by high-performance liquid chromatography and mass spectrometry 734(1999)169

Purification of horse immunoglobulin isotypes based on differential elution properties of isotypes from protein A and protein G columns 742(2000)327

Immunophilins

Three-step purification of a fragment of the large immunophilin FKBP52 737(2000)71

Impurities

Directly coupled HPLC–NMR and HPLC–NMR–MS in pharmaceutical research and development 748(2000)233

Inborn errors of metabolism

Inborn errors of metabolism discovered in Asian department of pediatrics and mental retardation research center 746(2000)41

Indinavir

Simultaneous determination of the HIV-protease inhibitors indinavir, nelfinavir, saquinavir and ritonavir in human plasma by reversed-phase high-performance liquid chromatography 727(1999)139

Determination of indinavir in human cerebrospinal fluid and plasma by solid-phase extraction and high-performance liquid chromatography with column switching 734(1999)63

High-performance liquid chromatographic method for the simultaneous determination of HIV-1 protease inhibitors indinavir, saquinavir and ritonavir in plasma of patients during highly active antiretroviral therapy 735(1999)41

Simultaneous determination of the HIV protease inhibitors indinavir, amprenavir, saquinavir, ritonavir, nelfinavir and the non-nucleoside reverse transcriptase inhibitor efavirenz by high-performance liquid chromatography after solid-phase extraction 740(2000)43

Indocyanine green

Non-covalent labeling of human serum albumin with indocyanine green: a study by capillary electrophoresis with diode laser-induced fluorescence detection 729(1999)55

Indoleamines

Reversed-phase high-performance liquid chromatography of catecholamines and indoleamines using a simple gradient solvent system and native fluorescence detection 744(2000)221

Separation and assay methods for melatonin and its precursors 747(2000)95

Indomethacin

Study of the solid-phase extraction of diclofenac sodium, indomethacin and phenylbutazone for their analysis in human urine by liquid chromatography 729(1999)139

Simultaneous determination of hydrocortisone, dexamethasone, indomethacin, phenylbutazone and oxyphenbutazone in equine serum by high-performance liquid chromatography 738(2000)17

Inosine monophosphate

Improved high-performance liquid chromatographic assay for the determination of “high-energy” phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

Inositol phosphates

High-performance thin-layer chromatography method for inositol phosphate analysis 736(1999)221

Insulin

New strategy for the design of ligands for the purification of pharmaceutical proteins by affinity chromatography 740(2000)17

Rapid simultaneous determination of glucagon and insulin by capillary electrophoresis immunoassays 742(2000)353

Partitioning of whey proteins, bovine serum albumin and porcine insulin in aqueous two-phase systems 743(2000)235

Development of two high-performance liquid chromatographic methods for the analysis and characterization of insulin and its degradation products in pharmaceutical preparations 749(2000)25

Inulin

Rapid and simple determination of inulin in biological fluids by high-performance liquid chromatography with light-scattering detection 744(2000)241

Iodixanol

On-line dialysis and quantitative high-performance liquid chromatography analysis of iodixanol in human, rat and monkey plasma 749(2000)135

***m*-Iodobenzylguanidine**

Bioanalysis of *m*-iodobenzylguanidine in plasma by high-performance liquid chromatography after derivatization with benzoil 730(1999)193

Ion-exchange HPLC

Separation and quantitation of metallothionein isoforms from liver of untreated rats by ion-exchange high-performance liquid chromatography and atomic absorption spectrometry 744(2000)273

Ion-suppression

Factors to consider in the development of generic bioanalytical high-performance liquid chromatographic–mass spectrometric methods to support drug discovery 748(2000)21

Iopipерidol

High-performance liquid chromatographic assay of the X-ray contrast agent iopipерidol in plasma and urine of rats and humans 729(1999)323

Iothalamate

Determination of iothalamate in rat urine, plasma, and tubular fluid by capillary electrophoresis 728(1999)143

High-performance liquid chromatographic determination of *p*-aminohippuric acid and iothalamate in human serum and urine: comparison of two sample preparation methods 740(2000)81

Irinotecan

Sensitive determination of irinotecan (CPT-11) and its active metabolite SN-38 in human serum using liquid chromatography–electrospray mass spectrometry 736(1999)175

Simple and rapid determination of irinotecan and its metabolite SN-38 in plasma by high-performance liquid-chromatography: application to clinical pharmacokinetic studies 740(2000)159

Iron chelators

Gradient ion-pair high-performance liquid chromatographic method for analysis of 3-hydroxypyridin-4-one iron chelators 730(1999)135

Isoascorbic acid

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Isobutyl nitrite

Analysis of isobutyl nitrite inhalant in rat and human blood: application for pharmacokinetic investigations 734(1999)83

Isoflurane

Solid-phase microextraction gas chromatographic–mass spectrometric method for the determination of inhalation anesthetics in urine 732(1999)115

Quantitative determination of isoflurane enantiomers in blood samples during and after surgery via headspace gas chromatography–mass spectrometry 735(1999)93

Reliable gas chromatographic–mass spectrometric method combined with a headspace autosampler for isoflurane determination in blood 742(2000)277

Isooctane

Conformational transition and mass transfer in extraction of proteins by AOT–alcohol–isooctane reverse micellar systems 743(2000)203

Isopeptide crosslinks

Rapid, single-step procedure for the identification of transglutaminase-mediated isopeptide crosslinks in amino acid digests 732(1999)65

Isoprene

Determination of isoprene in human expired breath using solid-phase microextraction and gas chromatography–mass spectrometry 739(2000)183

8-Isoprostaglandin F_{2α}

Tandem mass spectrometric quantification of 8-iso-prostaglandin F_{2α} and its metabolite 2,3-dinor-5,6-dihydro-8-iso-prostaglandin F_{2α} in human urine 744(2000)99

Rapid and sensitive quantification of 8-isoprostaglandin F_{2α} in human plasma and urine by liquid chromatography–electrospray ionization mass spectrometry 746(2000)17

Isoprostane

Smoking and oxidant stress: assay of isoprostane in human urine by gas chromatography–mass spectrometry 746(2000)11

Isoproturon

Immuno-affinity solid-phase extraction 745(2000)39

Isorhamnetin

High-performance liquid chromatographic determination of quercetin and isorhamnetin in rat tissues using β-glucuronidase and acid hydrolysis 738(2000)413

Isotopes

Factors to consider in the development of generic bioanalytical high-performance liquid chromatographic–mass spectrometric methods to support drug discovery 748(2000)21

Itraconazole

Liquid chromatographic–mass spectrometric determination of itraconazole and its major metabolite, hydroxyitraconazole, in dog plasma 745(2000)413

Ivabradine

Simultaneous determination of ivabradine and its metabolites in human plasma by liquid chromatography–tandem mass spectrometry 745(2000)261

K

Ketamine

Simultaneous analysis of ketamine and bupivacaine in plasma by high-performance liquid chromatography 728(1999)107

Gas chromatographic–mass spectrometric analysis of veterinary tranquilizers in urine: evaluation of method performance 734(1999)113

Ketobemidone

Validation of urine drug-of-abuse testing methods for ketobemidone using thin-layer chromatography and liquid chromatography–electrospray mass spectrometry 736(1999)103

α-Ketoisovalerate

Determination of extracellular and intracellular enrichments of [1-¹³C]-α-ketoisovalerate using enzymatically converted [1-¹³C]-valine standardization curves and gas chromatography–mass spectrometry 729(1999)211

Ketone bodies

Enantioselective analysis of ketone bodies in patients with β-ketothiolase deficiency, medium-chain acyl coenzyme A dehydrogenase deficiency and ketonemic vomiting 739(2000)313

3-Ketonorgestimate

Determination of norgestimate and its metabolites in human serum using high-performance liquid chromatography with tandem mass spectrometric detection 734(1999)247

Ketoprofen

Identification and confirmation of 3-hydroxy metabolite of ketoprofen in camels by gas chromatography–mass spectrometry and nuclear magnetic resonance spectroscopy 732(1999)299

17-Ketosteroid glucuronide

Quantification of urine 17-ketosteroid sulfates and glucuronides by high-performance liquid chromatography–ion trap mass spectroscopy 750(2001)81

17-Ketosteroid sulfate

Quantification of urine 17-ketosteroid sulfates and glucuronides by high-performance liquid chromatography–ion trap mass spectroscopy 750(2001)81

 β -Ketothiolase

Enantioselective analysis of ketone bodies in patients with β -ketothiolase deficiency, medium-chain acyl coenzyme A dehydrogenase deficiency and ketonemic vomiting 739(2000)313

Ketotifen

Gas chromatographic–mass spectrometric method for quantitative determination of ketotifen in human plasma after enzyme hydrolysis of conjugated ketotifen 732(1999)251

Kinesin

Three-step chromatographic purification procedure for the production of a His-tag recombinant kinesin overexpressed in *E. coli* 737(2000)143

Kinetics

Kinetics of binding properties of 5'-GMP with cisplatin under simulated physiological conditions by capillary electrophoresis 745(2000)211

KW-2149

Simultaneous determination of the peptide-mitomycin KW-2149 and its metabolites in plasma by high-performance liquid chromatography 739(2000)281

L

 β -Lactams

Simple and rapid analytical method for carbapenems using capillary zone electrophoresis 727(1999)219

 β -Lactamase

Hydrophobic character of surface regions and total hydrophobicity of four variants of chromosomal class C β -lactamase from *Pseudomonas aeruginosa* are identical. Chromatographic comparison of the hydrophobic character of the variants and the effect of focusing buffer composition on the separation of the variants by chromatofocusing with internal and external pH gradients 746(2000)161

 β -Lactamases

Technique for a rapid and efficient purification of the SHV-1 and PSE-2 β -lactamases 737(2000)261

Lactate dehydrogenase

Polymer displacement/shielding in protein chromatography 741(2000)103

Lactic acid

High-performance liquid chromatographic assay of (\pm)-lactic acid and its enantiomers in calf serum 727(1999)23

Lactoferrins

Various forms of mouse lactoferrins: purification and characterization 726(1999)45

 β -Lactoglobulin

Isolation and purification of serum and interfacial peptides of a trypsinolyzed β -lactoglobulin oil-in-water emulsion 744(2000)389

Lamivudine

Simultaneous determination of lamivudine and zidovudine concentrations in human seminal plasma using high-performance liquid chromatography and tandem mass spectrometry 742(2000)173

Lamotrigine

Determination of lamotrigine in human plasma by high-performance liquid chromatography 727(1999)113

Determination of lamotrigine in human serum by liquid chromatography 736(1999)295

Validated high-performance liquid chromatographic method for the determination of lamotrigine in human plasma 741(2000)289

Lanthanide

Sensitive high-performance liquid chromatographic assay for motexafin gadolinium and motexafin lutetium in human plasma 749(2000)145

Laser-induced fluorescence

Sensitive method for the determination of ambroxol in body fluids by capillary electrophoresis and fluorescence detection 742(2000)205

Lathosterol

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

LC-MS

Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments. Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry 748(2000)369

LC-MS-MS

Simultaneous determination of bosentan and its three major metabolites in various biological matrices and species using narrow bore liquid chromatography with ion spray tandem mass spectrometric detection 749(2000)67

LC-NMR

Application of liquid chromatography–nuclear magnetic resonance spectroscopy to the identification of natural products 748(2000)259

Leflunomide

Simultaneous separation by high-performance liquid chromatography of carbamoyl aspartate, carbamoyl phosphate and dihydroorotic acid 732(1999)487

Lenampicillin

Simple high-performance liquid chromatography determination of ampicillin in human serum using solid-phase extraction disk cartridges 727(1999)245

Leucine

Simultaneous determination of the enantiomers of leucine and [²H₇]leucine in plasma by capillary gas chromatography–mass spectrometry 735(1999)141

Determination of minute amounts of D-leucine in various brain regions of rat and mouse using column-switching high-performance liquid chromatography 744(2000)213

Leukeran

Rapid determination of the anti-cancer drug chlorambucil (LeukeranTM) and its phenyl acetic acid mustard metabolite in human serum and plasma by automated solid-phase extraction and liquid chromatography–tandem mass spectrometry 732(1999)173

Leukotrienes

Determination of leukotriene E₄ in human urine using liquid chromatography–tandem mass spectrometry 729(1999)279

Levamisole

Quantitative analysis of levamisole in porcine tissues by high-performance liquid chromatography combined with atmospheric pressure chemical ionization mass spectrometry 742(2000)283

Levetiracetam

Enantioselective analysis of levetiracetam and its enantiomer R- α -ethyl-2-oxo-pyrrolidine acetamide using gas chromatography and ion trap mass spectrometric detection 745(2000)325

Levormeloxifene

Assay for levormeloxifene, a selective estrogen receptor modulator, in human and monkey plasma employing high-performance liquid chromatography and solid-phase extraction 729(1999)315

Levorphanol

Quantitation of levorphanol in human plasma at subnanogram per milliliter levels using capillary gas chromatography with electron-capture detection 729(1999)173

Lidocaine

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

Evaluation of the programmed temperature vaporiser for large-volume injection of biological samples in gas chromatography 729(1999)199

Determination of lidocaine in plasma by direct solid-phase microextraction combined with gas chromatography 739(2000)175

High-performance liquid chromatography–tandem electrospray mass spectrometry for the determination of lidocaine and its metabolites in human plasma and urine 741(2000)175

Ligands

New strategy for the design of ligands for the purification of pharmaceutical proteins by affinity chromatography 740(2000)17

Lignins

Investigation of aqueous biphasic systems for the separation of lignins from cellulose in the paper pulping process 743(2000)127

d-Limonene

Gas chromatographic–mass spectrometric analysis of perillyl alcohol and metabolites in plasma 728(1999)85

Lipid A analogue

Quantitative determination of a potent lipopolysaccharide antagonist, E5564, in rat and dog plasma by high-performance liquid chromatography with fluorescence detection 736(1999)67

Lipids

Application of atmospheric pressure chemical ionization liquid chromatography–mass spectrometry in identification of lymph triacylglycerols 748(2000)425

Lipopolysaccharide antagonist

Quantitative determination of a potent lipopolysaccharide antagonist, E5564, in rat and dog plasma by high-performance liquid chromatography with fluorescence detection 736(1999)67

Lipopolysaccharides

Efficient method for preparation of highly purified lipopolysaccharides by hydrophobic interaction chromatography 732(1999)39

Lipoproteins

High-performance liquid chromatographic assay of hydroperoxide levels in oxidatively modified lipoproteins 731(1999)223

Liquid chromatography

A sensitive high-performance liquid chromatographic method using electrochemical detection for the analysis of olanzapine and desmethylolanzapine in plasma of schizophrenic patients using a new solid-phase extraction procedure 750(2001)137

Liver

Separation and quantitation of metallothionein isoforms from liver of untreated rats by ion-exchange high-performance liquid chromatography and atomic absorption spectrometry 744(2000)273

Loperamide

Quantitation of loperamide and *N*-demethyl-loperamide in human plasma using electrospray ionization with selected reaction ion monitoring liquid chromatography–mass spectrometry 744(2000)233

Lorazepam

Determination of lorazepam in plasma and urine as trimethylsilyl derivative using gas chromatography–tandem mass spectrometry 732(1999)509

Losartan

Method for the simultaneous determination of losartan and its major metabolite, EXP-3174, in human plasma by liquid chromatography–electrospray ionization tandem mass spectrometry 734(1999)325

Low density lipoproteins

Identification of aminoethylcysteine ketimine decarboxylated dimer in human plasma 732(1999)137

LSD

Liquid chromatography–mass spectrometry: potential in forensic and clinical toxicology 733(1999)93

Recent advances in chromatographic and mass spectrometric methods for determination of LSD and its metabolites in physiological specimens 733(1999)145

Luminarin

Quantitative high-performance liquid chromatographic determination of acrolein in plasma after derivatization with Luminarin® 3 739(2000)239

Lurtotecan

Liposomal lurtotecan (NX211): determination of total drug levels in human plasma and urine by reversed-phase high-performance liquid chromatography 738(2000)155

Lycopene

Determination of lycopene in tissues and plasma of rats by normal-phase high-performance liquid chromatography with photometric detection 739(2000)291

Lysine

Simple method for the simultaneous analysis of pipecolic acid and lysine by high-performance liquid chromatography and its application to rumen liquor and plasma of ruminants 735(1999)63

Lysine pyroglutamate

Determination of urea, allantoin and lysine pyroglutamate in cosmetic samples by hydrophilic interaction chromatography 742(2000)447

 α -Lysophosphatidylcholine

Quantitative determination of dipalmitoylphosphatidylcholine and palmitic acid in porcine lung surfactants used in the treatment of respiratory distress syndrome 744(2000)407

Lysozyme

Diffusion of lysozyme in gels and liquids. A general approach for the determination of diffusion coefficients using holographic laser interferometry 743(2000)151

L-756 423

Determination of L-756 423, a novel HIV protease inhibitor, in human plasma and urine using high-performance liquid chromatography with fluorescence detection 732(1999)425

L-775,606

Quantitative determination of L-775,606, a selective 5-hydroxytryptamine 1D agonist, in rat plasma using automated sample preparation and detection by liquid chromatography–tandem mass spectrometry 750(2001)13

M

M8

Determination of nelfinavir, a potent HIV protease inhibitor, and its active metabolite M8 in human plasma by high-performance liquid chromatography with photodiode-array detection 735(1999)159

Mabuterol

Multiresidue analysis of β_2 -agonists in human and calf urine using multimodal solid-phase extraction and high-performance liquid chromatography with electrochemical detection 726(1999)149

Magnesium

Chromatographic study of magnesium and calcium binding to immobilized human serum albumin 728(1999)167

On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography 750(2001)61

MAK33

Chromatographic purification of the C_H2 domain of the monoclonal antibody MAK33 737(2000)63

Malagashanine

High-performance liquid chromatographic assay of malagashanine in rat plasma and urine and its pharmacokinetic application 744(2000)121

Malarone

Simultaneous separation of atovaquone, proguanil and its metabolites on a mixed mode high-performance liquid chromatographic column 741(2000)189

Malonaldehyde

High-performance liquid chromatographic determination of 1,3-diethyl-2-thiobarbituric acid–malonaldehyde adduct in fish meat 726(1999)313

Malondialdehyde

Free malondialdehyde determination in rat brain tissue by capillary zone electrophoresis: evaluation of two protein removal procedures 740(2000)87

Measurement of free and bound malondialdehyde in plasma by high-performance liquid chromatography as the 2,4-dinitrophenylhydrazine derivative 742(2000)315

Mandelic acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Determination of mandelic acid enantiomers in urine by gas chromatography and electron-capture or flame ionisation detection 738(2000)39

Mapenterol

Multiresidue analysis of β_2 -agonists in human and calf urine using multimodal solid-phase extraction and high-performance liquid chromatography with electrochemical detection 726(1999)149

Marbofloxacin

Determination of marbofloxacin in plasma samples by high-performance liquid chromatography using fluorescence detection 729(1999)157

Mass spectrometry

Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments. Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry 748(2000)369

Mefenamic acid

Simultaneous analysis of anthranilic acid derivatives in pharmaceuticals and human urine by high-performance liquid chromatography with isocratic elution 744(2000)81

Mefloquine

Improved validated assay for the determination of mefloquine and its carboxy metabolite in plasma, serum and whole blood using solid-phase extraction and high-performance liquid chromatography 727(1999)159

Melatonin

Separation and assay methods for melatonin and its precursors 747(2000)95

Meloxicam

Development and validation of a new high-performance liquid chromatographic estimation method of meloxicam in biological samples 738(2000)431

Melphalan

Analysis of melphalan adducts of 2'-deoxynucleotides in calf thymus DNA hydrolysates by capillary high-performance liquid chromatography–electrospray tandem mass spectrometry 736(1999)43

Comparison between capillary and nano liquid chromatography–electrospray mass spectrometry for the analysis of minor DNA–melphalan adducts 748(2000)197

Memantine

Sensitive and selective liquid chromatographic assay of memantine in plasma with fluorescence detection after pre-column derivatization 729(1999)217

Membrane proteins

Novel, rapid purification of the membrane protein photosystem I by high-performance liquid chromatography on porous materials 737(2000)131

Capillary gel electrophoresis: separation of major erythrocyte membrane proteins 742(2000)411

New procedures for glycophorin A purification with high yield and high purity 750(2001)109

Membrane vesicles

Erratum to “Chromatographic approaches to liposomes, proteoliposomes and biomembrane vesicles”. [J. Chromatogr. B, 722 (1999) 103–120] 727(1999)249

Menthol

Sensitive and selective gas chromatographic methods for the quantitation of camphor, menthol and methyl salicylate from human plasma 729(1999)163

Mepivacaine

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

6-Mercaptopurine

Assay of 6-mercaptopurine and its metabolites in patient plasma by high-performance liquid chromatography with diode-array detection 732(1999)459

Metabolites

Simultaneous quantitative determination of cilostazol and its metabolites in human plasma by high-performance liquid chromatography 728(1999)251

Metalloprotease

New approach for separating *Bacillus subtilis* metalloprotease and α -amylase by affinity chromatography and for purifying neutral protease by hydrophobic chromatography 737(2000)277

Metallothionein

Biological significance of non-acetylated metallothionein 735(1999)17

Rapid identification of metallothionein isoforms in liver cytosol fraction by capillary zone electrophoresis using EDTA 742(2000)193

Separation and quantitation of metallothionein isoforms from liver of untreated rats by ion-exchange high-performance liquid chromatography and atomic absorption spectrometry 744(2000)273

Metallothionein isoforms

Identification of metallothionein isoforms on capillary zone electrophoresis by adding anti-metallothionein antibody 736(1999)185

Metandienone

Comparison of sensitivity between gas chromatography–low-resolution mass spectrometry and gas chromatography–high-resolution mass spectrometry for determining metandienone metabolites in urine 734(1999)179

Metanephries

High-performance liquid chromatographic assay for catecholamines and metanephries using fluorimetric detection with pre-column 9-fluorenylmethoxycarbonyl chloride derivatization 749(2000)179

Methadone

Novel high-performance liquid chromatographic and solid-phase extraction methods for quantitating methadone and its metabolite in spiked human urine 729(1999)19

Stereoselective quantification of methadone and its major oxidative metabolite, 2-ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine, in human urine using high-performance liquid chromatography 744(2000)165

Determination of methadone and its metabolites EDDP and EMDP in human hair by headspace solid-phase microextraction and gas chromatography–mass spectrometry 746(2000)255

Methamphetamine

Direct determination of methamphetamine enantiomers in urine by liquid chromatography with a strong cation-exchange precolumn and phenyl- β -cyclodextrin-bonded semi-microcolumn 729(1999)97

Simultaneous determination of dimethylamphetamine and its metabolites in rat hair by gas chromatography–mass spectrometry 741(2000)163

Determination of enantiomeric amphetamines as metabolites of illicit amphetamines and selegiline in urine by capillary electrophoresis using modified β -cyclodextrin 741(2000)221

Simultaneous chiral analysis of methamphetamine and related compounds by capillary electrophoresis 749(2000)111

Methanol

Analysis of methanol or formic acid in body fluids by headspace solid-phase microextraction and capillary gas chromatography 734(1999)155

Methionine

Determination of homocysteine and its related compounds by solid-phase microextraction–gas chromatography–mass spectrometry 727(1999)1

Determination of isotopic enrichments of [$1\text{-}^{13}\text{C}$]homocysteine, [$1\text{-}^{13}\text{C}$]methionine and [$^2\text{H}_3$ -methyl- $1\text{-}^{13}\text{C}$]methionine in human plasma by gas chromatography–negative chemical ionization mass spectrometry 738(2000)99

Convenient method of threonine, methionine and their related amino compounds by high-performance liquid chromatography and its application to rumen fluid 741(2000)279

Kinetics of binding properties of 5'-GMP with cisplatin under simulated physiological conditions by capillary electrophoresis 745(2000)211

Methionine enkephalin

Mass spectrometric analysis of neuropeptidergic systems in the human pituitary and cerebrospinal fluid (Review) 731(1999)3

Methotrexate

High-performance liquid chromatography of methotrexate for environmental monitoring of surface contamination in hospital departments and assessment of occupational exposure 726(1999)95

Liquid chromatographic analysis and preliminary pharmacokinetics of methotrexate in cancer patients co-treated with docetaxel 735(1999)111

Highly sensitive high-performance liquid chromatographic assay for methotrexate in the presence and absence of anti-methotrexate antibody fragments in rat and mouse plasma 736(1999)191

6-Methoxy-2-naphthylacetic acid

Extractionless determination of 6-methoxy-2-naphthylacetic acid, a major metabolite of nabumetone, in human plasma by high-performance liquid chromatography 740(2000)247

Methoxyresorufin

Determination of cytochrome P450 1A activities in mammalian liver microsomes by high-performance liquid chromatography with fluorescence detection 744(2000)399

2-(3-Methyl-4-aminophenyl)-benzothiazole

Analysis of 2-(3-methyl-4-aminophenyl)-benzothiazole (NSC 674495) in plasma by gas chromatography with mass-selective detection 732(1999)315

Methylarginines

Improved method for simultaneous determination of L-arginine and its mono- and dimethylated metabolites in biological samples by high-performance liquid chromatography 742(2000)199

Determination of arginine and methylated arginines in human plasma by liquid chromatography–tandem mass spectrometry 748(2000)157

N-Methylcarbamates

Reversed-phase liquid chromatographic column switching for the determination of N-methylcarbamates and some of their main metabolites in urine 738(2000)387

N'-Methyl-p-bromophenylacetylurea

Application of the high-performance liquid chromatographic method for separation, purification and characterisation of p-bromophenylacetylurea and its metabolites (1999)349

N'-Methyl-p-bromophenylacetylurea

Application of the high-performance liquid chromatographic method for separation, purification and characterisation of p-bromophenylacetylurea and its metabolites (1999)349

Methylecgonidine

Gas chromatographic–mass spectrometric detection of anhydroecgonine methyl ester (methylecgonidine) in human serum as evidence of recent smoking of crack 735(1999)127

Methylenedioxyamphetamine

Determination of “Ecstasy” components in alternative biological specimens 733(1999)137

Methylenedioxyated amphetamines

Simple and reliable high-performance liquid chromatography fluorimetric procedure for the determination of amphetamine-derived designer drugs 735(1999)243

3,4-Methylenedioxymethamphetamine

Simultaneous chiral analysis of methamphetamine and related compounds by capillary electrophoresis 749(2000)111

Methylhippuric acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

N^α-Methylhistamine

Analysis of N^α-methylhistamine by gas chromatography–mass spectrometry 739(2000)337

Methylmalonic acid

Optimization and performance of a rapid gas chromatography–mass spectrometry analysis for methylmalonic acid determination in serum and plasma 741(2000)231

6-Methylmercaptopurine

Thiopurine methyl transferase activity: new extraction conditions for high-performance liquid chromatographic assay 727(1999)235

Methyl 6-mercaptopurine nucleotides

High-performance liquid chromatographic determination of methyl 6-mercaptopurine nucleotides (Me6-MPN) in red blood cells: analysis of Me6-MPN per se or Me6-MPN derivative? 730(1999)273

Comments on “High-performance liquid chromatographic determination of methyl 6-mercaptopurine nucleotides in red blood cells” Reply to R. Boulieu and T. Dervieux 730(1999)275

Methylprednisolone

Isolation and identification of the C₆-hydroxy and C₂₀-hydroxy metabolites and glucuronide conjugate of methylprednisolone by preparative high-performance liquid chromatography from urine of patients receiving high-dose pulse therapy 726(1999)157

High-performance liquid chromatography analysis, preliminary pharmacokinetics, metabolism and renal excretion of methylprednisolone with its C₆ and C₂₀ hydroxy metabolites in multiple sclerosis patients receiving high-dose pulse therapy 732(1999)337

High-performance size-exclusion chromatographic analysis of dextran–methylprednisolone hemisuccinate in rat plasma 744(2000)293

Methyl salicylate

Sensitive and selective gas chromatographic methods for the quantitation of camphor, menthol and methyl salicylate from human plasma 729(1999)163

N-Methyl-D-aspartate

Determination of N-methyl-D-aspartate in tissues of bivalves by high-performance liquid chromatography 728(1999)41

17 α -Methyltestosterone

Stable isotope dilution analysis of human urinary metabolites of 17 α -methyltestosterone 741(2000)271

Methyltestosterone

Supercritical fluid extraction of methyltestosterone, nortestosterone and testosterone at low ppb levels from fortified bovine urine 726(1999)121

5-Methyltetrahydrofolate

High-performance liquid chromatographic assay for the determination of 5-methyltetrahydrofolate in human plasma 744(2000)307

1-Methyl-1,2,3,4-tetrahydroisoquinoline

Determination of 1-methyl-1,2,3,4-tetrahydroisoquinoline in mouse brain after treatment with haloperidol by gas chromatography–selected ion monitoring 731(1999)53

Metoprolol

Enantioselective analysis of metoprolol in plasma using high-performance liquid chromatographic direct and indirect separations: applications in pharmacokinetics 738(2000)27

Metrifonate

Development, validation and application of assays to quantify metrifonate and 2,2-dichlorovinyl dimethylphosphate in human body fluids 741(2000)257

Metronidazole

Simultaneous quantification of amoxycillin and metronidazole in plasma using high-performance liquid chromatography with photodiode array detection 731(1999)261

Microscale compartments

Analysis of small-scale biological compartments by capillary electrophoresis 745(2000)127

Midazolam

Sensitive and specific determination of midazolam and 1-hydroxymidazolam in human serum by liquid chromatography–electrospray mass spectrometry 734(1999)137

Minocycline

High-performance liquid chromatographic assay for minocycline in human plasma and parotid saliva 738(2000)357

Mirtazapone

Chiral determination of mirtazapine in human blood plasma by high-performance liquid chromatography 748(2000)439

Mobile phase

Factors to consider in the development of generic bioanalytical high-performance liquid chromatographic–mass spectrometric methods to support drug discovery 748(2000)21

Modafinil

Determination of the D- and L-enantiomers of modafinil in human plasma utilizing liquid–liquid extraction and high-performance liquid chromatography 730(1999)1

Monoacetylpolyamines

Rapid and simultaneous high-performance liquid chromatography assay of polyamines and monoacetylpolyamines in biological specimens 745(2000)431

Monoclonal antibodies

Separation of bovine serum albumin and its monoclonal antibody from their immunocomplexes by sodium dodecyl sulfate–capillary gel electrophoresis and its application in capillary electrophoresis-based immunoassay 731(1999)389

Chromatographic purification of the C_H2 domain of the monoclonal antibody MAK33 737(2000)63

Mono-N-dealkyldisopyramide

Enantioselective analysis of disopyramide and mono-N-dealkyldisopyramide in plasma and urine by high-performance liquid chromatography on an amylose-derived chiral stationary phase 744(2000)299

Monoethylglycinexylidide

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

High-performance liquid chromatography–tandem electrospray mass spectrometry for the determination of lidocaine and its metabolites in human plasma and urine 741(2000)175

Morphine

Immunoaffinity extraction of morphine, morphine-3-glucuronide and morphine-6-glucuronide from blood of heroin victims for simultaneous high-performance liquid chromatographic determination 726(1999)111

Highly sensitive gas chromatographic–tandem mass spectrometric method for the determination of morphine and codeine in serum and urine in the femtomolar range 727(1999)81

Detection and identification of morphine in urine extracts using thin-layer chromatography and tandem mass spectrometry 729(1999)341

Liquid chromatography–mass spectrometry: potential in forensic and clinical toxicology 733(1999)93

Simultaneous assay of morphine, morphine-3-glucuronide and morphine-6-glucuronide in human plasma using normal-phase liquid chromatography–tandem mass spectrometry with a silica

column and an aqueous organic mobile phase 735(1999)255

High-performance liquid chromatographic determination of morphine and its 3- and 6-glucuronide metabolites by two-step solid-phase extraction 742(2000)115

Quantitative analysis of morphine in human plasma by gas chromatography–negative ion chemical ionization mass spectrometry 744(2000)113

Immuno-affinity solid-phase extraction 745(2000)39

Morphine glucuronide

Immunoaffinity extraction of morphine, morphine-3-glucuronide and morphine-6-glucuronide from blood of heroin victims for simultaneous high-performance liquid chromatographic determination 726(1999)111

Morphine glucuronides

Simultaneous assay of morphine, morphine-3-glucuronide and morphine-6-glucuronide in human plasma using normal-phase liquid chromatography–tandem mass spectrometry with a silica column and an aqueous organic mobile phase 735(1999)255

High-performance liquid chromatographic determination of morphine and its 3- and 6-glucuronide metabolites by two-step solid-phase extraction 742(2000)115

Motexafin gadolinium

Sensitive high-performance liquid chromatographic assay for motexafin gadolinium and motexafin lutetium in human plasma 749(2000)145

Motexafin lutetium

Sensitive high-performance liquid chromatographic assay for motexafin gadolinium and motexafin lutetium in human plasma 749(2000)145

Motilin receptor agonist

Application of liquid chromatography–turbo ion spray tandem mass spectrometry for quantitative analysis of a potent motilin receptor agonist, EM574, and its metabolites in human plasma 734(1999)101

Moxifloxacin

Determination of moxifloxacin (BAY 12-8039) in plasma and lung tissue by high-performance liquid chromatography with ultraviolet detection using a fully automated extraction method with a new polymeric cartridge 742(2000)247

Muramic acid

New and simple procedure for the determination of muramic acid in chemically complex environments by gas chromatography–ion trap tandem mass spectrometry 738(2000)57

Mycoloylglycerol

Detection of mycoloylglycerol by thin-layer chromatography as a tool for the rapid inclusion of corynebacteria of clinical origin in the genus *Corynebacterium* 738(2000)181

Mycophenolic acid

Simple and sensitive high-performance liquid chromatographic method for the determination of mycophenolic acid in plasma 738(2000)169

Quantification of free mycophenolic acid by high-performance liquid chromatography–atmospheric pressure chemical ionisation tandem mass spectrometry 748(2000)151

Liquid chromatographic assay for simultaneous determination of abacavir and mycophenolic acid in human plasma using dual spectrophotometric detection 750(2001)155

N

Na,K-ATPase

Assay of ATPase and Na,K-ATPase activity using high-performance liquid chromatographic determination of ADP derived from ATP 744(2000)19

Nabumetone

Extractionless determination of 6-methoxy-2-naphthylacetic acid, a major metabolite of nabumetone, in human plasma by high-performance liquid chromatography 740(2000)247

NADH oxidase

Applications of aqueous two-phase partition to isolation of membranes from plants: A periodic NADH oxidase activity as a marker for right side-out plasma membrane vesicles 743(2000)369

Nalbuphine

High-performance liquid chromatographic method for the simultaneous determination of nalbuphine and its prodrug, seabacoyl dinalbuphine ester, in dog plasma and application to pharmacokinetic studies in dogs 746(2000)241

Naphthalene

Assay of 2-naphthol in human urine by high-performance liquid chromatography 734(1999)211

Naphthaquinone-4-sulphonate

Routine analysis of amphetamine class drugs as their naphthaquinone derivatives in human urine by high-performance liquid chromatography 735(1999)229

Naphthol

Naphthol- and resorcinol-based azo dyes as metal ion complexants in aqueous biphasic systems 743(2000)107

2-Naphthol

Assay of 2-naphthol in human urine by high-performance liquid chromatography 734(1999)211

1-Naphthylisothiocyanate

Semi-automatic liquid chromatographic analysis of pamidronate in urine after derivatization with 1-naphthylisothiocyanate 730(1999)95

Natural products

Directly coupled HPLC–NMR and HPLC–NMR–MS in pharmaceutical research and development 748(2000)233

Nefazodone

Determination of nefazodone and its pharmacologically active metabolites in human blood plasma and breast milk by high-performance liquid chromatography 730(1999)249

Nelarabine

High-performance liquid chromatography method for the determination and quantitation of arabinosylguanine triphosphate and fludarabine triphosphate in human cells 745(2000)421

Nelfinavir

Simultaneous determination of the HIV-protease inhibitors indinavir, nelfinavir, saquinavir and ritonavir in human plasma by reversed-phase high-performance liquid chromatography 727(1999)139

Determination of nelfinavir, a potent HIV protease inhibitor, and its active metabolite M8 in human plasma by high-performance liquid chromatography with photodiode-array detection 735(1999)159

Simultaneous determination of the HIV protease inhibitors indinavir, amprenavir, saquinavir, ritonavir, nelfinavir and the non-nucleoside reverse transcriptase inhibitor efavirenz by high-performance liquid chromatography after solid-phase extraction 740(2000)43

Simultaneous high-performance liquid chromatographic determination of the antiretroviral agents amprenavir, nelfinavir, ritonavir, saquinavir, delavirdine and efavirenz in human plasma 742(2000)453

Neuroaminidase inhibitor

Liquid chromatographic–tandem mass spectrometric method for the determination of the neuraminidase inhibitor zanamivir (GG167) in human serum 732(1999)383

Neurokinin-1 receptor antagonist

Liquid chromatographic–tandem mass spectrometric method for the analysis of a neurokinin-1 antagonist and its metabolite using automated solid-phase sample preparation and automated data handling and reporting 748(2000)55

Neuropeptides

Mass spectrometric analysis of neuropeptidergic systems in the human pituitary and cerebrospinal fluid (Review) 731(1999)3

Neuropeptide Y

Separation and identification of neuropeptide Y, two of its fragments and their degradation products using capillary electrophoresis–mass spectrometry 727(1999)53

Neurosteroids

Validation of an analytical procedure to measure trace amounts of neurosteroids in brain tissue by gas chromatography–mass spectrometry 739(2000)301

Studies on neurosteroids XII. Determination of enzymatically formed catechol estrogens and guaiacol estrogens by rat brains using liquid chromatography–mass spectrometry–mass spectrometry 748(2000)89

Neurotoxic compounds

Development of a method for sample preparation for subsequent identification and measurement of 1,2,3,4-tetrahydroisoquinolines and other potentially neurotoxic compounds by high-performance liquid chromatography with ultraviolet and fluorescence detection in blood plasma of Parkinson's disease patients 746(2000)283

High-performance liquid chromatographic separation and measurement of various biogenic compounds possibly involved in the pathomechanism of Parkinson's disease 746(2000)297

NeutrAvidin

Affinity partitioning of biotinylated mixed liposomes: effect of charge on biotin–NeutrAvidin interaction 743(2000)389

Nevirapine

Determination of nevirapine, an HIV-1 non-nucleoside reverse transcriptase inhibitor, in human plasma by reversed-phase high-performance liquid chromatography 744(2000)65

NE-100

Ultrasensitive determination of NE-100, a novel sigma ligand, in human plasma by liquid chromatography and electrospray ionization tandem mass spectrometry combined with a column-switching technique 730(1999)61

Nickel chelates

Purification of the membrane binding domain of cytochrome *b₅* by immobilised nickel chelate chromatography 737(2000)119

Nicorandil

Routine and sensitive method for determination of nicorandil in human plasma developed for liquid chromatography with ultraviolet and mass spectrometric detection 735(1999)103

Nicotinamide

Chromatographic and capillary electrophoretic methods for the analysis of nicotinic acid and its metabolites 747(2000)229

Nicotine

Improved highly sensitive method for determination of nicotine and cotinine in human plasma by high-performance liquid chromatography 742(2000)211

High-performance liquid chromatographic determination of cotinine in urine in isocratic mode 746(2000)115

Nicotinic acid

Chromatographic and capillary electrophoretic methods for the analysis of nicotinic acid and its metabolites 747(2000)229

Nicotinic acid mononucleotide

Reversed-phase high-performance liquid chromatography of nicotinic acid mononucleotide for measurement of quinolate phosphoribosyltransferase 749(2000)281

Nifedipine

Sensitive high-performance liquid chromatographic determination of nifedipine in cat plasma following improved sample treatment 729(1999)265

Nimesulide

Rapid and sensitive method for determination of nimesulide in human plasma by high-performance liquid chromatography 727(1999)241

Nitrate

Simple and rapid method for the measurement of nitrite and nitrate in human plasma and cerebrospinal fluid by capillary electrophoresis 727(1999)73

Interference by *S*-nitroso compounds with the measurement of nitrate in methods requiring reduction of nitrate to nitrite by cadmium 729(1999)375

Gas chromatographic–tandem mass spectrometric quantification of human plasma and urinary nitrate after its reduction to nitrite and derivatization to the pentafluorobenzyl derivative 731(1999)285

Determination of nitrate in blood by gas chromatography and gas chromatography–mass spectrometry 742(2000)363

Ion chromatographic method for simultaneous determination of nitrate and nitrite in human saliva 744(2000)433

Rapid determination of nitrite by reversed-phase high-performance liquid chromatography with fluorescence detection 746(2000)199

Nitric oxide synthase

Assessment of nitric oxide synthase activity in vitro and in vivo by gas chromatography–mass spectrometry 742(2000)143

Nitrite

Simple and rapid method for the measurement of nitrite and nitrate in human plasma and cerebrospinal fluid by capillary electrophoresis 727(1999)73

Interference by *S*-nitroso compounds with the measurement of nitrate in methods requiring reduction of nitrate to nitrite by cadmium 729(1999)375

Gas chromatographic–tandem mass spectrometric quantification of human plasma and urinary nitrate after its reduction to nitrite and derivatization to the pentafluorobenzyl derivative 731(1999)285

Ion chromatographic method for simultaneous determination of nitrate and nitrite in human saliva 744(2000)433

Rapid determination of nitrite by reversed-phase high-performance liquid chromatography with fluorescence detection 746(2000)199

Nitrobenzylcarbamate isopropylidene derivative

Determination of the GABA_B receptor agonist CGP 44532 (3-amino-2-hydroxypropylmethylphosphinic acid) in rat plasma after pre-column derivatization by micro-high-performance liquid chromatography combined with negative electrospray tandem mass spectrometry 748(2000)349

***S*-Nitrosoalbumin**

Measurement of *S*-nitrosoalbumin by gas chromatography–mass spectrometry. I. Preparation, purification, isolation, characterization and metabolism of *S*-[¹⁵N]nitrosoalbumin in human blood in vitro 726(1999)1

Measurement of *S*-nitrosoalbumin by gas chromatography–mass spectrometry. II. Quantitative determination of *S*-nitrosoalbumin in human plasma using *S*-[¹⁵N]nitrosoalbumin as internal standard 726(1999)13

***S*-Nitroso compounds**

Interference by *S*-nitroso compounds with the measurement of nitrate in methods requiring reduction of nitrate to nitrite by cadmium 729(1999)375

Electrospray ionization mass spectrometry of low-molecular-mass *S*-nitroso compounds and their thiols 742(2000)99

2-(2-Nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione

Determination of 2-(2-nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione in plasma by direct injection into a coupled column liquid chromatographic system 730(1999)177

Nitrous oxide

Solid-phase microextraction gas chromatographic–mass spectrometric method for the determination of inhalation anesthetics in urine 732(1999)115

Non-nucleoside reverse transcriptase inhibitors

Determination of twelve antiretroviral agents in human plasma sample using reversed-phase high-performance liquid chromatography 744(2000)227

Non-steroidal anti-inflammatory drugs

Development of predictive retention–activity relationship models of non-steroidal anti-inflammatory drugs by micellar liquid chromatography: comparison with immobilized artificial membrane columns 740(2000)59

4-Nonylphenol

Determination of 4-nonylphenol, nonylphenol monoethoxylate, nonylphenol diethoxylate and other alkylphenols in fish and shellfish by high-performance liquid chromatography with fluorescence detection 746(2000)305

Nonylphenol ethoxylate

Determination of 4-nonylphenol, nonylphenol monoethoxylate, nonylphenol diethoxylate and other alkylphenols in fish and shellfish by high-performance liquid chromatography with fluorescence detection 746(2000)305

11-Nor- Δ^9 -tetrahydrocannabinol-9-carboxylic acid

Determination of 11-nor- Δ^9 -tetrahydrocannabinol-9-carboxylic acid in urine using high-performance liquid chromatography and electrospray ionization mass spectrometry 732(1999)155

Norephedrine

Sensitive determination of ephedrine and norephedrine in human plasma samples using derivatization with 9-fluorenylmethyl chloroformate and liquid chromatography 744(2000)25

Norepinephrine

Simultaneous determination of catecholamines in rat brain tissue by high-performance liquid chromatography 731(1999)361

Determination of norepinephrine in small volume plasma samples by stable-isotope dilution gas chromatography–tandem mass spectrometry with negative ion chemical ionization 738(2000)319

High-performance liquid chromatographic analysis of muscular interstitial arginine and norepinephrine kinetics. A microdialysis study in rats 745(2000)279

Determination of urine catecholamines by capillary electrophoresis with dual-electrode amperometric detection 750(2001)33

Norethandrolone

17 α -Ethyl-5 β -estrane-3 α ,17 β -diol, a biological marker for the abuse of norethandrolone and ethylestrenol in slaughter cattle 728(1999)217

Norgestimate

Determination of norgestimate and its metabolites in human serum using high-performance liquid chromatography with tandem mass spectrometric detection 734(1999)247

Norgestrel

Determination of norgestimate and its metabolites in human serum using high-performance liquid chromatography with tandem mass spectrometric detection 734(1999)247

Nortestosterone

Supercritical fluid extraction of methyltestosterone, nortestosterone and testosterone at low ppb levels from fortified bovine urine 726(1999)121

Nortriptyline

Determination of nortriptyline in human serum by fully automated solid-phase extraction and on-line high-performance liquid chromatography in the presence of antipsychotic drugs 746(2000)233

Norverapamil

Achiral and chiral high-performance liquid chromatography of verapamil and its metabolites in serum samples 732(1999)395

NSC 674495

Analysis of 2-(3-methyl-4-aminophenyl)-benzothiazole (NSC 674495) in plasma by gas chromatography with mass-selective detection 732(1999)315

NSC 686288

Identification of the principal circulating metabolite of a synthetic 5,4'-diaminoflavone (NSC 686288), an antitumor agent, in the rat 741(2000)205

Nucleic acids

Monitoring hybridization during polymerase chain reaction 741(2000)3

Nucleoside monophosphates

Kinetics of binding properties of 5'-GMP with cisplatin under simulated physiological conditions by capillary electrophoresis 745(2000)211

Nucleoside reverse transcriptase inhibitors

Determination of twelve antiretroviral agents in human plasma sample using reversed-phase high-performance liquid chromatography 744(2000)227

Nucleosides

Erratum to "Determination of adenosine and deoxyadenosine in urine by high-performance liquid chromatography with column switching". [J. Chromatogr. B, 719 (1998) 55–61] 726(1999)317

Reversed-phase high-performance liquid chromatographic investigation of urinary normal and modified nucleosides of cancer patients 732(1999)307

Application of capillary electrophoresis in clinical chemistry: the clinical value of urinary modified nucleosides 745(2000)189

Nucleotides

Analysis of melphalan adducts of 2'-deoxynucleotides in calf thymus DNA hydrolysates by capillary high-performance liquid chromatography–electrospray tandem mass spectrometry 736(1999)43

High-performance thin-layer chromatography method for inositol phosphate analysis 736(1999)221

Separation of near-infrared fluorescent conjugates of dATP and related compounds by capillary electrophoresis 744(2000)315

NX1838

Detection and plasma pharmacokinetics of an anti-vascular endothelial growth factor oligonucleotide-aptamer (NX1838) in rhesus monkeys 732(1999)203

Nystatin

High-performance liquid chromatographic determination of liposomal nystatin in plasma and tissues for pharmacokinetic and tissue distribution studies 735(1999)51

O**[¹³C]octanoic acid**

Improved use of the [¹³C]octanoic acid breath test as intra-individual parameter to study the effect of a prokinetic drug on gastric emptying in preterm infants with oral feeding intolerance 750(2001)147

Ofloxacin

High-performance liquid chromatography separation and quantitation of ofloxacin enantiomers in rat microsomes 728(1999)151

Olanzapine

Determination of olanzapine in human breast milk by high-performance liquid chromatography with electrochemical detection 726(1999)203

Monitoring of olanzapine in serum by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 732(1999)257

A sensitive high-performance liquid chromatographic method using electrochemical detection for the analysis of olanzapine and desmethylolanzapine in plasma of schizophrenic patients using a new solid-phase extraction procedure 750(2001)137

Oleandrine

Poisoning by plant material: review of human cases and analytical determination of main toxins by high-performance liquid chromatography–(tandem) mass spectrometry 733(1999)181

Oligonucleotides

Allelic discrimination by denaturing high-performance liquid chromatography 739(2000)345

Detection of a mixed-backbone oligonucleotide (GEM 231) in liver and tumor tissues by capillary electrophoresis 750(2001)129

Oligosaccharides

Mutual separation of hinge-glycopeptide isomers bearing five *N*-acetylgalactosamine residues from normal human serum immunoglobulin A1 by capillary electrophoresis 728(1999)175

High resolution of oligosaccharide mixtures by ultrahigh voltage micellar electrokinetic capillary chromatography 745(2000)365

Olopatadine

Determination of olopatadine, a new antiallergic agent, and its metabolites in human plasma by high-performance liquid chromatography with electrospray ionization tandem mass spectrometry 731(1999)345

Olpadronate

Semi-automatic liquid chromatographic analysis of olpadronate in urine and serum using derivatization with (9-fluorenylmethyl)chloroformate 738(2000)331

Omeprazole

Determination of the enantiomers of omeprazole in blood plasma by normal-phase liquid chromatography and detection by atmospheric pressure ionization tandem mass spectrometry 734(1999)191

On-line preconcentration CE

Determination of immunoreactive gonadotropin-releasing hormone in serum and urine by on-line immunoaffinity capillary electrophoresis coupled to mass spectrometry 749(2000)197

Opiates

Simultaneous assay of morphine, morphine-3-glucuronide and morphine-6-glucuronide in human plasma using normal-phase liquid chromatography–tandem mass spectrometry with a silica column and an aqueous organic mobile phase 735(1999)255

***O*⁶-Benzylguanine**

Determination of 8-oxoguanine in human plasma and urine by high-performance liquid chromatography with electrochemical detection 731(1999)241

Organic compounds

Relative hydrophobicity of organic compounds measured by partitioning in aqueous two-phase systems 743(2000)187

Immunoaffinity solid-phase extraction for the trace-analysis of low-molecular-mass analytes in complex sample matrices 745(2000)15

Ornithine

Analysis of diagnostic metabolites by capillary electrophoresis–mass spectrometry 727(1999)43

Ortic acid

Determination of urinary orotic acid and uracil by capillary zone electrophoresis 734(1999)175

Oseltamivir

Development of a high-performance liquid chromatographic–mass spectrometric assay for the specific and sensitive quantification of Ro 64-0802, an anti-influenza drug, and its pro-drug, oseltamivir, in human and animal plasma and urine 745(2000)373

Ovoglycoprotein

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. VIII. Chiral recognition ability of partially and completely deglycosylated ovoglycoprotein 745(2000)149

Oxalic acid

Sensitive high-performance liquid chromatography method for the simultaneous determination of low levels of dichloroacetic acid and its metabolites in blood and urine 729(1999)271

Oxidized low density lipoprotein

High-performance liquid chromatographic assay of hydroperoxide levels in oxidatively modified lipoproteins 731(1999)223

4-Oxocyclophosphamide

Analysis of cyclophosphamide and five metabolites from human plasma using liquid chromatography–mass spectrometry and gas chromatography–nitrogen–phosphorus detection 732(1999)287

8-Oxoguanine

Determination of 8-oxoguanine in human plasma and urine by high-performance liquid chromatography with electrochemical detection 731(1999)241

Oxolinic acid

Determination of oxolinic acid in faeces and urine of turbot (*Scophthalmus maximus*) by high-performance liquid chromatography using fluorescence detection 749(2000)127

Oxybutynin

Extraction and determination of oxybutynin in human bladder samples by reversed-phase high-performance liquid chromatography 734(1999)163

Oxyphenbutazone

Simultaneous determination of hydrocortisone, dexamethasone, indomethacin, phenylbutazone and oxyphenbutazone in equine serum by high-performance liquid chromatography 738(2000)17

Oxytetracycline

Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments. Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry 748(2000)369

P

Paeoniflorin

Pharmacokinetic study of paeoniflorin in mice after oral administration of Paeoniae radix extract 735(1999)33

Palmitic acid

Quantitative determination of dipalmitoylphosphatidylcholine and palmitic acid in porcine lung surfactants used in the treatment of respiratory distress syndrome 744(2000)407

Pamidronate

Semi-automatic liquid chromatographic analysis of pamidronate in urine after derivatization with 1-naphthylisothiocyanate 730(1999)95

Pancreatistatin

Development and validation of a high-performance liquid chromatographic assay using solid-phase extraction for the novel antitumor agent pancreatistatin in human plasma 726(1999)249

Paroxetine

Simultaneous determination of citalopram, fluoxetine, paroxetine and their metabolites in plasma and whole blood by high-performance liquid chromatography with ultraviolet and fluorescence detection 734(1999)229

Determination of paroxetine levels in human plasma using gas chromatography with electron-capture detection 749(2000)275

Particle beam

Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments. Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry 748(2000)369

PDC-109 protein

Inhibition of acrosine-like protease activity by a lectin affinity chromatographic bovine seminal plasma fraction containing the PDC-109 and aSFP proteins 746(2000)141

Penicillin acylase

Partial purification of penicillin acylase from *Escherichia coli* in poly(ethylene glycol)–sodium citrate aqueous two-phase systems 734(1999)15

Preparation, evaluation and application of new pseudo-affinity chromatographic supports for penicillin acylase purification 739(2000)63

Penicillins

Simultaneous determination of seven penicillins in muscle, liver and kidney tissues from cattle and pigs by a multiresidue high-performance liquid chromatographic method 734(1999)307

Pentafluorophenylpropyl

Performance of a pentafluorophenylpropyl stationary phase for the electrospray ionization high-performance liquid chromatography–mass spectrometry–mass spectrometry assay of cocaine and its metabolite ecgonine methyl ester in human urine 748(2000)77

Pentazocine

Liquid chromatography–fast atom bombardment mass spectrometry for detection and determination of pentazocine in human tissues 731(1999)149

6-Pentyl- α -pyrone

Aroma compounds recovery from mycelial cultures in aqueous two-phase processes 743(2000)403

Peptide-mitomycin

Simultaneous determination of the peptide-mitomycin KW-2149 and its metabolites in plasma by high-performance liquid chromatography 739(2000)281

Peptides

Composition of the peptide fraction in human blood plasma: database of circulating human peptides 726(1999)25

Development of a gradient elution high-performance liquid chromatography assay with ultraviolet detection for the determination in plasma of the anticancer peptide [Arg⁶, D-Trp^{7,9}, MePhe⁸]-substance P (6–11) (antagonist G), its major metabolites and a C-terminal pyrene-labelled conjugate 732(1999)277

Purification of the c-erbB2/*neu* membrane-spanning segment: a hydrophobic challenge 737(2000)55

Application of Pluronic copolymer liquid crystals for the capillary electrophoretic separation of collagen type I cyanogen bromide fragments 739(2000)109

Capillary electrophoretic separation of proteins and peptides

using Pluronic liquid crystals and surface-modified capillaries 741(2000)37

External electric field control of electroosmotic flow in non-coated and coated fused-silica capillaries and its application for capillary electrophoretic separations of peptides 741(2000)43

Separation and detection techniques for peptides and proteins in stability research and bioanalysis 742(2000)401

Peptides partitioning in an aqueous dextran–polyethylene glycol two-phase system 743(2000)271

Partitioning of peptides and recombinant protein–peptide fusions in thermoseparating aqueous two-phase systems: effect of peptide primary structure 743(2000)295

Isolation and purification of serum and interfacial peptides of a trypsinolyzed β -lactoglobulin oil-in-water emulsion 744(2000)389

Integrated microchip-device for the digestion, separation and postcolumn labeling of proteins and peptides 745(2000)243

Comparison of different electrokinetic separation modes applicable to a model peptide mixture (collagen type I and III CNBr fragments) 745(2000)251

Determination of ionization efficiency of glycosylated and non-glycosylated peptides from the N-terminal of hemoglobin β -chain by electrospray ionization mass spectrometry 746(2000)83

Peptidyl-prolyl *cis*–*trans* isomerases

Three-step chromatographic purification of Cpr6, a cyclophilin from *Saccharomyces cerevisiae* 737(2000)295

Perflubron

Assay method for the perfluorooctyl bromide (perflubron) in rat blood by gas chromatography–mass spectrometry 734(1999)267

Perfluorodecalin

Determination of perfluorodecalin and perfluoro-*N*-methylcyclohexylpiperidine in rat blood by gas chromatography–mass spectrometry 745(2000)333

Perfluoro-*N*-methylcyclohexylpiperidine

Determination of perfluorodecalin and perfluoro-*N*-methylcyclohexylpiperidine in rat blood by gas chromatography–mass spectrometry 745(2000)333

Perfluorooctyl bromide

Assay method for the perfluorooctyl bromide (perflubron) in rat blood by gas chromatography–mass spectrometry 734(1999)267

Perillyl alcohol

Gas chromatographic–mass spectrometric analysis of perillyl alcohol and metabolites in plasma 728(1999)85

Permethrin

Simultaneous determination of pyridostigmine bromide, *N,N*-diethyl-*m*-toluamide, permethrin, and their metabolites in rat plasma and urine by high-performance liquid chromatography 749(2000)171

Peroxisomicine

High-performance liquid chromatographic determination of peroxisomicine A₁ (T-514) in genus *Karwinskia* 732(1999)405

Pesticides

Separation methods for amino group-possessing pesticides in biological samples 747(2000)241

Pharmaceuticals

Testing hair for pharmaceuticals 733(1999)231

Chromatographic assay of pharmaceutical compounds under column overloading 741(2000)55

Sensitive method for the determination of ambroxol in body fluids by capillary electrophoresis and fluorescence detection 742(2000)205

Capillary electrophoresis in the pharmaceutical industry: applications in discovery and chemical development 745(2000)177

Directly coupled HPLC–NMR and HPLC–NMR–MS in pharmaceutical research and development 748(2000)233

Phenamidine

Determination by capillary zone electrophoresis of berenil, phenamidine, diampron and dibromopropamidine in serum and urine 738(2000)293

Phenanthrene

Biomonitoring of environmental polycyclic aromatic hydrocarbon exposure by simultaneous measurement of urinary phenanthrene, pyrene and benzo[*a*]pyrene hydroxides 739(2000)225

1,10-Phenanthroline–copper(II) sulphate

Poly(ethylene glycol)-based aqueous biphasic systems: effect of temperature on phase equilibria and on partitioning of 1,10-phenanthroline–copper(II) sulphate complex 743(2000)231

Phenobarbital

Evaluation of the programmed temperature vaporiser for large-volume injection of biological samples in gas chromatography 729(1999)199

Immunoassay-based determination of phenobarbital using size-exclusion chromatography 749(2000)215

Phenolic flame-retardants

Determination of phenolic flame-retardants in human plasma using solid-phase extraction and gas chromatography–electron-capture mass spectrometry 750(2001)1

Phenothiazines

Gas chromatographic–mass spectrometric analysis of veterinary tranquilizers in urine: evaluation of method performance 734(1999)113

Phenprocoumon

Sensitive stereospecific determination of acenocoumarol and phenprocoumon in plasma by high-performance liquid chromatography 742(2000)131

Phenyl acetic acid mustard

Rapid determination of the anti-cancer drug chlorambucil (LeukeranTM) and its phenyl acetic acid mustard metabolite in human serum and plasma by automated solid-phase extraction and liquid chromatography–tandem mass spectrometry 732(1999)173

Phenylalanine

Neutral amino acids monitoring in phenylketonuric plasma microdialysates using micellar electrokinetic chromatography and laser-induced fluorescence detection 739(2000)247

Phenylbutazone

Study of the solid-phase extraction of diclofenac sodium, indomethacin and phenylbutazone for their analysis in human urine by liquid chromatography 729(1999)139

Simultaneous determination of hydrocortisone, dexamethasone, indomethacin, phenylbutazone and oxyphenbutazone in equine serum by high-performance liquid chromatography 738(2000)17

Phenylethylamines

Routine analysis of amphetamine class drugs as their naphthaquinone derivatives in human urine by high-performance liquid chromatography 735(1999)229

Phenylisothiocyanate

High-performance liquid chromatographic analysis of physiological amino acids in human brain tumors by pre-column derivatization with phenylisothiocyanate 730(1999)81

Phenylloxirane

Gas chromatography–electron capture determination of styrene-7,8-oxide enantiomers 749(2000)265

Phenytoin

Quantification of phenytoin and its metabolites in equine plasma and urine using high-performance liquid chromatography 746(2000)209

Pholedrine

High-performance liquid chromatographic determination of pholedrine in human serum using ion-pair extraction and amperometric detection 726(1999)309

Phosphatidylcholine

Quantitative determination of dipalmitoylphosphatidylcholine and palmitic acid in porcine lung surfactants used in the treatment of respiratory distress syndrome 744(2000)407

Phosphatidylinositol 4-kinase

Purification of rabbit lacrimal gland plasma membranes by aqueous two-phase affinity partitioning 743(2000)397

Phosphocreatine

Improved high-performance liquid chromatographic assay for the determination of “high-energy” phosphates in mammalian skeletal muscle. Application to a single-fibre study in man 730(1999)183

Phospholipids

Separation and quantification of cholesterol and major phospholipid classes in human semen by high-performance liquid chromatography and light-scattering detection 740(2000)101

Affinity partitioning of biotinylated mixed liposomes: effect of charge on biotin–NeutrAvidin interaction 743(2000)389

Quantitative determination of dipalmitoylphosphatidylcholine and palmitic acid in porcine lung surfactants used in the treatment of respiratory distress syndrome 744(2000)407

Effect of soybean oil and fish oil on individual molecular species of Atlantic salmon head kidney phospholipids determined by normal-phase liquid chromatography coupled to negative ion electrospray tandem mass spectrometry 748(2000)137

Phosphoramidate mustard

Analysis of cyclophosphamide and five metabolites from human plasma using liquid chromatography–mass spectrometry and gas chromatography–nitrogen–phosphorus detection 732(1999)287

Simple and selective determination of the cyclophosphamide metabolite phosphoramidate mustard in human plasma using high-performance liquid chromatography 745(2000)345

Photosystem I

Novel, rapid purification of the membrane protein photosystem I by high-performance liquid chromatography on porous materials 737(2000)131

Phycoerythrin

One-step purification of R-phycoerythrin from the red macroalga *Palmaria palmata* using preparative polyacrylamide gel electrophoresis 739(2000)117

High-performance liquid chromatography with on-line post-column immunoreaction detection of digoxin and its metabolites based on fluorescence energy transfer in the far-red spectral region 745(2000)305

Phytic acid

Phytic acid as an efficient low-molecular-mass displacer for anion-exchange displacement chromatography of proteins 741(2000)23

Phytosterols

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Pibutidine

Development and validation of a liquid chromatographic–tandem mass spectrometric method for the determination of pibutidine in human urine 740(2000)187

Pimobendan

Simple method for determination of the active metabolite of the inotropic drug pimobendan in rat liver microsomes 744(2000)189

Pipecolic acid

Simple method for the simultaneous analysis of pipecolic acid and lysine by high-performance liquid chromatography and its application to rumen liquor and plasma of ruminants 735(1999)63

Pirimicarb

Determination of metabolites of pirimicarb in human urine by gas chromatography–mass spectrometry 730(1999)229

Piroxicam

High-performance liquid chromatographic determination with amperometric detection of piroxicam in human plasma and tissues 729(1999)183

PK 11195

High-performance liquid chromatographic determination of [¹⁴C]1-(2-chlorophenyl)-*N*-methyl-*N*-(1-methylpropyl)-3-isoquinoline carboxamide in mouse plasma and tissue and in human plasma 736(1999)61

Plasma

Gas chromatographic–mass spectrometric determination of plasma saturated fatty acids using pentafluorophenyldimethylsilyl derivatization 742(2000)37

Pluronic F127

Application of Pluronic copolymer liquid crystals for the capillary electrophoretic separation of collagen type I cyanogen bromide fragments 739(2000)109

Polar metabolites

Use of directly coupled ion-exchange liquid chromatography–mass spectrometry and liquid chromatography–nuclear magnetic resonance spectroscopy as a strategy for polar metabolite identification 748(2000)295

Polyamines

Rapid and simultaneous high-performance liquid chromatography assay of polyamines and monoacetylpolyamines in biological specimens 745(2000)431

Polychlorinated biphenyls

Determination of polychlorinated biphenyls in human blood by solid-phase extraction including on-column lipid decomposition 734(1999)219

Polyclonal antibodies

Fractionation of polyclonal antibodies to fragments of a neuroreceptor using three increasingly chaotropic solvents 728(1999)49

Polyethylene

On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography 750(2001)61

Poly(ethylene glycol)–sodium citrate

Partial purification of penicillin acylase from *Escherichia coli* in poly(ethylene glycol)–sodium citrate aqueous two-phase systems 734(1999)15

Poly(ethylene glycol)

New aqueous two-phase system based on cashew-nut tree gum and poly(ethylene glycol) 743(2000)79

Solvatochromic studies in polyethylene glycol–salt aqueous biphasic systems 743(2000)137

Viscosity prediction of polyethylene glycol–dextran–water solutions used in aqueous two-phase systems 743(2000)181

Poly(ethylene glycol)-based aqueous biphasic systems: effect of temperature on phase equilibria and on partitioning of 1,10-phenanthroline–copper(II) sulphate complex 743(2000)231

Poly(ethylene oxide)

Improvements in DNA sequencing by capillary electrophoresis at elevated temperature using poly(ethylene oxide) as a sieving matrix 745(2000)221

Polymers

Separation and sensing based on molecular recognition using molecularly imprinted polymers 728(1999)1

Molecular imprinting for drug bioanalysis. A review on the application of imprinted polymers to solid-phase extraction and binding assay 739(2000)163

Polymer displacement/shielding in protein chromatography 741(2000)103

Analysis of compression of polymer mushrooms using self-consistent field theory 743(2000)31

Partitioning and diffusion of large molecules in fibrous structures 743(2000)169

Polynuclear aromatic compounds

Biomonitoring of environmental polycyclic aromatic hydrocarbon exposure by simultaneous measurement of urinary phenanthrene, pyrene and benzo[a]pyrene hydroxides 739(2000)225

Polyoma VP1

Purification of a viral coat protein by an engineered polyionic sequence 737(2000)77

Porcine haptoglobin

Simple high-performance liquid chromatographic purification procedure for porcine plasma haptoglobin 731(1999)395

Potassium

Capillary zone electrophoresis of potassium in human vitreous humour: validation of a new method 733(1999)273

On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography 750(2001)61

Praziquantel

Capillary electrophoresis–mass spectrometry, liquid chromatography–mass spectrometry and nanoelectrospray-mass spectrometry of praziquantel metabolites 748(2000)221

Pregnanolone

High-performance liquid chromatography of the neuroactive steroids alphaxalone and pregnanolone in plasma using dansyl hydrazine as fluorescent label: application to a pharmacokinetic-pharmacodynamic study in rats 745(2000)357

Probenecid

Complete two-dimensional separation for analysis of acidic compounds in plasma using column-switching reversed-phase liquid chromatography 726(1999)225

Procaine

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

Proenkephalin A

Mass spectrometric analysis of neuropeptidergic systems in the human pituitary and cerebrospinal fluid (Review) 731(1999)3

Proguanil

Simultaneous separation of atovaquone, proguanil and its metabolites on a mixed mode high-performance liquid chromatographic column 741(2000)189

Proopiomelanocortin

Mass spectrometric analysis of neuropeptidergic systems in the human pituitary and cerebrospinal fluid (Review) 731(1999)3

Propafenone

Determination of propafenone and its main metabolite 5-hydroxypropafenone in human serum with direct injection into a column-switching chromatographic system 726(1999)211

Determination of propafenone and its phase I and phase II metabolites in plasma and urine by high-performance liquid chromatography–electrospray ionization mass spectrometry 748(2000)113

Propitocaine

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

Propranolol

Two reproducible and sensitive liquid chromatographic methods to quantify atenolol and propranolol in human plasma and determination of their associated analytical error functions 738(2000)225

Direct high-performance liquid chromatographic determination of (*R*)- and (*S*)-propranolol in rat microdialysate using on-line column switching procedures 739(2000)151

Propranolol

Fast liquid chromatographic–mass spectrometric determination of pharmaceutical compounds 732(1999)445

Prostaglandin

Use of short high-performance liquid chromatography columns and tandem-mass spectrometry for the rapid analysis of a prostaglandin analog, fluprostenol, in rat plasma 741(2000)213

Prostaglandins

Simultaneous assay of prostaglandins and thromboxane in the cerebrospinal fluid by gas chromatography–mass spectrometry–selected ion monitoring 731(1999)73

Prostate-specific antigen

Serial lectin affinity chromatography demonstrates altered asparagine-linked sugar-chain structures of prostate-specific antigen in human prostate carcinoma 727(1999)9

Protease

New approach for separating *Bacillus subtilis* metalloprotease and α -amylase by affinity chromatography and for purifying neutral protease by hydrophobic chromatography 737(2000)277

Protease inhibitor

Sensitive liquid chromatographic assay for amprenavir, a human immunodeficiency virus protease inhibitor, in human plasma, cerebrospinal fluid and semen 742(2000)185

Protease inhibitors

Determination of twelve antiretroviral agents in human plasma sample using reversed-phase high-performance liquid

chromatography 744(2000)227

Protein-denaturing reagent

Quantitative liquid chromatographic–tandem mass spectrometric determination of reserpine in FVB/N mouse plasma using a “chelating” agent (disodium EDTA) for releasing protein-bound analytes during 96-well liquid–liquid extraction 742(2000)369

Protein G

Chromatographic behavior of mouse serum immunoglobulin G in protein G perfusion affinity chromatography 738(2000)149

Proteins

Erratum to “Chromatographic approaches to liposomes, proteoliposomes and biomembrane vesicles”. [J. Chromatogr. B, 722 (1999) 103–120] 727(1999)249

Effect of salt concentration on separation patterns in static capillary isoelectric focusing with imaging detection 729(1999)355

Mass spectrometric study on the protein chemical modification of uremic patients in advanced Maillard reaction 731(1999)131

Aptamer affinity chromatography: combinatorial chemistry applied to protein purification 731(1999)275

Enzyme-linked immunosorbent assay-based selection and optimization of elution buffer for TAG72-affinity chromatography 731(1999)299

Simultaneous determination of enantioselective plasma protein binding of aminohydantoins by ultrafiltration and chiral high-performance liquid chromatography 732(1999)31

Three-step purification of a fragment of the large immunophilin FKBP52 737(2000)71

Novel, rapid purification of the membrane protein photosystem I by high-performance liquid chromatography on porous materials 737(2000)131

Three-step chromatographic purification procedure for the production of a His-tag recombinant kinesin overexpressed in *E. coli* 737(2000)143

High-performance affinity chromatography and immobilized serum albumin as probes for drug- and hormone-protein binding 739(2000)39

Application of Pluronic copolymer liquid crystals for the capillary electrophoretic separation of collagen type I cyanogen bromide fragments 739(2000)109

Bisphenol derivative of allysine for high-performance liquid chromatographic analysis of allysine residue of proteins 739(2000)273

New strategy for the design of ligands for the purification of pharmaceutical proteins by affinity chromatography 740(2000)17

Phytic acid as an efficient low-molecular-mass displacer for anion-exchange displacement chromatography of proteins 741(2000)23

Capillary electrophoretic separation of proteins and peptides using Pluronic liquid crystals and surface-modified capillaries 741(2000)37

Polymer displacement/shielding in protein chromatography 741(2000)103

Separation and detection techniques for peptides and proteins in stability research and bioanalysis 742(2000)401

Stimuli-responsive separation of proteins using immobilized liposome chromatography 743(2000)85

Partitioning and diffusion of large molecules in fibrous structures 743(2000)169

Conformational transition and mass transfer in extraction of proteins by AOT–alcohol–isooctane reverse micellar systems 743(2000)203

Partitioning of whey proteins, bovine serum albumin and porcine insulin in aqueous two-phase systems 743(2000)235

Phase behavior and protein partitioning in aqueous two-phase systems of cationic–anionic surfactant mixtures 743(2000)327

Strategy for qualitative and quantitative analysis in proteomics based on signature peptides 745(2000)197

Integrated microchip-device for the digestion, separation and postcolumn labeling of proteins and peptides 745(2000)243

Use of non-porous reversed-phase high-performance liquid chromatography for protein profiling and isolation of proteins induced by temperature variations for Siberian permafrost bacteria with identification by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry and capillary electrophoresis–electrospray ionization mass spectrometry 748(2000)167

On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography 750(2001)61

Proteoliposomes

Erratum to ‘‘Chromatographic approaches to liposomes, proteoliposomes and biomembrane vesicles’’. [J. Chromatogr. B, 722 (1999) 103–120] 727(1999)249

Proteome

Reversed-phase high-performance liquid chromatographic assay for the adenovirus type 5 proteome 732(1999)411

Pseudohypericin

Determination of hypericin and pseudohypericin in pharmaceutical preparations by liquid chromatography with fluorescence detection 749(2000)57

Pullulanase

Pullulanase from the hyperthermophilic bacterium *Thermotoga maritima*: purification by β -cyclodextrin affinity chromatography 737(2000)245

Purine

Comparative determination of purine compounds in carotid plaque by capillary zone electrophoresis and high-performance liquid chromatography 728(1999)185

Pyrene

Extraction and high-performance liquid chromatographic separation of selected pyrene and benzo[*a*]pyrene sulfates and glucuronides: preliminary application to the analysis of smokers’ urine 728(1999)209

Biomonitoring of environmental polycyclic aromatic hydrocarbon exposure by simultaneous measurement of urinary phenanthrene, pyrene and benzo[*a*]pyrene hydroxides 739(2000)225

1-Pyrenyldiazomethane

Headspace solid-phase microextraction with 1-pyrenyldiazomethane in-fibre derivatisation for analysis of faecal short-chain fatty acids 730(1999)113

Pyridostigmine bromide

Simultaneous determination of pyridostigmine bromide, *N,N*-diethyl-*m*-toluamide, permethrin, and their metabolites in rat plasma and urine by high-performance liquid chromatography 749(2000)171

Pyroglutamate

Analysis of diagnostic metabolites by capillary electrophoresis–mass spectrometry 727(1999)43

Pyrroloquinoline quinone

Determination of pyrroloquinoline quinone by capillary zone electrophoresis 739(2000)101

Pyruvate kinase

Arsanilic acid–Sepharose chromatography of pyruvate kinase from KB cells 740(2000)109

Q

Quantitative analysis

Simultaneous determination of bosentan and its three major metabolites in various biological matrices and species using narrow bore liquid chromatography with ion spray tandem mass spectrometric detection 749(2000)67

Quercetin

Validated method for the quantitation of quercetin from human plasma using high-performance liquid chromatography with electrochemical detection 727(1999)179

High-performance liquid chromatographic determination of quercetin and isorhamnetin in rat tissues using β -glucuronidase and acid hydrolysis 738(2000)413

Quinamide

Solid-phase extraction and liquid chromatographic quantitation of quinamide in biological samples 746(2000)133

Quinolate phosphoribosyltransferase

Reversed-phase high-performance liquid chromatography of nicotinic acid mononucleotide for measurement of quinolate phosphoribosyltransferase 749(2000)281

Quinolones

Determination of quinolones in plasma samples by capillary electrophoresis using solid-phase extraction 742(2000)255

R

Raclopride

Determination of raclopride in human plasma by on-column focusing packed capillary liquid chromatography–electrospray ionisation mass spectrometry 748(2000)189

Ranitidine

Automated in-tube solid-phase microextraction–liquid chromatography–electrospray ionization mass spectrometry for the determination of ranitidine 731(1999)353

Rat

Separation and quantitation of metallothionein isoforms from liver of untreated rats by ion-exchange high-performance liquid chromatography and atomic absorption spectrometry 744(2000)273

Reboxetine

Automated determination of reboxetine by high-performance liquid chromatography with column-switching and ultraviolet detection 740(2000)135

Recombinant adeno-associated virus

Cation-exchange high-performance liquid chromatography of recombinant adeno-associated virus type 2 740(2000)195

Recombinant hepatitis B surface antigen

Comparison of different elution conditions for the immunopurification of recombinant hepatitis B surface antigen 735(1999)271

Recombinant human basic fibroblast growth factor

Preparative two-step purification of recombinant human basic fibroblast growth factor from high-cell-density cultivation of *Escherichia coli* 737(2000)25

Red cell membrane proteins

Immunopurification of the blood group RhD protein from human erythrocyte membranes 735(1999)207

Remacemide

Simultaneous determination of remacemide hydrochloride and desglycylremacemide (AR-R12495XX) in brain tissue by high-performance liquid chromatography 746(2000)325

Renin inhibitor

Automated quantitative determination of the new renin inhibitor CGP 60536 by high-performance liquid chromatography 738(2000)129

Reproterol

Excretion study of the β_2 -agonist reproterol in human urine 726(1999)141

Reserpine

Quantitative liquid chromatographic–tandem mass spectrometric determination of reserpine in FVB/N mouse plasma using a “chelating” agent (disodium EDTA) for releasing protein-bound analytes during 96-well liquid–liquid extraction 742(2000)369

Resorcinol

Naphthol- and resorcinol-based azo dyes as metal ion complexants in aqueous biphasic systems 743(2000)107

Restricted access

Enantioselective analysis of (*R*)- and (*S*)-atenolol in urine samples by a high-performance liquid chromatography column-switching setup 740(2000)219

Resveratrol

Liquid chromatography with multichannel electrochemical detection for the determination of *trans*-resveratrol in rat blood utilizing an automated blood sampling device 740(2000)129

Retigabine

Determination of retigabine and its acetyl metabolite in biological matrices by on-line solid-phase extraction (column switching) liquid chromatography with tandem mass spectrometry 748(2000)97

Retinol

Stability of individual carotenoids, retinol and tocopherols in human plasma during exposure to light and after extraction 729(1999)191

Simultaneous high-performance liquid chromatographic determination of retinol by fluorometry and of tocopherol by ultraviolet absorbance in the serum of newborns 729(1999)231

Simultaneous determination of serum retinol and α - and γ -tocopherol levels in type II diabetic patients using high-performance liquid chromatography with fluorescence detection 730(1999)265

Retroviruses

Recovery in aqueous two-phase systems of nanoparticulates applied as surrogate mimics for viral gene therapy vectors 743(2000)409

Reviews

Erratum to “Chromatographic approaches to liposomes, proteoliposomes and biomembrane vesicles”. [J. Chromatogr. B, 722 (1999) 103–120] 727(1999)249

RH1

Development and validation of a sensitive solid-phase extraction and high-performance liquid chromatographic assay for the novel bio-reductive anti-tumor agent RH1 in human and mouse plasma 729(1999)287

RhD

Immunopurification of the blood group RhD protein from human erythrocyte membranes 735(1999)207

Rhodanese

Rhodanese (thiosulfate:cyanide sulfurtransferase) from frog *Rana temporaria* 746(2000)315

Riboflavin

Analysis of riboflavin and riboflavin cofactor levels in plasma by high-performance liquid chromatography 739(2000)219

Riboflavin binding protein

Chromatographic investigation on the binding site characteristics of quail egg-white riboflavin binding protein as a chiral stationary phase 738(2000)343

Ribonucleosides

Detection and characterization of modified nucleosides in serum and urine of uremic patients using capillary liquid chromatography–frit-fast atom bombardment mass spectrometry 746(2000)51

Rifabutin

Specific high-performance liquid chromatography assay for determination of rifabutin plasma concentration following Extrelut column extraction 728(1999)233

Risperidone

Determination of risperidone and its major metabolite 9-hydroxyrisperidone in human plasma by reversed-phase liquid chromatography with ultraviolet detection 746(2000)173

Ritonavir

Simultaneous determination of the HIV-protease inhibitors indinavir, nelfinavir, saquinavir and ritonavir in human plasma by reversed-phase high-performance liquid chromatography 727(1999)139

High-performance liquid chromatographic method for the simultaneous determination of HIV-1 protease inhibitors indinavir, saquinavir and ritonavir in plasma of patients during highly active antiretroviral therapy 735(1999)41

Simultaneous determination of the HIV protease inhibitors indinavir, amprenavir, saquinavir, ritonavir, nelfinavir and the non-nucleoside reverse transcriptase inhibitor efavirenz by high-performance liquid chromatography after solid-phase extraction 740(2000)43

Simultaneous high-performance liquid chromatographic determination of the antiretroviral agents amprenavir, nelfinavir, ritonavir, saquinavir, delavirdine and efavirenz in human plasma 742(2000)453

RNA

Rapid method for relative gene expression determination in human tissues using automated capillary gel electrophoresis and multicolor detection 734(1999)47

Simultaneous accumulation of low-molecular-mass RNA at the interface along with accumulation of high-molecular-mass RNA on aqueous two-phase system partitioning 743(2000)421

RNA-aptamers

Purification of a highly modified RNA-aptamer. Effect of complete denaturation during chromatography on product recovery and specific activity 726(1999)237

Rodenticides

Use of an ion-pairing reagent for high-performance liquid chromatography–atmospheric pressure chemical ionization mass spectrometry determination of anionic anticoagulant rodenticides in body fluids 731(1999)155

Rofecoxib

Determination of rofecoxib, a cyclooxygenase-2 specific inhibitor, in human plasma using high-performance liquid chromatography with post-column photochemical derivatization and fluorescence detection 730(1999)221

Determination of Rofecoxib (MK-0966), a cyclooxygenase-2 inhibitor, in human plasma by high-performance liquid chromatography with tandem mass spectrometric detection 748(2000)31

Ropinirole

Separation and quantification of ropinirole and some impurities using capillary liquid chromatography 732(1999)437

Ropivacaine

Determination of free concentrations of ropivacaine and bupivacaine in plasma from neonates using small-scale equilibrium–dialysis followed by liquid chromatography–mass spectrometry 748(2000)321

Roxithromycin

Determination of erythromycin, clarithromycin, roxithromycin, and azithromycin in plasma by high-performance liquid chromatography with amperometric detection 738(2000)405

Determination of roxithromycin residues in the flounder muscle with electrospray liquid chromatography–mass spectrometry 746(2000)219

Ro 64-0802

Development of a high-performance liquid chromatographic–mass spectrometric assay for the specific and sensitive quantification of Ro 64-0802, an anti-influenza drug, and its pro-drug, oseltamivir, in human and animal plasma and urine 745(2000)373

S

S 16257

Simultaneous determination of ivabradine and its metabolites in human plasma by liquid chromatography–tandem mass spectrometry 745(2000)261

S100A1

Purification of the Ca²⁺-binding protein S100A1 from myocardium and recombinant *Escherichia coli* 737(2000)39

Salbutamol

Automated protein precipitation by filtration in the 96-well format 734(1999)257

Salmeterol

Rapid chiral high-performance liquid chromatographic assay for salmeterol and α -hydroxysalmeterol. Application to in vitro metabolism studies 729(1999)225

Salsolinol

Quantitative determination of endogenous tetrahydroisoquinoline salsolinol in peripheral blood mononuclear cells by gas chromatography–mass spectrometry 735(1999)299

Saquinavir

Simultaneous determination of the HIV-protease inhibitors indinavir, nelfinavir, saquinavir and ritonavir in human plasma by reversed-phase high-performance liquid chromatography 727(1999)139

High-performance liquid chromatographic method for the simultaneous determination of HIV-1 protease inhibitors indinavir, saquinavir and ritonavir in plasma of patients during highly active antiretroviral therapy 735(1999)41

Simultaneous determination of the HIV protease inhibitors indinavir, amprenavir, saquinavir, ritonavir, nelfinavir and the non-nucleoside reverse transcriptase inhibitor efavirenz by high-performance liquid chromatography after solid-phase extraction 740(2000)43

Simultaneous high-performance liquid chromatographic determination of the antiretroviral agents amprenavir, nelfinavir, ritonavir, saquinavir, delavirdine and efavirenz in human plasma 742(2000)453

SCH 56592

High-performance liquid chromatographic analysis of the anti-fungal agent SCH 56592 in dog serum 738(2000)93

SCH 66336

High-performance liquid chromatographic analysis of the anti-tumor agent SCH 66336 in cynomolgus monkey plasma and evaluation of its chiral inversion in animals 728(1999)133

SCH 66712

High-performance liquid chromatographic analysis of the D₄ receptor antagonist SCH 66712 in rat plasma 735(1999)11

Scopolamine

Determination of scopolamine in human serum and microdialysis samples by liquid chromatography–tandem mass spectrometry 750(2001)121

SDF-1 α

High-performance liquid chromatographic purification and capillary electrophoresis quantification of the chemokine stromal cell-derived factor-1 729(1999)369

SDZ-RAD

Automated, fast and sensitive quantification of drugs in blood by liquid chromatography–mass spectrometry with on-line extraction: immunosuppressants 748(2000)41

Sebacoyl dinalbuphine ester

High-performance liquid chromatographic method for the simultaneous determination of nalbuphine and its prodrug, sebacoyl dinalbuphine ester, in dog plasma and application to pharmacokinetic studies in dogs 746(2000)241

Secobarbital

Evaluation of the programmed temperature vaporiser for large-volume injection of biological samples in gas chromatography 729(1999)199

L-Selectin

Aptamer affinity chromatography: combinatorial chemistry applied to protein purification 731(1999)275

Selegiline

Determination of enantiomeric amphetamines as metabolites of illicit amphetamines and selegiline in urine by capillary electrophoresis using modified β -cyclodextrin 741(2000)221

Selenoprotein P

Comparison of different transition metal ions for immobilized metal affinity chromatography of selenoprotein P from human plasma 735(1999)85

Seocalcitol

Determination of the vitamin D analog EB 1089 (seocalcitol) in human and pig serum using liquid chromatography–tandem mass spectrometry 740(2000)117

Serotonin

Simultaneous determination of catecholamines in rat brain tissue by high-performance liquid chromatography 731(1999)361

Methods for the determination of seven selective serotonin reuptake inhibitors and three active metabolites in human serum using high-performance liquid chromatography and gas chromatography 742(2000)229

Clinical chemistry of serotonin and metabolites 747(2000)33

Quantitative determination of L-775,606, a selective 5-hydroxytryptamine 1D agonist, in rat plasma using automated sample preparation and detection by liquid chromatography–tandem mass spectrometry 750(2001)13

Serralyisin

Use of liquid chromatography–mass spectrometry coupling for monitoring the serralyisin-catalyzed hydrolysis of a peptide library 732(1999)271

Sesquiterpenoids

Qualitative and quantitative determination of sesquiterpenoids in *Achillea* species by reversed-phase high-performance liquid chromatography, mass-spectrometry and thin-layer chromatography 729(1999)361

Shikonin

Simultaneous analysis of shikimate-derived secondary metabolites in *Lithospermum erythrorhizon* cell suspension cultures by high-performance liquid chromatography 738(2000)3

Short-chain

Routine analysis of short-chain fatty acids for anaerobic bacteria identification using capillary electrophoresis and indirect ultraviolet detection 741(2000)89

Short-chain fatty acids

Headspace solid-phase microextraction with 1-pyrenyldiazomethane in-fibre derivatisation for analysis of faecal short-chain fatty acids 730(1999)113

Signature peptides

Strategy for qualitative and quantitative analysis in proteomics based on signature peptides 745(2000)197

Silicone

High-performance liquid chromatographic determination of 17 β -estradiol and 17 β -estradiol-3-acetate solubilities and diffusion coefficients in silicone elastomeric intravaginal rings 744(2000)157

Single-chain Fv antibodies

Three-step purification of bacterially expressed human single-chain Fv antibodies for clinical applications 737(2000)107

Sirolimus

Automated, fast and sensitive quantification of drugs in blood by liquid chromatography–mass spectrometry with on-line extraction: immunosuppressants 748(2000)41

Sitostanol

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Sitosterol

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Size-exclusion chromatography

Immunoassay-based determination of phenobarbital using size-exclusion chromatography 749(2000)215

Small sample volume

Analysis of small-scale biological compartments by capillary electrophoresis 745(2000)127

SN-38

Sensitive determination of irinotecan (CPT-11) and its active metabolite SN-38 in human serum using liquid chromatography–electrospray mass spectrometry 736(1999)175

Simple and rapid determination of irinotecan and its metabolite SN-38 in plasma by high-performance liquid-chromatography: application to clinical pharmacokinetic studies 740(2000)159

Sodium

On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography 750(2001)61

Sodium artesunate

Determination of sodium artesunate in plasma using ion-pairing high-performance liquid chromatography 744(2000)415

Sodium chloride

Bovine serum albumin partitioning in an aqueous two-phase system: Effect of pH and sodium chloride concentration 743(2000)255

Sodium sulphate

Drowning-out crystallisation of sodium sulphate using aqueous two-phase systems 743(2000)101

Solid-phase extraction

A sensitive high-performance liquid chromatographic method using electrochemical detection for the analysis of olanzapine and desmethylolanzapine in plasma of schizophrenic patients using a new solid-phase extraction procedure 750(2001)137

Soybean peroxidase

Purification of soybean peroxidase (*Glycine max*) by metal affinity partitioning in aqueous two-phase systems 743(2000)287

Soy protein

Isoelectric precipitation of soybean protein using carbon dioxide as a volatile acid 743(2000)357

Specific serotonin reuptake inhibitors

Methods for the determination of seven selective serotonin reuptake inhibitors and three active metabolites in human serum using high-performance liquid chromatography and gas chromatography 742(2000)229

Sphingolipidosis

Application of delayed extraction matrix-assisted laser desorption ionization time-of-flight mass spectrometry for analysis of sphingolipids in tissues from sphingolipidosis patients 731(1999)45

Sphingomyelin

Quantitative determination of dipalmitoylphosphatidylcholine and palmitic acid in porcine lung surfactants used in the treatment of respiratory distress syndrome 744(2000)407

Stanols

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Stanozolol

Testing of the anabolic stanozolol in human hair by gas chromatography–negative ion chemical ionization mass spectrometry 740(2000)265

Stavudine

Determination of stavudine in human plasma and urine by high-performance liquid chromatography using a reduced sample volume 746(2000)183

Steroids

Supercritical fluid extraction of methyltestosterone, nortestosterone and testosterone at low ppb levels from fortified bovine urine 726(1999)121

Isolation and identification of the C₆-hydroxy and C₂₀-hydroxy metabolites and glucuronide conjugate of methylprednisolone by preparative high-performance liquid chromatography from urine of patients receiving high-dose pulse therapy 726(1999)157

17 α -Ethyl-5 β -estrane-3 α ,17 β -diol, a biological marker for the abuse of norethandrolone and ethylestrenol in slaughter cattle 728(1999)217

Development of a urinary free cortisol assay using solid-phase extraction–capillary electrophoresis 730(1999)123

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Gas chromatographic–tandem mass spectrometric determination of anabolic steroids and their esters in hair. Application in doping control and meat quality control 735(1999)189

Simultaneous determination of endogenous and ¹³C-labelled cortisols and cortisones in human plasma by stable isotope dilution mass spectrometry 738(2000)119

Stable isotope dilution analysis of human urinary metabolites of 17 α -methyltestosterone 741(2000)271

High-performance liquid chromatography of the neuroactive steroids alphaxalone and pregnanolone in plasma using dansyl hydrazine as fluorescent label: application to a pharmacokinetic–pharmacodynamic study in rats 745(2000)357

Quantification of urine 17-ketosteroid sulfates and glucuronides by high-performance liquid chromatography–ion trap mass spectroscopy 750(2001)81

Stigmasterol

Precise quantitative determination of phytosterols, stanols, and cholesterol metabolites in human serum by capillary gas–liquid chromatography 732(1999)17

Styrene

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Purge-and-trap gas chromatographic determination of styrene in urine and blood. Application to exposed workers 741(2000)301

Gas chromatography-electron capture determination of styrene-7,8-oxide enantiomers 749(2000)265

Substance P

Assay of ezlopitant, a substance P receptor antagonist, and metabolites in biological matrices by gas chromatography with mass spectrometric detection: simultaneous analysis of a benzyl alcohol and alkene 744(2000)333

Narrow-bore high-performance liquid chromatography in combination with ionspray tandem mass spectrometry for the determination of the substance P receptor antagonist ezlopitant and its two active metabolites in plasma 748(2000)179

Succinyladenosine

Identification and determination of succinyladenosine in human cerebrospinal fluid 726(1999)53

Sugar alcohols

Gas chromatographic–mass spectrometric analysis of urinary sugar and sugar alcohols during pregnancy 731(1999)111

Sugars

Gas chromatographic–mass spectrometric analysis of urinary sugar and sugar alcohols during pregnancy 731(1999)111

Sulfated β -cyclodextrin

Non-aqueous capillary electrophoresis chiral separations with sulfated β -cyclodextrin 731(1999)187

Charged cyclodextrin-mediated sample stacking in micellar capillary electrophoresis. A simple method for enhancing the detection sensitivity of hydrophobic compounds 731(1999)369

Sulfonamide drugs

Simultaneous determination of bosentan and its three major metabolites in various biological matrices and species using narrow bore liquid chromatography with ion spray tandem mass spectrometric detection 749(2000)67

Sulfonamides

Fast liquid chromatographic–mass spectrometric determination of pharmaceutical compounds 732(1999)445

Sulphonamides

Screening of sulphonamides in egg using gas chromatography–mass-selective detection and liquid chromatography–mass spectrometry 729(1999)127

SYN-2836

Novel sample preparation method facilitating identification of urinary drug metabolites by liquid chromatography–tandem mass spectrometry 738(2000)259

S-1090

Determination of a new oral cephalosporin, S-1090, in human plasma and urine by direct injection high-performance liquid chromatography with ultraviolet detection and column switching 728(1999)125

T**Tachykinin**

Mass spectrometric analysis of neuropeptidergic systems in the human pituitary and cerebrospinal fluid (Review) 731(1999)3

Tacrolimus

Automated, fast and sensitive quantification of drugs in blood by liquid chromatography–mass spectrometry with on-line extraction: immunosuppressants 748(2000)41

TAG72

Enzyme-linked immunosorbent assay-based selection and optimization of elution buffer for TAG72-affinity chromatography 731(1999)299

Taurine

Ion chromatographic determination of taurine in medicine, nutrient capsule and human urine with electrochemical detection 730(1999)161

On-line derivatization for continuous and automatic monitoring of brain extracellular glutamate levels in anesthetized rats: a microdialysis study 734(1999)1

Taxane

High-performance liquid chromatographic assay for the determination of the novel taxane derivative IDN5109 in mouse plasma 736(1999)135

Taxol

Poisoning by plant material: review of human cases and analytical determination of main toxins by high-performance liquid chromatography–(tandem) mass spectrometry 733(1999)181

TCV-116

Determination of candesartan cilexetil, candesartan and a metabolite in human plasma and urine by liquid chromatography and fluorometric detection 731(1999)411

Tear fluid

Analysis of small-scale biological compartments by capillary electrophoresis 745(2000)127

Terazosin

Improved high-performance liquid chromatographic analysis of terazosin in human plasma 745(2000)439

Terbutaline

Simple method for determination of terbutaline plasma concentration by high-performance liquid chromatography 741(2000)307

Terfenadine

Analysis of hydroxylated and *N*-dealkylated metabolites of terfenadine in microsomal incubates by liquid chromatography–mass spectrometry 741(2000)145

Terfenadine alcohol

Analysis of hydroxylated and *N*-dealkylated metabolites of terfenadine in microsomal incubates by liquid chromatography–mass spectrometry 741(2000)145

Testosterone

Supercritical fluid extraction of methyltestosterone, nortestosterone and testosterone at low ppb levels from fortified bovine urine 726(1999)121

Determination of testosterone and 6 β -hydroxytestosterone by gas chromatography–selected ion monitoring–mass spectrometry for the characterization of cytochrome p450 3a activity 734(1999)73

Testosterone esters

Gas chromatographic–tandem mass spectrometric determination of anabolic steroids and their esters in hair. Application in doping control and meat quality control 735(1999)189

2,3,4,6-Tetra-*O*-acetyl- β -D-glycopyranosyl isothiocyanate

Reversed-phase high-performance liquid chromatographic analysis of atenolol enantiomers in rat hepatic microsome after chiral derivatization with 2,3,4,6-tetra-*O*-acetyl- β -D-glycopyranosyl isothiocyanate 742(2000)433

Tetracaine

Simultaneous determination of local anesthetics including ester-type anesthetics in human plasma and urine by gas chromatography–mass spectrometry with solid-phase extraction 726(1999)185

Tetracycline

Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments. Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry 748(2000)369

Tetracyclines

Simultaneous determination of residual tetracyclines in foods by high-performance liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry 732(1999)55

Automated residue analysis of tetracyclines and their metabolites in whole egg, egg white, egg yolk and hen's plasma utilizing a modified ASTED system 739(2000)191

2,3,5,6-Tetrafluoro-4-trifluoromethylaniline

Identification and quantification of metabolites of 2,3,5,6-tetrafluoro-4-trifluoromethylaniline in rat urine using ^{19}F nuclear magnetic resonance spectroscopy, high-performance liquid chromatography–nuclear magnetic resonance spectroscopy and high-performance liquid chromatography–mass spectrometry 748(2000)311

Tetrahydrocannabinol

On-line coupling of automated solid-phase extraction with high-performance liquid chromatography and electrochemical detection. Quantitation of oxidizable drugs of abuse and their metabolites in plasma and urine 731(1999)167

Tetrahydrocannabinolic acid

On-line coupling of automated solid-phase extraction with high-performance liquid chromatography and electrochemical detection. Quantitation of oxidizable drugs of abuse and their metabolites in plasma and urine 731(1999)167

Tetrahydroisoquinoliner

Quantitative determination of endogenous tetrahydroisoquinoline salsolinol in peripheral blood mononuclear cells by gas chromatography–mass spectrometry 735(1999)299

1,2,3,4-Tetrahydroisoquinolines

Development of a method for sample preparation for subsequent identification and measurement of 1,2,3,4-tetrahydroisoquinolines and other potentially neurotoxic compounds by high-performance liquid chromatography with ultraviolet and fluorescence detection in blood plasma of Parkinson's disease patients 746(2000)283

High-performance liquid chromatographic separation and measurement of various biogenic compounds possibly involved in the pathomechanism of Parkinson's disease 746(2000)297

Tetrahydrothiophene

Determination of tetrahydrothiophene formation as a probe of in vitro busulfan metabolism by human glutathione *S*-transferase A1-1: use of a highly sensitive gas chromatographic–mass spectrometric method 730(1999)25

Texaphyrin

Sensitive high-performance liquid chromatographic assay for motexafin gadolinium and motexafin lutetium in human plasma 749(2000)145

Thalidomide

Quantitative determination of thalidomide in human serum with high-performance liquid chromatography using protein precipitation with trichloroacetic acid and ultraviolet detection 734(1999)203

Theophylline

High-performance thin-layer chromatographic determination of theophylline in plasma 736(1999)289

Therapeutic drugs

Hair analysis for abused and therapeutic drugs 733(1999)161

Thermoseparating polymers

Model process for separation based on unfolding and refolding of chymotrypsin inhibitor 2 in thermoseparating polymer two-phase systems 743(2000)13

Polymer recycling in aqueous two-phase extractions using thermoseparating ethylene oxide–propylene oxide copolymers 743(2000)115

Partitioning of peptides and recombinant protein–peptide fusions in thermoseparating aqueous two-phase systems: effect of peptide primary structure 743(2000)295

2-Thiazolidinethione-4-carboxylic acid

Determination of urinary 2-thiazolidinethione-4-carboxylic acid after exposure to alkylene bisdithiocarbamates using gas chromatography–mass spectrometry 726(1999)85

Thiocoraline

Bioanalysis of thiocoraline, a new marine antitumoral depsipeptide, in plasma by high-performance liquid chromatography and fluorescence detection 726(1999)255

Thiocyanates

High-performance liquid chromatographic determination of cotinine in urine in isocratic mode 746(2000)115

Thiols

Determination of reduced and oxidized homocysteine and related thiols in plasma by thiol-specific pre-column derivatization and capillary electrophoresis with laser-induced fluorescence detection 735(1999)219

Electrospray ionization mass spectrometry of low-molecular-mass *S*-nitroso compounds and their thiols 742(2000)99

On-column derivatization for the analysis of homocysteine and other thiols by capillary electrophoresis with laser-induced fluorescence detection 744(2000)149

Thiopurine methyl transferase

Thiopurine methyl transferase activity: new extraction conditions for high-performance liquid chromatographic assay 727(1999)235

Thioredoxin

Purification of the membrane binding domain of cytochrome *b*₅ by immobilised nickel chelate chromatography 737(2000)119

Thioredoxin reductase

Expression of chymotrypsin(ogen) in the thioredoxin reductase deficient mutant strain of *Escherichia coli* AD494(DE3) and purification via a fusion product with a hexahistidine-tail 737(2000)213

Thiosulfate:cyanide sulfurtransferase

Rhodanese (thiosulfate:cyanide sulfurtransferase) from frog *Rana temporaria* 746(2000)315

ThioTEPA-mercapturate

Liquid chromatographic–mass spectrometric determination of the novel, recently identified thioTEPA metabolite, thioTEPA-mercapturate, in urine 732(1999)73

Threonine

Neutral amino acids monitoring in phenylketonuric plasma microdialysates using micellar electrokinetic chromatography and laser-induced fluorescence detection 739(2000)247

Convenient method of threonine, methionine and their related amino compounds by high-performance liquid chromatography and its application to rumen fluid 741(2000)279

Thrombin inhibitor

Determination of a novel thrombin inhibitor in human plasma and urine utilizing liquid chromatography with tandem mass spectrometric and ultraviolet detection 738(2000)83

Thromboxane

Simultaneous assay of prostaglandins and thromboxane in the cerebrospinal fluid by gas chromatography–mass spectrometry–selected ion monitoring 731(1999)73

Thymine

Derivatization of thymine and thymine photodimers with 4-bromomethyl-7-methoxycoumarin for fluorescence detection in high-performance liquid chromatography 731(1999)179

Tianeptine

High-performance liquid chromatographic determination of tianeptine in plasma applied to pharmacokinetic studies 748(2000)407

α-Tocopherol

Simple high-performance liquid chromatographic method for determination of α-tocopherol in human plasma 732(1999)227

Tocopherol

Stability of individual carotenoids, retinol and tocopherols in human plasma during exposure to light and after extraction 729(1999)191

Simultaneous high-performance liquid chromatographic determination of retinol by fluorometry and of tocopherol by ultraviolet absorbance in the serum of newborns 729(1999)231

Simultaneous determination of serum retinol and α - and γ -tocopherol levels in type II diabetic patients using high-performance liquid chromatography with fluorescence detection 730(1999)265

Tocotrienols

Simple high-performance liquid chromatographic method for the determination of tocotrienols in human plasma 735(1999)279

Tolcapone

Determination of the catechol-*O*-methyltransferase inhibitor tolcapone and three of its metabolites in animal and human plasma and urine by reversed-phase high-performance liquid chromatography with ultraviolet detection 730(1999)153

Direct analysis of nitrocatechol-type glucuronides in urine by capillary electrophoresis–electrospray ionisation mass spectrometry and tandem mass spectrometry 749(2000)253

Tolfenamic acid

Simultaneous analysis of anthranilic acid derivatives in pharmaceuticals and human urine by high-performance liquid chromatography with isocratic elution 744(2000)81

Tolterodine

Capillary solid-phase extraction–tandem mass spectrometry for fast quantification of free concentrations of tolterodine and two metabolites in ultrafiltered plasma samples 736(1999)247

Toluene

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Toluidines

Gas chromatographic method for the determination of toluidines in spiked urine samples 738(2000)427

Topotecan

Quantification of topotecan and its metabolite *N*-desmethyltopotecan in human plasma, urine and faeces by high-performance liquid chromatographic methods 727(1999)191

Toxic proteins

Polymeric adsorbent for removing toxic proteins from blood of patients with kidney failure 739(2000)73

Tramadol

Determination of tramadol by capillary gas chromatography with flame ionization detection. Application to human and rabbit pharmacokinetic studies 736(1999)89

Assay of tramadol in urine by capillary electrophoresis using laser-induced native fluorescence detection 745(2000)271

Enantiomeric determination of tramadol and its main metabolite *O*-desmethyltramadol in human plasma by liquid chromatography–tandem mass spectrometry 748(2000)65

Tramadol glucuronide

Direct determination of tramadol glucuronides in human urine by high-performance liquid chromatography with fluorescence detection 732(1999)185

Transferrin

Dynamically-coated capillaries allow for capillary electrophoretic resolution of transferrin sialoforms via direct analysis of human serum 742(2000)79

Transferrins

Improved method for carbohydrate-deficient transferrin determination in human serum by capillary zone electrophoresis 739(2000)81

Transglutaminase

Rapid, single-step procedure for the identification of transglutaminase-mediated isopeptide crosslinks in amino acid digests 732(1999)65

Transition metal ions

Comparison of different transition metal ions for immobilized metal affinity chromatography of selenoprotein P from human plasma 735(1999)85

Trenbolone

Determination of α - and β -trenbolone in bovine muscle and liver by liquid chromatography with fluorescence detection 739(2000)363

Triacylglycerol

Application of atmospheric pressure chemical ionization liquid chromatography–mass spectrometry in identification of lymph triacylglycerols 748(2000)425

Trichloroethylene

Simple method for rapid measurement of trichloroethylene and its major metabolites in biological samples 732(1999)145

Tricyclic amines

Fast liquid chromatographic–mass spectrometric determination of pharmaceutical compounds 732(1999)445

***N,N',N''*-Triethylenephosphoramide**

Liquid chromatographic–mass spectrometric determination of the novel, recently identified thioTEPA metabolite, thioTEPA-mercaptopurine, in urine 732(1999)73

Simultaneous determination of thioTEPA, TEPA and a novel, recently identified thioTEPA metabolite, monochloroTEPA, in urine using capillary gas chromatography 742(2000)335

***N,N',N''*-Triethylenethiophosphoramide**

Liquid chromatographic–mass spectrometric determination of the novel, recently identified thioTEPA metabolite, thioTEPA-mercaptopurine, in urine 732(1999)73

Simultaneous determination of thioTEPA, TEPA and a novel, recently identified thioTEPA metabolite, monochloroTEPA, in urine using capillary gas chromatography 742(2000)335

 ϵ -Trimethyllysine

Measurement of carnitine precursors, ϵ -trimethyllysine and γ -butyrobetaine in human serum by tandem mass spectrometry 731(1999)89

Trimethyl-*N*-oxide

Sensitive assay of trimethylamine *N*-oxide in liver microsomes by headspace gas chromatography with flame thermionic detection 734(1999)319

TS-962

Highly sensitive determination of TS-962 (HL-004), a novel acyl-CoA:cholesterol acyltransferase inhibitor, in rat and rabbit plasma by liquid chromatography and atmospheric pressure chemical ionization-tandem mass spectrometry combined with a column-switching technique 750(2001)99

Tumor necrosis factor receptor immunoglobulin G

Purification and quantitation of tumor necrosis factor receptor immunoadhesin using a combination of immunoaffinity and reversed-phase chromatography 728(1999)21

Tyrosine

Neutral amino acids monitoring in phenylketonuric plasma microdialysates using micellar electrokinetic chromatography and laser-induced fluorescence detection 739(2000)247

U**506U78**

High-performance liquid chromatography method for the determination and quantitation of arabinosylguanine triphosphate and fludarabine triphosphate in human cells 745(2000)421

Ubiquinol

Automated high-performance liquid chromatographic method with precolumn reduction for the determination of ubiquinol and ubiquinone in human plasma 726(1999)297

Ubiquinone

Automated high-performance liquid chromatographic method with precolumn reduction for the determination of ubiquinol and ubiquinone in human plasma 726(1999)297

UDP-GalNAc:polypeptide *N*-acetylgalactosaminyltransferase

Capillary zone electrophoresis and MALDI–mass spectrometry for the monitoring of in vitro *O*-glycosylation of a threonine/serine-rich MUC5AC hexadecapeptide 729(1999)65

Uracil

Determination of urinary orotic acid and uracil by capillary zone electrophoresis 734(1999)175

Measurement of plasma uracil using gas chromatography–mass spectrometry in normal individuals and in patients receiving inhibitors of dihydropyrimidine dehydrogenase 738(2000)249

Urea

Determination of urea, allantoin and lysine pyroglutamate in cosmetic samples by hydrophilic interaction chromatography 742(2000)447

Urease

Pilot study of gas chromatographic–mass spectrometric screening of newborn urine for inborn errors of metabolism after treatment with urease 731(1999)141

Purification of surface-associated urease from *Helicobacter pylori* 737(2000)202

Differential diagnosis of homocystinuria by urease treatment, isotope dilution and gas chromatography–mass spectrometry 742(2000)59

Differential diagnosis of homocystinuria by urease treatment, isotope dilution and gas chromatography–mass spectrometry 746(2000)103

Urethane

Effect of 2-deoxy-D-glucose on acetylcholine and histamine levels in gastric juice of pylorus-ligated rats anesthetized with urethane 742(2000)295

Uric acid

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

Quantification of allantoin, uric acid, xanthine and hypoxanthine in ovine urine by high-performance liquid chromatography and photodiode array detection 744(2000)129

Determination of uremic toxins in biofluids: creatinine, creatine, uric acid and xanthines 747(2000)217

Urinary steroids

Optimization of the high-performance liquid chromatographic separation of a complex mixture containing urinary steroids, boldenone and bolasterone: application to urine samples 742(2000)47

Urocanic acid

Determination of histidine and urocanic acid isomers in the human skin by high-performance capillary electrophoresis 749(2000)41

Analysis of histidine and urocanic acid isomers by reversed-phase high-performance liquid chromatography 750(2001)71

Urokinase

Affinity chromatography for purification of two urokinases from human urine 742(2000)71

Urokinase plasminogen activator

Refolding and purification of a urokinase plasminogen activator fragment by chromatography 737(2000)225

V

Valaciclovir

Rapid determination of valaciclovir and acyclovir in human biological fluids by high-performance liquid chromatography using isocratic elution 732(1999)47

Valine

Determination of extracellular and intracellular enrichments of [1-¹³C]-α-ketoisovalerate using enzymatically converted [1-¹³C]-valine standardization curves and gas chromatography–mass spectrometry 729(1999)211

Neutral amino acids monitoring in phenylketonuric plasma microdialysates using micellar electrokinetic chromatography and laser-induced fluorescence detection 739(2000)247

Determination of the dissociation constant of valine from acetohydroxy acid synthase by equilibrium partition in an aqueous two-phase system 743(2000)225

Valium

On-line deproteinization by adsorption of proteins on a polyethylene powder pre-column for the determination of Na, K, Mg and Ca in human serum by high-performance liquid chromatography 750(2001)61

Valproic acid

Metabolic profiling of valproic acid by cDNA-expressed human cytochrome P450 enzymes using negative-ion chemical ionization gas chromatography–mass spectrometry 742(2000)217

Vancomycin

Chiral separation by simultaneous use of vancomycin as stationary phase chiral selector and chiral mobile phase additive 745(2000)159

Vanillic acid

High-performance liquid chromatographic method for determination of vanillin and vanillic acid in human plasma, red blood cells and urine 726(1999)303

Vanillin

High-performance liquid chromatographic method for determination of vanillin and vanillic acid in human plasma, red blood cells and urine 726(1999)303

Vanilmandelic acid

Liquid chromatography–electrospray tandem mass spectrometry of acidic monoamine metabolites 744(2000)423

Verapamil

Achiral and chiral high-performance liquid chromatography of verapamil and its metabolites in serum samples 732(1999)395

Rapid and highly sensitive method for the determination of verapamil, [³H]₇verapamil and metabolites in biological fluids by liquid chromatography–mass spectrometry 738(2000)137

Development of an immobilized P-glycoprotein stationary phase for on-line liquid chromatographic determination of drug-binding affinities 739(2000)33

Vigabatrin

Fast isocratic high-performance liquid chromatographic assay method for the simultaneous determination of gabapentin and vigabatrin in human serum 746(2000)311

Vinorelbine

High-performance liquid chromatographic determination of vinorelbine in human plasma and blood: application to a pharmacokinetic study 748(2000)389

Viral coat proteins

Purification of a viral coat protein by an engineered polyionic sequence 737(2000)77

Viral proteins

Aggregation of recombinant hepatitis B surface antigen induced in vitro by oxidative stress 736(1999)153

Vitamins

Analysis of riboflavin and riboflavin cofactor levels in plasma by high-performance liquid chromatography 739(2000)219

Vitamin D

Determination of the vitamin D analog EB 1089 (seocalcitol) in human and pig serum using liquid chromatography–tandem mass spectrometry 740(2000)117

Vitamin K-dependent

Separation of human vitamin K-dependent coagulation proteins using hydrophobic interaction chromatography 736(1999)77

Vitamin K-dependent coagulation proteins

Separation of human vitamin K-dependent coagulation proteins using hydrophobic interaction chromatography 736(1999)77

Vitamin E

Simple high-performance liquid chromatographic method for determination of α -tocopherol in human plasma 732(1999)227

Vitamins

Rapid mass spectrometric analysis for ascorbate and related organic acids in small volumes of plasma for use in pediatric subjects 726(1999)79

Improved method for simultaneous determination of ascorbic acid and dehydroascorbic acid, isoascorbic acid and dehydroisoascorbic acid in food and biological samples 730(1999)101

Application of two-dimensional high-performance liquid chromatography–mass spectrometry with particle beam interface 730(1999)141

Simultaneous determination of serum retinol and α - and γ -tocopherol levels in type II diabetic patients using high-performance liquid chromatography with fluorescence detection 730(1999)265

Simple high-performance liquid chromatographic method for the determination of tocotrienols in human plasma 735(1999)279

Determination of pyrroloquinoline quinone by capillary zone electrophoresis 739(2000)101

Capillary electrophoretic separation of vitamins in sodium dodecyl sulfate containing buffers with lower aliphatic alcohols and *n*-hexane as organic modifiers 741(2000)67

Analysis of glutathionyl hemoglobin levels in diabetic patients by electrospray ionization liquid chromatography–mass spectrometry: effect of vitamin E administration 746(2000)91

Vitellogenin

Two-step purification method of vitellogenin from three teleost fish species: rainbow trout (*Oncorhynchus mykiss*), gudgeon (*Gobio gobio*) and chub (*Leuciscus cephalus*) 737(2000)3

Volatile hydrocarbons

Headspace solid-phase microextraction and gas chromatographic–mass spectrometric screening for volatile hydrocarbons in blood 748(2000)401

Volatile organic compounds

Method for liquid–liquid extraction of blood surrogates for assessing human exposure to jet fuel 728(1999)193

Variation in volatile organic compounds in the breath of normal humans 729(1999)75

Volatile salts

Partitioning behavior of amino acids in aqueous two-phase systems with recyclable volatile salts 743(2000)317

X

Xanthine

Quantification of allantoin, uric acid, xanthine and hypoxanthine in ovine urine by high-performance liquid chromatography and photodiode array detection 744(2000)129

Determination of uremic toxins in biofluids: creatinine, creatine, uric acid and xanthines 747(2000)217

Xylanolitic complex

Partitioning of xylanolitic complex from *Penicillium janthinellum* by an aqueous two-phase system 743(2000)339

Xylazine

Gas chromatographic–mass spectrometric analysis of veterinary tranquilizers in urine: evaluation of method performance 734(1999)113

Xylene

Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis. Comparison with high-performance liquid chromatography 730(1999)41

2,6-Xylidine

High-performance liquid chromatography–tandem electrospray mass spectrometry for the determination of lidocaine and its metabolites in human plasma and urine 741(2000)175

Z

Zanamivir

Liquid chromatographic–tandem mass spectrometric method for the determination of the neuraminidase inhibitor zanamivir (GG167) in human serum 732(1999)383

Zearalenol

Determination of zearalenone and its metabolites α - and β -zearalenol in beer samples by high-performance liquid chromatography–tandem mass spectrometry 738(2000)233

Zearalenone

Determination of zearalenone and its metabolites α - and β -zearalenol in beer samples by high-performance liquid chromatography–tandem mass spectrometry 738(2000)233

Zidovudine

Simultaneous determination of zidovudine and its monophosphate in mouse plasma and peripheral red blood cells by high-performance liquid chromatography 740(2000)281

Simultaneous determination of lamivudine and zidovudine concentrations in human seminal plasma using high-performance liquid chromatography and tandem mass spectrometry 742(2000)173

Zidovudine monophosphate

Simultaneous determination of zidovudine and its monophosphate in mouse plasma and peripheral red blood cells by high-performance liquid chromatography 740(2000)281

Zidovudine- β -D-glucuronide

Complete two-dimensional separation for analysis of acidic compounds in plasma using column-switching reversed-phase liquid chromatography 726(1999)225

Zofenopril

Assay of zofenopril and its active metabolite zofenoprilat by liquid chromatography coupled with tandem mass spectrometry 749(2000)287

Zofenoprilat

Assay of zofenopril and its active metabolite zofenoprilat by liquid chromatography coupled with tandem mass spectrometry 749(2000)287

Zolpidem

Determination of zolpidem in serum microsamples by high-performance liquid chromatography and its application to pharmacokinetics in rats 734(1999)299