

Available online at www.sciencedirect.com



JOURNAL OF CHROMATOGRAPHY B

Journal of Chromatography B, 824 (2005) 354-360

www.elsevier.com/locate/chromb

Keyword Index Vol. 824

Absolute molecular weight distribution: Mass fractions

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

Absolute molecular weight distribution

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

Acetate

Determination of testosterone concentrations in rat plasma using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry combined with ethyl oxime and acetyl ester derivatization 824(2005)258

N-Acetyl-S-(2-carbamoylethyl)-L-cysteine (AAMA)

Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC-ESI-MS/MS 824(2005)283

Acetylcodeine

The quantitative analysis of heroin, methadone and their metabolites and the simultaneous detection of cocaine, acetylcodeine and their metabolites in human plasma by high-performance liquid chromatography coupled with tandem mass spectrometry 824(2005)213

Acrylamide (AA)

Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC-ESI-MS/MS 824(2005)283

Aminoacetone

A novel HPLC procedure for detection and quantification of aminoacetone, a precursor of methylglyoxal, in biological samples 824(2005)116

3-Aminobenzanthrone

Assessment of 3-nitrobenzanthrone reductase activity in mammalian tissues by normal-phase HPLC with fluorescence detection 824(2005)229

Analysis in plasma

Determination of lorazepam in plasma from children by high-performance liquid chromatography with UV detection 824(2005)333

Analysis

Analysis of omeprazole, midazolam and hydroxy-metabolites in plasma using liquid chromatography coupled to tandem mass spectrometry 824(2005)71

Anisodamine

Liquid chromatography-tandem mass spectrometry analysis of anisodamine and its phase I and II metabolites in rat urine (Review) 824(2005)21

Ascorbic acid

High-throughput analysis of Vitamin C in human plasma with the use of HPLC with monolithic column and UV-detection 824(2005)132

Astragaloside IV

Quantitation of astragaloside IV in rat plasma by liquid chromatography-tandem mass spectrometry 824(2005)244

Astragalus membranaceous

Quantitation of astragaloside IV in rat plasma by liquid chromatography-tandem mass spectrometry 824(2005)244

Automated pre-column derivatization

A rapid and validated HPLC method to quantify sphingosine 1-phosphate in human plasma using solid-phase extraction followed by derivatization with fluorescence detection 824(2005)65

Benzo[a]pyrene

A novel, sensitive method for determining benzo[a]pyrene-diones using high-performance liquid chromatography with post-column zinc reduction 824(2005)166

Benzo[a]pyrene-dione

A novel, sensitive method for determining benzo[a]pyrene-diones using high-performance liquid chromatography with post-column zinc reduction 824(2005)166

BGC638

A liquid chromatographic-tandem mass spectrometric method for the determination of two selective thymidylate synthase inhibitors, BGC945 and BGC638, in mouse plasma 824(2005)181

BGC945

A liquid chromatographic–tandem mass spectrometric method for the determination of two selective thymidylate synthase inhibitors, BGC945 and BGC638, in mouse plasma 824(2005)181

Biological monitoring

Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC-ESI-MS/MS 824(2005)283

Broad distributed branched AP-glucans

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

Calphoglin

Identification of human calphoglin-induced phosphoglucomutase phosphorylation in *Escherichia coli* 824(2005)312

Capillary electrophoresis

Capillary electrophoresis of the electrochemical oxidation products of Nghydroxy-L-arginine at physiological pH 824(2005)308

Capillary isoelectric focusing

Optimization of protein identification from digests as analyzed by capillary isoelectric focusing-mass spectrometry 824(2005)189

Carotenoids

An improved HPLC method for determination of carotenoids in human serum 824(2005)99

Catechin

One-step purification of epigallocatechin gallate from crude green tea extracts by isocratic hydrogen bond adsorption chromatography on β -cyclodextrin substituted agarose gel media 824(2005)323

Chemiluminescence

Determination of tetracyclines residues in honey using high-performance liquid chromatography with potassium permanganate–sodium sulfite–β-cyclodextrin chemiluminescence detection 824(2005)57

Children

Determination of lorazepam in plasma from children by high-performance liquid chromatography with UV detection 824(2005)333

Chlorcyclizine

An LC-MS-MS method for the determination of cyclizine in human serum 824(2005)148

Cocaine

The quantitative analysis of heroin, methadone and their metabolites and the simultaneous detection of cocaine, acetylcodeine and their metabolites in human plasma by high-performance liquid chromatography coupled with tandem mass spectrometry 824(2005)213

Collagen

Quantification of hydroxyproline in small amounts of skin tissue using isocratic high performance liquid chromatography with NBD-F as fluorogenic reagent 824(2005)161

Curcumin

Tetrahydrocurcumin in plasma and urine: Quantitation by high performance liquid chromatography 824(2005)206

B-Cyclodextrin

One-step purification of epigallocatechin gallate from crude green tea extracts by isocratic hydrogen bond adsorption chromatography on β -cyclodextrin substituted agarose gel media 824(2005)323

Cyclizine

An LC–MS–MS method for the determination of cyclizine in human serum 824(2005)148

CYP2C19

Analysis of omeprazole, midazolam and hydroxy-metabolites in plasma using liquid chromatography coupled to tandem mass spectrometry 824(2005)71

Cytochrome P450

A novel, sensitive method for determining benzo[a]pyrene-diones using high-performance liquid chromatography with post-column zinc reduction 824(2005)166

Data analysis

Optimization of protein identification from digests as analyzed by capillary isoelectric focusing-mass spectrometry 824(2005)189

Dehydroascorbic acid

High-throughput analysis of Vitamin C in human plasma with the use of HPLC with monolithic column and UV-detection 824(2005)132

Derivatization, LC

Quantitative analysis of valienamine in the microbial degradation of validamycin A after derivatization with p-nitrofluorobenzene by reversed-phase high-performance liquid chromatography 824(2005)341

Derivatization

Determination of testosterone concentrations in rat plasma using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry combined with ethyl oxime and acetyl ester derivatization 824(2005)258

Designer drug

Studies on the metabolism and toxicological detection of the new designer drug 4'-methyl- α -pyrrolidinobutyrophenone (MPBP) in rat urine using gas chromatography-mass spectrometry 824(2005)81

Studies on the metabolism and toxicological detection of the designer drug 4-methylthioamphetamine (4-MTA) in human urine using gas chromatography-mass spectrometry 824(2005)123

Detection, LC

Quantitative analysis of valienamine in the microbial degradation of validamycin A after derivatization with *p*-nitrofluorobenzene by reversed-phase high-performance liquid chromatography 824(2005)341

Detection

Studies on the metabolism and toxicological detection of the new designer drug 4'-methyl- α -pyrrolidinobutyrophenone (MPBP) in rat urine using gas chromatography—mass spectrometry 824(2005)81

Drug analysis by capillary electrophoresis–electrospray ionisation mass spectrometry (CE–ESI-MS) $\,$

Recent studies on the electrospray ionisation mass spectrometric behaviour of selected nitrogen-containing drug molecules and its application to drug analysis using liquid chromatography–electrospray ionisation mass spectrometry (Review) 824(2005)1

Drug analysis by gas liquid chromatography-mass spectrometry (GLC-MS)

Recent studies on the electrospray ionisation mass spectrometric behaviour of selected nitrogen-containing drug molecules and its application to drug analysis using liquid chromatography–electrospray ionisation mass spectrometry(Review) 824(2005)1

Drug analysis by liquid chromatography–electrospray ionisation mass spectrometry (LC–ESI-MS)

Recent studies on the electrospray ionisation mass spectrometric behaviour of selected nitrogen-containing drug molecules and its application to drug analysis using liquid chromatography–electrospray ionisation mass spectrometry (Review) 824(2005)1

Drugs of small molecular mass

Recent studies on the electrospray ionisation mass spectrometric behaviour of selected nitrogen-containing drug molecules and its application to drug analysis using liquid chromatography–electrospray ionisation mass spectrometry (Review) 824(2005)1

EGCG

One-step purification of epigallocatechin gallate from crude green tea extracts by isocratic hydrogen bond adsorption chromatography on β -cyclodextrin substituted agarose gel media 824(2005)323

Eggs

Determination of nitrofuran metabolites in poultry muscle and eggs by liquid chromatography-tandem mass spectrometry 824(2005)30

Electron capture detection (ECD)

Use of pressurized liquid extraction (PLE)/gas chromatography–electron capture detection (GC–ECD) for the determination of biodegradation intermediates of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in soils 824(2005)277

Electrospray ionisation mass spectrometry (ESI-MS)

Recent studies on the electrospray ionisation mass spectrometric behaviour of selected nitrogen-containing drug molecules and its application to drug analysis using liquid chromatography-electrospray ionisation mass spectrometry (Review) 824(2005)1

Electrospray ionization mass spectrometry

Development and validation of a high performance liquid chromatography assay for 17α -methyltestosterone in fish feed 824(2005)107

Endogenous sorbitol and fructose

Quantitative determination of endogenous sorbitol and fructose in human erythrocytes by atmospheric-pressure chemical ionization LC tandem mass spectrometry 824(2005)36

Exercise

Urinary high performance reverse phase chromatography cortisol and cortisone analyses before and at the end of a race in elite cyclists 824(2005)51

Explosives

Use of pressurized liquid extraction (PLE)/gas chromatography–electron capture detection (GC–ECD) for the determination of biodegradation intermediates of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in soils 824(2005)277

Fish feed

Development and validation of a high performance liquid chromatography assay for 17α -methyltestosterone in fish feed 824(2005)107

FLU detection

A rapid and validated HPLC method to quantify sphingosine 1-phosphate in human plasma using solid-phase extraction followed by derivatization with fluorescence detection 824(2005)65

5-Fluorodihydrouracil

Sensitive MS/MS-liquid chromatography assay for simultaneous determination of tegafur, 5-fluorouracil and 5-fluorodihydrouracil in plasma 824(2005)153

5-Fluorouracil

Sensitive MS/MS-liquid chromatography assay for simultaneous determination of tegafur, 5-fluorouracil and 5-fluorodihydrouracil in plasma 824(2005)153

Fragmentation patterns

Recent studies on the electrospray ionisation mass spectrometric behaviour of selected nitrogen-containing drug molecules and its application to drug analysis using liquid chromatography–electrospray ionisation mass spectrometry (Review) 824(2005)1

Gas chromatography (GC)

Use of pressurized liquid extraction (PLE)/gas chromatography–electron capture detection (GC–ECD) for the determination of biodegradation intermediates of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in soils 824(2005)277

GC-MS

Studies on the metabolism and toxicological detection of the new designer drug 4'-methyl- α -pyrrolidinobutyrophenone (MPBP) in rat urine using gas chromatography-mass spectrometry 824(2005)81

Studies on the metabolism and toxicological detection of the designer drug 4-methylthioamphetamine (4-MTA) in human urine using gas chromatography-mass spectrometry 824(2005)123

Gel permeation

Quantification of rotavirus-like particles by gel permeation chromatography 824(2005)267

Genome

Identification of human calphoglin-induced phosphoglucomutase phosphorylation in *Escherichia coli* 824(2005)312

Glycidamide (GA)

Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC-ESI-MS/MS 824(2005)283

Heparan sulphate

Use of graphitised carbon negative ion LC-MS to analyse enzymatically digested glycosaminoglycans 824(2005)139

Heparin

Use of graphitised carbon negative ion LC-MS to analyse enzymatically digested glycosaminoglycans 824(2005)139

Heroin

The quantitative analysis of heroin, methadone and their metabolites and the simultaneous detection of cocaine, acetylcodeine and their metabolites in human plasma by high-performance liquid chromatography coupled with tandem mass spectrometry 824(2005)213

High performance liquid chromatography (HPLC)

Development and validation of a high performance liquid chromatography assay for 17α -methyltestosterone in fish feed 824(2005)107

Quantification of hydroxyproline in small amounts of skin tissue using isocratic high performance liquid chromatography with NBD-F as fluorogenic reagent 824(2005)161

A novel, sensitive method for determining benzo[a]pyrene-diones using high-performance liquid chromatography with post-column zinc reduction 824(2005)166

Determination of lorazepam in plasma from children by high-performance liquid chromatography with UV detection 824(2005)333

Honey

Determination of tetracyclines residues in honey using high-performance liquid chromatography with potassium permanganate–sodium sulfite– β -cyclodextrin chemiluminescence detection 824(2005)57

HPLC column-switching

A rapid and validated HPLC method to quantify sphingosine 1-phosphate in human plasma using solid-phase extraction followed by derivatization with fluorescence detection 824(2005)65

HPLO

Determination of tetracyclines residues in honey using high-performance liquid chromatography with potassium permanganate–sodium sulfite– β -cyclodextrin chemiluminescence detection 824(2005)57

An improved HPLC method for determination of carotenoids in human serum $\;\;824(2005)99\;\;$

A novel HPLC procedure for detection and quantification of aminoacetone, a precursor of methylglyoxal, in biological samples 824(2005)116

Assessment of 3-nitrobenzanthrone reductase activity in mammalian tissues by normal-phase HPLC with fluorescence detection 824(2005)229

Determination of rabeprazole and its active metabolite, rabeprazole thioether in human plasma by column-switching high-performance liquid chromatography and its application to pharmacokinetic study 824(2005)238

Quantification of rotavirus-like particles by gel permeation chromatography 824(2005)267

Determination of thiopental in urine sample with high-performance liquid chromatography using iodine-azide reaction as a postcolumn detection system 824(2005)327

HPLC-ESI-MS

Quantification of lipoic acid in plasma by high-performance liquid chromatography–electrospray ionization mass spectrometry 824(2005)249

HPLC-ESI-MS

High performance liquid chromatography-electrospray ionization mass spectrometric determination of tolterodine tartrate in human plasma 824(2005)92

HPTLC

Detection of mercaptopyridines and mercaptopyrimidines in planar chromatography with iodine-azide reaction as a detection system 824(2005)222

11β-HSD

Urinary high performance reverse phase chromatography cortisol and cortisone analyses before and at the end of a race in elite cyclists 824(2005)51

Human plasma

High performance liquid chromatography-electrospray ionization mass spectrometric determination of tolterodine tartrate in human plasma 824(2005)92

A rapid and validated HPLC method to quantify sphingosine 1-phosphate in human plasma using solid-phase extraction followed by derivatization with fluorescence detection 824(2005)65

Human red blood cells

Quantitative determination of endogenous sorbitol and fructose in human erythrocytes by atmospheric-pressure chemical ionization LC tandem mass spectrometry 824(2005)36

Human serum

An improved HPLC method for determination of carotenoids in human serum 824(2005)99

An LC–MS–MS method for the determination of cyclizine in human serum 824(2005)148

Hvaluronic acid

Use of graphitised carbon negative ion LC-MS to analyse enzymatically digested glycosaminoglycans 824(2005)139

Hydroxyproline

Quantification of hydroxyproline in small amounts of skin tissue using isocratic high performance liquid chromatography with NBD-F as fluorogenic reagent 824(2005)161

Indirect fluorescence detection

Estimation of pK_a values using microchip capillary electrophoresis and indirect fluorescence detection 824(2005)201

Iodine-azide reaction

Detection of mercaptopyridines and mercaptopyrimidines in planar chromatography with iodine-azide reaction as a detection system 824(2005)222

Determination of thiopental in urine sample with high-performance liquid chromatography using iodine-azide reaction as a postcolumn detection system 824(2005)327

Keratan sulphate

Use of graphitised carbon negative ion LC-MS to analyse enzymatically digested glycosaminoglycans 824(2005)139

LC/APCI-MS/MS

Quantitative determination of endogenous sorbitol and fructose in human erythrocytes by atmospheric-pressure chemical ionization LC tandem mass spectrometry 824(2005)36

LC/MS-MS

Sensitive MS/MS-liquid chromatography assay for simultaneous determination of tegafur, 5-fluorouracil and 5-fluorodihydrouracil in plasma 824(2005)153

LC

A new and simple solid-phase extraction method for LC determination of pyronaridine in human plasma 824(2005)45

LCMS

Analysis of omeprazole, midazolam and hydroxy-metabolites in plasma using liquid chromatography coupled to tandem mass spectrometry 824(2005)71

LC-MS/MS

Liquid chromatography-tandem mass spectrometry analysis of anisodamine and its phase I and II metabolites in rat urine (Review) 824(2005)21

LC-MS-MS

An LC–MS–MS method for the determination of cyclizine in human serum 824(2005)148

The quantitative analysis of heroin, methadone and their metabolites and the simultaneous detection of cocaine, acetylcodeine and their metabolites in human plasma by high-performance liquid chromatography coupled with tandem mass spectrometry 824(2005)213

Lipoic acid

Quantification of lipoic acid in plasma by high-performance liquid chromatography-electrospray ionization mass spectrometry 824(2005)249

Liquid chromatography

Determination of uric acid in urine, saliva and calcium oxalate renal calculi by high-performance liquid chromatography/mass spectrometry 824(2005)175

Liquid chromatography-atmosphere pressure chemical ionization mass spectrometry

Determination of testosterone concentrations in rat plasma using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry combined with ethyl oxime and acetyl ester derivatization 824(2005)258

Liquid chromatography-tandem mass spectrometry

A liquid chromatographic-tandem mass spectrometric method for the determination of two selective thymidylate synthase inhibitors, BGC945 and BGC638, in mouse plasma 824(2005)181

Liquid-liquid extraction

Determination of metformin in human plasma by high-performance liquid chromatography 824(2005)319

Lorazepam

Determination of lorazepam in plasma from children by high-performance liquid chromatography with UV detection 824(2005)333

Malaria

A new and simple solid-phase extraction method for LC determination of pyronaridine in human plasma 824(2005)45

Mass spectrometry

Determination of nitrofuran metabolites in poultry muscle and eggs by liquid chromatography-tandem mass spectrometry 824(2005)30

Determination of uric acid in urine, saliva and calcium oxalate renal calculi by high-performance liquid chromatography/mass spectrometry 824(2005)175

Optimization of protein identification from digests as analyzed by capillary isoelectric focusing-mass spectrometry 824(2005)189

Matrix effects

Quantitative determination of endogenous sorbitol and fructose in human erythrocytes by atmospheric-pressure chemical ionization LC tandem mass spectrometry 824(2005)36

Mercaptopyridines

Detection of mercaptopyridines and mercaptopyrimidines in planar chromatography with iodine-azide reaction as a detection system 824(2005)222

Mercaptopyrimidines

Detection of mercaptopyridines and mercaptopyrimidines in planar chromatography with iodine-azide reaction as a detection system 824(2005)222

Mercapturic acids

Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC-ESI-MS/MS 824(2005)283

Metabolism

Studies on the metabolism and toxicological detection of the new designer drug 4'-methyl- α -pyrrolidinobutyrophenone (MPBP) in rat urine using gas chromatography–mass spectrometry 824(2005)81

Studies on the metabolism and toxicological detection of the designer drug 4-methylthioamphetamine (4-MTA) in human urine using gas chromatography-mass spectrometry 824(2005)123

Metabolite

Liquid chromatography-tandem mass spectrometry analysis of anisodamine and its phase I and II metabolites in rat urine (Review) 824(2005)21

A novel, sensitive method for determining benzo[a]pyrene-diones using high-performance liquid chromatography with post-column zinc reduction 824(2005)166

Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC–ESI-MS/MS 824(2005)283

Metformin

Determination of metformin in human plasma by high-performance liquid chromatography 824(2005)319

17α -Methyltestosterone

Development and validation of a high performance liquid chromatography assay for 17α-methyltestosterone in fish feed 824(2005)107

Methadone

The quantitative analysis of heroin, methadone and their metabolites and the simultaneous detection of cocaine, acetylcodeine and their metabolites in human plasma by high-performance liquid chromatography coupled with tandem mass spectrometry 824(2005)213

4'-Methyl-α-pyrrolidinobutyrophenone

Studies on the metabolism and toxicological detection of the new designer drug 4'-methyl- α -pyrrolidinobutyrophenone (MPBP) in rat urine using gas chromatography-mass spectrometry 824(2005)81

Methylglyoxal

A novel HPLC procedure for detection and quantification of aminoacetone, a precursor of methylglyoxal, in biological samples 824(2005)116

4-Methylthioamphetamine

Studies on the metabolism and toxicological detection of the designer drug 4-methylthioamphetamine (4-MTA) in human urine using gas chromatography-mass spectrometry 824(2005)123

Microchip capillary electrophoresis

Estimation of pK_a values using microchip capillary electrophoresis and indirect fluorescence detection 824(2005)201

Midazolam

Analysis of omeprazole, midazolam and hydroxy-metabolites in plasma using liquid chromatography coupled to tandem mass spectrometry 824(2005)71

Mixed type branched AP-glucans

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

Mixed-mode solid-phase extraction

Quantitative determination of endogenous sorbitol and fructose in human erythrocytes by atmospheric-pressure chemical ionization LC tandem mass spectrometry 824(2005)36

Molar fractions

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

Monolithic column

High-throughput analysis of Vitamin C in human plasma with the use of HPLC with monolithic column and UV-detection 824(2005)132

MPBP

Studies on the metabolism and toxicological detection of the new designer drug 4'-methyl- α -pyrrolidinobutyrophenone (MPBP) in rat urine using gas chromatography-mass spectrometry 824(2005)81

4-MTA

Studies on the metabolism and toxicological detection of the designer drug 4-methylthioamphetamine (4-MTA) in human urine using gas chromatography–mass spectrometry 824(2005)123

N-(R/S)-Acetyl-S-(2-carbamoyl-2-hydroxyethyl)-L-cysteine (GAMA)

Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC-ESI-MS/MS 824(2005)283

Narrow distributed non-branched AP-glucans

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

NBD-F

Quantification of hydroxyproline in small amounts of skin tissue using isocratic high performance liquid chromatography with NBD-F as fluorogenic reagent 824(2005)161

Ng-hydroxy-arginine

Capillary electrophoresis of the electrochemical oxidation products of Nghydroxy-L-arginine at physiological pH 824(2005)308

Nitroarenes

Assessment of 3-nitrobenzanthrone reductase activity in mammalian tissues by normal-phase HPLC with fluorescence detection 824(2005)229

Nitrofuran metabolites

Determination of nitrofuran metabolites in poultry muscle and eggs by liquid chromatography-tandem mass spectrometry 824(2005)30

Nitroreductase

Assessment of 3-nitrobenzanthrone reductase activity in mammalian tissues by normal-phase HPLC with fluorescence detection 824(2005)229

p-Nitrofluorobenzene

Quantitative analysis of valienamine in the microbial degradation of validamycin A after derivatization with *p*-nitrofluorobenzene by reversed-phase high-performance liquid chromatography 824(2005)341

Omeprazole

Analysis of omeprazole, midazolam and hydroxy-metabolites in plasma using liquid chromatography coupled to tandem mass spectrometry 824(2005)71

Orthologue proteome

Identification of human calphoglin-induced phosphoglucomutase phosphorylation in *Escherichia coli* 824(2005)312

Oxidation

Capillary electrophoresis of the electrochemical oxidation products of Nghydroxy-L-arginine at physiological pH 824(2005)308

Oxim

Determination of testosterone concentrations in rat plasma using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry combined with ethyl oxime and acetyl ester derivatization 824(2005)258

Pharmacokinetics

High performance liquid chromatography-electrospray ionization mass spectrometric determination of tolterodine tartrate in human plasma 824(2005)92

Determination of lorazepam in plasma from children by high-performance liquid chromatography with UV detection 824(2005)333

Phosphoglucomutase

Identification of human calphoglin-induced phosphoglucomutase phosphorylation in *Escherichia coli* 824(2005)312

pK,

Estimation of pK_a values using microchip capillary electrophoresis and indirect fluorescence detection 824(2005)201

Plasma

A new and simple solid-phase extraction method for LC determination of pyronaridine in human plasma 824(2005)45

High-throughput analysis of Vitamin C in human plasma with the use of HPLC with monolithic column and UV-detection 824(2005)132

Tetrahydrocurcumin in plasma and urine: Quantitation by high performance liquid chromatography 824(2005)206

Post-column

A novel, sensitive method for determining benzo[a]pyrene-diones using high-performance liquid chromatography with post-column zinc reduction 824(2005)166

Poultry muscle

Determination of nitrofuran metabolites in poultry muscle and eggs by liquid chromatography-tandem mass spectrometry 824(2005)30

Pressurized liquid extraction (PLE)

Use of pressurized liquid extraction (PLE)/gas chromatography–electron capture detection (GC–ECD) for the determination of biodegradation intermediates of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in soils 824(2005)277

Protein analysis

Optimization of protein identification from digests as analyzed by capillary isoelectric focusing-mass spectrometry 824(2005)189

Protein phosphorylation

Identification of human calphoglin-induced phosphoglucomutase phosphorylation in *Escherichia coli* 824(2005)312

Proteoglycans

Use of graphitised carbon negative ion LC-MS to analyse enzymatically digested glycosaminoglycans 824(2005)139

Pyridylamination of glucans with terminal hemiacetal

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

Pyronaridine

A new and simple solid-phase extraction method for LC determination of pyronaridine in human plasma 824(2005)45

Rabeprazole thioether

Determination of rabeprazole and its active metabolite, rabeprazole thioether in human plasma by column-switching high-performance liquid chromatography and its application to pharmacokinetic study 824(2005)238

Rabeprazole

Determination of rabeprazole and its active metabolite, rabeprazole thioether in human plasma by column-switching high-performance liquid chromatography and its application to pharmacokinetic study 824(2005)238

RDX

Use of pressurized liquid extraction (PLE)/gas chromatography–electron capture detection (GC–ECD) for the determination of biodegradation intermediates of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in soils 824(2005)277

Renal calculi

Determination of uric acid in urine, saliva and calcium oxalate renal calculi by high-performance liquid chromatography/mass spectrometry 824(2005)175

Residues

Determination of tetracyclines residues in honey using high-performance liquid chromatography with potassium permanganate–sodium sulfite– β -cyclodextrin chemiluminescence detection 824(2005)57

Rotavirus-like particles

Quantification of rotavirus-like particles by gel permeation chromatography 824(2005)267

S1

Sensitive MS/MS-liquid chromatography assay for simultaneous determination of tegafur, 5-fluorouracil and 5-fluorodihydrouracil in plasma 824(2005)153

SEC-mass/molar

De facto molecular weight distributions of glucans by size-exclusion chromatography combined with mass/molar-detection of fluorescence labeled terminal hemiacetals 824(2005)295

Sequest

Optimization of protein identification from digests as analyzed by capillary isoelectric focusing-mass spectrometry 824(2005)189

Skir

Quantification of hydroxyproline in small amounts of skin tissue using isocratic high performance liquid chromatography with NBD-F as fluorogenic reagent 824(2005)161

Soil analysis

Use of pressurized liquid extraction (PLE)/gas chromatography–electron capture detection (GC–ECD) for the determination of biodegradation intermediates of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in soils 824(2005)277

Solid phase extraction

Urinary high performance reverse phase chromatography cortisol and cortisone analyses before and at the end of a race in elite cyclists 824(2005)51

Development and validation of a high performance liquid chromatography assay for 17α -methyltestosterone in fish feed 824(2005)107

SPE

A rapid and validated HPLC method to quantify sphingosine 1-phosphate in human plasma using solid-phase extraction followed by derivatization with fluorescence detection 824(2005)65

Sphingosine 1-phosphate

A rapid and validated HPLC method to quantify sphingosine 1-phosphate in human plasma using solid-phase extraction followed by derivatization with fluorescence detection 824(2005)65

SSAO

A novel HPLC procedure for detection and quantification of aminoacetone, a precursor of methylglyoxal, in biological samples 824(2005)116

Stripping voltammetry (SV)

Recent studies on the electrospray ionisation mass spectrometric behaviour of selected nitrogen-containing drug molecules and its application to drug analysis using liquid chromatography–electrospray ionisation mass spectrometry 824(2005)1

Tea polyphenols

One-step purification of epigallocatechin gallate from crude green tea extracts by isocratic hydrogen bond adsorption chromatography on β -cyclodextrin substituted agarose gel media 824(2005)323

Tégafur

Sensitive MS/MS-liquid chromatography assay for simultaneous determination of tegafur, 5-fluorouracil and 5-fluorodihydrouracil in plasma 824(2005)153

Testosterone

Determination of testosterone concentrations in rat plasma using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry combined with ethyl oxime and acetyl ester derivatization 824(2005)258

Tetracycline antibiotics

Determination of tetracyclines residues in honey using high-performance liquid chromatography with potassium permanganate-sodium sulfite-β-cyclodextrin chemiluminescence detection 824(2005)57

Tetrahydrocurcumin

Tetrahydrocurcumin in plasma and urine: Quantitation by high performance liquid chromatography 824(2005)206

Thiopental

Determination of thiopental in urine sample with high-performance liquid chromatography using iodine-azide reaction as a postcolumn detection system 824(2005)327

TLC

Detection of mercaptopyridines and mercaptopyrimidines in planar chromatography with iodine-azide reaction as a detection system 824(2005)222

Tolterodine tartrate

High performance liquid chromatography-electrospray ionization mass spectrometric determination of tolterodine tartrate in human plasma 824(2005)92

UFT

Sensitive MS/MS-liquid chromatography assay for simultaneous determination of tegafur, 5-fluorouracil and 5-fluorodihydrouracil in plasma 824(2005)153

Ultraviolet

Development and validation of a high performance liquid chromatography assay for 17α -methyltestosterone in fish feed 824(2005)107

Uric acid

Determination of uric acid in urine, saliva and calcium oxalate renal calculi by high-performance liquid chromatography/mass spectrometry 824(2005)175

Urine

Tetrahydrocurcumin in plasma and urine: Quantitation by high performance liquid chromatography 824(2005)206

Validamycin A

Quantitative analysis of valienamine in the microbial degradation of validamycin A after derivatization with p-nitrofluorobenzene by reversed-phase high-performance liquid chromatography 824(2005)341

Validation

A liquid chromatographic–tandem mass spectrometric method for the determination of two selective thymidylate synthase inhibitors, BGC945 and BGC638, in mouse plasma 824(2005)181

Valienamine

Quantitative analysis of valienamine in the microbial degradation of validamycin A after derivatization with *p*-nitrofluorobenzene by reversed-phase high-performance liquid chromatography 824(2005)341

Vitamin (

High-throughput analysis of Vitamin C in human plasma with the use of HPLC with monolithic column and UV-detection 824(2005)132

Zinc reduction

A novel, sensitive method for determining benzo[a]pyrene-diones using high-performance liquid chromatography with post-column zinc reduction 824(2005)166