

Keyword Index Vol. 829

Acetyl-11-keto- α -boswellic acid

High-performance liquid chromatographic determination of acetyl-11-keto- α -boswellic acid, a novel pentacyclic triterpenoid, in plasma using a fluorinated stationary phase and photodiode array detection: Application in pharmacokinetic studies 829(2005)144

Anabolic steroid

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

Antisense oligonucleotide

Determination of GTI-2040, a novel antisense oligonucleotide, in human plasma by using HPLC combined with solid phase and liquid–liquid extractions 829(2005)45

Arbutin

Microdialysis sampling coupled to on-line high-performance liquid chromatography for determination of arbutin in whitening cosmetics 829(2005)149

Assay technologies

Receptor–ligand binding assays: Technologies and Applications 829(2005)1

Boldenone

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

Bupropion

Validation of a liquid chromatography–mass spectrometry method to assess the metabolism of bupropion in rat everted gut sacs 829(2005)50

Cancer chemoprevention

Simultaneous determination of myristyl nicotinate, nicotinic acid, and nicotinamide in rabbit plasma by liquid chromatography–tandem mass spectrometry using methyl ethyl ketone as a deproteinization solvent 829(2005)123

Capillary

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

Capillary electrophoresis

Identification of kinetin and kinetin riboside in coconut (*Cocos nucifera* L.) water using a combined approach of liquid chromatography–tandem mass spectrometry, high performance liquid chromatography and capillary electrophoresis 829(2005)26

Chick chorioallantoic membrane

High-performance liquid chromatographic determination of acetyl-11-keto- α -boswellic acid, a novel pentacyclic triterpenoid, in plasma using a fluorinated stationary phase and photodiode array detection: Application in pharmacokinetic studies 829(2005)144

Clinical glycomics

Oligosaccharide analysis by capillary-scale high-pH anion-exchange chromatography with on-line ion-trap mass spectrometry 829(2005)136

Coconut water

Identification of kinetin and kinetin riboside in coconut (*Cocos nucifera* L.) water using a combined approach of liquid chromatography–tandem mass spectrometry, high performance liquid chromatography and capillary electrophoresis 829(2005)26

Cytochrome P450

Validation of a liquid chromatography–mass spectrometry method to assess the metabolism of bupropion in rat everted gut sacs 829(2005)50

Dalbergia odorifera

Identification and determination of major flavonoids in rat serum by HPLC–UV and HPLC–MS methods following oral administration of *Dalbergia odorifera* extract 829(2005)35

O-Desmethyl venlafaxine

Liquid chromatography–tandem mass spectrometry (LC–MS–MS) method for simultaneous determination of venlafaxine and its active metabolite *O*-desmethyl venlafaxine in human plasma 829(2005)75

Doping control

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

Doxil

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

Doxorubicin

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

Electrophoresis

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

Electrospray mass spectrometry

LC–MS solvent composition monitoring and chromatography alignment using mobile phase tracer molecules 829(2005)107

Exercise

Erratum to “Urinary high performance reverse phase chromatography cortisol and cortisone analyses before and at the end of a race in elite cyclists” 829(2005)154

Flavonoids

Identification and determination of major flavonoids in rat serum by HPLC–UV and HPLC–MS methods following oral administration of *Dalbergia odorifera* extract 829(2005)35

Formononetin

Identification and determination of major flavonoids in rat serum by HPLC–UV and HPLC–MS methods following oral administration of *Dalbergia odorifera* extract 829(2005)35

G_{M1}-gangliosidosis

Oligosaccharide analysis by capillary-scale high-pH anion-exchange chromatography with on-line ion-trap mass spectrometry 829(2005)136

GTI-2040

Determination of GTI-2040, a novel antisense oligonucleotide, in human plasma by using HPLC combined with solid phase and liquid–liquid extractions 829(2005)45

Gut sac

Validation of a liquid chromatography–mass spectrometry method to assess the metabolism of bupropion in rat everted gut sacs 829(2005)50

High-performance liquid chromatography

Microdialysis sampling coupled to on-line high-performance liquid chromatography for determination of arbutin in whitening cosmetics 829(2005)149

High-throughput screening

Receptor–ligand binding assays: Technologies and Applications 829(2005)1

Horse

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

HPAEC-PAD

Oligosaccharide analysis by capillary-scale high-pH anion-exchange chromatography with on-line ion-trap mass spectrometry 829(2005)136

HPLC-MS/MS

Simultaneous determination of HIV protease inhibitors amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir and saquinavir in human plasma by high-performance liquid chromatography–tandem mass spectrometry 829(2005)82

11 β -HSD

Erratum to “Urinary high performance reverse phase chromatography cortisol and cortisone analyses before and at the end of a race in elite cyclists” 829(2005)154

Human plasma

Quantification of NC100668, a new tracer for imaging of venous thromboembolism, in human plasma using reversed-phase liquid chromatography coupled with electrospray ionization ion-trap mass spectrometry 829(2005)91

Rapid and sensitive method for the determination of sertraline in human plasma using liquid chromatography–tandem mass spectrometry (LC–MS/MS) 829(2005)69

Liquid chromatography–tandem mass spectrometry (LC–MS–MS) method for simultaneous determination of venlafaxine and its active metabolite *O*-desmethyl venlafaxine in human plasma 829(2005)75

Hydroxybupropion

Validation of a liquid chromatography–mass spectrometry method to assess the metabolism of bupropion in rat everted gut sacs 829(2005)50

Imaging

Quantification of NC100668, a new tracer for imaging of venous thromboembolism, in human plasma using reversed-phase liquid chromatography coupled with electrospray ionization ion-trap mass spectrometry 829(2005)91

Inulin

Oligosaccharide analysis by capillary-scale high-pH anion-exchange chromatography with on-line ion-trap mass spectrometry 829(2005)136

Ion-trap

Quantification of NC100668, a new tracer for imaging of venous thromboembolism, in human plasma using reversed-phase liquid chromatography coupled with electrospray ionization ion-trap mass spectrometry 829(2005)91

Kinetin

Identification of kinetin and kinetin riboside in coconut (*Cocos nucifera* L.) water using a combined approach of liquid chromatography–tandem mass spectrometry, high performance liquid chromatography and capillary electrophoresis 829(2005)26

Kinetin riboside

Identification of kinetin and kinetin riboside in coconut (*Cocos nucifera* L.) water using a combined approach of liquid chromatography–tandem mass spectrometry, high performance liquid chromatography and capillary electrophoresis 829(2005)26

Laser-induced fluorescence

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

LC–MS

Validation of a liquid chromatography–mass spectrometry method to assess the metabolism of bupropion in rat everted gut sacs 829(2005)50

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

Quantification of NC100668, a new tracer for imaging of venous thromboembolism, in human plasma using reversed-phase liquid chromatography coupled with electrospray ionization ion-trap mass spectrometry 829(2005)91

LC–MS/MS

Identification of the major metabolites of resveratrol in rat urine by HPLC–MS/MS 829(2005)97

Rapid and sensitive method for the determination of sertraline in human plasma using liquid chromatography–tandem mass spectrometry (LC–MS/MS) 829(2005)69

Simultaneous determination of myristyl nicotinate, nicotinic acid, and nicotinamide in rabbit plasma by liquid chromatography–tandem mass spectrometry using methyl ethyl ketone as a deproteinization solvent 829(2005)123

LC–MS–MS

Liquid chromatography–tandem mass spectrometry (LC–MS–MS) method for simultaneous determination of venlafaxine and its active metabolite *O*-desmethyl venlafaxine in human plasma 829(2005)75

Liposome

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

Liquid chromatography

Identification of kinetin and kinetin riboside in coconut (*Cocos nucifera* L.) water using a combined approach of liquid chromatography–tandem mass spectrometry, high performance liquid chromatography and capillary electrophoresis 829(2005)26

LC–MS solvent composition monitoring and chromatography alignment using mobile phase tracer molecules 829(2005)107

Metabolism

Validation of a liquid chromatography–mass spectrometry method to assess the metabolism of bupropion in rat everted gut sacs 829(2005)50

Identification of the major metabolites of resveratrol in rat urine by HPLC–MS/MS 829(2005)97

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

(3R)-4'-methoxy-2',3,7-trihydroxy-isoflavanone

Identification and determination of major flavonoids in rat serum by HPLC–UV and HPLC–MS methods following oral administration of *Dalbergia odorifera* extract 829(2005)35

Micellar electrokinetic capillary chromatography

Micellar electrokinetic capillary chromatography reveals differences in intracellular metabolism between liposomal and free doxorubicin treatment of human leukemia cells 829(2005)115

Microdialysis sampling

Microdialysis sampling coupled to on-line high-performance liquid chromatography for determination of arbutin in whitening cosmetics 829(2005)149

Myristyl nicotinate

Simultaneous determination of myristyl nicotinate, nicotinic acid, and nicotinamide in rabbit plasma by liquid chromatography–tandem mass spectrometry using methyl ethyl ketone as a deproteinization solvent 829(2005)123

NC100668

Quantification of NC100668, a new tracer for imaging of venous thromboembolism, in human plasma using reversed-phase liquid chromatography coupled with electrospray ionization ion-trap mass spectrometry 829(2005)91

Nicotinamide

Simultaneous determination of myristyl nicotinate, nicotinic acid, and nicotinamide in rabbit plasma by liquid chromatography–tandem mass spectrometry using methyl ethyl ketone as a deproteinization solvent 829(2005)123

Nicotinic acid

Simultaneous determination of myristyl nicotinate, nicotinic acid, and nicotinamide in rabbit plasma by liquid chromatography–tandem mass spectrometry using methyl ethyl ketone as a deproteinization solvent 829(2005)123

On-line desalter

Oligosaccharide analysis by capillary-scale high-pH anion-exchange chromatography with on-line ion-trap mass spectrometry 829(2005)136

Pentacyclic triterpenoids

High-performance liquid chromatographic determination of acetyl-11-keto- α -boswellic acid, a novel pentacyclic triterpenoid, in plasma using a fluorinated stationary phase and photodiode array detection: Application in pharmacokinetic studies 829(2005)144

Pentafluorophenylpropyl-silica

High-performance liquid chromatographic determination of acetyl-11-keto- α -boswellic acid, a novel pentacyclic triterpenoid, in plasma using a fluorinated stationary phase and photodiode array detection: Application in pharmacokinetic studies 829(2005)144

Peptide

LC–MS solvent composition monitoring and chromatography alignment using mobile phase tracer molecules 829(2005)107

Pharmacokinetics

High-performance liquid chromatographic determination of acetyl-11-keto- α -boswellic acid, a novel pentacyclic triterpenoid, in plasma using a fluorinated stationary phase and photodiode array detection: Application in pharmacokinetic studies 829(2005)144

Plasma

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

Simultaneous determination of myristyl nicotinate, nicotinic acid, and nicotinamide in rabbit plasma by liquid chromatography–tandem mass spectrometry using methyl ethyl ketone as a deproteinization solvent 829(2005)123

Polyamide SPE

Identification of the major metabolites of resveratrol in rat urine by HPLC–MS/MS 829(2005)97

Protease inhibitors

Simultaneous determination of HIV protease inhibitors amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir and saquinavir in human plasma by high-performance liquid chromatography–tandem mass spectrometry 829(2005)82

Proteomics

LC–MS solvent composition monitoring and chromatography alignment using mobile phase tracer molecules 829(2005)107

Quantitative receptor assay

Receptor–ligand binding assays: Technologies and Applications 829(2005)1

Receptor–ligand binding

Receptor–ligand binding assays: Technologies and Applications 829(2005)1

Recombinant receptor expression

Receptor–ligand binding assays: Technologies and Applications 829(2005)1

Sativanone

Identification and determination of major flavonoids in rat serum by HPLC–UV and HPLC–MS methods following oral administration of *Dalbergia odorifera* extract 829(2005)35

Selected reaction monitoring

Identification of kinetin and kinetin riboside in coconut (*Cocos nucifera* L.) water using a combined approach of liquid chromatography–tandem mass spectrometry, high performance liquid chromatography and capillary electrophoresis 829(2005)26

Selective reaction monitoring (SRM)

Rapid and sensitive method for the determination of sertraline in human plasma using liquid chromatography–tandem mass spectrometry (LC–MS/MS) 829(2005)69

Sertraline

Rapid and sensitive method for the determination of sertraline in human plasma using liquid chromatography–tandem mass spectrometry (LC–MS/MS) 829(2005)69

Solid-phase extraction

Determination of GTI-2040, a novel antisense oligonucleotide, in human plasma by using HPLC combined with solid phase and liquid-liquid extractions 829(2005)45

Rapid and sensitive method for the determination of sertraline in human plasma using liquid chromatography-tandem mass spectrometry (LC-MS/MS) 829(2005)69

Erratum to "Urinary high performance reverse phase chromatography cortisol and cortisone analyses before and at the end of a race in elite cyclists" 829(2005)154

Tandem mass spectrometry

Identification of kinetin and kinetin riboside in coconut (*Cocos nucifera* L.) water using a combined approach of liquid chromatography-tandem mass spectrometry, high performance liquid chromatography and capillary electrophoresis 829(2005)26

Testosterone

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

Therapeutic drug monitoring

Simultaneous determination of HIV protease inhibitors amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir and saquinavir in human plasma by high-performance liquid chromatography-tandem mass spectrometry 829(2005)82

THG

Detection, quantification and confirmation of anabolic steroids in equine plasma by liquid chromatography and tandem mass spectrometry 829(2005)56

Tracer molecules

LC-MS solvent composition monitoring and chromatography alignment using mobile phase tracer molecules 829(2005)107

Trans-resveratrol

Identification of the major metabolites of resveratrol in rat urine by HPLC-MS/MS 829(2005)97

Validation

Microdialysis sampling coupled to on-line high-performance liquid chromatography for determination of arbutin in whitening cosmetics 829(2005)149

Venlafaxine

Liquid chromatography-tandem mass spectrometry (LC-MS-MS) method for simultaneous determination of venlafaxine and its active metabolite *O*-desmethyl venlafaxine in human plasma 829(2005)75

Venous thromboembolism

Quantification of NC100668, a new tracer for imaging of venous thromboembolism, in human plasma using reversed-phase liquid chromatography coupled with electrospray ionization ion-trap mass spectrometry 829(2005)91

Vestitone

Identification and determination of major flavonoids in rat serum by HPLC-UV and HPLC-MS methods following oral administration of *Dalbergia odorifera* extract 829(2005)35

Whitening cosmetics

Microdialysis sampling coupled to on-line high-performance liquid chromatography for determination of arbutin in whitening cosmetics 829(2005)149