

Keyword Index Vol. 844

Accuracy profile

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Aconitine

Characterization of metabolites and cytochrome P450 isoforms involved in the microsomal metabolism of aconitine 844(2006) 292

Acteoside

Determination of acteoside in *Cistanche deserticola* and *Boschniakia rossica* and its pharmacokinetics in freely-moving rats using LC-MS/MS 844(2006) 89

s-Adenosylhomocysteine

Quantitative analysis of s-adenosylmethionine and s-adenosylhomocysteine in neurulation-stage mouse embryos by liquid chromatography tandem mass spectrometry 844(2006) 112

s-Adenosylmethionine

Quantitative analysis of s-adenosylmethionine and s-adenosylhomocysteine in neurulation-stage mouse embryos by liquid chromatography tandem mass spectrometry 844(2006) 112

Algal dietary supplement

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Alzheimer's disease

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Antibody

Rapid antibody screening by membrane chromatographic immunoassay technique 844(2006) 163

Antigen

Rapid antibody screening by membrane chromatographic immunoassay technique 844(2006) 163

Aqueous biphasic extraction chromatography

The opposite effect of temperature on polyethylene glycol-based aqueous biphasic systems versus aqueous biphasic extraction chromatographic resins 844(2006) 23

Aqueous two-phase systems

The opposite effect of temperature on polyethylene glycol-based aqueous biphasic systems versus aqueous biphasic extraction chromatographic resins 844(2006) 23

Simplified two-stage method to B-phycoerythrin recovery from *Porphyridium cruentum* 844(2006) 39

Aqueous–aqueous two-phase systems

Aqueous–aqueous two-phase systems composed of low molecular weight of polyethylene glycols and dextrans for counter-current chromatographic purification of proteins 844(2006) 217

Artemia assay

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Automated blood sampling

Determination of acteoside in *Cistanche deserticola* and *Boschniakia rossica* and its pharmacokinetics in freely-moving rats using LC-MS/MS 844(2006) 89

Avermectins

Review of methodology for the determination of macrocyclic lactone residues in biological matrices 844(2006) 175

Azelnidipine

Simultaneous determination of azelnidipine and two metabolites in human plasma using liquid chromatography-tandem mass spectrometry 844(2006) 45

Baicalein

Liquid chromatography with tandem mass spectrometry for the simultaneous determination of baicalein, baicalin, oroxylin A and wogonin in rat plasma 844(2006) 261

Baicalin

Liquid chromatography with tandem mass spectrometry for the simultaneous determination of baicalein, baicalin, oroxylin A and wogonin in rat plasma 844(2006) 261

Biapenem

A simple and rapid determination of biapenem in plasma by high-performance liquid chromatography 844(2006) 148

Bioavailability

Measurement and pharmacokinetic study of plumbagin in a conscious freely moving rat using liquid chromatography/tandem mass spectrometry 844(2006) 1

Determination of acteoside in *Cistanche deserticola* and *Boschniakia rossica* and its pharmacokinetics in freely-moving rats using LC-MS/MS 844(2006) 89

Development and validation of a sensitive HPLC-ESI-MS/MS method for the direct determination of glucosamine in human plasma 844(2006) 119

Determination of echinacoside in rat serum by reversed-phase high-performance liquid chromatography with ultraviolet detection and its application to pharmacokinetics and bioavailability 844(2006) 308

Biomarker

Quantification of the urinary concentrations of parabens in humans by on-line solid phase extraction-high performance liquid chromatography–isotope dilution tandem mass spectrometry 844(2006) 53

Biomonitoring

Quantification of the urinary concentrations of parabens in humans by on-line solid phase extraction-high performance liquid chromatography–isotope dilution tandem mass spectrometry 844(2006) 53

Biotinidase

Determination of specific activities and kinetic constants of biotinidase and lipamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

Bovine plasma

Confirmation of diminazene diaceturate in bovine plasma using electrospray liquid chromatography–mass spectrometry 844(2006) 127

Bovine serum albumin

Hydrophobic interaction adsorption of whey proteins: Effect of temperature and salt concentration and thermodynamic analysis 844(2006) 6

B-phycoerythrin

Simplified two-stage method to B-phycoerythrin recovery from *Porphyridium cruentum* 844(2006) 39

Brain

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Buspiron

Quantification of a cytochrome P450 3A4 substrate, bupirone, in human plasma by liquid chromatography–tandem mass spectrometry 844(2006) 235

Camptothecin analogs

Validation of an HPLC method for analysis of DB-67 and its water soluble pro-drug in mouse plasma 844(2006) 15

Capillary electrophoresis

Characterization of proteinase activation dynamics by capillary electrophoresis conjugating with fluorescent protein-based probe 844(2006) 158

Carry over

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Caspase-2

Characterization of proteinase activation dynamics by capillary electrophoresis conjugating with fluorescent protein-based probe 844(2006) 158

Catecholamines

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Chemical fingerprint

Chemical fingerprint analysis of rhizomes of *Gymnadenia conopsea* by HPLC–DAD–MSⁿ 844(2006) 301

Chiral thiol

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Chromatography

Rapid antibody screening by membrane chromatographic immunoassay technique 844(2006) 163

A simple and sensitive bioanalytical assay for simultaneous determination of omeprazole and its three major metabolites in human blood plasma using RP-HPLC after a simple liquid–liquid extraction procedure 844(2006) 314

Column-switching

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Cotinine

Simple, rapid and sensitive assay method for simultaneous quantification of urinary nicotine and cotinine using gas chromatography–mass spectrometry 844(2006) 322

Counter-current chromatography

Aqueous–aqueous two-phase systems composed of low molecular weight of polyethylene glycols and dextrans for counter-current chromatographic purification of proteins 844(2006) 217

Cyanobacteria

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Cytochrome P450

Characterization of metabolites and cytochrome P450 isoforms involved in the microsomal metabolism of aconitine 844(2006) 292

Dark red kidney bean

A hemagglutinin with mitogenic activity from dark red kidney beans 844(2006) 213

DB-67

Validation of an HPLC method for analysis of DB-67 and its water soluble pro-drug in mouse plasma 844(2006) 15

Derivatisation

Review of methodology for the determination of macrocyclic lactone residues in biological matrices 844(2006) 175

Desmethylselegiline

Simultaneous determination of selegiline and desmethylselegiline in human body fluids by headspace solid-phase microextraction and gas chromatography–mass spectrometry 844(2006) 283

Determination

Optimization and validation of the direct HPLC method for the determination of moxifloxacin in plasma 844(2006) 104

Diminazene diaceturate

Confirmation of diminazene diaceturate in bovine plasma using electrospray liquid chromatography–mass spectrometry 844(2006) 127

DNA adducts

Quantification of *O*⁶-methyl and *O*⁶-ethyl deoxyguanosine adducts in C57BL/6N/*Tk*^{+/-} mice using LC/MS/MS 844(2006) 60

Dopamine

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Doping control

Short-term stability of testosterone and epitestosterone conjugates in urine samples: Quantification by liquid chromatography–linear ion trap mass spectrometry 844(2006) 168

Drug assay

Quantitative determination of nystatin in human plasma using LC–MS after inhalative administration in healthy subjects 844(2006) 84

Dynamics

Characterization of proteinase activation dynamics by capillary electrophoresis conjugating with fluorescent protein-based probe 844(2006) 158

ECD

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Echinacoside

Determination of echinacoside in rat serum by reversed-phase high-performance liquid chromatography with ultraviolet detection and its application to pharmacokinetics and bioavailability 844(2006) 308

Electrospray ionisation mass spectrometry

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography–tandem electrospray ionisation mass spectrometry 844(2006) 223

Electrospray ionization

Quantification of a cytochrome P450 3A4 substrate, buspirone, in human plasma by liquid chromatography–tandem mass spectrometry 844(2006) 235

Environmental tobacco smoke (ETS)

Simple, rapid and sensitive assay method for simultaneous quantification of urinary nicotine and cotinine using gas chromatography–mass spectrometry 844(2006) 322

Enzyme kinetics

Determination of specific activities and kinetic constants of biotinidase and lipamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

Epilepsy

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Epinephrine

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Epitestosterone

Short-term stability of testosterone and epitestosterone conjugates in urine samples: Quantification by liquid chromatography–linear ion trap mass spectrometry 844(2006) 168

O⁶-Ethyl-2'-deoxyguanosine

Quantification of O⁶-methyl and O⁶-ethyl deoxyguanosine adducts in C57BL/6N/Tk^{+/−} mice using LC/MS/MS 844(2006) 60

Exposure

Quantification of the urinary concentrations of parabens in humans by on-line solid phase extraction-high performance liquid chromatography–isotope dilution tandem mass spectrometry 844(2006) 53

Extraction

Review of methodology for the determination of macrocyclic lactone residues in biological matrices 844(2006) 175

Fluorescent protein

Characterization of proteinase activation dynamics by capillary electrophoresis conjugating with fluorescent protein-based probe 844(2006) 158

Garcinia extracts

Identification and quantification of two biologically active polyisoprenylated benzophenones xanthochymol and isoxanthochymol in *Garcinia* species using liquid chromatography–tandem mass spectrometry 844(2006) 67

GC–MS

Simple, rapid and sensitive assay method for simultaneous quantification of urinary nicotine and cotinine using gas chromatography–mass spectrometry 844(2006) 322

Genotyping

Isotachopheresis preconcentration integrated microfluidic chip for highly sensitive genotyping of the hepatitis B virus 844(2006) 32

Glucosamine

Development and validation of a sensitive HPLC–ESI-MS/MS method for the direct determination of glucosamine in human plasma 844(2006) 119

Glucosinolates

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography–tandem electrospray ionisation mass spectrometry 844(2006) 223

Glucosyltransferase

Aqueous–aqueous two-phase systems composed of low molecular weight of polyethylene glycols and dextrans for counter-current chromatographic purification of proteins 844(2006) 217

Glucuronide

Short-term stability of testosterone and epitestosterone conjugates in urine samples: Quantification by liquid chromatography–linear ion trap mass spectrometry 844(2006) 168

Gynadenia conopsea

Chemical fingerprint analysis of rhizomes of *Gynadenia conopsea* by HPLC–DAD–MSⁿ 844(2006) 301

Hemagglutinin

A hemagglutinin with mitogenic activity from dark red kidney beans 844(2006) 213

Hepatitis virus B

Isotachopheresis preconcentration integrated microfluidic chip for highly sensitive genotyping of the hepatitis B virus 844(2006) 32

Herbal medicine

Measurement and pharmacokinetic study of plumbagin in a conscious freely moving rat using liquid chromatography/tandem mass spectrometry 844(2006) 1

High-performance liquid chromatography

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

Histamine

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

Histamine absorption

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

Histamine metabolism

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

Histamine release

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

Histidine-tag

The effect of the hexahistidine-tag in the oligomerization of HSC70 constructs 844(2006) 328

HPLC

Development and validation of a sensitive HPLC–ESI-MS/MS method for the direct determination of glucosamine in human plasma 844(2006) 119

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Determination of echinacoside in rat serum by reversed-phase high-performance liquid chromatography with ultraviolet detection and its application to pharmacokinetics and bioavailability 844(2006) 308

A simple and sensitive bioanalytical assay for simultaneous determination of omeprazole and its three major metabolites in human blood plasma using RP-HPLC after a simple liquid–liquid extraction procedure 844(2006) 314

HPLC Fluorescence

Review of methodology for the determination of macrocyclic lactone residues in biological matrices 844(2006) 175

HPLC–DAD–MSⁿ

Chemical fingerprint analysis of rhizomes of *Gymnadenia conopsea* by HPLC–DAD–MSⁿ 844(2006) 301

HPLC-fluorimetric enzyme assay

Determination of specific activities and kinetic constants of biotinidase and lipamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

HSC70

The effect of the hexahistidine-tag in the oligomerization of HSC70 constructs 844(2006) 328

5-HT_{1B/1D} receptor agonist

Development and validation of a selective and robust LC–MS/MS method for high-throughput quantifying rizatriptan in small plasma samples: Application to a clinical pharmacokinetic study 844(2006) 268

Human

Quantification of the urinary concentrations of parabens in humans by on-line solid phase extraction-high performance liquid chromatography–isotope dilution tandem mass spectrometry 844(2006) 53

Human plasma

Quantification of a cytochrome P450 3A4 substrate, buspirone, in human plasma by liquid chromatography–tandem mass spectrometry 844(2006) 235

Hydrophobic interaction

Hydrophobic interaction adsorption of whey proteins: Effect of temperature and salt concentration and thermodynamic analysis 844(2006) 6

Rapid antibody screening by membrane chromatographic immunoassay technique 844(2006) 163

5-Hydroxyomeprazole

A simple and sensitive bioanalytical assay for simultaneous determination of omeprazole and its three major metabolites in human blood plasma using RP-HPLC after a simple liquid–liquid extraction procedure 844(2006) 314

Iberin

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography–tandem electrospray ionisation mass spectrometry 844(2006) 223

Identification

Chemical fingerprint analysis of rhizomes of *Gymnadenia conopsea* by HPLC–DAD–MSⁿ 844(2006) 301

Immobilized metal affinity chromatography (IMAC)

The effect of the hexahistidine-tag in the oligomerization of HSC70 constructs 844(2006) 328

Immunoassay

Rapid antibody screening by membrane chromatographic immunoassay technique 844(2006) 163

Inhaled aerosol

Quantitative determination of nystatin in human plasma using LC–MS after inhalative administration in healthy subjects 844(2006) 84

Intestine

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

Isolation

A hemagglutinin with mitogenic activity from dark red kidney beans 844(2006) 213

Isotachopheresis

Isotachopheresis preconcentration integrated microfluidic chip for highly sensitive genotyping of the hepatitis B virus 844(2006) 32

Isotherm

Hydrophobic interaction adsorption of whey proteins: Effect of temperature and salt concentration and thermodynamic analysis 844(2006) 6

Isothiocyanates

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography–tandem electrospray ionisation mass spectrometry 844(2006) 223

Isoxanthochymol

Identification and quantification of two biologically active polyisoprenylated benzophenones xanthochymol and isoxanthochymol in *Garcinia* species using liquid chromatography–tandem mass spectrometry 844(2006) 67

***Lactobacillus casei* (Shirota)**

Determination of specific activities and kinetic constants of biotinidase and lipamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

β-Lactoglobulin

Hydrophobic interaction adsorption of whey proteins: Effect of temperature and salt concentration and thermodynamic analysis 844(2006) 6

LC–ESI–MS

Determination of total tiopronin in human plasma by LC–ESI–MS using tris (2-carboxy-ethyl) phosphine as reducing reagent and methyl acrylate as derivatization reagent for the thiol group 844(2006) 153

LC–MS

Quantitative determination of nystatin in human plasma using LC–MS after inhalative administration in healthy subjects 844(2006) 84

LC–MS/MS

Determination of acteoside in *Cistanche deserticola* and *Boschniakia rossica* and its pharmacokinetics in freely-moving rats using LC–MS/MS 844(2006) 89

Review of methodology for the determination of macrocyclic lactone residues in biological matrices 844(2006) 175

Liquid chromatography with tandem mass spectrometry for the simultaneous determination of baicalein, baicalin, oroxylin A and wogonin in rat plasma 844(2006) 261

Development and validation of a selective and robust LC-MS/MS method for high-throughput quantifying rizatriptan in small plasma samples: Application to a clinical pharmacokinetic study 844(2006) 268

LEW rat

Determination of specific activities and kinetic constants of biotinidase and lipamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

Lipoamidase (lipoyl-X hydrolase)

Determination of specific activities and kinetic constants of biotinidase and lipamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

Liquid chromatography

Identification and quantification of two biologically active polyisoprenylated benzophenones xanthochymol and isoxanthochymol in *Garcinia* species using liquid chromatography-tandem mass spectrometry 844(2006) 67

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography-tandem electrospray ionisation mass spectrometry 844(2006) 223

Quantification of a cytochrome P450 3A4 substrate, buspirone, in human plasma by liquid chromatography-tandem mass spectrometry 844(2006) 235

Liquid chromatography-mass spectrometry

Short-term stability of testosterone and epitestosterone conjugates in urine samples: Quantification by liquid chromatography-linear ion trap mass spectrometry 844(2006) 168

Mass spectrometry

Quantification of *O*⁶-methyl and *O*⁶-ethyl deoxyguanosine adducts in C57BL/6N/*Tk*^{+/-} mice using LC/MS/MS 844(2006) 60

Development and validation of a sensitive HPLC-ESI-MS/MS method for the direct determination of glucosamine in human plasma 844(2006) 119

Confirmation of diminazene diacetate in bovine plasma using electrospray liquid chromatography-mass spectrometry 844(2006) 127

Measurement uncertainty

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Membrane

Rapid antibody screening by membrane chromatographic immunoassay technique 844(2006) 163

Mercapturic acid pathway

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography-tandem electrospray ionisation mass spectrometry 844(2006) 223

Metabolism

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography-tandem electrospray ionisation mass spectrometry 844(2006) 223

Characterization of metabolites and cytochrome P450 isoforms involved in the microsomal metabolism of aconitine 844(2006) 292

Metabolites

TopCount coupled to ultra-performance liquid chromatography for the profiling of radiolabeled drug metabolites in complex biological samples 844(2006) 96

Characterization of metabolites and cytochrome P450 isoforms involved in the microsomal metabolism of aconitine 844(2006) 292

Method validation

Validation of an HPLC method for analysis of DB-67 and its water soluble prodrug in mouse plasma 844(2006) 15

Methyl acrylate

Determination of total tiopronin in human plasma by LC-ESI-MS using tris (2-carboxy-ethyl) phosphine as reducing reagent and methyl acrylate as derivatization reagent for the thiol group 844(2006) 153

1-Methylhistamine

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

***O*⁶-Methyl-2'-deoxyguanosine**

Quantification of *O*⁶-methyl and *O*⁶-ethyl deoxyguanosine adducts in C57BL/6N/*Tk*^{+/-} mice using LC/MS/MS 844(2006) 60

Microcystin (MC)

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Microcystis

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Microfluidic chip

Isotachopheresis preconcentration integrated microfluidic chip for highly sensitive genotyping of the hepatitis B virus 844(2006) 32

Microsome

Characterization of metabolites and cytochrome P450 isoforms involved in the microsomal metabolism of aconitine 844(2006) 292

Milbemycins

Review of methodology for the determination of macrocyclic lactone residues in biological matrices 844(2006) 175

Molecularly imprinted polymer

Molecularly imprinted solid-phase extraction for rapid screening of mycophenolic acid in human plasma 844(2006) 142

Molecularly imprinted solid-phase extraction

Molecularly imprinted solid-phase extraction for rapid screening of mycophenolic acid in human plasma 844(2006) 142

Mouse Bioassay

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Mouse embryo

Quantitative analysis of s-adenosylmethionine and s-adenosylhomocysteine in neurulation-stage mouse embryos by liquid chromatography tandem mass spectrometry 844(2006) 112

Moxifloxacin

Optimization and validation of the direct HPLC method for the determination of moxifloxacin in plasma 844(2006) 104

Multidimensional

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Multiple reaction monitoring

Identification and quantification of two biologically active polyisoprenylated benzophenones xanthochymol and isoxanthochymol in *Garcinia* species using liquid chromatography–tandem mass spectrometry 844(2006) 67

Mutagenesis

The effect of the hexahistidine-tag in the oligomerization of HSC70 constructs 844(2006) 328

Mycophenolic acid

Molecularly imprinted solid-phase extraction for rapid screening of mycophenolic acid in human plasma 844(2006) 142

Neurochemistry

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Nicotine

Simple, rapid and sensitive assay method for simultaneous quantification of urinary nicotine and cotinine using gas chromatography–mass spectrometry 844(2006) 322

NMDA receptor

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Norepinephrine

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Nystatin

Quantitative determination of nystatin in human plasma using LC–MS after inhalative administration in healthy subjects 844(2006) 84

Oligomerization

The effect of the hexahistidine-tag in the oligomerization of HSC70 constructs 844(2006) 328

Omeprazole

A simple and sensitive bioanalytical assay for simultaneous determination of omeprazole and its three major metabolites in human blood plasma using RP-HPLC after a simple liquid–liquid extraction procedure 844(2006) 314

Omeprazole sulfide

A simple and sensitive bioanalytical assay for simultaneous determination of omeprazole and its three major metabolites in human blood plasma using RP-HPLC after a simple liquid–liquid extraction procedure 844(2006) 314

Omeprazole sulfone

A simple and sensitive bioanalytical assay for simultaneous determination of omeprazole and its three major metabolites in human blood plasma using RP-HPLC after a simple liquid–liquid extraction procedure 844(2006) 314

On-line SPE

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Oroxylin A

Liquid chromatography with tandem mass spectrometry for the simultaneous determination of baicalein, baicalin, oroxylin A and wogonin in rat plasma 844(2006) 261

Osteoarthritis

Development and validation of a sensitive HPLC–ESI-MS/MS method for the direct determination of glucosamine in human plasma 844(2006) 119

Paraben

Quantification of the urinary concentrations of parabens in humans by on-line solid phase extraction-high performance liquid chromatography–isotope dilution tandem mass spectrometry 844(2006) 53

Pentafluorophenylpropyl

Quantitative analysis of s-adenosylmethionine and s-adenosylhomocysteine in neurulation-stage mouse embryos by liquid chromatography tandem mass spectrometry 844(2006) 112

Pharmacokinetics

Measurement and pharmacokinetic study of plumbagin in a conscious freely moving rat using liquid chromatography/tandem mass spectrometry 844(2006) 1

Determination of acteoside in *Cistanche deserticola* and *Boschniakia rossica* and its pharmacokinetics in freely-moving rats using LC–MS/MS 844(2006) 89

Determination of total tiopronin in human plasma by LC–ESI–MS using tris (2-carboxy-ethyl) phosphine as reducing reagent and methyl acrylate as derivatization reagent for the thiol group 844(2006) 153

Development and validation of a selective and robust LC–MS/MS method for high-throughput quantifying rizatriptan in small plasma samples: Application to a clinical pharmacokinetic study 844(2006) 268

Determination of echinacoside in rat serum by reversed-phase high-performance liquid chromatography with ultraviolet detection and its application to pharmacokinetics and bioavailability 844(2006) 308

Phenylethanoid glycoside

Determination of echinacoside in rat serum by reversed-phase high-performance liquid chromatography with ultraviolet detection and its application to pharmacokinetics and bioavailability 844(2006) 308

Plasma

Optimization and validation of the direct HPLC method for the determination of moxifloxacin in plasma 844(2006) 104

Determination of total tiopronin in human plasma by LC–ESI–MS using tris (2-carboxy-ethyl) phosphine as reducing reagent and methyl acrylate as derivatization reagent for the thiol group 844(2006) 153

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography–tandem electrospray ionisation mass spectrometry 844(2006) 223

Plumbagin

Measurement and pharmacokinetic study of plumbagin in a conscious freely moving rat using liquid chromatography/tandem mass spectrometry 844(2006) 1

Plumbago zeylanica L

Measurement and pharmacokinetic study of plumbagin in a conscious freely moving rat using liquid chromatography/tandem mass spectrometry 844(2006) 1

Protein activation

Characterization of proteinase activation dynamics by capillary electrophoresis conjugating with fluorescent protein-based probe 844(2006) 158

Protein phosphatase assay

Potential of a simple solid-phase extraction method coupled to analytical and bio-analytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Proteins

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Aqueous–aqueous two-phase systems composed of low molecular weight of polyethylene glycols and dextrans for counter-current chromatographic purification of proteins 844(2006) 217

Radiochemical detection

TopCount coupled to ultra-performance liquid chromatography for the profiling of radiolabeled drug metabolites in complex biological samples 844(2006) 96

RAM

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Rat plasma

Liquid chromatography with tandem mass spectrometry for the simultaneous determination of baicalein, baicalin, oroxylin A and wogonin in rat plasma 844(2006) 261

Recovery

Simplified two-stage method to B-phycoerythrin recovery from *Porphyridium cruentum* 844(2006) 39

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Restricted access materials

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Reversed phase

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Risk analysis

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Rizatriptan

Development and validation of a selective and robust LC–MS/MS method for high-throughput quantifying rizatriptan in small plasma samples: Application to a clinical pharmacokinetic study 844(2006) 268

RP-HPLC

Optimization and validation of the direct HPLC method for the determination of moxifloxacin in plasma 844(2006) 104

Sample preconcentration

Isotachopheresis preconcentration integrated microfluidic chip for highly sensitive genotyping of the hepatitis B virus 844(2006) 32

Schizophrenia

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Screening

Rapid antibody screening by membrane chromatographic immunoassay technique 844(2006) 163

SEC protein assay

Determination of specific activities and kinetic constants of biotinidase and lipoamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

Selectivity

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Selegiline

Simultaneous determination of selegiline and desmethylselegiline in human body fluids by headspace solid-phase microextraction and gas chromatography–mass spectrometry 844(2006) 283

Serine racemase

Determination of D-serine and related neuroactive amino acids in human plasma by high-performance liquid chromatography with fluorimetric detection 844(2006) 278

Solid-phase extraction (SPE)

Potential of a simple solid-phase extraction method coupled to analytical and bioanalytical methods for an improved determination of microcystins in algal samples 844(2006) 134

Solid-phase microextraction (SPME)

Simultaneous determination of selegiline and desmethylselegiline in human body fluids by headspace solid-phase microextraction and gas chromatography–mass spectrometry 844(2006) 283

SPE

Review of methodology for the determination of macrocyclic lactone residues in biological matrices 844(2006) 175

Stability

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Succinate derivative ester

Chemical fingerprint analysis of rhizomes of *Gymnadenia conopsea* by HPLC–DAD–MSⁿ 844(2006) 301

Sulfate

Short-term stability of testosterone and epitestosterone conjugates in urine samples: Quantification by liquid chromatography–linear ion trap mass spectrometry 844(2006) 168

Sulforaphane

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography–tandem electrospray ionisation mass spectrometry 844(2006) 223

Symmetry

Evaluation of stationary phases for 2-dimensional HPLC of proteins. Part 1. Validation of commercial RP-columns 844(2006) 204

Tandem mass spectrometry

Identification and quantification of two biologically active polyisoprenylated benzophenones xanthochymol and isoxanthochymol in *Garcinia* species using liquid chromatography–tandem mass spectrometry 844(2006) 67

Quantitative analysis of s-adenosylmethionine and s-adenosylhomocysteine in neurulation-stage mouse embryos by liquid chromatography tandem mass spectrometry 844(2006) 112

Quantification of a cytochrome P450 3A4 substrate, buspirone, in human plasma by liquid chromatography–tandem mass spectrometry 844(2006) 235

Temperature effect

The opposite effect of temperature on polyethylene glycol-based aqueous biphasic systems versus aqueous biphasic extraction chromatographic resins 844(2006) 23

Testosterone

Short-term stability of testosterone and epitestosterone conjugates in urine samples: Quantification by liquid chromatography–linear ion trap mass spectrometry 844(2006) 168

Thermodynamic parameters

Hydrophobic interaction adsorption of whey proteins: Effect of temperature and salt concentration and thermodynamic analysis 844(2006) 6

Thermodynamics

The opposite effect of temperature on polyethylene glycol-based aqueous biphasic systems versus aqueous biphasic extraction chromatographic resins 844(2006) 23

Tiopronin

Determination of total tiopronin in human plasma by LC–ESI–MS using tris (2-carboxy-ethyl) phosphine as reducing reagent and methyl acrylate as derivatization reagent for the thiol group 844(2006) 153

Tissue homogenate

Determination of specific activities and kinetic constants of biotinidase and lipoamidase in LEW rat and *Lactobacillus casei* (Shirota) 844(2006) 240

TopCount

TopCount coupled to ultra-performance liquid chromatography for the profiling of radiolabeled drug metabolites in complex biological samples 844(2006) 96

Tris (2-carboxy-ethyl) phosphine

Determination of total tiopronin in human plasma by LC–ESI–MS using tris (2-carboxy-ethyl) phosphine as reducing reagent and methyl acrylate as derivatization reagent for the thiol group 844(2006) 153

Two metabolites

Simultaneous determination of azelnidipine and two metabolites in human plasma using liquid chromatography–tandem mass spectrometry 844(2006) 45

Ultrafiltration

A simple and rapid determination of biapenem in plasma by high-performance liquid chromatography 844(2006) 148

Ultra-performance liquid chromatography

TopCount coupled to ultra-performance liquid chromatography for the profiling of radiolabeled drug metabolites in complex biological samples 844(2006) 96

Urine

Quantification of the urinary concentrations of parabens in humans by on-line solid phase extraction–high performance liquid chromatography–isotope dilution tandem mass spectrometry 844(2006) 53

Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography–tandem electrospray ionisation mass spectrometry 844(2006) 223

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Simple, rapid and sensitive assay method for simultaneous quantification of urinary nicotine and cotinine using gas chromatography–mass spectrometry 844(2006) 322

Ussing chamber

High-performance liquid chromatographic method for the determination of histamine and 1-methylhistamine in biological buffers 844(2006) 335

Validation

Performances of a multidimensional on-line SPE-LC-ECD method for the determination of three major catecholamines in native human urine: Validation, risk and uncertainty assessments 844(2006) 251

Wogonin

Liquid chromatography with tandem mass spectrometry for the simultaneous determination of baicalein, baicalin, oroxylin A and wogonin in rat plasma 844(2006) 261

Xanthochymol

Identification and quantification of two biologically active polyisoprenylated benzophenones xanthochymol and isoxanthochymol in *Garcinia* species using liquid chromatography–tandem mass spectrometry 844(2006) 67