

This article was downloaded by: [Renmin University of China]

On: 13 October 2013, At: 10:22

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Coordination Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gcoo20>

Synthesis, characterization, and antimicrobial activities of two mononuclear metal(II) complexes with (Z)-3-hydroxy-4-(3-hydroxy-3-phenylacryloyl)phenyl benzoate

Suo-Ping Xu ^a, Ban-Feng Ruan ^b, Qi-Yu Pan ^c & Ren-Tao Hu ^c

^a School of Chemistry and Chemical Engineering, Xuzhou Normal University, Xuzhou 221116, P. R. China

^b School of Medical Engineering, Hefei University of Technology, Hefei 230009, P. R. China

^c Anhui Research Institute of Chemical Industry, Hefei 230039, P. R. China

Published online: 14 Jul 2011.

To cite this article: Suo-Ping Xu, Ban-Feng Ruan, Qi-Yu Pan & Ren-Tao Hu (2011) Synthesis, characterization, and antimicrobial activities of two mononuclear metal(II) complexes with (Z)-3-hydroxy-4-(3-hydroxy-3-phenylacryloyl)phenyl benzoate, *Journal of Coordination Chemistry*, 64:14, 2489-2496, DOI: [10.1080/00958972.2011.601459](https://doi.org/10.1080/00958972.2011.601459)

To link to this article: <http://dx.doi.org/10.1080/00958972.2011.601459>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis, characterization, and antimicrobial activities of two mononuclear metal(II) complexes with (*Z*)-3-hydroxy-4-(3-hydroxy-3-phenylacryloyl)phenyl benzoate

SUO-PING XU[†], BAN-FENG RUAN^{*‡}, QI-YU PAN[§] and REN-TAO HU[§]

[†]School of Chemistry and Chemical Engineering, Xuzhou Normal University, Xuzhou
221116, P. R. China

[‡]School of Medical Engineering, Hefei University of Technology, Hefei 230009, P. R. China
[§]Anhui Research Institute of Chemical Industry, Hefei 230039, P. R. China

(Received 26 January 2011; in final form 8 June 2011)

Two mononuclear complexes with a β -diketone ligand (*Z*)-3-hydroxy-4-(3-hydroxy-3-phenylacryloyl)phenyl benzoate (**L**), $[\text{CoL}_2(\text{CH}_3\text{CH}_2\text{OH})_2]$ (**1**), and $[\text{MnL}_2(\text{CH}_3\text{CH}_2\text{OH})_2]$ (**2**) were prepared. Both complexes were characterized by X-ray crystallography, confirming that the central metal(II) are coordinated by four oxygens from two **L** and two oxygens from two ethanols. Both complexes were assayed for *in vitro* antibacterial (*Bacillus subtilis*, *Staphylococcus aureus*, *Streptococcus faecalis*, *Pseudomonas aeruginosa*, *Escherichia coli*, and *Enterobacter cloacae*) activities and showed better antimicrobial activity against Gram positive strains than Gram negative strains.

Keywords: β -Diketone ligand; Mononuclear complexes; Antibacterial activity

1. Introduction

β -Diketones are important intermediates in organic synthesis. The high reactivity of β -diketones such as 1,4-diphenyl-butane-1,3-dione, 1,4-dithiohene-butane-1,3-dione, 1,4-bis-(benzo(1,3)dio-butane-1,3-dione, 1,4-bis-(3-fluoro-4-methoxy-phenyl)-butane-1,3-dione, 3-hydroxy-1,3-diphenyl-propenone, and 1,3-di-furan-2-yl-3-hydroxy-propenone have been reported [1–4]. β -Diketones and their derivatives also have a wide range of applications in heat stabilizers, luminescence, catalysis, solvent extraction, and pharmaceuticals [5–11]. There were many methods for the preparation of β -diketones, but most of them could not provide satisfactory yields for a long time. Therefore, development of direct and efficient procedures for this class of compounds has been the target of synthetic organic chemistry. Recently, an operationally simple and new approach to synthesize β -diketones from zinc-mediated C–C bond sigmatropic rearrangement has been reported [12]. Herein, one bidentate β -diketone, (*Z*)-3-hydroxy-4-(3-hydroxy-3-phenylacryloyl)phenyl benzoate (**L**), was synthesized in good yield by microwave assistance and two mononuclear complexes were obtained by

*Corresponding author. Email: bf_ruan@163.com

reacting **L** with $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ and $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ in ethanol:acetone (1:1). Both complexes were assayed for antibacterial activities against three Gram positive bacterial strains (*Bacillus subtilis*, *Staphylococcus aureus*, and *Streptococcus faecalis*) and three Gram negative bacterial strains (*Escherichia coli*, *Pseudomonas aeruginosa*, and *Enterobacter cloacae*) by the 3-(4,5-dimethyl-2-triazyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) method.

2. Experimental


2.1. Materials and instruments

All chemicals were of reagent grade and used as received. UV spectra were recorded on a U-3000 spectrophotometer. IR spectra were recorded on a Nexus 870 FT-IR. ESI-MS spectra were recorded on a Mariner System 5304 mass spectrometer. Elemental analyses were performed on a CHN-O-Rapid instrument and were within $\pm 0.4\%$ of the theoretical values. Melting points were measured on a Boetius micro melting point apparatus.

2.2. Preparation of **L**, **1**, and **2**

Compound **L** was designed and synthesized from 2,4-dihydroxyacetophenone and benzoyl chloride in acetone by microwave assistance. The ligand was reacted with $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ and $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ in ethanol:acetone (1:1), respectively, and afforded bis(benzoic acid 3-hydroxy-4-(3-hydroxy-3-phenyl-acryloyl)-phenyl ester)-bis-ethanol-cobalt(II) (**1**) and bis(benzoic acid 3-hydroxy-4-(3-hydroxy-3-phenyl-acryloyl)-phenyl ester)-bis-ethanol-manganese(II) (**2**) (scheme 1).

2.2.1. Preparation of **L.** K_2CO_3 (20 g) was slowly added to a round-bottom-flask containing 2,4-dihydroxyacetophenone (3.04 g, 0.02 mol) and benzoyl chloride (5.6 g,



Scheme 1. Synthesis of **L**, **1**, and **2**.

0.04 mol) dissolved in acetone (50 mL). The mixture was microwave-irradiated (90 W) for 2 h and then precipitated. After filtration, the yellow solid was washed with acetone (100 mL) and water (200 mL), dried, and recrystallized from ethanol/acetone (1/1). Yield: 80%, m.p.: 241–243°C (lit. 240–243°C [13]). UV (λ nm): 377.5; 256.5. Selected IR data (cm^{-1} , KBr): 3421.5 (m), 1734.8 (s), 1599.3 (s), 1569.3 (s), 1508.1 (s), 1260.2 (s), 1205.3 (m), 1138.5 (s), 1062.8 (m), 1023.3 (m), 767.0 (m), 710.2 (s); $^1\text{H-NMR}$ (CDCl_3) δ ppm: 15.46 (s, 1H), 12.23 (s, 1H), 8.20 (d, $J=7.1$ Hz, 2H), 7.95 (d, $J=6.8$ Hz, 2H), 7.86 (d, $J=8.8$ Hz, 1H), 7.66 (d, $J=14.8$ Hz, 1H), 7.55 (d, $J=12.6$ Hz, 4H), 6.91 (d, $J=2.4$, 1H), 6.85 (d, $J=11.2$ Hz, 1H), 6.81 (s, 1H). ESI-MS: 361.10 ($\text{C}_{22}\text{H}_{17}\text{O}_5^+$, $[\text{M}+\text{H}]^+$). Anal. Calcd for $\text{C}_{22}\text{H}_{16}\text{O}_5$ (%): C, 73.33; H, 4.48. Found (%): C, 73.52; H, 4.45.

2.2.2. Compound 1. Compound **L** (0.360 g, 1 mmol) was added to an ethanol/acetone (1/1) solution (5 mL) of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.237 g, 1 mmol) and stirred at room temperature for 15 min to give a clear solution. After standing for 10–15 days, the precipitates were separated by filtration, washed with ethanol thrice, and dried. Single crystals of **1** were obtained by slowly evaporating the methanol/ethyl ether solution. Yield: 75%, m.p.: 250–252°C. UV (λ nm): 373.0; 255.5. IR (cm^{-1} , KBr): 3400.5 (m), 3125.2 (m), 1735.4 (s), 1595.9 (s), 1558.7 (s), 1507.8 (s), 1483.2 (s), 1433.5 (s), 1292.6 (m), 1240.4 (s), 1205.3 (s); 1138.8 (s), 1062.0 (s), 1023.5 (m), 768.6 (m), 706.0 (s). Anal. Calcd for $\text{C}_{48}\text{H}_{42}\text{O}_{12}\text{Co}$ (%): C, 66.28; H, 4.87. Found (%): C, 66.32; H, 4.84.

2.2.3. Compound 2. Prepared accordingly using **L** and $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$. Yield: 81%, m.p.: 227–229°C. UV (λ nm): 377.5; 255.0. IR (cm^{-1} , KBr): 3385.3 (m), 3138.2 (m), 1733.6 (s), 1595.6 (s), 1559.9 (s), 1507.1 (s), 1483.4 (s), 1430.9 (m), 1290.8 (s), 1252.7 (s), 1205.3 (s); 1139.6 (s), 1062.3 (s), 1023.0 (m), 769.2 (m), 705.6 (s). Anal. Calcd for $\text{C}_{48}\text{H}_{42}\text{O}_{12}\text{Mn}$ (%): C, 65.59; H, 4.89. Found (%): C, 65.43; H, 4.82.

2.3. Crystal structure determinations and refinements

The crystallographic data for **1** and **2** were collected on a Bruker Smart 1000 CCD area detector diffractometer equipped with Mo-K α ($\lambda=0.71073$ Å) radiation using ω -scan mode. Empirical absorption correction was applied to the data. Unit cell dimensions were obtained with least-squares refinements, and all structures were solved by direct methods with SHELXL-97. All non-hydrogen atoms were located from the trial structure and then refined anisotropically. All hydrogens were generated in idealized positions. All calculations were performed with SHELXL-97 programs [14]. Other relevant parameters of the crystal structure are listed in table 1.

2.4. Antimicrobial activity

The antibacterial activity of **1** and **2** was tested against *B. subtilis*, *S. aureus*, *S. faecalis*, *P. aeruginosa*, *E. coli*, and *E. cloacae* using MTT medium. The MICs of the test complexes were determined by a colorimetric method using the dye MTT [15]. A stock solution of the synthesized complex ($50 \mu\text{g mL}^{-1}$) in DMSO was prepared and graded quantities of the test complexes were incorporated in a specified quantity of sterilized

Table 1. Crystallographic and experimental data for **1** and **2**.

	1	2
Empirical formula	C ₄₈ H ₄₂ O ₁₂ Co	C ₄₈ H ₄₂ O ₁₂ Mn
Crystal system	Triclinic	Triclinic
Formula weight	869.75	865.76
Space group	<i>P</i> $\bar{1}$	<i>P</i> $\bar{1}$
Unit cell dimensions (Å, °)		
<i>a</i>	7.2070(10)	7.1380(10)
<i>b</i>	9.4301(13)	9.377(4)
<i>c</i>	16.5509(18)	16.578(2)
α	106.360(2)	74.1510(10)
β	95.4320(10)	84.2010(10)
γ	90.9180(10)	89.363(2)
Volume (Å ³), <i>Z</i>	1073.4(2), 1	1061.8(5), 1
Calculated density (g cm ⁻³)	1.346	1.354
Absorption coefficient (mm ⁻¹)	0.464	0.376
<i>F</i> (000)	453	451
Data/restraints/parameters	3728/0/296	3698/0/287
θ range for data collection (°)	2.25–25.02	2.26–25.021
<i>R</i> _{int}	0.0393	0.0293
Final <i>R</i> indices [<i>I</i> > 2σ(<i>I</i>)]	<i>R</i> ₁ = 0.0646, <i>wR</i> ₂ = 0.1203	<i>R</i> ₁ = 0.0579, <i>wR</i> ₂ = 0.1329
Largest difference peak and hole (e Å ⁻³)	0.326 and -0.396	0.275 and -0.226
Reflections collected/unique	5624/3728	5599/3698

$$R_1 = \sum ||F_o| - |F_c|| / \sum |F_o|, \quad {}^b wR_2 = [\sum [w(F_o^2 - F_c^2)^2] / \sum [w(F_o^2)^2]]^{1/2}.$$

Table 2. MICs of the compounds.

Compound	MICs (μg mL ⁻¹)					
	Gram positive			Gram negative		
	<i>B. subtilis</i>	<i>S. aureus</i>	<i>S. faecalis</i>	<i>S. faecalis</i>	<i>E. coli</i>	<i>E. cloacae</i>
1	6.25	12.5	6.25	12.5	12.5	6.25
2	12.5	6.25	6.25	12.5	12.5	12.5
L	25	12.5	12.5	25	25	25
PenicillinG	1.562	1.562	1.562	6.25	6.25	3.125
KanamycinB	0.39	1.562	3.125	3.125	3.125	1.562

liquid medium (MH medium for antibacterial activity). A specified quantity of the medium containing the complex was poured into microtitration plates. Suspension of the microorganism was prepared to contain approximately 10⁵ cfu mL⁻¹ and applied to microtitration plates with serially diluted complexes in DMSO to be tested and incubated at 37°C for 24 h. After the minimum inhibitory concentrations (MICs) were visually determined on each of the microtitration plates, 50 μL of PBS (Phosphate Buffered Saline 0.01 mol L⁻¹, pH 7.4: Na₂HPO₄ · 12H₂O 2.9 g, KH₂PO₄ 0.2 g, NaCl 8.0 g, KCl 0.2 g, distilled water 1000 mL) containing 2 mg mL⁻¹ of MTT was added to each well. Incubation was continued at room temperature for 4–5 h. The content of each well was removed, and 100 μL of isopropanol containing 5% 1 mol L⁻¹ HCl was added to extract the dye. After 12 h of incubation at room temperature, the optical density (OD) was measured with a microplate reader at 570 nm. The observed MICs are presented in table 2.




Figure 1. Molecular structure of **1** (all hydrogens and carbons of EtOH have been omitted for clarity).




Figure 2. Molecular structure of **2** (all hydrogens and carbons of EtOH have been omitted for clarity).


3. Results and discussion

Complexes of formula $[L_2M(CH_3CH_2OH)_2]$ ($M=Co$ and Mn) were prepared as described in section 2, in moderate yields (75–85%). IR spectra of all three complexes show four bands at $1480\text{--}1600\text{ cm}^{-1}$, characteristic of the mixed modes of vibrations arising from normal coordinates having contributions from $\nu_{(C=O)}$ and $\nu_{(C=C)}$ of β -diketone groups [16]. The UV spectra of the complexes display intense absorptions at $255.0\text{--}258.5\text{ nm}$ ($\pi \rightarrow \pi^*$) and $368.0\text{--}377.5\text{ nm}$ ($n \rightarrow \pi^*$).

The structures of **1** and **2** were confirmed by single-crystal X-ray diffraction and are shown in figures 1 and 2, respectively; bond distances and angles are provided in table 3. Both complexes are electronically neutral mononuclear compounds. The central metals (Co and Mn), on an inversion center, are in pseudo octahedral coordination geometry

Table 3. Selected bond lengths (Å) and angles (°) of **1** and **2**.

1			
Co–O1	2.014(3)	Co–O2	2.033(2)
Co–O1A	2.014(3)	Co–O2A	2.033(2)
Co–O6	2.184(3)	Co–O6A	2.184(3)
O1–Co–O2	88.3(1)	O1–Co–O6A	85.5(1)
O1–Co–O2A	91.7(1)	O1–Co–O1A	180.0(0)
O1–Co–O6	94.5(1)		
2			
Mn–O1	2.084(2)	Mn–O2	2.118(2)
Mn–O1A	2.084(2)	Mn–O2A	2.118(2)
Mn–O6	2.245(3)	Mn–O6A	2.245(3)
O1–Mn–O2	84.4(9)	O1–Mn–O6	85.1(9)
O1–Mn–O6A	94.8(9)		

Figure 3. 1-D chains of **1** formed by intermolecular H bonds (O–H···O).

with two ethanols occupying both axial positions and oxygen donors from two β -diketone fragments binding in equatorial positions; each bis- β -diketonate is essentially planar [17]. The general M–O (M = Co, Mn) bond lengths are in the range 2.014(3)–2.245(3) Å, unexceptional and similar to the corresponding bonds in other cobalt and manganese diketonate complexes [18–20]. As shown in figures 3 and 4,




Figure 4. 1-D chains of **2** formed by intermolecular H bonds (O–H \cdots O).

intermolecular H-bonds (O–H \cdots O) formed between adjacent molecules lead to 1D chain structures for **1** and **2**, respectively. The H \cdots O distances are 2.018(3) and 1.967(2) Å and the OH \cdots O angles are 167.9(2) $^\circ$ and 163.7(2) $^\circ$ for **1** and **2**, respectively.

From MIC values (table 2), the complexes were more toxic towards Gram positive strains than Gram negative strains when compared to the positive controls penicillin and kanamycin, respectively. The reason may be the difference in the structures of the cell walls [21]. The walls of the Gram negative cells are more complex than those of Gram positive cells. Lipopolysaccharides form an outer lipid membrane and contribute to the complex antigenic specificity of Gram negative cells. Anti-microbial activity of complexes is due to either killing the microbes or inhibiting their multiplication by blocking their active sites [22]. Since the molecular structures are quite similar, the antibacterial activities of **1** and **2** are quite similar.

Acknowledgments

The work was co-financed by grants (Projects 30772627) from the National Natural Science Foundation of China and (Projects 09KJD150005) from Province Natural Science Foundation of Jiangsu, China.

References

- [1] L. Gong, R. Leung-Toung, T.T. Tidwell. *J. Org. Chem.*, **55**, 3634 (1990).
- [2] M. Sakkarapalayam, S.A. Indrapal. *J. Org. Chem.*, **71**, 349 (2006).

- [3] A.R. Katritzky, Z.-Q. Wang, M.-Y. Wang, C.R. Wilkerson, C.D. Hall, N.G. Akhmedov. *J. Org. Chem.*, **69**, 6617 (2004).
- [4] G. Blaskó, L. Xun, G.A. Cordell. *J. Nat. Prod.*, **51**, 60 (1988).
- [5] K. Singletary, C. MacDonald, M. Iovinelli, C. Fisher, M. Wallig. *Carcinogenesis*, **19**, 1039 (1998).
- [6] R. Kumar, Y.C. Joshi. *J. Serb. Chem. Soc.*, **73**, 937 (2008).
- [7] X. Zhang, Z.-C. Li, N. Xu, K.-B. Li, S. Lin, F.-Z. Lu, F.-S. Du, F.-M. Li. *Tetrahedron Lett.*, **47**, 2623 (2006).
- [8] D. Teixeira, T. Lalot, M. Brigodiot, E. Maréchal. *Macromolecules*, **32**, 70 (1999).
- [9] A. Joshi, S. Verma, R.B. Gaur, R.R. Sharma. *Bioinorg. Chem. Appl.*, **3**, 201 (2005).
- [10] C.L. Modenbach, F.R. Fronczek, E.W. Berg, T.C. Taylor. *Inorg. Chem.*, **22**, 4083 (1983).
- [11] A.D. Garnovskii, B.I. Kharisov, L.M. Blanco, D.A. Garnovskii, A.S. Burlov, I.S. Vasilchenko, G.I. Bondarenko. *J. Coord. Chem.*, **46**, 365 (1999).
- [12] L. Li, P.-J. Cai, D. Xu, Q.-X. Guo, S. Xue. *J. Org. Chem.*, **72**, 8131 (2007).
- [13] L.-J. Tang, S.-F. Zhang, J.-Z. Yang, W.-T. Gao, J. Cui. *Org. Prep. Proced. Int.*, **36**, 453 (2004).
- [14] G.M. Sheldrick. *SHELXTL V5.1 Software Reference Manual*, Bruker AXS, Inc., Madison, WI, USA (1997).
- [15] J. Meletiadis, J.F. Meis, J.W. Mouton, J.P. Donnelly, P.E. Verweij. *J. Clin. Microbiol.*, **38**, 2949 (2000).
- [16] K. Nakamoto. *Infrared Spectra of Inorganic and Coordination Compounds*, Wiley, New York (1964).
- [17] K. Dong, J. Sun, B.-F. Ruan, H.-B. Gong. *Acta Cryst.*, **E66**, m1290 (2010).
- [18] J.-M. Li, J.-Z. Li, H.-Q. Zhang, L.-Y. Xu. *Inorg. Chem. Commun.*, **13**, 573 (2010).
- [19] J.K. Clegg, D.J. Bray, K. Gloe, K. Gloe, M.J. Hayter, K.A. Jolliffe, G.A. Lawrance, G.V. Meehan, J.C. McMurtrie, L.F. Lindoy, M. Wenzel. *Dalton Trans.*, 1719 (2007).
- [20] G. Aromi, P. Gamez, J. Krzystek, H. Kooijman, A.L. Spek, E.J. MacLean, S.J. Teat, H. Nowell. *Inorg. Chem.*, **46**, 2519 (2007).
- [21] R.V. Singh, P. Chaudhary, S. Chauhan, M. Swami. *Spectrochim. Acta, Part A*, **72**, 260 (2009).
- [22] M. Nath, S. Pokharia, X. Song, G. Eng, M. Gielen, M. Kemmer, M. Biesemans, R. Willem, D. De Vos. *Appl. Organomet. Chem.*, **17**, 305 (2003).