


This article was downloaded by: [Renmin University of China]

On: 13 October 2013, At: 10:26

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Coordination Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gcoo20>

E/Z isomerism in monoalkylated derivatives of $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ containing 2,4-dinitrophenylhydrazone substituents

Oguejiofo T. Ujam^a, Sarah M. Devoy^a, William Henderson^a, Alistair L. Wilkins^a & Brian K. Nicholson^a

^a Department of Chemistry, University of Waikato, Private Bag 3105, Hamilton, New Zealand

Published online: 09 Aug 2011.

To cite this article: Oguejiofo T. Ujam, Sarah M. Devoy, William Henderson, Alistair L. Wilkins & Brian K. Nicholson (2011) E/Z isomerism in monoalkylated derivatives of $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ containing 2,4-dinitrophenylhydrazone substituents, *Journal of Coordination Chemistry*, 64:16, 2782-2790, DOI: [10.1080/00958972.2011.606573](https://doi.org/10.1080/00958972.2011.606573)

To link to this article: <http://dx.doi.org/10.1080/00958972.2011.606573>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

***E/Z* isomerism in monoalkylated derivatives of [Pt₂(μ-S)₂(PPh₃)₄] containing 2,4-dinitrophenylhydrazone substituents**

OGUEJIOFO T. UJAM, SARAH M. DEVOY, WILLIAM HENDERSON*,
ALISTAIR L. WILKINS and BRIAN K. NICHOLSON

Department of Chemistry, University of Waikato, Private Bag 3105,
Hamilton, New Zealand

(Received 11 May 2011; in final form 6 July 2011)

Alkylation of [Pt₂(μ-S)₂(PPh₃)₄] with 2,4-dinitrophenylhydrazone-functionalized alkylating agents XC₆H₄C(=NNHC₆H₃(NO₂)₂)CH₂Br (X = H, Ph) gives monoalkylated cations [Pt₂(μ-S){μ-SCH₂C(=NNHC₆H₃(NO₂)₂)C₆H₄X}(PPh₃)₄]⁺. An X-ray diffraction study on [Pt₂(μ-S){μ-SCH₂C(=NNHC₆H₃(NO₂)₂)Ph}(PPh₃)₄]BPh₄ shows the crystal to be the *Z* isomer, with the phenyl ring and NHC₆H₃(NO₂)₂ groups mutually *trans*. ¹H- and ³¹P{¹H} NMR spectroscopic methods indicate a mixture of *Z* (major) and *E* (minor) isomers in solution, which slowly convert mainly to the *E* isomer. Reaction of [Pt₂(μ-S)₂(PPh₃)₄] with the dinitrophenylhydrazone of chloroacetone [ClCH₂C(=NNH(C₆H₃(NO₂)₂)Me)] and NaBPh₄ gives [Pt₂(μ-S){μ-SCH₂C(=NNHC₆H₃(NO₂)₂)Me}(PPh₃)₄]BPh₄, which exists as a single (*E*) isomer.

Keywords: Platinum complexes; Alkylation reactions; Electrospray ionization mass spectrometry; X-ray crystal structure; Dinitrophenylhydrazones

1. Introduction

Alkylation and arylation reactions of the platinum(II) sulfide complex [Pt₂(μ-S)₂(PPh₃)₄] (1) [1, 2] are a versatile means of generating thiolate ligands on dinuclear platinum cores [3, 4]. Using this methodology, a wide range of functionality can be introduced. Recently, we reported the synthesis and characterization of a range of monoalkylated complexes [Pt₂(μ-S)(μ-SR)(PPh₃)₄]⁺ from alkyl halides RX, where R contained a wide variety of C=O and C=N containing moieties, such as semicarbazones and oximes [5]. The alkylation chemistry has also been further extended to an investigation of the factors that promote dialkylation of the {Pt₂S₂} core using activated alkylating agents of the type RC(O)CH₂X (X = Cl, Br) [6].

In this contribution, we report a detailed study of the monoalkylation chemistry of some analogous alkylating agents that contain the 2,4-dinitrophenylhydrazone group. The formation of 2,4-dinitrophenylhydrazone derivatives of aldehydes and ketones is a

*Corresponding author. Email: w.henderson@waikato.ac.nz

classical derivatization strategy for their qualitative and quantitative determination, despite the fact that the outcome is usually quantitatively complicated by the formation of *E* and *Z* isomers [7, 8]. We previously reported a scoping study of alkylation chemistry of [Pt₂(μ-S)₂(PPh₃)₄] monitored using the technique of electrospray ionization mass spectrometry (ESI-MS) [9], which included a very preliminary study of some of the dinitrophenylhydrazones investigated herein [10]. However, these complexes were not isolated and characterized, and their initial identification was based solely on MS evidence; this article now reports the detailed synthesis and characterization of these new dinitrophenylhydrazone derivatives together with an investigation of isomerization.

2. Results and discussion

[Pt₂(μ-S)₂(PPh₃)₄] (**1**) reacts smoothly in methanol with the 2,4-dinitrophenylhydrazone derivative of BrCH₂C(O)Ph [i.e. BrCH₂C{=NNHC₆H₃(NO₂)₂}Ph] in an analogous manner to reactions with other related monofunctional alkylating agents [5]. The progress of the reaction was indicated by the dissolution of the sparingly soluble **1** and replacement of the [1 + H]⁺ ion by the molecular cation of the product in the ESI mass spectrum. From the resulting solution, the product was isolated as its tetraphenylborate salt [Pt₂(μ-S){μ-SCH₂C{=NNHC₆H₃(NO₂)₂}Ph}(PPh₃)₄]BPh₄ (**2a**·BPh₄), by the addition of excess NaBPh₄ to the reaction mixture. Similarly, the reaction of [Pt₂(μ-S)₂(PPh₃)₄] with the dinitrophenylhydrazone of BrCH₂C(O)C₆H₄Ph-4 gave [Pt₂(μ-S){μ-SCH₂C{=NNHC₆H₃(NO₂)₂}C₆H₄Ph}(PPh₃)₄]PF₆ (**2b**·PF₆), by precipitation with excess NH₄PF₆.

Crystals of **2a**·BPh₄ suitable for a single-crystal X-ray diffraction study were obtained by recrystallization from dichloromethane–diethyl ether, and a structure determination was carried out to fully characterize the compound. The molecular structure of the cation is given in figure 1 together with the atom numbering scheme, while selected bond lengths and angles are summarized in table 1. The structure determination confirms the product as a monoalkylated derivative of [Pt₂(μ-S)₂(PPh₃)₄] with structural features very similar to those of the corresponding phenacyl compound [Pt₂(μ-S){μ-SCH₂C(O)Ph}(PPh₃)₄]⁺ [5]. The complex has the typically folded butterfly {Pt₂S₂} core showing a dihedral angle between the two PtS₂ coordination planes of 129.22°, which compares well with 133.8° in [Pt₂(μ-S){μ-SCH₂C(O)Ph}(PPh₃)₂]⁺. As normal in complexes of this type, the Pt(1)–P(4) and Pt(2)–P(2) bonds are longer than Pt(1)–P(3) and Pt(2)–P(1), due to P(2) and P(4) being *trans* to the higher *trans*-influence [11] sulfide. Pt–S(sulfide) bond distances are notably shorter [Pt(1)–S(1) 2.3319(7), Pt(2)–S(1) 2.3379(7) Å] compared to Pt–S(thiolate) bond distances [Pt(1)–S(2) 2.3843(7), Pt(2)–S(2) 2.3541(7) Å].

The alkyl substituent on S(2) adopts the typical axial position, analogous to [Pt₂(μ-S){μ-SCH₂C(O)Ph}(PPh₃)₂]⁺ [5]. The hydrazone has the NHC₆H₃(NO₂)₂ group and phenyl substituents mutually *trans*, formally making it the *Z* isomer on the basis of the Cahn–Ingold–Prelog priority rules [12]. The hydrazone N–N bond is positioned directly above Pt(2). The phenyl ring on C(2) is directed away from the {Pt₂S₂} core, while the NNHC₆H₃(NO₂)₂ group is positioned to one side of the core.




Figure 1. ORTEP diagram of the cation of $[\text{Pt}_2(\mu\text{-S})\{\mu\text{-SCH}_2\text{C}(\text{=NNHC}_6\text{H}_3(\text{NO}_2)_2)\text{Ph}\}(\text{PPh}_3)_4]\text{BPh}_4$ (**2a**· BPh_4). Thermal ellipsoids are at the 50% probability level. Phenyl rings of the triphenylphosphine ligands and the BPh_4^- counterions have been omitted for clarity.

Table 1. Selected bond lengths (Å) and angles (°) for $[\text{Pt}_2(\mu\text{-S})\{\mu\text{-SCH}_2\text{C}(\text{=NNHC}_6\text{H}_3(\text{NO}_2)_2)\text{Ph}\}(\text{PPh}_3)_4]\text{BPh}_4$ (**2a**· BPh_4); estimated standard deviations are in parentheses.

Pt(1)–S(1)	2.3319(7)	Pt(2)–S(1)	2.3379(7)
Pt(1)–S(2)	2.3843(7)	Pt(2)–S(2)	2.3541(7)
Pt(1)–P(3)	2.2657(7)	Pt(2)–P(1)	2.2880(7)
Pt(1)–P(4)	2.3060(7)	Pt(2)–P(2)	2.2940(7)
S(2)–C(1)	1.854(3)	C(1)–C(2)	1.502(4)
C(2)–N(1)	1.301(4)	N(1)–N(2)	1.379(4)
Pt(1)⋯Pt(2)	3.2162(2)	S(1)⋯S(2)	3.075(1)
S(1)–Pt(1)–S(2)	81.37(2)	S(1)–Pt(2)–S(2)	81.89(2)
P(3)–Pt(1)–S(1)	91.12(3)	P(1)–Pt(2)–S(1)	84.66(3)
P(3)–Pt(1)–P(4)	97.46(3)	P(2)–Pt(2)–S(2)	94.87(2)
P(4)–Pt(1)–S(2)	90.07(2)	Pt(1)–S(1)–Pt(2)	87.06(2)
P(1)–Pt(2)–P(2)	98.62(3)	Pt(1)–S(2)–Pt(2)	85.49(2)
C(1)–S(2)–Pt(1)	98.56(10)	C(1)–S(2)–Pt(2)	107.37(10)
C(2)–C(1)–S(2)	114.7(2)	N(1)–C(2)–C(1)	124.4(3)
N(1)–C(2)–C(11)	116.0(3)	C(2)–N(1)–N(2)	117.8(3)

The hydrazone NH interacts with an oxygen of the *ortho* nitro group forming a six-membered hydrogen-bonded ring. Such hydrogen-bond interactions in dinitrophenylhydrazones are well-known [13]. The $\text{H}\cdots\text{O}$ distance of 2.095(3) Å and the $\text{N}\cdots\text{O}$ separation of 2.672(4) Å fall well within the suggested ranges for $\text{N}\cdots\text{H}\cdots\text{O}$ and $\text{N}\cdots\text{O}$

distances of 1.60–2.40 Å [14] and 2.57–3.22 Å [15], respectively. The N–H bond also appears to be directed toward the free sulfide, S(1), raising the possibility of an N–H···S hydrogen bond [16, 17]. The sulfide centers of [Pt₂(μ-S)₂(PPh₃)₄] have been recently shown to form O–H···S hydrogen-bonded adducts with alcohols [18], while [(Ph₃P)₄Pt₂(μ-S){μ-SCH₂C(O)NHC₆H₄NHC(O)CH₂S}(μ-S)Pt₂(PPh₃)₄]²⁺ (containing an *ortho*-phenylenediamide group) was also recently proposed to have an N–H···S interaction based on an N···S separation of 3.297 Å [19]. N···S distances in hydrogen-bonded systems are normally in the range 3.3–3.5 Å [20, 21]. However, the N···S separation in **2a**·BPh₄ is long [3.570(3) Å], and any N–H···S interaction will therefore be very weak.

The positive ion ESI mass spectra of the phenyl-substituted dinitrophenylhydrazones **2a**·BPh₄ and **2b**·PF₆ gave single ions at *m/z* 1802 and 1876, respectively, suggesting the formation of pure products. However, the ³¹P{¹H} NMR spectra of freshly prepared solutions of the compounds in CDCl₃ gave almost identical spectra that showed the presence of two different isomers, as illustrated in figure 2(a) for **2b**·PF₆. These were subsequently shown (*vide infra*) to be the *E* and *Z* geometric isomers at the C=N bond (scheme 1). For **2b**, the major *Z* isomer shows two overlapping multiplets for two inequivalent PPh₃ ligands, showing ¹J(PtP) values of 3344 Hz (for PPh₃ *trans* to thiolate) and 2602 Hz (for PPh₃ *trans* to sulfide). The initially minor *E* isomer of **2b** gave better separated multiplets for the two PPh₃ environments with similar ¹J(PtP) coupling constants to the *Z* isomer [3411 and 2639 Hz]. The ¹J(PtP) coupling constants for **2a** and **2b** are similar to those previously reported for the ketone compound [Pt₂(μ-S){μ-SCH₂C(O)Ph}(PPh₃)₂]BPh₄ (3386 and 2630 Hz) [5].

The ¹H-NMR spectrum of the initially prepared solution of **2b**·PF₆ confirmed the presence of two isomers. In order to determine that the crystallographically characterized *Z* isomer was the major isomer in the initially prepared solution, further ¹H-NMR studies were carried out. The hydrazone NH protons are easily identified in the ¹H-NMR spectrum of the initial solution for the major and minor isomers at δ 10.68 and 11.19, by comparison with other dinitrophenylhydrazones, e.g. CH₃C(=NNHC₆H₃(NO₂)₂)CH₃ (δNH 11.3) [22]. The spectrum also showed signals due to the SCH₂ protons at δ 4.40 and 3.65 for the major and minor isomers, respectively. Assignment of the NH and SCH₂ signals to the individual *E* and *Z* isomers was made on the basis of ROESY NMR data. Thus the SCH₂ resonance at δ 4.40 and the NH resonance at δ 10.68 (for the initial major isomer) exhibited mutual ROESY correlations, consistent with the close proximity of these protons in the *Z* isomer (scheme 1). However, the SCH₂ proton resonance in the initial minor isomer (at δ 3.65) showed a ROESY correlation to a nearby aromatic proton at δ 6.79 and not to the NH proton at δ 11.19. Similarly, this NH proton showed a ROESY correlation to an aromatic proton at δ 6.79 and not to the SCH₂ protons at δ 3.65. These correlations are consistent with the relationship between the NH and SCH₂ protons as given in the *E* isomer (scheme 1). Analysis of the starting alkylating agent BrCH₂C(=NNHC₆H₃(NO₂)₂)C₆H₄Ph by ¹H-NMR indicated that the major component had the same spatial arrangement as observed in the initially major *Z* isomer of **2b**.

When the solution of **2b**·PF₆ used for NMR analysis was stored at –18°C for 6 months, there was reversal of the relative amounts of the two isomers in both the ³¹P{¹H}- and ¹H-NMR spectra, with the initially minor *E* isomer now being the major species (figure 2b). After a longer period (9 months) complete conversion had still not occurred. The conversion of the initial *Z* isomer to the *E* isomer in solution is





Figure 2. $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of **2b**-PF₆ in CDCl₃: (a) freshly prepared solution; peaks due to the initially dominant *Z* isomer are indicated by stars, while those for the *E* isomer are indicated by circles; (b) after storage at -18°C for 6 months.



Scheme 1. Structures of the *Z* and *E* isomers of the cations $[\text{Pt}_2(\mu\text{-S})\{\mu\text{-SCH}_2\text{C}(\text{=NNHC}_6\text{H}_3(\text{NO}_2)_2)\text{C}_6\text{H}_4\text{X}\}(\text{PPh}_3)_4]^+$ ($\text{X}=\text{H}$ **2a** or Ph **2b**), showing the spatial relationship between the NH and SCH₂ protons.

presumably driven by steric interactions, as found in previous studies of the *E*-*Z* isomerization of dinitrophenylhydrazones derived from substituted acetophenones [23]. In the *E* isomer, the steric bulk of the 2,4-dinitrophenyl group projects further away from the {Pt₂S₂} core. The observation of this isomerization process suggests that any N-H...S hydrogen bond interaction (only possible in the *Z* isomer) must indeed be very weak.


Finally, observation of both *E*/*Z* isomers appears to be confined to aryl ketone derivatives. Thus, reaction of **1** with dinitrophenylhydrazone of chloroacetone [ClCH₂C(=NNH(C₆H₃(NO₂)₂))Me] proceeded rapidly, giving a bright yellow-orange solution from which the salt [Pt₂(μ-S){μ-SCH₂C(=NNHC₆H₃(NO₂)₂))Me}(PPh₃)₄]BPh₄, **3**·BPh₄ was isolated by the addition of NaBPh₄. The product showed a single ion in its ESI mass spectrum, at *m/z* 1740, due to the product cation. The NMR spectra of this product also showed only a single isomer, in contrast to the aromatic-substituted dinitrophenylhydrazones described above. Thus, two ³¹P resonances were observed for the two inequivalent types of PPh₃, showing coupling constants of 3342 and 2613 Hz. The ¹H-NMR spectrum showed a single NH resonance at δ 10.56, as well as single CH₂ and CH₃ resonances at δ 3.73 and 0.47, respectively. ¹H-ROESY NMR data showed enhancement of the CH₃ but not the NH proton on irradiation of the CH₂ protons, while irradiation of the NH proton produced an enhancement only in the methyl protons. This places the NH and CH₃ groups mutually *cis*, making the complex the *E* isomer (scheme 2).

In conclusion, this study has further extended the alkylation chemistry of [Pt₂(μ-S)₂(PPh₃)₄] to include alkylating agents containing 2,4-dinitrophenylhydrazone groups and demonstrated the occurrence of hydrazone *E*-*Z* isomerization in some derivatives.

3. Experimental

[Pt₂(μ-S)₂(PPh₃)₄] (**1**) was prepared by the literature procedure [18, 24]. The dinitrophenylhydrazone alkylating agents BrCH₂C(=NNHC₆H₃(NO₂)₂)C₆H₄Ph, BrCH₂C(=NNHC₆H₃(NO₂)₂)Ph, and ClCH₂C(=NNHC₆H₃(NO₂)₂)Me were prepared by the standard procedure for the conversion of ketones into their 2,4-dinitrophenylhydrazones [25], by the reaction of BrCH₂C(O)C₆H₄Ph (2-bromo-4'-phenylacetophenone, Aldrich), BrCH₂C(O)Ph (phenacyl bromide, BDH), or ClCH₂C(O)CH₃ (chloroacetone, BDH) with 2,4-dinitrophenylhydrazine (BDH) in acidified ethanol solution.

ESI mass spectra were recorded on a VG Platform II instrument. Solutions of approximate concentration *ca.* 0.1 mg mL⁻¹ were prepared by dissolving a small quantity of sample in a few drops of either dichloromethane or acetonitrile, followed by dilution to *ca.* 1.5 mL with methanol. Confirmation of species was facilitated by comparison of observed and calculated isotope distribution patterns, the latter obtained from the *Isotope* program [26]. *m/z* values are of the most abundant isotopomer in the isotope envelope of the ion. Elemental analyses were obtained by the Campbell Microanalytical Laboratory at the University of Otago, Dunedin, New Zealand. NMR spectra were recorded in CDCl₃ solution on a Bruker Avance DRX300 spectrometer using XWIN-NMR software version 3.0. IR spectra were recorded as KBr discs using a



Scheme 2. Structure of the cation $[\text{Pt}_2(\mu\text{-S})\{\mu\text{-SCH}_2\text{C}\{=\text{NNHC}_6\text{H}_3(\text{NO}_2)_2\}\text{CH}_3\}(\text{PPh}_3)_4]^+$ **3** as the *E* isomer observed by $^1\text{H-NMR}$ studies.

Digilab FTS2000 Scimitar Series FT-IR with Digilab Merlin Version 3.4 software. Melting points were recorded on a Reichert Thermovar apparatus, and are uncorrected.

3.1. Synthesis of $[\text{Pt}_2(\mu\text{-S})\{\mu\text{-SCH}_2\text{C}\{=\text{NNHC}_6\text{H}_3(\text{NO}_2)_2\}\text{Ph}\}(\text{PPh}_3)_4]\text{BPh}_4$ (**2a·BPh₄**)

$[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ (149 mg, 0.099 mmol) and $\text{BrCH}_2\text{C}\{=\text{NNHC}_6\text{H}_3(\text{NO}_2)_2\}\text{Ph}$ (76 mg, 0.201 mmol) in methanol (40 mL) were stirred for 1 h giving a cloudy yellow-orange solution. The solution was filtered and NaBPh_4 (73 mg, 0.214 mmol) added to the filtrate to give a yellow-orange precipitate. The product was filtered, washed successively with water (10 mL), methanol (10 mL), and diethyl ether (10 mL) and dried to give **2a·BPh₄** (121 mg, 58%). Found: C, 61.4; H, 4.3; N, 2.6. $\text{C}_{110}\text{H}_{91}\text{N}_4\text{BO}_4\text{P}_4\text{Pt}_2\text{S}_2$ requires C, 62.2; H, 4.3; N, 2.6%. M.p. 168–172°C. IR $\nu(\text{NH})$ 3285; $\nu(\text{C}=\text{N})$ 1614; $\nu(\text{NO}_2)$ 1512, 1335 cm^{-1} . ESI-MS (cone voltage 20 V) m/z 1802 ($[\text{M}]^+$, 100%). $^{31}\text{P}\{^1\text{H}\}$ NMR: *Z* isomer, δ 24.4 [m, $^1\text{J}(\text{PtP})$ 3341 and 2602]; *E* isomer, δ 23.3 [m, $^1\text{J}(\text{PtP})$ 2648] and 22.6 [m, $^1\text{J}(\text{PtP})$ 3424]. Selected $^1\text{H-NMR}$ data: *Z* isomer, δ 10.5 (s, NH), 4.45 (s, br, SCH_2); *E* isomer, δ 11.1 (s, NH), 3.60 (s, br, SCH_2).

3.2. Synthesis of $[\text{Pt}_2(\mu\text{-S})\{\mu\text{-SCH}_2\text{C}\{=\text{NNHC}_6\text{H}_3(\text{NO}_2)_2\}\text{C}_6\text{H}_4\text{Ph}\}(\text{PPh}_3)_4]\text{PF}_6$ (**2b·PF₆**)

$[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ (100 mg, 0.067 mmol) and $\text{BrCH}_2\text{C}\{=\text{NNHC}_6\text{H}_3(\text{NO}_2)_2\}\text{C}_6\text{H}_4\text{Ph}$ (36.4 mg, 0.080 mmol) in methanol (25 mL) were stirred for 1 h giving a clear, deep yellow-orange solution. After filtration to remove a small amount of insoluble matter, NH_4PF_6 (80 mg, 0.491 mmol) was added to the filtrate. Water (10 mL) was added to assist precipitation, the product was filtered, washed with diethyl ether (40 mL), and dried to give **2b·PF₆** (91.5 mg, 68%). The product was recrystallized by vapor diffusion of diethyl ether into a dichloromethane solution of the complex. Found: C, 54.2; H, 3.7; N, 2.7. $\text{C}_{92}\text{H}_{75}\text{N}_4\text{F}_6\text{O}_4\text{P}_5\text{Pt}_2\text{S}_2$ requires C, 54.6; H, 3.7; N, 2.8%. IR $\nu(\text{C}=\text{N})$ 1613;

$\nu(\text{NO}_2)$ 1501, 1333 cm⁻¹. ESI-MS (cone voltage 20 V) m/z 1876 ([M]⁺, 100%). ³¹P{¹H} NMR: *Z* isomer, δ 24.5 [m, ¹J(PtP) 3344 and 2602]; *E* isomer, δ 23.1 [m, ¹J(PtP) 2639] and 22.5 [m, ¹J(PtP) 3411]. Selected ¹H-NMR data: *Z* isomer, δ 10.68 (s, NH), 4.40 (s, br, SCH₂); *E* isomer, δ 11.19 (s, NH), 3.65 (s, br, SCH₂).

3.3. Synthesis of [Pt₂(μ-S){μ-SCH₂C{=NNHC₆H₃(NO₂)₂Me}(PPh₃)₄]BPh₄ (3·BPh₄)

[Pt₂(μ-S)₂(PPh₃)₄] (150 mg, 0.100 mmol) and ClCH₂C{=NNH(C₆H₃(NO₂)₂)Me} (54 mg, 0.198 mmol) in methanol (40 mL) were stirred at room temperature for 30 min giving a clear, bright yellow-orange solution. NaBPh₄ (70 mg, 0.205 mmol) was added to give a bright yellow precipitate. The product was filtered, washed successively with water (10 mL), methanol (10 mL), and petroleum spirits (10 mL), and dried to give 3·BPh₄ as a bright yellow solid (139 mg, 68%). Found: C, 61.1; H, 4.4; N, 2.8. C₁₀₅H₈₉N₄BO₄P₄Pt₂S₂ requires C, 61.2; H, 4.4; N, 2.7%. M.p. decomposition >152°C. IR $\nu(\text{NH})$ 3287; $\nu(\text{C}=\text{N})$ 1616; $\nu(\text{NO}_2)$ 1509, 1335 cm⁻¹. ESI-MS (cone voltage 20 V) m/z 1740 ([M]⁺, 100%). ³¹P{¹H} NMR, δ 24.9 [m, ¹J(PtP) 3342] and 23.6 [m, ¹J(PtP) 2613]. Selected ¹H-NMR data, δ 10.56 (s, NH), 3.73 [s, br, 2H, SCH₂, ³J(PtH) 33] and 0.47 (s, 3H, CH₃).

3.4. X-ray crystal structure determination of [Pt₂(μ-S){μ-SCH₂C{=NNHC₆H₃(NO₂)₂Ph}(PPh₃)₄]BPh₄ (2a·BPh₄)

Crystals (orange blocks) were grown by vapor diffusion of diethyl ether into a dichloromethane solution of the complex at room temperature. Intensity data and unit cell dimensions were obtained on a Bruker SMART CCD diffractometer at the University of Auckland.

The structure was solved for the positions of the Pt atoms using the Patterson methods option of SHELXS-97 [27]. The structures of the cation and anion were developed routinely. Later difference maps showed significant electron density associated with disordered solvents in the lattice, possibly two CH₂Cl₂ and one ether molecules. These could not be sensibly modeled, so they were eliminated using the SQUEEZE procedure of PLATON [28] and were not included in the refinement.

Full-matrix least-squares refinement (SHELXL-97) [29] was based on F_o^2 with all non-hydrogen atoms anisotropic and hydrogen in calculated positions converged with $R_1=0.0247$ ($I \geq 2\sigma(I)$), $wR_2=0.0596$ (all data), goodness-of-fit = 1.068. A final difference map showed no feature greater than 1.039 and -0.932 e Å⁻³. All calculations were carried out by the SHELX-97 suite of programs.

3.5. Crystal data for C₁₁₀H₉₁BN₄O₄P₄Pt₂S₂

$M_r=2121.86$; triclinic; space group *P*-1; $a=14.4408(2)$ Å; $b=16.5455(3)$ Å; $c=22.9006(2)$ Å; $\alpha=89.355(1)^\circ$, $\beta=72.784(1)^\circ$, $\gamma=78.209(1)^\circ$; $V=5109.12(12)$ Å³; $Z=2$, $T=83(2)$ K, $\lambda(\text{Mo-K}\alpha)=0.71073$ Å; $\mu(\text{Mo-K}\alpha)=2.891$ mm⁻¹; $d_{\text{calcd}}=1.379$ g cm⁻³; 48,989 reflections collected; 20,704 unique ($R_{\text{int}}=0.0224$).

Supplementary material

CCDC no. 824691 contains the supplementary crystallographic data for this article. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre, via www.ccdc.cam.ac.uk/data_request/cif.

Acknowledgments

We thank the University of Waikato for financial support of this study and Dr Tania Groutso (University of Auckland) for collection of the X-ray data set. We also thank Pat Gread for assisting with mass spectrometry instrumentation.

References

- [1] S.-W.A. Fong, T.S.A. Hor. *J. Chem. Soc., Dalton Trans.*, 639 (1999).
- [2] J. Chatt, D.M.P. Mingos. *J. Chem. Soc. A*, 1243 (1970).
- [3] B.J. Deadman, W. Henderson, B.K. Nicholson, L.E. Petchell, S.L. Rose, T.S.A. Hor. *Inorg. Chim. Acta*, **363**, 637 (2010).
- [4] S.H. Chong, L.L. Koh, W. Henderson, T.S.A. Hor. *Chem. Asian J.*, 264 (2006).
- [5] O.T. Ujam, S.M. Devoy, W. Henderson, B.K. Nicholson, T.S.A. Hor. *Inorg. Chim. Acta*, **363**, 3558 (2010).
- [6] O.T. Ujam, W. Henderson, B.K. Nicholson, T.S.A. Hor. *Inorg. Chim. Acta* (in press). DOI: 10.1016/j.ica.2011.06.023.
- [7] S. Uchiyama, M. Ando, S. Aoyagi. *J. Chromatogr. A*, **996**, 95 (2003).
- [8] F. Ramirez, A.F. Kirby. *J. Am. Chem. Soc.*, **76**, 1037 (1954).
- [9] W. Henderson, J.S. McIndoe. *Mass Spectrometry of Inorganic, Coordination and Organometallic Compounds*, Wiley & Sons, Chichester (2005).
- [10] W. Henderson, S.H. Chong, T.S.A. Hor. *Inorg. Chim. Acta*, **359**, 3440 (2006).
- [11] T.G. Appleton, H.C. Clark, L.E. Manzer. *Coord. Chem. Rev.*, **10**, 335 (1973).
- [12] J.G. Smith. *Organic Chemistry*, Section 5.6, McGraw-Hill, New York (2006).
- [13] F. Kaberia, B. Vickery, G.R. Willey, M.G.B. Drew. *J. Chem. Soc. Perkin II*, 1622 (1980).
- [14] G.A. Jeffrey, W. Saenger. *Hydrogen Bonding in Biological Structures*, Springer-Verlag, Berlin (1991).
- [15] G.H. Stout, L.H. Jensen. *X-ray Structure Determination*, McMillan, London (1968).
- [16] K. Baba, T. Okamura, H. Yamamoto, T. Yamamoto, N. Ueyama. *Inorg. Chem.*, **47**, 2837 (2008).
- [17] Z. Yu, M. Wang, P. Li, W. Dong, F. Wang, L. Sun. *Dalton Trans.*, 2400 (2008).
- [18] R. Ugo, G. La Monica, S. Cenini, A. Segre, F. Conti. *J. Chem. Soc.*, 522 (1971).
- [19] O.T. Ujam, W. Henderson, B.K. Nicholson, C.M. Fitchett. *Inorg. Chim. Acta* (in press). DOI: 10.1016/j.ica.2011.05.010.
- [20] M.A. Walters, C.L. Roche, A.L. Rheingold, S.W. Kassel. *Inorg. Chem.*, **44**, 3777 (2005).
- [21] W.P. Chung, J.C. Dewan, M.A. Walters. *Inorg. Chem.*, **30**, 4280 (1991).
- [22] K. Miyamoto, N. Tada, M. Ochiai. *J. Am. Chem. Soc.*, **129**, 2772 (2007).
- [23] J.P. Idoux, J.A. Sikorski. *J. Chem. Soc. Perkin II*, 921 (1972).
- [24] W. Henderson, S. Thwaite, B.K. Nicholson, T.S.A. Hor. *Eur. J. Inorg. Chem.*, 5119 (2008).
- [25] O.L. Brady. *J. Chem. Soc.*, 757 (1931).
- [26] L.J. Arnold. *J. Chem. Educ.*, **69**, 811 (1992).
- [27] G.M. Sheldrick. *SHELXS-97 Program for Solution of Crystal Structures*, University of Göttingen, Germany (1997).
- [28] A.L. Spek. *J. Appl. Cryst.*, **36**, 7 (2003).
- [29] G.M. Sheldrick. *SHELXL-97 Program for the Refinement of Crystal Structures*, University of Göttingen, Germany (1997); G.M. Sheldrick. *Acta Crystallogr.*, **A64**, 112 (2008).