

CONTACT RIEMANNIAN SUBMANIFOLDS

MASAFUMI OKUMURA

Introduction

In a previous paper [3] the author studied a submanifold of codimension 2, which inherits a contact Riemannian structure from the enveloping contact Riemannian manifold.

In the present paper, the author generalizes the results obtained in [3] to submanifolds of codimension greater than 2. In § 1 we recall first of all the definition of contact Riemannian manifolds and some identities which hold in such manifolds, and in § 2 we give some formulas which hold for submanifolds in a Riemannian manifold. After these preliminaries, § 3 contains some identities which hold for submanifolds in a contact Riemannian manifold. In § 4 we define the notion of contact Riemannian submanifolds in the same way as given in [3]. In § 5 we define an F -invariant submanifold and study the relations between contact Riemannian submanifolds and F -invariant submanifolds.

§ 6 is devoted to a condition for a submanifold to be a contact Riemannian manifold. In the last section, § 7, we introduce the notion of normal contact submanifolds in a normal contact manifold, and obtain a condition for a contact Riemannian manifold to be a normal contact manifold.

1. Contact Riemannian manifolds

A $(2n + 1)$ -dimensional differentiable manifold \bar{M} is said to have a *contact structure* and called a *contact manifold* if there exists a 1-form $\tilde{\eta}$, to be called the *contact form*, on \bar{M} such that

$$(1.1) \quad \tilde{\eta} \wedge (d\tilde{\eta})^n \neq 0$$

everywhere on \bar{M} , where $d\tilde{\eta}$ is the exterior derivative of $\tilde{\eta}$, and the symbol \wedge denotes the exterior multiplication.

In terms of local coordinate $\{y^i\}$ of \bar{M} the contact form $\tilde{\eta}$ is expressed as

$$(1.2) \quad \tilde{\eta} = \eta_i dy^i.$$

Since, according to (1.1), the 2-form $d\tilde{\eta}$ is of rank $2n$ everywhere on \bar{M} , we can find a unique vector field ξ^r on \bar{M} satisfying

Communicated by K. Yano, December 11, 1968.

$$(1.3) \quad \eta_{\lambda}^{\xi^{\lambda}} = 1, \quad (d\tilde{\eta})_{\lambda}^{\xi^{\lambda}} = 0.$$

It is well known that there exists a positive definite Riemannian metric $\bar{g}_{\lambda\mu}$ such that the (1, 1)-tensor F_{λ}^{ξ} , defined by

$$(1.4) \quad 2\bar{g}_{\lambda\xi}F_{\mu}^{\xi} = (d\tilde{\eta})_{\mu\lambda},$$

satisfies the conditions

$$(1.5) \quad F_{\lambda}^{\xi}F_{\mu}^{\lambda} = -\delta_{\mu}^{\xi} + \tilde{\eta}_{\mu}^{\xi},$$

$$(1.6) \quad \tilde{\eta}_{\xi}F_{\lambda}^{\xi} = 0,$$

$$(1.7) \quad \bar{g}_{\lambda\mu}\tilde{\xi}^{\mu} = \tilde{\eta}_{\lambda},$$

$$(1.8) \quad \bar{g}_{\lambda\xi}F_{\nu}^{\lambda}F_{\mu}^{\xi} = \bar{g}_{\nu\mu} - \tilde{\eta}_{\nu}\tilde{\eta}_{\mu}.$$

(S. Sasaki [4], Y. Hatakeyama [1]). The set $(F_{\lambda}^{\xi}, \tilde{\xi}^{\lambda}, \tilde{\eta}_{\lambda}, \bar{g}_{\lambda\xi})$ satisfying (1.1), (1.3), (1.5) and (1.7) is called a *contact Riemannian* (or *metric*) *structure*, and the manifold with such a structure is called a *contact Riemannian* (or *metric*) *manifold*.

If in a contact Riemannian manifold the tensor, defined by

$$(1.9) \quad N_{\mu\xi} = F_{\mu}^{\nu}(\partial_{\nu}F_{\lambda}^{\xi} - \partial_{\lambda}F_{\nu}^{\xi}) - F_{\lambda}^{\nu}(\partial_{\nu}F_{\mu}^{\xi} - \partial_{\mu}F_{\nu}^{\xi}) \\ + \partial_{\lambda}\tilde{\xi}^{\xi}\tilde{\eta}_{\mu} - \partial_{\mu}\tilde{\xi}^{\xi}\tilde{\eta}_{\lambda},$$

where $\partial_{\nu} = \partial/\partial y^{\nu}$ vanishes everywhere on \tilde{M} , then the structure is said to be *normal*, and the manifold is called a *normal contact manifold* or a *Sasakian manifold*. In a normal contact manifold we have

$$(1.10) \quad \tilde{V}_{\mu}\tilde{\eta}_{\lambda} = F_{\mu\lambda},$$

$$(1.11) \quad \tilde{V}_{\mu}F_{\lambda\xi} = \tilde{\eta}_{\lambda}\bar{g}_{\mu\xi} - \tilde{\eta}_{\xi}\bar{g}_{\mu\lambda},$$

where \tilde{V} denotes the covariant differentiation with respect to the Riemannian metric \bar{g} . Conversely, if (1.11) holds, the manifold is a normal contact manifold (Y. Hatakeyama, Y. Ogawa, and S. Tanno [2]).

2. Submanifolds in a Riemannian manifold

Let M be an m -dimensional oriented differentiable manifold and ι be an immersion of M into an $(m+k)$ -dimensional oriented Riemannian manifold \tilde{M} . In terms of local coordinates (x^1, \dots, x^m) of M and (y^1, \dots, y^{m+k}) of \tilde{M} the immersion ι is locally expressed by $y^{\kappa} = y^{\kappa}(x^1, \dots, x^m)$, $\kappa = 1, \dots, m+k$. If we put $B_i^{\kappa} = \partial_{\lambda}y^{\kappa}$, $\partial_i = \partial/\partial x^i$, then B_i^{κ} are m local vector fields in M spanning the tangent space at each point of M . A Riemannian metric g on M is

naturally induced from the Riemannian metric \bar{g} on \bar{M} by the immersion in such a way that

$$(2.1) \quad g_{ji} = \bar{g}_{\lambda\kappa} B_j^\lambda B_i^\kappa.$$

Since M and \bar{M} are both orientable, in each coordinate neighborhood U of $p \in M$, we can choose k fields of mutually orthogonal unit normal vectors N_A^ϵ ($A = 1, \dots, k$) of M at each point of U in such a way that $(N_1^\epsilon, \dots, N_k^\epsilon, B_i^\epsilon)$ is positively oriented in \bar{M} , provided that the frame $(B_i^\epsilon, i = 1, \dots, m)$ is so in M .

Let H_{Aj^i} ($A = 1, \dots, k$) be the second fundamental tensors, and L_{AB^i} the third fundamental tensors of the immersion ι . Then we have the following Gauss and Weingarten equations:

$$(2.2) \quad \nabla_j B_i^\epsilon = \sum_{A=1}^k H_{Aj^i} N_A^\epsilon,$$

$$(2.3) \quad \nabla_j N_A^\epsilon = -H_{Aj^i} B_i^\epsilon + \sum_{B=1}^k L_{AB^j} N_B^\epsilon,$$

where ∇_j is the so-called van der Waerden-Bortolotti covariant differentiation, where $\nabla_j B_i^\epsilon$ and $\nabla_j N_A^\epsilon$ are defined respectively by

$$\begin{aligned} \nabla_j B_i^\epsilon &= \partial_j B_i^\epsilon - \left\{ \begin{matrix} h \\ j \ i \end{matrix} \right\} B_h^\epsilon + \left\{ \begin{matrix} \kappa \\ \lambda \ \mu \end{matrix} \right\} B_j^\lambda B_i^\mu, \\ \nabla_j N_A^\epsilon &= \partial_j N_A^\epsilon + \left\{ \begin{matrix} \kappa \\ \lambda \ \mu \end{matrix} \right\} B_j^\mu N_A^\lambda \quad (A = 1, \dots, k), \end{aligned}$$

$\left\{ \begin{matrix} i \\ j \ k \end{matrix} \right\}$ and $\left\{ \begin{matrix} \kappa \\ \lambda \ \mu \end{matrix} \right\}$ being the Christoffel's symbols of M and \bar{M} respectively.

3. Submanifolds in a contact Riemannian manifold

Let \bar{M} be a $(2n + 1)$ -dimensional contact Riemannian manifold with a contact Riemannian structure $(F_i^\epsilon, \bar{\xi}^\epsilon, \bar{\eta}_\lambda, \bar{g}_{\lambda\kappa})$ and M a $(2m + 1)$ -dimensional submanifold in \bar{M} . The transform $F_i^\epsilon B_i^\lambda$ of the tangent vector field B_i^ϵ by F_i^ϵ can be represented as a sum of its tangential part and its normal part, that is,

$$(3.1) \quad F_i^\epsilon B_i^\lambda = f_i^h B_h^\epsilon + \sum_A f_{Ai} N_A^\epsilon.$$

In the same way, we can put

$$(3.2) \quad F_i^\epsilon N_A^\lambda = h^i B_i^\epsilon + \sum_B h_{AB} N_B^\epsilon, \quad A = 1, \dots, 2(n - m).$$

From these two equations we have

$$(3.3) \quad h_i = -f_i,$$

$$(3.4) \quad h_{AB} = -h_{BA}.$$

On the other hand, ξ^* being tangent to \tilde{M} is expressed as a linear combination of B_i^* and N_A^* . Hence we can put

$$(3.5) \quad \xi^* = u^h B_h^* + \sum_A u_A N_A^*,$$

which implies

$$(3.6) \quad u_i = \tilde{\eta}_i B_i^*,$$

$$(3.7) \quad u_A = \tilde{\eta}_A N_A^*.$$

Transforming both members of (3.1) by F_i^* and making use of (1.5), (3.1), (3.2), (3.3) and (3.5), we find

$$\begin{aligned} -B_i^* + u_i u^j B_j^* + \sum_B u_i u_B N_B^* &= (f_i^h f_h^j + \sum_A f_i^j f_j^A) B_j^* \\ &\quad + \sum_B (f_i^h f_h^B + \sum_A f_i^A h_{AB}) N_B^*, \end{aligned}$$

which implies

$$(3.8) \quad f_i^h f_h^j = -\delta_i^j + u_i u^j + \sum_A f_i^j f_j^A,$$

$$(3.9) \quad f_i^h f_h^A = u_A u_i - \sum_B f_i^B h_{BA}.$$

Transforming again both members of (3.2) by F_i^* and taking account of (1.5), (3.1), (3.2), (3.3) and (3.5), we obtain

$$\begin{aligned} u_A u^j B_j^* - N_A^* + \sum_B u_A u_B N_B^* &= -(f^i f_i^j + \sum_B h_{AB} f_j^B) B_j^* \\ &\quad + \sum_B (-f^i f_i^B + \sum_C h_{AC} h_{CB}) N_B^*, \end{aligned}$$

which implies

$$(3.10) \quad f^i f_i^j = -\sum_B h_{AB} f_j^B - u_A u^j,$$

$$(3.11) \quad f^i f_i^A = \delta_{AB} - u_A u_B + \sum_C h_{AC} h_{CB}.$$

On the other hand, conditions (1.6) and (1.3) can be rewritten respectively as

$$F_i^* \xi^i = F_i^* (u^i B_i^* + \sum_A u_A N_A^*) = 0,$$

$$\tilde{\eta}_i \xi^i = (u^i B_{i^*} + \sum_A u_A N_{A^*}) (u^j B_j^* + \sum_B u_B N_B^*) = 1,$$

from which we easily have

$$(3.12) \quad u^i f_i^h = \sum_A u_A f_A^h,$$

$$(3.13) \quad \bar{u}_A^i f_i = - \sum_B u_B h_{BA},$$

$$(3.14) \quad u^i u_i = 1 - \sum_A u_A^2.$$

Let \bar{M} be a normal contact manifold. Differentiating (3.1) covariantly and making use of (1.11), (3.2) and (3.5), we obtain

$$\begin{aligned} u_i B_j^i - g_{ji}(u^h B_h^i + \sum_A u_A N_A^i) + \sum_A H_{Aji}(-f_A^h B_h^i + \sum_B h_{AB} N_B^i) \\ = \nabla_j f_i^h B_h^i + \sum_A (f_i^h H_{Ajh} N_A^i + \nabla_j f_i^h N_A^i - f_i^h H_{Aji} B_h^i \\ + \sum_B f_i L_{BAj} N_A^i), \end{aligned}$$

which implies

$$(3.15) \quad \nabla_j f_i^h = u_i g_{jh} - u_h g_{ji} - \sum_A (f_h H_{Ahi} - f_i H_{Ajh}),$$

$$(3.16) \quad \nabla_j f_i = -u_A g_{ji} + \sum_B (H_{Bji} h_{BA} - f_i L_{BAj}) - f_i^h H_{Ajh}.$$

Differentiating (3.2) covariantly and making use of (1.11), (3.1), (3.2) and (3.5), we have

$$\begin{aligned} u_A B_j^i - H_{Aji}(f_h^i B_h^i + \sum_B f_h N_B^i) + \sum_B L_{ABj}(-f_B^i B_i^i + \sum_C h_{BC} N_C^i) \\ = -\nabla_j f_A^h B_h^i - \sum_B (f_A^i H_{Bji} - \nabla_j h_{AB}) N_B^i \\ + \sum_B h_{AB}(-H_{Bji} B_i^i + \sum_C L_{BCj} N_C^i), \end{aligned}$$

which implies

$$\begin{aligned} \nabla_j f_A^i = -u_A \delta_j^i + H_{Aji} f_h^i + \sum_B (h_{AB} H_{Bji} - L_{ABj} f_B^i), \\ (3.17) \quad \nabla_j h_{AC} = f_A^i H_{Cji} - f_i^h H_{Aji} + \sum_B (L_{ABj} h_{BC} - L_{BCj} h_{AB}). \end{aligned}$$

Differentiating (3.5) covariantly and using (1.10) which holds in a normal contact manifold, we find

$$\begin{aligned} f_j^i B_i^i + \sum_A f_j N_A^i = \nabla_j u^i B_i^i + \sum_A u^i H_{Aji} N_A^i \\ + \sum_A \{ \nabla_j u_A N_A^i + u_A(-H_{Aji} B_i^i + \sum_B L_{BAj} N_B^i) \}, \end{aligned}$$

which implies

$$(3.18) \quad \nabla_j u^i = f_j^i + \sum_A u_A H_{A j}^i,$$

$$(3.19) \quad \nabla_j u_A = f_j^i - u^i H_{A j}^i - \sum_A u_B L_{B A j}.$$

4. Contact Riemannian submanifolds

Let \tilde{M} be a $(2n + 1)$ -dimensional contact Riemannian manifold, and M a $(2m + 1)$ -dimensional orientable differentiable submanifold in \tilde{M} . We define a 1-form u on M by

$$(4.1) \quad u = u_i dx^i = \tilde{\eta}_i B_i^j dx^j,$$

in terms of the contact form $\tilde{\eta} = \tilde{\eta}_i dy^i$.

Definition 4.1. Let g_{ji} be the induced Riemannian metric of M , and u the 1-form defined by (4.1). If there exists a pair of positive constants t and c such that $\eta = tu$ and $G_{ji} = cg_{ji}$ constitute a contact Riemannian structure on M , then we call the submanifold M a *contact Riemannian submanifold* of \tilde{M} .

Since (η, G) is a contact metric structure in a contact Riemannian submanifold M , the linear mapping $\phi_j^i: T(M) \rightarrow T(M)$ and the vector field ξ^i defined respectively by

$$(4.2) \quad 2\phi_j^h G_{hi} = \partial_j \eta_i - \partial_i \eta_j, \quad \eta_i = G_{ji} \xi^j$$

satisfy the conditions

$$(4.3) \quad \eta_i \xi^i = 1,$$

$$(4.4) \quad \phi_j^i \xi^j = 0, \quad \eta_i \phi_j^i = 0,$$

$$(4.5) \quad \phi_j^h \phi_h^i = -\delta_j^i + \eta_j \xi^i.$$

Directly from Definition 4.1 we have

Proposition 4.2. Let M be a contact Riemannian submanifold in \tilde{M} , and $'M$ a contact Riemannian submanifold in M . Then $'M$ is a contact Riemannian submanifold in \tilde{M} .

Proposition 4.3. Let M be a contact Riemannian submanifold of \tilde{M} , and $'M$ a submanifold of M . If $'M$ is a contact Riemannian submanifold of \tilde{M} , then $'M$ is also a contact Riemannian submanifold of M .

Proposition 4.4. Let M be a contact Riemannian submanifold of a contact Riemannian manifold \tilde{M} . If the dimension of M is greater than the codimension of M in \tilde{M} , then we have

$$(4.6) \quad \phi_j^i = f_j^i,$$

$$(4.7) \quad u^i = \xi^i.$$

Proof. From the definitions of ξ^i, η_i, G_{ji} we have

$$(4.8) \quad \xi^j = G^{ji}\eta_i = \frac{t}{c}g^{ji}u_i = \frac{t}{c}u^j,$$

from which

$$(4.9) \quad 1 = \eta_j\xi^j = tu_j\frac{t}{c}u^j = \frac{t^2}{c}u_iu^i,$$

$$(4.10) \quad u_iu^i = c/t^2.$$

On the other hand, the two equations

$$\begin{aligned} 2f_{ji} &= 2B_j^{\lambda}B_i^{\kappa}F_{\lambda\kappa} = B_j^{\lambda}B_i^{\kappa}(\tilde{V}_{\lambda}\tilde{\eta}_{\mu} - \tilde{V}_{\mu}\tilde{\eta}_{\lambda}) = \nabla_ju_i - \nabla_iu_j, \\ 2\phi_{ji} &= \partial_j\eta_i - \partial_i\eta_j = t(\nabla_ju_i - \nabla_iu_j) \end{aligned}$$

imply $f_{ji} = (1/t)\phi_{ji}$ and hence

$$(4.11) \quad f_j^h = g^{hi}f_{ji} = \frac{c}{t}G^{hi}\phi_{ji} = \frac{c}{t}\phi_j^h.$$

Since f_j^h, ϕ_j^h satisfy (3.8) and (4.5) respectively, (4.11) together with (4.10) implies

$$(4.12) \quad -\delta_j^h + u^hu_j + \sum_A \frac{f^h}{A} \frac{f_j}{A} = \frac{c^2}{t^2} \left(-\delta_j^h + \frac{t^2}{c}u_ju^h \right).$$

We assume now that there is a point p in M , at which the $2(n-m)+1$ vectors u^i, f_A^i ($A=1, \dots, 2(n-m)$) are linearly dependent. Then we can find a vector $v^i(p)$ orthogonal to the subspace spanned by u^i and f_A^i ($A=1, \dots, 2(n-m)$), since M is of dimension greater than $2(n-m)$. Transforming this vector $v^i(p)$ by (4.12), we get $v^h(p) = (c/t)^2v^h(p)$, that is, $(c/t)^2 = 1$, which together with (4.8) and (4.11) implies the Proposition.

Next we suppose that u^i and f_A^i ($A=1, \dots, 2(n-m)$) are linearly independent at any point of M . Then (3.12), (4.4) and (4.8) imply $\sum_A u_A f_A^h = f_j^h u^j = (c/t)\phi_j^h(c/t)\xi^j = 0$. Since f_A^h 's are linearly independent, we have, in this case,

$$(4.13) \quad u_A = 0 \quad (A=1, \dots, 2(n-m)),$$

which and (3.1) give

$$(4.14) \quad u^i f_i = 0.$$

Transforming f_A^j by (4.12), we have

$$(4.15) \quad -f^h + \sum_A f_j f^j f^h = -\frac{c^2}{t^2} f^h$$

because of (4.14). Substituting (3.11) into (4.15) we get $\sum_{B,C} h_{AC} h_{CB} f^h = -(c/t)^2 f^h$ implying

$$(4.16) \quad \sum_C h_{AC} h_{CB} = -\frac{c^2}{t^2} \delta_{BA},$$

and consequently

$$(4.17) \quad \sum_{A,C} h_{AC} h_{CA} = -\frac{c^2}{t^2} \sum_A \delta_{AA} = -2(n-m) \frac{c^2}{t^2}.$$

Furthermore, from (4.11) we obtain $(c/t)^2 \phi_i^h \phi_h^j = -\delta_i^j + u_i u^j + \sum_A f_i f^j$, which yields

$$-2m \frac{c^2}{t^2} = -2m - 1 + u_i u^i + 2(n-m) + \sum_{A,C} h_{AC} h_{CA}$$

because of (3.11). On the other hand, $u_A = 0$ and (3.14) imply $u_i u^i = 1$. Thus we have, from the equation obtained above,

$$(4.18) \quad -2m \frac{c^2}{t^2} = 2(n-2m) + \sum_{A,C} h_{AC} h_{CA}.$$

Combining (4.17) and (4.18), we have $(t/c)^2 = 1$, which completes the proof.

Corollary 4.5. $G_{ji} = (u, u^r)^{-1} g_{ji}$, $\eta_i = (u, u^r)^{-1} u_i$.

5. F -invariant submanifolds

F -invariant submanifolds of a contact Riemannian manifold are recently studied in [5]. In this section we show that any F -invariant submanifold is a contact Riemannian submanifold.

Definition 5.1. Let \bar{M} be a $(2n+1)$ -dimensional contact Riemannian manifold. A $(2m+1)$ -dimensional submanifold M of \bar{M} is called an F -invariant submanifold if the tangent space of M is invariant under the action of F_t^ϵ .

Proposition 5.2. Let M be a $(2m+1)$ -dimensional submanifold of a contact Riemannian manifold \bar{M} . In order that M be an F -invariant submanifold it is necessary and sufficient that

$$(5.1) \quad \sum_C h_{AC} h_{CB} = -\delta_{AB}.$$

Proof. We first assume M to be F -invariant, and then by (3.1) show that

$$F_t^\epsilon B_i^\lambda = f_i^h B_h^\epsilon, \quad F_t^\epsilon N_A^\lambda = \sum_B h_{BA} N_B^\epsilon,$$

or equivalently $f_A^i = 0$ ($A = 1, \dots, 2(n - m)$). Consequently, we have $u_i u_A = 0$ because of (3.9). If there is a point p on M , where $u_i(p) = 0$, then (3.8) implies $f_j^i f_i^h = -\delta_j^h$, which means that the tangent space at p is even-dimensional, contradicting our assumption. Hence we have $u_A = 0$ in M . Therefore we have $\sum_C h_{AC} h_{CB} = -\delta_{AB}$ by virtue of (3.11). Next, we assume that M is a submanifold of \bar{M} satisfying the condition (5.1). Then, by means of (3.11), we have $f_A^i f_{Bi} + u_A u_B = 0$, and therefore $\sum_A f_A^i f_{Ai} + u_A^2 = 0$. Thus we get $f_A^i = 0$, $u_A = 0$, which show that M is F -invariant.

Proposition 5.3. *If M is a $(2m + 1)$ -dimensional F -invariant submanifold of \bar{M} . Then M is necessarily a contact Riemannian submanifold of \bar{M} .*

Proof. Since M is F -invariant, as seen in the proof of Proposition 5.2 we have $f_A^i = 0$, $u_A = 0$ ($A = 1, \dots, 2(n - m)$). Therefore, (3.8) and (3.14) imply $f_i^h f_h^j = -\delta_i^j + u_i u^j$, $u_i u^i = 1$. If we now put $\eta = u$, $G_{ji} = g_{ji}$ then we find

$$\begin{aligned} \nabla_j \eta_i - \nabla_i \eta_j &= \nabla_j u_i - \nabla_i u_j = \nabla_j (\tilde{\eta}_\kappa B_i^\kappa) - \nabla_i (\tilde{\eta}_\kappa B_j^\kappa) \\ &= B_i^\kappa B_j^\lambda \tilde{\nabla}_\lambda \tilde{\eta}_\kappa - B_i^\lambda B_j^\kappa \tilde{\nabla}_\lambda \tilde{\eta}_\kappa + \sum_A (H_{Aji} N_A^\kappa - H_{Aik} N_A^\kappa) \tilde{\eta}_\kappa \\ &= B_i^\kappa B_j^\lambda (\tilde{\nabla}_\lambda \tilde{\eta}_\kappa - \tilde{\nabla}_\kappa \tilde{\eta}_\lambda) = 2f_{ji}, \end{aligned}$$

which means that the (η, G) is a contact Riemannian structure on M . Thus the proof is complete.

6. Conditions for a submanifold to be a contact Riemannian submanifold

In this section we states a condition for a submanifold M in a contact Riemannian manifold \bar{M} to be a contact Riemannian submanifold. Since for this purpose we have to use Proposition 4.4 so that we always assume in this section that the dimension of M is greater than the codimension of M in \bar{M} . First we have

Proposition 6.1. *Let \bar{M} be a $(2n + 1)$ -dimensional contact Riemannian manifold. In order that a submanifold M in \bar{M} be a contact Riemannian submanifold it is necessary and sufficient that the relations*

$$(6.1) \quad u_r u^r = \text{const.} \neq 0,$$

$$(6.2) \quad f_j^i f_h^j = -\delta_h^i + (u_r u^r)^{-1} u_h u^i$$

be both valid.

Proof. Let M be a contact Riemannian submanifold of \bar{M} . Then from Proposition 4.4 it follows that $f_j^i = \phi_j^i$ and consequently

$$(6.3) \quad f_i^h f_h^j = \phi_i^h \phi_h^j = -\delta_i^j + \eta_i \xi^j = -\delta_i^j + t u_i u^j.$$

On the other hand, we have $\eta_i \xi^i = tu_i \xi^i = tu_i u^i = 1$, which implies

$$(6.4) \quad u_i u^i = \frac{1}{t} = \text{const. .}$$

Combining (6.3) and (6.4), we get (6.1) and (6.2).

Conversely, if (6.1) and (6.2) are both valid, putting

$$\eta_i = (u_r u^r)^{-1} u_i, \quad G_{ji} = (u_r u^r)^{-1} g_{ji},$$

we have

$$\begin{aligned} \eta_i \xi^i &= (u_r u^r)^{-1} u_i G^{ik} \eta_k = (u_r u^r)^{-1} u_i u^i = 1, \\ f_i^j f_h^i &= -\delta_h^j + (u_r u^r)^{-1} u_h u^j = -\delta_h^j + \eta_h \xi^j. \end{aligned}$$

Thus $(f_j^i, \eta_i, G^{ji} \eta_j, G_{ji})$ is an almost contact Riemannian structure on M . By virtue of (6.1) and (1.4) we now have

$$\begin{aligned} \nabla_j \eta_i - \nabla_i \eta_j &= (u_r u^r)^{-1} (\nabla_j u_i - \nabla_i u_j) \\ &= (u_r u^r)^{-1} (\nabla_j (B_i^l \tilde{\eta}_l) - \nabla_i (B_j^l \tilde{\eta}_l)) \\ &= (u_r u^r)^{-1} (B_i^l B_j^\mu \tilde{\nabla}_\mu \tilde{\eta}_l - B_i^\mu B_j^l \tilde{\nabla}_\mu \tilde{\eta}_l + \sum_A (H_{Aji} - H_{Aij}) N_A^k \tilde{\eta}_k) \\ &= (u_r u^r)^{-1} B_i^l B_j^\mu (\tilde{\nabla}_\mu \tilde{\eta}_l - \tilde{\nabla}_l \tilde{\eta}_\mu) = 2(u_r u^r)^{-1} B_j^\mu B_i^l F_{\mu l} \\ &= 2(u_r u^r)^{-1} f_{ji} = 2G_{ih} f_j^h, \end{aligned}$$

which shows that (η, G) is a contact Riemannian structure on M .

Proposition 6.2. *Let \tilde{M} be a contact Riemannian manifold. In order that a submanifold M in \tilde{M} be a contact Riemannian submanifold, it is necessary and sufficient that the following relations be both valid:*

$$(6.5) \quad u_r u^r = \text{const. ,}$$

$$(6.6) \quad f_A^i = -(u_r u^r)^{-1} \sum_B u_B h_{BA} u^i .$$

Proof. Let M be a contact Riemannian submanifold in \tilde{M} . Then from Proposition 6.1, we have (6.5). On putting

$$(6.7) \quad f_A^i = P_A u^i + P_A^i \quad (A = 1, \dots, 2(n-m)),$$

where P_A^i are vectors orthogonal to u^i , if we transvect (6.7) with u_i , we get $f_A^i u_i = u_i u^i P_A$, which together with (3.13) implies

$$(6.8) \quad P_A = (u_r u^r)^{-1} f_A^i u_i = -(u_r u^r)^{-1} \sum_B u_B h_{BA} .$$

Substituting (6.8) into (6.7), we have

$$(6.9) \quad f_A^t = -(u_r u^r)^{-1} \sum_B u_B h_{BA} u^i + P_A^i,$$

which implies $f_A^i f_{Bi} = (u_r u^r)^{-1} \sum_{C,D} u_D h_{DA} u_C h_{CB} + P_A^i P_{Bi}$ and consequently

$$(6.10) \quad \sum_A f_A^i f_{Ai} = (u_r u^r)^{-1} \sum_{A,B,C} u_B h_{BA} u_C h_{CA} + \sum_A P_A^i P_{Ai}.$$

On the other hand, since M is a contact Riemannian submanifold, from (3.9) we have $u^i f_i^h f_{Ah} = (u_i u^i) u_A - \sum_B f_{Bi} u^i h_{BA} = 0$. Substituting (3.13) into the above equation, we get

$$(6.11) \quad (u_i u^i) u_A = - \sum_{B,C} u_C h_{CB} h_{BA}.$$

Then a combination of (6.10) and (6.11) gives

$$(6.12) \quad \sum_A f_A^i f_{Ai} = \sum_A (u_A^2 + P_A^i P_{Ai}).$$

However, by virtue of (3.8) we obtain $\sum_A f_A^i f_{Ai} = f_{ji} f^{ij} + 2m + 1 - u_i u^i$, which reduces to

$$(6.13) \quad \sum_A f_A^i f_{Ai} = 1 - u_i u^i = \sum_A u_A^2$$

because of (3.14) since M is a contact Riemannian submanifold. Comparing (6.12) with (6.13), we have $\sum_A P_A^i P_{Ai} = 0$, that is, $P_A^i = 0$ ($A = 1, \dots, 2(n-m)$). Hence we obtain (6.6).

Conversely, if the submanifold satisfies (6.5) and (6.6), according to (3.8) we get

$$(6.14) \quad \begin{aligned} f_i^h f_h^j &= -\delta_i^j + u_i u^j + \sum_A f_{Ai} f_A^j \\ &= -\delta_i^j + u_i u^j + (u_r u^r)^{-2} \sum_{A,B,C} u_B h_{BA} u_C h_{CA} u_i u^j. \end{aligned}$$

Since f_{ji} is skew symmetric, the condition (6.6) implies $f_i^h u^i f_{Ah} = (u_i u^i) u_A - \sum_B f_{Bi} u^i h_{BA} = 0$ because of (3.9). Substituting (3.13) into the above equation, we get

$$(6.15) \quad \sum_{B,C} u_C h_{CB} h_{BA} = -(u_i u^i) u_A.$$

Therefore (6.14) reduces to

$$\begin{aligned} f_i^h f_h^j &= -\delta_i^j + u_i u^j + (u_r u^r)^{-1} \sum_B u_B^2 u_i u^j \\ &= -\delta_i^j + (u_r u^r)^{-1} (u_r u^r + \sum_B u_B^2) u_i u^j \\ &= -\delta_i^j + (u_r u^r)^{-1} u_i u^j. \end{aligned}$$

Thus the conditions stated in Proposition 6.1 are satisfied, and the proof is complete.

7. Contact Riemannian submanifolds in a normal contact manifold

Let \bar{M} be a normal contact manifold. In this section we define the notion of a normal contact submanifold M in \bar{M} . After deriving a condition for M to be a normal contact submanifold in \bar{M} , we show that any $(2m + 1)$ -dimensional F -invariant submanifold M in \bar{M} is a normal contact submanifold.

Definition 7.1. Let \bar{M} be a normal contact manifold, and M a contact Riemannian submanifold in \bar{M} . If the induced contact structure of M in \bar{M} is normal, the submanifold M is called a *normal contact submanifold*.

Proposition 7.2. Let M be a normal contact submanifold in \bar{M} , and $'M$ a normal contact submanifold in M . Then $'M$ is a normal contact submanifold in \bar{M} .

Proof. Since M and $'M$ are normal contact submanifolds respectively in \bar{M} and M , there exist two pairs of positive constants (t, c) and (t', c') . Then, as we have seen in § 4, $'M$ becomes a contact Riemannian submanifold in \bar{M} with respect to the pair $(t't, c'c)$. We denote these contact metric structures on M in \bar{M} and on $'M$ in M respectively by (η_i, G_{ji}) and (η_a, G_{ba}) , and denote the contact metric structure on $'M$ in \bar{M} by (η_a, G_{ba}) . Then we have

$$\begin{aligned} 2'\phi_{ba} &= \partial_b'\eta_a - \partial_a'\eta_b = tt'B_b^j B_a^i (\partial_i \tilde{\eta}_j - \partial_j \tilde{\eta}_i) \\ &= tt'B_b^j B_j^i B_a^k B_i^l (\partial_l \tilde{\eta}_k - \partial_k \tilde{\eta}_l) = tt'B_b^j B_a^i (\partial_j u_i - \partial_i u_j) \\ &= t'B_b^j B_a^i (\partial_j \eta_i - \partial_i \eta_j) = \partial_b \eta_a - \partial_a \eta_b = 2\phi_{ba}, \end{aligned}$$

and therefore

$$\begin{aligned} \nabla_c'\phi_{ba} &= \nabla_c \phi_{ba} = \eta_b G_{ca} - \eta_a G_{cb} \\ &= t'c'(\eta_i B_b^j G_{jh} B_c^i B_a^h - \eta_h B_a^h G_{ji} B_c^j B_b^i) \\ &= t't'cc'(\tilde{\eta}_i B_b^j \tilde{g}_{\mu\nu} B_c^\mu B_a^\nu - \tilde{\eta}_\nu B_a^\nu \tilde{g}_{\mu\lambda} B_c^\mu B_b^\lambda) \\ &= \eta_b' G_{ca} - \eta_a' G_{ba}, \end{aligned}$$

which proves by virtue of (1.11) that the structure (η_b, G_{ab}) is normal.

Proposition 7.3. Let M be a contact Riemannian submanifold of a normal contact manifold \bar{M} , and suppose that the dimension of M is greater than the codimension of M in \bar{M} . In order that M be a normal contact submanifold in \bar{M} it is necessary and sufficient that

$$(7.1) \quad \sum_A P_A H_{Aji} = H g_{ji} + K u_j u_i$$

hold, where

$$(7.2) \quad P_A = -(u_r u^r)^{-1} \sum_B u_B h_{BA},$$

and H and K are suitable scalar functions defined on M .

Remark. As it is easily checked, the left hand member of (7.1) is independent of the choice of the unit normal vectors to M .

Proof of Proposition 7.3. Let M be a normal contact submanifold in \bar{M} . Then by the definition of normality we have

$$N_{ji}{}^h = f_j{}^r (\nabla_r f_i{}^h - \nabla_i f_r{}^h) - f_i{}^r (\nabla_r f_j{}^h - \nabla_j f_r{}^h) + \eta_j \nabla_i u^h - \eta_i \nabla_j u^h = 0$$

because of Proposition 4.4. Substituting (3.15) and (3.18) into the above equation and taking account of (4.4), (6.6), Proposition 4.4 and Corollary 4.5, we find

$$(7.3) \quad \begin{aligned} N_{ji}{}^h &= f_j{}^r u_i (\delta_r^h + \sum_A P_A H_A{}^h{}_r) - f_i{}^r u_j (\delta_r^h + \sum_A P_A H_A{}^h{}_r) \\ &+ (u_r u^r)^{-1} \{ (f_i^h + \sum_A u_A H_A{}^h{}_i) u_j - (f_j^h + \sum_A u_A H_A{}^h{}_j) u_i \} = 0. \end{aligned}$$

On the other hand, we know that the vector field ξ^i is a Killing vector field if the contact Riemannian structure is normal. Thus, from (3.18) and (4.7), we have

$$(7.4) \quad \sum_A u_A H_{Aji} = 0.$$

Substituting (7.4) into (7.3) and taking account of (3.14), we obtain

$$N_{ji}{}^h = \{ \sum_A P_A H_{Aji}{}^h - (u_r u^r)^{-1} \sum_A u_A^2 \delta_r^h \} (f_j{}^r u_i - f_i{}^r u_j) = 0,$$

and therefore $\sum_A P_A H_{Aji} = (u_r u^r)^{-1} \sum_A u_A^2 g_{ji} + K u_j u_i$, which proves the necessity of the given condition.

Conversely, suppose that in a contact Riemannian submanifold M in \bar{M} the condition (7.1) holds. Differentiating

$$(7.5) \quad f_i = \frac{P u_i}{A}$$

covariantly, we get $\nabla_j f_{Ai} = \nabla_j P_A u_i + P_A \nabla_j u_i$. Substituting (3.16) and (3.18) into the above equation, we find

$$\begin{aligned} & - u_A g_{ji} + \sum_B (H_{Bji} h_{BA} - f_i L_{BAj}) - f_i^h H_{Ajh} \\ & = \nabla_j P_A u_i + P_A (f_{ji} + \sum_B u_B H_{Bji}), \end{aligned}$$

which together with (7.5) implies

$$\begin{aligned}
& - \sum_A P_A u_A g_{ji} + \sum_{B,A} (H_{Bji} h_{BA} P_A - P P u_i L_{BAj}) - \sum_A P_A H_{Aji} f_i^h \\
& = \sum_A u_i P_A \nabla_j P_A + \sum_A \frac{P^2}{A} (f_{ji} + \sum_B u_B H_{Bji}) .
\end{aligned}$$

Transvecting this with f^{ji} and making use of (7.1), we get $-f^{ji} f_i^h (H g_{jh} + K u_j u_h) = 2m \sum_A P_A^2$ from which $H = \sum_A P_A^2$. Therefore (7.1) reduces to

$$(7.6) \quad \sum_A P_A H_{Aji} = \sum_A P_A^2 g_{ji} + K u_j u_i .$$

Substituting (7.6) into the left hand member of (7.3), we find

$$\begin{aligned}
(7.7) \quad N_{ji}^h &= (f_j^h u_i - f_i^h u_j) (1 + \sum_A \frac{P^2}{A} - (u_r u^r)^{-1}) \\
&= (u_r u^r)^{-1} (u_r u^r + u_r u^r \sum_A \frac{P^2}{A} - 1) (f_j^h u_i - f_i^h u_j) .
\end{aligned}$$

On the other hand, (7.2) and (6.11) imply

$$\sum_A P_A^2 = (u_r u^r)^{-2} \sum_{B,C} u_B h_{BA} u_C h_{CA} = (u_r u^r)^{-1} \sum_C u_C^2 .$$

Thus, from (3.14) and (7.7) it follows that $N_{ji}^h = 0$, which completes the proof of the sufficiency.

Corollary 7.4. *Let M be a contact Riemannian submanifold in a normal contact manifold \bar{M} . If M is a totally geodesic or a totally umbilical submanifold in \bar{M} , then M is a normal contact submanifold.*

As we have mentioned in the previous paper [3], every totally umbilical submanifold M in a normal contact manifold \bar{M} is not a normal contact submanifold. In [3] we have proved that a normal contact submanifold of codimension 2 in a normal contact manifold of constant curvature is either an F -invariant submanifold or a totally umbilical submanifold. However, if the codimension is greater than 2 we cannot prove this fact, because by Proposition 7.2, for example, an F -invariant submanifold M in a totally umbilical submanifold \bar{M} is also a normal contact submanifold in \bar{M} . In general, a normal contact submanifold in a normal contact manifold is neither F -invariant nor totally umbilical.

Proposition 7.5. *An F -invariant submanifold in a normal contact manifold is a normal contact submanifold.*

Proof. Since the submanifold is F -invariant, it follows that $f_A^i = 0$, $u_A = 0$ ($A = 1, \dots, 2(n-m)$). Consequently we have $u_i u^i = 1$ because of (3.14). Substituting these into the left hand member of (7.3), we find

$$N_{ji}^h = (1 - (u_r u^r)^{-1}) (f_j^h u_i - f_i^h u_j) = 0 ,$$

which completes the proof.

Bibliography

- [1] Y. Hatakeyama, *On the existence of Riemann metrics associated with a 2-form of rank $2r$* , Tôhoku Math. J. **14** (1962) 162–166.
- [2] Y. Hatakeyama, Y. Ogawa & S. Tanno, *Some properties of manifolds with contact metric structure*, Tôhoku Math. J. **15** (1962) 42–48.
- [3] M. Okumura, *On contact metric immersion*, Kōdai Math. Sem. Rep. **20** (1968) 389–409.
- [4] S. Sasaki, *On differentiable manifolds with certain structures which are closely related to almost contact structure. I*, Tôhoku Math. J. **12** (1960) 459–476.
- [5] K. Yano & S. Ishihara, *Invariant submanifold of almost contact manifold*, Kōdai Math. Sem. Rep. **21** (1969) 350–364.

SAITAMA UNIVERSITY, JAPAN