

FUNCTIONS OF TRANSITION FOR CERTAIN KÄHLER MANIFOLDS

CARL VERHEY

1. Introduction

In [1] Adler has shown that Kähler metrics can be classified by geometric conditions of the image of an isometry into certain Grassmannians. In this paper, we find a necessary condition on the isometry which will guarantee that the original metric was in fact a Hodge metric. (The cohomology class of the fundamental form of the metric belongs to an integral cohomology class.)

Some standard conventions are observed. Differentiable will mean differentiable of class C^∞ . If φ is a mapping, φ_* will denote the induced map in tangent spaces. Lower case letters will denote the Lie algebra, upper case letters the Lie group. For example, $\mathfrak{o}(n)$ will denote the Lie algebra of the orthogonal group $O(n)$. Finally, if g is an element of a matrix group, g^t will denote the transpose of g .

The results of this paper are part of the author's Ph. D. thesis, which was written at Purdue University under A. W. Adler.

The following material can be found in [1]. We include it here for the sake of completeness.

By a modification of Nash's theorem on isometric imbeddings in Euclidean space, it can be shown that every k -dimensional Riemannian manifold M can be isometrically imbedded in S^{k+p-1} (the unit sphere in E^{k+p}) where p is a large positive integer depending on K but not on M .

Let $B_{0(2n)}^+ = O(2n+p)/O(2n) \times O(p-1)$. Then $B_{0(2n)}^+$ can be considered as the set of all pairs (P_1, P_2) , where P_1 is a $2n$ -plane in $E(2n+p)$ through the origin, and P_2 is a vector in $E(2n+p)$ orthogonal to P_1 . Let F be an isometric imbedding of a $2n$ -dimensional Riemannian manifold M into S^{2n+p-1} . Each point $F(m)$ of $F(M)$ defines an element of $B_{0(2n)}^+$ (i.e., a pair (P_1, P_2)) as follows: P_1 is to be the tangent space of $F(M)$ at $F(m)$ translated to the origin in $E(2n+p)$, and P_2 is to be the position vector of $F(m)$. The mapping $\pi: F(M) \rightarrow B_{0(2n)}^+$ defined by $\pi(m) = (P_1, P_2)$ is called the spherical image mapping; on composition with F , it determines a map f of M into $B_{0(2n)}^+$.

Let B' be the bundle of orthonormal bases over M . Then B' is the space of all $(2n+1)$ -tuples (m, e_1, \dots, e_{2n}) , where m is a point in M , and e_1, \dots, e_{2n} is an orthonormal basis for M_m . Define a mapping


Received September 19, 1968.

$$g: B' \rightarrow V_{2n,p}^+ = 0(2n + p)/0(p - 1)$$

by


$$g(m: e_1, \dots, e_{2n}) = [F(m), F'_*(e_1), \dots, F'_*(e_{2n})],$$

where $F'_*(e_i)$ denotes the vector derived from $F_*(e_i)$ by parallel translation to the origin in E^{2n+p} . The following diagram is the commutative



where $\lambda, \tau,$ and α are the natural mappings.

Let M be a hermitian manifold. Then the bundle B' of orthonormal bases of M is reducible to a principal $U(n)$ -bundle B over M , and the condition that M be Kähler is equivalent to the existence of a torsionless connection on B . Let $B_{U(n)}^+ = 0(2n + p)/U(n) \times 0(p - 1)$. Then there is a mapping \bar{f} of M into $B_{U(n)}^+$ such that the following diagram commutes:



where τ, τ^1, ∂ are the natural mappings.

Let x be a tangent vector to $0(2n + p)$, and denote by X the element of $0(2n + p)$ defined by x . Define $w(x)$ to be the projection of X into $\mathfrak{o}(2n)$. Since the 1-form w is horizontal over $V_{2n,p}^+$ and right invariant under the action of $0(p - 1)$ there is a $\mathfrak{o}(2n)$ -valued 1-form w^1 on $V_{2n,p}^+$ such that $\sigma^*(w^1) = w$. A vector x on $B_{U(n)}^+$ is said to be H -horizontal if there is a vector y on $V_{2n,p}^+$ with $x = \tau^*(y)$ and $w^1(y) = 0$.

The importance of the notion of H -horizontality is seen in the following theorem:

Theorem 1. *A $2n$ -dimensional Riemannian manifold M is a Kähler manifold if and only if it admits a mapping g into $B_{U(n)}^+$ such that:*

- (a) $g(M)$ is H -horizontal,
- (b) the projection of $g(M)$ into $B_{0(2n)}^+$ is the spherical image of the projection of $g(M)$ into S^{2n+p-1} .

A $2n$ -dimensional submanifold M of $B_{U(n)}^+$ is said to be a K -manifold if it satisfies the following two conditions:

- (a) M is H -horizontal, that is, every tangent vector of M is H -horizontal.
- (b) The projection of M into $B_{0(2n)}^+$ is the image under the spherical map of the projection of M into S^{2n+p-1} .

By Theorem 1, every K -manifold can be identified with a Kähler manifold. In fact, it can be shown that every K -manifold \underline{M} induces a partial Hermitian metric h in $B_{U(n)}^+$. Let $\hat{\Omega}$ denote the fundamental form of h , and Ω be the restriction of $\hat{\Omega}$ to \underline{M} . Then Ω is the fundamental form of the natural Kähler metric and complex structure on \underline{M} , and we have

Theorem 2. *Let M be a $2n$ -dimensional manifold with Riemannian metric r .*

1. *r is the real part of a Kähler metric of an almost complex structure on M if and only if M admits a differentiable isometric imbedding f onto a K -manifold. In case such an f exists, it is in fact a homeomorphic isometry with respect to the natural Kähler metric and complex structure of $f(M)$.*

2. *If r is the real part of a Kähler metric of a complex analytic structure on M , then $f^*(\hat{\Omega})$ is the fundamental form of the metric.*

Let x be a tangent vector of $0(2n + p)$, and denote by X the element of $o(2n + p)$ defined by x . Define 1-forms w_0 and w' as follows:

$w_0(x)$ is to be the projection of X into $u(n)$, the Lie algebra of $U(n)$, and $w'(x)$ is to be the projection of X into $o(2n + p - 1)$. Identify $o(2n + p)$ with the space of $(2n + p) \times (2n + p)$ skew symmetric real matrices, and denote by $w_{i,j}$ the 1-form which assigns to each matrix its (i, j) th entry, $1 \leq i, j \leq 2n + p$. Let

$$\text{trace Im } X = \sum_{k=1}^n w_{k, n+k}(w_0(X)) = \sum_{k=1}^n w_{k, n+k}(X) .$$

Note that trace Im is invariant under the action of $U(n) \times 0(p - 1)$. Finally let Ω' denote the curvature form of w' .

Let M be a compact complex analytic manifold with a Kähler metric h , and f be a differential isometric imbedding of M into a k -manifold. Then the 2-forms trace Im dw_0 , trace Im $w' \wedge w'$, and trace Im Ω' are horizontal over $f(M)$. Since they are also invariant under the right action of $U(n) \times 0(p - 1)$, they induce 2-forms on $f(M)$. Denote these 2-forms by Trace Im dw_0 , Trace Im $w' \wedge w'$, and Trace Im Ω' , respectively.

Proposition 1. (a) $(1/2\pi)\underline{f}^*(\text{Trace Im } dw_0)$ is the first Chern form of the Kähler metric on M .

(b) $(1/2\pi)\underline{f}^*(\text{Trace Im } \Omega')$ is the fundamental form of the Kähler metric on M .

(c) $\text{Trace Im } \Omega' = \text{Trace Im } dw_0 + \text{Trace Im } w' \wedge w'$.

The fact that $w = w_0$ on $f^{-1}[\underline{f}(M)]$ implies that $\text{Trace Im } w' \wedge w' = \sum_{i=1}^n \sum_{\alpha=2+1}^{2n+p-1} w_{i\alpha} \wedge w_{i+n\alpha}$. We will denote this form by Ω^\perp .

2. A condition

A K -manifold M' contained in $B_{U(n)}^+$ will be said to be special if each m' of M' has a neighborhood $V(m')$ which admits a cross-section σ_v into $\delta^{-1}(V)$ such that $d(\sigma_v^* w^\perp) = 0$, where w^\perp denotes the $\mathfrak{o}(2n + p - 1)$ valued 1-form $w' - w$.

A complex analytic manifold M together with a Kähler metric $K(\cdot, \cdot)$ on M will be said to be a special Kähler manifold if it admits an isometric imbedding F into S^{2n+p-1} (for some p) such that $\underline{f}(M)$ is a special K -manifold. Let D denote covariant differentiation with respect to the connection w' on $\mathfrak{o}(2n + p)$ as a bundle over S^{2n+p-1} .

Proposition 2. Let $(M, K(\cdot, \cdot))$ be a special Kähler manifold, F be as prescribed, and $m \in M$. Then there is an orthonormal basis e_1, \dots, e_{2n+p-1} of vector fields tangent to S^{2n+p-1} and defined on some neighborhood $F(U(m))$ of $F(m)$ such that:

- (a) e_1, \dots, e_{2n} is a basis for the tangent space of $F(U)$,
- (b) $e_{2n+1}, \dots, e_{2n+p-1}$ is a basis for the orthonormal complement to the tangent space of $F(U)$,

(c) $dw_{i,\alpha}^{u,e} = 0$ for $i = 1, \dots, 2n; \alpha = 2n + 1, \dots, 2n + p - 1$ where $w_{i,\alpha}^{u,e}(x) = \langle D_x e_i, e_\alpha \rangle$.

The converse is also true.

Proof. Given a cross-section σ_u on a neighborhood $\underline{f}(U)$ of a point $\underline{f}(m)$, one gets an orthonormal basis for vector fields tangent to S^{2n+p-1} and defined in a neighborhood $F(U)$ of $F(m)$ satisfying (a) and (b). Conversely, such an orthonormal basis e_1, \dots, e_{2n+p-1} gives a cross-section $\sigma_u(\underline{f}(m)) = \{F(m), e_1, \dots, e_{2n+p-1}\}$ defined on the neighborhood $\underline{f}(U)$ of $\underline{f}(m)$. So it suffices to show that $d(\sigma_u^*(w^\perp)) = 0$ if and only if $dw_{i,\alpha}^{u,e} = 0$ for all $i = 1, \dots, 2n; \alpha = 2n + 1, \dots, 2n + p - 1$. But this is immediate since, in fact, $w_{i,\alpha}^{u,e} = \beta^* \sigma_u^*((w^\perp)_{i\alpha})$, this last statement being the equivalence of the Cartan and bundle definitions of a connection.

3. The isomorphism between de Rahm and Čech cohomology for special K -manifolds

Let M be a special K -manifold contained in $B_{U(n)}^+$, m a point of M , and

$U(m)$ a neighborhood of m in M admitting a cross-section $\sigma_{u(m)}$ into $0(2n + p)$ such that $\sigma_{u(m)}^*w^\perp$ is closed. Then $\mathcal{U} = \{U(m); m \in M\}$ is an open covering of M . Let $\mathcal{V} = \{V_s; s \in S\}$ be a locally finite (differentiably) simple refinement of the covering \mathcal{U} , [3]. Since \mathcal{V} is a refinement of \mathcal{U} , each V_s is contained in some member of the covering \mathcal{U} . Hence, for each s in S , there is a cross-section σ_s defined on V_s such that $d(\sigma_s^*w^\perp) = 0$. Since each V_s is simply connected, there are functions $h_{i\alpha}^s, i = 1, \dots, 2n; \alpha = 2n + 1, \dots, 2n + p - 1$, such that $dh_{i\alpha}^s = \sigma_s^*w_{i\alpha}$ on V_s . Let h^s be the skew symmetric $(2n + p) \times (2n + p)$ matrix whose (i, α) th entry is $h_{i\alpha}^s$ for $i = 1, \dots, 2n; \alpha = 2n + 1, \dots, 2n + p - 1$ and whose remaining entries above the diagonal are zero. Let h^s be the skew symmetric matrix defined on $\delta^{-1}(V_s)$ by $h^s = \underline{h}^s \circ \delta$, where δ is the natural projection of $0(2n + p)$ onto $B_{u(n)}^+$.

Lemma 1. On $\sigma_s(V_s), dh^s = w^\perp$.

Proof. $\sigma_s^*dh^s = \sigma_s^*(d(\underline{h}^s \circ \delta)) = \sigma_s^*(\delta^*d\underline{h}^s) = (\delta \circ \sigma_s)^*(d\underline{h}^s) = \sigma_s^*w^\perp$.

Let R_g denote right translation along the fiber for $\delta^{-1}(V_s)$ by an element g of $U(n) \times 0(p - 1)$. Then $\delta \circ R_g = \delta$, for all g in $U(n) \times 0(p - 1)$. For each b in $\delta^{-1}(V_s)$ define $g_s(b)$ to be the element of $U(n) \times 0(p - 1)$ such that $R_{g_s(b)}(b) = \sigma_s \circ \delta(b)$, that is, $g_s(b)$ is to be the element of $U(n) \times 0(p - 1)$ such that right translation by $g_s(b)$ carries b to the point of the cross-section $\sigma_s(V_s)$ lying in the same fiber as b .

Lemma 2. Let b be any point of $\delta^{-1}(V_s)$. Then

$$w_b^\perp = g_s^t(b)(R_{g_s(b)}^*(dh^s(\delta_s \circ \delta(b))))g_s(b).$$

Proof. Since $b \in \delta^{-1}(V_s)$, b can be written as $(m, f_1, \dots, f_{2n+p-1})$, where $m = \alpha \circ \delta \circ \delta(b), f_1, \dots, f_{2n}$ is an orthonormal basis for the vector fields tangent to $\alpha \circ \delta(M)$ on some neighborhood of the point m , and $f_{2n+1}, \dots, f_{2n+p-1}$ is an orthonormal basis for the orthogonal complement to the tangent space of M on this neighborhood. Here, as before, $\alpha \circ \delta$ denotes the natural projection of $B_{U(n)}^+$ onto S^{2n+p-1} . Let e_1, \dots, e_{2n+p-1} be the orthonormal vector fields defined by the section σ_b . Then by definition of $g_s(b)$, we have

$$(f_1(m), \dots, f_{2n+p-1}(m)) = (e_1^{(m)}, \dots, e_{2n+p-1}^{(m)})g_s(b),$$

where $m = \alpha \circ \delta \circ \delta(b)$. Let x be any tangent vector of $\delta^{-1}(V_s)$ at the point b and $\underline{x} = (\alpha \circ \delta \circ \delta)_*x$. Then

$$\begin{aligned} (w_{i\alpha}(x))_b &= \langle f_i, D_{\underline{x}}f_\alpha \rangle = \left\langle \sum_{k=1}^{2n} g_{ki}e_k, D_{\underline{x}} \sum_{\beta=2n+1}^{2n+p-1} g_{\beta\alpha}e_\beta \right\rangle \\ &= \sum_{k=1}^{2n} \sum_{\beta=2n+1}^{2n+p-1} g_{ki} \langle e_k, D_{\underline{x}}(g_{\beta\alpha}e_\beta) \rangle = \sum_{k=1}^{2n} \sum_{\beta=2n+1}^{2n+p-1} g_{ki} \langle e_k, \underline{x}(g_{\beta\alpha})e_\beta + g_{\beta\alpha}D_{\underline{x}}e_\beta \rangle \\ &= \sum_{k=1}^{2n} \sum_{\beta=2n+1}^{2n+p-1} g_{ki} \langle e_k, D_{\underline{x}}e_\beta \rangle g_{\beta\alpha} = \sum_{k=1}^{2n} \sum_{\beta=2n+1}^{2n+p-1} g_{ki} [w_{k\beta}(R_{g_s(b)}^*(x))]_{\sigma_s \delta(b)} g_{\beta\alpha}. \end{aligned}$$

where $1 \leq k \leq 2n$, $2n - 1 \leq \beta \leq 2n + p - 1$. Since by Lemma 1, $\sigma_s^* dh^s \equiv \sigma_s^* w^\perp$, the assertion now follows.

Define an operator T on pairs A_1, A_2 of $(2n + p) \times (2n + p)$ matrices by $T(A_1, A_2) = \text{trace Im } [A_1, A_2]$ where $[A_1, A_2]$ denotes the matrix $A_1 A_2 - A_2 A_1$. Finally, define a 1-form α_s on $\delta^{-1}(V_s)$ by

$$\alpha_s(X)_b = T[g_s^t(b)h^s(b)g_s(b), w_b^\perp(X)]$$

for any tangent vector X of $\delta^{-1}(V_s)$ at the point b . Recall that Ω^\perp denotes the 2-form $\sum_{\alpha=2n+1}^{2n+p-1} \sum_{i=1}^n w_{i\alpha} \wedge w_{i+n\alpha} = \text{trace Im } w^\perp \wedge w^\perp$.

Proposition 3. *On $\delta^{-1}(V_s)$, $d\alpha_s/2 = \Omega^\perp$.*

Proof. Let b be any point of $\delta^{-1}(V_s)$. Then

$$w_b^\perp = g_s^t(b)(R_{g_s^s(b)}^*(dh_{\sigma_s^s(b)}^s))g_s(b).$$

Since trace Im is right invariant under the action of $U(n) \times O(p - 1)$, we have

$$\begin{aligned} \alpha_s &= T(g^t h^s g_s, w^\perp) = \text{trace Im } (g_s^t h^s g_s w^\perp - w^\perp g_s^t h^s g_s) \\ &= \text{trace Im } (g_s^t h^s g_s g_s^t (R_{g_s^s(b)}^* dh^s) g_s - g_s^t (R_{g_s^s(b)}^* dh^s) g_s g_s^t h^s g_s) \\ &= \text{trace Im } (h^s(b) R_{g_s^s(b)}^* dh_{\sigma_s^s(b)}^s - R_{g_s^s(b)}^* dh_{\sigma_s^s(b)}^s h^s), \end{aligned}$$

which becomes, in consequence of $h^s(\sigma_s \delta(b)) = h^s(b)$,

$$\alpha_s = \text{trace Im } (h^s(b) dh_b^s - dh_b^s h^s(b)).$$

Thus

$$\begin{aligned} d\alpha_s &= \text{trace Im } (dh_b^s \wedge dh_b^s + dh_b^s dh_b^s) \\ &= 2 \text{trace Im } (R_{g_s^s(b)}^* (dh_{\sigma_s^s(b)}^s \wedge dh_{\sigma_s^s(b)}^s)) \\ &= 2R_{g_s^s(b)}^* \text{trace Im } (w_{\sigma_s^s(b)}^\perp \wedge w_{\sigma_s^s(b)}^\perp) = 2\Omega_b^\perp, \end{aligned}$$

since the 2-form Ω_b^\perp is invariant under the right action of $U(n) \times O(p - 1)$.

Now let r and s be elements of the index set S such that $V_r \cap V_s$ is not empty. For each m in $V_r \cap V_s$, let $g_{rs}(m)$ be the element of $U(n) \times O(p - 1)$ such that

$$\sigma_s(m) = R_{g_{rs}(m)}(\sigma_r(m)).$$

If b is any point in $\delta^{-1}(V_r \cap V_s)$, then $g_s(b)g_r^t(b) = g_{rs}(\delta(b))$.

Lemma 3. *On $V_r \cap V_s$, the 1-form*

$$T(h^s g_{sr}, h^r dg_{rs}^t) + T(h^s dg_{rs}, h^r g_{rs}^t)$$

is closed.

Proof. By an application of Lemma 3 we have $dh^r = g_{rs}^t dh^s g_{rs}$. Thus $dg_{rs}^t dh^s g_{rs} = g_{rs}^t dh^s dg_{rs}$, and

$$\begin{aligned} & d(T(\underline{h}^s g_{rs}, \underline{h}^r dg_{rs}^t) + T(\underline{h}^s dg_{rs}, \underline{h}^r g_{rs}^t)) \\ &= d \operatorname{trace} \operatorname{Im} (\underline{h}^s g_{rs} \underline{h}^r dg_{rs}^t - \underline{h}^r dg_{rs}^t \underline{h}^s g_{rs}) \\ & \quad + d \operatorname{trace} \operatorname{Im} (-\underline{h}^r g_{rs}^t \underline{h}^s dg_{rs} + \underline{h}^s dg_{rs} \underline{h}^r g_{rs}^t) \\ &= \operatorname{trace} \operatorname{Im} (d(\underline{h}^s g_{rs} \underline{h}^r dg_{rs}^t + \underline{h}^s dg_{rs} \underline{h}^r g_{rs}^t) \\ & \quad - \operatorname{trace} \operatorname{Im} (d(\underline{h}^r dg_{rs}^t \underline{h}^s g_{rs} + \underline{h}^r g_{rs}^t \underline{h}^s dg_{rs})). \end{aligned}$$

The first term above is equal to

$$\begin{aligned} & \operatorname{trace} \operatorname{Im} (d\underline{h}^s g_{rs} \underline{h}^r dg_{rs}^t + \underline{h}^s g_{rs} d\underline{h}^r dg_{rs}^t + d\underline{h}^s dg_{rs} \underline{h}^r g_{rs}^t - \underline{h}^s dg_{rs} d\underline{h}^r g_{rs}^t) \\ &= \operatorname{trace} \operatorname{Im} ((-dg_{rs}^t d\underline{h}^s g_{rs} + g_{rs}^t d\underline{h}^s dg_{rs})(\underline{h}^r)) \\ & \quad + \operatorname{trace} \operatorname{Im} (\underline{h}^s (g_{rs} d\underline{h}^r dg_{rs}^t - dg_{rs} d\underline{h}^r g_{rs}^t)) = 0. \end{aligned}$$

Similarly,

$$\operatorname{trace} \operatorname{Im} (d(\underline{h}^r dg_{rs}^t \underline{h}^s g_{rs} + \underline{h}^r g_{rs}^t \underline{h}^s dg_{rs})) = 0.$$

Since $V_r \cap V_s$ is simply connected, there exists a function f_{rs}^1 such that

$$df_{rs}^1 = T(\underline{h}^s g_{rs}, \underline{h}^r dg_{rs}^t) + T(\underline{h}^s dg_{rs}, \underline{h}^r g_{rs}^t)$$

on $V_r \cap V_s$. Define a function f_{rs}^2 on $V_r \cap V_s$ by $f_{rs}^2 = T(\underline{h}^s g_{rs}, \underline{h}^r g_{rs}^t)$ and let $\underline{\alpha}_s$ (resp. $\underline{\alpha}_r$) be the 1-form $\sigma_s^*(\alpha_s)$ (resp. $\sigma_r^*(\alpha_r)$).

Proposition 4. On $V_r \cap V_s$, $\underline{\alpha}_s - \underline{\alpha}_r = d(f_{rs}^2 - f_{rs}^1)$.

$$\begin{aligned} \text{Proof.} \quad \underline{\alpha}_s - \underline{\alpha}_r &= T(\underline{h}^s, d\underline{h}^s) - T(\underline{h}^r, d\underline{h}^r) \\ &= T(\underline{h}^s, g_{rs} d\underline{h}^r g_{rs}^t) - T(\underline{h}^r, g_{rs}^t d\underline{h}^s g_{rs}) \\ &= T(\underline{h}^s, g_{rs} d\underline{h}^r g_{rs}^t) + T(g_{rs}^t d\underline{h}^s g_{rs}, \underline{h}^r). \end{aligned}$$

Since $T = \operatorname{trace} \operatorname{Im} ([,])$, and $\operatorname{trace} \operatorname{Im}$ is invariant under the right action of $U(n) \times O(p-1)$, we have:

$$\begin{aligned} \underline{\alpha}_s - \underline{\alpha}_r &= T(\underline{h}^s g_{rs}, d\underline{h}^r g_{rs}^t) + T(d\underline{h}^s g_{rs}, \underline{h}^r g_{rs}^t) \\ &= d(T(\underline{h}^s g_{rs}, \underline{h}^r g_{rs}^t)) - T(\underline{h}^s g_{rs}, \underline{h}^r dg_{rs}^t) - T(\underline{h}^s dg_{rs}, \underline{h}^r g_{rs}^t) \\ &= d(f_{rs}^2 - f_{rs}^1). \end{aligned}$$

Now let r, s , and t be any elements of the index set S such that $V_r \cap V_s \cap V_t$ is not empty, and let

$$a_{rst} = (f_{rs}^2 - f_{rt}^2 + f_{st}^2) - (f_{rs}^1 - f_{rt}^1 + f_{st}^1).$$

Let $\{a\}$ denote the cohomology class of $\hat{H}^2(M, R)$ of which $[(1/4\pi)a_{rst}]$ is a representative, and $\{\text{trace Im } w^\perp \wedge w^\perp\}$ the cohomology class of the 2-form $\text{trace Im } w^\perp \wedge w^\perp$ in $H^2(M)$.

Theorem 3. *Let M be a special K -manifold, and $[a]$ as defined above. If ϕ denotes the isomorphism of $H^2(M)$ onto $H^2(V, R)$, then*

$$\phi(\{\text{trace Im } w^\perp \wedge w^\perp\}) = \{a\}.$$

Proof. The assertion follows immediately from Propositions 3 and 4.

4. A sufficient condition for a special Kähler manifold to be a Hodge manifold

Theorem 4. *Let $(M, K(\cdot, \cdot))$ be a special Kähler manifold. Suppose moreover that the matrices h^s can be chosen so that $dg_{rs}h^r g_{rs}^t + g_{rs}h^r dg_{rs}^t$ vanishes whenever r and s are elements of the index set S such that $V_r \cap V_s$ is not empty. Then $K(\cdot, \cdot)$ is a Hodge metric.*

The remainder of this section is devoted to the proof of this theorem.

Lemma 1. *Under the conditions of Theorem 4 the functions f_{rs}^1 can be chosen to be identically zero.*

Proof. By definition,

$$\begin{aligned} df_{rs}^1 &= T(h^s g_{rs}, h^r dg_{rs}^t) + T(h^s dg_{rs}, h^r g_{rs}^t) \\ &= T(h^s, g_{rs} h^r dg_{rs}^t + dg_{rs} h^r g_{rs}^t) = 0 \end{aligned}$$

on $V_r \cap V_s$. Thus f_{rs}^1 is a constant and, in fact, can be chosen to be zero.

Define constant matrices c_{rs} by $c_{rs} = h^s - g_{rs} h^r g_{rs}^t$ for all r, s in the index set S such that $V_r \cap V_s$ is not empty.

Lemma 2. *If r, s, t are elements of the index set S such that $V_r \cap V_s \cap V_t$ is not empty, then*

$$c_{rt} + g_{rt} c_{sr} g_{rt}^t - c_{st} = 0.$$

Proof. We have

$$h^r = g_{sr} h^s g_{sr}^t + c_{sr},$$

$$h^t = g_{st} h^s g_{st}^t + c_{st},$$

$$h^t = g_{rt} h^r g_{rt}^t + d_{rt},$$

so,

$$\begin{aligned} c_{rt} + g_{rt} c_{sr} g_{rt}^t - c_{st} &= g_{st} h^s g_{st}^t - g_{rt} h^r g_{rt}^t + g_{rt} c_{rs} g_{rt}^t \\ &= g_{rt} (g_{sr} h^s g_{rs}^t - h^r) g_{rt}^t + g_{rt} c_{sr} g_{rt}^t \\ &= -g_{rt} (c_{sr}) g_{rt}^t + g_{rt} c_{sr} g_{rt}^t = 0. \end{aligned}$$

Now fix any element s of the index set S and for each r in S such that $V_r \cap V_s$ is not empty, define $\hat{h}^r = h^r - c_{sr}$.

Lemma 3. *Under the same hypothesis as in Lemma 2,*

$$g_{rt}\hat{h}^r g_{rt} = \hat{h}^t .$$

Proof.
$$\begin{aligned} g_{rt}\hat{h}^r g_{rt}^t &= g_{rt}(h^r - c_{sr})g_{rt}^t = g_{rt}h^r g_{rt}^t - g_{rt}c_{sr}g_{rt}^t \\ &= g_{rt}h^r g_{rt}^t + c_{rt} - c_{st} = h^t - c_{st} = \hat{h}^t . \end{aligned}$$

Let \tilde{S} denote all elements r of the index set S such that $V_r \cap V_s$ is not empty. If u is an element of S such that $V_u \cap V_r$ is not empty for some r in \tilde{S} , define $\hat{h}^u = h^u - \hat{c}_{ru}$, where \hat{c}_{ru} is defined by $\hat{c}_{ru} = h^u - g_{ru}\hat{h}^r g_{ru}^t$.

Lemma 4. *\hat{h}^u is well defined.*

Proof. We must show that if r and t are elements of \tilde{S} such that $V_u \cap V_r$ and $V_u \cap V_t$ are not empty, then $h^u - \hat{c}_{ru} = h^u - \hat{c}_{tu}$, that is, $\hat{c}_{ru} = \hat{c}_{tu}$. But, as before, $\hat{c}_{ru} + g_{ru}\hat{c}_{tr}g_{ru}^t - \hat{c}_{tu} = 0$, and, by Lemma 3, $\hat{c}_{tr} = 0$. Hence the lemma follows.

Continuing this process defines matrices \hat{h}^r for each r in the index set S in the same connected component as V_s such that $g_{rt}\hat{h}^r g_{rt}^t = \hat{h}^t$ for all r and t with $V_r \cap V_t$ not empty. Doing this for every connected component gives matrices \hat{h}^r for each r in S such that $g_{rt}\hat{h}^r g_{rt}^t = \hat{h}^t$ for all r and t such that $V_r \cap V_t$ is not empty.

Lemma 5. *The cohomology class $[a]$ vanishes.*

Proof. Since $d\hat{h}^s = dh^s$ for all s , and trace Im is invariant under the right action of $U(n) \times O(p-1)$, a representative of $[a]$ is

$$\begin{aligned} a_{rst} &= T(\hat{h}^s g_{rs}, \hat{h}^r g_{rs}^t) - T(\hat{h}^t g_{rt}, \hat{h}^r g_{rt}^t) + T(\hat{h}^t g_{st}, \hat{h}^s g_{st}^t) \\ &= T(g_{rs}^t \hat{h}^s g_{rs}, \hat{h}^r) - T(g_{rt}^t \hat{h}^t g_{rt}, \hat{h}^r) + T(g_{st}^t \hat{h}^t g_{st}, \hat{h}^s) . \end{aligned}$$

Since $T(\hat{h}^r, \hat{h}^r) = \text{trace Im}([\hat{h}^r, \hat{h}^r]) = 0$ for all r in S , we have

$$\begin{aligned} a_{rst} &= T(g_{rs}^t \hat{h}^s g_{rs} - \hat{h}^r, \hat{h}^r) - T(g_{rt}^t \hat{h}^t g_{rt} - \hat{h}^r, \hat{h}^r) + T(g_{st}^t \hat{h}^t g_{st} - \hat{h}^s, \hat{h}^s) \\ &= 0 \end{aligned}$$

by the definition of h . Thus the cohomology class $[a]$ vanishes.

Lemma 6. *If $\{a\}$ vanishes, then $K(\cdot, \cdot)$ is a Hodge metric.*

Proof. By Proposition 1, $(1/2\pi)\Omega^\perp = \Omega - c_1$, where c_1 is the first Chern form of M , and Ω is the fundamental form of the metric $K(\cdot, \cdot)$. By Theorem 3, $\phi(\Omega^\perp) = \{a\}$. Thus $\{a\}$ vanishes, so that the first Chern form and the fundamental form of the metric are cohomologous. Since the first Chern form is integral, the assertion follows.

References

- [1] A. Adler, *Classifying spaces for Kähler Metrics*, Math. Ann. **152** (1963) 164–184.
- [2] K. Kodaira, *Kähler varieties of restricted type*, Ann. of Math. **60** (1954) 28–48.
- [3] A. Weil, *Sur les théorèmes de Riemann-Roch*, Comment. Math. Helv. **26** (1952) 119–145.
- [4] ———, *Variétés kählériennes*, Hermann, Paris, 1958.

UNIVERSITY OF SOUTHERN CALIFORNIA