

## A FORMULA FOR THE RADIAL PART OF THE LAPLACE-BELTRAMI OPERATOR

SIGURDUR HELGASON

Let  $V$  be a manifold and  $H$  a Lie transformation group of  $V$ . Suppose  $Du = 0$  is a differential equation on  $V$ , both the differential operator  $D$  and the function  $u$  assumed invariant under  $H$ . Then the differential equation will involve several inessential variables, a fact which may render general results about differential operators rather ineffective for the differential equation at hand. Thus although  $D$  may not be an elliptic operator it might become one after the inessential variables are eliminated (cf. [3, p. 99]).

This viewpoint leads to the general definition (cf. [7]) of the transversal part and radial part of a differential operator on  $V$  given in §§ 2 and 3. The radial part has been constructed for many special differential operators in the literature; see for example [1], [3], [4], [5], [8] for Lie groups, Lie algebras and symmetric spaces, [9], [6] for some Lorentzian manifolds. Our main result, formula (3.3) in Theorem 3.2, includes various known examples worked out by computations suited for each individual case. See Harish-Chandra [4, p. 99] for the Laplacian on a semisimple Lie algebra, Berezin [1] and Harish-Chandra [3, § 8] for the Laplacian on a semisimple Lie group, and Harish-Chandra [5, § 7] and Karpelevič [8, § 15] for the Laplacian on a symmetric space. The author is indebted to J. Lepowsky for useful critical remarks.

**Notation.** If  $V$  is a manifold and  $v \in V$ , then the tangent space to  $V$  at  $v$  will be denoted  $V_v$ ; the differential of a differentiable mapping  $\varphi$  of one manifold into another is denoted  $d\varphi$ . We shall use Schwartz' notation  $\mathcal{E}(V)$  (resp.  $\mathcal{D}(V)$ ) for the space of complex-valued  $C^\infty$  functions (resp.  $C^\infty$  functions of compact support) on  $V$ . Composition of differential operators  $D_1, D_2$  is denoted  $D_1 \circ D_2$ .

### 2. The transversal part of a differential operator

Let  $V$  be a manifold satisfying the second axiom of countability, and  $H$  a Lie transformation group of  $V$ . If  $h \in H$ ,  $v \in V$ , let  $h \cdot v$  denote the image of  $v$  under  $H$  and let  $H^v$  denote the isotropy subgroup of  $H$  at  $v$ . Let  $\mathfrak{h}$  denote the Lie algebra of  $H$ . If  $X \in \mathfrak{h}$ , let  $X^+$  denote the vector field on  $V$  induced by  $X$ , i.e.,

---

Received November 3, 1971. Supported in part by National Science Foundation Grant GP-22928.

$$(2.1) \quad (X^+f)(v) = \left\{ \frac{d}{dt} f(\exp tX \cdot v) \right\}_{t=0}, \quad f \in \mathcal{D}(V), \quad v \in V.$$

A  $C^\infty$  function  $f$  on an open subset of  $V$  is said to be *locally invariant* if  $X^+f = 0$ .

**Lemma 2.1.** *Suppose  $W \subset V$  is a submanifold such that for each  $w \in W$  the tangent spaces at  $w$  satisfy the condition:*

$$(2.2) \quad V_w = W_w + (H \cdot w)_w \quad (\text{direct sum}).$$

Let  $w_0 \in W$ . Then there exists an open relatively compact neighborhood  $W_0$  of  $w_0$  in  $W$  and a relatively compact submanifold  $B \subset H$ ,  $e \in B$  such that the natural projection  $\pi: H \rightarrow H/H^{w_0}$  is a diffeomorphism of  $B$  onto an open neighborhood  $U_0$  of  $\pi(e)$  in  $H/H^{w_0}$  and such that the mapping  $\eta: (b, w) \rightarrow b \cdot w$  is a diffeomorphism of  $B \times W_0$  onto an open neighborhood  $V_0$  of  $w_0$  in  $V$ .

*Proof.* Let  $\mathfrak{h}^0$  denote the Lie algebra of  $H^{w_0}$ , and  $\mathfrak{n} \subset \mathfrak{h}$  any subspace complementary to  $\mathfrak{h}^0$ . Then the mapping  $\varphi: (X, w) \rightarrow \exp X \cdot w$  of  $\mathfrak{n} \times W$  into  $V$  is regular at  $(0, w_0)$ . In fact, since  $(d\varphi)_{(0, w_0)}$  fixes  $W_{w_0}$ , it suffices to prove

$$(2.3) \quad (d\varphi)_{(0, w_0)}(\mathfrak{n} \times 0) = (H \cdot w_0)_{w_0}.$$

This however is clear from dimensionality considerations. Now the lemma follows from the standard fact that if  $\mathfrak{n}_0$  is a sufficiently small neighborhood of 0 in  $\mathfrak{n}$ , then  $\exp$  is a diffeomorphism of  $\mathfrak{n}_0$  onto a submanifold  $B \subset H$  diffeomorphic under  $\pi$  to an open neighborhood of  $w_0$  in  $H/H^{w_0}$ .

It was pointed out to me by R. Palais that the local integration of involutive distributions (Chevalley [3, p. 89]) shows that a submanifold  $W$  satisfying (2.2) always exists.

Now let us assume that  $V$  has a Riemannian structure  $g$  invariant under the action of  $H$ . Assuming furthermore that all the orbits of  $H$  have the same dimension, we shall with each differential operator  $D$  on  $V$  associate a new differential operator  $D_T$  on  $V$  which acts "transversally to the orbits".

Fix  $s_0 \in V$  and let  $S$  denote the orbit  $H \cdot s_0$ . For each  $s \in S$  consider the geodesics in  $V$  starting at  $s$ , perpendicular to  $S$ . If we take sufficiently short pieces of these geodesics, their union is a submanifold  $S_s^\perp$  of  $V$ . Shrinking  $S_s^\perp$  if necessary we may assume that it satisfies transversality condition (2.2) for  $W$ . Take  $w_0$  as  $s_0$ , and let  $W_0$ ,  $B$  and  $V_0$  be as in the lemma. For  $f \in \mathcal{E}(V)$  (or even for functions defined on  $V_0$ ) we define a new function  $f_{s_0}$  on  $V_0$  by

$$f_{s_0}(b \cdot w) = f(w), \quad b \in B, \quad w \in W_0.$$

We then define  $D_T$  by

$$(2.4) \quad (D_T f)(s_0) = (Df_{s_0})(s_0), \quad s_0 \in V.$$

Since  $B \cdot w$  is a neighborhood of  $w$  in the orbit  $H \cdot w$ , and since  $D$  decreases supports, the choice of  $B$  above is immaterial, and (2.4) is indeed a valid definition; the operator  $D_T$  decreases supports and is therefore a differential operator, which we call *the transversal part of  $D$* .

**Theorem 2.2.** *Let  $V$  be a Riemannian manifold,  $H$  a Lie transformation group of isometries of  $V$ , all orbits assumed to have the same dimension. Let  $S$  be any  $H$ -orbit and let  $\bar{f}$  denote restriction of a function  $f$  to  $S$ . Then the Laplace-Beltrami operators  $L = L_V$  and  $L_S$  on  $V$  and  $S$ , respectively, satisfy*

$$(2.5) \quad (Lf)^- = L_S \bar{f} + (L_T f)^- \quad f \in \mathcal{E}(V) .$$

*Proof.* Let  $(y_1, \dots, y_r)$  be any coordinate system on  $B$  such that  $y_1(e) = \dots = y_r(e) = 0$ , and let  $w \rightarrow (z_{r+1}(w), \dots, z_n(w))$  be a coordinate system on  $W_0$  such that the geodesics forming  $S_{s_0}^\perp$  correspond to the straight lines through 0. Then we define a coordinate system  $(x_1, \dots, x_n)$  on  $V_0$  by

$$\begin{aligned} &(x_1(b \cdot w), \dots, x_r(b \cdot w), x_{r+1}(b \cdot w), \dots, x_n(b \cdot w)) \\ &= (y_1(b), \dots, y_r(b), z_{r+1}(w), \dots, z_n(w)) . \end{aligned}$$

The Laplace-Beltrami operator is given by

$$L = \sum_{p,q=1}^n g^{pq} (\partial_{pq} - \sum_t I_{pq}^t \partial_t) ,$$

where  $\partial_p = \partial/\partial x_p$ ,  $\partial_{pq} = \partial^2/\partial x_p \partial x_q$ ,  $g^{pq}$  is the inverse of the matrix  $g_{pq} = g(\partial_p, \partial_q)$ , and  $I_{pq}^t$  is the Christoffel symbol

$$I_{pq}^r = \frac{1}{2} \sum_s g^{rs} (\partial_q g_{ps} + \partial_p g_{qs} - \partial_s g_{pq}) .$$

Suppose  $\psi \in \mathcal{E}(V_0)$  satisfies the condition

$$(2.6) \quad \psi(x_1, \dots, x_n) \equiv \psi(0, \dots, 0, x_{r+1}, \dots, x_n) ,$$

or equivalently

$$\psi(b \cdot w) = \psi(w) , \quad b \in B , \quad w \in W_0 .$$

Then

$$(2.7) \quad \psi = \psi_{s_0} , \quad (L\psi)(s_0) = (L_T \psi)(s_0) .$$

On the other hand, suppose  $\varphi \in \mathcal{E}(V_0)$  satisfies

$$(2.8) \quad \varphi(x_1, \dots, x_n) \equiv \varphi(x_1, \dots, x_r, 0, \dots, 0) ,$$

or equivalently

$$\varphi(b \cdot w) = \varphi(b \cdot s_0), \quad b \in B, \quad w \in W_0.$$

For each set of real numbers  $a_{r+1}, \dots, a_n$ , not all 0, the curve

$$t \rightarrow (x_1(s_0), \dots, x_r(s_0), a_{r+1}t, \dots, a_n t)$$

is a geodesic in  $V$ . The differential equation for geodesics

$$\ddot{x}_i + \sum_{p,q} \Gamma_{pq}^i \dot{x}_p \dot{x}_q = 0$$

(dot denoting differentiation with respect to  $t$ ) therefore shows that

$$\Gamma_{\alpha\beta}^i(s_0) = 0, \quad 1 \leq i \leq n, \quad r+1 \leq \alpha, \beta \leq n.$$

Since the geodesic is perpendicular to  $S$  at  $s_0$ ,

$$(2.9) \quad g_{i\alpha}(s_0) = g^{i\alpha}(s_0) = 0, \quad \text{for } 1 \leq i \leq r, \quad r+1 \leq \alpha \leq n.$$

It follows that

$$(L\varphi)(s_0) = \sum_{1 \leq i, j \leq r} g^{ij}(\partial_{i,j}\varphi - \sum_{1 \leq k \leq r} \Gamma_{ij}^k \partial_k \varphi)(s_0).$$

But by (2.9),  $\Gamma_{ij}^k(s_0)$  is the same for  $S$  and for  $V$ , so

$$(2.10) \quad (L\varphi)(s_0) = (L_S \bar{\varphi})(s_0).$$

But

$$L(\varphi\psi) = \varphi L\psi + 2g(\text{grad } \varphi, \text{grad } \psi) + \psi L\varphi,$$

where for any  $f \in \mathcal{E}(V_0)$ ,

$$\text{grad } f = \sum_{p,q} g^{pq}(\partial_p f) \partial_q.$$

Hence (2.9) implies

$$(2.11) \quad L(\varphi\psi)(s_0) = \varphi(s_0)(L\psi)(s_0) + \psi(s_0)(L\varphi)(s_0).$$

But  $\varphi_{s_0}$  is a constant function, so by (2.4) and (2.7)

$$\varphi_{s_0}(L\psi)(s_0) = L((\varphi\psi)_{s_0})(s_0) = (L_T(\varphi\psi))(s_0).$$

Similarly, since  $\bar{\psi}$  is a constant function, (2.10) implies

$$\psi(s_0)(L\varphi)(s_0) = L_S(\bar{\varphi}\bar{\psi})(s_0).$$

This gives formula (2.5) for the function  $f = \varphi\psi$ , and since the linear combinations of such products form a dense subspace of  $\mathcal{D}(V_0)$  the theorem follows by approximation.

**Remark.** The theorem remains true with the same proof if  $V$  is a manifold with a pseudo-Riemannian structure  $g$  provided  $g$  is nonsingular on  $S$ .

### 3. The radial part of a differential operator

Again let  $V$  be a manifold satisfying the second axiom of countability, and  $H$  a Lie transformation group of  $V$ . Suppose  $W \subset V$  is a submanifold satisfying transversality condition (2.2) in Lemma 2.1.

**Lemma 3.1.** *Let  $D$  be a differential operator on  $V$ . Then there exists a unique differential operator  $\Delta(D)$  on  $W$  such that*

$$(3.1) \quad (Df)^- = \Delta(D)\bar{f}$$

for each locally invariant function  $f$  on an open subset of  $V$ , the bar denoting restriction to  $W$ .

*Proof.* Let  $w_0 \in W$  and select  $W_0, B$  and  $V_0$  as in Lemma 2.1. If  $\varphi \in \mathcal{E}(W_0)$ , we define  $f$  on  $V_0$  by

$$f(b \cdot w) = \varphi(w), \quad b \in B, \quad w \in W_0.$$

The mapping  $\varphi \rightarrow (Df)^-$  gives an operator  $D_{w_0, W_0, B}$  of  $\mathcal{E}(W_0)$  into itself. It is now an easy matter to verify that the linear transformation  $\Delta(D)$  given by

$$(\Delta(D)\psi)(w_0) = (D_{w_0, W_0, B}\psi)(w_0)$$

is a well-defined differential operator on  $\mathcal{E}(W)$ , with the properties stated in the lemma.

The operator  $\Delta(D)$  is called the *radial part* of  $D$ . We shall now give a formula for the radial part of the Laplace-Beltrami operator on  $V$  under a strengthening of transversality assumption (2.2); in fact we assume that each  $H$ -orbit intersects  $W$  just once and orthogonally.

**Theorem 3.2.** *Suppose  $V$  is a Riemannian manifold,  $H$  a closed unimodular subgroup of the Lie group of all isometries of  $V$  (with the compact open topology). Let  $W \subset V$  be a submanifold satisfying the condition: For each  $w \in W$ ,*

$$(3.2) \quad (H \cdot w) \cap W = \{w\}; \quad V_w = (H \cdot w)_w \oplus W_w,$$

where  $\oplus$  denotes orthogonal direct sum. Let  $L_V$  and  $L_W$  denote the Laplace-Beltrami operators on  $V$  and  $W$ , respectively. Then

$$(3.3) \quad \Delta(L_V) = \delta^{-\frac{1}{2}}L_W \circ \delta^{\frac{1}{2}} - \delta^{-\frac{1}{2}}L_W(\delta^{\frac{1}{2}}),$$

where the function  $\delta$  is the volume element ratio in (3.8) below.

*Proof.* Let  $V^*$  denote the subset  $H \cdot W$  of  $V$ . Since the mapping  $(h, w) \rightarrow h \cdot w$  of  $H \times W$  into  $V$  has (by (3.2)) a surjective differential at each point,  $V^*$  is an open subset of  $V$ . Since  $H$  is closed, the isotropy subgroup  $H^w$  at each

point  $w \in W$  is compact and the orbit  $H \cdot w$  is closed; if we fix a left invariant Haar measure on  $H$  and a Haar measure on  $H^w$  (with total measure 1), we obtain in a standard way an  $H$ -invariant measure  $d\hat{h}$  on each orbit  $H \cdot w = H/H^w$ . Denoting by  $dv$  and  $dw$  the Riemannian measures on  $V$  and  $W$ , respectively, we shall prove that there exists a function  $\delta \in \mathcal{E}(W)$  such that

$$(3.4) \quad \int_{V^*} F(v)dv = \int_W \delta(w) \left( \int_{H \cdot w} F(h \cdot w) d\hat{h} \right) dw, \quad F \in \mathcal{D}(V^*).$$

Let  $w_0 \in W$ . Because of the second part of (3.2) there exist a coordinate neighborhood  $W_0$  of  $w_0$  in  $W$ , a vector subspace  $\mathfrak{m} \subset \mathfrak{h}$  of dimension  $\dim V - \dim W$  and a neighborhood  $\mathfrak{m}_0$  of 0 in  $\mathfrak{m}$  such that the map

$$\eta: (X, w) \rightarrow \exp X \cdot w$$

is a diffeomorphism of  $\mathfrak{m}_0 \times W_0$  onto an open neighborhood  $V_0$  of  $w_0$  in  $V$ . Let  $(x_1, \dots, x_r)$  be a Cartesian coordinate system on  $\mathfrak{m}$ , and  $(x_{r+1}, \dots, x_n)$  an arbitrary coordinate system on  $W_0$ . In the formulas below let  $1 \leq i, j \leq r$ ,  $r+1 \leq \alpha, \beta \leq n$ . Let the coordinate system  $(x_1, \dots, x_n)$  on  $V_0$  be determined by

$$x_i(\exp X \cdot w) = x_i(X), \quad x_\alpha(\exp X \cdot w) = x_\alpha(w).$$

Let  $g$  denote the Riemannian structure of  $V$ , and put  $g_{pq} = g(\partial_p, \partial_q)$  as usual, so that

$$dv = \bar{g}^{\frac{1}{2}} dx_1 \cdots dx_n, \quad dw = \bar{\gamma}^{\frac{1}{2}} dx_{r+1} \cdots dx_n,$$

where

$$(3.5) \quad \bar{g} = |\det((g_{pq})_{1 \leq p, q \leq n})|, \quad \bar{\gamma} = |\det(g_{\alpha\beta})|.$$

Because of the orthogonality in (3.2) we have

$$(3.6) \quad g_{i\alpha}(w) = 0, \quad w \in W_0.$$

But if  $h = \exp X$  ( $X \in \mathfrak{m}_0$ ) then our choice of coordinates implies for the differential  $dh$ ,

$$dh \left( \frac{\partial}{\partial x_\alpha} \right)_w = \left( \frac{\partial}{\partial x_\alpha} \right)_{h \cdot w}, \quad dh \left( \frac{\partial}{\partial x_i} \right)_w = \sum_{j=1}^r a_{ij} \left( \frac{\partial}{\partial x_j} \right)_{h \cdot w},$$

where  $a_{ij} \in \mathbf{R}$ . Hence  $g_{\alpha\beta}(h \cdot w) = g_{\alpha\beta}(w)$  and using (6),  $g_{i\alpha}(h \cdot w) = 0$ ; consequently

$$(3.7) \quad \bar{g}(h \cdot w) = \det(g_{ij})(h \cdot w) \bar{\gamma}(w).$$

However

$$|\{\det(g_{ij})\}^{\frac{1}{2}}(h \cdot w)| dx_1 \cdots dx_r(h \cdot w)$$

is just the Riemannian volume element  $d\sigma_w$  on the orbit  $H \cdot w$ . Thus, if  $F \in \mathcal{D}(V_0)$  we obtain from the Fubini theorem and (3.7) that

$$\int_V F(v) dv = \int_W \bar{\gamma}^{\frac{1}{2}}(w) \left( \int_{H \cdot w} F(p) d\sigma_w(p) \right) dx_{r+1} \cdots dx_n(w).$$

But  $d\sigma_w$  is invariant under  $H$ , so it must be a scalar multiple of  $d\dot{h}$ ,

$$(3.8) \quad d\sigma_w = \delta(w) d\dot{h}.$$

This proves (3.4) for all  $F \in \mathcal{D}(V_0)$ ; then it holds also if  $F$  has support inside  $h \cdot V_0$  for some  $h \in H$ . But as  $w_0$  runs through  $W$ , the sets  $h \cdot V_0$  form a covering of  $V^*$ . Passing to a locally finite refinement and a corresponding partition of unity, (3.4) follows for all  $F \in \mathcal{D}(V^*)$ .

Let  $\hat{F}(w)$  denote the inner integral in (3.4), so that

$$(3.9) \quad \hat{F}(w) = \int_{H \cdot w} F(h \cdot w) d\dot{h}.$$

It is a routine matter to verify that the mapping  $F \rightarrow \hat{F}$  is surjective, i.e.,

$$(3.10) \quad \mathcal{D}(V^*) \cdot = \mathcal{D}(W).$$

For the determination of  $\Delta(L_V)$  we first observe that

$$(3.11) \quad \Delta(L_V) = L_W + \text{lower order terms.}$$

This is clear from the coordinate expression for  $L_V$  together with (3.6) if we also note that the vector fields  $\partial/\partial x_i$  are tangential to the  $H$ -orbits. Next we recall that  $L_V$  is symmetric with respect to  $dv$ , i.e.,

$$(3.12) \quad \int_V (L_V f_1)(v) f_2(v) dv = \int_V f_1(v) (L_V f_2)(v) dv$$

for all  $f_1, f_2 \in \mathcal{D}(V^*)$ . But then this relation holds for all  $f_2 \in \mathcal{E}(V^*)$ . In particular we can use it on  $f_2$  invariant under  $H$ . Applying (3.4) to the left hand side of (3.12) we obtain

$$(3.13) \quad \int_W \delta(w) f_2(w) \left( \int_{H \cdot w} (L_V f_1)(h \cdot w) d\dot{h} \right) dw.$$

But for each  $v \in V$  the isotropy subgroup  $H^v$  is compact, so by invariance of  $L_V$

$$(L_V)_v \left( \int_{H^v} f_1(h \cdot v) dh \right) = \int_{H^v} (L_V f_1)(h \cdot v) dh.$$

Now putting here  $v = w$  we get the inner integral in (3.13) equal to  $(\Delta(L_V)f_1)(w)$ ; thus the left hand side of (3.12) is

$$\int_W (\Delta(L_V)f_1)(w)\bar{f}_2(w)\delta(w)dw ,$$

the bar denoting restriction to  $W$ . But using the  $H$ -invariance of  $L_Vf_2$ , formula (3.4) and the definition of radial part, the right hand side of (3.12) reduces to

$$\int_W f_1(w)(\Delta(L_V)\bar{f}_2)(w)\delta(w)dw .$$

But in view of (3.10) the functions  $f_1$  (and of course the  $\bar{f}_2$ ) fill up  $\mathcal{D}(W)$ , so the equality of the two last expressions implies that  $\Delta(L_V)$  is symmetric with respect to  $\delta(w)dw$ . Now since  $L_W$  is symmetric with respect to  $dw$ , a simple computation shows that the composition  $\delta^{-\frac{1}{2}}L_W \circ \delta^{\frac{1}{2}}$  is symmetric with respect to  $\delta(w)dw$  and it clearly agrees with  $L_W$  up to lower order terms. Thus by (3.11) the symmetric operators  $\Delta(L_V)$  and  $\delta^{-\frac{1}{2}}L_W \circ \delta^{\frac{1}{2}}$  agree up to an operator of order  $\leq 1$ . But this operator, being symmetric, must be a function, and now (3.3) follows by applying the operators to the function 1.

It is of interest to generalize Theorem 3.2 to pseudo-Riemannian manifolds  $V$ . If  $V$  has a pseudo-Riemannian structure  $g$ , which for each  $w \in W$  is non-degenerate on the closed orbit  $H \cdot w$ , and if each  $H^w$  ( $w \in W$ ) is compact, then Theorem 3.2 remains valid. In fact, the isotropy group  $H^v$  is then compact for each  $v \in V^*$ , so no change is necessary in the proof.

When a semisimple Lie group  $H$  acts on its Lie algebra by the adjoint representation, the regular elements of a Cartan subalgebra constitute a transversal submanifold  $W$  where the isotropy subgroup  $H^w$  is the same for all  $w \in W$ . This then provides an example for the following variation of Theorem 3.2.

**Theorem 3.3.** *Let the assumptions be as in Theorem 3.2 except that  $V$  has only a pseudo-Riemannian structure  $g$ . Then formula (3.3) remains valid if we further assume that*

- (i) *for each  $w \in W$  the orbit  $H \cdot w$  is closed and  $g$  is non-degenerate on it,*
- (ii)  *$H^w$  is the same for all  $w \in W$ , and its Lie algebra is its own normalizer in the Lie algebra of  $H$ .*

*Proof.* Put  $H^0 = H^w$  ( $w \in W$ ) and  $\dot{h} = hH^0$ , and fix an  $H$ -invariant measure  $d\dot{h}$  on the coset space  $H/H^0$ . Such a measure exists since each orbit  $H \cdot w$  has an  $H$ -invariant measure  $d\sigma_w$  defined as above. If  $\gamma$  is a geodesic in  $V$  tangential to  $W$  at  $w$  then  $\gamma$  is left fixed by each  $h \in H_0$ . Thus (ii) implies  $\gamma \subset W$  so  $W$  is a totally geodesic submanifold of  $V$ . Defining  $\delta$  by (3.8) the only part of the proof above which requires change is the justification of the formula

$$(3.14) \quad \int_{H \cdot w} (L_V f_1)(h \cdot w)d\dot{h} = (\Delta(L_V)f_1)(w) .$$



For this we use Theorem 2.2 and the subsequent remark to split  $L_V$  into its "orbital part" and transversal part. The orbital part gives integral 0 over  $H \cdot w$ , so in the integral (3.14) we can replace  $L_V$  by its transversal part  $L_{V,T}$ . Putting  $f_1^h(w) = f_1(h \cdot w)$  for  $h \in H$ , we have, by the  $H$ -invariance of  $L_{V,T}$ ,

$$(L_{V,T}f_1)(h \cdot w) = (L_{V,T}(f_1^h))(w),$$

which, by the definition of transversal part and radial part, equals  $\Delta(L_V)(\overline{f_1^h})(w)$ ,  $W$  being totally geodesic. But then the left hand side of (3.14) equals

$$\int_{H/H^0} \Delta(L_V)_w(f_1(\dot{h} \cdot w)) d\dot{h},$$

which equals  $(\Delta(L_V)f_1)(w)$  because now  $\dot{h}$  and  $w$  are independent variables. This proves (3.14) and therefore also Theorem 3.3.

### References

- [1] F. A. Berezin, *Laplace operators on semisimple Lie groups*, Trudy Moskov. Mat. Obšč. **6** (1957) 371–463.
- [2] C. Chevalley, *Theory of Lie groups*. I, Princeton University Press, Princeton, 1946.
- [3] Harish-Chandra, *The characters of semisimple Lie groups*, Trans. Amer. Math. Soc. **83** (1956) 98–163.
- [4] —, *Differential operators on a semisimple Lie algebra*, Amer. J. Math. **79** (1957) 87–120.
- [5] —, *Spherical functions on a semisimple Lie group*. I, Amer. J. Math. **80** (1958) 241–310.
- [6] S. Helgason, *Differential operators on homogeneous spaces*, Acta Math. **102** (1959) 239–299.
- [7] —, *Radon-Fourier transforms on symmetric spaces and related group representations*, Bull. Amer. Math. Soc. **71** (1965) 757–763.
- [8] F. I. Karpelevič, *The Geometry of geodesics and the eigenfunctions of the Beltrami-Laplace operator on symmetric spaces*, Trudy Moskov. Mat. Obšč. **14** (1963) 48–185.
- [9] P. D. Methéé, *Sur les distributions invariantes dans le groupe des rotations de Lorentz*, Comment. Math. Helv. **28** (1954) 1–49.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY