

ALMOST HERMITIAN MANIFOLDS WITH CONSTANT HOLOMORPHIC SECTIONAL CURVATURE

SUMIO SAWAKI & KOUEI SEKIGAWA

1. Introduction

B. Smyth proved in his thesis [6] the following

Theorem. *Let M be a complex hypersurface of a Kählerian manifold \tilde{M} of constant holomorphic sectional curvature \tilde{c} . If M is of complex dimension ≥ 2 , the following statements are equivalent:*

- (i) M is totally geodesic in \tilde{M} ,
- (ii) M is of constant holomorphic sectional curvature,
- (iii) M is an Einstein manifold and at one point of M all sectional curvatures of M are $\geq \frac{1}{4}\tilde{c}$ (resp. $\leq \frac{1}{4}\tilde{c}$) when $\tilde{c} \geq 0$ (resp. ≤ 0).

One of the purposes of the present paper is to generalize this theorem to almost Hermitian manifolds, and another is to prove that an F -space of constant holomorphic sectional curvature is Kählerian. Here by an F -space we mean an almost Hermitian manifold M satisfying $R(X, Y) \cdot F = 0$ for any vector fields X and Y on M , where the endomorphism $R(X, Y)$ operates on the almost complex structure tensor F as a derivation at each point of M .

In § 2, we shall state the differential-geometric properties of a complex hypersurface of an almost Hermitian manifold satisfying a certain condition and a generalization of the equivalence of the first two statements of Smyth's result. We proceed in § 3 to study the same properties of $*O$ -spaces and K -spaces, and to state a generalization of the result of Smyth. In § 4 we shall prove some theorems for F -spaces of constant holomorphic sectional curvature. In §§ 2 and 3, by a complex hypersurface we mean a connected almost complex hypersurface.

2. Complex hypersurfaces of an almost Hermitian manifold

Let \tilde{M} be an almost Hermitian manifold of complex dimension $n + 1$, and denote the almost complex structure and the Hermitian metric of \tilde{M} by F and g respectively. Moreover, let M be a complex hypersurface of \tilde{M} , i.e., suppose that there exists a complex analytic mapping $f: M \rightarrow \tilde{M}$. Then for each $x \in M$ we identify the tangent space $T_x(M)$ with $f_*(T_x(M)) \subset T_{f(x)}(\tilde{M})$ by means of f_* . Since $f^* \circ g = g'$ and $F \circ f_* = f_* \circ F'$ where g' and F' are the Hermitian

metric and the almost complex structure of M respectively, g' and F' are respectively identified with the restrictions of the structures g and F to the subspace $f_*(T_x(M))$.

As is well known, we can choose the following special neighborhood $U(x)$ of x for a neighborhood $\tilde{U}(f(x))$ of $f(x)$. Let $\{\tilde{U}; \tilde{x}^i\}$ ($i = 1, \dots, 2n + 2$) be a system of coordinate neighborhoods of \tilde{M} . Then $\{U; x^i\}$ is a system of coordinate neighborhoods of M such that $x^{2n+1} = x^{2n+2} = 0$ where $x^i = \tilde{x}^i \circ f$.

By $\tilde{\nabla}$ we always mean the Riemannian covariant differentiation on \tilde{M} and by ξ a differentiable unit vector field normal to M at each point of $U(x)$.

If X and Y are vector fields on the neighborhood $U(x)$, we may write

$$(2.1) \quad \tilde{\nabla}_X Y = \nabla_X Y + h(X, Y)\xi + k(X, Y)F\xi,$$

where $\nabla_X Y$ denotes the component of $\tilde{\nabla}_X Y$ tangent to M .

Lemma 2.1. (i) ∇ is the covariant differentiation of the almost Hermitian manifold M .

(ii) h and k are symmetric covariant tensor fields of degree 2 on $U(x)$.

Proof. Making use of (2.1), we have

$$\begin{aligned} \tilde{\nabla}_{f_1 X}(f_2 Y) &= f_1 \tilde{\nabla}_X(f_2 Y) = f_1(Xf_2)Y + f_1 f_2 \tilde{\nabla}_X Y \\ &= f_1(Xf_2)Y + f_1 f_2 \nabla_X Y + f_1 f_2 h(X, Y)\xi + f_1 f_2 k(X, Y)F\xi, \\ \tilde{\nabla}_{f_1 X}(f_2 Y) &= \nabla_{f_1 X}(f_2 Y) + h(f_1 X, f_2 Y)\xi + k(f_1 X, f_2 Y)F\xi, \end{aligned}$$

where X and Y are vector fields on $U(x)$, and f_1 and f_2 are differentiable functions on $U(x)$. From the above two equations, we have

$$h(f_1 X, f_2 Y) = f_1 f_2 h(X, Y), \quad k(f_1 X, f_2 Y) = f_1 f_2 k(X, Y),$$

which show that h and k are tensor fields on $U(x)$.

Thus, since $\nabla_X Y$ becomes a vector fields, from (2.1) it follows that ∇ is a covariant differentiation on $U(x)$.

Next, from

$$\begin{aligned} \tilde{\nabla}_X Y &= \nabla_X Y + h(X, Y)\xi + k(X, Y)F\xi, \\ \tilde{\nabla}_Y X &= \nabla_Y X + h(Y, X)\xi + k(Y, X)F\xi, \\ [X, Y]_{\tilde{M}} &= [X, Y]_M, \end{aligned}$$

we have

$$\tilde{T}(X, Y) = T(X, Y) + \{h(X, Y) - h(Y, X)\}\xi + \{k(X, Y) - k(Y, X)\}F\xi,$$

where \tilde{T} (resp. T) is the torsion tensor of the connection on \tilde{M} (resp. $U(x)$) with respect to $\tilde{\nabla}$ (resp. ∇). Since $\tilde{T} = 0$, it follows that $T = 0$ and h and k are symmetric.

From $\tilde{\nabla}g = 0$ we have easily $\nabla g = 0$. Hence the proof is completed.

The identities $g(\xi, \xi) = 1$ and $g(F\xi, F\xi) = 1$ imply $g(\tilde{\nabla}_X \xi, \xi) = 0$ and $g(\tilde{\nabla}_X(F\xi), F\xi) = 0$ respectively. Therefore we may put

$$(2.2) \quad \tilde{\nabla}_X \xi = -A(X) + s(X)F\xi ,$$

$$(2.3) \quad \tilde{\nabla}_X(F\xi) = -B(X) + t(X)\xi ,$$

where $A(X)$ and $B(X)$ are tangent to M .

Lemma 2.2. (i) A, B and s, t are tensor fields on $U(x)$ of type (1.1) and (0,1) respectively.

(ii) A and B are symmetric with respect to g , and satisfy

$$(2.4) \quad h(X, Y) = g(AX, Y) ,$$

$$(2.5) \quad k(X, Y) = g(BX, Y)$$

for any vector fields X and Y .

Proof. For any vector field X and any differentiable function f on $U(x)$, we have

$$f\tilde{\nabla}_X \xi = \tilde{\nabla}_{fX} \xi = -A(fX) + s(fX)F\xi = -fA(X) + fs(X)F\xi ,$$

from which it follows that $A(fX) = fA(X)$, $s(fX) = fs(X)$. Thus A and s are tensor fields on $U(x)$. For ξ and any vector field Y on $U(x)$, we have $g(Y, \xi) = 0$ and therefore

$$g(\tilde{\nabla}_X Y, \xi) + g(Y, \tilde{\nabla}_X \xi) = 0 ,$$

in which substitution of (2.1) and (2.2) gives (2.4). However, since h is symmetric, from (2.4) it follows that $g(AX, Y) = g(X, AY)$ which shows that A is symmetric. Similarly the properties of B are verified.

Now let M be a complex hypersurface satisfying the condition

$$(2.6) \quad h(X, Y) = k(X, FY)$$

for any vector fields X and Y on $U(x)$ at every point $x \in M$. It is easily verified that the condition (2.6) is independent of the choice of mutually orthogonal unit vectors ξ and $F\xi$ normal to M .

Lemma 2.3. In a complex hypersurface M of \tilde{M} satisfying (2.6), we have

$$(i) \quad FA = -AF , \quad FB = -BF ,$$

$$(ii) \quad FA \text{ and } FB \text{ are symmetric with respect to } g,$$

$$(iii) \quad B = FA .$$

Proof. By virtue of (2.4) and (2.6), for any vector fields X and Y we have

Applying \tilde{V}_x to this equation and making use of (2.2) and (2.3), we obtain

$$(2.10) \quad \begin{aligned} \tilde{V}_x \tilde{V}_Y W &= \nabla_x \nabla_Y W - h(Y, W)A(X) - k(Y, W)B(X) \\ &+ \{h(X, \nabla_Y W) + X(h(Y, W)) + k(Y, W)t(X)\}\xi \\ &+ \{k(X, \nabla_Y W) + X(k(Y, W)) + h(Y, W)s(X)\}F\xi, \\ \tilde{V}_{[X, Y]} W &= \nabla_{[X, Y]} W + h([X, Y], W)\xi + k([X, Y], W)F\xi. \end{aligned}$$

Substitution of (2.10) in

$$\begin{aligned} &\tilde{R}(X, Y)W - R(X, Y)W \\ &= \tilde{V}_x \tilde{V}_Y W - \tilde{V}_Y \tilde{V}_x W - \tilde{V}_{[X, Y]} W - (\nabla_x \nabla_Y W - \nabla_Y \nabla_x W - \nabla_{[X, Y]} W) \end{aligned}$$

gives easily

$$\begin{aligned} \tilde{R}(X, Y, Z, W) &= R(X, Y, Z, W) - \{g(AX, Z)h(Y, W) - g(AY, Z)h(X, W)\} \\ &- \{g(BX, Z)k(Y, W) - g(BY, Z)k(X, W)\}, \end{aligned}$$

or (2.9) by (2.4), (2.5) and (2.6).

Lemma 2.6. *Let M be a complex hypersurface of \tilde{M} and satisfy the condition (2.6).*

(i) *If p is 2-plane tangent to M at a point of $U(x)$, then*

$$(2.11) \quad \begin{aligned} \tilde{K}(p) &= K(p) - \{g(AX, X)g(AY, Y) - g(AX, Y)^2\} \\ &- \{g(FAX, X)g(FAY, Y) - g(FAX, Y)^2\}, \end{aligned}$$

where X, Y form an orthonormal basis of p , and $\tilde{K}(p)$ (resp. $K(p)$) is the sectional curvature of p considered as a 2-plane tangent to \tilde{M} (resp. M).

(ii) *If X is a unit vector tangent to M at a point of $U(x)$, then*

$$(2.12) \quad \tilde{H}(X) = H(X) + 2\{g(AX, X)^2 + g(FAX, X)^2\},$$

where $\tilde{H}(X)$ (resp. $H(X)$) is the holomorphic sectional curvature in \tilde{M} (resp. M).

Proof. (i) is immediate on replacing Z and W in the Gauss equation by X and Y respectively, and making use of the fact that A and FA are symmetric. (ii) is also immediate on replacing Y by FX in (2.11) and making use of the fact that $FA = -AF$.

Proposition 2.7. *Let M be a complex hypersurface of \tilde{M} of constant holomorphic sectional curvature \tilde{c} . If M is of complex dimension ≥ 2 and satisfies the condition (2.6), then at each point of M there exists a holomorphic plane whose sectional curvature in M is \tilde{c} , and therefore if M is of constant holomorphic sectional curvature c , then $c = \tilde{c}$.*

Proof. Let $\{e_1, \dots, e_n, Fe_1, \dots, Fe_n\}$ be an orthonormal basis in Lemma 2.4. Since $n \geq 2$, there exist λ_i and λ_j ($i \neq j$) defined in Lemma 2.4.

In the case where $\lambda_i > 0$ and $\lambda_j > 0$, we set

$$X = (\lambda_i + \lambda_j)^{-\frac{1}{2}}(\sqrt{\lambda_j}e_i + \sqrt{\lambda_i}Fe_j).$$

Then

$$AX = \frac{\sqrt{\lambda_j}\lambda_i e_i - \sqrt{\lambda_i}\lambda_j Fe_j}{(\lambda_i + \lambda_j)^{\frac{1}{2}}}, \quad FAX = \frac{\sqrt{\lambda_j}\lambda_i Fe_i + \sqrt{\lambda_i}\lambda_j e_j}{(\lambda_i + \lambda_j)^{\frac{1}{2}}},$$

so that

$$(2.13) \quad g(AX, X) = 0, \quad g(FAX, X) = 0.$$

In the case where $\lambda_i < 0$ and $\lambda_j > 0$, and in the case where $\lambda_i < 0$ and $\lambda_j < 0$, we set, respectively,

$$X = \frac{\sqrt{\lambda_j}e_i + \sqrt{-\lambda_i}e_j}{(\lambda_j - \lambda_i)^{\frac{1}{2}}}, \quad X = \frac{\sqrt{-\lambda_j}e_i + \sqrt{-\lambda_i}Fe_j}{(-\lambda_j - \lambda_i)^{\frac{1}{2}}},$$

so that we can also obtain (2.13).

Consequently, from (2.12) and (2.13) we have $\tilde{c} = H(\tilde{X}) = H(X)$ which proves the proposition.

Theorem 2.8. *Let M be a complex hypersurface of \tilde{M} of constant holomorphic sectional curvature \tilde{c} . If M is of complex dimension ≥ 2 and satisfies the condition (2.6), then the following statements are equivalent:*

- (i) M is totally geodesic in \tilde{M} ,
- (ii) M is of constant holomorphic sectional curvature.

Proof. If M is totally geodesic, then A vanishes on M , and therefore from (2.12) it follows that M is of constant holomorphic sectional curvature \tilde{c} . Conversely, if M is of constant holomorphic sectional curvature c , then by virtue of Proposition 2.7 we have, for any unit vector X tangent to M , $\tilde{c} = H(\tilde{X}) = H(X)$, which reduces (2.12) to $g(AX, X)^2 + g(FAX, X)^2 = 0$, so that $A = 0$, that is, M is totally geodesic.

3. *O-spaces and K-spaces

An almost Hermitian manifold \tilde{M} is called an *O-space (or quasi-Kählerian manifold) [3] or a K-space (or Tachibana space or nearly Kähler manifolds) [7] according as

$$(3.1) \quad \tilde{V}_X(F)Y + \tilde{V}_{FX}(F)FY = 0,$$

or

$$(3.2) \quad \tilde{V}_X(F)Y + \tilde{V}_Y(F)X = 0 \quad (\text{or equivalently } \tilde{V}_X(F)X = 0)$$

holds for any vector fields X and Y on \tilde{M} . It is well-known that a K-space is an *O-space.

First of all, let M be a complex hypersurface of an $*O$ -space \tilde{M} . Then for any vector fields X and Y on $U(x) \subset M$ we have

$$\tilde{V}_x(FY) = F\tilde{V}_xY + \tilde{V}_x(F)Y, \quad \tilde{V}_{FX}(FFY) = F\tilde{V}_{FX}(FY) + \tilde{V}_{FX}(F)FY.$$

Adding these equations and making use of (3.1) we obtain

$$(3.3) \quad \tilde{V}_x(FY) - \tilde{V}_{FX}Y = F(\tilde{V}_xY + \tilde{V}_{FX}(FY)).$$

Substituting (2.1) in (3.3) gives immediately

$$(3.4) \quad \nabla_x(FY) - \nabla_{FX}Y - F\nabla_xY - F\nabla_{FX}(FY) = 0,$$

$$(3.5) \quad h(X, FY) - h(FX, Y) = -k(X, Y) - k(FX, FY),$$

$$(3.6) \quad k(X, FY) - k(FX, Y) = h(X, Y) + h(FX, FY).$$

In consequence of

$$(3.7) \quad \nabla_x(F)FY = -F\nabla_x(F)Y,$$

(3.4) reduces to

$$\nabla_x(F)Y + \nabla_{FX}(F)FY = 0,$$

which shows that M is also an $*O$ -space.

Since the left hand side of (3.5) is skew-symmetric in X, Y and the right hand side is symmetric in X, Y due to the symmetry of h and k , we have

$$h(X, FY) = h(FX, Y), \quad k(X, Y) + k(FX, FY) = 0.$$

Similarly, from (3.6) follow

$$k(X, FY) = k(FX, Y), \quad h(X, Y) + h(FX, FY) = 0,$$

which are equivalent to the above two equations.

Hence we have

Lemma 3.1. *A complex hypersurface M of an $*O$ -space \tilde{M} is also an $*O$ -space, and satisfies*

$$(3.8) \quad h(X, FY) = h(FX, Y),$$

$$(3.9) \quad k(X, FY) = k(FX, Y).$$

Next, let M be a complex hypersurface of a K -space \tilde{M} . Then for vector fields X and Y on $U(x) \subset M$ we have

$$\tilde{V}_x(FY) = F\tilde{V}_xY + \tilde{V}_x(F)Y, \quad \tilde{V}_y(FY) = F\tilde{V}_yX + \tilde{V}_y(F)X.$$

Adding these equations and making use of (3.2) we obtain

$$(3.10) \quad \tilde{F}_x(FY) + \tilde{F}_Y(FX) = F(\tilde{F}_xY + \tilde{F}_YX) .$$

Substituting (2.1) in (3.10) gives readily

$$(3.11) \quad \nabla_x(FY) + \nabla_Y(FX) = F\nabla_xY + F\nabla_YX ,$$

$$(3.12) \quad h(X, FY) + h(FX, Y) = -2k(X, Y) ,$$

$$(3.13) \quad k(X, FY) + k(FX, Y) = -2h(X, Y) .$$

(3.11) reduces to

$$\nabla_x(F)Y + \nabla_Y(F)X = 0 ,$$

which shows that M is also a K -space. (3.12) and (3.8) imply $h(X, FY) = -k(X, Y)$, i.e., $h(X, Y) = k(X, FY)$, which is equivalent to $B = FA$ by the remark in § 2. From (3.13) we shall get the same result.

Consequently, we have

Lemma 3.2. *A complex hypersurface M of a K -space \tilde{M} is also a K -space, and satisfies*

$$h(X, Y) = k(X, FY) \quad (\text{or equivalently } B = FA) .$$

Recently, Gray [1] proved

Lemma 3.3. *In a K -space \tilde{M} of constant holomorphic sectional curvature \tilde{c} at a point $x \in \tilde{M}$, we have*

$$(3.15) \quad \tilde{K}(p) = \frac{1}{4}\tilde{c}\{1 + 3g(FX, Y)^2\} + \frac{3}{4}\|\tilde{F}_x(F)Y\|^2 ,$$

where p is a 2-plane spanned by any two orthonormal vectors $X, Y \in T_x(\tilde{M})$.

Making use of these Lemmas, we can prove

Theorem 3.4. *Let M be a complex hypersurface of a K -space \tilde{M} with constant holomorphic sectional curvature \tilde{c} . If M is of complex dimension ≥ 3 , then the following statements are equivalent:*

- (i) M is totally geodesic in \tilde{M} ,
- (ii) M is of constant holomorphic sectional curvature,
- (iii) at every point $x \in M$, all the sectional curvatures of M satisfy

$$(3.16) \quad K(p) \geq \frac{1}{4}\tilde{c}\{1 + 3g(FX, Y)^2\} ,$$

where p is a 2-plane spanned by any two orthonormal vectors $X, Y \in T_x(M)$.

Proof. Since, by Lemma 3.2, K -space satisfies (2.6), the fact that (i) is equivalent to (ii) is nothing but Theorem 2.8 (i). Next, if M is of constant holomorphic sectional curvature c , then $c = \tilde{c}$ by Proposition 2.7, and therefore by Lemma 3.3 we have, for any orthonormal vectors $X, Y \in T_x(M)$ at every point $x \in M$,

$$(3.17) \quad K(p) = \frac{1}{4}\tilde{c}\{1 + 3g(FX, Y)^2\} + \frac{3}{4}\|\tilde{V}_X(F)Y\|^2,$$

which implies (3.16).

Finally, we shall prove that (iii) implies (i). Substituting (3.15) in (2.11), and making use of (3.16) we can easily obtain

$$(3.18) \quad \frac{3}{4}\|\tilde{V}_X(F)Y\|^2 + \{g(AX, X)g(AY, Y) - g(AX, Y)^2\} \\ + \{g(FAX, X)g(FAY, Y) - g(FAX, Y)^2\} \geq 0.$$

Now let $\{e_1, \dots, e_n, Fe_1, \dots, Fe_n\}$ be an orthonormal basis given in Lemma 2.4, and set

$$X = (e_i + Fe_i)/\sqrt{2}, \quad Y = (e_i - Fe_i)/\sqrt{2}.$$

Since

$$AX = \lambda_i(e_i - Fe_i)/\sqrt{2}, \quad AY = \lambda_i(e_i + Fe_i)/\sqrt{2}, \\ FAX = \lambda_i(Fe_i + e_i)/\sqrt{2}, \quad FAY = \lambda_i(Fe_i - e_i)/\sqrt{2},$$

we have

$$g(AX, X) = 0, \quad g(FAX, X) = \lambda_i \\ g(FAY, Y) = -\lambda_i, \quad g(FAX, Y) = 0, \quad g(AX, Y) = \lambda_i.$$

Moreover, from $Y = -FX$, (3.2) and (3.7) we have

$$\tilde{V}_X(F)Y = -\tilde{V}_X(F)FX = F\tilde{V}_X(F)X = 0.$$

Thus (3.18) reduces to $\lambda_i = 0$ ($i = 1, \dots, n$), which together with Lemma 2.4 implies that A is identically zero at each point of M , so that M is totally geodesic in \tilde{M} .

Remark. It is well-known that in a K -space M of constant holomorphic sectional curvature \tilde{c} , $\tilde{c} > 0$ [8]. Hence from (3.17) we have

$$(3.19) \quad K(p) \geq \frac{1}{4}\tilde{c}.$$

However, the authors do not know whether M is totally geodesic or not if (3.19) holds.

4. F -spaces

Recall that an almost Hermitian manifold M of dimension $2n$ is called an F -space if $R(X, Y) \cdot F = 0$ holds for any vector fields X and Y on M . Of course, a Kählerian manifold is an F -space, and an almost Kählerian manifold or a K -space satisfying $R(X, Y) \cdot F = 0$ is Kählerian [5]. However, an example

of a nonkählerian $*O$ -space satisfying $R(X, Y) \cdot F = 0$ has been recently given by Yanamoto [9].

Now for an F -space M of constant holomorphic sectional curvature c we have (cf. [2, pp. 165–166])

$$(4.1) \quad \begin{aligned} R(X, Y, Z, W) = & \frac{1}{4}c\{g(X, Z)g(Y, W) - g(X, W)g(Y, Z) \\ & + g(X, FZ)g(Y, FW) - g(X, FW)g(Y, FZ) \\ & + 2g(X, FY)g(Z, FW)\}, \end{aligned}$$

where X, Y, Z and W are any tangent vectors at a point of M , since $R(X, Y) \cdot F = 0$ means that

$$R(X, Y, Z, W) = R(X, Y, FZ, FW) = R(FX, FY, Z, W).$$

On replacing Z and W in (4.1) by mutually orthogonal unit vectors X and Y respectively, we obtain

$$K(p) = \frac{1}{4}c\{1 + 3g(X, FY)^2\}.$$

Hence we have the following theorem which is a generalization of the corresponding result in a Kählerian manifold [10].

Theorem 4.1. *An F -space M of constant holomorphic sectional curvature c is an Einstein space. When $c \neq 0$, the sectional curvature $K(p)$ of a 2-plane p spanned by any two orthonormal vectors X and Y in M satisfies the inequalities:*

$$\frac{1}{4}c \leq K(p) \leq c \quad \text{for } c > 0, \quad \frac{1}{4}c \geq K(p) \geq c \quad \text{for } c < 0,$$

where the equality $\frac{1}{4}c = K(p)$ occurs when $g(X, FY) = 0$, and $K(p) = c$ occurs when $g(X, FY) = \pm 1$.

Proof. It is sufficient to prove the first assertion of the theorem. Let $R_{jih}{}^k$, g_{ji} and $F_j{}^i$ be the local components of R , g and F respectively, and put $R_{jihk} = g_{ka}R_{jih}{}^a$ and $F_{ji} = g_{ia}F_j{}^a$. Then (4.1) can be written as

$$(4.2) \quad R_{jihk} = -\frac{1}{4}c(g_{jh}g_{ik} - g_{jk}g_{ih} + F_{hj}F_{ki} - F_{kj}F_{hi} + 2F_{ij}F_{kh}).$$

Transvecting (4.2) with g^{ih} we have

$$(4.3) \quad R_{jk} = \frac{1}{2}(n+1)cg_{jk},$$

so that our space is Einsteinian. q.e.d.

Applying $\nabla_b \nabla_a$ to (4.2), we have

$$\begin{aligned} \nabla_b \nabla_a R_{jihk} = & -\frac{1}{4}c\{(\nabla_b \nabla_a F_{hj})F_{ki} + F_{hj}\nabla_b \nabla_a F_{ki} - (\nabla_b \nabla_a F_{kj})F_{hi} \\ & - F_{kj}\nabla_b \nabla_a F_{hi} + 2(\nabla_b \nabla_a F_{ij})F_{kh} + 2F_{ij}\nabla_b \nabla_a F_{kh}\} \end{aligned}$$

$$(4.4) \quad \begin{aligned} & -\frac{1}{4}c\{(\nabla_a F_{hj})\nabla_b F_{ki} + (\nabla_b F_{hj})\nabla_a F_{ki}\} \\ & + \frac{1}{4}c\{(\nabla_a F_{kj})\nabla_b F_{hi} + (\nabla_b F_{kj})\nabla_a F_{hi}\} \\ & - \frac{1}{2}c\{(\nabla_a F_{ij})\nabla_b F_{kh} + (\nabla_b F_{ij})\nabla_a F_{kh}\}. \end{aligned}$$

Since $R(X, Y) \cdot F = 0$ means that $\nabla_b \nabla_a F_{hj}$ is symmetric in a, b , the right hand side of (4.4) is symmetric in a, b . Thus from (4.4) we have

Lemma 4.2. *In an F-space of constant holomorphic sectional curvature, we have*

$$\nabla_b \nabla_a R_{jihk} - \nabla_a \nabla_b R_{jihk} = 0, \quad \text{i.e., } R(X, Y) \cdot R = 0.$$

Next, calculating the square of both sides of (4.2) we have

$$R_{jihk} R^{jihk} = 2c^2 n(n+1)$$

and therefore

$$(4.5) \quad R_{jihk} R^{jihk} = 2R^2/[n(n+1)],$$

since $C = R/[n(n+1)]$ from (4.3). Hence we obtain

Lemma 4.3. *In an F-space of constant holomorphic sectional curvature, the length of the tensor R_{jihk} is constant.*

On the other hand, the following two lemmas are known.

Lemma 4.4 (Lichnerowicz [4], Yano [10]). *In a Riemannian manifold, we have*

$$\begin{aligned} \Delta(R_{jihk} R^{jihk}) &= 2(\nabla_s R_{jihk})\nabla^s R^{jihk} - 4R^{jihk}\nabla_j(\nabla_h R_{ik} - \nabla_k R_{ih}) \\ &\quad - 4R^{jihk}H^s_{ihk, sj}, \end{aligned}$$

where Δ and $H^s_{jihk, st} X^s Y^t$ are the Laplacian and the components of $R(X, Y) \cdot R$ respectively.

Lemma 4.5 (Sawaki [5]). *An almost Hermitian manifold M is Kählerian if it satisfies:*

$$(i) \quad R(X, Y) \cdot F = 0, \quad \nabla_Z R(X, Y) \cdot F = 0$$

for any vector fields X, Y and Z on M ,

(ii) *the rank of the Ricci form is maximum.*

Making use of the above results, we can prove

Theorem 4.6. *If M is an F-space of nonzero constant holomorphic sectional curvature, then M is Kählerian.*

Proof. By virtue of Theorem 4.1, Lemma 4.2 and Lemma 4.3, from Lemma 4.4 we have $\nabla_s R_{jihk} = 0$, so that M is locally symmetric. Thus from Lemma 4.5 it follows that M is Kählerian.

Bibliography

- [1] A. Gray, *Nearly Kähler manifolds*, J. Differential Geometry **4** (1970) 283–309.
- [2] S. Kobayashi & K. Nomizu, *Foundations of differential geometry*, Vol. II, John Wiley, New York, 1969.
- [3] S. Kotō, *Some theorems on almost Kählerian spaces*, J. Math. Soc. Japan **12** (1960) 422–433.
- [4] A. Lichnerowicz, *Géométrie des groupes de transformations*, Dunod, Paris, 1958.
- [5] S. Sawaki, *Sufficient conditions for an almost Hermitian manifold to be Kählerian*, Hokkaido Math. J. (1972).
- [6] B. Smyth, *Differential geometry of complex hypersurfaces*, Thesis, Brown University, 1966.
- [7] S. Tachibana, *On almost-analytic vectors in certain almost-Hermitian manifolds*, Tôhoku Math. J. **11** (1959) 351–363.
- [8] K. Takamatsu, *Some properties of constant scalar curvature*, Bull. Fac. Ed. Kanazawa Univ. **19** (1968) 25–27.
- [9] H. Yanamoto, *On orientable hypersurface of R^7 satisfying $R(X,Y) \cdot F=0$* , Res. Rep. Nagaoka Tech. College **8** (1972) 9–14.
- [10] K. Yano, *Differential geometry on complex and almost complex spaces*, Pergamon, Oxford, 1965.

NIIGATA UNIVERSITY, NIIGATA, JAPAN