

CONFORMALITY OF RIEMANNIAN MANIFOLDS TO SPHERES

KRISHNA AMUR & V. S. HEGDE

1. Introduction

Let M be an orientable smooth Riemannian manifold of dimension n with Riemannian metric g_{ij} . Let ∇ be the covariant differentiation operator on M , and K_{hijk}, K_{ij}, r be the Riemann curvature tensor, Ricci curvature tensor, and scalar curvature tensor of M respectively. Let X denote the infinitesimal conformal transformation on M so that we have

$$(1.1) \quad (\mathcal{L}_X g)_{ij} = \nabla_i X_j + \nabla_j X_i = 2\rho g_{ij},$$

where ρ is a function, and \mathcal{L}_X denotes the Lie differentiation with respect to X . Assuming that $\mathcal{L}_X r = 0$ Yano, Obata, Hsiung-Mugridge, Hsiung-Stern (see [1], [2], [6], [8]) have studied the condition for a Riemannian n -manifold M to be conformal to an n -sphere. The purpose of this paper is to relax the condition $\mathcal{L}_X r = 0$ further, that is, to assume $\mathcal{L}_{D_\rho} \mathcal{L}_X r = 0$, and to obtain conditions for M to be conformal to an n -sphere where D_ρ is the vector field associated with the 1-form $d\rho$. Towards this end we prove the following theorems.

Theorem 1.1. *If a compact orientable smooth Riemannian manifold M of dimension $n > 2$ admitting an infinitesimal conformal transformation $X: \mathcal{L}_X g = 2\rho g, \rho \neq \text{constant}$ with $\mathcal{L}_{D_\rho} \mathcal{L}_X r = 0$ satisfies $\int_M \left(A_{ij} \rho^i \rho^j + \frac{\alpha}{n^2} \mathcal{L}_X \mathcal{L}_{D_\rho} r \right) dv \geq 0$ where $A_{ij} = K_{ij} - (\alpha r/n) g_{ij}$ and $\alpha = 1$, then M is conformal to an n -sphere.*

Theorem 1.2. *Let M be an orientable smooth Riemannian manifold of dimension $n > 2$ admitting an infinitesimal conformal transformation X satisfying (1.1) such that $\rho \neq \text{constant}$, and $\mathcal{L}_{D_\rho} \mathcal{L}_X r = 0$. Then M is conformal to an n -sphere if $\mathcal{L}_X \mathcal{L}_{D_\rho} r \geq 0$ and $\mathcal{L}_X |G|^2 = 0$ where $G_{ij} = K_{ij} - (r/n) g_{ij}$.*

Theorem 1.3. *Let M be an orientable smooth Riemannian manifold of dimension $n > 2$ admitting an infinitesimal conformal transformation X satisfying (1.1) such that $\rho \neq \text{constant}$ and $\mathcal{L}_{D_\rho} \mathcal{L}_X r = 0$. Then M is conformal to an n -sphere if $\mathcal{L}_X \mathcal{L}_{D_\rho} r \geq 0$ and $\mathcal{L}_X |W|^2 = 0$ where W is a tensor defined in § 2.*

It is shown in § 5 that when $\mathcal{L}_x r = 0$, Theorems 1.1 and 1.2 reduce to those of Yano [6], and Theorem 1.3 reduces to that of Hsiung and Stern [2]. Also it is proved that when $r = \text{constant}$, the condition $\alpha = 1$ in Theorem 1.1 may be replaced by $\alpha \geq 1$, and the manifold M would then be isometric to a sphere. The following known theorems are needed in the proofs of our theorems.

Theorem 1.4 (Obata [3]). *If a complete Riemannian manifold M of dimension $n \geq 2$ admits a nonconstant function ρ such that $\nabla_i \nabla_j \rho = -c^2 \rho g_{ij}$ where c is a positive constant, then M is isometric to an n -sphere of radius $1/c$.*

Theorem 1.5 (Tashiro [4]). *If a complete Riemannian manifold M of dimension $n \geq 2$ admits a nonconstant function ρ such that $\nabla_i \nabla_j \rho + (1/n)\Delta \rho g_{ij} = 0$, then M is conformal to an n -sphere.*

2. Notations and formulas

The raising and lowering of the indices are, as usual, carried out respectively with g^{ij} and g_{ij} . The tensors thus obtained are called associated tensors. Let S, T be covariant tensors of order s with local components $S_{i_1 \dots i_s}$ and $T_{i_1 \dots i_s}$ respectively. The associated contravariant components of T are $T^{i_1 \dots i_s}$. We define the inner product of S and T by $S_{i_1 \dots i_s} T^{i_1 \dots i_s}$ and denote it by $\langle S, T \rangle$. If $S = T$ we write $|S|^2$ for $\langle S, S \rangle$. For the sake of easy reference we list some known formulas; for details see Yano [7]:

$$(2.1) \quad \mathcal{L}_x r = 2(n-1)\Delta \rho - 2r\rho,$$

$$(2.2) \quad \mathcal{L}_x g^{ij} = -2\rho g^{ij},$$

$$(2.3) \quad \mathcal{L}_x K_{hijk} = 2\rho K_{hijk} - g_{hk} \nabla_j \rho_i + g_{hj} \nabla_i \rho_k - g_{ij} \nabla_h \rho_k + g_{ik} \nabla_h \rho_j,$$

$$(2.4) \quad \mathcal{L}_x K_{ij} = g_{ij} \Delta \rho - (n-2) \nabla_i \rho_j,$$

$$(2.5) \quad \nabla_k \nabla_i Y^j - \nabla_i \nabla_k Y^j = K_{kih}^j Y^h, \quad g^{kj} (\nabla_k \nabla_i Y_j - \nabla_i \nabla_k Y_j) = K_i^h Y_h,$$

where Δ is the Laplace-Beltrami operator on M , and Y is any differentiable vector field on M . If the associated 1-form of a vector field Y is ξ , the components of ΔY and $\Delta \xi$ are given by

$$(2.6) \quad \Delta Y: -g^{kj} \nabla_k \nabla_j Y^i + K_h^i Y^h, \quad \Delta \xi: -g^{kj} \nabla_k \nabla_j Y_i + K_i^h Y_h.$$

If d is the exterior differentiation operator on M , and f is any function on M , then we denote the associated vector field of the 1-form df by Df .

Write $f_i = \nabla_i f$, and $f^i = g^{ij} f_j$, and define the tensors Z and W by

$$(2.7) \quad Z_{hijk} = K_{hijk} - \frac{r}{n(n-1)} (g_{hk} g_{ij} - g_{hj} g_{ik}),$$

$$(2.8) \quad \begin{aligned} W_{hijk} = & aZ_{hijk} + b_1g_{hk}G_{ij} - b_2g_{hj}G_{ik} + b_3g_{ij}g_{hk} \\ & - b_4g_{ik}G_{hj} + b_5g_{hi}G_{jk} - b_6g_{jk}G_{hi}, \end{aligned}$$

where a, b_1, \dots, b_6 are any constants.

3. Lemmas

Lemma 3.1. *Let M be a compact orientable Riemannian manifold of dimension $n \geq 2$. For any vector field Y and a differentiable function f we have*

$$\int_M (\nabla_i Y^i) dv = 0, \quad \int_M \Delta f dv = 0.$$

The first equation is the well known Green's formula, and the second follows as a consequence of the first.

Lemma 3.2 (Yano and Sawaki [9]). *Let M be a compact oriented Riemannian manifold of dimension $n > 2$ admitting an infinitesimal non-isometric conformal transformation X satisfying (1.1). Then for any function f on M we have*

$$\int_M \rho f dv = -\frac{1}{n} \int_M \mathcal{L}_X f dv.$$

Lemma 3.3. *For a manifold M having the same properties as in Lemma 3.2, we have*

$$(3.1) \quad \int_M (\Delta \rho)^2 dv = \int_M \rho^i \nabla_i \Delta \rho dv = \int_M (K_{ij} \rho^j - g^{kj} \nabla_k \nabla_j \rho_i) \rho^i dv.$$

Furthermore, if $r = \text{constant}$, then

$$(3.2) \quad \int_M (\Delta \rho)^2 dv = \frac{r}{n-1} \int_M \rho^i \rho_i dv$$

Proof. $\nabla_i (\rho^i \Delta \rho) = \rho^i \nabla_i \Delta \rho - (\Delta \rho)^2 = (K_{ij} \rho^j - g^{kj} \nabla_k \nabla_j \rho_i) \rho^i - (\Delta \rho)^2$ by (2.5). Integrating and using Lemma 3.1 we get (3.1).

Setting $\mathcal{L}_X r = 0$ in (2.1) and using the result in (a) we obtain (3.2).

Lemma 3.4. *Let M be a manifold having the same properties as in Lemma 3.2 and satisfying the condition $\mathcal{L}_{D_\rho} \mathcal{L}_X r = 0$. Then*

$$(3.3) \quad \int_M (r \rho^i \rho_i) dv = (n-1) \int_M (\Delta \rho)^2 dv + \frac{1}{n} \int_M \mathcal{L}_X \mathcal{L}_{D_\rho} r dv.$$

Furthermore, if $\mathcal{L}_X r = 0$, then

$$(3.4) \quad \frac{1}{n} \int_M \mathcal{L}_x \mathcal{L}_{D\rho} r dv = \int_M r \rho_i \rho^i dv - \frac{1}{n-1} \int_M r^2 \rho^2 dv .$$

Proof. From (2.1) we have

$$\begin{aligned} 0 &= \mathcal{L}_{D\rho} \mathcal{L}_x r = 2\mathcal{L}_{D\rho}((n-1)\Delta\rho - \rho r) \\ &= 2[(n-1)\rho^i \nabla_i \Delta\rho - \rho \rho^i \nabla_i r - r \rho_i \rho^i] . \end{aligned}$$

Integrating and using Lemmas 3.2 and 3.3 we get (3.3). If $\mathcal{L}_x r = 0$, then $(n-1)\Delta\rho = \rho r$. Substituting this in (3.3) we obtain (3.4).

4. Proofs of Theorems

Proof of Theorem 1.1. For an arbitrary vector field Y , by writing $\nabla^j = g^{ji} \nabla_i$ and using (2.5) we find that

$$\begin{aligned} &\nabla^j \left(\nabla_j Y_i + \nabla_i Y_j - \frac{2\alpha}{n} g_{ij} \nabla_l Y^l \right) Y^i \\ &= \left(g^{jk} \nabla_k \nabla_j Y_i + \nabla_i \nabla_j Y^j + K_{jih}{}^j Y^h - \frac{2\alpha}{n} \nabla_i \nabla_l Y^l \right) Y^i + \frac{2}{n} \alpha (1 - \alpha) (\nabla_l Y^l)^2 \\ &\quad + \frac{1}{2} \left(\nabla_j Y_i + \nabla_i Y_j - \frac{2\alpha}{n} g_{ij} \nabla_l Y^l \right) \left(\nabla^j Y^i + \nabla^i Y^j - \frac{2\alpha}{n} g^{ij} \nabla_l Y^l \right) . \end{aligned}$$

Putting $Y^i = \rho^i$, integrating the above equation, using Lemmas 3.1 and 3.3, and setting $K_{ij} = A_{ij} + (r\alpha/n)g_{ij}$ we get

$$\begin{aligned} &\int_M A_{ij} \rho^i \rho^j dv + \frac{1}{n} (-n + 2\alpha - \alpha^2) \int_M (\Delta\rho)^2 dv + \frac{\alpha}{n} \int_M r \rho_i \rho^i dv \\ &\quad + \int_M \left[\nabla \nabla \rho + \frac{\alpha}{n} g \Delta \rho^2 \right] dv = 0 . \end{aligned}$$

Substituting (3.3) in the above equation and simplifying we obtain finally

$$(4.1) \quad \int_M \left(A_{ij} \rho^i \rho^j + \frac{\alpha}{n^2} \mathcal{L}_x \mathcal{L}_{D\rho} r \right) dv + \int_M \left[\nabla \nabla \rho + \frac{1}{n} (1 + \sqrt{(\alpha-1)(n-1)}) g \Delta \rho^2 \right] dv = 0 .$$

Hence Theorem 1.1 follows from Theorem 1.5 and the integral formula (4.1).

Proof of Theorem 1.2. From (2.2) and (2.4) we easily get

$$(4.2) \quad \langle G, \nabla \nabla \rho \rangle = -\frac{2\rho}{n-2} |G|^2 - \frac{1}{2(n-2)} \mathcal{L}_x |G|^2 .$$

On the other hand,

$$(4.3) \quad \nabla^i(G_{ij}\rho^i\rho^j) = G_{ij}\rho^i\rho^j + \rho\langle G, \nabla\nabla\rho\rangle + \frac{n-2}{2n}\rho(\rho^i\nabla_i r).$$

Multiply (4.2) by ρ and integrate, integrate (4.3), and eliminate $\int_M \rho\langle G, \nabla\nabla\rho\rangle dv$ from the two resulting equations so that we have the integral formula

$$(4.4) \quad \int_M \left(G_{ij}\rho^i\rho^j + \frac{1}{n^2}\mathcal{L}_x\mathcal{L}_{D_\rho}r \right) dv \\ = \frac{2}{n-2} \int_M \left((\rho^2|G|^2 + \frac{1}{4}\rho\mathcal{L}_x|G|^2) \right) dv + \frac{1}{2n} \int_M \mathcal{L}_x\mathcal{L}_{D_\rho}r dv.$$

Hence Theorem 1.2 follows from Theorem 1.1 and the integral formula (4.4).

Proof of Theorem 1.3. From (2.7), (2.8), (2.3), (2.4) and (2.2) we get (for details see [2])

$$(4.5) \quad \langle \mathcal{L}_x W, W \rangle = 2\rho|W|^2 - c\langle G, \nabla\nabla\rho \rangle,$$

where c is a constant given by

$$\frac{c-4a^2}{n-2} = 2a \sum_{i=1}^4 b_i + \left(\sum_{i=1}^6 (-1)^{i-1} b_i \right)^2 \\ - 2(b_1 b_3 + b_2 b_4 - b_5 b_6) + (n-1) \sum_{i=1}^6 b_i^2.$$

Here $c \geq 0$. Use of (2.2) yields

$$(4.6) \quad \mathcal{L}_x|W|^2 = 2\langle \mathcal{L}_x W, W \rangle - 8\rho|W|^2$$

Thus from (4.3), (4.5) and (4.6) we obtain

$$(4.7) \quad c \int_M \left(G_{ij}\rho^i\rho^j + \frac{1}{n^2}\mathcal{L}_x\mathcal{L}_{D_\rho}r \right) dv \\ = 2 \int_M \rho^2|W|^2 dv + \frac{1}{2} \int_M \rho\mathcal{L}_x|W|^2 dv + \frac{c}{2n} \int_M \mathcal{L}_x\mathcal{L}_{D_\rho}r dv.$$

Hence Theorem 1.3 follows from Theorem 1.1 and the integral formula (4.7).

5. Special cases

1. Let $\alpha = 1$ and $\mathcal{L}_x r = 0$. The condition for conformality in Theorem 1.1 reduces, by (3.4), to

$$\int_M \left(K_{ij} \rho^i \rho^j - \frac{r^2 \rho^2}{n(n-1)} \right) dv \geq 0.$$

Also we have

$$\mathcal{L}_x |G|^2 = \mathcal{L}_x |R|^2, \quad \mathcal{L}_x |W|^2 = \alpha^2 \mathcal{L}_x |K|^2 + \frac{c - 4\alpha^2}{n-2} \mathcal{L}_x |R|^2,$$

where $|K|^2 = K_{hijk} K^{hijk}$ and $|R|^2 = K_{ij} K^{ij}$. The condition $\mathcal{L}_x \mathcal{L}_{D_\rho} r \geq 0$ for M implies by (3.4) that

$$\int_M \left(r \rho_i \rho^i - \frac{r^2 \rho^2}{n-1} \right) dv \geq 0.$$

With these, Theorem, 1.1 and 1.2 reduce to results due to Yano [6], and Theorem 1.3 reduces to that due to Hsiung and Stern [2].

2. Let $\alpha \geq 1$ and $r = \text{constant}$. From (4.1) it follows that M is isometric to a sphere if

$$\int_M A_{ij} \rho^i \rho^j dv \geq 0;$$

when $\alpha = 1$, this is a known condition [5]

$$\int_M G_{ij} \rho^i \rho^j dv \geq 0$$

for M to be isometric to a sphere.

References

- [1] C. C. Hsiung & L. R. Mugridge, *Conformal changes of metrics on a Riemannian manifold*, Math. Z. **119** (1971) 179-187.
- [2] C. C. Hsiung & L. W. Stern, *Conformality and isometry of Riemannian manifolds to spheres*, Trans. Amer. Math. Soc. **163** (1972) 65-73.
- [3] M. Obata, *Certain conditions for a Riemannian manifold to be isometric with a sphere*, J. Math. Soc. Japan **14** (1962) 333-340.
- [4] Y. Tashiro, *Complete Riemannian manifolds and some vector fields*, Trans. Amer. Math. Soc. **117** (1965) 251-275.
- [5] K. Yano, *On Riemannian manifolds with constant scalar curvature admitting a conformal transformation group*, Proc. Nat. Acad. Sci. U.S.A. **55** (1966) 472-476.
- [6] —, *On Riemannian manifolds admitting an infinitesimal conformal transformation*, Math. Z. **113** (1970) 205-214.
- [7] —, *Integral formulas in Riemannian geometry*, Marcel Dekker, New York, 1970.
- [8] K. Yano & M. Obata, *Conformal changes of Riemannian metrics*, J. Differential Geometry **4** (1970) 53-72.
- [9] K. Yano & S. Sawaki, *Riemannian manifolds admitting a conformal transformation group*, J. Differential Geometry **2** (1968) 161-184.