

A SEMI-LOCAL COMBINATORIAL FORMULA FOR THE SIGNATURE OF A $4k$ -MANIFOLD

ANDREW RANICKI & DENNIS SULLIVAN

Let M be a compact oriented and triangulated manifold of dimension $4k$. If we orient each simplex, we obtain geometric bases for the chains and co-chains, which are thus identified

$$C^i = C_i .$$

The classical boundary and coboundary operators

$$\delta : C_i \rightarrow C_{i-1} , \quad d : C_i \rightarrow C_{i+1}$$

are transposes of one another. We shall describe below a symmetric transformation

$$* : C_i \rightarrow C_{4k-i} .$$

If M has no boundary, we then form the symmetric transformation

$$\begin{pmatrix} * & \partial \\ d & 0 \end{pmatrix} : C \rightarrow C ,$$

where $C = C_{2k} \oplus C_{2k+1}$. If M has a boundary, we form the transformation

$$\begin{pmatrix} * & \partial \\ d & 0 \end{pmatrix} : C \cdot \rightarrow C \cdot ,$$

where $C \cdot$ is the d -sub (or ∂ -quotient) space of C consisting of the chains which vanish (or have no coefficient) on the boundary. In either case we have

Theorem. signature $M = \text{signature} \begin{pmatrix} * & \partial \\ d & 0 \end{pmatrix}$.

The operation $*$ is defined as follows: say two oriented simplices σ and τ are complementary with sign $*(\sigma, \tau)$ if

- (i) σ and τ are of complementary dimensions,
- (ii) σ and τ span a $4k$ -simplex η ,
- (iii) orientation $\sigma \cdot \text{orientation } \tau = *(\sigma, \tau) \cdot \text{orientation } \eta$.

Then let

Communicated by W. P. A. Klingenberg, June 7, 1974.

$$*: C_i \rightarrow C_{4k-i}; \sigma \mapsto i!(4k-i)! \sum_{\tau} *(\sigma, \tau)\tau,$$

where τ ranges over the complementary simplices of σ . This is a combinatorial analogue of the $*$ operator on a Riemannian manifold. We are indebted to Walter Neumann for pointing out that the factor $i!(4k-i)!$ is required here for (i) below to be satisfied. However, the factor may be omitted in the statement of the theorem, because the matrix identity

$$\begin{pmatrix} \lambda & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} A & B \\ B^* & 0 \end{pmatrix} \begin{pmatrix} \lambda & 0 \\ 0 & 1 \end{pmatrix} = \lambda \begin{pmatrix} \lambda A & B \\ B^* & 0 \end{pmatrix}$$

shows that

$$\text{signature} \begin{pmatrix} A & B \\ B^* & 0 \end{pmatrix} = \text{signature} \begin{pmatrix} \lambda A & B \\ B^* & 0 \end{pmatrix} \in \mathbb{Z}$$

for any strictly positive number λ .

Note that $\begin{pmatrix} * & \partial \\ d & 0 \end{pmatrix}$ is local in character. The value of this transformation on a chain depends only on the simplicial geometry near the chain. However, our formula is not solely a function of the geometry around each vertex (plus a boundary term if M is not closed) which is the real goal here. (Such a formula is known in Riemannian geometry, using curvature and the Thom-Hirzebruch signature formula.)

We shall describe the proof in the closed case. The proof in the bounded case is identical, after replacing C by $C \cdot$.

There are three steps:

(i) Establish a commutative diagram

$$\begin{array}{ccccc} C_{2k-1} & \xrightarrow{d} & C_{2k} & \xrightarrow{d} & C_{2k+1} \\ * \downarrow & & \downarrow * & & \downarrow * \\ C_{2k+1} & \xrightarrow{\partial} & C_{2k} & \xrightarrow{\partial} & C_{2k-1} \end{array}$$

yielding a transformation

$$*: H^{2k} \rightarrow H_{2k}.$$

(ii) Identify this transformation with the cup product pairing

$$(x, y) \mapsto \int_M x \cup y$$

on H^{2k} defining the signature of M . (Recall that the signature of a manifold with boundary is defined by this procedure using $H^{2k}(M, \partial M)$. Thus $C \cdot$ is

relevant in the bounded case.)


(iii) Appeal to the algebraic lemma below to obtain

$$\text{signature } (H^{2k}, *) = \text{signature } \left(C_{2k} \oplus C_{2k+1}, \begin{pmatrix} * & \partial \\ d & 0 \end{pmatrix} \right) \in \mathbf{Z}.$$

We describe two ways to establish (i) and (ii). First one can directly establish (i) by a combinatorial argument and then identify $*$ with the cup product by appealing to Whitney's discussion [4, p. 363] and [5]. This method is elementary but not completely conceptual. Perhaps a more satisfying method (and one which is useful in a variety of contexts) is to use Whitney's embedding of the cochains on a complex into the forms on the complex. For an elementary cochain σ with vertices x_0, x_1, \dots, x_r Whitney defines ([4, p. 229], see also [3]) the differential form

$$\omega_\sigma = r! \sum_{i=0}^r (-1)^i x_i dx_0 \wedge dx_1 \wedge \dots \wedge \widehat{dx}_i \wedge \dots \wedge dx_r,$$

thinking of the x_i as barycentric coordinates with nonzero values in the stars of vertices of σ . The form ω_σ is nonzero only in the open star of σ , and its integral over any (tiny) simplex σ' in the star has appealing geometric interpretation; namely, consider the figure:



Then

$$\int_{\sigma'} \omega_\sigma = \frac{\text{volume } \sigma' * \tau}{\text{volume } \sigma * \tau},$$

where $\sigma' \subset \text{interior } \sigma * \tau$.

The linear extension of this formula embeds cochains into the forms on the polyhedron, it commutes with d , and it yields an isomorphism between simplicial cohomology and de Rham cohomology (see [4], [5] for real coefficients and [2] for rational coefficients).

The cup product on cohomology is represented by the exterior product of forms. For example, if σ and τ are of complementary dimensions, then $\omega_\sigma \wedge \omega_\tau$ is nonzero precisely when σ and τ span a $4k$ -simplex η . We can calculate $\int_\eta \omega_\sigma \wedge \omega_\tau$ to obtain a nonzero number with the sign of $*(\sigma, \tau)$ (see the

appendix at the end of this paper). This shows (ii), namely, the *-pairing from diagram (ii) is a positive multiple of the cup-product pairing. The commutativity of diagram (i) follows by integrating the equation among differential forms

$$d(\omega \wedge \eta) = d\omega \wedge \eta \pm \omega \wedge d\eta$$

over the manifold. In the closed case we find

$$*d = \pm \partial^* .$$

(The other relation $*\partial = \pm d^*$ is quite false.) In the bounded case we also find this relation among the chains which vanish on the boundary C . Thus (i) and (ii) are proved.

Algebraic lemma. *Let*

$$\begin{array}{ccccc} U & \xrightarrow{f} & V & \xrightarrow{g} & W \\ \theta \downarrow & & \downarrow \varphi & & \downarrow \theta^* \\ W^* & \xrightarrow{g^*} & V^* & \xrightarrow{f^*} & U^* \end{array}$$

be a commutative diagram of finite-dimensional vector spaces and morphisms with

$$gf = 0 \in \text{Hom}(U, W), \quad \varphi^* = \varphi \in \text{Hom}(V, V^*).$$

Then the bilinear symmetric form $(H, \bar{\varphi})$ induced by φ on the homology space $H = \ker g / \text{im } f$ is such that

$$\text{signature}(H, \bar{\varphi}) = \text{signature}\left(V \oplus W^*, \begin{pmatrix} \varphi & g^* \\ g & 0 \end{pmatrix}\right) \in \mathbf{Z}.$$

Proof. The signature $\sigma(V, \varphi) \in \mathbf{Z}$ of (V, φ) is (by definition) the signature of the associated nondegenerate form $(V/\ker \varphi, \bar{\varphi})$.

The form (Z, ξ) to which φ restricts on $Z = \ker g$ has the same associated nondegenerate form as $(H, \bar{\varphi})$, so that

$$\sigma(H, \bar{\varphi}) = \sigma(Z, \xi) \in \mathbf{Z}.$$

Choosing direct complements X, Y to Z, fX in V, W respectively, we can write

$$\begin{aligned} \varphi &= \begin{pmatrix} \xi & \psi^* \\ \psi & \xi \end{pmatrix}: V = Z \oplus X \rightarrow Z^* \oplus X^*, \\ f &= \begin{pmatrix} 0 & \iota \\ 0 & 0 \end{pmatrix}: V = Z \oplus X \rightarrow W = fX \oplus Y, \end{aligned}$$

where $\iota \in \text{Hom}(X, fX)$ is an isomorphism. Then there is defined an isomorphism of forms

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \psi & \frac{1}{2}\xi & \iota^* & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} : \left(V \oplus W^*, \begin{pmatrix} \varphi & g^* \\ g & 0 \end{pmatrix} \right) = \left(Z \oplus X \oplus fX^* \oplus Y^*, \begin{pmatrix} \xi & \psi^* & 0 & 0 \\ \psi & \xi & \iota^* & 0 \\ 0 & \iota & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \right) \\ \rightarrow (Y, \xi) \oplus \left(X \oplus X^*, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \oplus (Y^*, 0),$$

and so

$$\begin{aligned} \sigma \left(V \oplus W^*, \begin{pmatrix} \varphi & g^* \\ g & 0 \end{pmatrix} \right) &= \sigma(Z, \xi) + \sigma \left(X \oplus X^*, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) + \sigma(Y^*, 0) \\ &= \sigma(Z, \xi) = \sigma(H, \bar{\varphi}) \in \mathbf{Z}, \end{aligned}$$

proving the lemma *a*.

The above lemma comes from an algebraic theory of surgery (see [1]) in the following special case. Let $(C, d, *)$ be a $4k$ -dimensional cochain complex (C, d) with a symmetric pairing

$$* : (C, d) \rightarrow (C, \partial)$$

to the dual chain complex, involving the usual dimension shift:

$$\begin{array}{ccccccc} C^0 & \xrightarrow{d} & C^1 & \xrightarrow{d} & \dots & \xrightarrow{d} & C^{2k} & \xrightarrow{d} & \dots & \xrightarrow{d} & C^{4k} \\ * \downarrow & & \downarrow * & & & & \downarrow * & & & & \downarrow * \\ C_{4k} & \xrightarrow{\partial} & C_{4k-1} & \xrightarrow{\partial} & \dots & \xrightarrow{\partial} & C_{2k} & \xrightarrow{\partial} & \dots & \xrightarrow{\partial} & C_0 \end{array}$$

The cohomology $H^{2k+1}(C)$ is killed by replacing $(C, d, *)$ with the triple $(C', d', *)$ defined by

$$\begin{array}{ccccccc} C^0 & \xrightarrow{d} & C^1 & \xrightarrow{d} & \dots & \xrightarrow{d} & C^{2k-1} & \xrightarrow{\begin{pmatrix} d \\ \varepsilon^* \end{pmatrix}} & C^{2k} \oplus Z \oplus Z^* & \xrightarrow{(d \ e \ 0)} & C^{2k+1} & \xrightarrow{d} & \dots & \xrightarrow{d} & C^{4k} \\ * \downarrow & & \downarrow * & & & & \downarrow * & & \downarrow \begin{pmatrix} * & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} & & \downarrow * & & & & \downarrow * \\ C_{4k} & \xrightarrow{\partial} & C_{4k-1} & \xrightarrow{\partial} & \dots & \xrightarrow{\partial} & C_{2k+1} & \xrightarrow{\begin{pmatrix} \partial \\ \varepsilon \end{pmatrix}} & C_{2k} \oplus Z^* \oplus Z & \xrightarrow{(\partial \ 0 \ *e)} & C_{2k-1} & \xrightarrow{\partial} & \dots & \xrightarrow{\partial} & C_0 \end{array}$$

where $e \in \text{Hom}(Z, C^{2k+1})$ is the inclusion of $Z = \ker(d: C^{2k+1} \rightarrow C^{2k+2})$ and $\varepsilon = e^* \in \text{Hom}(C_{2k+1}, Z^*)$ is the dual. The nondegenerate forms associated with $(H^{2k}(C'), *)$ and $\left(C^{2k} \oplus C_{2k+1}, \begin{pmatrix} * & \hat{\partial} \\ d & 0 \end{pmatrix}\right)$ are isomorphic, so that they have the same signature. The algebraic lemma is thus a verification of

$$\text{signature}(H^{2k}(C'), *) = \text{signature}(H^{2k}(C), *) \in Z,$$

that is, “cobordant complexes have the same signature”.

Appendix: The calculation of $\int_{\eta} \omega_{\varepsilon} \wedge \omega$.

Let σ, τ be complementary oriented simplices in the triangulation of M . Write x_0, x_1, \dots, x_r for the vertices of σ , and y_0, y_1, \dots, y_s for those of τ , with

$$x_0 = y_0, \quad r + s = 4k.$$

Let η be the $4k$ -dimensional simplex spanned by σ and τ , with the vertices z_0, z_1, \dots, z_{4k} given by

$$z_i = \begin{cases} x_0 = y_0, & i = 0, \\ x_i, & 0 < i \leq r, \\ y_{i-r}, & r < i \leq 4k. \end{cases}$$

We have the differential forms

$$\begin{aligned} \omega_{\varepsilon} &= r! \sum_{i=0}^r (-)^i x_i dx_0 \wedge \dots \wedge \widehat{dx_i} \wedge \dots \wedge dx_r, \\ \omega_{\varepsilon} &= s! \sum_{j=0}^s (-)^j y_j dy_0 \wedge \dots \wedge \widehat{dy_j} \wedge \dots \wedge dy_s, \end{aligned}$$

and also define

$$\omega = dz_1 \wedge dz_2 \wedge \dots \wedge dz_{4k}.$$

The relation

$$\sum_{i=0}^{4k} z_i = 1$$

holds in η (and indeed defines η), so that

$$\sum_{i=0}^{4k} dz_i = 0.$$

Using these relations, we obtain

$$\begin{aligned}
 & \left(\sum_{i=0}^r (-)^i x_i dx_0 \wedge \cdots \wedge \widehat{dx}_i \wedge \cdots \wedge dx_r \right) \\
 & \quad \wedge \left(\sum_{j=0}^s (-)^j y_j dy_0 \wedge \cdots \wedge \widehat{dy}_j \wedge \cdots \wedge dy_s \right) \\
 & = x_0 y_0 \omega + (x_0 dx_1 \wedge \cdots \wedge dx_r) \\
 & \quad \wedge \left(\sum_{j=1}^s (-)^j y_j dy_0 \wedge \cdots \wedge \widehat{dy}_j \wedge \cdots \wedge dy_s \right) \\
 & \quad + \left(\sum_{i=1}^r (-)^i x_i dx_0 \wedge \cdots \wedge \widehat{dx}_i \wedge \cdots \wedge dx_r \right) \\
 & \quad \wedge (y_0 dy_1 \wedge \cdots \wedge dy_s) \\
 & = \left(x_0 y_0 + x_0 \left(\sum_{j=1}^s y_j \right) + \left(\sum_{i=1}^r x_i \right) y_0 \right) \omega \\
 & = z_0 \left(\sum_{i=0}^{4k} z_i \right) \omega = z_0 \omega .
 \end{aligned}$$

It follows that

$$\int_{\tau} \omega_\sigma \wedge \omega_\tau = r! s! \int_{\tau} z_0 \omega = *(\sigma, \tau) \frac{r! s!}{(4k+1)!} .$$

References

- [1] A. A. Ranicki, *Geometric L-theory*, to appear.
- [2] D. Sullivan, *Differential forms and the topology of manifolds*, Proc. Conf. on Manifolds (Tokyo), 1973.
- [3] A. Weil, *Sur les théorèmes de de Rham*, Comment. Math. Helv. **26** (1952) 119–145.
- [4] H. Whitney, *Geometric integration theory*, Princeton University Press, Princeton, 1956.
- [5] —, *On products in a complex*, Ann. of Math **39** (1938) 397–432.

INSTITUT DES HAUTES ETUDES SCIENTIFIQUES
BURES-SUR-YVETTE