

EXISTENCE OF GENERALIZED SYMMETRIC RIEMANNIAN SPACES OF ARBITRARY ORDER

OLDŘICH KOWALSKI

A Riemannian symmetric space is a Riemannian manifold (M, g) with the following properties: for each $x \in M$ there is a (unique) isometry J_x on M such that

- (a) x is an isolated fixed point of J_x ,
- (b) $(J_x)^2 = \text{identity}$.

It is also easy to show the following property: for every two points $x, y \in M$ we have

- (c) $J_x \circ J_y = J_z \circ J_x$, where $z = J_x(y)$.

The following is a direct generalization of the previous situation.

Definition. A Riemannian k -symmetric space ($k \geq 2$) is a Riemannian manifold (M, g) on which a family $\{s_x\}_{x \in M}$ of isometries exists with the following properties:

- (a) Each $x \in M$ is an isolated fixed point of the corresponding s_x ,
- (b) $(s_x)^k = \text{identity}$ for all $x \in M$, and k is the minimum number of this property,
- (c) for every $x, y \in M$, $s_x \circ s_y = s_z \circ s_x$, where $z = s_x(y)$.

In fact, Ledger and Obata [3] have proved that for every $k > 2$ there is a k -symmetric Riemannian space which is not symmetric. The purpose of this paper is to strengthen the previous result in the following sense: *for every $k > 2$ there is a k -symmetric Riemannian space which is not l -symmetric for $l = 2, \dots, k - 1$.* (Such a Riemannian space is said to be *generalized symmetric of order k* ; see [2]). In our further considerations we shall make full use of the original construction by Ledger and Obata.

1. Let $M = G/H$ be a homogeneous Riemannian space. As usual, we suppose G acting effectively on the coset space G/H . Thus the Lie group G can be considered as a group of isometries on M . Let $\pi: G \rightarrow M$ denote the canonical projection.

Proposition 1. *Let G admit an automorphism σ such that*

- (i) $H = G^\sigma = \text{the fixed point set of } \sigma$,
- (ii) $\sigma^k = \text{identity}$,
- (iii) *the transformation s of M determined by $\pi \circ \sigma = s \circ \pi$ is an isometry.*

Then M is a Riemannian k -symmetric space.

Proof. For $x \in M$ define a transformation s_x of M by the formula $s_x = g \circ s \circ g^{-1}$, where $g \in \pi^{-1}(x)$. Then s_x is independent of the choice of g . In fact, for each $h \in H$ we have $L_h \circ \sigma \circ L_{h^{-1}} = \sigma$ and $\pi \circ L_h = h \circ \pi$. Hence $(h \circ s \circ h^{-1}) \circ \pi = h \circ (s \circ \pi) \circ L_{h^{-1}} = h \circ (\pi \circ \sigma) \circ L_{h^{-1}} = \pi \circ (L_h \circ \sigma \circ L_{h^{-1}}) = \pi \circ \sigma$, and consequently, $h \circ s \circ h^{-1} = s$. Thus for $g' = gh$ we obtain $g' \circ s \circ g'^{-1} = g \circ s \circ g^{-1}$.

It is clear that $(s_x)^k = \text{identity}$ for each $x \in M$. We have to prove that x is an isolated fixed point of s_x . For, it is sufficient to show that the initial point $o \in M$, $o = \pi(H)$, is an isolated fixed point of s . Condition (iii) implies that $s_{*o} \circ \pi_{*e} = \pi_{*e} \circ \sigma_{*e}$ on the tangent space G_e . Let $X \in M_o$ be such that $s_{*o}(X) = X$, and let $\tilde{X} \in G_e$ be a lift of X . Then $\pi_{*e}(\sigma_{*e}(\tilde{X})) = \pi_{*e}(\tilde{X})$, and hence $\sigma_{*e}(\tilde{X}) = \tilde{X} + \tilde{Z}$, where $\tilde{Z} \in H_e$. Now $\sigma_{*e}(\tilde{Z}) = \tilde{Z}$, and $(\sigma_{*e})^k(\tilde{X}) = \tilde{X} + k\tilde{Z} = \tilde{X}$ because $(\sigma_{*e})^k = \text{identity}$. Thus $\tilde{Z} = 0$ and \tilde{X} is a fixed vector of σ_{*e} . We deduce $\tilde{X} \in H_e$ and $X = 0$. Because s_{*o} has no nonzero fixed vectors and s is an isometry of M , we conclude that o is an isolated fixed point of s .

Finally, we have to prove the formula $s_x \circ s_y = s_z \circ s_x$, $z = s_x(y)$. For this purpose we shall identify the elements of G with the corresponding transformations of M . Then we deduce $s \circ g \circ s^{-1} = \sigma(g)$. Put $s_x = g \circ s \circ g^{-1}$, $s_y = g' \circ s \circ (g')^{-1}$, where $x = g(o)$ and $y = g'(o)$. Then $(g \circ s \circ g^{-1} \circ g' \circ s^{-1})(o) = s_x(g'(o)) = s_x(y)$. On the other hand, $g \circ s \circ g^{-1} \circ g' \circ s^{-1} = g \circ \sigma(g^{-1}g') = g''$ belongs to G . Consequently, $s_x \circ s_y = g \circ s \circ g^{-1} \circ g' \circ s \circ (g')^{-1} = g'' \circ s \circ (g'')^{-1} \circ g \circ s \circ g^{-1} = s_{s_x(y)} \circ s_x$.

2. We shall recall here a class of Riemannian manifolds constructed by Ledger and Obata (see [3]). Let G be a compact connected nonabelian Lie group, G^{k+1} the direct product of G with itself $(k+1)$ -times, and ΔG^{k+1} the diagonal of G^{k+1} . Consider the action of G^{k+1} on G^k given by

$$(x_1, \dots, x_{k+1})(y_1, \dots, y_k) = (x_1 y_1 x_{k+1}^{-1}, \dots, x_k y_k x_{k+1}^{-1}).$$

Then G^{k+1} acts on G^k transitively and effectively, and ΔG^{k+1} is the isotropy group at the identity $o = (e, \dots, e)$ of G^k . We get a diffeomorphism between G^k and the coset space $G^{k+1}/\Delta G^{k+1}$. Each tangent vector at the identity of G^k can be written in a unique way in the form (X_1, \dots, X_k) , where $X_1, \dots, X_k \in G_e$.

Now let Φ be an $\text{Ad}(G)$ -invariant inner product on G_e , and let $\Phi^{[k]}$ be the $\text{Ad}(\Delta G^{k+1})$ -invariant inner product on $(G^k)_o$ defined by

$$\begin{aligned} \Phi^{[k]}((X_1, \dots, X_k), (X_1, \dots, X_k)) \\ = \sum_{i=1}^k \Phi(X_i, X_i) + \sum_{i < j} \Phi(X_i - X_j, X_i - X_j). \end{aligned}$$

An alternative definition of $\Phi^{[k]}$ is the following: for $i = 1, \dots, k$ and $X \in G_e$ let $X^{(i)}$ denote the vector $(X_1, \dots, X_k) \in (G^k)_o$ such that $X_i = X$ and $X_j = 0$ for $j \neq i$. Then $\Phi^{[k]}(X^{(i)}, Y^{(i)}) = k\Phi(X, Y)$, and $\Phi^{[k]}(X^{(i)}, Y^{(j)}) = -\Phi(X, Y)$ for $i \neq j$.

The inner product $\bar{\Phi}^{[k]}$ can be extended, by the left translations of G^{k+1} , to a Riemannian metric on G^k denoted also by $\bar{\Phi}^{[k]}$. Then $G^{k+1}/\Delta G^{k+1}$ becomes a homogeneous Riemannian manifold $(G^k, \bar{\Phi}^{[k]})$.

Let σ be an automorphism of G^{k+1} defined by the rule $\sigma(x_1, \dots, x_{k+1}) = (x_{k+1}, x_1, \dots, x_k)$. Then σ satisfies all the conditions of Proposition 1, where we write $k + 1, G^{k+1}, \Delta G^{k+1}, G^k$ instead of k, G, H, M respectively. In particular, condition (iii) can be verified as follows: consider the transformation s of G^k determined by $\pi \circ \sigma = s \circ \pi$. Then for any $X \in G_e$ we deduce easily $s_{*0}(X^{(i)}) = X^{(i+1)}$ for $i = 1, \dots, k - 1, s_{*0}(X^{(k)}) = -(X^{(1)} + \dots + X^{(k)})$, and $\bar{\Phi}^{[k]}(s_{*0}X^{(i)}, s_{*0}Y^{(j)}) = \bar{\Phi}^{[k]}(X^{(i)}, Y^{(j)})$ for $i, j = 1, \dots, k$. Thus the Riemannian manifold $(G^k, \bar{\Phi}^{[k]})$ is $(k + 1)$ -symmetric.

3. In the remainder of this paper we shall specialize the class of manifolds $(G^k, \bar{\Phi}^{[k]})$ in a proper way.

Proposition 2. Consider a homogeneous Riemannian manifold $(G^k, \bar{\Phi}^{[k]})$ such that

- (a) G is simple,
- (b) G^{k+1} is the component of unity of the full isometry group $I(G^k, \bar{\Phi}^{[k]})$.

Then $(G^k, \bar{\Phi}^{[k]})$ is not l -symmetric for any $l < k + 1$.

Proof. Let r be an isometry of $(G^k, \bar{\Phi}^{[k]})$ with the isolated fixed point $o = (e, \dots, e)$ such that $r^l = \text{identity}$. Define an automorphism $\tilde{\rho}$ of the group $I(G^k, \bar{\Phi}^{[k]})$ by the formula $\tilde{\rho}(g) = r \circ g \circ r^{-1}$. Then the restriction of $\tilde{\rho}$ to G^{k+1} is an automorphism ρ of G^{k+1} . We can easily see that $\pi \circ \rho = r \circ \pi$.

Now G^{k+1} is a direct product of simple subgroups $G^{*(i)}, i = 1, \dots, k + 1$, all of them being canonically isomorphic to the group G . Then the automorphism $\rho: G^{k+1} \rightarrow G^{k+1}$ induces a permutation ν of the indices $1, \dots, k + 1$ such that $\rho(G^{*\nu(i)}) = G^{*(i)}, i = 1, \dots, k + 1$. Denoting by φ_i the restriction of ρ to $G^{*\nu(i)}$, we get $\rho(g_1, \dots, g_{k+1}) = (\varphi_1(g_{\nu(1)}), \dots, \varphi_{k+1}(g_{\nu(k+1)}))$. In particular, $\rho(g, \dots, g) = (\varphi_1(g), \dots, \varphi_{k+1}(g))$. Because $\rho(\Delta G^{k+1}) \subset \Delta G^{k+1}$, we obtain $\varphi_1 = \varphi_2 = \dots = \varphi_{k+1}$ under the canonical identification $G^{*(1)} = \dots = G^{*(k+1)} = G$, and therefore a unique automorphism $\varphi: G \rightarrow G$ such that $\rho(g_1, \dots, g_{k+1}) = (\varphi(g_{\nu(1)}), \dots, \varphi(g_{\nu(k+1)}))$. Denote by $d\rho$ (respectively, $d\varphi$) the induced automorphism of the Lie algebra \mathfrak{g}^{k+1} (respectively, \mathfrak{g}). Then $d\rho(X_1, \dots, X_{k+1}) = (d\varphi(X_{\nu(1)}), \dots, d\varphi(X_{\nu(k+1)})), X_1, \dots, X_{k+1} \in \mathfrak{g}$.

Now let us recall the following result by Borel and Mostow, [1].

Lemma. A semi-simple automorphism A of a nonsolvable Lie algebra \mathfrak{g} leaves fixed an element X such that $\text{ad } X$ is not nilpotent.

$d\varphi$ is a semi-simple automorphism of \mathfrak{g} because $(d\varphi)^l = \text{identity}$. Let $X \neq 0$ be a fixed vector of $d\varphi$ and suppose $l < k + 1$. Then the permutation ν contains a cycle (i_1, \dots, i_m) of length $m < k + 1$. Consider the vector $Z = (X_1, \dots, X_{k+1}) \in \mathfrak{g}^{k+1}$ such that $X_i = X$ for $i = i_1, \dots, i_m$ and $X_i = -X$ otherwise. Clearly, $d\rho(Z) = Z$. Now we can identify \mathfrak{g}^{k+1} with the tangent space $(G^{k+1})_e$ and $d\rho$ with the tangent map ρ_{*e} . We have $\pi_{*e} \circ \rho_{*e} = r_{*0} \circ \pi_{*e}$, and thus the projection $\pi_{*e}(Z) \in (G^k)_0$ is a fixed vector with respect to r_{*0} .

Moreover, $Z \in (G^{k+1})_e$ is not tangent to the submanifold ΔG^{k+1} and hence $\pi_{*e}(Z) \neq 0$, a contradiction. This completes the proof.

Proposition 3. For $G = SO(3)$ and $\Phi(X, Y) = -\frac{1}{2} \text{tr}(\text{ad } X \circ \text{ad } Y)$ the conditions of Proposition 2 are satisfied.

Proof. In the following, the elements of \mathfrak{g} (respectively, \mathfrak{g}^k) are considered as left invariant vector fields on G (respectively, G^k). First of all, there is a basis $\{X_1, X_2, X_3\}$ of \mathfrak{g} such that $[X_1, X_2] = X_3$, $[X_2, X_3] = X_1$, $[X_3, X_1] = X_2$. We have $\Phi(X_\alpha, X_\beta) = \delta_{\alpha\beta}$ for $\alpha, \beta = 1, 2, 3$, and the vectors $X_\alpha^{(i)}$, $\alpha = 1, 2, 3$, $i = 1, \dots, k$, form a basis of \mathfrak{g}^k . Now recall formulas (14) of [3]: for $X, Y \in \mathfrak{g}$

$$\begin{aligned} \nabla_{X^{(i)}} Y^{(j)} &= \frac{1}{2(k+1)} \{ [X, Y]^{(j)} - [X, Y]^{(i)} \} \quad \text{for } i \neq j, \\ \nabla_{X^{(i)}} Y^{(i)} &= \frac{1}{2} [X, Y]^{(i)}. \end{aligned}$$

A routine calculation shows the following properties of the curvature tensor R of $\Phi^{[k]}$:

$$(1) \quad \begin{aligned} &R(X_\alpha^{(i)}, X_\beta^{(j)})X_\gamma^{(k)} = 0 \text{ whenever } \alpha \neq \beta \neq \gamma \text{ or } \alpha = \beta = \gamma, \\ &R(X_\alpha^{(i)}, X_\beta^{(j)})X_\alpha^{(k)} \text{ and } R(X_\alpha^{(i)}, X_\alpha^{(j)})X_\beta^{(k)} \text{ belong to the subspace} \\ &\text{generated by } X_\beta^{(i)}, X_\beta^{(j)}, X_\beta^{(k)}. \end{aligned}$$

Let H_0 be the component of the unity of the isotropy group of $I(G^k, \Phi^{[k]})$ at the origin o , and denote the corresponding Lie algebra by \mathfrak{h}_0 . Then \mathfrak{h}_0 has a faithful isotropy representation by endomorphisms of $\mathfrak{g}^k = (G^k)_o$. Clearly, the necessary condition for $A \in \mathfrak{h}_0$ is that $A(\Phi^{[k]}) = A(R) = 0$, where A acts as a derivation on the tensor algebra of \mathfrak{g}^k .

Let $A \in \mathfrak{h}_0$ and set

$$(2) \quad AX_\alpha^{(i)} = \sum_{\beta=1}^3 \sum_{j=1}^k a_{(j)\alpha}^{(i)\beta} X_\beta^{(j)}, \quad i = 1, \dots, k, \alpha = 1, 2, 3.$$

The relation $(A\Phi^{[k]})(X_\alpha^{(i)}, X_\beta^{(j)}) = 0$ implies

$$(3) \quad k(a_{(i)\alpha}^{(i)\beta} + a_{(j)\beta}^{(j)\alpha}) - \sum_{l \neq i} a_{(l)\alpha}^{(i)\beta} - \sum_{l \neq j} a_{(l)\beta}^{(j)\alpha} = 0.$$

Further, we can calculate easily

$$R(X_\alpha^{(i)}, X_\beta^{(i)})X_\alpha^{(i)} = -\frac{1}{4}X_\beta^{(i)} \quad \text{for } \alpha \neq \beta.$$

Consider the relation $(AR)(X_\alpha^{(i)}, X_\beta^{(i)})X_\alpha^{(i)} = 0$, i.e.,

$$(4) \quad \begin{aligned} -\frac{1}{4}AX_\beta^{(i)} &= R(AX_\alpha^{(i)}, X_\beta^{(i)})X_\alpha^{(i)} + R(X_\alpha^{(i)}, AX_\beta^{(i)})X_\alpha^{(i)} \\ &\quad + R(X_\alpha^{(i)}, X_\beta^{(i)})AX_\alpha^{(i)}. \end{aligned}$$

Let us substitute (2) in (4) and consider a vector $X_\gamma^{(j)}$, where $\gamma \neq \alpha, \beta$ and

$j \neq i$. This vector enters into the left-hand side with the coefficient $-\frac{1}{4}a_{(j)\beta}^{(i)\gamma}$. According to (1), there is only one term on the right-hand side the evaluation of which can involve $X_r^{(j)}$, namely, the term $R(X_\alpha^{(i)}, a_{(j)\beta}^{(i)\gamma}X_r^{(j)})X_\alpha^{(i)}$. Now

$$R(X_\alpha^{(i)}, a_{(j)\beta}^{(i)\gamma}X_r^{(j)})X_\alpha^{(i)} = a_{(j)\beta}^{(i)\gamma}[(k + 2)X_r^{(i)} - X_r^{(j)}]/[4(k + 1)^2].$$

Comparing the coefficients at $X_r^{(j)}$ we finally get $a_{(j)\beta}^{(i)\gamma} = 0$. Thus we have proved

$$(5) \quad a_{(j)\beta}^{(i)\alpha} = 0 \quad \text{for } i \neq j, \alpha \neq \beta.$$

Substituting in (3) we get

$$(6) \quad a_{(i)\alpha}^{(i)\beta} + a_{(j)\beta}^{(j)\alpha} = 0 \quad \text{for } \alpha \neq \beta.$$

In particular, for $i = j$ we obtain

$$(7) \quad a_{(i)\alpha}^{(i)\beta} + a_{(i)\beta}^{(i)\alpha} = 0,$$

and hence

$$(8) \quad a_{(1)\alpha}^{(1)\beta} = a_{(2)\alpha}^{(2)\beta} = \dots = a_{(k)\alpha}^{(k)\beta} \quad \text{for } \alpha \neq \beta.$$

Now let us compare the coefficients at $X_\beta^{(j)}$, $j \neq i$, in the relation (4). $X_\beta^{(j)}$ enters into the left-hand side with the coefficient $-\frac{1}{4}a_{(j)\beta}^{(i)\beta}$. As for the right-hand side, $X_\beta^{(j)}$ can be involved only in the evaluations of the terms $R(a_{(j)\alpha}^{(i)\alpha}X_\alpha^{(j)}, X_\beta^{(i)})X_\alpha^{(i)}$, $R(X_\alpha^{(i)}, a_{(j)\beta}^{(i)\beta}X_\beta^{(j)})X_\alpha^{(i)}$, $R(X_\alpha^{(i)}, X_\beta^{(i)})(a_{(j)\alpha}^{(i)\alpha}X_\alpha^{(j)})$. After routine calculations we obtain

$$(9) \quad (3k + 2)a_{(j)\alpha}^{(i)\alpha} + (k^2 + 2k)a_{(j)\beta}^{(i)\beta} = 0.$$

Writing these relations for $(\alpha, \beta) = (1, 2), (2, 3), (3, 1)$ respectively, we obtain finally

$$(10) \quad a_{(j)\alpha}^{(i)\alpha} = 0 \quad \text{for } i \neq j, \alpha = 1, 2, 3.$$

Having $i = j$ and $\alpha = \beta$ in (3), we deduce from (10)

$$(11) \quad a_{(i)\alpha}^{(i)\alpha} = 0, \quad \alpha = 1, 2, 3, i = 1, \dots, k.$$

If we summarize (5), (10) and then (7), (8), (11), we can see that $\mathfrak{h}_0 \subset \Delta([\mathfrak{so}(3)]^k)$. On the other hand, the group $G^{k+1} = SO(3)^{k+1}$ is contained in $I(G^k, \Phi^{[k]})$ so that $\Delta(SO(3)^{k+1})$ is contained in H_0 . Thus $H_0 = \Delta(SO(3)^{k+1})$, and consequently $SO(3)^{k+1}$ is the component of the unity of $I(G^k, \Phi^{[k]})$ as required. Hence we can conclude our paper with

Theorem. *For each integer $k \geq 2$ there exists a compact generalized symmetric Riemannian space (M, g) of order k such that the component of the unity of the full isometry group $I(M, g)$ is semi-simple.*

References

- [1] A. Borel & G. D. Mostow, *On semi-simple automorphisms of Lie algebras*, An. of Math. **61** (1955) 389–405.
- [2] O. Kowalski, *Riemannian manifolds with general symmetries*, Math. Z. **136** (1974) 137–150.
- [3] A. J. Ledger & M. Obata, *Affine and Riemannian s-manifolds*, J. Differential Geometry **2** (1968) 451–459.
- [4] J. A. Wolf & A. Gray, *Homogeneous spaces defined by Lie group automorphisms. I*, J. Differential Geometry **2** (1968) 77–114.

CHARLES UNIVERSITY, PRAGUE