

## ANTI-HOLOMORPHIC AUTOMORPHISMS OF THE EXCEPTIONAL SYMMETRIC DOMAINS

H. A. JAFFEE

### Introduction

A serious fault of the theory described in [8] and called "real forms of hermitian symmetric spaces" was the lack of information about the exceptional symmetric domains. This gap has been filled, and the new results are given here.

Let me now express my thanks to Professor Kuga for having posed the problems of [7], [8], and to Professors Borel, Helgason, and Langlands for several enlightening discussions about the present work. In particular, while the idea behind the "hard part" of Lemma (2.4) is my own, Borel is to be credited for adding the necessary rigor to my original argument. Various improvements in my original paper were suggested by the referee, especially the use of Theorem (2.10) in § 4.

### 1. The problem

Let  $X$  denote a hermitian symmetric space of noncompact type (for short, symmetric domain),  $x_0 \in X$  a base point,  $\mathcal{C}$  the set of anti-holomorphic involutive automorphisms of  $X$ , and  $\mathcal{C}_0 = \{\sigma \in \mathcal{C} \mid \sigma(x_0) = x_0\}$ . If  $G^h$  is the group of holomorphic automorphisms of  $X$ , and  $K^h$  the isotropy group at  $x_0$ , we have  $X \approx G^h/K^h$ . For any  $\sigma \in \mathcal{C}$ , we call  $X^\sigma = \{x \in X \mid \sigma(x) = x\}$  the *real form* of  $X$  associated to  $\sigma$ .  $G^h$  acts by conjugation on  $\mathcal{C}$ , and  $K^h$  preserves  $\mathcal{C}_0$ . We call the quotient  $\mathcal{C}/G^h$  the set of *complex conjugations* of  $X$ . Representing a conjugation by a  $\sigma \in \mathcal{C}$  (or even  $\mathcal{C}_0$  by Remark 2.3) we associate a real form to each conjugation. Another representative  $\sigma'$  for the same conjugation is  $G^h$ -conjugate to  $\sigma$ , hence the real form associated to a conjugation is well determined, up to isometry.

In Theorems (4.3) and (4.4) we give  $\mathcal{C}/G^h$  and the associated real forms for the two exceptional symmetric domains. The theorem in [8] on the conjugations of a symmetric domain without exceptional factor now applies with no restriction. It follows that, in general, distinct conjugations have nonisometric real forms; we know no *a priori* reason for this.

The next section (§ 2) applies more generally than just to the exceptional sym-

metric domains. Thus it also provides a sketch of a derivation of the (major) results of [8], although these results were first obtained by much more *ad hoc* procedures.

## 2. Preliminaries

**(2.0)** In § 1, we defined the set of complex conjugations  $\mathcal{C}/G^h$  of  $X$ ; the conjugations *fixing the base point*  $x_0$  are  $\mathcal{C}_0/K^h$ . There is a hermitian symmetric space  $X_u$  of compact type associated to  $X$  "by duality" (see [4] for details). Let  $G_u^h$  be the (compact) group of holomorphic isometries of  $X_u$ , and  $K_u^h$  the isotropy subgroup at some chosen base point of  $X_u$  ( $= G_u^h/K_u^h$ ),  $\mathcal{C}_u$  the set of anti-holomorphic involutive isometries of  $X_u$ ,  $\mathcal{C}_u^*$  the subset of  $\mathcal{C}_u$  with fixed points, and  $\mathcal{C}_u^0$  the subset fixing the base point. By results of Harish-Chandra and Borel [4, pp. 311–322], we may assume the following has been done. The domain  $X$  is holomorphically embedded as a bounded open subset of a  $\mathbf{C}$ -linear subspace  $p_-$  in the complexification of the Lie algebra of  $G^h$ , with the point  $x_0$  going to the origin, and so that the isometries of  $X$  fixing  $x_0$  are restrictions to  $X$  of  $\mathbf{R}$ -linear automorphisms of  $p_-$ . (Elements of  $K^h$  are  $\mathbf{C}$ -linear, and elements of  $\mathcal{C}_0$  conjugate linear.) We embed  $p_-$  as a Zariski open [2, § 4.3 (4)] in  $X_u$ , with  $x_0$  ( $= 0 \in p_-$ ) going to the base point of  $X_u$ , and isometries of  $X_u$  which fix  $x_0$  preserve the embedded  $p_-$  and the domain  $X$  inside it; we thus make identifications  $K^h = K_u^h$ ,  $\mathcal{C}_0 = \mathcal{C}_u^0$ .

We choose a  $\sigma_0 \in \mathcal{C}_0 = \mathcal{C}_u^0$ ; the group  $\text{Gal} = \{1, \sigma_0\}$  acts by conjugation on  $G^h$ , on  $K^h = K_u^h$ , and on  $G_u^h$ .  $G^h$  acts by conjugation on  $\mathcal{C}$ , and  $K^h$  preserves  $\mathcal{C}_0$ ;  $G_u^h$  acts by conjugation on  $\mathcal{C}_u$  and preserves the subsets  $\mathcal{C}_u^*$  and  $\mathcal{C}_u^{\text{def}'n} = \mathcal{C}_u - \mathcal{C}_u^*$  of  $\mathcal{C}_u$ , and  $K_u^h$  preserves the subset  $\mathcal{C}_u^0$ . An easy translation of the definitions (see, e.g., [9, p. I-56]) gives

**(2.1) Proposition.** *There are canonical identifications (after the choice of  $\sigma_0$ ):*

$$(2.1.1) \quad \mathcal{C}/G^h = H^1(\text{Gal}, G^h) ;$$

$$(2.1.2) \quad \mathcal{C}_0/K^h = H^1(\text{Gal}, K^h) ;$$

$$(2.1.3) \quad \mathcal{C}_u/G_u^h = H^1(\text{Gal}, G_u^h) ;$$

$$(2.1.4) \quad \mathcal{C}_u^*/G_u^h = \text{a subset } H^1_*(\text{Gal}, G_u^h) \text{ of } H^1(\text{Gal}, G_u^h) ;$$

$$(2.1.5) \quad \mathcal{C}_u^{\text{def}'n}/G_u^h = H^1_{\beta}(\text{Gal}, G_u^h) \underset{\text{def}'n}{=} H^1(\text{Gal}, G_u^h) - H^1_*(\text{Gal}, G_u^h) .$$

We will use the abbreviations  $H^1(G^h)$  for  $H^1(\text{Gal}, G^h)$ , etc.

**(2.2) Theorem.** *The diagrams below are identical by Proposition (2.1). The maps  $\iota_1$  and  $\iota_2$  are bijections. Hence  $\mathcal{C}/G^h$  is bijective to both  $\mathcal{C}_0/K^h$  and  $\mathcal{C}_u^*/G_u^h$ .*

$$(2.2.1) \quad \begin{array}{ccc} & & H_u^h(G_u^h) \subset H^1(G_u^h) \\ & \nearrow \iota_1 & \downarrow \parallel \\ H^1(K^h) & \xrightarrow{\approx} & \\ & \searrow \iota_2 & H^1(G^h) \end{array}$$

$$(2.2.2) \quad \begin{array}{ccc} & & \mathcal{C}_u^*/G_u^h \subset \mathcal{C}_u/G_u^h \\ & \nearrow \iota_1 & \downarrow \parallel \\ \mathcal{C}_0/K^h & \xrightarrow{\approx} & \\ & \searrow \iota_2 & \mathcal{C}/G^h \end{array}$$

(2.3) **Remark.** The proof of Theorem (2.2) requires a lemma, but first we note several things. If  $\sigma$  has a fixed point  $x$ , and  $gx = x_0$ , then  $g\sigma g^{-1}$  fixes  $x_0$ . The surjectivity of  $\iota_1$  and  $\iota_2$  then follows from the fact that  $G^h$  and  $G_u^h$  are transitive on  $X$  and  $X_u$ , respectively (and is a restatement of the fact that a  $\sigma \in \mathcal{C}$  or  $\mathcal{C}_u^*$  is  $G^h$ - or  $G_u^h$ -conjugate to a  $\sigma' \in \mathcal{C}_0$ ).

(2.4) **Lemma.** *Let  $\sigma \in \mathcal{C}_0$  or  $\mathcal{C}_u^*$ . Then the fixed point sets  $X^\sigma$  and  $X_u^\sigma$  of  $\sigma$  on  $X$  and  $X_u$  are connected.*

*Proof.* For  $X$ , let  $x_1$  and  $x_2$  be fixed by  $\sigma$ , and  $\gamma$  the unique geodesic segment joining  $x_1$  and  $x_2$ . Since  $\sigma$  is an isometry  $\gamma$  must be preserved, and since the endpoints are fixed,  $\gamma$  is fixed pointwise.

For  $X_u$ , Remark (2.3) says that without loss of generality we may assume  $\sigma \in \mathcal{C}_0$ ; as explained above,  $\sigma$  preserves a Zariski open set, say  $C$ , centered at  $x_0$  and isomorphic to  $\mathbf{C}^n$ . The restriction of  $\sigma$  to  $C$  decomposes as a direct sum  $(+1) \oplus (-1)$ , where the  $(+1)$ -eigenspace  $D$  is isomorphic to  $\mathbf{R}^n$ , and Zariski dense in  $C$ . Let  $x_1$  be any fixed point of  $\sigma$  in  $X_u$ , outside of  $C$ .  $\sigma$  preserves another Zariski open  $C'$ , centered at  $x_1$  and isomorphic to  $\mathbf{C}^n$ . Let  $D'$  be the fixed point set of  $\sigma$  in  $C'$ . We have  $D \cup D' \subset x_u^\sigma$ , and will show that  $D \cap D'$  is non-empty. But  $C \cap C' = U$  is Zariski open in, say,  $C$ . Therefore  $D \cap U$  is non-empty; hence  $D \cap C'$  and also  $D \cap D'$ . This proves the lemma.

*Proof of Theorem (2.2).* We have already the surjectivity of  $\iota_1$  and  $\iota_2$ . We shall prove the injectivity of  $\iota_1$ , the proof for  $\iota_2$  being completely analogous. By [9, Cor. 1, pp. 1-65],  $\ker(\iota_1)$  may be identified with the quotient of  $(X_u)^\sigma$  by  $(G_u^h)^\sigma =$  the centralizer of  $\sigma_0$  in  $G_u^h$ . We have  $x_0 \in (X_u)^\sigma$ ; let  $x_1 \in (X_u)^\sigma$  and (using Lemma (2.4)) let  $\gamma$  be a geodesic (in  $(X_u)^\sigma$ ) from  $x_0$  to  $x_1$ . By [4, Th. 3.3, p. 173] we identify  $\gamma$  with a ray in the Lie algebra of  $(G_u^h)^\sigma$ , orthogonal to that of  $K_u^h$ , and by exponentiation obtain an element of  $(G_u^h)^\sigma$  which transforms  $x_0$  to  $x_1$ . This shows  $\ker(\iota_1)$  is trivial. Now  $\iota_1$  is also the canonical map  $\mathcal{C}_0/K^h \rightarrow \mathcal{C}_u^*/G_u^h$ , so we may repeat the argument for any other  $\sigma \in \mathcal{C}_0$  (applying Lemma (2.4) to each) to get that all fibres of  $\iota_1$  are trivial. This proves Theorem (2.2).

The following theorem of de Siebenthal implies that there are only a finite number of real forms of a symmetric domain.

(2.5) **Lemma (de Siebenthal)** [10, pp. 57-58]. *Let  $G$  be a compact Lie group,  $G_0$  the identity component,  $\sigma_0 \in G$ , and  $T$  a maximal torus of the centralizer  $(G_0)^{\sigma_0}$*

of  $\sigma_0$  in  $G_0$ . Given any  $\sigma$  in the component of  $G$  containing  $\sigma_0$ , there are a  $t \in T$  and a  $g \in G_0$  with  $\sigma = g(\sigma_0 t)g^{-1}$ .

**(2.6) Remark.** If  $\sigma_0$  and  $\sigma$  are involutions, then  $t^2 = 1$ .

**(2.7) Theorem.** The set  $\mathcal{C}_u/G_u^h = H^1(G_u^h)$  is finite.

*Proof.* Let  $G$  be the compact group generated by  $G_u^h$  and  $\sigma_0$ . If  $\sigma \in \mathcal{C}_u$  is in the same component of  $G$  as  $\sigma_0$ , then by Lemma (2.5)  $\sigma$  is  $G_0$ -conjugate, and therefore  $G_u^h$ -conjugate, to some  $\sigma_0 t$ , and  $t^2 = 1$  by (2.6). If  $l = \text{rk}(T)$  with  $T$  defined in Lemma (2.5), then there are only  $2^l$  such elements  $t$ . If  $G$  has only two components, then we would have  $\text{card}(\mathcal{C}_u/G_u^h) \leq 2^l$ . If not, let  $\sigma_1 \in \mathcal{C}_u$  be an involution in a different component of  $G$  than  $\sigma_0$ . Then  $G$  is also generated by  $G_u^h$  and  $\sigma_1$ , and we can apply the above argument to the component of  $\sigma_1$ . Clearly, this process will bound  $\text{card}(\mathcal{C}_u/G_u^h)$  by a (finite) sum of powers of 2, and finishes the proof.

**(2.8) Corollary.** The number of complex conjugations  $c(X) = \text{card}(\mathcal{C}/G^h)$  of a symmetric domain  $X$  is finite.

*Proof.* By Theorem (2.2),  $\mathcal{C}/G^h$  is isomorphic to the subset  $\mathcal{C}_u^*/G_u^h$  of  $\mathcal{C}_u/G^h$ . We will make extensive use of the classification of symmetric spaces, together with Theorem 2.10 below, to determine the isometry types of the real forms  $X^\sigma$ . Similar techniques were used in [5].

For now, let  $X$  denote a hermitian symmetric space which is purely non-Euclidean,  $\sigma$  an isometry,  $X^\sigma$  the set of fixed points of  $\sigma$ , and let  $x_0 \in X^\sigma$ . Let  $G(X)$  be the identity component of the isometry group of  $X$ ,  $K(X) \subset G(X)$  the isotropy subgroup at  $x_0$ , and  $G(X)^\sigma$  and  $K(X)^\sigma$  the respective centralizers of  $\sigma$ . Let  $G(X^\sigma)$  and  $K(X^\sigma)$  be the full isometry and isotropy groups for  $X^\sigma$ , with base point  $x_0$ . Denote by  $\mathfrak{G}(X)$ ,  $\mathfrak{K}(X)$ , etc. the Lie algebras of  $G(X)$ ,  $K(X)$ , etc.

If  $g \in G(X)^\sigma$  and  $y \in X^\sigma$ , then  $g(y) \in X^\sigma$ . By restriction, we associate an isometry  $\rho(g)$  of  $X^\sigma$  to  $g$ . Denote by  $\rho: G(X)^\sigma \rightarrow G(X^\sigma)$  and  $d\rho: \mathfrak{G}(X)^\sigma \rightarrow \mathfrak{G}(X^\sigma)$  the homomorphisms thus defined. If  $s \in G(X)$  is the symmetry in  $X$  at a point  $y \in X^\sigma$ , then  $\sigma s \sigma^{-1}$  has differential  $-1$  at  $y$ ; thus  $s \in G(X)^\sigma$ , and  $\rho(s)$  is a geodesic reflection for  $X^\sigma$  at  $y$ . This shows that  $X^\sigma$  itself is a globally symmetric space, and a symmetric subspace of  $X$ .

**(2.9) Proposition.** With the above notation, the pair  $(d\rho(\mathfrak{G}(X)^\sigma), d\rho(\mathfrak{K}(X)^\sigma))$  is an orthogonal involutive Lie subalgebra (in the sense of [11, p. 235]) of the orthogonal involutive Lie algebra of the component of  $X^\sigma$  containing  $x_0$ .

*Proof.* The point is that the orbit of  $x_0$  under  $G(X)^\sigma$  is the entire component of  $X^\sigma$  containing  $x_0$ . This follows from [4, Theorem 3.3] as in the proof of Theorem (2.2). If  $X$  is of noncompact type, this fact is [12, Theorem 2.4.1]; the author is thankful to the referee for pointing this out.

Now if  $\sigma$  is an antiholomorphic involution,  $X^\sigma$  is connected by Lemma (2.4). Moreover  $\dim_{\mathbb{R}} X^\sigma = \dim_{\mathbb{C}} X$  since multiplication by the complex structure  $J$  of  $X$ , along  $X^\sigma$ , defines an isomorphism between the tangent and normal bundles to  $X^\sigma$ . For the same reason, a holomorphic isometry of  $X$  is the identity if its restriction to  $X^\sigma$  is the identity. This implies that the maps  $\rho$  and  $d\rho$  above are

injective, as far as we are concerned in this paper.

**(2.10) Theorem.** *Let  $X$  be a hermitian symmetric space as above, and  $\sigma$  an antiholomorphic involutive isometry. If  $\mathfrak{G}(X)^\sigma$  is semi-simple, or of the form  $\mathbf{R}^1 \times$  semi-simple, then the orthogonal Lie algebra of  $X^\sigma$  is isomorphic to  $(\mathfrak{G}(X)^\sigma, \mathfrak{K}(X)^\sigma)$ .*

*Proof.* By injectivity of  $d\rho$ , the subalgebra in Proposition (2.9) is isomorphic to  $(\mathfrak{G}(X)^\sigma, \mathfrak{K}(X)^\sigma)$ . In the semi-simple case, [11, Lemma 8.2.3] says that this subalgebra is maximal, which gives the required statement. In the other case, the classification of orthogonal Lie algebras [11, Theorem 8.2.4], and of the Euclidean ones [11, Theorem 8.2.10], gives the same thing.

### 3. Descriptions of EIII and EVII

There are two irreducible hermitian symmetric spaces with exceptional isometry groups; they are the Riemannian symmetric spaces EIII and EVII of (complex) dimensions 16 and 27. Their Lie algebras are given [4, p. 354] in the noncompact form as  $(e_{6(-14)}, so(10) + \mathbf{R})$  and  $(e_{7(-25)}, e_6 + \mathbf{R})$ , and the respective ranks are 2 and 3. The groups  $K^h$  and  $G^h$  are connected in each case [3, Théorème H].

The space EVII is described in [1, pp. 525–527] as a symmetric subspace of  $\mathfrak{h}_{28}$ , the Siegel space of genus 28. The action of the group  $G^h$  is given there. The same picture is given in [5], where a bounded version (cf. § 2.0 above) is also given. We will denote this bounded symmetric domain by  $Z$ . The group  $K^h$  for  $Z$  is described using Jordan algebras. If  $\alpha$  is the exceptional central simple reduced (nondivision) Jordan algebra ([6, p. 80] for the definition) of dimension 27 over  $\mathbf{R}$ ,  $\text{Aut}(\alpha)$  is the compact Lie group  $F_4$ . Then  $K^h$  is a subgroup of  $GL(\alpha \otimes \mathbf{C})$ :  $K^h = \{\exp(iL(x)) \cdot V \mid V \in \text{Aut}(\alpha) \text{ and } L(x) \text{ is left multiplication by an } x \in \alpha\}$ . The center  $T^1$  of  $K^h$  is  $\{\exp(iL(\xi)) \mid 1 = \text{identity of } \alpha, \xi \in \mathbf{R}\}$ , and  $K^h = T^1 \cdot E_6$  (see [11, p. 315]); the intersection  $T^1 \cap E_6 = (\varepsilon)$  is the center of  $E_6$ , which is cyclic of order 3 [11, Cor. 8.9.28] generated by, say,  $\varepsilon$ .

The space EIII is a holomorphic subdomain of  $Z$ . Let, for example,  $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

be a (primitive) idempotent of  $\alpha$  (see [5], [6] for details). The Peirce-decomposition  $\alpha = \alpha_0 + \alpha_1 + \alpha_{1/2}$  is associated, which is the eigenspace decomposition for the idempotent. Define  $a \in F_4 \subset K^h$  as

$$+1 \text{ on } \alpha_0 + \alpha_1, \quad \text{and } -1 \text{ on } \alpha_{1/2}.$$

Let  $\iota$  denote the geodesic reflection of  $Z$  at  $0 \in \alpha \otimes \mathbf{C}$ ;  $\iota(z) = -z$ . The space  $Y$  of fixed points of  $\iota \circ a$  on  $Z$  is shown in [5] to be isomorphic to the hermitian symmetric space EIII.

For determining  $\mathcal{C}/G^h$  and the associated real forms for the two symmetric domains described above, we shall use an explicitly constructed element  $\sigma_0 \in \mathcal{C}(Z)$ . Define  $\sigma_0$  by  $z \mapsto -\bar{z}$  in the nonbounded versions of [1] and [5] and by

$z \mapsto \bar{z}$  in the bounded version  $Z$ . If  $0 \in Z$  denotes the base point referred to above, we have  $\sigma_0(0) = 0$ . In the notation of [1],  $\sigma_0$  is represented by the matrix  $\left( \begin{array}{c|c} 1_{28} & 0 \\ \hline 0 & -1_{28} \end{array} \right)$ . Conjugation by  $\sigma_0$  on  $K^h$  is the involutive automorphism  $\pi(\sigma_0)$ :

$$\exp(iL(x)) \cdot V \mapsto \exp(-iL(x)) \cdot V,$$

so the centralizer  $(K^h)^{\sigma_0}$  has identity component  $F_4$ .

#### 4. Computations of $\mathcal{C}/G^h$ and the real forms

We first compute the set of complex conjugations  $\mathcal{C}/G^h$  for the space  $Z$  (cf. § 3) of type EVII. By (2.2.2), this set is bijective to  $\mathcal{C}_0/K^h$ .  $\mathcal{C}_0$  is the set of elements of order 2 in the component  $K^h \cdot \sigma_0$  of  $K = K^h \cup K^h \cdot \sigma_0$ , where  $\sigma_0 \in \mathcal{C}_0$  is defined at the end of § 3.  $\mathcal{C}_0/K^h$  is thus the set of  $K^h$ -conjugacy classes of elements of order 2 in  $K^h \cdot \sigma_0$ . The automorphism  $\pi(\sigma_0)$  of  $K^h = T^1 \cdot E_6$  leaves the “factors” invariant. Thus the subset  $\tilde{A} = E_6 \cup E_6 \cdot \sigma_0$  of  $K$  is a subgroup. Define  $\pi: \tilde{A} \rightarrow \text{Aut}(E_6)$  through the adjoint action (hence the symbolism  $\pi(\sigma_0)$ ).  $\text{Aut}(E_6)/\text{Inn}(E_6) \approx Z/2Z$  by [11, Cor. 8.11.3], and  $\pi(\sigma_0)$  fixes a subgroup ( $F_4$ ) of nonmaximal rank in  $E_6$ , so  $\pi(\sigma_0)$  is an “outer” automorphism. Hence  $\pi$  is surjective, with  $\ker(\pi) = (\varepsilon)$ .

**(4.1) Proposition.** *Any element  $\lambda e \cdot \sigma_0 \in K^h \cdot \sigma_0$  ( $\lambda \in T^1, e \in E_6$ ) is  $K^h$ -conjugate to  $e \cdot \sigma_0$ .*

*Proof.* Conjugate  $\lambda e \cdot \sigma_0$  by any  $\sqrt{\lambda}^{-1}$  in  $T^1$ . Then use  $\sigma_0 \sqrt{\lambda} = \sqrt{\lambda}^{-1} \sigma_0$ .

**(4.2) Proposition.** *The elements  $\sigma_1 = e_1 \cdot \sigma_0$  and  $\sigma_2 = e_2 \cdot \sigma_0$  are  $K^h$ -conjugate if  $\pi(\sigma_1)$  and  $\pi(\sigma_2)$  are ad  $(E_6)$ -conjugate in  $\text{Aut}(E_6)$ .*

*Proof.* Suppose  $\exists a \in \text{ad}(E_6)$  with  $\pi(\sigma_1) = a\pi(\sigma_2)a^{-1}$ . Lift  $a$  to any  $\tilde{a} \in \pi^{-1}(a)$  and obtain  $\sigma_1 \varepsilon^i = \tilde{a} \sigma_2 \tilde{a}^{-1}$  for some  $\varepsilon^i \in (\varepsilon)$ . Now apply Proposition (4.1) to  $\sigma_1 \varepsilon^i = \varepsilon^{-i} \sigma_1$  with  $\lambda = \varepsilon^{-i}$ .

Computing  $\mathcal{C}/G^h$  now reduces to finding the ad  $(E_6)$ -conjugacy classes of outer involutive automorphisms of  $E_6$ . These are determined in [11, p. 288] as part of the classification of symmetric spaces of  $E_6$ . Each such automorphism is ad  $(E_6)$ -conjugate to either  $\pi(\sigma_0)$  or another automorphism, say  $\pi(\sigma_1)$ , and the Lie algebras of the centralizers in  $E_6$  of  $\sigma_0$  and  $\sigma_1$  are<sup>1</sup> respectively  $f_4$  and  $sp(4)$ . By Proposition (4.2),  $\sigma_0$  and  $\sigma_1$  represent all conjugations of  $Z$ ; strictly speaking as yet they might be  $K^h$ -conjugate. We will find that the associated real forms are nonisometric, hence  $\sigma_0$  and  $\sigma_1$  are not  $G^h$ -conjugate.

As in § 2, extend each of  $\sigma_0$  and  $\sigma_1$  to the compact dual  $Z_u$  of  $Z$ . Let  $Z_u^{\sigma_i}$  ( $i = 0, 1$ ) be the respective fixed point sets. The centralizers  $G(Z_u)^{\sigma_i}$  are such that the quotients  $G(Z_u)/G(Z_u)^{\sigma_i}$  are symmetric spaces of  $G(Z_u) = \text{ad}(E_7)$ . By classification of these [11, table, p. 285] the only possibilities for  $\mathfrak{G}(Z_u)^{\sigma_i}$  are

<sup>1</sup>The hidden fact here is that  $\exists \sigma_1 \in \pi^{-1}(\pi(\sigma_1))$  with  $\sigma_1^2 = 1$ .  $F_4 \subset \tilde{A}$ , and  $F_4 \subset$  centralizer of  $\sigma_0$ , so  $\pi|_{F_4}$  is an isomorphism. By [11, p. 288],  $\pi(\sigma_1) = \pi(\varphi) \circ \pi(\sigma_0)$  with  $\pi(\varphi) \in \pi(F_4)$  and  $\pi(\varphi)^2 = 1$ . The unique lift  $\varphi \in \pi^{-1}(\pi(\varphi))$  then satisfies  $\varphi^2 = 1$ ,  $\varphi \sigma_0 = \sigma_0 \varphi$ . Now  $\sigma_1 = \varphi \circ \sigma_0$  is as required.

(1)  $\mathbf{R} \times e_6$ , (2)  $su(8)$ , (3)  $so(12) \times su(2)$ .

By Theorem (2.10), the orthogonal Lie algebra  $\mathfrak{G}(Z_u^{\sigma_i})$  is isomorphic to one of these. Since  $\dim Z_u^{\sigma_i} = 27$ ,  $\text{rk}(Z_u^{\sigma_i}) \leq 3$ , and  $\mathfrak{K}(Z_u^{\sigma_i}) \approx f_4$  or  $sp(4)$  ( $i = 0, 1$ ), we see that the classification of symmetric spaces with Lie algebras (1), (2), or (3) uniquely determines  $\mathfrak{G}(Z_u^{\sigma_i})$ . Namely,  $\mathfrak{G}(Z_u^{\sigma_0}) = (\mathbf{R} \times e_6, f_4)$ ,  $\mathfrak{G}(Z_u^{\sigma_1}) = (su(8), sp(4))$ . By duality, we get the real forms  $Z^{\sigma_0}$  and  $Z^{\sigma_1}$ .

**(4.3) Theorem.** *There are 2 complex conjugations of the symmetric domain  $Z = (e_{7(-26)}, e_6 + \mathbf{R})$ . The associated real forms are  $\mathbf{R} \times (e_{6(-26)}, f_4)$ , and  $(su^*(8), sp(4))$  both of rank 3.*

We next compute the set  $\mathcal{C}/G^h$  for the domain  $Y$  (see § 3) of type EIII. Let  $Y_u$  be the compact dual. By (2.2.2) we have  $\mathcal{C}/G^h \approx \mathcal{C}_u^*/G_u^h$ . It is shown in [11, p. 316] that every isometry of  $Y_u$  has a fixed point; thus  $\mathcal{C}_u^* = \mathcal{C}_u$ . Now  $\mathcal{C}/G^h \approx \mathcal{C}_u/G_u^h$ . The identity component  $Is_0$  of the isometry group  $Is$  of  $Y_u$  is isomorphic to  $\text{ad}(E_6)$  by [11, Theorem 8.7.9].  $Is(Y_u)$  has 2 components [3, Theorem H] and some element of  $Is - Is_0$  gives rise to an outer automorphism of  $E_6$  [11, p. 316]. Hence there is an isomorphism  $\psi: \text{Aut}(E_6) \cong Is(Y_u)$ . Now  $\psi$  defines an isomorphism between  $\mathcal{C}_u/G_u^h$  and the  $\text{ad}(E_6)$ -conjugacy classes of outer involutive automorphisms of  $E_6$ . As quoted before, the latter are represented by  $\pi(\sigma_0)$  and  $\pi(\sigma_1)$ , so that  $\tau_0 = \psi(\pi(\sigma_0))$  and  $\tau_1 = \psi(\pi(\sigma_1))$  represent the complex conjugations of  $Y$  under  $\mathcal{C}_u/G_u^h \cong \mathcal{C}/G^h$ .

We will find the associated real forms on  $Y$  by working on  $Y_u$  and then dualizing. The centralizers  $\mathfrak{G}(Y_u)^{\tau_i}$  ( $i = 0, 1$ ) are isomorphic to  $f_4$  and  $sp(4)$ . Hence by Theorem (2.10) the orthogonal Lie algebra  $\mathfrak{G}(Y_u^{\tau_i})$  is isomorphic to  $f_4$  and  $sp(4)$  for  $i = 0, 1$ . The classification of these orthogonal involutive Lie algebras, together with  $\dim Y_u^{\tau_i} = 16$ ,  $\text{rk}(Y_u^{\tau_i}) \leq 2$ , again uniquely determines  $\mathfrak{G}(Y_u^{\tau_i})$ . Namely,  $\mathfrak{G}(Y_u^{\tau_0}) = (f_4, so(9))$ ,  $\mathfrak{G}(Y_u^{\tau_1}) = (sp(4), sp(2) \times sp(2))$ . By duality, we have

**(4.4) Theorem.** *There are 2 complex conjugations of the symmetric domain  $Y = (e_{6(-14)}, so(10) + \mathbf{R})$ . The associated real forms are  $(f_{4(-20)}, so(9))$  of rank 1, and  $(sp(2, 2), sp(2) \times sp(2))$  of rank 2.*

## 5. Retrospections

One of the conjugations of  $Z$  is represented by the involution  $\sigma_0$  which was defined explicitly in § 3. We would like explicit definitions also for  $\sigma_1$ ,  $\tau_0$ ,  $\tau_1$  whose real forms are given in Theorems (4.3) and (4.4). For  $Z$ ,  $F_4$  is a subgroup of  $K^h$ . An involution  $\beta \in F_4$  (the "quaternion" case) is defined in [6, Theorem 13];  $\beta$  and  $\sigma_0$  commute. The author suspects that  $\beta \circ \sigma_0$  represents the same conjugation of  $Z$  as  $\sigma_1$ .

$\sigma_0$  commutes with the defining isometry  $\iota \circ \alpha$  of  $Y$ , hence leaves  $Y$  invariant. It is not difficult to show using [6] that the restriction of  $\sigma_0$  to  $Y$  represents the same conjugation as  $\tau_0$ . The author suspects that  $\beta \circ \sigma_0$  also leaves  $Y$  invariant, and that the restriction is conjugate to  $\tau_1$ .

## References

- [ 1 ] W. L. Baily, Jr., *An exceptional arithmetic group and its Eisenstein series*, Ann. of Math. **91** (1970) 512–549.
- [ 2 ] A. Borel & J. Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. No. 27 (1965) 55–150.
- [ 3 ] E. Cartan, *Sur les domaines bornés homogènes de l'espace de  $n$  variables complexes*, Abh. Math. Sem. Univ. Hamburg **11** (1935) 116–162.
- [ 4 ] S. Helgason, *Differential geometry and symmetric spaces*, Academic Press, New York, 1962.
- [ 5 ] U. Hirzebruch, *Über Jordan-Algebren und beschränkte symmetrische Gebiete*, Math. Z. **94** (1966) 387–390.
- [ 6 ] N. Jacobson, *Some groups of transformations defined by Jordan algebras. II*, J. Reine Angew. Math. **204** (1960) 74–98.
- [ 7 ] H. Jaffee, *Real forms of hermitian symmetric spaces and real algebraic varieties*, Thesis, State University of New York at Stony Brook, 1974.
- [ 8 ] —, *Real forms of hermitian symmetric spaces*, Bull. Amer. Math. Soc. **81** (1975), 456–458.
- [ 9 ] J. P. Serre, *Cohomologie galoisienne*, Lecture Notes in Math. Vol. 5, Springer, Berlin, 1965.
- [ 10 ] J. de Siebenthal, *Sur les groupes de Lie compacts non connexes*, Comment. Math. Helv. **31** (1956) 41–89.
- [ 11 ] J. A. Wolf, *Spaces of constant curvature*, McGraw-Hill, New York, 1967.
- [ 12 ] V. Ozols, *Critical points of the displacement function of an isometry*, J. Differential Geometry **3** (1969) 411–432.

INSTITUTE FOR ADVANCED STUDY