

## VECTOR FIELDS OF A FINITE TYPE G-STRUCTURE

A. M. AMORES

### 0. Introduction

Let  $M$  be a connected manifold,  $g$  a Riemannian metric on  $M$ , and  $\mathcal{F}$  either the set of Killing vector fields or the set of conformal vector fields. The following theorems are known.

**(0.1) Theorem.** *If  $U \subset M$  is open and  $X, Y \in \mathcal{F}$ , then  $X|_U = Y|_U$  implies  $X = Y$  on the whole of  $M$ .*

**(0.2) Theorem.** *If  $M$  and  $g$  are analytic,  $M$  is simply connected, and  $X$  is a Killing (resp. conformal) field on  $U$ , open subset of  $M$ , then there is a unique extension of  $X$  to an analytic Killing (resp. conformal) field defined on the whole of  $M$ .*

These theorems were proved in [4] for the Killing case and in [3] for the conformal case. The aim of this paper is to generalize them, when  $\mathcal{F}$  is taken to be set of vector fields of a finite type  $G$ -structure. The precise definitions and statements of the theorems are in §2 and §3. §4 is devoted to proving some auxiliary results on fields on a parallelisable manifold. When no precision is made about the differentiability class of a manifold or map, it will be understood that the definition or result works for both the category of manifolds of class infinity and real analytic manifolds.

### 1. Parallelism fields

Let  $m = \dim M$ , and  $\pi$  be a parallelism on  $M$ ; that is, a 1-exterior form on  $M$  with values in  $R^m$  such that for all  $x \in M$ ,  $\pi(x) : TM(x) \rightarrow R^m$  is an isomorphism. Suppose that  $X$  is a vector field on  $M$ , and  $\{\psi_t : t \in R\}$  the corresponding pseudogroup of diffeomorphisms. Then we say that  $X$  is a parallelism field if for all  $t \in R$ ,  $\psi_t^* \pi = \pi$ , or, equivalently, if  $L_X \pi = 0$ . Let  $(u^1, \dots, u^m)$  be a coordinate system on  $U$ . If  $X$  is a field on  $U$  and  $c : I \rightarrow U$

---

Received November 22, 1976. The author would like to thank Dr. A. J. Ledger for his kindness and help in writing this paper.

is a smooth curve,  $I$  being an interval, we write

$$X = \sum X^i \frac{\partial}{\partial u^i}, \quad c^i(t) = u^i \circ c(t) \quad \text{for } t \in I, 1 \leq i \leq m,$$

$$\pi = \sum_{i,j} (\pi_j^i du^j) e_i,$$

where  $\{e_1, \dots, e_m\}$  is the canonical basis of  $R^m$ .

**(1.1) Lemma.** *If  $X$  is a parallelism field on  $U$ , then the curve  $t \rightarrow (X^1 \circ c(t), \dots, X^m \circ c(t))$  is a solution of the linear system*

$$\frac{dx^i}{dt} = \sum_j \mathcal{Q}_j^i(t) x^j \quad 1 \leq i \leq m,$$

where

$$\mathcal{Q}_j^i(t) = - \sum_{h,k} \rho_h^i(c(t)) \frac{\partial \pi_k^h(c(t))}{\partial u^j} \frac{dc^k(t)}{dt},$$

and  $(\rho_j^i(x))$  is the inverse matrix of  $\pi_j^i(x)$  for all  $x \in U$ .

*Proof.* It is just an easy computation, if we write the equation  $L_X \pi \cdot c'(t) = 0$  in local coordinates,  $c'(t) \in TM(c(t))$  being the velocity of  $c$  at the point  $t$ .

**(1.2) Lemma.** *If  $X, Y$  are parallelism fields on  $M$ , and  $X(x_0) = Y(x_0)$  for some  $x_0 \in M$ , then  $X = Y$  on  $M$ .*

*Proof.* Let  $x_1$  be an arbitrary point of  $M$ , and  $c: [0, 1] \rightarrow M$  a smooth curve such that  $c(0) = x_0, c(1) = x_1$ . We prove that  $X = Y$  on  $c([0, 1])$ ; hence  $X(x_1) = Y(x_1)$ . Certainly,  $X(c(0)) = Y(c(0))$ . The idea—quite standard—is to show that if  $X = Y$  on  $c([0, \tau])$ , with  $0 \leq \tau < 1$ , there is  $\varepsilon > 0$  such that  $X = Y$  on  $c([0, \tau + \varepsilon])$ , and this is done by using (1.1). If  $(u; U)$  is a coordinate system around  $c(\tau)$ , there is  $\varepsilon > 0$  such that  $c([\tau - \varepsilon, \tau + \varepsilon]) \subset U$ , and the curves  $(X^j \circ c)$  and  $(Y^j \circ c)$  defined on  $(\tau - \varepsilon, \tau + \varepsilon)$  are solutions of the system (1.1). Since they coincide for  $t = \tau$ , they are equal on their domain of definition. This proves  $X = Y$  on  $c([0, \tau + \varepsilon])$ .

**(1.3) Lemma.** *Let  $M$  be analytic, and  $(u; U)$  a coordinate system such that  $u(U) \subset R^m$  is convex. Then any parallelism field  $X$  defined on an open connected subset  $V$  of  $U$  can be extended to a unique parallelism field  $Y$  on  $U$ .*

*Proof.* The uniqueness of the extension follows from (1.2) or, more easily, from the fact that if two analytic vector fields coincide on  $V$ , they must coincide in the connected component of  $V$  in the domain of definition.

Choose  $x_0 \in V$ . Define  $c_x: [0, 1] \rightarrow U$  for  $x \in U$  as the curve determined by the condition  $u(c_x(t)) = (1-t)u(x_0) + tu(x)$ . The map  $U \times [0, 1] \rightarrow U, (x, t) \rightarrow c_x(t)$  is analytic. Clearly  $c_x$  is a curve joining  $x_0$  and  $x$ . Substitute  $c$  for  $c_x$  in the formula for  $\mathcal{Q}_j^i(t)$  in (1.1). One gets a family of analytic maps

$\mathcal{Q}_x^i(x, t)$ , and we have a linear system  $S_x$  of equations depending on a parameter  $x$ . For each  $x \in U$  the solution  $(\alpha_x^i)$  with initial condition  $\alpha_x^i(0) = u^i(x_0)$  is defined for  $t = 1$ . We define  $Y = \sum Y^i \partial / \partial u^i$  by the formula  $Y^i(x) = \alpha_x^i(1)$ , and show that  $Y$  is the required extension.

There is a neighborhood  $W$  of  $x_0$  with the following property: If  $x \in W$ , then  $c_x(t) \in V$  for all  $t \in [0, 1]$ . Using (1.1) and the uniqueness of the solution we get for  $x \in W$ :  $X^i(x) = X^i(c_x(1)) = \alpha_x^i(1) = Y^i(x)$ . Therefore  $X|_W = Y|_W$ , and this implies, since our fields are analytic, that  $X = Y|_V$ . The field  $Y$  is a parallelism field because  $L_Y \pi|_V = L_X \pi = 0$  implies, using analyticity once more, that  $L_Y \pi = 0$  on  $U$ .

### 2. The uniqueness theorem

Let  $p: \mathcal{Q} \rightarrow M$  be a  $G$ -structure, and  $P$  the corresponding pseudogroup of transformations. By definition a diffeomorphism  $f: U \rightarrow V$ ;  $U, V$  open subsets of  $M$ , is in  $P$  if and only if the natural lift  $f_*$  to the frame bundle sends  $\mathcal{Q}|_U$  into  $\mathcal{Q}|_V$ . If  $f \in P$ , we denote this natural restriction of  $f_*$  by  $f_0$ , and we still call it the natural lift. If  $X$  is a vector field on  $M$ , and  $\{\psi_t; t \in R\}$  the corresponding pseudogroup induced by  $X$ , we say that  $X$  is an  $\mathcal{Q}$ -field if for all  $t \in R, \psi_t \in P$ . If this is the case,  $X$  has a natural lift to a field  $X_0$  on  $\mathcal{Q}$  which projects on  $X$ . The pseudogroup determining  $X_0$  is just  $\{(\psi_t)_0; t \in R\}$ . We denote the set of  $\mathcal{Q}$ -fields by  $\mathfrak{F}$ . If  $U \subset M$  is open, then  $\mathfrak{F}_U$  will denote the set of  $\mathcal{Q}|_U$ -fields. Let  $\theta$  be the canonical 1-form on  $\mathcal{Q}$  with values in  $R^m$ . It is well known that  $f_0^* \theta = \theta$  for  $f \in P$ , and  $L_{X_0} \theta = 0$  for  $X \in \mathfrak{F}$ .

We now quote some facts about Sternberg prolongations. The reader interested in details should go to [1], whose notation we keep as much as possible. If  $\mathfrak{G}$  is the Lie algebra of  $G$ , we denote by  $\mathfrak{G}_k$  the  $k$ th prolongation of  $\mathfrak{G}$ , and write  $E_k = R^m \oplus \mathfrak{G} \oplus \mathfrak{G}_1 \oplus \dots \oplus \mathfrak{G}_k$ .

We collect the necessary facts in the following theorem:

**(2.1) Theorem.** *There is a sequence of manifolds  $\mathcal{Q}_k$  ( $k \geq 0$ ), maps  $p_k: \mathcal{Q}_k \rightarrow \mathcal{Q}_{k-1}$  ( $k \geq 1$ ) and groups  $G_k$  ( $k \geq 0$ ) such that the following hold:*

- (a)  $\mathcal{Q}_0 = \mathcal{Q}, G_0 = G$ , and  $G_k$  is isomorphic to the vector group  $\mathfrak{G}_k$  for  $k \geq 1$ .
- (b)  $p_k: \mathcal{Q}_k \rightarrow \mathcal{Q}_{k-1}$  is a  $G_k$ -structure. All the maps  $p_k$  admit global sections; hence, these principal bundles are trivial.
- (c) If  $\theta_k$  is the canonical 1-form on  $\mathcal{Q}_k$ , then  $\theta_k$  takes values in  $E_{k-1}$ .
- (d) If  $X \in \mathfrak{F}$ , one can define inductively a lift  $X_k$  of  $X$  to a field in  $\mathcal{Q}_k$  for each  $k \geq 0$ .  $X_0$  is defined as in the paragraph above, and  $X_k = (X_{k-1})_0$  for  $k \geq 1$ .

We give two more elementary lemmas; the first is a simple exercise, the second is in [2, VI.2.1].

**(2.2) Lemma.** *Let  $q: X \rightarrow Y$  be a quotient map of topological spaces (this is the case if  $q$  is continuous, open and onto). If  $q^{-1}(y)$  is connected for all  $y \in Y$ , and  $Z \subset Y$  is open and connected, then  $q^{-1}Z$  is connected.*

**(2.3) Lemma.** *If  $X_0$  is the natural lift of  $X \in \mathcal{F}$ , then it has the following properties:*

- (a) For all  $g \in G$ ,  $g_*X_0 = X_0$ ,
- (b)  $L_{X_0}\theta = 0$ ,
- (c)  $X_0$  projects on  $X$ .

*Conversely, if  $Y$  is a field on an open subset  $U$  of  $\mathcal{Q}$ , satisfying (a) and (b), then  $Y$  is projectable onto a field  $X$  on  $pU$  and  $Y = X_0$  on  $U$ .*

We get from this lemma that if  $X \in \mathcal{F}$ , then  $L_{X_k}\theta_k = 0$  for all  $k \geq 0$ . We make from now on the hypothesis that  $\mathcal{G}$  is of finite type; hence there is  $k \geq 1$  such that  $\mathcal{G}_{k-1} \neq 0$  and  $\mathcal{G}_k = 0$ . In this case  $\theta_k$  is a parallelism on  $\mathcal{Q}_k$ , and  $X_k$  is a parallelism field for  $\theta_k$  if  $X \in \mathcal{F}$ .

**(2.4) Proposition.** *If  $M$  is connected, and  $X, Y \in \mathcal{F}$  are such that for some  $a_k \in \mathcal{Q}_k$ ,  $X_k(a_k) = Y_k(a_k)$ , then  $X = Y$ .*

*Proof.* We get from (2.1)(a), (2.1)(b) and (2.2) that the connected components of  $\mathcal{Q}_k$  are the sets  $(p_1 \circ \dots \circ p_k)^{-1}C = C_k$  where  $C$  is a component of  $\mathcal{Q}$ . If  $a_k \in C_k$ , then  $X_k = Y_k$  on  $C_k$  by (1.2). Since these fields project on  $X$  and  $Y$  and  $(p \circ p_1 \circ \dots \circ p_k)C_k = M$ , we get  $X = Y$ .

**(2.5) Theorem.** *If  $M$  is connected, and  $X, Y \in \mathcal{F}$  are such that  $X|U = Y|U$  for some open  $U \subset M$ , then  $X = Y$ .*

*Proof.* By definition of a  $k$ -lift, if  $X|U = Y|U$  then  $X_k = Y_k$  on  $(p \circ \dots \circ p_k)^{-1}U \subset \mathcal{Q}_k$ , and the theorem follows from (2.4).

This generalizes (0.1) since the Lie algebras of the orthogonal group and the conformal group are of finite type [1, I.2].

### 3. The extension theorem

**(3.1) Proposition.** *Let the structure  $\mathcal{Q}$  be analytic, and  $Z$  a vector field on an open connected subset  $W$  of  $\mathcal{Q}_k$ . Let  $V \subset M$  be open, and  $X \in \mathcal{F}_V$  such that  $X_k = Z$  on  $W \cap (p \circ \dots \circ p_k)^{-1}V$ . Then  $Z$  is projectable on a field  $Y \in \mathcal{F}_V$ , with  $U = (p \circ \dots \circ p_k)W$ ,  $Y = X$  on  $U \cap V$ , and  $Y_k|W = Z$ .*

*Proof.* Consider the 1-forms  $L_Z\theta_k$  and  $L_{X_k}\theta_k$ ; they coincide on  $W \cap (p \circ \dots \circ p_k)^{-1}V$ , and by (2.3)(b) the second form is 0 there. Thus the analytic form  $L_Z\theta_k = 0$  on  $W$  which is connected. Analogously one proves that for all  $g_k \in G_k$ ,  $(g_k)_*Z - Z = 0$  on  $W$ . Using (2.3) once more, we get that  $Z$  projects on a field  $Z_1$  defined on  $p_k(W) \subset \mathcal{Q}_{k-1}$ , and  $(Z_1)_0 = Z$  on  $W$ . It is easy to construct with these ideas a sequence of fields  $Z_h$  defined on  $(p_{k-h+1} \circ \dots \circ p_k)W \subset \mathcal{Q}_{k-h}$  which coincide with  $X_{k-h}$  on

$(p_{k-h+1} \circ \dots \circ p_k)W \cap (p \circ \dots \circ p_{k-h})^{-1}V$ , and on the common domain of definition  $Z_h$  equals the  $p$ -lift of  $Z_{h+p}$ . This construction can be carried down to  $M$  with the convention  $\mathcal{Q}_{-1} = M$  and  $p_0 = p$ . It is immediate that the field  $Y = Z_{k+1}$  has the required properties.

**(3.2) Proposition.** *Suppose that  $\mathcal{Q}$  is analytic of finite type  $k$ . Let  $(u; U)$  be a chart such that  $u(U) \subset R^m$  is convex. If  $V \subset U$  is open and connected, then any  $X \in \mathcal{F}_V$  has a unique extension to a field  $Y \in \mathcal{F}_U$ .*

*Proof.* The uniqueness is clear from analyticity or (2.5). We prove the existence, assuming first that  $G$  is connected. Take a chart  $(u'; U')$  on  $G$  such that  $u'(U')$  is convex. We get easily from (2.1)(a) and (2.1)(b) that there is an open set  $W \subset \mathcal{Q}_k$  diffeomorphic to the convex set  $u(U) \times u'(U') \times \mathcal{G}_1 \times \dots \times \mathcal{G}_k$  which projects onto  $U$ . On the other hand  $(p \circ \dots \circ p_k)^{-1}V \subset \mathcal{Q}_k$  is connected by (2.2), since  $G$  is connected. Now by applying (1.3) we obtain a field  $Z$  on  $W$  equal to  $X_k$  on  $W \cap (p \circ \dots \circ p_k)^{-1}V$ , which is connected. By (3.1)  $Z$  projects on  $Y$  defined on  $U$ , and so is the required extension.

If  $G$  is arbitrary, let  $C \subset \mathcal{Q}$  be a connected component of  $\mathcal{Q}$ , and let  $H$  be the connected component of the identity in  $G$ . It is clear that  $C$  is an  $H$ -structure on  $M$  of finite type  $k$ . If  $\mathcal{F}'$  is the set of  $C$ -fields, we proved in the preceding paragraph that there is a field  $Y \in \mathcal{F}'_U$  which extends  $X$ . We only need to show that  $Y \in \mathcal{F}_U$ . Let  $\{\psi_t; t \in R\}$  be pseudogroup of  $Y$ . Then  $Y \in \mathcal{F}_U$  if for all  $t \in R$  and  $a \in \mathcal{Q}$ ,  $(\psi_t)_0 a \in \mathcal{Q}$ . Writing  $a = cg$  with  $c \in C$  and  $g \in G$  we get  $(\psi_t)_0 a = (\psi_t)_0 (cg) = ((\psi_t)_0 c)g \in Cg \subset \mathcal{Q}$ . This ends the proof.

**(3.3) Theorem.** *(Generalization of (0.2)). Let  $M$  be a connected simply connected manifold, and  $\mathcal{Q}$  an analytic  $G$ -structure on  $M$  of finite type. If  $U$  is an open connected subset of  $M$  and  $X \in \mathcal{F}_U$ , then  $X$  has a unique extension to a field  $Y \in \mathcal{F}$ .*

*Proof.* The uniqueness of the extension follows from analyticity or (2.5). The idea for proving the existence of the extension is a standard one in algebraic topology, and therefore we just give a sketch of the proof. Fix  $x_0 \in U$ . For each  $x_1 \in M$  choose a continuous curve  $c: [0, 1] \rightarrow M$  with  $c(0) = x_0$  and  $c(1) = x_1$ . One shows: (a) There are a neighborhood  $N$  of  $c([0, 1])$  and a field  $Z \in \mathcal{F}_N$  which coincides with  $X$  in a neighborhood of  $x_0$ . (b) If  $c_0, c_1$  are curves joining  $x_0$  and  $x_1$  and if  $Z_0, Z_1$  are fields constructed as in (a), then  $Z_0 = Z_1$  on a neighborhood of  $x_1$ . It follows from (a) and (b) that if we define the field  $Y$  on  $M$  by  $Y(x_1) = Z_0(x_1) = Z_1(x_1)$ , then  $Y$  is well defined and  $Y \in \mathcal{F}$ .

To prove (a) one considers the set  $S$  of  $s \in [0, 1]$  such that there are a neighborhood  $M$  of  $c([0, s])$  and a field  $Z \in \mathcal{F}_N$  which coincides with  $X$  in a

neighborhood of  $x_0$ . We want to show that  $S = [0, 1]$ . This follows from the fact that  $0 \in S$ ,  $S$  is an interval open in  $[0, 1]$ , and  $\sup S \in S$  by (3.2).

The proof of (b) is analogous. If  $(s, t) \rightarrow c_s(t)$  is a homotopy between  $c_0$  and  $c_1$ , consider  $S$ , the set of  $s \in [0, 1]$  such that there are a neighborhood  $N$  of  $\{c_s(t): 0 \leq r \leq s, 0 \leq t \leq 1\}$  and a field  $Z \in \mathcal{F}_N$  which coincides with  $X$  on a neighborhood of  $x_0$ . One shows that  $0 \in S$ ,  $S$  is an interval open in  $[0, 1]$  and  $\sup S \in S$ . This last fact requires (3.2) for its proof. It follows then that  $S = [0, 1]$  proving (b).

**Remark.** Our main results (2.5) and (3.3) are also valid when  $M$  is the family of infinitesimal transformations of a linear connection  $\omega$  on a  $G$ -structure  $A$ . If  $\theta$  is the fundamental form on  $A$ , then  $\pi = \theta \oplus \omega$  is a parallelism on  $A$  with values on  $R^m \oplus \mathcal{G}$ , and the natural lift of  $X$  is a field of the parallelism  $\pi$  [1]. The reader may check easily that the methods of proof of (2.5) and (3.3) work in this new situation.

### References

- [1] S. Kobayashi, *Transformation groups in differential geometry*, Springer, Berlin, 1972.
- [2] S. Kobayashi & K. Nomizu, *Foundations of differential geometry*, Vol. 1, Interscience, New York, 1963.
- [3] A. J. Ledger & M. Obata, *Compact Riemannian manifolds with essential groups of conformorphisms*, *Trans. Amer. Math. Soc.* **150** (1970), 645-651.
- [4] K. Nomizu, *On local and global existence of Killing vector fields*, *Ann. of Math.* **72** (1960) 105-120.

UNIVERSIDAD DE COMPLUTENSE, MADRID