

CR-SUBMANIFOLDS OF A KAEHLER MANIFOLD. II

BANG-YEN CHEN

1. Introduction

A submanifold N in a Kaehler manifold \tilde{M} is called a CR -submanifold if (1) the maximal complex subspace \mathfrak{D}_x of the tangent space $T_x\tilde{M}$ containing in T_xN , $x \in N$, defines a differentiable distribution on N , and (2) the orthogonal complementary distribution \mathfrak{D}^\perp of \mathfrak{D} is a totally real distribution, i.e., $J\mathfrak{D}_x^\perp \subseteq T_x^\perp N$, $x \in N$, where J denotes the almost complex structure of \tilde{M} , and $T_x^\perp N$ the normal space of N in \tilde{M} at x .

In the first part of this series, we have obtained several fundamental results for CR -submanifolds. In the present part, we shall continue our study on such submanifolds. In particular, we prove that (a) the holomorphic distribution \mathfrak{D} of any CR -submanifold in a Kaehler manifold is minimal (Proposition 3.9); (b) every leaf of the holomorphic distribution of a mixed foliate proper CR -submanifold in a complex hyperbolic space H^m is Einstein-Kaehlerian (Proposition 4.4); and (c) every CR -submanifold with semi-flat normal connection in CP^m is either an anti-holomorphic submanifold in some totally geodesic CP^{h+p} of CP^m or a totally real submanifold (Theorem 5.11).

2. Preliminaries

Let \tilde{M}^m be a complex m -dimensional Kaehler manifold with complex structure J , and N be a real n -dimensional ($n \geq 2$) Riemannian manifold isometrically immersed in \tilde{M}^m . We denote by \langle, \rangle the metric tensor of \tilde{M}^m as well as that induced on N . Let ∇ and $\tilde{\nabla}$ be the covariant differentiations on N and \tilde{M} respectively. Then the Gauss and Weingarten formulas for N are given respectively by

$$(2.1) \quad \tilde{\nabla}_X Y = \nabla_X Y + \sigma(X, Y),$$

$$(2.2) \quad \tilde{\nabla}_X \xi = -A_\xi X + D_X \xi,$$

for any vector fields X, Y tangent to N , and ξ normal to N , where σ is the second fundamental form, and D the normal connection.

For any vector X tangent to N and ξ normal to N we put

$$(2.3) \quad JX = PX + FX,$$

$$(2.4) \quad J\xi = t\xi + f\xi$$

where PX and $t\xi$ (respectively, FX and $f\xi$) are the tangential (respectively, normal) components of JX and $J\xi$ respectively.

In the following we shall denote by $\tilde{M}^m(c)$ a complex m -dimensional complex-space-form of constant holomorphic sectional curvature c . We have

$$(2.5) \quad \begin{aligned} \tilde{R}(X, Y)Z = \frac{c}{4} \{ &\langle Y, Z \rangle X - \langle X, Z \rangle Y + \langle JY, Z \rangle JX \\ &- \langle JX, Z \rangle JY + 2\langle X, JY \rangle JZ \}. \end{aligned}$$

We denote by R and R^\perp the curvature tensors associated with ∇ and D respectively. A submanifold N is said to be flat (respectively, to have flat normal connection) if $R \equiv 0$ (respectively, $R^\perp \equiv 0$). For any vector fields X, Y, Z, W in the tangent bundle TN , and ξ, η in the normal bundle $T^\perp N$, the equations of Gauss, Codazzi and Ricci are given respectively by

$$(2.6) \quad \begin{aligned} R(X, Y; Z, W) = \tilde{R}(X, Y; Z, W) + \langle \sigma(X, W), \sigma(Y, Z) \rangle \\ - \langle \sigma(X, Z), \sigma(Y, W) \rangle, \end{aligned}$$

$$(2.7) \quad \begin{aligned} \tilde{R}(X, Y; Z, \xi) = \langle D_X \sigma(Y, Z) - \sigma(\nabla_X Y, Z) - \sigma(Y, \nabla_X Z), \xi \rangle \\ - \langle D_Y \sigma(X, Z) - \sigma(\nabla_Y X, Z) - \sigma(X, \nabla_Y Z), \xi \rangle, \end{aligned}$$

$$(2.8) \quad \tilde{R}(X, Y; \xi, \eta) = R^\perp(X, Y; \xi, \eta) - \langle [A_\xi, A_\eta] X, Y \rangle,$$

where $R(X, Y; Z, W) = \langle R(X, Y)Z, W \rangle, \dots$, etc.

Definition 2.1. A submanifold N of a Kaehler manifold \tilde{M} is called a *CR-submanifold* if there is a differentiable distribution $\mathfrak{D}: x \rightarrow \mathfrak{D}_x \subseteq T_x N$ on N satisfying the following conditions:

(a) \mathfrak{D} is holomorphic, i.e., $J\mathfrak{D}_x = \mathfrak{D}_x$ for each $x \in N$, and

(b) the complementary orthogonal distribution $\mathfrak{D}^\perp: x \rightarrow \mathfrak{D}_x^\perp \subseteq T_x N$ is totally real, i.e., $J\mathfrak{D}_x^\perp \subseteq T_x^\perp N$ for each $x \in N$.

If $\dim \mathfrak{D}_x^\perp = 0$ (respectively, $\dim \mathfrak{D}_x = 0$), N is called a *complex* (respectively, *totally real*) submanifold. A *CR-submanifold* is said to be *proper* if it is neither complex nor totally real.

For a *CR-submanifold* N we shall denote by ν the orthogonal complementary subbundle of $J\mathfrak{D}^\perp$ in $T^\perp N$. We have

$$(2.9) \quad T^\perp N = J\mathfrak{D}^\perp \oplus \nu, \quad \nu_x = T_x^\perp N \cap J(T_x^\perp N).$$

A subbundle μ of the normal bundle is said to be parallel if $D_x \xi \in \mu$ for any vector $X \in TN$ and section ξ in μ .

A CR-submanifold N in a Kaehler manifold \tilde{M} is said to be anti-holomorphic if $T_x^\perp N = J\mathfrak{D}_x^\perp$, $x \in N$.

3. Some basic lemmas

First we recall some basic lemmas for later use.

Lemma 3.1 [4]. *Let N be a CR-submanifold of a Kaehler manifold \tilde{M} . Then we have*

$$(3.1) \quad \langle \nabla_U Z, X \rangle = \langle JA_{JZ}U, X \rangle,$$

$$(3.2) \quad A_{JZ}W = A_{JW}Z,$$

$$(3.3) \quad A_{J\xi}X = -A_\xi JX$$

for any vector fields U tangent to N , X in \mathfrak{D} , Z, W in \mathfrak{D}^\perp , and ξ in ν .

Lemma 3.2 [4]. *The totally real distribution \mathfrak{D}^\perp of any CR-submanifold in a Kaehler manifold is integrable.*

Lemma 3.3 [1], [2], [4]. *Let N be a CR-submanifold of a Kaehler manifold \tilde{M} . Then the holomorphic distribution \mathfrak{D} is integrable if and only if*

$$(3.4) \quad \langle \sigma(X, JY), JZ \rangle = \langle \sigma(JX, Y), JZ \rangle$$

for any vectors X, Y in \mathfrak{D} , and Z in \mathfrak{D}^\perp .

Lemma 3.4 [2]. *Let N be a CR-submanifold in a Kaehler manifold \tilde{M} . Then the leaves of \mathfrak{D}^\perp are totally geodesic in \tilde{M} if and only if*

$$(3.5) \quad \langle \sigma(\mathfrak{D}, \mathfrak{D}^\perp), J\mathfrak{D}^\perp \rangle = \{0\}.$$

Lemma 3.5. *Let N be a CR-submanifold in a Kaehler manifold \tilde{M} . We have the following statements:*

(a) *If the leaves of \mathfrak{D}^\perp are totally geodesic in \tilde{M} , then*

$$(3.6) \quad \sigma(\mathfrak{D}^\perp, \mathfrak{D}^\perp) = \{0\}, \quad \langle \sigma(\mathfrak{D}, \mathfrak{D}^\perp), J\mathfrak{D}^\perp \rangle = \{0\},$$

$$(3.7) \quad \tilde{H}_B(X, Z) = 2\|\sigma(X, Z)\|^2 + 2\langle A_{JZ}JX, JA_{JZ}X \rangle$$

for any unit vectors X in \mathfrak{D} , and Z in \mathfrak{D}^\perp , where \tilde{H}_B denotes the holomorphic bisectional curvature of \tilde{M} .

(b) *If (3.6) holds, the leaves of \mathfrak{D}^\perp are totally geodesic in \tilde{M} .*

Proof. Let N be a CR-submanifold in a Kaehler manifold \tilde{M} . Then \mathfrak{D}^\perp is integrable (Lemma 3.2). Let N^\perp be a leaf of \mathfrak{D}^\perp . We denote by σ^\perp and σ'' the second fundamental form of N^\perp in \tilde{M} and N , respectively. We have

$$\sigma^\perp(Z, W) = \sigma''(Z, W) + \sigma(Z, W)$$

for any vectors Z, W in \mathfrak{D}^\perp . Thus, by Lemma 3.4, the leaves of \mathfrak{D}^\perp are totally geodesic in \tilde{M} , if and only if (3.6) holds.

Assume that the leaves of \mathfrak{D}^\perp are totally geodesic in \tilde{M} . For any vector fields X, Y in \mathfrak{D} and Z, W in \mathfrak{D}^\perp , equation (2.7) of Codazzi and (3.5) give

$$\begin{aligned} \tilde{R}(X, Y; Z, JW) &= \langle D_X\sigma(Y, Z) - \sigma(Y, \nabla_X Z), JW \rangle \\ &\quad - \langle D_Y\sigma(X, Z) - \sigma(X, \nabla_Y Z), JW \rangle, \\ &= \langle \sigma(X, Z), J\tilde{\nabla}_Y W \rangle - \langle \sigma(Y, Z), J\tilde{\nabla}_X W \rangle \\ &\quad + \langle A_{JW}X, \nabla_Y Z \rangle - \langle A_{JW}Y, \nabla_X Z \rangle \\ &= \langle \sigma(X, Z), J\sigma(Y, W) \rangle - \langle \sigma(Y, Z), J\sigma(X, W) \rangle \\ &\quad + \langle A_{JW}X, \nabla_Y Z \rangle - \langle A_{JW}Y, \nabla_X Z \rangle. \end{aligned}$$

Thus by applying (3.5) and Lemma 4.1 we find

$$\begin{aligned} \tilde{R}(X, Y; Z, JW) &= \langle \sigma(X, Z), \sigma(JY, W) \rangle - \langle \sigma(Y, Z), \sigma(JX, W) \rangle \\ &\quad + \langle A_{JW}X, JA_{JZ}Y \rangle - \langle A_{JW}Y, JA_{JZ}X \rangle, \end{aligned}$$

from which we obtain (3.7).

Corollary 3.6. *Let N be a proper anti-holomorphic submanifold in $\mathbb{C}P^{h+p}$. If the leaves of \mathfrak{D}^\perp are totally geodesic in $\mathbb{C}P^{h+p}$, then the holomorphic distribution is not integrable.*

This corollary follows from Lemmas 3.4 and 3.5.

For the holomorphic distribution \mathfrak{D} , we have

Lemma 3.7. *Let N be a CR-submanifold in a Kaehler manifold \tilde{M} . Then*

(1) *the holomorphic distribution is integrable, and its leaves are totally geodesic in N if and only if*

$$(3.8) \quad \langle \sigma(\mathfrak{D}, \mathfrak{D}), J\mathfrak{D}^\perp \rangle = \{0\},$$

(2) *the holomorphic distribution is integrable, and its leaves are totally geodesic in \tilde{M} if and only if*

$$(3.9) \quad \sigma(\mathfrak{D}, \mathfrak{D}) = \{0\}.$$

Proof. Let N be a CR-submanifold in a Kaehler manifold \tilde{M} . If (3.8) holds, then also (3.4). Thus the holomorphic distribution \mathfrak{D} is integrable (Lemma 3.3). Moreover, from (2.1), (2.2) and (2.3) we have

$$\begin{aligned} \langle \nabla_X Z, JY \rangle &= \langle \tilde{\nabla}_X Z, JY \rangle = -\langle \tilde{\nabla}_X JZ, Y \rangle \\ &= -\langle A_{JZ}X, Y \rangle = -\langle \sigma(X, Y), JZ \rangle = 0 \end{aligned}$$

for any vector fields X, Y in \mathfrak{D} , and Z in \mathfrak{D}^\perp . Thus the leaves of \mathfrak{D} are totally geodesic in N . The converse of this has been proved in [4].

Statement (2) follows from statement (1) and the following identity

$$\sigma^T(X, Y) = \sigma'(X, Y) + \sigma(X, Y)$$

for any vectors X, Y in \mathfrak{D} , where σ' and σ^T are the second fundamental forms of any leaf N^T of \mathfrak{D} in N and \tilde{M} respectively.

Let \mathfrak{H} be a differentiable distribution on a CR-submanifold N ($\mathfrak{H}: x \rightarrow \mathfrak{H}_x \subseteq T_x N, x \in N$). We put

$$(3.10) \quad \mathring{\sigma}(X, Y) = (\nabla_X Y)^\perp$$

for any vector fields X, Y in \mathfrak{H} , where $(\nabla_X Y)^\perp$ denotes the component of $\nabla_X Y$ in the orthogonal complementary distribution \mathfrak{H}^\perp in N . Then the Frobenius theorem gives the following

Lemma 3.8. *The distribution \mathfrak{H} is integrable if and only if $\mathring{\sigma}$ is a symmetric on $\mathfrak{H} \times \mathfrak{H}$.*

Let X_1, \dots, X_r be an orthonormal basis in \mathfrak{H} . We put

$$\mathring{H} = \frac{1}{r} \sum_{i=1}^r \mathring{\sigma}(X_i, X_i).$$

Then \mathring{H} is a well-defined vector field on N (up to sign). We call \mathring{H} the *mean-curvature vector of the distribution \mathfrak{H}* .

A distribution \mathfrak{H} on N is said to be *minimal* if the mean curvature vector \mathring{H} of \mathfrak{H} vanishes identically, and \mathfrak{H} is said to be *totally geodesic* if $\mathring{\sigma} \equiv 0$.

Proposition 3.9. *Let N be a CR-submanifold of a Kaehler manifold \tilde{M} . Then*

- (a) *the holomorphic distribution \mathfrak{D} is minimal, and*
- (b) *the distribution \mathfrak{D} is totally geodesic if and only if \mathfrak{D} is integrable, and its leaves are totally geodesic in N .*

Proof. Let N be a CR-submanifold of a Kaehler manifold \tilde{M} . For any vector fields X in \mathfrak{D} , and Z in \mathfrak{D}^\perp , Lemma 3.1 gives

$$(3.11) \quad \langle Z, \nabla_X X \rangle = \langle A_{JZ} X, JX \rangle.$$

Thus we have

$$(3.12) \quad \langle Z, \nabla_{JX} JX \rangle = -\langle A_{JZ} X, JX \rangle.$$

Combining (3.11) and (3.12) we obtain

$$(3.13) \quad \langle \nabla_X X + \nabla_{JX} JX, Z \rangle = 0.$$

This implies statement (a). Statement (b) follows from (3.10) and Lemma 3.8.

4. Mixed foliate CR-submanifolds

Definition 4.1. A CR-submanifold is said to be *mixed totally geodesic* if $\sigma(\mathfrak{D}, \mathfrak{D}^\perp) = \{0\}$.

Definition 4.2. A CR-submanifold N in a Kaehler manifold \tilde{M} is said to be *mixed foliate*, if it is mixed totally geodesic, and its holomorphic distribution is integrable.

In [2], Bejancu, Kon and Yano proved that there is no mixed foliate proper CR-submanifold in $\tilde{M}^m(c)$ with $c > 0$. In [4] the author proved that a CR-submanifold in C^m is mixed foliate if and only if N is a CR-product (for anti-holomorphic case, see [2]).

In this section, we shall study mixed foliate CR-submanifolds in a complex hyperbolic space H^m . For simplicity, we assume that H^m is a complex m -dimensional complex hyperbolic space with constant holomorphic sectional curvature -4 .

Lemma 4.1. *Let N be a mixed foliate CR-submanifold in H^m . Then for any unit vectors $X \in \mathfrak{D}$ and $Z \in \mathfrak{D}^\perp$,*

$$(4.1) \quad \|A_{JZ}X\| = 1,$$

$$(4.2) \quad \|\sigma\|^2 \geq 2hp,$$

where $h = \dim_{\mathbb{C}} \mathfrak{D}$, and $p = \dim_{\mathbb{R}} \mathfrak{D}^\perp$. The equality sign in (4.2) holds if and only if (a) the leaves of \mathfrak{D}^\perp are totally geodesic in H^m , and (b) $\text{Im } \sigma = J\mathfrak{D}^\perp$.

Proof. Let N be a mixed foliate CR-submanifold in H^m . Then Lemma 9.1 of [4] gives

$$(4.3) \quad \tilde{H}_B(X, Z) = -2\|A_{JZ}X\|^2,$$

for any unit vectors X in \mathfrak{D} , and Z in \mathfrak{D}^\perp . This gives (4.1).

Inequality (4.2) follows immediately from (4.1). From (4.1) it is clear that $\|\sigma\| = 2hp$ if and only if we have

$$(4.4) \quad \text{Im } \sigma = J\mathfrak{D}^\perp,$$

$$(4.5) \quad A_{J\mathfrak{D}^\perp} \mathfrak{D}^\perp = \{0\}.$$

The lemma thus follows from Lemma 3.5.

Let N be a mixed foliate CR-submanifold in H^m , and N^T a leaf of the holomorphic distribution \mathfrak{D} . Then N^T is a Kaehler submanifold of H^m . We denote by σ^T, D^T, \dots , etc. the second fundamental form, the normal connection, \dots , etc. for N^T in H^m , and by σ', D', \dots , etc. the corresponding quantities for N^T in N . Then we have

$$(4.6) \quad \sigma^T(X, Y) = \sigma'(X, Y) + \sigma(X, Y)$$

for X, Y in TN^T . For any Z in \mathfrak{D}^\perp , this implies

$$(4.7) \quad \langle A_Z^T X, Y \rangle = \langle J\sigma^T(X, Y), JZ \rangle = \langle \sigma(JX, Y), JZ \rangle = \langle A_{JZ} JX, Y \rangle,$$

$$(4.8) \quad \langle A_{JZ}^T X, Y \rangle = \langle \sigma(X, Y), JZ \rangle = \langle A_{JZ} X, Y \rangle.$$

Because N is mixed foliate, these give

$$(4.9) \quad A_Z^T X = A_{JZ} JX, \quad A_{JZ}^T X = A_{JZ} X.$$

Moreover, for any unit vector fields X in \mathfrak{D} , and Z in \mathfrak{D}^\perp , we have that

$$(4.10) \quad J \nabla_X Z = \tilde{\nabla}_X JZ = -A_{JZ} X + D_X JZ,$$

so that

$$(4.11) \quad D_X JZ = F \nabla_X Z.$$

From

$$\tilde{\nabla}_X JZ = -A_{JZ}^T X + D_X^T JZ$$

we also get

$$(4.12) \quad D_X^T JZ = D_X JZ.$$

Let η be any normal vector field in ν (for the definition of ν , see (2.9)) and X, Y any tangent vector fields in \mathfrak{D} , (2.5), (4.11) and (4.12) imply

$$(4.13) \quad \tilde{R}(X, Y; JZ, \eta) = 0,$$

$$(4.14) \quad R_T^\perp(X, Y; JZ, \eta) = 0.$$

Combining these with equation (2.7) of Codazzi we obtain

$$(4.15) \quad [A_{JZ}^T, A_\eta^T] = 0 \quad \text{for } \eta \in \nu, z \in \mathfrak{D}^\perp.$$

Because N^T is a Kaehler submanifold, $A_{J\eta}^T = JA_\eta^T = -A_\eta^T J$. Thus by using (4.15) we have

$$0 = A_{JZ}^T A_\eta^T - A_\eta^T A_{JZ}^T = J(A_\eta^T A_{JZ}^T + A_{JZ}^T A_\eta^T).$$

Since J is nonsingular, this gives

$$(4.16) \quad A_\eta^T A_{JZ}^T + A_{JZ}^T A_\eta^T = 0.$$

Combining (4.15) and (4.16) we have

$$(4.17) \quad A_{JZ}^T A_\eta^T = 0.$$

Because N is mixed foliate, $A_{JZ} \mathfrak{D} \subseteq \mathfrak{D}$ for any Z in \mathfrak{D}^\perp . Thus using Lemma 4.1 and (4.9) we get

$$(4.18) \quad \|A_Z^T X\| = \|A_{JZ}^T X\| = 1$$

for any unit vectors X in TN^T , and Z in \mathfrak{D}^\perp . By linearity, this implies

$$(4.19) \quad \langle A_{JZ}^T X, A_{JZ}^T Y \rangle = 0$$

for orthogonal vectors X, Y in TN^T . From (4.18) and (4.19) we find

$$(4.20) \quad A_Z^T, A_{JZ}^T \in \in O(2h).$$

In particular, A_{JZ}^T is nonsingular. Thus we have, in consequence of (4.17), $A_\eta^T = 0$ for any vector η in ν . Since N is mixed foliate, (2.1) and (2.2) give

$$-A_Z^T X + D_X^T Z = \tilde{\nabla}_X Z = \nabla_X Z = -A_Z' X + D_X' Z.$$

from which we find $D_X^T Z = D'_X Z$. This shows that the normal subbundle $\mathfrak{D}^\perp|_{N^T}$ is a parallel subbundle of the normal bundle of N^T in H^m . Therefore we have

$$(4.21) \quad R_T^\perp(X, Y; Z, JW) = 0$$

for any vector fields X, Y in TN^T , and Z, W in $\mathfrak{D}^\perp|_{N^T}$. Let Z_1, \dots, Z_p be an orthonormal basis of \mathfrak{D}^\perp_x , $x \in N^T$. (2.5), (4.21) and the Ricci equation for N^T in H^m give

$$(4.22) \quad [A_{Z_\alpha}^T, A_{JZ_\beta}^T] = 0 \text{ for } \alpha \neq \beta, \alpha, \beta = 1, \dots, p.$$

Since $A_{JZ}^T J = -JA_{JZ}^T$, (4.20) shows that A_{JZ}^T has two eigenvalues 1 and -1 with the same multiplicity h . We put

$$V_1 = \{X \in T_x N \mid A_{JZ_1}^T X = X\}.$$

Thus, for any $X \in V_1$, (4.22) gives

$$A_{JZ_1}^T A_{Z_\alpha}^T X = A_{Z_\alpha}^T A_{JZ_1}^T X = A_{Z_\alpha}^T X, \quad \alpha = 2, \dots, p.$$

Moreover, for any unit vector X in V_1 , (4.22) implies that $A_{Z_\alpha}^T X$, $\alpha = 2, \dots, p$ lie in V_1 , which are orthonormal by (4.18). Consequently, we obtain $p \leq h + 1$.

From (4.22), we may also get

$$A_{Z_\alpha}^T A_{Z_\beta}^T + A_{Z_\beta}^T A_{Z_\alpha}^T = 0 \quad \text{for } \alpha \neq \beta.$$

From the equation of Gauss and (2.5), the sectional curvature K of N satisfies

$$(4.23) \quad K(X, Z) = -1 + \langle \sigma(X, X), \sigma(Z, Z) \rangle$$

for any unit vectors X in \mathfrak{D} , and Z in \mathfrak{D}^\perp . Since N is mixed foliate, we also have

$$K(JX, Z) = -1 - \langle \sigma(X, X), \sigma(Z, Z) \rangle.$$

Combining this with (4.23) gives

$$K(X, Z) + K(JX, Z) = -2.$$

By summarizing the above facts we can state the next lemma.

Lemma 4.2. *Let N be a mixed foliate CR-submanifold in H^m . Then*

- (a) $D_X^T JZ = D_X JZ = F \nabla_X Z$,
- (b) $D_X^T Z = D'_X Z = -tD_X JZ$,
- (c) $\text{Im } \sigma^T = \mathfrak{D}^\perp \oplus J\mathfrak{D}^\perp$,
- (d) $A_Z^T, A_{JZ}^T \in O(2h)$,
- (e) $p \leq h + 1$,
- (f) $A_Z^T A_W^T + A_W^T A_Z^T = 0$,
- (g) $K(X, Z) + K(JX, Z) = -2$, for any unit vector field X in TN^T , and orthonormal vector fields Z, W in \mathfrak{D}^\perp .

From Lemma 4.2 and Proposition 3 of [2] we have the following.

Lemma 4.3. *Let N be a mixed foliate proper CR-submanifold of $\tilde{M}^m(c)$, $c \neq 0$. Then $c < 0$ and $p > 1$.*

Proof. Let N be a mixed foliate proper CR-submanifold of $\tilde{M}^m(c)$, $c \neq 0$. Then Proposition 3 of [2] implies $c < 0$. If $p = 1$, then, for any unit vector field Z in \mathfrak{D}^\perp , statement (b) of Lemma 4.2 implies $D_X^T Z = D_X' Z = 0$. Hence, Z is a parallel normal vector field of the complex submanifold N^T in $\tilde{M}^m(c)$, $c < 0$. This contradicts a theorem of Chen and Ogiue [5].

Proposition 4.4. *Let N be a mixed foliate proper CR-submanifold of H^m . Then*

(a) *each leaf N^T of \mathfrak{D} lies in a complex $(h + p)$ -dimensional totally geodesic complex submanifold H^{h+p} of H^m ,*

(b) *each leaf N^T is an Einstein-Kaehler submanifold of H^{h+p} with Ricci tensor given by*

$$(4.24) \quad S^T(X, Y) = -2(h + p + 1)\langle X, Y \rangle,$$

(c) $h + 1 \geq p \geq 2$; $h \geq 2$, and

(d) *the leaves of \mathfrak{D}^\perp are totally geodesic in N .*

Proof. Lemma 4.2 implies that the first normal space $\text{Im } \sigma^T$ is nothing but $\mathfrak{D}^\perp \oplus J\mathfrak{D}^\perp$. Since $\mathfrak{D}^\perp \oplus J\mathfrak{D}^\perp$ is a parallel normal subbundle of the normal bundle of N^T in H^m , by a theorem of Chen and Ogiue [5], N^T lies in a complex $(h + p)$ -dimensional totally geodesic submanifold H^{h+p} of H^m . Thus (a) is proved.

Since N^T is a Kaehler submanifold of H^m , equation (2.8) of Gauss gives

$$S^T(X, Y) = -2(h + 1)\langle X, Y \rangle - \sum \langle A_{\xi_\alpha}^T X, A_{\xi_\alpha}^T Y \rangle,$$

where ξ_α 's form an orthonormal basis of $T^\perp N^T$. Thus by Lemmas 4.1 and 4.2 we obtain

$$S^T(X, X) = -2(h + p + 1)\langle X, X \rangle,$$

which implies (4.24).

If $h = \dim_{\mathbb{C}} \mathfrak{D} = 1$, then from statement (b) it follows that N^T is of constant curvature $-2(p + 2)$. Since N^T is a Kaehler submanifold of H^m , a theorem of Calabi [4] gives that $p = 0$. This is a contradiction. The remaining part of this proposition follows from Lemmas 3.4 and 4.3.

Theorem 4.5. *Let N be a mixed foliate CR-submanifold of H^m . If $\dim_{\mathbb{R}} N \leq 5$, then N is either a complex submanifold or a totally real submanifold.*

This theorem follows immediately from statement (c) of Proposition 4.4.

Remark 4.1. The author believes that Theorem 4.5 holds for any mixed foliate CR-submanifold of H^m . However, he is unable to prove it at this moment.

5. Semi-flat normal connection

First we recall the following definition [6].

Definition 5.1. A CR-submanifold N in a complex-space-form $\tilde{M}^m(c)$ is said to have *semi-flat normal connection* if its normal curvature tensor R^\perp satisfies

$$(5.1) \quad R^\perp(X, Y; \xi, \eta) = \frac{c}{2} \langle X, PY \rangle \langle J\xi, \eta \rangle$$

for any vectors X, Y in TN , and ξ, η in $T^\perp N$.

The main purpose of this section is to classify CR-submanifolds with semi-flat normal connection.

Lemma 5.1. A CR-submanifold N in a complex-space-form $\tilde{M}^m(c)$ has semi-flat normal connection if and only if

$$(5.2) \quad \langle [A_\xi, A_\eta]X, Y \rangle = \frac{c}{4} \{ \langle JX, \xi \rangle \langle JY, \eta \rangle - \langle JX, \eta \rangle \langle JY, \xi \rangle \}$$

for any vectors X, Y in TN , and ξ, η in $T^\perp N$.

This lemma follows from Definition 5.1 and the equation of Ricci.

From Lemma 5.1 we obtain the following.

Lemma 5.2. Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$. Then

$$(5.3) \quad \langle [A_\xi, A_\eta]X, U \rangle = 0,$$

$$(5.4) \quad \langle [A_\xi, A_\eta]Z, W \rangle = \frac{c}{4} \{ \langle JZ, \xi \rangle \langle JW, \eta \rangle - \langle JZ, \eta \rangle \langle JW, \xi \rangle \}$$

for any vectors U in TN , X in \mathfrak{D} , Z, W in \mathfrak{D}^\perp , and ξ, η in $T^\perp N$.

Moreover, we also have

Lemma 5.3. Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$. Then

$$(5.5) \quad A_{\nu} \mathfrak{D} = \{0\},$$

$$(5.6) \quad \langle A_{J\mathfrak{D}^\perp} \mathfrak{D}, A_{\nu} \mathfrak{D}^\perp \rangle = \{0\},$$

where $\nu_x = T_x^\perp N \cap J(T_x^\perp N)$, $x \in N$.

Proof. From Lemmas 3.1 and 5.2 we have

$$0 = \langle [A_\xi, A_{J\xi}]X, JX \rangle = -\|A_\xi JX\|^2 - \|A_\xi X\|^2$$

for any vectors X in \mathfrak{D} , and ξ in ν . Thus we get (5.5). Formula (5.6) follows from (5.4) and (5.5).

Lemma 5.4 is an immediate consequence of Lemma 5.3.

Lemma 5.4. Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$. If there is a ξ in ν such that $A_\xi \mathfrak{D}^\perp = \mathfrak{D}^\perp$, then N is mixed totally geodesic.

From Lemma 5.2 we have

Lemma 5.5. *Let N be a CR-submanifold with semi-flat normal connection. Then*

$$(5.7) \quad \|A_{JZ}W\|^2 = \frac{\epsilon}{4} + \langle A_{JZ}Z, A_{JW}W \rangle$$

for orthonormal vectors Z, W in \mathfrak{D}^\perp .

Proof. For orthonormal vectors Z and W in \mathfrak{D}^\perp , Lemma 5.2 gives

$$\frac{\epsilon}{4} = \langle [A_{JZ}, A_{JW}]Z, W \rangle = \langle A_{JZ}W, A_{JW}W \rangle - \langle A_{JZ}Z, A_{JW}W \rangle.$$

Thus by using Lemma 3.1 we obtain (5.7).

Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$. By Lemma 5.3 we obtain $A_\nu \mathfrak{D} = \{0\}$. Define an endomorphism

$$\tilde{A}_\xi: \mathfrak{D}_x^\perp \rightarrow \mathfrak{D}_x^\perp$$

by

$$(5.8) \quad \tilde{A}_\xi Z = A_\xi Z$$

for any vectors ξ in ν_x , and Z in \mathfrak{D}_x^\perp . Then \tilde{A}_ξ is self-adjoint.

Let $\lambda_1, \dots, \lambda_r$ be the distinct eigenvalues of \tilde{A}_ξ , and V_1, \dots, V_r the corresponding eigenspaces. Then we have

$$(5.9) \quad \mathfrak{D}_x^\perp = V_1 \oplus \dots \oplus V_r, \langle V_i, V_j \rangle = 0 \quad \text{for } i \neq j.$$

Lemma 5.6. *Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$, $c \neq 0$. Then, for any ξ in ν , \tilde{A}_ξ is proportional to the identity endomorphism.*

Proof. Under the hypothesis, Lemma 5.2 implies

$$(5.10) \quad \langle A_\xi W, A_{JZ}Y \rangle = \langle A_\xi Y, A_{JZ}W \rangle$$

for any vectors ξ in ν , and Y, Z, W in \mathfrak{D}^\perp . If \tilde{A}_ξ is not proportional to the identity endomorphism, $r \geq 2$. Let $Z = W = Z_i \in V_i$, $Y = Z_j \in V_j$, for $i \neq j$. Then (5.10) and Lemma 3.1 imply

$$(5.11) \quad \langle A_{JZ_j}Z_i, Z_i \rangle = 0.$$

By linearity we have

$$(5.12) \quad \langle A_{JZ_j}V_i, V_i \rangle = \{0\} \quad \text{for } i \neq j.$$

Putting $W = Z_i \in V_i$, $Y = Z_j \in V_j$ and $Z = Z_k \in V_k$ for $i \neq j$, (5.10) gives

$$\lambda_i \langle A_{JZ_k}Z_j, Z_i \rangle = \lambda_j \langle A_{JZ_k}Z_j, Z_i \rangle \quad \text{for } i \neq j,$$

which implies

$$(5.13) \quad A_{JZ_k}V_j \subseteq \mathfrak{D} \oplus V_j.$$

On the other hand, by Lemma 5.3 we obtain

$$0 = \langle A_{JZ_k}X, A_\xi Z_j \rangle = \lambda_j \langle A_{JZ_k}Z_j, X \rangle$$

for any vectors X in \mathfrak{D} , $Z_j \in V_j$, and $Z_k \in V_k$. This shows that $A_{JZ_k}V_j \subseteq \mathfrak{D}^\perp$ if $\lambda_j \neq 0$. Combining this with (5.13) yields

$$(5.14) \quad A_{JZ_k}V_j \subseteq V_j \quad \text{whenever } \lambda_j \neq 0.$$

From (5.12) and (5.14) we get

$$(5.15) \quad A_{JZ_k}V_i = 0 \text{ if } j \neq i \text{ and } \lambda_i \neq 0.$$

Since A_ξ has at least two distinct eigenvalues, we may assume that $\lambda_1 \neq 0$. From (5.7) of Lemma 5.5 and (5.15) we have

$$(5.16) \quad 0 = \|A_{JZ_2}Z_1\|^2 = \frac{c}{4} + \langle A_{JZ_2}Z_2, A_{JZ_1}Z_1 \rangle.$$

On the other hand, Lemma 3.1 and (5.12) imply

$$0 = \langle A_{JZ_i}Z_i, Z_i \rangle = \langle A_{JZ_i}Z_i, Z_j \rangle \quad \text{for } i \neq j.$$

Combining this with (5.14) we find

$$(5.17) \quad A_{JZ_i}Z_i \in \mathfrak{D} \oplus V_i.$$

Since $A_{JZ_1}Z_1 \in V_1$ by (5.14), equations (5.16) and (5.17) give $c = 0$. This is a contradiction.

From Lemmas 5.3 and 5.6 we immediately have the following.

Lemma 5.7. *Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$, $c \neq 0$. Then for any $x \in N$, there is a unit normal vector $\bar{\eta} \in \nu_x$ such that*

$$(5.18) \quad A_{\bar{\eta}}X = 0, \quad A_{\bar{\eta}}Z = \lambda Z,$$

$$(5.19) \quad A_\xi = 0$$

for any vectors X in \mathfrak{D}_x , Z in \mathfrak{D}_x , and ξ in ν_x with $\langle \xi, \bar{\eta} \rangle = 0$.

Lemma 5.8. *Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$, $c \neq 0$. If λ is nowhere zero on N , then N is mixed foliate.*

Proof. Under the hypothesis, Lemmas 5.4 and 5.7 imply that N is mixed totally geodesic.

For any vector fields X, Y in \mathfrak{D} , Z in \mathfrak{D}^\perp , and ξ in $T^\perp N$, equation (2.9) of Codazzi gives

$$\begin{aligned} \tilde{R}(X, Y; Z, \xi) &= \langle \sigma([X, Y], Z), \xi \rangle \\ &\quad + \langle \sigma(X, \nabla_Y Z) - \sigma(Y, \nabla_X Z), \xi \rangle. \end{aligned}$$

In particular, if we choose ξ to be the vector $\bar{\eta}$ of Lemma 5.7, we can reduce this to

$$0 = \langle \sigma([X, Y], Z), \bar{\eta} \rangle = \lambda \langle [X, Y], Z \rangle$$

by applying (2.6) and Lemma 5.7. Since $\lambda \neq 0$, this shows that the holomorphic distribution is integrable.

Lemma 5.9. *Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$, $c \neq 0$.*

(1) *Then λ is constant, and for any vectors X, Y in TN and Z in \mathfrak{D}^\perp we have*

$$(5.20) \quad \begin{aligned} F(R(X, Y)Z) &= \sigma(X, P\nabla_Y Z) - \sigma(Y, P\nabla_X Z) \\ &\quad + \lambda^2\{\langle Y, Z \rangle FX - \langle X, Z \rangle FY\}, \end{aligned}$$

$$(5.21) \quad D_X JZ = F\nabla_X Z + \lambda\langle X, Z \rangle J\bar{\eta},$$

(2) *If $\lambda = 0$, then N lies in a totally geodesic complex submanifold $\tilde{M}^{h+p}(c)$ of $\tilde{M}^m(c)$ as an anti-holomorphic submanifold.*

(3) *If $\lambda \neq 0$, then N is a mixed foliate CR-submanifold with $fD\bar{\eta} = 0$.*

Proof. For any vectors X, Y in TN , and Z in \mathfrak{D}^\perp , we have

$$-A_{JZ}X + D_X JZ = J\nabla_X Z + \sigma(X, Z).$$

Thus

$$(5.22) \quad D_X JZ = F\nabla_X Z + f\sigma(X, Z).$$

By applying Lemma 5.7, this gives

$$(5.23) \quad D_X JZ = F\nabla_X Z + \lambda\langle X, Z \rangle J\bar{\eta}.$$

Therefore by considering the normal component of $\tilde{\nabla}_X D_Y JZ$ we obtain

$$(5.24) \quad \begin{aligned} D_X D_Y JZ &= D_X(F\nabla_Y Z) + X(\lambda\langle Y, Z \rangle)J\bar{\eta} \\ &\quad - \lambda^2\langle Y, Z \rangle FX + \lambda\langle Y, Z \rangle fD_X \bar{\eta}. \end{aligned}$$

On the other hand, by equation (3.9) of [4] and Lemma 8.1 of [4] we have

$$D_X(F\nabla_Y Z) = f\sigma(X, \nabla_Y Z) - \sigma(X, P\nabla_Y Z) + F(\nabla_X \nabla_Y Z).$$

Substituting this into (5.24) we obtain

$$\begin{aligned} D_X D_Y JZ &= f\sigma(X, \nabla_Y Z) - \sigma(X, P\nabla_Y Z) + F(\nabla_X \nabla_Y Z) \\ &\quad + X(\lambda\langle Y, Z \rangle)J\bar{\eta} - \lambda\langle Y, Z \rangle\{FX - fD_X \bar{\eta}\}. \end{aligned}$$

Thus the normal curvature tensor R^\perp is given by

$$\begin{aligned} R^\perp(X, Y)JZ &= F(R(X, Y)Z) + f\sigma(X, \nabla_Y Z) - f\sigma(Y, \nabla_X Z) \\ &\quad - \sigma(X, P\nabla_Y Z) + \sigma(Y, P\nabla_X Z) - \lambda\langle [X, Y], Z \rangle J\bar{\eta} \\ &\quad + \{X(\lambda\langle Y, Z \rangle) - Y(\lambda\langle X, Z \rangle)\}J\bar{\eta} \\ &\quad - \lambda^2\{\langle Y, Z \rangle FX - \langle X, Z \rangle FY\} \\ &\quad + \lambda\{\langle Y, Z \rangle fD_X \bar{\eta} - \langle X, Z \rangle fD_Y \bar{\eta}\}. \end{aligned}$$

By applying Lemma 5.7 this gives

$$\begin{aligned}
 R^\perp(X, Y)JZ &= F(R(X, Y)Z) - \lambda\{\langle PX, P\nabla_Y Z \rangle - \langle PY, P_X Z \rangle\}J\bar{\eta} \\
 &\quad - \sigma(X, P\nabla_Y Z) + \sigma(Y, P\nabla_Y Z) \\
 &\quad + \{(X\lambda)\langle Y, Z \rangle - (Y\lambda)\langle X, Z \rangle\}J\bar{\eta} \\
 (5.25) \quad &\quad - \lambda^2\{\langle Y, Z \rangle FX - \langle X, Z \rangle FY\} \\
 &\quad + \lambda\{\langle Y, Z \rangle fD_X \bar{\eta} - \langle X, Z \rangle fD_Y \bar{\eta}\}.
 \end{aligned}$$

It follows from Lemma 5.7 that both $\sigma(X, P\nabla_Y Z)$ and $\sigma(Y, P\nabla_X Z)$ lie in $J\mathcal{D}^\perp$. Since $R^\perp(X, Y)JZ = 0$ by (5.1), equation (5.25) gives (5.20) and

$$(5.26) \quad (X\lambda)\langle Y, Z \rangle - (Y\lambda)\langle X, Z \rangle = \lambda\{\langle PX, P\nabla_Y Z \rangle - \langle PY, P\nabla_X Z \rangle\},$$

$$(5.27) \quad \lambda\{\langle Y, Z \rangle fD_X \bar{\eta} - \langle X, Z \rangle fD_Y \bar{\eta}\} = 0.$$

If N is a complex submanifold of $\tilde{M}^m(c)$, then $\mathcal{D} = TN$ and $\nu = T^\perp N$. Lemma 5.5 shows that N is a totally geodesic complex submanifold of $\tilde{M}^m(c)$.

Now we assume that N is *not* a complex submanifold. We have $\dim_{\mathbf{R}} \mathcal{D}^\perp = p > 0$.

Case (a). If $\mu \equiv 0$, then we have $\text{Im } \sigma \subseteq J\mathcal{D}^\perp$. Moreover, for any vector fields X in TN , Z in \mathcal{D}^\perp , and ξ in ν , Lemma 5.7 gives

$$0 = \langle \sigma(X, Z), \xi \rangle = \langle \tilde{\nabla}_X JZ, J\xi \rangle = \langle D_X JZ, J\xi \rangle.$$

Since this is true for all ξ in ν , $J\mathcal{D}^\perp$ is a parallel normal subbundle. Because the first normal spaces of N lie in $J\mathcal{D}^\perp$, the fundamental theorem of submanifolds shows that N lies in a totally geodesic complex submanifold $\tilde{M}^{h+p}(c)$ of $\tilde{M}^m(c)$. In this case, N is an anti-holomorphic submanifold of $\tilde{M}^{h+p}(c)$.

Case (b). If $\lambda \not\equiv 0$, then $N' = \{x \in N \mid \lambda(x) \neq 0\}$ is an open nonempty subset of N . Lemma 5.8 tells us that each component of N' is a mixed foliate CR-submanifold $\tilde{M}^m(c)$, $c \neq 0$.

If $c > 0$, then N is totally real (Lemma 4.3). Thus (5.26) gives

$$(5.28) \quad (X\lambda)\langle Y, Z \rangle - (Y\lambda)\langle X, Z \rangle = 0,$$

for any vectors X, Y in TN , and Z in \mathcal{D}^\perp . Because $\dim_{\mathbf{R}} \mathcal{D}_x^\perp = \dim_{\mathbf{R}} N \geq 2$ and λ^2 is differentiable, (5.28) implies that λ is a nonzero constant on N . Thus by (5.27) we get $fD\bar{\eta} = 0$.

If $c < 0$, then Proposition 4.4 and Lemma 5.8 show that $\dim_{\mathbf{R}} \mathcal{D}_x^\perp = p > 1$. Thus for any unit vector Z in \mathcal{D}^\perp there exists a unit vector W in \mathcal{D}^\perp so that $\langle Z, W \rangle = 0$. From (5.26) we find

$$(5.29) \quad Z(\lambda^2) = 0 \quad \text{for } Z \in \mathcal{D}^\perp.$$

Let X and Z be any unit vector fields in \mathfrak{D} and \mathfrak{D}^\perp respectively. Then (5.26) gives

$$(5.30) \quad X(\lambda)^2 = 2\lambda^2 \langle X, \nabla_Z Z \rangle.$$

On the other hand, for such X and Z we have

$$\langle X, \nabla_Z Z \rangle = \langle JX, \tilde{\nabla}_Z JZ \rangle = -\langle A_{JZ}Z, JX \rangle = -\langle \sigma(Z, JX), JZ \rangle.$$

Thus by using (5.30), Lemma 5.8, and the continuity of λ^2 we get $X(\lambda^2) \equiv 0$ for any vector X in \mathfrak{D} . Combining this with (5.29), we conclude that λ is a nonzero constant on N . The equation $fD\bar{\eta} = 0$ then follows from (5.27).

Lemma 5.10. *Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$, $c \neq 0$. If $\lambda \neq 0$, then the sectional curvature of N satisfies*

$$(5.31) \quad K(Z \wedge W) = \lambda^2$$

for any orthonormal vectors Z, W in \mathfrak{D}^\perp .

Proof. Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$, $c \neq 0$. If $\lambda \neq 0$, then N is mixed foliate (Lemma 5.8). For any vector U in TN , $PU \in \mathfrak{D}$. Thus for any orthonormal vectors Z, W in \mathfrak{D}^\perp , (5.20) of Lemma 5.9 gives

$$F(R(Z, W)Z) = -\lambda^2 FW.$$

From this we obtain (5.31).

Now we give the following classification theorem.

Theorem 5.11. *Let N be a CR-submanifold in a complex-space-form $\tilde{M}^m(c)$, $c \neq 0$. Then N has semi-flat normal connection in $\tilde{M}^m(c)$ if and only if N is one of the following:*

- (1) a totally geodesic complex submanifold $\tilde{M}^h(c)$,
- (2) a flat totally real submanifold of a totally geodesic complex submanifold $\tilde{M}^p(c)$ of $\tilde{M}^m(c)$,
- (3) a proper anti-holomorphic submanifold with flat normal connection in a totally geodesic complex submanifold $\tilde{M}^{h+p}(c)$ of $\tilde{M}^m(c)$,
- (4) a space of positive constant sectional curvature immersed in a totally geodesic complex submanifold $\tilde{M}^{p+1}(c)$ of $\tilde{M}^m(c)$ with flat normal connection as a totally real submanifold.

Proof. Let N be a CR-submanifold with semi-flat normal connection in $\tilde{M}^m(c)$, $c \neq 0$. If N is a complex submanifold of $\tilde{M}^m(c)$, N is a totally geodesic complex submanifold of $\tilde{M}^m(c)$ (Lemma 5.5). Thus N is itself a complex-space-form $\tilde{M}^h(c)$.

Assume that N is not a complex submanifold of $\tilde{M}^m(c)$. Then $p > 0$, and there exists a unit normal vector field $\bar{\eta}$ satisfies (5.18) and (5.19) for some constant λ (Lemmas 5.7 and 5.8).

If $\lambda = 0$ and N is totally real, (5.20) shows that N is flat.

If $\lambda = 0$ and N is neither complex nor totally real, then N lies in a totally geodesic complex submanifold $\tilde{M}^{h+p}(c)$ as an anti-holomorphic submanifold (Lemma 5.9). In this case, (5.1) implies that N has flat normal connection.

If $\lambda \neq 0$, Lemma 5.9 gives

$$(5.32) \quad D_X \bar{\eta} \in J\mathfrak{D}^\perp$$

for any vector X in TN . On the other hand, Lemma 5.7 also gives

$$(5.33) \quad D_X J\bar{\eta} = \tilde{\nabla}_X J\bar{\eta} = -JA_{\bar{\eta}}X + JD_X \bar{\eta}.$$

From Lemma 5.7 and (5.32) we see that $A_{\bar{\eta}}X \in \mathfrak{D}^\perp$, $JD_X \bar{\eta} \in TN$. Thus (5.33) gives

$$(5.34) \quad D\bar{\eta} \equiv 0.$$

Now, since N is mixed foliate (Lemma 5.8), the holomorphic distribution is integrable. Let N^T be a leaf of \mathfrak{D} . Denote by A^T and D^T the second fundamental tensor and normal connection of N^T in $\tilde{M}^m(c)$ as before. Then we have

$$-A_{\bar{\eta}}^T X + D_X^T \bar{\eta} = \tilde{\nabla}_X \bar{\eta} = -A_{\bar{\eta}} X + D_X \bar{\eta} = 0 \text{ for } X \in TN^T$$

by virtue of (5.34) and Lemma 5.7. This shows that $\bar{\eta}|_{N^T}$ is parallel in the normal bundle of N^T in $\tilde{M}^m(c)$. This contradicts a theorem of [5] unless N is totally real in $\tilde{M}^m(c)$. If N is totally real, N is of positive constant sectional curvature λ^2 (Lemma 5.10), and N has flat normal connection (Definition 5.1).

From (5.33) and (5.34) we find

$$(5.35) \quad D_X J\bar{\eta} = -JA_{\bar{\eta}}X \in J\mathfrak{D}^\perp$$

for any vector X in TN . Therefore by (5.21) of Lemma 5.9, (5.34) and (5.35), we see that $\mu = J\mathfrak{D}^\perp \oplus \text{Span}\{\bar{\eta}, J\bar{\eta}\}$ is a parallel normal subbundle, and $\mu \supseteq \text{Im } \sigma$. From these we conclude that N lies in a totally geodesic complex submanifold $M^{p+1}(c)$ of $\tilde{M}^m(c)$ as a totally real submanifold with flat normal connection.

The converse of this is trivial.

Remark 5.1. From Lemma 5.9 it follows that the assumption of compactness in Theorem 2 of [7] can be omitted.

References

- [1] A. Bejancu, *CR-submanifolds of a Kaehler manifold*. I, Proc. Amer. Math. Soc. **69** (1978) 135–142.
- [2] A. Bejancu, M. Kon & K. Yano, *CR-submanifolds of a complex space form*, J. Differential Geometry **16** (1981) 137–145.

- [3] E. Calabi, *Metric Riemann surfaces*, Annals of Math. Studies, No. 30, Princeton University Press, Princeton, 1953, 77–85.
- [4] B. Y. Chen, *CR-submanifolds of a Kaehler manifold. I*, J. Differential Geometry **16** (1981) 305–322.
- [5] B. Y. Chen & K. Ogiue, *Some extrinsic results for Kaehler submanifolds*, Tamkang J. Math. **4** (1973) 207–213.
- [6] K. Yano & M. Kon, *CR-sous-variétés d'un espace projectif complexe*, C. R. Acad. Sci. Paris **288** (1979) 515–517.
- [7] K. Yano, M. Kon & I. Ishihara, *Anti-invariant submanifolds with flat normal connection*, J. Differential Geometry **13** (1978) 577–588.

MICHIGAN STATE UNIVERSITY