

## A REGULARITY THEORY FOR HARMONIC MAPS

RICHARD SCHOEN & KAREN UHLENBECK

### 0. Introduction

In this paper we develop a regularity theory for energy minimizing harmonic maps into Riemannian manifolds. Let  $u: M^n \rightarrow N^k$  be a map between Riemannian manifolds of dimension  $n$  and  $k$ . It was shown by C. B. Morrey [17] in 1948 that if  $n = 2$ , then an energy minimizing harmonic map is Hölder continuous (and smooth if  $M$  and  $N$  are smooth). Since that time results have been found under special assumptions on  $N$ . Eells and Sampson [5] proved in 1963 that if  $N$  is compact and has nonpositive curvature, then every homotopy class of maps from a closed manifold  $M$  into  $N$  has a smooth harmonic representative. In the case where the image of the map is contained in a convex ball of  $N$ , there is a complete existence and regularity theory due to Hildebrandt and Widman [15] as well as Hildebrandt, Kaul and Widman [13]. Recently Giaquinta and Giusti obtained results for the case in which the image lies in a coordinate chart [9], [10].

In this paper we show that a bounded, energy minimizing map  $u: M^n \rightarrow N^k$  is regular (in the interior) except for a closed set  $\mathfrak{S}$  of Hausdorff dimension at most  $n - 3$ . We also show  $\mathfrak{S}$  is discrete for  $n = 3$ . Moreover, we derive techniques (see Theorem IV) for lowering the dimension of  $\mathfrak{S}$  under the condition that certain smooth harmonic maps of spheres into  $N$  are trivial. This can be checked in some interesting cases, for example if  $N$  has nonpositive curvature or if the image of the map lies in a convex ball of  $N$ , we show  $\mathfrak{S} = \emptyset$  and any minimizing harmonic map into such a manifold is smooth. Using our methods, it is possible to reduce the dimension of  $\mathfrak{S}$  if  $N$  is a sphere or Lie group by studying harmonic spheres in  $N$ . Our methods work for functionals which are the energy plus lower order terms, and thus have direct bearing on the question of the existence of global Coulomb gauges in nonabelian gauge theories.

We point out that there is a strong historical precedent for partial regularity results in problems involving elliptic systems (see Almgren [1], De Giorgi [3],

Giusti and Miranda [11] and Morrey [19]). Moreover, the types of singularities which arise in this problem have been observed in connection with other elliptic systems by De Giorgi [4] and others [11], and explicitly for harmonic maps by Hildebrandt, Kaul and Widman [13]. We also observe that the method we use for reducing the dimension of  $\mathfrak{S}$  is taken from H. Federer [7].

The problem which we deal with in this paper is to prove the regularity of vector-valued functions which minimize the ordinary Dirichlet integral subject to a family of smooth nonlinear (manifold) constraints. We use comparison maps as was done by Morrey for  $n = 2$ . The major difference is that for  $n > 2$  one cannot localize the problem in the image. Finding comparison maps which satisfy the constraints is a major technical difficulty of this paper. If one assumes that the image is a bounded domain in Euclidean space (without constraints), the comparison construction is straightforward. Our methods work for a restricted class of functionals because we need a scaling inequality (see Proposition 2.4) to construct maps satisfying the constraints. In a second paper we carry out a similar program to obtain complete boundary regularity for solutions to the Dirichlet problem. The reason is that in this case the obstruction to regularity is a harmonic map of a hemisphere which is constant on the boundary, and one can show that such a map is trivial. This generalizes previous results by Hildebrandt and Widman [15] and R. Hamilton [12] on the boundary value problem.

### 1. Statement of results

Let  $M^n$  and  $N^k$  be Riemannian manifolds of dimension  $n$  and  $k$ . For technical reasons we assume that  $N \subset R^k$  is isometrically embedded in Euclidean space. We assume throughout that  $M$  is compact, possibly with boundary, and that  $N$  is an open manifold. For  $r \geq 0$  let  $C^r(M, N)$  be the space of maps  $u: M \rightarrow N$  which have continuous derivatives through order  $r$ , so that  $C^r(M, N) \subseteq C^r(M, R^k)$  is a Banach submanifold. Likewise let  $C^{r,\alpha}(M, N)$  for  $\alpha \in (0, 1]$  denote the subset of  $C^r(M, N)$  whose  $r$ th derivatives are Hölder continuous with exponent  $\alpha$ .

In order to discuss harmonic maps from  $M$  to  $N$ , we work in the separable Hilbert space  $L_1^2(M, R^k)$ , the set of maps  $u: M \rightarrow R^k$  whose component functions have first derivatives in  $L^2$ . By  $L_{1,0}^2(M, R^k)$  we mean those  $L_1^2$  maps which are zero on  $\partial M$ . Define

$$L_1^2(M, N) = \{u \in L_1^2(M, R^k): u(x) \in N \text{ a.e. } x \in M\}.$$

Note that if  $\dim M = 1$ , then  $L_1^2(M, N)$  is a Hilbert submanifold of  $L_1^2(M, R^k)$ , but this is not so for  $\dim M > 1$ . The set  $L_1^2(M, N)$  inherits strong and weak

topologies from  $L^2_1(M, R^k)$ . Moreover, the space  $L^2_1(M, N)$  is a strongly closed subset with the additional property that the set  $\{u \in L^2_1(M, N) : \|u\|_{1,2} \leq C\}$  is weakly compact in  $L^2_1(M, R^k)$ .

For  $u \in L^2_1(M, R^k)$ , the energy functional is given by

$$E(u) = \int_M \langle du(x), du(x) \rangle dV = \int_M e(u).$$

Here the Lagrangian  $e(u)$  is given in local coordinates by

$$e(u) = \sum_{\alpha, \beta} \sum_i g^{\alpha\beta} \frac{\partial u^i}{\partial x^\beta} \frac{\partial u^i}{\partial x^\alpha} (\det g_{\gamma\delta})^{1/2} dx,$$

where  $g_{\alpha\beta}$  is the metric tensor of  $M$ . The norm on  $L^2_1(M, R^k)$  is then given by

$$\|u\|_{1,2}^2 = E(u) + \int_M \sum_i (u^i(x))^2 dV,$$

where  $dV$  is the volume element of  $M$ . A harmonic map  $u: M \rightarrow N$  is a weak solution of the Euler-Lagrange equation for  $E$  in  $L^2_1(M, N)$  (see (2.1)).

For certain applications of our results we will need to consider critical points of functionals with additional lower order terms. Let  $\tilde{E}(u) = E(u) + V(u)$  where  $V(u) = \int_M v(u)$ . In local coordinates,  $v(u)$  will have the form

$$v(u) = \left[ \sum_i \sum_\alpha \gamma_i^\alpha(x, u(x)) \frac{\partial u^i}{\partial x^\alpha}(x) + \Gamma(x, u(x)) \right] dV.$$

More invariantly, if  $\Theta$  is an open neighborhood of  $N$  in  $R^k$ , then  $\gamma \in C^r(M \times \Theta, T^*M \otimes R^k)$  and  $\Gamma \in C^r(M \times \Theta, R)$ . By an  $\tilde{E}$ -minimizing map we mean a map  $u \in L^2_1(M, N)$  such that  $\tilde{E}(u) \leq \tilde{E}(w)$  for any map  $w \in L^2_1(M, N)$  with  $(u - w) \in L^2_{1,0}(M, R^k)$ . Throughout the paper we will assume that the metric on  $M$  is  $C^2$  and  $\gamma, \Gamma \in C^r$  for  $r \geq 2$ . Our first result is a "sufficiently small" type result. The explicit dependence on parameters is derived in detail in Theorem 3.1. Here  $B_\sigma(a)$  is the geodesic ball about  $a$  in  $M$ .

**Theorem I.** *Let  $u \in L^2_1(B_\sigma(a), N)$  be an  $\tilde{E}$ -minimizing map such that  $u(x) \in N_0$  a.e. for some compact subset  $N_0 \subseteq N$ . If  $\sigma$  and  $\sigma^{2-n}\tilde{E}(u)$  are sufficiently small, then  $u$  is Hölder continuous on  $B_{\sigma/2}(a)$ .*

It is well known, but difficult to find in the literature (see [2]), that  $u$  is smooth in the interior of  $B_{\sigma/2}(a)$  once we have that  $u$  is continuous there.

For an  $\tilde{E}$ -minimizing map  $u$ , a point  $x \in M$  is a regular point if  $u$  is continuous in a neighborhood of  $x$ . Let  $\mathfrak{R} = \mathfrak{R}(u)$  be the set of all regular

points and  $\mathfrak{S} = \mathfrak{S}(u)$  be the complement of  $\mathfrak{R}$  in the interior of  $M$ . The singular set  $\mathfrak{S}$  is then obviously a closed subset of  $\text{int}(M)$ . The reader should refer to [6,2.10.2] for a discussion of Hausdorff dimension.

**Theorem II.** *Let  $u \in L^2_1(M, N)$  be  $\tilde{E}$ -minimizing with  $u(x) \in N_0$  a.e. for a compact set  $N_0 \subseteq N$ . It then follows that  $\dim(\mathfrak{S} \cap \text{int } M) \leq n - 3$  where  $n = \dim M$  and  $\dim A$  is the Hausdorff dimension of a set  $A$ . If  $n = 3$ , then  $\mathfrak{S}$  is a discrete set of points.*

We state the next results as a theorem only to motivate our results. Recall that a harmonic map is a weak solution to the Euler-Lagrange equations for  $E$  on  $L^2_1(M, N)$ . Suppose  $u \in L^2_{1,\text{loc}}(R^n, N)$  is a map such that  $\partial u / \partial r = 0$  a.e. Then there is a map  $w: S^{n-1} \rightarrow N$  such that  $u(x) = w(x/|x|)$ , and it is easy to see that  $u$  is harmonic if and only if  $w$  is harmonic. In fact  $E(u|B_r(0)) = (n-2)^{-1}r^{n-2}E(w)$ . Moreover, the map  $u$  has a singularity at 0 if and only if  $w$  is not a constant map.

**Theorem III.** *Let  $u \in L^2_1(M, N)$  be an  $\tilde{E}$ -minimizing map and let  $z \in \mathfrak{S} \cap \text{int } M$ . There exists a sequence  $\sigma_i \in R^+$ ,  $\sigma_i \rightarrow 0$  such that the maps  $u_i \in L^2_1(B_1(0), N)$ ,  $u_i(x) = u(\exp_z \sigma_i x)$  converge to  $u \in L^2_1(B_1(0), N)$ . The map  $u$  is a nonconstant harmonic map satisfying  $u(x) = w(x/|x|)$ ,  $w \in L^2_1(S^{n-1}, N)$  harmonic.*

A homogeneous harmonic map with an isolated singularity at 0 will be referred to as a *tangent map* (TM). A tangent map which is  $\tilde{E}$ -minimizing on compact subsets of  $R^n$  is a *minimizing tangent map* (MTM).

**Theorem IV.** *Suppose there is an integer  $l \geq 3$  such that every MTM from  $R^j \rightarrow N$  is trivial,  $3 \leq j \leq l$ . Then if  $u \in L^2_1(M, N)$  is  $\tilde{E}$ -minimizing with  $u(x) \in N_0$  a.e., then  $\dim(\mathfrak{S} \cap \text{int } M) \leq n - l - 1$ . If  $n = l + 1$ , then  $\mathfrak{S}$  is a discrete set of points, and if  $n < l + 1$ ,  $\mathfrak{S} = \emptyset$ .*

This theorem has the following corollary which is closely related to the work of Eells and Sampson [5] and Hildebrandt, Kaul and Widman [13].

**Corollary.** *If the sectional curvature of  $N$  is nonpositive or if  $u(M)$  is contained in a strictly convex ball of  $N$ , then  $\mathfrak{S} = \emptyset$ ; that is, any  $\tilde{E}$ -minimizing map  $u \in L^2_1(M, N)$  is smooth.*

*Proof.* To prove the corollary from Theorem IV, it suffices to show that any tangent map  $R^j \rightarrow N$  for  $j \geq 3$  is trivial; that is, any smooth harmonic  $u: S^{j-1} \rightarrow N$  is trivial. But this is elementary because if  $N$  has nonpositive curvature, we can lift  $u$  to a map  $\tilde{u}: S^{j-1} \rightarrow \tilde{N}$  where  $\tilde{N}$  is the universal cover of  $N$ . Since the square of the distance function  $\rho$  to a point is strictly convex we have that  $\rho^2 \circ u$  is a subharmonic function on  $S^{j-1}$  which is hence constant. Thus  $u$  is constant. The same argument works if  $u(M)$  is contained in a convex ball of  $N$ . This proves the corollary.

2. The Euler-Lagrange equation and scaling inequalities

We compute the Euler-Lagrange equation for  $\tilde{E}$  and show that  $\tilde{E}$ -minimizing maps are weak solutions of this equation. As in the previous section,  $N_0$  will be compact subset of  $N$ .

**Lemma 2.1.** *If  $u$  is  $\tilde{E}$ -minimizing on  $M$  and  $u(x) \in N_0$  a.e., then  $u$  satisfies the formal Euler-Lagrange equations for  $E$ . These equations have the form*

$$(2.1) \quad \Delta_M u - A(du, du) + \sum_{i,\alpha} B_{i,\alpha}(x, u(x)) \frac{\partial u^i}{\partial x^\alpha} + C(x, u(x)) = 0,$$

where  $A, B, C$  are smooth in their arguments, and  $A$  is quadratic in  $du$ .

*Proof.* We will say that a map is stationary for  $\tilde{E}$  if  $d\tilde{E}(u(t))/dt|_{t=0}$  for any differentiable curve of maps  $u: (-\varepsilon, \varepsilon) \rightarrow L^2_1(M, R^k)$  where  $u(t) \in L^2_1(M, N)$ ,  $u(0) = u$ , and  $u(t) - u \in L^2_{1,0}(M, R^k)$ . Of course it is true that an  $\tilde{E}$ -minimizing map is stationary. We will show that the space of admissible variations  $\{\psi = u'(0): u(t) \text{ satisfies the above condition}\}$  is large enough so that  $u$  satisfies the Euler-Lagrange equation for  $\tilde{E}$ . Note that it does not make sense to discuss differentiable curves in  $L^2_1(M, N)$  per se because  $L^2_1(M, N)$  does not have a local smooth structure.

Let  $\Theta$  be an open neighborhood of  $N$  in  $R^k$  such that the map  $\Pi: \Theta \rightarrow N$ , given by  $\Pi(y) =$  nearest point in  $N$  to  $y$ , is a smooth fibration. Since  $N_0$  is a compact set,  $\Theta$  contains a uniform neighborhood of  $N_0$ . Thus for  $t$  sufficiently small and any  $C^\infty$   $R^k$ -valued function  $\varphi$  which is zero on  $\partial M$ , we can define

$$u(t)(x) = \pi(u(x) + t\varphi(x)).$$

Observe that for  $x \in M$  the curve  $t \mapsto f(t) = u(t)(x)$  is smooth for small  $t$  and  $f'(t) = \varphi(X)d\Pi(u(x) + t\varphi(x))$  is uniformly bounded. Moreover,  $du(t)(x) = (du(x) + t d\varphi(x)) \cdot d\pi(u(x) + t\varphi(x))$  is smooth in  $t$  for almost all  $x$ . We find

$$\|u'(t)\|_{1,2} = \|\varphi(x) \cdot d\Pi(u(x) + t\varphi(x))\|_{1,2} \leq c(\varphi)[1 + \|du\|_{1,2}],$$

and  $u(t)$  is differentiable in  $L^2_1(M, R^k)$ . Note that

$$u'(t)(x)|_{t=0} = \varphi(x) \cdot d\Pi_{u(x)},$$

where  $d\Pi_y$  is the tangential projection  $R^k \rightarrow T_y N$  for  $y \in N$ . Let

$$\mathcal{E}(u) = -2\Delta_M u - d_M^* \gamma \circ u + (d_N \gamma \cdot du) + d_N \Gamma$$

be the expression for the Euler-Lagrange equation of the unconstrained problem. Then we have

$$\begin{aligned} \frac{d}{dt} \tilde{E}(u(t))|_{t=0} &= \int_M \langle \mathcal{E}(u), \Pi_{u(x)}(\varphi(x)) \rangle dV \\ &= \int_M \langle \Pi_{u(x)}(\mathcal{E}(u)), \varphi(x) \rangle dV = 0 \end{aligned}$$

for all  $\varphi$ . We find that, in a distributional sense,  $d\Pi_{u(x)}\mathfrak{E}(u(x)) = 0$ . Because

$$d\Pi_{u(x)}(\Delta_M u) = \Delta_M u - d^2\pi_{u(x)}(du, du) = \Delta_M u - A(du, du),$$

( $A$  is the second fundamental form of  $N$  in  $R^k$ ), the equations can be put in the required form. We are not really interested in the exact form of the lower order terms. This completes the proof of Lemma 2.1.

Our theorems will be proved by covering  $M$  with geodesic coordinate balls and proving regularity for  $\tilde{E}$ -minimizing maps on balls. Let  $B_1 = B_1^n(0)$  be the unit ball in  $R^n$ . For  $\Lambda > 0$ , let  $\mathfrak{F}_\Lambda$  denote the class of functionals  $\tilde{E}$  on  $B_1$  with metric  $g_{\alpha\beta}$  such that  $g_{\alpha\beta}(0) = \delta_{\alpha\beta}$  and lower order terms satisfying, for  $x \in B_1$ ,  $u \in N_0$ ,

$$(2.2) \quad \sum_{\alpha, \beta, \tau} \left| \frac{\partial}{\partial x^\tau} g_{\alpha\beta}(x) \right| + |\gamma(x, u)| + |d_u \gamma(x, u)| \\ + |\Gamma(x, u)|^{1/2} + |d_u \Gamma(x, u)|^{1/2} \leq \Lambda.$$

If  $u$  is  $\tilde{E}$ -minimizing for  $\tilde{E} \in \mathfrak{F}_\Lambda$  and  $u(x) \in N_0$  a.e., we say  $u \in \mathfrak{J}_\Lambda$ . The lower order terms are handled by showing that  $\Lambda$  is a dimensional constant which shrinks with the radius of a coordinate ball. In fact, if  $\tilde{E}$  is a functional and  $B_\sigma(\rho)$  is a geodesic ball in  $M$  of radius  $\sigma$  centered at  $p \in M$ , we define a functional  $\tilde{E}^{p, \sigma}$  on  $B_1$  by setting

$$(2.3) \quad \tilde{E}^{p, \sigma}(w) = \int_{B_1} (|dw|_{g_\sigma}^2 + \sigma(dw \cdot \gamma(y, w)) + \sigma^2 \Gamma(y, w)) g_\sigma^{1/2} dy \\ = \sigma^{2-n} \tilde{E}_{B_\sigma(\rho)}(u),$$

where  $w(y) = u(\sigma y)$  and  $g_\sigma(y) = g(\sigma y)$ . Since  $M$  and  $N_0$  are compact, we can choose  $\Lambda$  so that  $\tilde{E}^{p, \sigma} \in \mathfrak{F}_\Lambda$  for all  $p$  and some  $\sigma > 0$ . It follows that if  $\tilde{E}^{p, \sigma} \in \mathfrak{F}_\Lambda$ , then  $\tilde{E}^{p, \lambda\sigma} \in \mathfrak{F}_{\lambda\Lambda}$  for any  $\lambda \in (0, 1]$ . We state the following.

**Lemma 2.2.** *Given  $\Lambda > 0$ , there exists  $\sigma_0 > 0$  such that for  $0 < \sigma \leq \sigma_0$  and  $p \in M$ , if  $u$  is  $\tilde{E}$ -minimizing, then  $w(y) = u(\exp_p \sigma y)$  is  $\tilde{E}^{p, \sigma}$ -minimizing where  $\tilde{E}^{p, \sigma} \in \mathfrak{F}_\Lambda$ .*

Thus we restrict our attention to  $\tilde{E} \in \mathfrak{F}_\Lambda$  where  $\Lambda$  is small. Let  $E$  be the energy functional in the Euclidean metric on  $B_1$ . Let  $E_\sigma, \tilde{E}_\sigma$  denote energies taken over  $B_\sigma$ ,  $0 < \sigma \leq 1$ . The inequalities

$$|E_\alpha(u) - \tilde{E}_\sigma(u)| \leq c\Lambda(\sigma E_\alpha(u) + \sigma^{n/2} E_\alpha(u)^{1/2} + \Lambda\sigma^n) \\ \leq \frac{3}{2}c\Lambda(\sigma E_\alpha(u) + \sigma^{n-1})$$

are straightforward provided  $\Lambda\sigma \leq 1$ . Consequently for  $\sigma \in (0, 1]$  we have

$$\begin{aligned} \tilde{E}_\sigma(u) &\leq (1 + \bar{c}\Lambda\sigma)E_\alpha(u) + \bar{c}\Lambda\sigma^{n-1}, \\ E_\alpha(u) &\leq (1 + \bar{c}\Lambda\sigma)\tilde{E}_\sigma(u) + \bar{c}\Lambda\sigma^{n-1}, \end{aligned}$$

provided  $c\Lambda \leq \frac{1}{2}$ . We have the following.

**Lemma 2.3.** *If  $\Lambda$  is sufficiently small, and  $u$  is  $\tilde{E}$ -minimizing for  $\tilde{E} \in \mathfrak{F}_\Lambda$  (i.e.,  $u \in \mathfrak{H}_\Lambda$ ), then there exists a constant  $c = c(n) > 0$  such that for  $\sigma \in (0, 1]$*

$$E_\alpha(u) \leq (1 + c\Lambda\sigma)E_\alpha(w) + c\Lambda\sigma^{n-1}$$

for any  $w \in L^2_1(B_1, N)$  with  $w = u$  on  $B_1 \sim B_\sigma$ .

*Proof.* Since  $\tilde{E}_\sigma(u) \leq \tilde{E}_\sigma(w)$ , this follows directly from the above inequalities.

We can now prove the first basic inequality of the paper. We use the notation

$$E_\sigma^x(u) = \int_{B_\sigma(x)} |du|^2(y) dy.$$

**Proposition 2.4.** *Let  $u \in \mathfrak{H}_\Lambda$  for  $\Lambda$  sufficiently small. Then we have*

$$\sigma^{2-n}E_\sigma^x(u) \leq c[\rho^{2-n}E_\rho^x(u) + \Lambda\rho]$$

for  $x \in B_{1/2}$ ,  $0 < \sigma \leq \rho \leq \frac{1}{2}$ .

*Proof.* By rescaling as discussed above, we can work on  $B_1$  instead of  $B_{1/2}(x)$ . This will introduce at worst an extra multiplicative factor. For almost all  $\sigma \in (0, 1]$  we have  $\int_{|x|=\sigma} |du|^2 d\xi < \infty$  where  $\xi$  is a variable on the sphere. Introduce the comparison map

$$\begin{aligned} v_\sigma(x) &= u(x), & |x| \geq \sigma, \\ v_\sigma(x) &= u(\sigma x/|x|), & |x| \leq \sigma. \end{aligned}$$

Since the result is trivial for  $n = 2$ , we assume  $n > 2$ . Denote by  $|d_\xi u|^2$  the tangential energy along the spheres  $|x| = r$ , so that  $|du|^2 = |d_\xi u|^2 + |\partial u/\partial r|^2$ . We compute

$$\begin{aligned} E_\alpha(v_\sigma) &= (n-2)^{-1}\sigma \int_{|x|=\sigma} |d_\xi u|^2 d\xi \\ &= (n-2)^{-1}\sigma \left( \frac{d}{d\sigma} E_\alpha(u) - \int_{|x|=\sigma} \left| \frac{\partial u}{\partial r} \right|^2 d\xi \right). \end{aligned}$$

From Lemma 2.3 we get, with  $\bar{c} = c\Lambda$ ,

$$\begin{aligned} E_\alpha(u) &\leq (1 + \bar{c}\sigma)E_\alpha(v_\sigma) + \bar{c}\sigma^{n-1} \\ &\leq (n-2)^{-1}\sigma(1 + \bar{c}\sigma) \left[ \frac{d}{d\sigma} E_\alpha(u) - \int_{|x|=\sigma} \left| \frac{\partial u}{\partial r} \right|^2 d\xi \right] + \bar{c}\sigma^{n-1}. \end{aligned}$$

This implies

$$(2.4) \quad 0 \leq \sigma^{2-n} \int_{|x|=\sigma} \left| \frac{\partial u}{\partial r} \right|^2 d\xi \leq \frac{d}{d\sigma} \left[ (1 + \bar{c}\sigma)^{n-2} \sigma^{2-n} E_\alpha(u) \right] + \bar{c}.$$

Since  $E_\alpha(u)$  is a nondecreasing function, we can integrate this inequality from  $\sigma$  to  $\rho$

$$(2.5) \quad (1 + \bar{c}\sigma)^{n-2} \sigma^{2-n} E_\alpha(u) \leq (1 + \bar{c}\rho)^{n-2} \rho^{2-n} E_\rho(u) + \bar{c}(\rho - \sigma),$$

where we have discarded the radial derivative term. This implies directly the conclusion of Proposition 2.4. Note that  $\bar{c}$  is a constant times  $\Lambda$ .

If we take the radial derivative term into consideration in the above argument we can prove more. Note that if we set  $u_\lambda(x) = u(\lambda x)$  for  $\lambda \in (0, 1]$ , then as in (2.3) we have  $u_\lambda \in \mathcal{H}_{\lambda\Lambda}$  for  $u \in \mathcal{H}_\Lambda$ , and

$$(2.6) \quad E_1(u_\lambda) = \lambda^{2-n} E_\lambda(u).$$

**Lemma 2.5.** *There is a sequence  $\lambda(i) \rightarrow 0$ ,  $\lambda(i) \in (0, 1]$ , such that  $u_{\lambda(i)}$  converge weakly in  $L^2_1(B_1, N)$  to a limiting map  $u_0 \in L^2_1(B_1, N)$ . The map  $u_0$  is a harmonic map satisfying  $\partial u_0 / \partial r = 0$  a.e. in  $B_1$ .*

*Proof.* From the previous result and (2.6),  $E_1(u_\lambda)$  is bounded for  $\lambda \in (0, 1]$ , and therefore we get a weakly convergent sequence  $u_{\lambda(i)} \rightarrow u_0 \in L^2_1(B_1, N)$ . Since  $u_{\lambda(i)} \in \mathcal{H}_{\lambda(i)\Lambda}$ , it satisfies Euler equations of the form (2.1).

It follows easily that  $u_0$  satisfies the Euler-Lagrange equation for  $E$  and hence  $u_0$  is harmonic. To see that  $\partial u_0 / \partial r = 0$  a.e., first note that (2.5) implies the existence of a number  $L_0$  with

$$(2.7) \quad L_0 = \lim_{\sigma \rightarrow 0} \sigma^{2-n} E_\alpha(u) = \lim_{\sigma \rightarrow 0} E_1(u_\sigma).$$

If we integrate (2.4) from 0 to  $\lambda$  keeping the radial derivative term, we have

$$\int_{B_\lambda} r^{2-n} \left| \frac{\partial u}{\partial r} \right|^2 dx \leq \left[ (1 + \bar{c}\lambda)^{n-2} \lambda^{2-n} E_\lambda(u) - L_0 \right] + \bar{c}\lambda.$$

By a change of variables

$$\int_{B_1} r^{2-n} \left| \frac{\partial u_\lambda}{\partial r} \right|^2 dx = \int_{B_\lambda} r^{2-n} \left| \frac{\partial u}{\partial r} \right|^2 dx.$$

Therefore we have

$$\lim_{\lambda \rightarrow 0} \int_{B_1} r^{2-n} \left| \frac{\partial u_\lambda}{\partial r} \right|^2 dx = 0,$$

which implies that  $\partial u_0 / \partial r = 0$  a.e., for any weak limit  $u_0$ . This proves Lemma 2.5.



In §4 we will show that this convergence is actually norm convergence. To do that will require some preliminary regularity results. The first is (see Theorem 3.1)

**Regularity Estimate 2.6.** *There exists  $\bar{\epsilon} > 0$  depending only on  $n$  and  $N_0 \subseteq N$  such that if  $u \in \mathcal{H}_\Lambda$ ,  $\Lambda \leq \epsilon$ , and  $E_1(u) \leq \bar{\epsilon}$ , then  $u$  is Hölder continuous on  $B_{1/2}$  and satisfies  $|u(x) - u(y)| \leq c|x - y|^\alpha$  for  $x, y \in B_{1/2}$  where  $c, \alpha > 0$  depend only on  $n, N_0$ .*

We first note that Theorem I follows immediately from this result by rescaling, see Lemma 2.2. Secondly, we can immediately prove the corollary.

**Corollary 2.7.** *If  $u \in L^2_1(B_1, N)$  is in  $\mathcal{H}_\Lambda$  and  $\mathcal{S}$  is the singular set of  $u$ , then  $\mathcal{H}^{n-2}(\mathcal{S} \cap B_{1/2}) = 0$ . More generally, if  $u \in L^2_1(M, N)$  is  $\bar{E}$ -minimizing, then  $\mathcal{H}^{n-2}(\mathcal{S} \cap \text{int } M) = 0$ .*

*Proof.* By rescaling, the second statement follows from the first. For  $x \in \mathcal{S} \cap B_{1/2}$ , choose normal coordinates  $y$  centered at  $x$ . Let  $u_{x,\lambda}(y) = u(\exp_x \lambda y)$ . Then the maps  $u_{x,\lambda}$  are in  $\mathcal{H}_{\lambda\Lambda}$  (see Lemma 2.2). By the regularity estimate we have

$$(2.8) \quad \bar{\epsilon} \leq E_1(u_{x,\lambda}) = \lambda^{2-n} \int_{B_\lambda(x)} |du|^2 dx$$

for all  $x \in \mathcal{S} \cap B_{1/2}$ ,  $\lambda\Lambda \leq \bar{\epsilon}$ . For  $\delta \in (0, \bar{\epsilon}/\Lambda)$ , let  $\{B_\delta(x_1), \dots, B_\delta(x_l)\}$  be a maximal family of  $l = l(\delta)$  disjoint balls of radius  $\delta$  with center  $x_i \in \mathcal{S} \cap B_{1/2}$ . By maximality,  $\mathcal{S} \cap B_{1/2} \subseteq \cup_{j=1}^l B_{2\delta}(x_j)$ . Applying (2.8) on each ball and summing, we get

$$(2.9) \quad l\delta^{n-2} \leq \bar{\epsilon}^{-1} \int_{\cup_i B_\delta(x_i)} |du|^2 dx \leq \bar{\epsilon}^{-1} E(u).$$

Since  $\mathcal{S} \cap B_{1/2} \subseteq \cup_i B_{2\delta}(x_i)$ , we see that  $\mathcal{H}^{n-2}(\mathcal{S} \cap B_{1/2}) \leq cE(u)$ . In particular  $\mathcal{H}^n(\cup_i B_\delta(x_i)) \leq \bar{c}\delta^2 E(u)$ , and by the dominated convergence theorem

$$\lim_{\delta \rightarrow 0} \int_{\cup_i B_\delta(x_i)} |du|^2 dx = 0.$$

Using this in (2.9) then shows  $\mathcal{H}^{n-2}(\mathcal{S} \cap B_{1/2}) = 0$ .

### 3. The $\epsilon$ -regularity theorem

In this section we prove regularity of minimizing maps under the assumption that the energy is small. Precisely we prove

**Theorem 3.1.** *There exists a constant  $\bar{\epsilon} = \bar{\epsilon}(n, N_0)$  such that if  $u \in \mathcal{H}_\Lambda$ ,  $\Lambda \leq \bar{\epsilon}$ , and  $E_1(u) \leq \bar{\epsilon}$ , then  $u$  is Hölder continuous on  $B_{1/2}$  and  $|u(x) - u(y)| \leq c|x - y|^\alpha$  for  $x, y \in B_{1/2}$  where  $\alpha = \alpha(n) > 0$  and  $c = c(n, N_0)$ .*

We will prove this theorem by establishing energy decay estimates on small balls. We first show that if  $\bar{\epsilon}$  is chosen small, it is possible to approximate  $u$  by smooth maps into  $N$ . To see this we let  $\varphi: R^n \rightarrow R^+$  be a smooth radial mollifying function so that  $\text{support}(\varphi) \subseteq B_1$  and  $\int_{R^n} \varphi(x) dx = 1$ . We then note that if  $u^* = \int_{B_1} \varphi(x)u(x) dx$ , we can apply a version of the Poincaré inequality to assert

$$\int_{B_1} |u - u^*|^2 dx \leq c_1 E_1(u) \leq c_1 \bar{\epsilon}.$$

(Throughout this section  $c_1, c_2, \dots$  will denote constants depending only on  $n, N_0$ .) This inequality implies in particular that  $u^*$  lies near many values of  $u(x)$  for  $x \in B_1^n$ . Hence in particular we see that  $\text{dist}(u^*, N) \leq c_2(\bar{\epsilon})^{1/2}$ . This inequality gains in power when it is combined with the scaling inequality Proposition 2.4, for we can apply it on the ball  $B_h^n(x)$  for any  $x \in B_{1/2}^n$ ,  $0 < h \leq \frac{1}{4}$ . That is, we apply it to the scaled map  $u_{x,h}: B_1^n \rightarrow N$  given by

$$u_{x,h}(y) = u(x - hy).$$

By Proposition 2.4 we have

$$E_1(u_{x,h}) = h^{2-n} E_{B_h(x)}(u) \leq c_3 E_1(u) + c_3 \bar{\epsilon} \leq c_4 \bar{\epsilon},$$

provided  $x \in B_{1/2}$ ,  $h \in (0, \frac{1}{4}]$ . Thus if we set

$$u^{(h)}(x) = \int_{B_1^n} \varphi(y)u(x - hy) dy = \int_{B_1^n} \varphi^{(h)}(x - z)u(z) dz$$

where  $\varphi^{(h)}(x) = h^{-n}\varphi(x/h)$ , we have

$$(3.1) \quad \text{dist}(u^{(h)}(x), N_0) \leq c_5 \bar{\epsilon}^{1/2}$$

for any  $x \in B_{1/2}$ ,  $h \in (0, \frac{1}{4}]$ . Let  $\mathcal{O}$  be a normal neighborhood of  $N$  in  $R^k$ , and let  $\Pi: \mathcal{O} \rightarrow N$  denote the smooth nearest point projection map. Since  $N_0$  is compact,  $\mathcal{O}$  contains a uniform neighborhood of  $N_0$ . By (3.1), if  $\bar{\epsilon}$  is chosen small we will have  $u^{(h)}(x) \in \mathcal{O}$  for all  $x \in B_{1/2}$ , and we can define a smooth map  $u_h: B_{1/2} \rightarrow N$  by  $u_h(x) = \Pi \circ u^{(h)}(x)$ . We note the following result.

**Lemma 3.2.** *Let  $\bar{h} = \bar{\epsilon}^{1/4}$ , and suppose  $h \in (0, \bar{h}]$ . Then we have*

$$\int_{B_{1/2}} |du^{(h)}|^2 dx \leq c_6 E_1(u),$$

$$\sup_{x \in B_{1/2}} |u^{(\bar{h})}(x) - u^{(\bar{h})}(0)|^2 \leq c_6 \bar{\epsilon}^{1/2}.$$

*Proof.* The first inequality is standard. To prove the second, observe

$$\begin{aligned} |du^{(\bar{h})}|^2(x) &= \left| \int_{B_1} \varphi^{(\bar{h})}(x-y) du(y) \right|^2 dy \\ &\leq \int_{B_1} \varphi^{(\bar{h})}(x-y) |du|^2(y) dy. \end{aligned}$$

By Proposition 2.4 this implies

$$|du^{(\bar{h})}|^2(x) \leq c_7 \bar{h}^{-n} E_{B_{\bar{h}}}(u) \leq c_8 \bar{h}^{-2} \bar{\varepsilon} = c_8 \bar{\varepsilon}^{1/2}$$

for any  $x \in B_{1/2}$ . This implies the second inequality and completes the proof of Lemma 3.2.

In order to compare our approximating maps to  $u$ , we must force them to agree with  $u$  on the boundary of some small ball. To achieve this we observe that inequality (3.1) is a pointwise inequality, that is,  $x$  is fixed and  $h$  arbitrarily small. Thus we can choose  $h = h(x)$  and the inequality still holds for each  $x$ . Let  $\tau = \bar{\varepsilon}^{1/8}$ , and suppose  $\theta \in (\tau, \frac{1}{4})$ . We choose  $h = h(r)$ ,  $r = |x|$  to be a nonincreasing smooth function of  $r$  satisfying

$$(3.2) \quad h(r) = \bar{h} \text{ for } r \leq \theta, h(\theta + \tau) = 0, |h'(r)| \leq 2\bar{\varepsilon}^{1/8}.$$

We can then set

$$u^{(h(x))}(x) = \int_{B_1} \varphi^{(h(x))}(x-y) u(y) dy,$$

and by (3.1),  $u_{h(x)}(x) = \Pi \circ u^{(h(x))}(x)$ . We can prove the following result.

**Lemma 3.3.** For  $\theta \in (\tau, \frac{1}{4}]$ , the map  $u_h$  is in  $L^2_1(B_{1/2}, N)$  and satisfies  $u_h = u$  on  $B_{1/2} \sim B_{\theta+\tau}$ , and

$$\int_{B_{\theta+\tau} \sim B_\theta} |du_h|^2 dx \leq c_9 \int_{B_{\theta+2\tau} \sim B_{\theta-\tau}} |du|^2 dx.$$

*Proof.* Since  $\Pi$  is a smooth map, it suffices to prove the lemma for  $u^{(h)}$  instead of  $u_h$ . We first note that by a change of variable,  $u^{(h)}(x)$  can be written

$$u^{(h)}(x) = \int_{B_1} \varphi(y) u(x - h(x)y) dy.$$

From this expression it is clear that  $u^{(h)}$  is smooth if  $u$  is smooth. We first consider a smooth  $u: B_{1/2} \rightarrow R^k$ . If  $\Omega$  is a domain compactly contained in  $B_{1-\tau}$ , then we compute

$$\frac{\partial u^{(h)}}{\partial x^\alpha} = \int_{B_1} \varphi(y) \left[ \frac{\partial u}{\partial x^\alpha}(x - hy) - \frac{\partial h}{\partial x^\alpha} y \cdot \nabla u(x - hy) \right] dy.$$

Thus it follows that

$$\int_{\Omega} |du^{(h)}|^2 dx \leq c_{10} \int_{\Omega} \int_{B_1} \varphi(y) |du|^2(x - hy) dy dx.$$

For  $x \in \Omega$ , we let  $z = x - h(x)y$ . This defines a map  $F_y: \Omega \rightarrow \Omega_{\tau}$ , where we let  $\Omega_{\tau} = \{x: \text{dist}(x, \Omega) < \tau\}$ . By (3.2) we see that  $F_y$  is a diffeomorphism onto  $F_y(\Omega)$  with Jacobian approximately one. Thus by change of variables we can estimate

$$\int_{\Omega} |du|^2(x - h(x)y) dx \leq 2 \int_{\Omega_{\tau}} |du|^2 dx.$$

Therefore we have

$$(3.3) \quad \int_{\Omega} |du^{(h)}|^2 dx \leq c_{11} \int_{\Omega_{\tau}} |du|^2 dx.$$

Now if  $u_i$  is a sequence of smooth maps  $B_{3/4} \rightarrow R^k$  converging strongly to  $u$  in  $L^2_1(B_{3/4}, N)$ , then (3.3) implies

$$\int_{B_{1/2}} |du_i^{(h)} - du_j^{(h)}|^2 dx \leq c_{11} \int_{B_{3/4}} |du_i - du_j|^2 dx,$$

and hence  $\{u_i^{(h)}\}$  is Cauchy in  $L^2_1(B_{1/2}, R^k)$ . From this it follows that  $\lim_{i \rightarrow \infty} u_i^{(h)} = u^{(h)}$ , and  $u^{(h)} = u$  on  $B_{1/2} \sim B_{\theta+\tau}$ . We can then apply (3.3) with  $\Omega = B_{\theta+\tau} \sim B_{\theta}$  to get the conclusion of Lemma 3.3.

In order to prove Theorem 3.1, it suffices by Morrey's Lemma [16,2.4.1] to prove that

$$r^{2-n} E_{B_r(x)}(u) \leq c_{12} r^{2\alpha}$$

for any  $x \in B_{1/2}$ ,  $r \in (0, \frac{1}{4}]$ . We will prove that if  $\bar{\epsilon}$  is sufficiently small, then we have

$$(3.4) \quad r^{2-n} E_r(u) \leq c_{12} r^{2\alpha}$$

for  $r \in (0, \frac{1}{2}]$ . The previous estimate can then be gotten by reapplying (3.4) with varying center point. We now state a result which is a discrete version of (3.4).

**Proposition 3.4.** *There exists  $\bar{\epsilon} = \bar{\epsilon}(n, N_0) > 0$  such that if  $u \in \mathcal{H}_{\Lambda}$ ,  $\Lambda \leq \bar{\epsilon}$ , and  $E_1(u) \leq \bar{\epsilon}$ , then we have*

$$\bar{\theta}^{2-n} E_{\bar{\theta}}(u) + \bar{\theta} \Lambda \leq \frac{1}{2} (E_1(u) + \Lambda)$$

for some  $\bar{\theta} = \bar{\theta}(n, N_0) \in (0, 1)$ .

*Proof of Theorem 3.1.* We show how Theorem 3.1 follows from Proposition 3.4. We will prove (3.4) by an iterative procedure. Observe that the scaled map

$u_{\bar{\theta}}(x) = u(\bar{\theta}x)$  lies in  $\mathcal{H}_{\bar{\theta}\Lambda}$  and from Proposition 3.4 we have

$$E_1(u_{\bar{\theta}}) = \bar{\theta}^{2-n} E_{\bar{\theta}}(u) \leq \bar{\epsilon}.$$

Thus Proposition 3.4 is applicable to  $u_{\bar{\theta}}$ , and we get

$$\bar{\theta}^{2-n} E_{\bar{\theta}}(u_{\bar{\theta}}) + \bar{\theta}^2 \Lambda \leq \frac{1}{2} (E_1(u_{\bar{\theta}}) + \bar{\theta} \Lambda).$$

But this is the same as

$$(\bar{\theta}^2)^{2-n} E_{\bar{\theta}^2}(u) + \bar{\theta}^2 \Lambda \leq \frac{1}{2} (\bar{\theta}^{2-n} E_{\bar{\theta}}(u) + \bar{\theta} \Lambda).$$

Applying Proposition 3.4 on the right we get

$$(\bar{\theta}^2)^{2-n} E_{\bar{\theta}^2}(u) + \bar{\theta}^2 \Gamma L \leq (\frac{1}{2})^2 (E_1(u) + \Lambda).$$

Repeating this argument  $i$  times we get

$$(\bar{\theta}^i)^{2-n} E_{\bar{\theta}^i}(u) + \bar{\theta}^i \Lambda \leq 2^{-i} (E_1(u) + \Lambda)$$

for any nonnegative integer  $i$ . Given any  $r \in (0, 1)$ , there is an integer  $i$  so that  $r \in [\bar{\theta}^{i+1}, \bar{\theta}^i]$ . Setting  $\alpha = (\log 2)/(2 \log \bar{\theta}^{-1})$ , we have

$$(\bar{\theta}^i)^{2-n} E_r(u) + r \Lambda \leq (\bar{\theta}^i)^{2\alpha} (E_1(u) + \Lambda)$$

for any  $r \in (0, 1)$ . This implies

$$r^{2-n} E_r(u) \leq \bar{\theta}^{2-n-2\alpha} r^{2\alpha} (E_1(u) + \Lambda),$$

which verifies (3.4). As we have observed, this proves Theorem 3.1.

*Proof of Proposition 3.4.* Let  $v$  be the solution of the linear Dirichlet problem

$$\begin{aligned} \Delta v &= 0, & \text{in } B_{1/2}, \\ v &= u^{(\bar{h})} & \text{on } \partial B_{1/2}. \end{aligned}$$

Thus  $v: B_{1/2} \rightarrow R^k$  is a smooth harmonic map. We observe the following properties of  $v$ . First we note that by Lemma 3.2,  $u^{(\bar{h})}(B_{1/2}) \subseteq B_{c\bar{\epsilon}^{1/4}}^k(u^{(\bar{h})}(0))$ . Thus it follows that the image of  $v$  is also contained in this ball. In particular we have

$$(3.5) \quad \sup_{B_{1/2}} |v - u^{(\bar{h})}|^2 \leq c_{13} \bar{\epsilon}^{1/2}.$$

The mean value inequality for subharmonic functions implies the result

$$\sup_{B_{1/2}} |dv|^2 \leq c_{14} \int_{B_{1/2}} |dv|^2.$$

Since  $v$  minimizes energy on  $B_{1/2}$  for its boundary, we have

$$\int_{B_{1/2}} |dv|^2 \leq \int_{B_{1/2}} |du^{(\bar{h})}|^2 \leq E_1(u)$$

by Lemma 3.2. Therefore, we have

$$(3.6) \quad \sup_{B_{1/4}} |dv|^2 \leq c_{15} E_1(u).$$

For any  $\theta \in (0, \frac{1}{4}]$  we can estimate

$$(3.7) \quad \begin{aligned} \theta^{2-n} E_\theta(u_{\bar{h}}) &\leq c_{16} \theta^{2-n} E_\theta(u^{(\bar{h})}) \\ &\leq 2c_{16} \theta^{2-n} \int_{B_\theta} \{ |d(u^{(\bar{h})} - v)|^2 + |dv|^2 \} dx, \end{aligned}$$

where we have used the inequality (3.1) and the smoothness of  $\Pi$ . By (3.6) we see that

$$(3.8) \quad \theta^{2-n} \int_{B_\theta} |dv|^2 \leq c_{17} \theta^2 E_1(u)$$

for any  $\theta \in (0, \frac{1}{4})$ . Integrating by parts we have

$$\int_{B_{1/2}} |d(u^{(\bar{h})} - v)|^2 = - \int_{B_{1/2}} (u^{(\bar{h})} - v) \cdot \Delta(u^{(\bar{h})} - v).$$

Using (3.5) and the harmonic property of  $v$  we get

$$(3.9) \quad \int_{B_{1/2}} |d(u^{(\bar{h})} - v)|^2 \leq c_{18} \bar{\epsilon}^{1/4} \int_{B_{1/2}} |\Delta u^{(\bar{h})}|.$$

The Euler equation (2.1) for  $u$  tells us

$$\begin{aligned} \Delta u^{(\bar{h})}(x) &= \int_{\mathbb{R}^n} [\Delta_x \varphi^{(\bar{h})}(x-y)] u(y) dy \\ &= \int_{\mathbb{R}^n} [\Delta_y \varphi^{(\bar{h})}(x-y)] u(y) dy \\ &= \int_{\mathbb{R}^n} \varphi^{(\bar{h})}(x-y) [A(du, du) - B du - C] dy. \end{aligned}$$

From the form of  $B, C$  we conclude

$$|\Delta u^{(\bar{h})}(x)| \leq c_{19} \int_{\mathbb{R}^n} \varphi^{(\bar{h})}(x-y) [|du|^2 + \Lambda] dy.$$

Integrating over  $x \in B_{1/2}$  we finally have

$$\int_{B_{1/2}} |\Delta u^{(\bar{h})}| \leq c_{19} (E_1(u) + \Lambda).$$

Combining this with (3.7), (3.8), and (3.9) gives

$$(3.10) \quad \theta^{2-n} E_\theta(u_{\bar{h}}) \leq c_{20} \theta^{2-n} \bar{\epsilon}^{1/4} (E_1(u) + \Lambda) + c_{20} \theta^2 E_1(u)$$

for any  $\theta \in (0, \frac{1}{4})$ .

Let  $\gamma_n \in (0, 1/16]$  be a number to be chosen depending only on  $n$ , and let  $\bar{\theta} = \bar{\varepsilon}^{\gamma_n}$ . Let  $p$  be the greatest integer less than or equal to  $\bar{\theta}/(3\tau)$  where  $\tau = \bar{\varepsilon}^{1/8}$  and write

$$[\bar{\theta}, \bar{\theta} + 3p\tau] = \bigcup_{i=1}^p I_i, \quad |I_i| = 3\tau,$$

where each  $I_i$  is a closed interval of length  $3\tau$ . Since  $\gamma_n \leq \frac{1}{16}$ , we have  $p \geq \frac{1}{3}\bar{\varepsilon}^{-1/16} - 1$ . We have

$$\int_{r \in [\theta, \theta + 3p\tau]} |du|^2 dx = \sum_{i=1}^p \int_{r \in I_i} |du|^2 dx \leq E_1(u).$$

Thus we can choose an interval  $I_j$  for some  $j$  with  $1 \leq j \leq p$  such that

$$(3.11) \quad \int_{r \in I_j} |du|^2 dx \leq p^{-1}E_1(u) \leq c_{21}\bar{\varepsilon}^{1/16}E_1(u).$$

Let  $\theta$  be the number such that  $I_j = [\theta - \tau, \theta + 2\tau]$ , and let  $h(x)$  be as in Lemma 3.3. Thus  $u_{h(x)}(x) \in L^2_1(B_{1/2}, N)$  and satisfies  $u_h = u$  for  $r \geq \theta + \tau$ , and

$$\int_{r \in [\theta, \theta + \tau]} |du_h|^2 \leq c_{22} \int_{r \in I_j} |du|^2.$$

Thus by (3.11) we have

$$(3.12) \quad \int_{r \in [\theta, \theta + \tau]} |du_h|^2 \leq c_{23}\bar{\varepsilon}^{1/16}E_1(u).$$

By Lemma 2.3 we have

$$E_{\theta+\tau}(u) \leq c_{24}E_{\theta+\tau}(u_h) + c_{24}\Lambda\bar{\theta}^{n-1},$$

since  $\theta + \tau \leq 2\bar{\theta}$ . By (3.12) this implies

$$E_{\theta}(u) \leq c_{25}E_{\theta}(u_h) + c_{25}\bar{\varepsilon}^{1/16}E_1(u) + c_{25}\Lambda\bar{\theta}^{n-1}.$$

Combining this with (3.10) and using  $\theta \in [\bar{\theta}, 2\bar{\theta}]$  we have

$$\bar{\theta}^{2-n}E_{\theta}(u) \leq c_{26}(\bar{\theta}^{2-n}\bar{\varepsilon}^{1/16} + \bar{\theta}^2)(E_1(u) + \Lambda).$$

Since  $\bar{\theta} = \bar{\varepsilon}^{\gamma_n}$  this gives

$$\bar{\theta}^{2-n}E_{\bar{\theta}}(u) \leq c_{27}(\bar{\varepsilon}^{1/16-\gamma_n(n-2)} + \bar{\varepsilon}^{2\gamma_n})(E_1(u) + \Lambda).$$

We choose  $\gamma_n = \min\{[32(n-2)]^{-1}, 64^{-1}\}$  and hence

$$\bar{\theta}^{2-n}E_{\bar{\theta}}(u) \leq c_{27}\bar{\varepsilon}^{2\gamma_n}(E_1(u) + \Lambda).$$

This implies

$$\bar{\theta}^{2-n}(E_{\bar{\theta}}(u) + \theta\Lambda) \leq (c_{27} + 1)\bar{\varepsilon}^{2\gamma_n}(E_1(u) + \bar{\theta}\Lambda).$$

Choosing  $\bar{\varepsilon}$  so small that  $(c_{27} + 1)\bar{\varepsilon}^{2\gamma_n} \leq \frac{1}{2}$  finishes the proof of Proposition 3.4.

#### 4. Extension and compactness theorems

In this section we study convergence of  $\bar{E}$ -minimizing maps. This study involves the construction of lots of comparison maps. Our basic tool is the cone-type comparison already used in the proof of Proposition 2.4. For a fixed point  $u^* \in R^k$ , we introduce the notation

$$W_\sigma(u) = \int_{B_\sigma} |u - u^*|^2 dx.$$

This notation will be used throughout this section. Recall that  $E_\Omega(u)$  denotes the energy taken over a region  $\Omega \subseteq R^n$ ; likewise  $W_\Omega(u) = \int_\Omega |u - u^*|^2 dx$ . Let  $C_\sigma^n = B_\sigma^{n-1} \times [-\sigma, \sigma]$  be the cylinder of height and diameter  $2\sigma$ .

**Lemma 4.1.** *Let  $u \in L_1^2(\partial C_\sigma^n, N)$  be given such that  $u(x, -\sigma) = u_1(x)$ ,  $u(x, \sigma) = u_2(x)$  for  $x \in B_\sigma^{n-1}$  with  $u_1, u_2 \in L_1^2(B_\sigma^{n-1}, N)$ . Suppose also that  $u(x, t) = \circ u(x)$  for  $(x, t) \in S_\sigma^{n-2} \times [-\sigma, \sigma]$ . In particular we have  $u^1 = u^2 = \circ u$  on  $\partial B_\sigma^{n-1} = S_\sigma^{n-2}$  with  $\circ u \in L_1^2(S_\sigma^{n-2}, N)$ . Then there exists an extension  $\bar{u} \in L_1^2(C_\sigma^n, N)$ ,  $\bar{u} = u$  on  $\partial C_\sigma^n$ , satisfying the inequalities*

$$E(\bar{u}) \leq c\sigma(E_\sigma(u_1) + E_\sigma(u_2) + \sigma E(\circ u)),$$

$$W(\bar{u}) \leq c\sigma(W_\sigma(u_1) + W_\sigma(u_2) + \sigma W(\circ u)).$$

*Proof.* By scaling the domain we assume  $\sigma = 1$ . The easiest proof of this lemma is to observe that there exists a bi-Lipschitz homeomorphism  $f: \partial B_1^n \rightarrow \partial C_1^n$  which extends to a bi-Lipschitz homeomorphism  $\tilde{f}: B_1^n \rightarrow C_1^n$  where  $\tilde{f}(x) = |x|f(x/|x|)$ . Let  $\Pi: B_1^n \setminus \{0\} \rightarrow \partial B_1^n$  be the radial projection, i.e.,  $\Pi(x) = x/|x|$ . Define a projection map  $\hat{\Pi}: C_1^n \setminus \{(0, 0)\} \rightarrow \partial C_1^n$  by  $\hat{\Pi} = f \circ \Pi \circ \tilde{f}^{-1}$ . Then define  $\bar{u} = u \circ \hat{\Pi}$ . As in Proposition 2.4

$$E(\bar{u} \circ \tilde{f}) \leq (n-2)^{-1} E(u \circ f).$$

Due to Lipschitz equivalence

$$E(\bar{u}) \leq KE(\bar{u} \circ \tilde{f}), \quad E(u \circ f) \leq KE(u)$$

with constant  $K$  depending on the Lipschitz constants of  $\tilde{f}^{-1}$  and  $f$ . These inequalities and a similar argument for  $W$  yields the results of Lemma 4.1.

Since our main extension lemma, Lemma 4.3, will be proved by induction on dimension, we first give the result for  $n = 2$  where the proof is quite different and the result much simpler.



**Lemma 4.2.** *If  $u \in L^2_1(S^1_\sigma, N)$  and  $E(u)W(u) \leq \delta_1^2$  for a number  $\delta_1 = \delta_1(N_0)$ , then there exists  $\bar{u} \in L^2_1(B^2_\sigma, N)$  with  $\bar{u}|_{\partial B^2_\sigma} = u$  and*

$$E_\sigma(\bar{u}) \leq c_1(E(u)W(u))^{1/2}, \quad W_\sigma(\bar{u}) \leq c_1\sigma W(u).$$

*Proof.* As usual we take  $\sigma = 1$ . Let  $\delta^2 = E(u)W(u)$  so that  $\delta^2 \leq \delta_1^2$ . If  $S^1$  is parametrized by  $\theta \in [0, 2\pi)$ , then we have

$$|u(\theta) - u^*|^2 \leq 2 \int_0^{2\pi} |u(\theta) - u^*| |u'(\theta)| d\theta \leq 2\delta.$$

Thus if  $\delta_1$  is small, then  $\min |u - u^*|$  is small. Let  $\bar{u}$  be the (Morrey [18,5.4]) harmonic map minimizing  $E$  for boundary values given by  $u$ . Since  $\delta_1$  is small, the boundary values of  $\bar{u}$  lie in a convex ball (see Hildebrandt, Kaul and Widman [13]), so it follows that  $\|u - u^*\|_\infty \leq c_2\delta^{1/2}$ . The Euler-Lagrange equation for  $\bar{u}$  is

$$\Delta \bar{u} = A_{\bar{u}}(d\bar{u}, d\bar{u}).$$

This implies the inequality (in weak form)

$$\begin{aligned} \frac{1}{2} \Delta |\bar{u} - u^*|^2 - |d\bar{u}|^2 &= \langle \bar{u} - u^*, A_{\bar{u}}(d\bar{u}, d\bar{u}) \rangle \\ &\geq -\|\bar{u} - u^*\|_\infty \|A\|_\infty |d\bar{u}|^2. \end{aligned}$$

Thus if  $\delta_1$  is small, we have  $\Delta |\bar{u} - u^*|^2 \geq 0$ , so by the mean value inequality  $W(\bar{u}) \leq \frac{1}{2}W(u)$ .

To get the estimate on  $E(\bar{u})$ , we compare  $\bar{u}$  to  $\Pi \circ v$ , where  $v: B^2_1 \rightarrow R^k$  is the solution of the linear Dirichlet problem with boundary values  $u$  and  $\Pi$  is projection from a normal neighborhood of  $N_0$  onto  $N$ . By standard  $H_{1/2}$  norm estimates,  $E(v) \leq c\delta^{1/2}$ . Since  $\bar{u}$  is minimizing, we have

$$E(\bar{u}) \leq E(\Pi \circ v) \leq c_3E(v) \leq c_4\delta^{1/2}.$$

This proves Lemma 4.2.

We now prove a higher dimensional version of this.

**Lemma 4.3.** *For  $n \geq 2$  there exists  $\delta = \delta(n, N_0)$  and a constant  $q = q(n)$  such that if  $\epsilon \in (0, 1)$  is given and  $u \in L^2_1(\partial B^n_\sigma, N_0)$  satisfies  $\sigma^{4-2n}E(u)W(u) \leq \delta^2\epsilon^q$  (note that  $W$  depends also on a fixed vector  $u^* \in R^k$ ), then there exists  $\bar{u} \in L^2_1(B^n_\sigma, N)$ ,  $\bar{u}|_{\partial B^n_\sigma} = u$  such that*

$$\begin{aligned} E(\bar{u}) &\leq c_5(\epsilon\sigma E(u) + \epsilon^{-q}\sigma^{-1}W(u)), \\ W(\bar{u}) &\leq c_5\epsilon^{-q}\sigma W(u). \end{aligned}$$

*Proof.* First note that Lemma 4.2 implies Lemma 4.3 for  $n = 2$  with  $q(n) = 1$  which proves the first step of our induction. Also, by rescaling we take  $\sigma = 1$ . The following lemma is part of our construction. We leave its proof until later.

**Lemma 4.4.** *Let  $\sigma \in (0, \frac{1}{2})$ ,  $A_\sigma = S^{n-1} \times [-\sigma, \sigma]$ . Assume Lemma 4.3 is true for  $n - 1$ , and let  $v \in L^2_1(S^{n-1}, N)$  satisfy  $E(v)W(v) \leq \sigma^{2n-4}(\delta')^2$  where  $\delta' = \delta'(n - 1, N_0)$  depends on the constants arising from Lemma 4.3 for  $n - 1$ . Then there exists a combinatorial constant  $\alpha = \alpha(n) < 1$ , a constant  $K = K(n)$ , and a map  $\bar{v} \in L^2_1(A_\sigma, N)$ ,  $\bar{v}|_{S^{n-1} \times \{\sigma\}} = v$ ,  $\bar{v}|_{S^{n-1} \times \{-\sigma\}} = v'$  where  $v' \in L^2_1(S^{n-1}, N)$  such that*

$$\begin{aligned} E(\bar{v}) &\leq K\sigma E(v) + K\sigma^{-1}W(v), \\ W(\bar{v}) &\leq K\sigma W(v), \\ E(v') &\leq \sigma E(v) + K\sigma^{-2}W(v), \\ W(v') &\leq KW(v). \end{aligned}$$

We assume for now that Lemma 4.4 is true and proceed with the proof of Lemma 4.3. Let  $\varepsilon \in (0, 1)$  be given and choose an integer  $s$  with  $\alpha^s \approx \varepsilon$  ( $\alpha = \alpha(n)$  given in Lemma 4.4) and a cylinder of height  $2\sigma = \varepsilon$ . Consider the  $s$  disjoint annuli  $A_{i,\sigma}$  given by

$$A_{i,\sigma} = \{x \in B_1 : 1 - 2i\sigma \leq |x| \leq 1 - 2(i - 1)\sigma\}$$

for  $i = 1, \dots, s$ . Apply Lemma 4.4 on each of the  $A_{i,\sigma}$  by taking  $v = v_1 = u$  on the outer boundary of  $A_{1,\sigma}$  and at the  $i$ th step  $v = v_i = v'_{i-1}$  where  $v'_{i-1}$  is the value on the inner boundary obtained via Lemma 4.4 at the previous step. Note that as long as  $2s\sigma = \varepsilon s < \frac{1}{2}$ , each annulus  $A_{i,\sigma}$  is uniformly equivalent (Lipschitz) to  $S^{n-1} \times [-\sigma, \sigma]$ . In order to apply Lemma 4.4 we must have  $E(v_i)W(v_i) \leq \sigma^{2n-4}(\delta')^2$ , but from the  $(i - 1)$ -st application we get

$$\begin{aligned} E(v_i) &\leq \alpha E(v_{i-1}) + K\sigma^{-2}W(v_{i-1}), \\ W(v_i) &\leq KW(v_{i-1}). \end{aligned}$$

By iteration this gives (provided  $k \geq 2$ )

$$\begin{aligned} E(v_i) &\leq \alpha^{i-1}E(u) + 2\sigma^{-2}K^iW(u), \\ W(v_i) &\leq K^{i-1}W(u). \end{aligned}$$

Thus we may continue  $s$  times provided

$$(4.1) \quad \begin{aligned} K^s E(u)W(u) &\leq 2^{-1}\sigma^{2n-4}(\delta')^2, \\ 2\sigma^{-2}K^{2s-1}W^2(u) &\leq \sigma^{2n-4}(\delta')^2 2^{-1}. \end{aligned}$$

We check these inequalities by noting

$$K^s = \alpha^{s \ln K / \ln \alpha} \approx \varepsilon^{\ln K / \ln \alpha}, \quad \sigma = \varepsilon/2,$$

so that the first inequality is equivalent to

$$E(u)W(u) \leq c_6 \varepsilon^{2n-4 - \ln K / \ln \alpha} (\delta')^2,$$

that is, taking  $q = 2n - 4 - \ln K / \ln \alpha$ ,  $\delta(n, N_0) = c_6^{1/2} \delta'$  in the hypothesis of Lemma 4.3 for  $n$ . Now if the second inequality of (4.1) fails and the first holds, we have  $K^s E(u) W(u) \leq 2\sigma^{-2} K^{2s-1} W^2(u)$  which implies  $E(u) \leq 2\sigma^{-2} K^{s-1} W(u)$ . In this case we do not need to apply Lemma 4.4 at all since we can extend  $u$  homogeneously into  $B_1$  by  $\bar{u}(x) = u(x/|x|)$ , and we have

$$\begin{aligned} E(\bar{u}) &\leq \varepsilon E(u) + 2\sigma^{-2} K^{s-1} W(u) \\ &\leq \varepsilon E(u) + c_7 \varepsilon^{-2 + \ln K / \ln \alpha} W(u) \end{aligned}$$

as required.

We can thus assume that (4.1) is true and that we have applied Lemma 4.4  $s$  times. We then let  $\bar{u}|_{A_{i,\sigma}} = \bar{v}_i$ , the extension obtained at the  $i$ th state from Lemma 4.4. We extend  $\bar{u}$  into  $B_{1-s\varepsilon}$  by setting  $\bar{u}(x) = v'_s((1-s\varepsilon)x/|x|)$ . This gives a map  $\bar{u} \in L^2_1(B_1, N)$  with  $\bar{u}|_{\partial B_1} = u$ . By Lemma 4.4 we have

$$\begin{aligned} E(\bar{u}|_{A_{i,\sigma}}) &= E(\bar{v}_i) \leq K(\sigma E(v_i) + \sigma^{-1} W(v_i)) \\ &\leq K(\sigma \alpha^{i-1} E(u) + 3\sigma^{-1} K^i W(u)), \\ W(\bar{u}|_{A_{i,\sigma}}) &= W(\bar{u}_i) \leq K\sigma W(v_i) \leq \sigma K^i W(u). \end{aligned}$$

In the ball  $B_{1-s\varepsilon}^n$

$$\begin{aligned} E(\bar{u}|_{B_{1-s\varepsilon}^n}) &\leq E(v_s) \leq \alpha^{s-1} E(u) + 2\sigma^{-2} K^s W(u), \\ W(\bar{u}|_{B_{1-s\varepsilon}^n}) &\leq W(v_s) \leq K^{s-1} W(u). \end{aligned}$$

Adding these up we get

$$\begin{aligned} E(\bar{u}) &\leq \left( \alpha^{s-1} + K\sigma \sum_{i=1}^s \alpha^{i-1} \right) E(u) + c_8 \sigma^{-2} K^s W(u), \\ W(\bar{u}) &\leq c_8 K^s W(u). \end{aligned}$$

Since  $\alpha^s + K\sigma \sum_{i=1}^s \alpha^{i-1} \leq (1 + 2K(1 - \alpha)^{-1})\varepsilon = c_9 \varepsilon$  and  $K^s = \varepsilon^{\ln K / \ln \alpha}$ , we get the conclusion of Lemma 4.3.

*Proof of Lemma 4.4.* It remains to prove that Lemma 4.4 follows from Lemma 4.3 for  $n - 1$ . Let  $x_1, \dots, x_q$  be a maximal array of points on  $S^{n-1}$  satisfying  $|x_i - x_j| \geq \sigma$  for  $i \neq j$ . By maximality,  $S^{n-1} \subseteq \cup_{i=1}^q B_\sigma^{n-1}(x_i)$ . For each  $i \leq q$ , let  $\sigma_i \in [\sigma, 2\sigma]$  be chosen so that the restriction  $v_i = v|_{\partial B_{\sigma_i}^{n-1}(x_i)}$  lies in  $L^2_1(\partial B_{\sigma_i}^{n-1}(x_i), N)$  and

$$\begin{aligned} E(v_i) &\leq c_{10} \sigma^{-1} \int_{B_{2\sigma}^{n-1}(x_i)} |dv|^2 d\mu, \\ W(v_i) &\leq c_{10} \sigma^{-1} \int_{B_{2\sigma}^{n-1}(x_i)} |v - u^*|^2 d\mu. \end{aligned}$$

Let  $B_i = B_{\sigma_i}^{n-1}(x_i) \subseteq S^{n-1}$ . The existence of such  $\sigma_i$  follows from Fubini's theorem. We have

$$\sigma_i^{6-2n} E(v_i) W(v_i) \leq c_{11} \sigma^{4-2n} E(v) W(v).$$

Thus we can apply Lemma 4.3, for  $\epsilon$  to be chosen, on  $B_i$  provided

$$(4.2) \quad c_{11} \sigma^{4-2n} E(v) W(v) \leq (\bar{\delta})^2 \epsilon^{\bar{q}},$$

where  $\bar{\delta} = \delta(n-1, N_0)$ ,  $\bar{q} = q(n-1)$ . Let  $v'_i \in L_1^2(B_i, N)$  be given by Lemma 4.3 satisfying

$$\begin{aligned} E(v'_i) &\leq c_{12} (\epsilon \sigma E(v_i) + \epsilon^{-\bar{q}} \sigma^{-1} W(v_i)), \\ W(v'_i) &\leq c_{12} \epsilon^{-\bar{q}} \sigma W(v_i). \end{aligned}$$

From the choice of  $\sigma_i$  this implies

$$(4.3) \quad \begin{aligned} E(v'_i) &\leq c_{13} \left( \epsilon \int_{B_{2\sigma}^{n-1}(x_i)} |dv|^2 d\mu + \epsilon^{-\bar{q}} \sigma^{-2} \int_{B_{2\sigma}^{n-1}(x_i)} |v - u^*|^2 d\mu \right), \\ W(v'_i) &\leq c_{13} \epsilon^{-\bar{q}} \int_{B_{2\sigma}^{n-1}(x_i)} |v - u^*|^2 d\mu. \end{aligned}$$

We also observe that since any point  $x \in S^{n-1}$  is contained in a bounded number (depending only on  $n$ ) of balls  $B_{2\sigma}^{n-1}(x_i)$ , we have

$$(4.4) \quad \begin{aligned} \sum_{i=1}^q \int_{B_{2\sigma}^{n-1}(x_i)} |dv|^2 d\mu &\leq c_{14} E(v), \\ \sum_{i=1}^q \int_{B_{2\sigma}^{n-1}(x_i)} |v - u^*|^2 d\mu &\leq c_{14} W(v). \end{aligned}$$

From the choice of  $x_1, \dots, x_q$  we see that there is a fixed integer  $I_n$  and families  $\mathfrak{B}_1, \dots, \mathfrak{B}_{I_n}$  of balls such that

$$\bigcup_{j=1}^{I_n} \mathfrak{B}_j = \{B_i : i = 1, \dots, q\},$$

and each  $\mathfrak{B}_j$  is comprised of a family of disjoint balls. Since  $S^{n-1} \subseteq \bigcup_{i=1}^q B_i$ , we have  $\sum_{i=1}^q E(v|B_i) \geq E(v)$ . Thus for some  $\mathfrak{B}_j$ , say  $\mathfrak{B}_1$ , we have

$$(4.5) \quad \sum_{B_i \in \mathfrak{B}_1} E(v|B_i) \geq I_n^{-1} E(v).$$

Let  $\Theta = \bigcup_{B_i \in \mathfrak{B}_1} B_i$ , and define the extension  $\bar{v}$  on  $(S^{n-1} \sim \Theta \times [-\sigma, \sigma])$  by  $\bar{v}(x, t) = v(x)$ . On each cylinder  $B_i \times [-\sigma, \sigma]$ , apply Lemma 4.1 to get  $\bar{v} \in L^2_1(B_i \times [-\sigma, \sigma], N)$  satisfying  $\bar{v}(x, \sigma) = v(x)$ ,  $\bar{v}(x, -\sigma) = v'_i(x)$ , and  $\bar{v}(x, t) = v(x)$  for  $(x, t) \in (\partial B_i) \times [-\sigma, \sigma]$ . We have

$$\begin{aligned} E(\bar{v} | B_i \times [-\sigma, \sigma]) &\leq c_{15}\sigma(E(v | B_i) + E(v'_i) + \sigma E(v_i)), \\ W(\bar{v} | B_i \times [-\sigma, \sigma]) &\leq c_{15}\sigma(W(v | B_i) + W(v'_i) + \sigma W(v_i)), \\ E(\bar{v} | (S^{n-1} \sim \Theta) \times [-\sigma, \sigma]) &= 2\sigma E(v | S^{n-1} \sim \Theta), \\ W(\bar{v} | (S^{n-1} \sim \Theta) \times [-\sigma, \sigma]) &= 2\sigma W(v | S^{n-1} \sim \Theta). \end{aligned}$$

Therefore we have by (4.3) and (4.4)

$$(4.6) \quad \begin{aligned} E(\bar{v}) &\leq c_{16}\sigma E(v) + c_{16}\varepsilon^{-\bar{q}}\sigma^{-1}W(v), \\ W(\bar{v}) &\leq c_{16}\varepsilon^{-\bar{q}}W(v). \end{aligned}$$

Let  $v' = \bar{v} | S^{n-1} \times \{-\sigma\}$ , so that by (4.3), (4.4), and (4.5)

$$\begin{aligned} E(v') &\leq E(v | S^{n-1} \sim \Theta) + \sum_{i=1}^q E(v'_i) \\ &\leq (1 - I_n^{-1})E(v) + c_{17}\varepsilon E(v) + c_{17}\varepsilon^{-\bar{q}}\sigma^{-2}W(v), \\ W(v') &\leq W(v) + \sum_{i=1}^q W(v'_i) \leq c_{17}\varepsilon^{-\bar{q}}W(v). \end{aligned}$$

We now fix  $\varepsilon$  so small that  $\alpha(n) = (1 - I_n^{-1} + c_{17}\varepsilon) < 1$ . This fixes all constants  $E(v)W(V) \leq \sigma^{2n-4}(\delta')^2$  with  $(\delta')^2 = c_{11}^{-1}(\bar{\delta})^2\varepsilon^{\bar{q}}$ . This completes the proof of Lemma 4.4.

The main application of Lemma 4.4 is to give a significant improvement of Theorem 3.1. We now prove this.

**Proposition 4.5.** *Given  $B > 0$  there exists a constant  $\varepsilon_0 = \varepsilon_0(n, N_0, B)$  such that if  $u \in \mathcal{K}_\Lambda$ ,  $\Lambda \leq \varepsilon_0$ ,  $E_1(u) \leq B$ , and  $W_1(u) \leq \varepsilon_0$ , then  $u$  is Hölder continuous on  $B_{1/2}$  and  $|u(x) - u(y)| \leq c|x - y|^\alpha$  for  $x, y \in B_{1/2}$  where  $\alpha = \alpha(n) \geq 0$  and  $c = c(n, N_0)$ .*

*Proof.* By Fubini's theorem, there exists  $\sigma \in [\frac{3}{4}, 1]$  such that

$$W(u | \partial B_\sigma) = \int_{\partial B_\sigma} |u - u^*|^2 d\xi \leq 8W_1(u) \leq 8\varepsilon_0,$$

$$E(u | \partial B_\sigma) \leq 8E_1(u) \leq 8B.$$

Applying Lemma 4.3 on  $B_\sigma$ , there exists  $\bar{u} \in L^2_1(B_\sigma, N_0)$  with  $\bar{u} | \partial B_\sigma = u$  such that if  $W(u | \partial B_\sigma) \leq 8^{-1}\sigma^{2n-4}\delta^2\varepsilon^q B^{-1}$

$$E_\sigma(\bar{u}) \leq 8c_5(\varepsilon\sigma B + \varepsilon^{-q}\sigma^{-1}\varepsilon_0).$$

Since  $u \in \mathcal{H}_\lambda$ , we can apply Lemma 2.3 to get

$$E_\sigma(u) \leq (1 + c\Lambda\sigma)E_\sigma(\bar{u}) + c\Lambda\sigma^{n-1} \leq c_{18}(\varepsilon B + \varepsilon^{-q}\varepsilon_0).$$

Fix  $\varepsilon$  so small that  $c_{18}\varepsilon B \leq 2^{-1}\bar{\varepsilon}\sigma^{n-2}$  where  $\bar{\varepsilon}$  is given in Theorem 3.1. If  $\varepsilon_0$  is so small that  $c_{18}\varepsilon^{-q}\varepsilon_0 \leq 2^{-1}\bar{\varepsilon}\sigma^{n-2}$  and  $8\varepsilon_0 B \leq 8^{-1}\sigma^{2n-4}\delta^2\varepsilon^q$ , then we have  $\sigma^{2-n}E_\sigma(u) \leq \bar{\varepsilon}$ ,  $\sigma\Lambda \leq \bar{\varepsilon}$  so we can apply Theorem 3.1 to assert that  $u$  is Hölder continuous on  $B_{\sigma/2}$ . Proposition 3.5 now follows.

We now look at weak limits of minimizing maps. We cannot show that these are again minimizing. Our first result is a compactness theorem which says that weak convergence is actually strong. The key ingredient in its proof is Proposition 4.5.

**Proposition 4.6.** *Let  $\{u_i\} \subseteq \mathcal{H}_\Lambda$  be a weakly convergent (in  $L_1^2$ ) sequence with limit  $u_0$  such that  $E_1(u_i) \leq c_{19}$ . Then  $u_0$  is locally Hölder continuous outside a closed set  $\mathcal{S}_0$  with  $\mathcal{H}^{n-2}(\mathcal{S}_0) = 0$ . Moreover,  $u_i$  converges to  $u_0$  in  $L_1^2$ -norm on  $B_{1/2}$  and uniformly on compact subsets of  $\bar{B}_{1/2} \sim \mathcal{S}_0$ .*

*Proof.* Since the  $u_i$  have uniformly bounded energy, we may assume they converge in  $L^2$ -norm to  $u_0$ . By lower semi-continuity  $E_1(u_0) \leq c_{19}$ . Let  $\mathcal{S}_0$  be the singular set of  $u_0$ . We prove that  $\mathcal{S}_0$  is small by noting that if  $W_\sigma^x(u_0) < \varepsilon_0\sigma^n$  for some  $u^* \in R^k$  and  $\varepsilon_0 > 0$  ( $\varepsilon_0$  given by Proposition 4.5), by  $L^2$ -convergence we have  $W_\sigma^x(u_i) < \varepsilon_0\sigma^n$  for large  $i$ . By Proposition 2.4, there exists  $B > 0$  such that  $\sigma^{2-n}E_\sigma^x(u_i) \leq B$  for all  $i$ . Moreover, if Proposition 4.5 is applied on a ball of radius  $\sigma$ , then the assumption on  $\Lambda$  is  $\Lambda \leq \varepsilon_0\sigma^{-1}$  which is automatically satisfied for  $\sigma$  small. Thus we can apply Proposition 4.5 to  $u_i$  giving us a uniform Hölder estimate on  $u_i$  in  $B_{\sigma/2}(x)$ , and hence  $u_i$  converges uniformly on  $B_{\sigma/2}(x)$  to  $u_0$  and  $u_0$  is Hölder continuous there. In particular, by the Poincaré inequality, if  $\sigma^{2-n}E_\sigma^x(u_0)$  is small, then  $u_0$  is Hölder continuous on  $B_{\sigma/2}(x)$ . Thus by the argument given in Corollary 2.7 we have  $\mathcal{H}^{n-2}(\mathcal{S}_0) = 0$ . We have also shown that  $u_i$  converges uniformly to  $u_0$  on compact subsets of  $\bar{B}_{1/2} \sim \mathcal{S}_0$ .

To prove the  $L_1^2$ -convergence of  $u_i$  to  $u_0$ , we now observe that we can cover  $\mathcal{S}_0 \cap B_{1/2}$  by a family of balls  $\{B_{r_i}(x_i)\}$  such that  $\sum_i r_i^{n-2} < \varepsilon$  for any  $\varepsilon > 0$ . If  $\Theta = \cup_i B_{r_i}(x_i)$ , we can estimate by Proposition 2.4

$$(4.7) \quad E_\Theta(u_j) \leq \sum_i E_{r_i}^{x_i}(u_j) \leq c_{20} \sum_i r_i^{n-2} < c_{20}\varepsilon$$

for any  $j$ . On the other hand, we have shown uniform convergence of  $u_j$  to  $u_0$  on  $B_{1/2} \sim \Theta$  so subtracting the Euler equations (2.1) for  $u_j$  and  $u_k$ , multiplying by  $u_j - u_k$  and putting in a cutoff function we easily get

$$\int_{B_{1/2} \sim \Theta} |d(u_j - u_k)|^2 dx \leq c(\Theta, \Lambda) \sup_{\bar{B}_{1/2} \sim \Theta} |u_j - u_k|.$$

Therefore we have from 4.7

$$\int_{B_{1/2}} |d(u_j - u_k)|^2 dx \leq c_{21}\epsilon + c(\Theta, \Lambda) \sup_{\bar{B}_{1/2-\epsilon}} |u_j - u_k|.$$

Thus  $\{u_j\}$  is a Cauchy sequence in  $L^2_1(B_{1/2}, N_0)$  which therefore converges in  $L^2_1$ -norm to  $u_0$ . This completes the proof of Proposition 4.6.

Let  $\mathcal{H}_{\Lambda, B}$  denote the set of maps  $u \in \mathcal{H}_\Lambda$  with  $E_1(u) \leq B$ . Let  $\bar{\mathcal{H}}_{\Lambda, B}$  denote the closure of  $\mathcal{H}_{\Lambda, B}$  taken in  $L^2_1(B_1, N_0)$ . We now prove a strong version of Lemma 2.5.

**Proposition 4.7.** *Given  $u \in \bar{\mathcal{H}}_{\Lambda, B}$  and  $x_0 \in B_{1/2}$ , there is a sequence  $\lambda(i) \rightarrow 0$ ,  $\lambda(i) \in (0, \frac{1}{2}]$ , such that the maps  $u_{x_0, \lambda(i)} \in L^2_1(B_1, N_0)$  defined by  $u_{x_0, \lambda(i)}(x) = u(\lambda(i)(x - x_0))$  converge in  $L^2_1$ -norm on  $B_1^n$  to a harmonic map  $u_0 \in \mathcal{H}_{0, B}$  satisfying  $\partial u_0 / \partial r = 0$  a.e. on  $B_1$ . Moreover, the convergence is uniform on compact subsets of  $\bar{B}_1 \sim \mathcal{S}_0$ .*

*Proof.* Since  $u$  is a strong limit of minimizing maps, we get inequality (2.4) satisfied for  $u$  and hence by Lemma 2.5 there is a sequence  $\lambda(i) \rightarrow 0$  such that  $u_{x_0, \lambda(i)}$  converges weakly to a harmonic map  $u_0$  satisfying  $\partial u_0 / \partial r = 0$  a.e. Since  $u_{x_0, \lambda(i)} \in \bar{\mathcal{H}}_{\lambda(i)\Lambda, B}$ , it follows that there is  $\tilde{u}_i \in \mathcal{H}_{\lambda(i)\Lambda, B}$  with  $\|\tilde{u}_i - u_{x_0, \lambda(i)}\|_{1,2} < i^{-1}$ . From this we see that  $\tilde{u}_i$  converges weakly to  $u_0$  and hence strongly by Proposition 4.6. Therefore,  $\{u_{x_0, \lambda(i)}\}$  converges strongly to  $u_0$  and we have proven Proposition 4.7.

### 5. Dimension reduction of $\mathcal{S}$

We prove Theorems II and IV simultaneously in this section by adapting to our setting a dimension reducing argument of H. Federer [7]. We first need some preliminary results.

**Lemma 5.1.** *Suppose  $l \geq 3$  and  $u \in L^2_{1,loc}(R^l, N)$  satisfies  $g, u / \partial x^l = 0$  a.e. Then there exists  $u_0 \in L^2_{1,loc}(R^{l-1}, N)$  such that  $u(x', x^l) = u_0(x')$  a.e.  $x' \in R^{l-1}$ . If  $u$  is  $E$ -minimizing on each compact subset of  $R^l$ , then  $u_0$  is  $E$ -minimizing on each compact subset of  $R^{l-1}$ .*

*Proof.* Suppose on the contrary that  $v: B_\sigma^{l-1} \rightarrow N_0$  satisfies  $v = u_0$  on  $\partial B_\sigma^{l-1}$  and  $E_\sigma(v) \leq E_\sigma(u_0) - \eta$  for some  $\eta > 0$ . Let  $\lambda \gg 0$  be a large number and consider a map  $\bar{v}: B_\sigma^{l-1} \times [-\lambda - 2\sigma, \lambda + 2\sigma] \rightarrow N_0$  satisfying  $\bar{v}(x', x^l) = v(x')$  for  $|x^l| \leq \lambda$  and constructed by Lemma 4.1 on  $B_\sigma^{l-1} \times [-\lambda - 2\sigma, -\lambda]$  and  $B_\sigma^{l-1} \times [\lambda, \lambda + 2\sigma]$  so that  $\bar{v} = u$  on  $\partial(B_\sigma^{l-1} \times [-\lambda - 2\sigma, \lambda + 2\sigma])$  and  $E(\bar{v}) \leq 2\lambda E_\sigma(v) + c$  where  $c$  depends on  $\sigma, u, v$ . By the minimizing property of  $u$  in  $R^l \sim \mathcal{S}$  we have

$$(2\lambda + 4\sigma)E_\sigma(u) \leq 2\lambda E_\sigma(v) + c.$$

Choosing  $\lambda$  large we contradict the inequality  $E_\sigma(v) \leq E_\sigma(u) - \eta$ . This proves Lemma 5.1.

The next result guarantees that, in a very simple situation, limits of minimizing maps are minimizing.

**Lemma 5.2.** *Let  $u_0 \in L^2_{1,\text{loc}}(R^l, N_0)$  with  $l \geq 3$  be a harmonic map with an isolated singularity at 0 such that  $u_0$  satisfies  $\partial u_0 / \partial r = 0$  a.e. Suppose  $u \in L^2_{1,\text{loc}}(R^n, N_0)$ ,  $n \geq l$ , is given by  $u(x', x'') = u_0(x')$ ,  $x' \in R^l$ ,  $x'' \in R^{n-l}$ . Suppose there is a sequence  $u_i \in \mathcal{H}_{\Lambda_i, B}$  such that  $u_i$  converges to  $u$  in  $L^2_1(B_1, N_0)$  and  $\Lambda_i \rightarrow 0$ . Then both  $u, u_0$  are  $E$ -minimizing on compact subsets of  $R^n, R^l$ . In particular,  $u_0$  is a minimizing tangent map (MTM).*

*Proof.* We first show that  $u$  is minimizing. Note that because of the homogeneity of  $u$  it suffices to prove the  $u$  is  $E$ -minimizing on  $B'_1 \times B_1^{n-l}$ . We first prove a preliminary inequality. Suppose  $v_0 \in L^2_1(B'_1, N_0)$  such that  $v_0 = u_0$  on  $\partial B'_1$ . We will modify  $v_0$  to make it agree with  $u$  near the origin. For any  $\delta > 0$ , define a map  $v_\delta \in L^2_1(B'_1, N)$  by

$$v_\delta(x) = \begin{cases} v(x) & \text{for } |x| \geq \delta, \\ v\left(\delta \frac{x}{|x|}\right) & \text{for } |x| \leq \delta. \end{cases}$$

We then have

$$(5.1) \quad E_\delta(v_\delta) \leq \delta \int_{\partial B'_\delta} |dv_0|^2 dx.$$

For any  $\varepsilon \in (0, 2^{-1}\delta)$ , define  $v_{\delta,\varepsilon}: B'_1 \rightarrow N$  by

$$v_{\delta,\varepsilon}(r, \xi) = \begin{cases} v_\delta(r, \xi) & \text{for } r \geq 2\varepsilon, \\ u_0(r, \xi) & \text{for } r \leq \varepsilon, \\ v_\delta(\rho(r), \xi) & \text{for } \varepsilon < r < 2\varepsilon \end{cases}$$

where  $r, \xi$  are polar coordinates in  $R^l$  and  $\rho(r)$  is the linear function

$$\rho(r) = (2 - 2\varepsilon) - \varepsilon^{-1}(1 - 2\varepsilon)r.$$

Since  $u_0(r, \xi)$  is independent of  $r$  and  $v_0 = u_0$  on  $\partial B'_1$ , we see that  $v_{\delta,\varepsilon} \in L^2_1(B'_1, N_0)$  and satisfies  $v_{\delta,\varepsilon} = u_0$  on  $\partial B'_1$ ,  $v_{\delta,\varepsilon} = u_0$  in  $B_\varepsilon$ . We now estimate

$$E_1(v_{\delta,\varepsilon}) \leq E_1(v_\delta) + E_\varepsilon(u_0) + \int_{B'_{2\varepsilon} \sim B_\varepsilon} |dv_{\delta,\varepsilon}|^2 dx.$$

By (5.1) and the homogeneity of  $u$  this implies

$$(5.2) \quad \begin{aligned} E_1(v_{\delta,\varepsilon}) &\leq E_1(v_0) + \delta \int_{\partial B'_\delta} |dv_0|^2 + \varepsilon^{l-2} E_1(u_0) \\ &\quad + \int_{B'_{2\varepsilon} \sim B'_\varepsilon} |dv_{\delta,\varepsilon}|^2 dx. \end{aligned}$$



We estimate the last term in two parts

$$\int_{B'_{2\epsilon} \sim B'_\epsilon} \left| \frac{\partial v_{\delta,\epsilon}}{\partial r} \right|^2 dx \leq \epsilon^{l-2} \int_\epsilon^{2\epsilon} \int_{S^{l-1}} \left| \frac{\partial v_\delta}{\partial \rho} \right|^2 (\rho, \xi) r^{-1} d\xi dr.$$

By change of variable this gives

$$\int_{B'_{2\epsilon} \sim B'_\epsilon} \left| \frac{\partial v_{\delta,\epsilon}}{\partial r} \right|^2 dx \leq \epsilon^{l-2} \int_{2\epsilon}^1 \int_{S^{l-1}} \left| \frac{\partial v_\delta}{\partial \rho} \right|^2 (\rho, \xi) d\xi d\rho.$$

From the definition of  $v_\delta$  this implies (since  $\partial v_\delta / \partial r = 0$  on  $|x| \leq \delta$ )

$$(5.3) \quad \int_{B'_{2\epsilon} \sim B'_\epsilon} \left| \frac{\partial v_{\delta,\epsilon}}{\partial r} \right|^2 dx \leq \epsilon^{l-2} \delta^{1-l} \int_{B_1 \sim B_\delta} \left| \frac{\partial v_0}{\partial r} \right|^2 dx.$$

We also can estimate

$$\int_{B_{2\epsilon} \sim B_\epsilon} r^{-2} |d_\xi v_{\delta,\epsilon}|^2 dx \leq c_1 \epsilon^{l-3} \int_\epsilon^{2\epsilon} \int_{S^{l-1}} |d_\xi v_\delta|^2 (\rho, \xi) d\xi dr.$$

Changing variables we get

$$(5.4) \quad \int_{B_{2\epsilon} \sim B_\epsilon} r^{-2} |d_\xi v_{\delta,\epsilon}|^2 dx \leq c_2 \epsilon \int_{B_1 \sim B_{2\epsilon}} r^{-2} |d_\xi v_\delta|^2 dx.$$

Combining (5.2), (5.3) and (5.4) we get

$$(5.5) \quad E_1(v_{\delta,\epsilon}) \leq (1 + c_2 \epsilon + \epsilon^{l-2} \delta^{1-l}) E_1(v_0) + c_3 \delta \int_{\partial B'_\delta} |dv_0|^2 + \epsilon^{l-2} E_1(u_0).$$

Now, given a map  $v: B'_1 \times B_1^{n-1} \rightarrow N_0$  which agrees with  $u$  on the boundary, we apply (5.5) on each  $R^l$  slice (with  $x''$  fixed) and integrate over  $B_1^{n-l}$ . We use the notation  $|dW|^2 = |d'W|^2 + |d''W|^2$  for a map  $W(x', x'')$ . We have from (5.5)

$$(5.6) \quad \begin{aligned} \int_{B'_1 \times B_1^{n-l}} |d'v_{\delta,\epsilon}|^2 dx' dx'' &\leq (1 + c_2 \epsilon + \epsilon^{l-2} \delta^{1-l}) \int_{B'_1 \times B_1^{n-l}} |d'v|^2 \\ &\leq c_3 \delta \int_{\partial B'_\delta \times B_1^{n-l}} |d'v|^2 + \epsilon^{l-2} \int_{B'_1 \times B_1^{n-l}} |du|^2. \end{aligned}$$

From the definition of  $v_{\delta,\epsilon}$  we can estimate

$$\int_{B'_{2\epsilon}} |d''v_{\delta,\epsilon}|^2 dx' \leq c_4 \epsilon^{l-1} \int_\epsilon^{2\epsilon} \int_{S^{l-1}} |d''v_\delta|^2 (\rho, \xi, x'') d\xi dr.$$

Changing variables we get

$$\int_{B'_{2\epsilon}} |d''v_{\delta,\epsilon}|^2 dx' \leq c_4 \epsilon \left( \int_{B'_1} |d''v|^2 dx' + \int_{B'_\delta} |d''v_\delta|^2 dx' \right).$$

We also see that  $\int_{B_\delta} |d''v_\delta|^2 dx' \leq \delta \int_{\partial B_\delta} |d''v|^2$ . Combining with (5.6) we then have

$$\int_{B'_1 \times B_1^{\eta-1}} |dv_{\delta,\epsilon}|^2 dx' dx'' \leq (1 + c_5 \epsilon + \epsilon'^{-2} \delta^{1-l}) \int_{B'_1 \times B_1^{\eta-1}} |dv|^2 dx' dx'' + c_5 \delta \int_{\partial B'_\delta \times B_1^{\eta-1}} |dv|^2 + \epsilon'^{-2} \int_{B'_1 \times B_1^{\eta-1}} |du|^2 dx' dx''.$$

We can choose  $\delta$  so that

$$\delta \int_{\partial B'_\delta \times B_1^{\eta-1}} |dv|^2 \leq 2 \int_{B'_\delta \times B_1^{\eta-1}} |dv|^2 dx' dx''.$$

Thus we can choose  $\epsilon, \delta$  small so that

$$(5.7) \quad \int_{B'_1 \times B_1^{\eta-1}} |dv_{\delta,\epsilon}|^2 dx \leq \int_{B'_1 \times B_1^{\eta-1}} |dv|^2 dx + \eta$$

for any given  $\eta > 0$ . The point is we have from the above construction  $v_{\delta,\epsilon} = u$  on  $\partial(B'_1 \times B_1^{\eta-1})$  as well as  $v_{\delta,\epsilon} = u$  on  $B'_\delta \times B_1^{\eta-1}$ , a neighborhood of  $\mathfrak{S} = \{0\} \times B_1^{\eta-1}$ . Since by Proposition 4.6,  $u_i$  converges uniformly to  $u$  away from  $\mathfrak{S}$ , it is obvious that  $u$  minimizes on each compact subset of  $R^n \sim \mathfrak{S}$ . Therefore we have

$$\int_{B'_1 \times B_1^{\eta-1}} |du|^2 \leq \int_{B'_1 \times B_1^{\eta-1}} |dv_{\delta,\epsilon}|^2.$$

By (5.7), since  $\eta$  is arbitrarily small we have

$$\int_{B'_1 \times B_1^{\eta-1}} |du|^2 \leq \int_{B'_1 \times B_1^{\eta-1}} |dv|^2,$$

and  $u$  is minimizing. Applying Lemma 5.1 successively, we get that  $u_0$  is minimizing, and this finishes Lemma 5.2.

We now return to Hausdorff measure, and define for  $E \subseteq R^n, s \geq 0$ ,

$$\varphi^s(E) = \inf \left\{ \sum r_i^s : E \subseteq \bigcup_i B_{r_i}(x_i) \right\}.$$

Following [7], we observe that for any  $E$

$$(5.8) \quad \varphi^s(E) = 0 \Leftrightarrow \mathfrak{H}^s(E) = 0.$$

We also need the following density result (see [6,2.10.19(2)])

$$(5.9) \quad \overline{\lim}_{\lambda \rightarrow 0} \lambda^{-s} \varphi^s(E \cap B_\lambda^n(x)) \geq c_6 > 0$$

for  $\varphi^s$  a.e.  $x \in E$ . We need the following result on the behaviour of  $\varphi^s$  under weak convergence.

**Lemma 5.3.** *Suppose  $u_i$  is a sequence in  $\mathcal{H}_\Lambda$  which converges weakly to  $u$  in  $L^2_1(B^n_1, N)$ . If  $\mathcal{S}_i, \mathcal{S}$  denote the singular sets of  $u_i, u$  respectively, then we have*

$$\varphi^s(\mathcal{S} \cap B^n_{1/2}) \geq \overline{\lim}_{i \rightarrow 0} \varphi^s(\mathcal{S}_i \cap B^n_{1/2})$$

for any  $s \geq 0$ .

*Proof.* For any  $\epsilon > 0$ , let  $\{B_{r_i}(x_i)\}$  be a covering of  $\mathcal{S} \cap B^n_{1/2}$  by balls satisfying

$$\sum_i r_i^s \leq \varphi^s(\mathcal{S} \cap B^n_{1/2}) + \epsilon.$$

Now the set  $K = \overline{B^n_{1/2}} \sim \cup_i B^n_{r_i}(x_i)$  is compact subset of  $\overline{B^n_{1/2}} \sim \mathcal{S}$ , so by Proposition 4.6 it follows that for  $j$  sufficiently large, the map  $u_j$  is smooth on  $K$ . Thus we have

$$\mathcal{S}_j \cap B^n_{1/2} \subseteq \bigcup_i B^n_{r_i}(x_i)$$

for  $j$  large. In particular we have

$$\varphi^s(\mathcal{S}_j \cap B^n_{1/2}) \leq \varphi^s(\mathcal{S} \cap B^n_{1/2}) + \epsilon$$

for any  $\epsilon > 0, j$  large. This gives the conclusion of Lemma 5.3.

*Proof of Theorems II and IV.* Suppose  $u \in L^2_1(M, N)$  is  $\tilde{E}$ -minimizing with singular set  $\mathcal{S} \subset \text{int } M$ . Let  $0 \leq s < n - 2$  be such that  $\varphi^s(\mathcal{S}) > 0$ . Then by (5.9) we can choose  $p_0 \in \mathcal{S}$  such that

$$(5.10) \quad \lim_{\lambda_i \rightarrow 0} \lambda_i^{-s} \varphi^s(\mathcal{S} \cap B^n_{\lambda_i/2}) > 0$$

for a sequence  $\lambda_i \rightarrow 0$ , where  $B_\lambda$  is taken in normal coordinates  $x$  centered at  $p_0$ . We look at the scaled maps  $u_\lambda(x) = u(\lambda x)$ . By Proposition 4.7 we can choose a subsequence of  $\lambda_i$ , call it  $\lambda_i$ , so that  $u_{\lambda_i}$  converges weakly in  $L^2_1(B^n_1, N)$  to a harmonic map  $u_0$ , strongly in  $L^2_1(B^n_{1/2}, N)$ , where  $u_0$  satisfies  $\partial u_0 / \partial r = 0$  a.e. If  $\mathcal{S}_\lambda$  denotes the singular set of  $u_\lambda$  in  $B^n_1$ , we clearly have  $\mathcal{S}_\lambda \cap B^n_{1/2} = \{x/\lambda : x \in \mathcal{S} \cap B^n_{\lambda/2}\}$  and hence  $\varphi^s(\mathcal{S}_\lambda \cap B^n_{1/2}) = \lambda^{-s} \varphi^s(\mathcal{S} \cap B^n_{\lambda/2})$ . Thus (5.10) implies

$$\lim_{\lambda_i \rightarrow 0} \varphi^s(\mathcal{S}_{\lambda_i} \cap B^n_{1/2}) > 0.$$

Thus by Lemma 5.3 we have

$$\varphi^s(\mathcal{S}_0 \cap B^n_{1/2}) > 0.$$

Since  $\partial u_0 / \partial r = 0$  a.e., we have  $\lambda \mathcal{S}_0 \subseteq \mathcal{S}_0$  for any  $\lambda \geq 0$ , and there are two possibilities: either we have  $s \leq 0$ , or we can choose a point  $x_1 \in \mathcal{S}_0 \cap \partial B^n_{1/2}$  by (5.9) such that

$$\overline{\lim}_{\lambda \rightarrow 0} \lambda^{-s} \varphi^s(\mathcal{S}_0 \cap B^n_\lambda(x_1)) > 0.$$

We choose Euclidean coordinates centered at  $x_1$  so that  $x^1$  is radial at  $x_1$ . Repeating the above argument at  $x_1$  we get a radially independent harmonic map  $u_1 \in L^2_{1,\text{loc}}(R^n, N_0)$  with  $\varphi^s(\mathcal{S}_1 \cap B_1^n) > 0$ . Since  $u_0$  satisfied  $\partial u_0 / \partial r = 0$ , it follows that  $\partial u_1 / \partial x^1 = 0$  a.e. If  $s - 1 \leq 0$ , we stop. Otherwise, there is a point  $x_2 \in \mathcal{S}_1 \cap \partial B_1^{n-1}$ ,  $R^{n-1} = \{(0, x^2, \dots, x^n)\}$  and we repeat the argument at  $x_2$ . If we repeat this procedure  $m$  times, we get harmonic maps  $u_j \in L^2_{1,\text{loc}}(R^n, N_0)$  for  $j = 1, \dots, m$  such that  $u_j|_{B_1^n} \in \overline{\mathcal{H}}_{\Lambda, B}$  for suitable  $B$  (see Proposition 4.7) and  $\partial u_j / \partial r = \partial u_j / \partial x^\alpha = 0$  a.e.  $\alpha = 1, \dots, j$ . Also we would have  $\varphi^s(\mathcal{S}_j \cap B_1^n) > 0$  for  $j = 1, \dots, m$ . We can repeat the argument until we have  $s - m \leq 0$ . In order to have constructed  $u_m$ , we must have had  $s - m + 1 > 0$ . Since  $s < n - 2$ , and  $m$  is an integer we then have  $m \leq n - 2$ . If  $m = n - 2$ , then we would have  $\mathcal{S}_m \supseteq R^{n-2} = \{(x^1, \dots, x^{n-2}, 0, 0)\}$  contradicting the fact that  $\mathcal{H}^{n-2}(\mathcal{S}_m) = 0$ . Therefore we have  $m \leq n - 3$ , and hence  $\varphi^t(\mathcal{S}_m \cap B_1^n) = 0$  for  $t > n - 3$ . Since  $\varphi^s(\mathcal{S}_m \cap B_1^n) > 0$ , we have  $s \leq n - 3$ , and since  $s$  can be any number smaller than  $\dim \mathcal{S}$  we have shown  $\dim \mathcal{S} \leq n - 3$ .

If we make the additional assumption that for  $j = 1, \dots, l$  we have no nontrivial MTM from  $R^j \rightarrow N_0$ , then we can say more. If  $m = n - 3$ , then we have  $u_m \in L^2_{1,\text{loc}}(R^n, N_0)$  such that  $u_m|_{B_1^n} \in \overline{\mathcal{H}}_{\Lambda, B}$ , and  $u_m(x', x'') = \tilde{u}_m(x'')$  for  $x' \in R^{n-3}$ ,  $x'' \in R^3$  where  $\tilde{u}_m$  has an isolated singularity at  $x'' = 0$ . Therefore, by Lemma 5.2,  $\tilde{u}_m \in L^2_{1,\text{loc}}(R^3, N_0)$  is an MTM and hence trivial by assumption. Thus we had  $m \leq n - 4$ . We can repeat the same reasoning for  $m = n - 4, \dots, n - l$  and we conclude that  $m \leq n - l - 1$  which then implies  $s \leq n - l - 1$  for any  $s < \dim \mathcal{S}$  and hence  $\dim \mathcal{S} \leq n - l - 1$ .

Finally, suppose we had  $n = l + 1$  and  $\mathcal{S} \neq \emptyset$ . If  $p_0 \in \mathcal{S}$  and  $u_0 \in L^2_{1,\text{loc}}(R^n, N_0)$  is a blown-up harmonic map at  $p_0$ , then the above argument shows that  $u_0$  has singular set  $\mathcal{S}_0 = \{0\}$  and  $u_0$  is an MTM. If there were a sequence  $p_i \in \mathcal{S}$  with  $p_i \rightarrow p_0$ , then we could choose  $\lambda(i) = 4 \text{dist}(p_i, p_0)$  and consider the scaled maps  $u_{\lambda(i)} \in L^2_1(B_1^n, N_0)$ . By the choice of  $\lambda(i)$ , we have  $\mathcal{S}_{\lambda(i)} \cap \partial B_{1/4}^n \neq \emptyset$  for each  $i$ . Since the limit  $u_0$  has an isolated singularity at 0, this contradicts Proposition 4.7. Therefore  $\mathcal{S}$  is discrete. The same argument shows that, in general, for  $n = 3$  either  $\mathcal{S} = \emptyset$  or  $\mathcal{S}$  is discrete. This completes the proofs of both Theorems II and IV.

### References

- [1] F. J. Almgren, Jr., *Existence and regularity almost everywhere of solutions to elliptic variational problems among surfaces of varying topological type and singularity structure*, Ann. of Math. **87** (1968) 321–391.
- [2] H. J. Borchers & W. J. Garber, *Analyticity of solutions of the  $O(N)$  non-linear  $\sigma$ -model*, Comm. Math. Phys. **71** (1980) 299–309.

- [3] E. De Giorgi, *Frontiere orientate di misura minima*, Seminario Mat. Scuola Norm. Sup. Pisa, 1966.
- [4] ———, *Un esempio di estremali discontinue per un problema variazionale di tipo ellittico*, Boll. Un. Mat. Ital. (5) 4 (1968).
- [5] J. Eells & J. H. Sampson, *Harmonic mappings of Riemannian manifolds*, Amer. J. Math. 86 (1964) 109–160.
- [6] H. Federer, *Geometric measure theory*, Springer, New York, 1969.
- [7] ———, *The singular sets of area minimizing rectifiable currents with codimension one and of area minimizing flat chains modulo two with arbitrary codimension*, Bull. Amer. Math. Soc. 79 (1970) 761–771.
- [8] M. Giaquinta, *Multiple integrals in the calculus of variations and non-linear elliptic systems*, Preprint 443, Bonn, 1981, 25–212.
- [9] M. Giaquinta & E. Giusti, *On the regularity of the minima of variational integrals*, Acta Math., to appear.
- [10] ———, *On the singular set of the minima of certain quadratic functionals*, Preprint 453, Bonn.
- [11] E. Giusti & M. Miranda, *Sulla regolarità delle soluzioni deboli di una classe di sistemi ellittici quasilineari*, Arch. Rational Mech. Anal. 31 (1968) 173–184.
- [12] R. Hamilton, *Harmonic maps of manifolds with boundary*, Lecture Notes in Math. Vol. 471, Springer, New York, 1975.
- [13] S. Hildebrandt, H. Kaul & K. O. Widman, *An existence theorem for harmonic mappings of Riemannian manifolds*, Acta Math. 138 (1970) 550–569.
- [14] S. Hildebrandt & K. O. Widman, *Some regularity for quasilinear elliptic systems of second order*, Math. Z. 142 (1975) 67–86.
- [15] ———, *On the Hölder continuity of weak solutions of quasilinear elliptic systems of second order*, Ann. Scuola Norm. Sup. Pisa (IV) 4 (1977) 145–178.
- [16] O. Ladyzhenskaya & N. Ural'tseva, *Linear and quasi-linear elliptic equations*, Academic Press, New York, 1968.
- [17] C. B. Morrey, Jr., *The problem of Plateau on a Riemannian manifold*, Ann. of Math. 49 (1948) 807–851.
- [18] ———, *Multiple integrals in the calculus of variations*, Springer, New York, 1966.
- [19] ———, *Partial regularity results for non-linear elliptic systems*, J. Math. Mech. 17 (1968) 649–670.

UNIVERSITY OF CALIFORNIA, BERKELEY,  
UNIVERSITY OF ILLINOIS, CHICAGO CIRCLE