

## THE DIRICHLET PROBLEM AT INFINITY FOR MANIFOLDS OF NEGATIVE CURVATURE

MICHAEL T. ANDERSON

This paper is concerned with the existence of bounded harmonic functions on simply connected manifolds  $N^n$  of negative curvature. It has been conjectured for some time with such manifolds admit a wealth of bounded harmonic functions provided the sectional curvature  $K_N$  satisfies  $-a^2 \leq K_N \leq -b^2$ , for some constants,  $a, b > 0$ , or even if  $K_N \leq -b^2 < 0$ ; see [7], [18]. Justification for this comes from the fact that the model space  $H^n(-1)$ , the space form of curvature  $-1$ , admits many bounded harmonic functions; in fact, there is a Poisson integral representation 'at infinity' in  $H^n(-1)$ . (Similar results hold in more general noncompact symmetric spaces, cf. [12].) Furthermore, in case  $n = 2$  the Ahlfors-Schwarz Lemma [1] shows that  $N^2$  is conformally the unit disc provided  $K_N \leq -b^2 < 0$ , so that the model  $H^2(-1)$  provides full information in this case.

It is natural to consider a Dirichlet problem at infinity for the Laplace-Beltrami operator  $\Delta$  on  $N^n$ ; there is a well-known compactification  $\bar{N}^n = N^n \cup S^{n-1}(\infty)$  of  $N^n$  giving a homeomorphism of  $(N^n, S^{n-1}(\infty))$  with the Euclidean pair  $(B^n, S^{n-1})$ . One can then state the

**Asymptotic Dirichlet problem for  $\Delta$ .** Given a continuous function  $\rho$  on  $S^{n-1}(\infty)$ , find  $f \in C^\infty(N^n) \cup C^0(\bar{N}^n)$  satisfying

$$\Delta f = 0, \quad f|_{S^{n-1}(\infty)} = \rho.$$

The main result of this paper is given by the following theorem (Theorem 3.2).

**Theorem.** *Let  $N^n$  be a complete simply connected Riemannian manifold with sectional curvature  $K_N$  satisfying  $-a^2 \leq K_N \leq -b^2$ , where  $a^2 \geq b^2$  are arbitrary positive constants. Then the asymptotic Dirichlet problem for  $\Delta$  is uniquely solvable, for any  $\rho \in C^0(S^{n-1}(\infty))$ .*

In particular, it follows that  $N^n$  has a large class of bounded harmonic functions. Using this one may show for instance that there are smooth proper

harmonic maps  $F$  from  $N^n$  onto the Euclidean unit ball (or any other convex domain in  $\mathbf{R}^n$ ), inducing a homeomorphism  $S^{n-1}(\infty) \rightarrow S^{n-1} = \partial B^n$ . For an open neighborhood of metrics sufficiently close to the hyperbolic metric,  $f$  will in fact be a diffeomorphism on  $N^n$ .

There have been a number of nonexistence results along the lines of the above theorem concerning generalizations of the Liouville theorem to Riemannian manifolds. Yau [17] proved that on any complete Riemannian manifold there are no globally defined harmonic functions in  $L^p$  for any  $1 < p < \infty$ . Greene and Wu [8] proved there are no bounded harmonic functions on manifolds  $N^n \approx \mathbf{R}^n$  for which the exponential map from some point is a quasi-isometry. Further, Yau [16] proved that if  $N^n$  has nonnegative Ricci curvature, then there are no bounded harmonic functions on  $N^n$ . In the opposite direction, Choi [4] has recently obtained existence results for spherically symmetric metrics and also in dimension 2, in the case of negative curvature.

The proof of the theorem is based on the classical Perron method of solving the Dirichlet problem. Recall the success of the method hinges upon the existence of barrier functions, that is, subharmonic functions  $B_x: N^n \rightarrow \mathbf{R}$  for  $x \in S^{n-1}(\infty)$ , such that  $B_x \leq 0$  and  $\lim_{y \rightarrow x} B_x(y) = 0$ . Now manifolds  $N^n$  with  $K_N \leq 0$ ,  $\pi_1(N) = 0$  admit a wealth of convex, thus subharmonic functions. However, none of the familiar constructions of such functions give rise to barrier functions, since their behavior at infinity becomes too trivial; consider for instance Busemann functions or distance functions to complete geodesics. Thus our major contribution is the construction of global convex sets having nontrivial asymptotic behavior; from this we deduce the existence of barrier functions for  $\Delta$ .

An outline of the contents of the paper is as follows. After presenting preliminary background material in §0, we discuss in §1 the Perron method, asymptotic maximum principle for harmonic functions and a characterization of the solvability of the Dirichlet problem in terms of convexity conditions at infinity (Theorem 1.4); the material for this section draws heavily on the work of Choi [4]. In §2 we construct a large family of global convex domains in  $N^n$  with controlled behavior at infinity; we refer to §2 for an outline of the construction. This is applied to give the solution to the Dirichlet problem in §3 (Theorem 3.2). We also show that the convex hull  $C(S)$  of closed sets  $S \subset S^{n-1}(\infty)$  is well behaved; in fact  $C(S) \cap S^{n-1}(\infty) = S$ , as is the case for hyperbolic space. This is useful in constructing barriers for complete minimal submanifolds in  $N^n$  and harmonic maps into  $N^n$ . In §4 we introduce harmonic measure on  $S^{n-1}(\infty)$  and a Poisson integral representation of harmonic functions on  $S^{n-1}(\infty)$ ; besides allowing one to construct larger classes of

harmonic functions, one obtains in this fashion a satisfying correspondence between the space of bounded harmonic functions on  $N^n$  and  $L^\infty(S^{n-1}(\infty), \mu)$ ; see Theorem 4.3. Finally, §5 closes with various extensions and remarks.

The author would like to thank Andrejs Treibergs for many helpful conversations during the formative stages of the work; also Richard Schoen for his interest in the work and H. I. Choi for writing [4]. Finally, the author is indebted to M. Gromov for providing the idea for a simpler and better proof of Proposition 2.2; this removed a technical hypothesis required in an earlier version.

We note that Sullivan [15] has recently obtained a proof of the asymptotic Dirichlet problem by quite different methods.

### 0. Preliminaries

Throughout this paper,  $N^n$  will denote a Cartan-Hadamard manifold, that is, a simply connected manifold of nonpositive curvature. The standard model for such spaces is the hyperbolic space form  $H^n(-\lambda^2)$  of constant sectional curvature  $-\lambda^2$ . The *sphere at infinity*  $S^{n-1}(\infty)$  of  $N^n$  is defined to be the set of asymptote classes of geodesic rays; two rays  $\gamma_1, \gamma_2: [0, \infty) \rightarrow N^n$  define the same asymptote class if  $\lim_{t \rightarrow \infty} \text{dist}_N(\gamma_1(t), \gamma_2(t)) < \infty$ . There is a natural topology on  $\overline{N^n} = N^n \cup S^{n-1}(\infty)$ , called the *cone topology*, given as follows: for any origin  $\emptyset \in N^n$ , choose  $v \in T_\emptyset N^n$  and let  $C(v, \delta)$  be the cone around  $v$  of angle  $\delta$ , i.e.,

$$C(v, \delta) = \{x \in N^n \cup S^{n-1}(\infty) : \angle_\emptyset(v, T_{\overline{\emptyset x}}) < \delta\},$$

where  $T_{\overline{\emptyset x}}$  denotes the tangent vector to the geodesic ray  $\overline{\emptyset x}$  through  $\emptyset$  and  $x$ , and  $\angle_\emptyset$  indicates angle in  $TN^n$ . Let  $T(v, \delta, r)$  be the truncated cone of radius  $r$ , i.e.,  $T(v, \delta, r) = C(v, \delta) \setminus B_\emptyset(r)$ ,  $B_\emptyset(r)$  the geodesic  $r$ -ball around  $\emptyset$ . Eberlein and O'Neill [5] have shown that the family  $T(v, \delta, r)$  for  $v \in T_\emptyset N^n$ ,  $\delta > 0$ ,  $r > 0$ , together with the balls  $B_q(r)$ ,  $q \in N^n$ , forms a local basis for the cone topology on  $\overline{N^n}$ ; it turns out this topology is independent of the choice of  $\emptyset$ . In this topology,  $\overline{N^n}$  is homeomorphic to a closed ball  $\overline{B}$  in  $\mathbf{R}^n$ ,  $S^{n-1}(\infty)$  being homeomorphic to the boundary sphere  $S^{n-1} \subset \mathbf{R}^n$ . In fact, if  $\eta: [0, 1] \rightarrow [0, \infty]$  is any homeomorphism, the map  $E_\eta: D_1 \subset T_\emptyset N^n \rightarrow N^n$  given by  $E_\eta(v) = \exp \eta(|v|) \cdot v$  is a homeomorphism of the unit disc  $D_1 \subset T_\emptyset N^n$  onto  $N^n$ , inducing a homeomorphism of the sphere  $S_1 = \partial D_1$  onto  $S^{n-1}(\infty)$ . We note that in general there is no natural (independent of  $\emptyset$ ) differentiable structure on  $S^{n-1}(\infty)$ . Especially in §4, we use the above homeomorphism to identify  $S^{n-1}(1)$  with  $S^{n-1}(\infty)$ .

We introduce some notation which will be used throughout the paper. Let  $\overline{xy}$  denote the geodesic ray determined by  $x$  and  $y$  in  $\overline{N^n}$ . For a given  $v \in T_o N^n$ ,  $x_v \in S^{n-1}(\infty)$  will denote the asymptote class determined by the ray  $\exp tv$ ,  $t \geq 0$ . Geodesic spheres of radius  $r$  will be denoted by  $S(r)$  or  $S_p(r)$  if  $p$  is the center of  $S(r)$ ; similarly geodesic balls are denoted by  $B(r)$  or  $B_p(r)$ . The notation above for cones and truncated cones will be kept throughout the paper. In addition, we adhere to the usual notation in Riemannian geometry and partial differential equations; our main references are [2] and [6] on these matters.

### 1. Dirichlet problem at infinity

In this section, we discuss the Perron method and barrier functions for the solution of the following Dirichlet problem. Much of the material in this section is contained in the work of Choi [4].

**1.0. Dirichlet problem at infinity for  $\Delta$ .** Let  $N^n$  be a simply connected manifold of nonpositive curvature. Given a continuous function  $\rho \in C^0(S^{n-1}(\infty))$ , find  $f \in C^\infty(N^n) \cup C^0(\overline{N^n})$  such that

$$f = 0 \quad \text{in } N^n, \quad f|_{S^{n-1}(\infty)} = \rho.$$

We recall that the topology on  $\overline{N^n}$  is given by the cone topology, and note that convergence in this topology is much stronger than radial convergence (convergence along rays). For example, the function  $f(x, y) = x \cdot y$  is harmonic in the upper half plane (with hyperbolic metric); along all geodesic rays emanating from  $i = (0, 1)$ ,  $f$  converges to 0 on  $S^1(\infty)$ . Nevertheless,  $f$  does not have continuous boundary values on  $S^1(\infty)$  in the cone topology. The classical Phragmen-Lindelöf principle illustrates more precisely the difference between the two topologies.

Using the natural identifications  $S_o^{n-1}(1) \approx S_o^{n-1}(t)$ , for any  $t$ , given by the exponential map, it is easy to see that  $f \in C^0(\overline{N^n})$  has asymptotic boundary values  $\rho$  if and only if the restrictions  $f_t = f|_{S_o^{n-1}(t)}$ , pulled back to functions on  $S_o^{n-1}(1)$ , converge to  $\rho \in C^0(S_o^{n-1}(1))$ .

The following maximum principle is a simple consequence of the definitions.

**Proposition 1.1.** (a) *Let  $f: N^n \rightarrow \mathbf{R}$  be a subharmonic function such that*

$$\overline{\lim}_{x \rightarrow x_\infty} f(x) \leq 0, \quad \text{for any } x_\infty \in S^{n-1}(\infty).$$

*Then  $f \leq 0$  on  $N^n$ .*

(b) If  $f$  is a subharmonic function on  $N^n$ ,  $g$  is a superharmonic function on  $N^n$ , and

$$\overline{\lim}_{x \rightarrow x_\infty} f(x) \leq \underline{\lim}_{x \rightarrow x_\infty} g(x), \text{ for any } x_\infty \in S^{n-1}(\infty),$$

then  $f \leq g$  on  $N^n$ .

*Proof.* We leave this to the reader, or see Choi [4].

Let  $S_\rho$  be the set of subfunctions on  $N^n$  relative to  $\rho$ : that is, given  $\rho \in C^0(S^{n-1}(\infty))$ ,  $S_\rho$  is the set of  $C^0$  subharmonic functions  $v: N^n \rightarrow \mathbf{R}$  such that  $\overline{\lim}_{x \rightarrow x_\infty} v(x) \leq \rho(x_\infty)$ . Clearly,  $S_\rho$  is nonempty. Let  $u(x) = \sup_{v \in S_\rho} v(x)$ ; it is well known that  $u$  is a globally defined harmonic function on  $N^n$ . The function  $u$  defined in this manner is a candidate for the solution of the Dirichlet problem; to show that  $u$  achieves the required boundary values, one needs to construct appropriate barrier functions.

**Definition 1.2.** Let  $v \in T_0N^n$  be a unit vector, and suppose  $\delta > 0$ . Then  $\beta = \beta(v, \delta): N^n \rightarrow \mathbf{R}$  is called a *barrier function* at  $v$  with angle  $\delta$  if

- (1)  $\beta$  is subharmonic,
- (2)  $\beta \leq 0$  and  $\lim_{x \rightarrow x_v} \beta(x) = 0$ ,
- (3)  $\exists \mu > 0$  such that  $\lim_{x \rightarrow x_w} \beta(x) \leq -\mu$  for any  $w \in T_0N^n$  with  $\angle_0(v, w) > \delta$ .

This is a natural analogue of the classical barrier concept for domains in  $\mathbf{R}^n$ ; see [11, 2.6.2] or [4, 2.6]. One then has

**Theorem 1.3.** Suppose there exist barrier functions  $\beta = \beta(v, \delta)$  with arbitrarily small angle  $\delta$  at any  $v \in T_0N^n$ . Then the Dirichlet problem 1.0 at infinity is uniquely solvable for any  $\rho \in C^0(S^{n-1}(\infty))$ .

*Proof.* Let  $u(x)$  be the Perron solution relative to  $\rho \in C^0(S^{n-1}(\infty))$  defined above. We show that  $\lim_{x \rightarrow x_\infty} u(x) = \rho(x_\infty)$  for any  $x_\infty \in S^{n-1}(\infty)$ , which implies the theorem. Fix any  $v \in T_0N^n$  and choose  $\epsilon > 0$ . Let  $\delta > 0$  be such that  $|\rho(x_v) - \rho(x_w)| < \epsilon$  if  $\angle_0(v, w) \leq \delta$ , and let  $\beta = \beta(v, \delta)$  be a barrier function at  $v$ , angle  $\delta$ . If  $\mu$  is given by (3) above, choose  $k$  so that  $\mu k \geq 2M$ , where  $M = \sup |\rho|$  on  $S^{n-1}(\infty)$ .

It is easy to see that  $\psi_1(x) \equiv \rho(x_v) - \epsilon + k\beta(x)$  is a subfunction, and  $\psi_2(x) \equiv \rho(x_v) + \epsilon - k\beta(x)$  is a superfunction relative to  $\rho$ ; since the proofs are almost identical, we verify this claim for  $\psi_2$ . It is clear that  $\psi_2(x)$  is superharmonic on  $N^n$ . To show that  $\lim_{x \rightarrow x_w} \psi_2(x) \geq \rho(x_w)$  for any  $w \in T_0N^n$ , suppose first that  $\angle_0(x_v, x_w) \leq \delta$ . Then  $|\rho(x_v) - \rho(x_w)| \leq \epsilon$  and since  $\beta \leq 0$ ,  $\psi_2(x) \geq \rho(x_w)$ , whenever  $x \in C(v, \delta)$ . If  $\angle_0(x_v, x_w) > \delta$ , we have

$$\underline{\lim}_{x \rightarrow x_w} \psi_2(x) = \rho(x_v) + \epsilon - k \cdot \overline{\lim}_{x \rightarrow x_w} \beta(x) \geq \rho(x_v) + \epsilon + k\mu \geq \rho(x_w),$$

as required. From the definition of  $u$  as the Perron solution and by Proposition 1.1(b), we find  $\psi_1(x) \leq u(x) \leq \psi_2(x)$ , or  $|u(x) - \rho(x_v)| \leq \varepsilon - k\beta(x)$ . Since  $\beta(x) \rightarrow 0$  as  $x \rightarrow x_v$  and  $\varepsilon$  is arbitrary, it follows that  $\lim_{x \rightarrow x_v} u(x) = \rho(x_v)$  as desired. This proves the solvability of the Dirichlet problem for arbitrary  $\rho \in C^0(S^{n-1}(\infty))$ . Uniqueness follows from the maximum principle (Proposition 1.1(a)) in the usual fashion. q.e.d.

It is well known that Cartan-Hadamard manifolds  $N^n$  possess a wealth of convex functions; typical examples are distance functions to a point or to a geodesic, or the mean square distance to a compact submanifold. In case  $K_N \leq -c^2 < 0$ , such convex functions may be reparametrized to give bounded subharmonic functions; see e.g. [4]. Thus one may hope to obtain the existence of barrier functions from suitable convex functions or sets, as is the case for bounded domains in  $\mathbf{R}^n$ .

**Theorem 1.4 (Choi).** *Suppose for any distinct  $x, y \in S^{n-1}(\infty)$  there exist disjoint open neighborhoods  $V_x, V_y$  of  $x, y$  in  $\overline{N^n}$  such that  $V_x \cap N^n$  and  $V_y \cap N^n$  are strictly convex. Then if  $K_N \leq -1$ , the Dirichlet problem at infinity for  $\Delta$  is uniquely solvable for any  $\rho \in C^0(S^{n-1}(\infty))$ .*

*Proof.* By Theorem 1.3 we need to construct a barrier  $\beta = \beta(v, \delta)$  for  $v \in T_0 N^n$  and any small  $\delta > 0$ . Given  $x_v \in S^{n-1}(\infty)$ , let  $S(v, \delta) = \{x_w \in S^{n-1}(\infty) : \angle_0(x_v, x_w) = \delta\}$ . Since  $S(v, \delta)$  is compact, we may cover it by a finite number of convex open sets  $\{V_{w_i}\}_{i=1}^m$  such that

$$V_{w_i} \cap V_{x_v} = \emptyset, \quad \text{for all } i,$$

where  $V_{x_v}$  is an open neighborhood of  $x_v$  in  $\overline{N^n}$ ; this much follows from the hypothesis of the theorem. Let  $\Omega = N^n - \bigcup_1^m V_{w_i}$ , and let  $s_i: \Omega \rightarrow \mathbf{R}^+$  be the distance function from  $V_{w_i}$ . By an approximation theorem of Greene and Wu [9, Proposition 2.2], we may assume each  $\partial V_{w_i}$ , and thus each  $s_i$  is a smooth function. Using the second variational formula one may show that

$$\text{Hess}(s_i) \geq \tanh\left(\frac{s_i}{2}\right) \cdot H_0,$$

where  $H_0 = ds^2 - ds_i \otimes ds_i$ ,  $ds^2$  is the metric on  $N^n$ ; [4]. From this it follows easily that

$$\Delta \tanh \frac{s_i}{2} \geq 0,$$

on  $\Omega$ . Now let  $\bar{\beta} = \sum_{i=1}^m \tanh(s_i/2) - m$ ; clearly  $\bar{\beta}$  is subharmonic, nonpositive and  $\lim_{x \rightarrow x_\infty} \bar{\beta}(x) = 0$  for any  $x_\infty \in V_{x_v} \cap S^{n-1}(\infty)$ . Choose  $R > 0$  so large that  $\Omega$  disconnects as the disjoint union  $\Omega_1 \cup \Omega_2$  outside  $B_\delta(R)$ . Let  $\Omega_1$  be the

set of  $q \in \Omega$  such that  $\text{dist}_N(\emptyset, q) > R$  and  $\angle_{\emptyset}(\overline{\emptyset q}, v) < \delta$ . Let  $\overline{\Omega} = \{x \in \Omega_1: \overline{\beta}(x) \geq c\}$ , where  $-m < c < 0$  is a fixed constant, sufficiently close to 0. Define  $\beta$  on  $N^n$  by

$$\beta(x) = \begin{cases} \overline{\beta}(x), & x \in \overline{\Omega}, \\ c, & x \in (\overline{\Omega})^c. \end{cases}$$

Then  $\beta$  is a barrier at  $v$ , of angle  $\delta$  ( $\mu = -c$ ). Since  $v$  and  $\delta$  are arbitrary, the result follows. q.e.d.

It appears that none of the standard constructions of convex functions on manifolds  $N^n$  of negative curvature give rise to convex sets satisfying the conditions of Theorem 1.4. In fact, there are no examples of convex sets  $C$  in general  $N^n$  such that  $C \cap S^{n-1}(\infty) \neq S^{n-1}(\infty)$  is a set with nonempty interior in the cone topology. Thus our aim in the next section is to construct convex sets with nontrivial behavior at infinity.

### 2. Construction of convex sets

In this section we will construct unbounded convex domains in manifolds of negative curvature, having prescribed asymptotic behavior. Recall that horoballs  $H_x$  in  $N^n$  are strictly convex sets intersecting  $S^{n-1}(\infty)$  at a single point  $x$ ; the construction is based on the idea that locally one may produce larger convex sets containing  $H_x$  (due to the strict convexity): this is done in Step I below. In Step II we use an iteration procedure to construct global convex domains  $\mathcal{C}$ . In Step III by using comparison with negative space forms we show convergence and are able to control the behavior of  $\mathcal{C}$  at infinity.

Throughout this section  $N^n$  will denote a simply connected Cartan-Hadamard manifold of sectional curvature  $K_N$  satisfying  $-a^2 \leq K_N \leq -b^2$ ; using a homothety of the metric, we may assume  $b = 1$ .

**Step I.** An important feature of the spaces  $N^n$  is the convexity of large geodesic spheres. In fact, if  $\Pi_R$  denotes the second fundamental form (with respect to inward normal) of a geodesic  $R$ -sphere in  $N^n$ , then standard arguments involving Jacobi fields show that

$$(2.1) \quad I \leq \coth R \leq \Pi_R \leq a \coth aR \cdot I,$$

where  $I$  denotes the identity matrix.

In order to find local convex expansions of spheres we use the following lemma.

**Lemma 2.1.** *Given any  $p \in N^n$ , there is an  $f_p \in C^\infty(N^n, \mathbf{R})$  such that  $0 \leq f_p \leq 1, f_p(p) = 0, f \equiv 1$  outside  $B_p(1)$  and*

$$(2.2) \quad |df_p| < C_1, \quad |D_{ij}^2 f_p| < C_2,$$

where  $C_1, C_2$  are constants depending only on  $a^2$ .

*Proof.* Let  $\rho$  be the distance function from  $p$ , and consider functions  $f_p = \phi(\rho)$  where  $\phi: \mathbf{R}^+ \rightarrow \mathbf{R}$  satisfies  $\phi(0) = 0, \phi(t) = 1, \text{ for } t \geq 1$ . Then  $df_p = \phi' \cdot d\rho$ , so that  $|df_p| = |\phi'|$ . A simple computation shows that  $D^2\phi(\rho) = \phi'' d\rho \otimes d\rho + \phi' D^2\rho$ ; thus

$$|D_{ij}^2 \phi(\rho)| \leq |\phi''| + |\phi'| a \coth a\rho.$$

Choosing  $\phi$  appropriately, e.g., as a fixed approximation to the characteristic function  $\chi_{[1/2, \infty]}$ , one easily obtains the estimates (2.2). *q.e.d.*

Now given a fixed origin  $\emptyset$ , consider the functions  $\rho_\emptyset + \epsilon f_p$  where  $\rho_\emptyset$  is the distance function from  $\emptyset$ ; these may be considered as local perturbations of  $\rho_\emptyset$ , provided  $\rho_\emptyset(p) \geq 10$  say.

**Proposition 2.2.** *There is an  $\epsilon > 0$  depending only on  $a^2$  such that the sublevel sets of  $\rho_\emptyset + \epsilon f_p$  are totally convex subsets of  $N^n$ .*

*Proof.* Note that the level sets of  $\rho_\emptyset + \epsilon f_p$  are smooth submanifolds of  $N^n$ ; since  $N^n$  has negative curvature, it is sufficient to show they have positive definite second fundamental form  $\Pi_{\epsilon f}$ . For a given tangent vector  $X$  we have

$$\Pi_{\epsilon f}(X, X) = \langle \nabla_X N, X \rangle = \frac{1}{|d(\rho_\emptyset + \epsilon f_p)|} \cdot D^2(\rho_\emptyset + \epsilon f_p)(X, X).$$

We have  $D^2(\rho_\emptyset) \geq I$  by (2.1) and  $D^2(\epsilon f_p) \leq \epsilon C_2$  by (2.2); choosing  $\epsilon$  sufficiently small then gives the result.

**Remark 2.3.** It is clear that Proposition 2.2 remains valid when  $f_p = \phi \circ \rho$  is replaced by  $\phi_R \circ \rho$  where  $\phi_R(t) = \phi(R \cdot t)$ ; then  $\epsilon$  depends on  $R$  as well as  $a^2$ . In order to simplify the computations in Step III, we will choose  $R$  to satisfy the following requirement. Let  $S_\emptyset(p)$  be the sphere around  $\emptyset$  containing  $p$ , and  $S_\emptyset(\epsilon)$  the concentric spheres with radius  $\text{dist}(\emptyset, p) + \epsilon$ . Then the level set of  $\rho_\emptyset + \epsilon f_p$  through  $\underline{p}$  should equal  $S_\emptyset(\epsilon)$  outside the intrinsic ball of radius 1 in  $S_\emptyset(\epsilon)$  centered at  $\underline{op} \cap S_\emptyset(\epsilon)$  (instead of the extrinsic ball of radius 1 centered at  $p$ ). It is not difficult to see that for  $\text{dist}(\emptyset, p) \geq C$ , e.g.,  $\text{dist}(\emptyset, p) \geq 10$ , such an  $R$  may be found independent of  $\emptyset, p$ .

**Step II.** We now construct global convex domains, using iteration of the local result contained in Proposition 2.2. Thus let  $S = S_\emptyset(R)$  be a fixed geodesic  $R$ -sphere and let  $S(t) = S_\emptyset(R + t)$ . Choose  $p \in S$  and define  $C_1$  to be the sublevel set of  $\rho_\emptyset + \epsilon f_p$  with  $p \in \partial C_1$ ; here and in what follows,  $\epsilon$  and  $f_p$  are defined by Proposition 2.2 and Remark 2.3. One may view  $C_1$  as the ball




$B(\epsilon) = B_0(R + \epsilon)$  with a ‘scallop’ cut out around  $p$ ;  $C_1$  is a totally convex domain.

One may proceed to cut out successively larger ‘scallops’ from successively larger spheres as follows. Let  $T_1$  be the ‘seam’ of  $C_1$ :  $T_1 = \partial(C_1 \cap S(\epsilon))$ . Let  $U_{T_1}$  be the collection of points on  $S_1 = S(\epsilon)$  whose outward intrinsic distance to  $T_1$  is  $\leq 1$ ; note this includes all points of  $S_1$  in the interior of  $T_1$  (the component containing  $\overline{op}$ ). For any  $q \in U_{T_1}$ , define the convex set  $C_2(q)$  to be the sublevel set of  $\rho_0 + \epsilon f_q$  such that  $q \in \partial C_2(q)$ , and set

$$C'_2 = \bigcap_{q \in U_{T_1}} C_2(q).$$

It is clear that  $C'_2$  is totally convex. Let  $C_2 = C'_2 \setminus (B(\epsilon) \setminus C_1)$ .

Note that  $C_1 \subset B_1 = B(\epsilon)$ , and  $B(\epsilon) \setminus C_1$  is the ‘scallop’ bored out of  $B(\epsilon)$ . It is easy to see that  $C_2$  is also totally convex. In fact let  $x, y \in \partial C_2$  and suppose not both  $x$  and  $y$  are in  $\partial C_1$ . Then  $\gamma = \overline{xy}$  is obviously contained in  $C'_2$ ; suppose  $\gamma$  entered the scallop  $B(\epsilon) \setminus C_1$ . By convexity of  $C_1$ , at least one end  $z$  of  $\gamma \cap B(\epsilon)$  does not lie in  $\partial C_1$ . However, then one of the two geodesic arcs  $zx$  or  $zy$  must intersect the complement of  $C_2(q)$  for some  $q$ , contradicting the convexity of  $C_2(q)$ .



It is now clear how we proceed inductively; the ‘seam’  $T_i$  of  $C_i$ ,  $T_i = \partial(C_i \cap S(i\epsilon))$ , serves to construct  $C'_{i+1}$  as

$$C'_{i+1} = \bigcap_{q \in U_{T_i}} C_{i+1}(q),$$

where  $C_{i+1}(q)$ ,  $U_{T_i}$  are defined as before. Then  $C_{i+1}$  is given by  $C_{i+1} = C'_{i+1} - (B_i - C_i)$ . We thus have a nested sequence of totally convex sets

$$C_1 \subset C_2 \subset \dots \subset C_k \subset C_{k+1} \subset \dots$$

Let  $\mathcal{C} = \bigcup_{i=1}^{\infty} C_i$ ; this is the desired global totally convex set. Note that we construct such  $\mathcal{C}$  for any  $p \in S$  and any  $S$  (of radius  $\geq 10$ ).

**Step III.** Having constructed global convex domains  $\mathcal{C}$  in  $N^n$ , we need to show that  $\mathcal{C}$  has ‘nontrivial’ asymptotic behavior. In particular, we need to control the size of  $\mathcal{C} \cap S^{n-1}(\infty)$ . The study of  $\mathcal{C} \cap S^{n-1}(\infty)$  uses comparison

of  $\mathcal{C}$  with models in the spaces  $H^n(-1)$  and  $H^n(-a^2)$ . It turns out it is actually sufficient to compare  $\mathcal{C}$  with models in  $H^2(-1)$  and  $H^2(-a^2)$ . Thus choose an origin  $o$  in  $H^2(-1)$  and  $H^2(-a^2)$  together with  $R$ -spheres  $S(R)$  centered at  $o$ , and let  $p \in S(R)$ . We construct the models  $\bar{\mathcal{C}} \subset H^2(-1)$  exactly according to the prescription given for  $\mathcal{C}$  in Steps I and II above (using the same or any equivalent  $f$ ). Similarly, construct models  $\underline{\mathcal{C}} \subset H^2(-a^2)$ , but replacing the intrinsic distances 1 by  $\frac{1}{2}$  everywhere.

We measure the asymptotic behavior of  $\mathcal{C}$  (and  $\bar{\mathcal{C}}, \underline{\mathcal{C}}$ ) by means of the angle at  $o$  from the ray  $\overline{op}$ ; in other words, for any  $x \in \mathcal{C} \subset S^{n-1}(\infty)$ , consider  $\angle_o(\overline{op}, \overline{ox})$ . Let

$$\alpha_{\mathcal{C}} = \sup\{\angle_o(\overline{op}, \overline{ox}) : x \in \mathcal{C} \cap S^{n-1}(\infty)\},$$

$$\beta_{\mathcal{C}} = \inf\{\angle_o(\overline{op}, \overline{ox}) : x \in \mathcal{C} \cap S^{n-1}(\infty)\}.$$

Similarly we define  $\alpha_{\bar{\mathcal{C}}}, \beta_{\bar{\mathcal{C}}}$  and  $\alpha_{\underline{\mathcal{C}}}, \beta_{\underline{\mathcal{C}}}$ ; it is easy to see that in fact  $\alpha_{\bar{\mathcal{C}}} = \beta_{\bar{\mathcal{C}}}$  and  $\alpha_{\underline{\mathcal{C}}} = \beta_{\underline{\mathcal{C}}}$ .

**Lemma 2.4.** *In the above notation, we have*

$$(2.5) \quad \alpha_{\mathcal{C}} \leq \alpha_{\bar{\mathcal{C}}}, \quad \beta_{\mathcal{C}} \leq \beta_{\underline{\mathcal{C}}}.$$

*Proof.* (i)  $\alpha_{\mathcal{C}} \leq \alpha_{\bar{\mathcal{C}}}$ . Referring to Step II in the construction of  $\mathcal{C}$ , let  $x_N \in T_N$  be a sequence such that  $x_N \rightarrow x_\infty$  and  $\angle_o(\overline{ox_\infty}, \overline{op}) = \alpha_{\mathcal{C}}$ . For each fixed  $N$ , let  $x_N^i \in T_i$  be chosen inductively so that  $x_N^N = x_N$  and  $\text{dist}_{S(i)}(x_N^i, (x_N^{i+1})') \leq 2$  where  $(x_N^{i+1})'$  is the normal geodesic projection of  $x_N^{i+1}$  onto  $S(i) = S(R + i\epsilon)$ . Consider the geodesic hinge  $x_N^i \overline{ox_N^{i+1}}$ ; set  $\alpha_N^i = \angle_o(\overline{ox_N^i}, \overline{ox_N^{i+1}})$  and note that  $l(\overline{ox^i}) = R + \epsilon$  for any  $N$ . Now consider analogous hinges in  $\bar{\mathcal{C}} \subset H^2(-1)$ . In this case, each  $T_i$  consists of two points  $\{\chi_i, \xi_i\}$ ; we let  $\chi_i \in T_i$  be chosen so that  $\{T_{\overline{o\chi_i}}, T_{\overline{op}}\}$  is an oriented basis for  $T_o(H^2(-1))$ . Let  $\bar{\alpha}^i = \angle_o(\overline{op}, \overline{o\chi_i})$ ; we have  $l(\overline{o\chi_i}) = R + i\epsilon$  and  $\text{dist}_{S(i)}(\chi_i, \chi_{i+1}) = 2$ . It now follows from the Rauch comparison theorem (see [2, 1.28, 1.30]), that

$$\alpha_N^i \leq \bar{\alpha}^i, \quad \text{for all } i, \text{ any } N.$$

Thus we have shown

$$\alpha_{\mathcal{C}} \leq \lim_{N \rightarrow \infty} \sum_{i=1}^N \alpha_N^i \leq \sum_{i=1}^{\infty} \bar{\alpha}^i = \alpha_{\bar{\mathcal{C}}},$$

where the last equality follows from the fact that  $H^2(-1)$  is 2-dimensional.

(ii)  $\beta_{\underline{\mathcal{C}}} \leq \beta_{\mathcal{C}}$ . Again we choose  $x_\infty \in \mathcal{C} \cap S^{n+1}(\infty)$  realizing  $\beta_{\mathcal{C}}$ . Let  $P \subset T_0 N^n$  be the 2-plane spanned by the vectors  $\{T_{\overline{op}}, T_{\overline{ox_\infty}}\}$ , and let  $\mathfrak{P} = \exp_0 P$ . Choose  $x_i \in T_i \cap \mathfrak{P}$  so that  $\text{dist}_{S(i)}(x_i, (x_{i+1})') \geq 1$  where  $(x_{i+1})'$  is as in (i); it is clear that such choices of  $x_i$  always exist and  $x_i \rightarrow x_\infty$ . We let  $\beta_i = \angle_o(\overline{ox_i}, \overline{ox_{i+1}})$

and recall  $l(\overline{o\chi_i}) = R + i\varepsilon$ . Now compare this data with the model  $\mathcal{C} \subset H^2(-a^2)$ ; choose  $\chi_i \subset T_i \subset \mathcal{C}$  as in (i) and set  $\beta_i = \angle_o(\overline{op}, \overline{o\chi_i})$  in  $H^2(-a^2)$ . Since  $l(o\chi_i) = R + i\varepsilon$  and  $\text{dist}_{S_i}(\chi_i, (\chi_{i+1})') = 1$  (recall 1 is replaced by  $\frac{1}{2}$  in  $\mathcal{C}$ ), it follows by Rauch comparison as above that

$$\underline{\beta}_i \leq \beta_i, \quad \text{for all } i.$$

Since all angles  $\beta_i$  are measured in  $P$ , we have  $\beta_{\mathcal{C}} = \sum_{i=1}^{\infty} \beta_i$ , which gives

$$\beta_{\mathcal{C}} = \sum_{i=1}^{\infty} \underline{\beta}_i \leq \sum_{i=1}^{\infty} \beta_i = \beta_{\mathcal{C}}. \quad \text{q.e.d.}$$

We are now ready to obtain estimates for  $\alpha_{\mathcal{C}}$  and  $\beta_{\mathcal{C}}$  in terms of  $\varepsilon, a, R$ . For the calculations we use the Poincaré model for  $H^2(-\lambda^2)$ ; recall the hyperbolic metric on the Euclidean ball  $B^2(1)$  is given by

$$ds_{\lambda}^2 = \frac{4}{\lambda^2(1-r^2)^2} ds_E^2,$$

where  $ds_E^2$  is the Euclidean metric. One easily computes that at Euclidean distance  $r \in (0, 1)$ , the hyperbolic distance  $d_{\lambda}$  is given by

$$d_{\lambda} = \frac{1}{\lambda} \log \frac{1+r}{1-r}.$$

Also, the intrinsic hyperbolic distance  $S_{\lambda}$  on  $S(R)$  is  $S_{\lambda} = 2S_E/[\lambda(1-r^2)]$ , for  $S_E$  the intrinsic Euclidean distance.

Now given  $T_i \subset \mathcal{C}$  or  $\bar{\mathcal{C}}$  in  $H^2(-\lambda^2)$  for  $\lambda^2 = 1, a^2$ , we measure the position of  $T_i$  with respect to  $o$  and the ray  $\overline{op}$  by means of ‘polar coordinates’:  $T_i = (l_i, \theta_i)$ , where  $l_i$  is the radius of the Euclidean sphere with  $T_i \in S(l_i)$ , and  $\theta_i$  is the hyperbolic angle at  $o$  between  $\overline{op}$  and  $\overline{oT_i}$ . We have  $T_0 = (l_0, 0)$  where  $l_0 = (e^{\lambda R} - 1)/(e^{\lambda R} + 1)$ , since  $T_0 = p$  is on the geodesic  $R$ -sphere centered at  $o$ . By construction,  $l_k$  is given by the formula

$$k\varepsilon = \frac{1}{\lambda} \int_{l_0}^{l_k} \frac{2}{1-t^2} dt.$$

Writing  $l_k = l + \mu_k$ , one finds that

$$\mu_k = \frac{(1+l_0)[e^{k\lambda\varepsilon} - 1]}{1 + e^{k\lambda\varepsilon}((1+l_0)/(1-l_0))}.$$

To compute  $\theta_k$ , recall that

$$S_{\lambda} = \frac{2}{\lambda} \left( \frac{1}{1-r^2} \right) \cdot S_E;$$

by construction we require  $S_\lambda = \text{dist}_{S(i)}(\chi_i, (\chi_{i+1})') = 2\sigma$ , where  $\sigma = 1$  in  $H^2(-1)$  and  $\sigma = \frac{1}{2}$  in  $H^2(-a^2)$ . This gives  $\theta_1 = S_\varepsilon/l_0 = \lambda\sigma(1 - l_0^2)/l_0$  and generally

$$\theta_k = \frac{\lambda\sigma(1 - l_{k-1}^2)}{l_{k-1}}.$$

For the coordinates of  $T_k$ , we then have

$$T_k = \left( l_0 + \mu_k, \lambda\sigma \sum_{i=0}^{k-1} \frac{1 - l_i^2}{l_i} \right).$$

Set  $\Omega_\lambda(R, \varepsilon) = \lim_{k \rightarrow \infty} \theta_k$ . Then  $\mu_k \rightarrow 1 - l_0$  as  $k \rightarrow \infty$  and a lengthy but straightforward computation shows that

$$\begin{aligned} (2.6) \quad \Omega_\lambda(R, \varepsilon) &= 4(1 - l_0^2)\lambda\sigma \cdot \sum_{i=0}^{\infty} \frac{1}{(1 + l_0)^2 e^{i\lambda\varepsilon} - (1 - l_0)^2 e^{-i\lambda\varepsilon}} \\ &= 4\lambda\sigma \sum_{i=0}^{\infty} \frac{1}{e^{2\lambda\varepsilon} e^{i\lambda\varepsilon} - e^{-i\lambda\varepsilon}}. \end{aligned}$$

It is clear that this series converges uniformly on compact sets in both variables  $R > 0$  and  $\varepsilon > 0$ ; thus  $\Omega_\lambda$  is a continuous function of  $R$  and  $\varepsilon$ . In particular, setting  $\lambda = 1$  and  $\lambda = a$ , and substituting in the value for  $\sigma$  above, we obtain bounds on  $\alpha_{\mathbb{C}}$  and  $\beta_{\mathbb{C}}$ :

$$\begin{aligned} (2.7) \quad \alpha_{\mathbb{C}} &= 4 \sum_{i=0}^{\infty} \frac{1}{e^{2R} e^{ie} - e^{-ie}}, \\ \beta_{\mathbb{C}} &= 2a \sum_{i=0}^{\infty} \frac{1}{e^{2aR} e^{iae} - e^{-iae}}. \end{aligned}$$

Of course, we are only interested in the case when  $\beta_{\mathbb{C}} \leq \alpha_{\mathbb{C}}$ ; this occurs for example for  $R$  satisfying  $(a - 1)R \geq \ln(a/2)$ ; so  $R \geq 1$  is sufficient.

Recall that  $\varepsilon$  is determined solely in terms of the constant  $a$  from Proposition 2.2. The estimates (2.7) in conjunction with Lemma 2.4 provide estimates for the behavior of  $\mathbb{C}$  at infinity. In the next section, these will be used to discuss the convexity of  $S^{n-1}(\infty)$  and the solution of the Dirichlet problem.

### 3. Convexity of $S^{n-1}(\infty)$ : Solution of the Dirichlet problem

We use the results of §2 to discuss the convexity of  $N^n$  at infinity. First, we prove the existence of arbitrarily ‘small’ convex neighborhoods for  $x \in S^{n-1}(\infty)$  in the cone topology; this leads to the solution of the Dirichlet

problem at infinity for  $\Delta$ . We also prove an important property of the convex hull of closed sets  $S \subset S^{n-1}(\infty)$ , namely,  $\mathcal{C}(S) \cap S^{n-1}(\infty) = S$ ; this property is well known in hyperbolic space  $H^n(-1)$ .

We continue to assume that  $N^n$  satisfies  $-a^2 \leq K_N \leq -1$ .

**Theorem 3.1.** *Given any  $v \in T_0(N^n)$  and  $\delta > 0$ , there are convex domains  $K_{v,\delta}(l)$  in  $N^n$  satisfying*

- (i)  $K_{v,\delta}(l) \subset T(v, \delta, l)$ , for  $l \geq \bar{l}$ , where  $\bar{l}$  depends continuously on  $\delta$  and  $a$ ,
- (ii)  $\mathcal{C}(v, \delta') \cap S^{n-1}(\infty) \subset K_{v,\delta}(l) \cap S^{n-1}(\infty)$ , where  $\delta' > 0$  depends continuously on  $a$  and  $l$ .

*Proof.* Let  $x_v \in S^{n-1}(\infty)$  be the asymptote class determined by  $v$ , and  $\overline{ox_v}$  the ray from  $o$  to  $x$ , and let  $o_l = S_o(l) \cap \overline{ox_v}$  for  $l > 0$ . Consider the spheres  $S_o(R)$  of radius  $R < l$  around  $o_l$  and set  $p_R = S_o(R) \cap \overline{ox_v}$ . Define  $\mathcal{C}(p_R, o_l)$  to be the convex domain constructed in §2 determined by the center  $o_l$  and point  $p_R$ ; we will show that for appropriate choices of  $R$  and  $l$ ,  $K_{v,\delta}(l) \equiv \mathcal{C}(p_R, o_l)$  satisfies (i) and (ii).

(i) Given any  $\delta > 0$ , we claim there is an  $\bar{l}$  such that  $l \geq \bar{l}$  implies  $\mathcal{C}(p_R, o_l) \subset T(v, \delta, l)$  for any  $R_0(a) \leq R \leq R_1(a)$ , where  $R_0, R_1$  are fixed constants depending only on  $a$ . To show this, recall that  $\beta_{\mathcal{C}} = \inf\{\angle_{o_l}(\overline{o_l p_R}, \overline{o_l x}): x \in \mathcal{C} \cap S^{n-1}(\infty)\}$ , and  $\beta_{\mathcal{C}} \geq \beta_{\mathcal{C}}$  by Lemma 2.4. Consider the geodesic triangle  $oo_l x_\infty$  where  $x_\infty \in S^{n-1}(\infty)$  realizes  $\beta_{\mathcal{C}}$ , and let  $\Omega = \angle_o(\overline{oo_l}, \overline{ox_\infty})$ . Let  $oo_l \chi_\infty$  be a similar triangle in  $H^2(-1)$  where  $\text{dist}(o, o_l) = l$ ,  $\chi_\infty \in S^1(\infty)$  and  $\angle_{o_l}(\overline{o_l o}, \overline{o_l \chi_\infty}) = \beta_{\mathcal{C}}$ . Setting  $\bar{\Omega} = \angle_o(\overline{oo_l}, \overline{o\chi_\infty})$  in  $H^2(-1)$ , the Rauch comparison theorem applied to the two triangles implies that

$$\Omega \leq \bar{\Omega}.$$

Now  $\beta_{\mathcal{C}}$  depends on  $a$  and  $R$  according to (2.7); in particular, there are constants  $R_0(a)$  and  $R_1(a)$  so that  $R_0(a) \leq R \leq R_1(a)$  implies that  $\pi/2 \leq \beta_{\mathcal{C}} \leq 3\pi/4$ , independent of  $l$ . It follows by elementary hyperbolic geometry that  $\bar{\Omega}$  can be made arbitrarily small by choosing  $l$  sufficiently large; thus there exists  $\bar{l}$  so that  $\Omega \leq \bar{\Omega} < \delta$  for  $l \geq \bar{l}$ .

We have proved that  $\mathcal{C}(p_R, o_l) \cap S^{n-1}(\infty) \subset C_o(v, \delta) \cap S^{n-1}(\infty)$  for  $l \geq \bar{l}$ ,  $R_0(a) \leq R \leq R_1(a)$ . By examining the construction of  $\mathcal{C}(p_R, o_l)$  in §2, this is easily seen to imply that

$$\mathcal{C}(p_R, o_l) \subset T(v, \delta, l - R_1(a)), \quad l \geq \bar{l}.$$

(ii) To see that  $K_{v,\delta}(l)$  intersects  $S^{n-1}(\infty)$  in a set of nonempty interior, choose  $y_\infty \in S^{n-1}(\infty)$  realizing  $\alpha_{\mathcal{C}} = \sup\{\angle_{o_l}(\overline{o_l p_R}, \overline{o_l x}): x \in \mathcal{C} \cap S^{n-1}(\infty)\}$ , where  $\mathcal{C} = \mathcal{C}(p_R, o_l)$  as in (i). Consider the geodesic triangle  $oo_l y_\infty$  and let  $\omega = \angle_o(\overline{oo_l}, \overline{oy_\infty})$ . We form the comparison triangle  $oo_l \xi_\infty$  in  $H^2(-a^2)$  where  $\text{dist}(o, o_l) = l$  and  $\angle_{o_l}(\overline{o_l o}, \overline{o_l \xi_\infty}) = \alpha_{\mathcal{C}}$  in  $H^2(-a^2)$ . Let  $\underline{\omega} = \angle_o(\overline{oo_l}, \overline{o\xi_\infty})$ ; since

$\alpha_c \leq \alpha_{\bar{c}}$  by Lemma 2.4, Rauch comparison applied to the pair of triangles gives

$$\underline{\omega} \leq \omega.$$

This shows that  $C_o(v, \underline{\omega}) \cap S^{n-1}(\infty) \subset \mathcal{C}(p_R, o_l) \equiv K_{v,\delta}(l)$ . It is clear that  $\delta' \equiv \underline{\omega} > 0$  depends continuously on  $a, l$  and  $R$ . Since  $R$  is bounded by  $R_0(a) \leq R \leq R_1(a)$ ,  $\delta'$  depends only on  $a$  and  $l$ . q.e.d.

As a consequence of Theorem 3.1, we deduce our main theorem on the solvability of the Dirichlet problem.

**Theorem 3.2** (*Dirichlet problem at infinity*). *Let  $N^n$  be a complete simply connected Riemannian manifold of sectional curvature  $K_N$  satisfying  $-a^2 \leq K_N \leq -1$ , where  $a^2 \geq 1$  is an arbitrary constant. Then the Dirichlet problem at infinity for  $\Delta, (1.0)$ , is uniquely solvable for any  $\rho \in C^0(S^{n-1}(\infty))$ .*

*Proof.* By Theorem 1.4 we need to prove that for pairs  $x, y \in S^{n-1}(\infty)$ ,  $x \neq y$ , there exist disjoint open sets  $V_x, V_y$  in  $\bar{N}^n$  relative to the cone topology so that  $V_x \cap N$  and  $V_y \cap N$  are convex. Let  $v, w \in T_0N^n$  be chosen so that  $x, y$  are the asymptote classes of the rays determined by  $v, w$ . Choose  $\delta > 0$  so that  $C_o(v, \delta) \cap C_o(w, \delta) = \emptyset$ . By Theorem 3.1 we may choose convex domains  $K_x$  and  $K_y$  so that  $K_x \subset C_o(v, \delta)$ ,  $K_y \subset C_o(w, \delta)$  and  $K_x \cap S^{n-1}(\infty), K_y \cap S^{n-1}(\infty)$  contain the intersection of  $\delta'$ -cones around  $v, w$  with  $S^{n-1}(\infty)$ . In particular, the domains  $K_x, K_y$  satisfy the above conditions. q.e.d.

Given a set  $S \subset \bar{N}^n$ , we define the *convex hull*  $\mathcal{C}(S)$  of  $S$  to be the smallest geodesically closed set in  $\bar{N}^n$  containing  $S$ . The manifold  $N^n$  is said to satisfy the *Visibility property* if given any distinct  $x, y \in S^{n-1}(\infty)$ , there is a complete geodesic  $\gamma$  in  $N^n$  asymptotic to  $x$  and  $y$ ; see [5]. As a simple special case,  $N^n$  is Visibility if  $K_N \leq -b^2 < 0$ . The Visibility manifolds are the natural class of manifolds in which to study the convexity of  $S^{n-1}(\infty)$ . A characteristic property of the convexity of the model space  $H^n(-1)$  at infinity is the fact  $\mathcal{C}(S) \cap S^{n-1}(\infty) = S$  for any closed set  $S \subset S^{n-1}(\infty)$ . We generalize this to other Riemannian manifolds as follows.

**Theorem 3.3.** *Let  $N^n$  be a complete simply connected manifold satisfying the conditions of Theorem 3.2. If  $S$  is a closed set in  $S^{n-1}(\infty)$ , then*

$$(3.3) \quad \mathcal{C}(S) \cap S^{n-1}(\infty) = S.$$

*Proof.* We note that it is sufficient to prove the existence of ‘large’ convex sets  $\mathcal{C}_{v,\delta}$  for any  $\delta > 0, v \in T_0N^n$  such that

$$S^{n-1}(\infty) / (C(v, \delta) \cap S^{n-1}(\infty)) \subset \mathcal{C}_{v,\delta} \cap S^{n-1}(\infty),$$

but  $x_v \notin \mathcal{C}_{v,\delta} \cap S^{n-1}(\infty)$ . For given such, let  $x \in S^{n-1}(\infty) \setminus S$ ; then there exists  $\delta$  and  $v$  with  $x = x_v \in C_o(v, \delta) \cap S^{n-1}(\infty)$  satisfying  $C_o(v, \delta) \cap S^{n-1}(\infty) \subset S^{n-1}(\infty) \setminus S$ . Choose  $\mathcal{C}_{v,\delta}$  as above; it follows that  $x \notin \mathcal{C}_{v,\delta}$  yet  $S \subset \mathcal{C}_{v,\delta}$ . Since  $\mathcal{C}(S) \subset \mathcal{C}_{v,\delta}$ , we have  $x \notin \mathcal{C}(S)$  as required.

The existence of  $\mathcal{C}_{v,\delta}$  follows from the results of §2; we choose  $\mathcal{O}$  and  $p \in S_{\mathcal{O}}(R)$  so that  $v$  is the unit tangent vector to  $\overline{\mathcal{O}p}$ . Choose  $R$  so large that  $\alpha_{\mathcal{O}} < \delta$ ; this is possible for any  $\delta > 0$  by combining Lemma 2.4 with the estimate (2.7). On the other hand, we have the estimate  $\beta_{\mathcal{O}} \geq \beta_{\mathcal{O}} \geq \delta'(a, \delta, R) > 0$  again by (2.7) and Lemma 2.4. Thus  $x_v \notin \mathcal{C}_{v,\delta}$ , and so  $\mathcal{C}_{v,\delta}$  satisfies the required properties.

**Remark.** The property (3.3) is useful in providing barriers for systems of partial differential equations satisfying certain maximum principles; in particular, it can be applied to the study of complete minimal submanifolds in  $N^n$  and harmonic maps of complete manifolds into  $N^n$ . Gromov [10, 3.2] has also called attention to property (3.3), partly in regard to generalizing the theory of Kleinian groups.

#### 4. Harmonic measure and general boundary values

In this section we generalize our results on solvability of the Dirichlet problem to more general boundary values; we begin by introducing a Poisson integral representation for globally defined harmonic functions via the harmonic measure on  $S^{n-1}(\infty)$ . Theorem 4.3 then gives a satisfying relation between the class of bounded harmonic functions on  $N^n$  and the class of  $L^\infty$  functions on  $S^{n-1}(\infty)$ .

Given  $f \in C^0(S^{n-1}(\infty))$ , let  $P[f]$  denote the unique harmonic extension of  $f$  into  $N^n$ ; thus  $P[f]$  is the harmonic function on  $N^n$  with asymptotic boundary values  $f$  on  $S^{n-1}(\infty)$ . For each  $x \in N^n$ , define a linear functional  $L_x$  on  $C^0(S^{n-1}(\infty))$  by

$$(4.1) \quad L_x(f) = P[f](x).$$

Note that since  $|P[f](x)| \leq \max_{S^{n-1}(\infty)} |f|$  by the maximum principle (Proposition 1.1(a)),  $L_x$  is a bounded linear functional of norm 1. Again the maximum principle shows that  $f \geq 0 \Rightarrow L_x(f) \geq 0$  so that  $L_x$  is a positive functional. Applying the Riesz representation theorem gives the existence of a regular positive Borel measure  $\mu_x$  on  $S^{n-1}(\infty)$  such that

$$P[f](x) = \int_{S^{n-1}(\infty)} f d\mu_x,$$

for any  $x \in N^n, f \in C^0(S^{n-1}(\infty))$ . We note that the above remarks show that  $S^{n-1}(\infty)$  is the Šilov boundary of  $N^n$  in the sense of harmonic analysis. The measure  $\mu_x$  is called the *harmonic measure* of  $S^{n-1}(\infty)$  at  $x$ ; clearly  $\mu_x(S^{n-1}(\infty)) = 1$  for all  $x$ .

The following result gives a means of constructing harmonic functions with more general behavior at infinity.

**Theorem 4.1.** *Let  $f \in S^{n-1}(\infty) \rightarrow \mathbf{R}$  be  $\mu_x$ -integrable for some  $x \in N^n$ . Then  $f$  is  $\mu_x$ -integrable for all  $x \in N^n$ , and the function  $P[f]$  given by*

$$(4.2) \quad P[f](x) = \int_{S^{n-1}(\infty)} f \cdot d\mu_x$$

*is a smooth harmonic function on  $N^n$ .*

The function  $P[f]$  defined by (4.2) is called the *harmonic extension* of  $f$ . In particular, if  $E \subset S^{n-1}(\infty)$  is a Borel set, then  $\mu_x(E)$  is harmonic in  $x$ .

*Proof.* The proof is a straightforward adaptation of the same result for bounded regular domains in  $\mathbf{R}^n$ ; see [11, §3.6]. We sketch the argument for  $f$  upper-semicontinuous and bounded. Choose a sequence of continuous functions  $\{f_n\}$  decreasing monotonically to  $f$  as  $n \rightarrow \infty$ . The corresponding sequence of harmonic extensions  $P[f_n]$  given by (4.2) decrease to a limit  $u(x)$ , and  $u$  is harmonic in  $N^n$ ; this follows easily from the Harnack-type convergence of harmonic functions in  $N^n$ . It is not difficult to see that  $u$  is independent of  $\{f_n\}$ . By monotone convergence,  $\int_{S^{n-1}(\infty)} f_n d\mu_x \rightarrow \int_{S^{n-1}(\infty)} f d\mu_x$  for any  $x$ , so that  $u = P[f]$  is harmonic. To prove (4.2) for Borel measurable  $f$ , one uses the fact that  $f$  is the monotone limit of upper- and lower-semicontinuous functions; see [11] for details. *q.e.d.*

We now show that the harmonic extension  $P[f]$  of  $f$  has the correct boundary values, at least in certain cases.

Given a continuous function  $u: N^n \rightarrow \mathbf{R}$ , let  $u_t$  denote the restriction to  $S^{n-1}(t)$ , and  $u^t$  the pullback of  $u_t$  to  $S^{n-1}(1)$  via the exponential map. If  $v: S^{n-1}(\infty) \rightarrow \mathbf{R}$ , we also view  $v: S^{n-1}(1) \rightarrow \mathbf{R}$  via the homeomorphism  $\eta: S^{n-1}(1) \rightarrow S^{n-1}(\infty)$  (see §0). Finally, if  $\lambda$  denotes the measure induced by the volume form on  $S^{n-1}(1)$ , we view  $\lambda$  as a measure on  $S^{n-1}(t)$  for  $t \in [1, \infty]$  via the above homeomorphisms.

**Theorem 4.2.** *Let  $f \in L^p(S^{n-1}(\infty), \mu_x) \cap L^r(S^{n-1}(\infty), \lambda)$ ,  $1 < r < \infty$ , and let  $P[f]$  denote the harmonic extension of  $f$ . Then  $\|f - P[f]^t\|_r \rightarrow 0$  as  $t \rightarrow \infty$  in  $L^r(S^{n-1}(1), \lambda)$ . In particular,  $P[f]_t \rightarrow f$  almost everywhere in the cone topology on  $N^n$ .*

*Proof.* Since  $1 < r < \infty$ ,  $C^0$  is dense in  $L^r$ . Let  $f_n \in C^0(S^{n-1}(\infty))$  be chosen so that  $f_n \rightarrow f$  in  $L^r(S^{n-1}(\infty), \lambda) \cap L^p(S^{n-1}(\infty), \mu)$ , and set  $u_n = P[f_n]$ . By the solution to the Dirichlet problem, Theorem 3.2,  $(u_n)_t$  converges to  $f_n$  uniformly in the cone topology so that  $u_n^t \rightarrow f_n$  uniformly on  $S^{n-1}(1)$ . Let  $u = P[f]$ . Then  $u_n \rightarrow u$  uniformly in  $N^n$  since

$$\begin{aligned} |u(x) - u_n(x)| &\leq \int_{S^{n-1}(\infty)} |f - f_n| d\mu_x \\ &\leq \left( \int_{S^{n-1}(\infty)} |f - f_n|^p d\mu_x \right)^{1/p} = \|f - f_n\|_{p, \mu}. \end{aligned}$$



We have

$$\|f - u^t\|_{r,\lambda} \leq \|f - f_n\|_{r,\lambda} + \|f_n - u_n^t\|_{r,\lambda} + \|u_n^t - u^t\|_{r,\lambda};$$

the above remarks show that each term is arbitrarily small for  $t, n$  sufficiently large. q.e.d.

We are interested in the converse of the above theorems, i.e., given harmonic functions  $u$  on  $N^n$ , when does  $u$  have boundary values in  $L^p$ ? For the case of bounded harmonic functions, we have the following theorem.

**Theorem 4.3.** *Let  $\beta$  denote the Banach space of bounded harmonic functions on  $N^n$  under the sup norm. Then the linear mapping*

$$P: L^\infty(S^{n-1}(\infty), \mu) \rightarrow \beta, \quad P(f) = P[f]$$

is a norm-nonincreasing isomorphism onto  $\beta$ . Further  $\|P[f]^t - f\|_p \rightarrow 0$  as  $t \rightarrow \infty$  for any  $1 < p < \infty$  provided  $f \in L^\infty(S^{n-1}(\infty), \lambda)$ .

*Proof.* It is clear that  $P$  is linear and 1-1.  $P$  is norm-nonincreasing since

$$|P[f](x)| = \int_{S^{n-1}(\infty)} f d\mu_x \leq \|f\|_\infty \cdot \int_{S^{n-1}(\infty)} d\mu_x = \|f\|_\infty.$$

Thus the major task lies in showing  $P$  is surjective; the proof of this is a variation of the proof by Ullrich in the case of the Bergmann ball; see [13, §4.3].

There is a natural action of  $SO(n)$  on  $S^{n-1}(t)$  induced by the linear action of  $SO(n)$  on the Euclidean  $t$ -spheres in  $TN^n$ ; of course, the action on  $S^{n-1}(t)$  is not by isometries. Let  $dg$  denote Haar measure on  $SO(n)$  and choose a continuous nonnegative function  $h: SO(n) \rightarrow \mathbf{R}$  such that  $\int_{SO(n)} h dg = 1$ . Given  $F(z): N^n \rightarrow \mathbf{R}$  harmonic, define

$$(4.3) \quad G(z) = \int_{SO(n)} F(gz) \cdot h(g) dg.$$

We claim there is a fixed constant  $k$ , depending only on the geometry of  $N$  such that

$$(4.4) \quad \int_{SO(n)} |F'(g\xi)|^p dg \leq k \cdot \int_{S^{n-1}(1)} |F'(x)|^p d\lambda,$$

for any  $\xi \in S^{n-1}(1)$ . To see this we have

$$\begin{aligned} \int_{S^{n-1}(1)} \left[ \int_{SO(n)} |F'(g\eta)|^p dg \right] d\lambda &= \int_{SO(n)} \left[ \int_{S^{n-1}(1)} |F'(g\eta)|^p d\lambda \right] dg \\ &= \int_{SO(n)} \left[ \int_{S^{n-1}(1)} |F'(\eta)|^p d((g^{-1})^*\lambda) \right] dg \\ &\leq k \int_{SO(n)} \left[ \int_{S^{n-1}(1)} |F(\eta)|^p d\lambda \right] dg \\ &= k \cdot \int_{S^{n-1}(1)} |F(\eta)|^p d\lambda, \end{aligned}$$

where  $k = \sup_g \{ \sup_x [h_g(x) : x \in S^{n-1}(1)] \}$ ,  $h_g =$  Radon-Nikodym derivative of  $(g^{-1})^*\lambda$  with respect to  $\lambda$ . On the other hand, since  $SO(n)$  acts transitively on  $S^{n-1}(1)$ ,

$$\int_{S^{n-1}(1)} \left[ \int_{SO(n)} |F'(t\eta)|^p dg \right] d\lambda = \text{vol } S^{n-1}(1) \cdot \int_{SO(n)} |F'(g\xi)|^p dg,$$

for any  $\xi \in S^{n-1}(1)$ ; this proves (4.4). By the Hölder inequality applied to (4.3), using (4.4) we have

$$(4.5) \quad |G(z)| \leq k \cdot M_p \|h\|_q,$$

where  $p, q$  are conjugate indices, and  $M_p = \sup_t \|F^t\|_p$ ; in particular,

$$(4.6) \quad |G(z)| \leq k \cdot M_\infty.$$

These estimates show that  $\{G^t\}$ ,  $t \in (1, \infty)$ , is an equicontinuous family on  $S^{n-1}(1)$ . In fact, given  $\varepsilon > 0$ , choose a neighborhood  $U$  of  $I \subset SO(n)$  such that

$$|h(g) - h(gg_0^{-1})| < \varepsilon$$

for  $g_0 \in U$ ,  $g \in SO(n)$ . Choose  $\delta > 0$  so that if  $\xi, \eta \in S^{n-1}(1)$  and  $\text{dist}(\xi, \eta) < \delta$ , then  $\xi = g_0\eta$  for  $g_0 \in U$ . Since  $G(g_0z) = \int_{SO(n)} F(gz)h(gg_0^{-1}) dg$ , one finds

$$|G^t(\xi) - G^t(\eta)| \leq \int_{SO(n)} |F^t(g\eta)| \cdot |h(gg_0^{-1}) - h(g)| dg \leq M_1 \varepsilon,$$

provided  $\text{dist}(\xi, \eta) < \delta$ . By (4.6),  $\{G^t\}$  is uniformly bounded, so it follows that there is a sequence  $\{t_i\} \rightarrow \infty$  such that  $\{G^{t_i}\}$  converges uniformly to  $g \in C^0(S^{n-1}(\infty))$ .

Now replace  $h$  by  $h_j$  in the definition of  $G$ , and let  $\text{supp}\{h_j\}$  shrink to  $I \in SO(n)$ . Then  $G_j$  converges pointwise to  $F$ . Consider the sequence  $\{g_j\}$ ; by (4.6) and the uniform convergence of  $G_j^t$  to  $g_j$ ,  $\|g_j\|_{p,\mu} < k \cdot M_\infty$  for all

$1 < p \leq \infty$ . Thus there is an  $f \in L^\infty(S^{n-1}(\infty), \mu)$  such that  $\{g_j\}$  has a subsequence, call it  $\{g_j\}$ , converging to  $f$  in the weak \*-topology of  $L^\infty$ . The function

$$P[f](x) = \int_{S^{n-1}(\infty)} f d\mu_x$$

is well defined and harmonic in  $N^n$ ; we will show that  $F = P[f]$ . To do this we first prove that

$$(4.7) \quad F^t - P[f]^t \xrightarrow{t \rightarrow \infty} 0 \text{ weakly in } L^\infty(S^{n-1}(\infty), \lambda)^* \cap L^\infty(S^{n-1}(\infty), \mu)^*.$$

Clearly,

$$(4.8) \quad F^t - P[f]^t = (F^t - G_j^t) + (G_j^t - P[g_j]^t) + (P[g_j]^t - P[f]^t),$$

and we will analyze each of these terms separately. Let  $\sigma$  denote either of the measures  $\lambda$  or  $\mu$ .

(i)  $(F^t - G_j^t)$ : We compute, for  $l \in L^1(S^{n-1}(1), \sigma)$ ,

$$\begin{aligned} & \int_{S^{n-1}(1)} (F^t(z) - G_j^t(z))l(z) d\sigma \\ &= \int_{S^{n-1}(1)} \left[ \int_{SO(n)} (F^t(z) - F^t(gz)) \cdot l \cdot h_j dg \right] d\sigma \\ &= \int_{SO(n)} \left[ \int_{S^{n-1}(1)} (F^t(z) - F^t(gz)) \cdot l d\sigma \right] h_j dg. \end{aligned}$$

Note that since  $F$  is bounded on  $N$ , there are a sequence  $\{t_i\} \rightarrow \infty$  and  $\phi \in L^\infty(S^{n-1}(1), \sigma)$  such that  $\{F^{t_i}\} \rightarrow \phi$  in the weak \*-topology on  $L^\infty$ . Thus

$$\begin{aligned} & \lim_{t \rightarrow \infty} \int_{S^{n-1}(1)} (F^{t_i}(z) - G_j^{t_i}(z))l(z) d\sigma \\ &= \int_{SO(n)} \left[ \int_{S^{n-1}(1)} (\phi(z) - \phi(gz))l d\sigma \right] \cdot h_j dg. \end{aligned}$$

This shows that, given any  $\epsilon > 0$  and  $l \in L^1(\sigma)$ , there is a  $J$  such that for  $j \geq J$ , the last term is bounded, in absolute value, by  $\epsilon$ . The same statement holds for any sequence  $\{t_j\} \rightarrow \infty$ .

In the next case, we can estimate in the strong  $L^p$ -norm.

$$\begin{aligned} (ii) \quad \lim_{t \rightarrow \infty} \|G_j^t - P[g_j]^t\|_{p,\sigma}^p &= \lim_{t \rightarrow \infty} \int_{S^{n-1}(1)} |G_j^t(z) - P[g_j]^t(z)|^p d\sigma \\ &\leq \lim_{t \rightarrow \infty} \int_{S^{n-1}(1)} |G_j^t(z) - g_j(z)|^p d\sigma \\ &\quad + \lim_{t \rightarrow \infty} \int_{S^{n-1}(\infty)} |P[g_j]^t(z) - g_j(z)|^p d\sigma. \end{aligned}$$

Since  $G_j^t \rightarrow g_j$  and  $P[g_j]^t \rightarrow g_j$  uniformly on  $S^{n-1}(1)$ , the last two terms vanish.

(iii) To estimate  $\lim_{t \rightarrow \infty} (P[g_j]^t \rightarrow P[f]^t)$  in the weak \*-topology, we just note that  $P[g_j]^t \rightarrow g_j$  uniformly for fixed  $j$ , and  $P[f]^t \rightarrow f$  in  $L^p$ -norm,  $1 < p < \infty$ ; further  $g_j \rightarrow f$  weakly in  $L^\infty(S^{n-1}(1), \sigma)$ .

Combining the estimates in (i), (ii) and (iii) with (4.8) gives the desired (4.7). Finally, since  $F - P[f]$  is harmonic, bounded and  $(F - P[f])^t \rightarrow 0$  weakly as  $t \rightarrow \infty$  we claim that  $F = P[f]$ . Thus let  $\psi = F - P[f]$  and let  $P_t^x d\lambda$  denote harmonic measure on  $S^{n-1}(t)$  at  $x$ . Then we have

$$\begin{aligned} \psi(x) &= \int_{S^{n-1}(1)} \psi^t(y) P_t^x(y) d\lambda \\ &= \int_{S^{n-1}(1)} \psi^t(y) \cdot d\mu^x(y) + \int_{S^{n-1}(1)} \psi^t(y) \cdot [P_t^x(y) d\lambda - d\mu^x(y)]. \end{aligned}$$

By assumption the first term converges to 0 as  $t \rightarrow \infty$ , and by general principles the second term also converges to 0 since  $\|\psi\|_\infty$  is finite.

This shows  $F = P[f]$  as desired. Finally, the fact that  $\|P[f]^t - f\|_p \rightarrow 0$  as  $t \rightarrow \infty$  follows immediately from Theorem 4.2.

## 5. Concluding remarks

1. It is not difficult to see that Theorem 3.2 remains valid provided the conditions on the curvature hold outside some compact set  $B$  of  $N^n$ ; in other words  $a^2 \leq K_N \leq -1$  on  $N^n - B$ , where  $B$  is any compact set in  $N^n$ , and  $N^n - B$  is diffeomorphic to  $\mathbf{R}^n - \bar{B}$ ,  $\bar{B}$  being a closed ball. This is in line with the philosophy of Greene-Wu that much of the function theory on Cartan-Hadamard manifolds should be determined by asymptotic conditions only.

2. Let  $0 \in N^n$  and let  $\partial/\partial x_i$  be global normal coordinates for  $N^n$  around 0. Setting  $g_{ij} = \langle \partial/\partial x_i, \partial/\partial x_j \rangle$ , the Laplace-Beltrami operator takes the form

$$\Delta u = \frac{1}{\sqrt{g}} \sum_{i,j} \frac{\partial}{\partial x_i} \left( \sqrt{g} g^{ij} \frac{\partial u}{\partial x_j} \right),$$

where  $g = \det(g_{ij})$ , and  $g^{ij} = (g_{ij})^{-1}$ . Then Theorem 3.2 may be interpreted as showing that this operator admits many bounded solutions in  $\mathbf{R}^n$ , provided  $a^{ij} = \sqrt{g} g^{ij}$  satisfies certain curvature conditions—certain bounds on 2nd and 3rd derivatives of  $a^{ij}$ .

## References

- [1] L. V. Ahlfors, *Sur le type d'une surface de Riemann*, C. R. Acad. Sci. Paris **201** (1935) 30–32.
- [2] J. Cheeger & D. Ebin, *Comparison theorems in Riemannian geometry*, North Holland, Amsterdam, 1975.
- [3] S. Y. Cheng & S. T. Yau, *Differential equations on Riemannian manifolds and their geometric applications*, Comm. Pure Appl. Math. **28** (1975), 333–354.
- [4] H. I. Choi, Thesis, University of California, Berkeley, 1982.
- [5] P. Eberlein & B. O'Neill, *Visibility manifolds*, Pacific J. Math. **46** (1973) 45–109.
- [6] D. Gilbarg & N. Trudinger, *Elliptic partial differential equations of second order*, Springer, New York, 1977.
- [7] R. E. Greene & H. Wu, *Analysis on noncompact Kähler manifolds*, Proc. Sympos. Pure Math. Part 2, Vol. 30, Amer. Math. Soc., 1977, 69–100.
- [8] ———, *Function theory on manifolds which possess a pole*, Lecture Notes in Math. Vol. 699, Springer, New York, 1979.
- [9] ———,  *$C^\infty$  Approximations of convex, subharmonic and plurisubharmonic functions*, Ann. Sci. École Norm. Sup. **12** (1979) 47–84.
- [10] M. Gromov, *Hyperbolic manifolds, groups and actions*, Proc. Stony Brook Conf. on Riemann Surfaces, Annals of Math. Studies, No. 97, Princeton University Press, Princeton, 1981.
- [11] W. K. Hayman & P. B. Kennedy, *Subharmonic functions*, Vol. 1, Academic Press, New York, 1976.
- [12] A. Koranyi, *Harmonic functions on symmetric spaces; symmetric spaces*, Edited by W. M. Boothby and G. L. Weiss, Marcel Dekker, New York, 1972, 379–412.
- [13] W. Rudin, *Function theory in the unit ball of  $\mathbf{C}^n$* , Springer, New York, 1980.
- [14] Y. T. Siu & S. T. Yau, *Complete Kähler manifolds with non-positive curvature of faster than quadratic decay*, Ann. of Math. **105** (1977) 225–264.
- [15] D. Sullivan, *The Dirichlet problem at infinity for a negatively curved manifold*, preprint, Institut des Hautes Études Scientifiques.
- [16] S. T. Yau, *Harmonic functions on complete Riemannian manifolds*, Comm. Pure Appl. Math. **28** (1975) 201–228.
- [17] ———, *Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry*, Indiana Univ. Math. J. **25** (1976) 659–670.
- [18] ———, *Survey on partial differential equations in differential geometry*, Annals of Math. Studies, Vol. 102, Ed. S. T. Yau, Princeton University Press, Princeton, 1982.

RICE UNIVERSITY